Perez et al. Resubmission

A novel, ataxic mouse model of Ataxia Telangiectasia caused by a clinically relevant nonsense mutation

Perez, Harvey ^{‡1}; Abdallah, May, F. ^{‡1}; Chavira, Jose, I. ^{‡1}; Egeland, Martin, T. ¹; Vo, Karen, L. ¹; Buechsenschuetz, Callan, L. ¹; Sanghez, Valentina ¹; Kim, Jeannie, L. ¹; Pind, Molly ², Nakamura, Kotoka ³, Hicks, Geoffrey, G. ²; Gatti, Richard, A. ³; Madrenas, Joaquin ^{1,5}; Iacovino, Michelina ^{1,4}; McKinnon, Peter, J. ⁶; Mathews, Paul, J. ^{*1,7}

1. The Lundquist Institute for Biomedical Innovation, Harbor-UCLA Medical Center, Torrance, CA
2. Department of Biochemistry & Medical Genetics, Max Rady College of Medicine, University of Manitoba
3. Department of Pathology & Laboratory Medicine, David Geffen School of Medicine, University of California, Los Angeles, CA
4. Department of Pediatrics, Harbor-UCLA Medical Center, Torrance, CA
5. Department of Medicine, Harbor-UCLA Medical Center, Torrance, CA
6. Center for Pediatric Neurological Disease Research, St. Jude Translational Neuroscience, St. Jude Children's Research Hospital, Memphis, TN, USA
7. Department of Neurology, Harbor-UCLA Medical Center, Torrance, CA

[‡]Contributed equally

^{*}Corresponding Author

Abstract

Ataxia Telangiectasia (A-T) and ataxia with ocular apraxia type 1 (AOA1) are devastating neurological disorders caused by null mutations in the genome stability genes, A-T mutated (*ATM*) and aprataxin (*APTX*), respectively. Our mechanistic understanding and therapeutic repertoire for treating these disorders is severely lacking, in large part due to the failure of prior animal models with similar null mutations recapitulating the characteristic loss of motor coordination (i.e., ataxia) and associated cerebellar defects. By increasing genotoxic stress, through the insertion of null mutations in both the *Atm* (nonsense) and *Aptx* (knockout) genes in the same animal, we have generated a novel mouse model that for the first time progressively develops a severe ataxic phenotype associated with atrophy of the cerebellar molecular layer. We find biophysical properties of cerebellar Purkinje neurons are significantly perturbed (e.g., reduced membrane capacitance, lower action potential thresholds, etc.), while properties of synaptic inputs remain largely unchanged. These perturbations significantly alter

Perez et al. Resubmission

Purkinje neuron neural activity, including a progressive reduction in spontaneous action potential firing frequency that correlates with both cerebellar atrophy and ataxia over the animal's first year of life. Double mutant mice also exhibit a high predisposition to developing cancer (thymomas) and immune abnormalities (impaired early thymocyte development and T-cell maturation), symptoms characteristic of the disorder A-T. Lastly, by inserting a clinically relevant nonsense type null mutation in *Atm*, we demonstrate that small molecule readthrough (SMRT) compounds can restore ATM production, indicating their potential as a future A-T therapeutic.

1.0 Introduction

Ataxia Telangiectasia (A-T) is a rare (1 in ~100,000) (Swift et al. 1986), autosomal recessive genetic disorder characterized by cancer predisposition, immune deficiency, and a highly penetrant progressive and severe ataxia linked to cerebellar atrophy (Rothblum-Oviatt et al. 2016; Boder and Sedgwick 1958; Levy and Lang 2018). A-T patients typically die in their second and third decade of life (Crawford et al. 2006) from lymphatic cancers, respiratory infections, or debilitating ataxia—unfortunately, survivability has not dramatically changed since the 1950s (Micol et al. 2011; Rothblum-Oviatt et al. 2016). While disease progression and cause of death vary widely across patients, the highly penetrant progressive decline in motor coordination is reported as having the greatest negative impact on a patient's quality of life (Jackson et al. 2016). Care is generally palliative, directed at reducing, limiting, or eliminating cancers or infections. No long-term therapies are available for treating the ataxia and associated cerebellar dysfunction and atrophy.

A-T is caused by deficiency or dysfunction of the ATM (A-T mutated) protein (Savitsky et al. 1995). Premature termination codon (PTC) causing nonsense mutations account for up to a half of known cases with missense and deletions also contributing (Concannon and Gatti 1997; Sandoval et al. 1999). ATM is a serine/threonine PIKK family kinase that plays a key role in the DNA damage response (DDR), protecting cells from the tens of thousands of DNA lesions incurred each day (Lindahl and Barnes 2000; Kastan and Bartek 2004; Shiloh and Ziv 2013). In the active monomeric form, ATM phosphorylates several key proteins halting the production of new DNA (cell cycle arrest) (Ando et al.

Perez et al. Resubmission

2012), and then, depending on severity of the damage, initiating DNA repair or programmed cell death (i.e., apoptosis) (Ando et al. 2012; Rashi-Elkeles et al. 2006). Several downstream DDR pathway targets of ATM have been identified, including p53, CHK2, BRCA1, SMC1, and NBS1 (Matsuoka et al. 2007). ATM's role in DNA repair is also implicated in normal immune system development, where it is proposed to contribute to the recombination of natural DNA splicing that occurs during gene rearrangement in T- and B-lymphocyte maturation (Chao, Yang, and Xu 2000; Matei, Guidos, and Danska 2006; Vacchio et al. 2007; Schubert, Reichenbach, and Zielen 2002). Although its roles are still emerging, ATM has also been implicated in oxidative stress homeostasis (Guo et al. 2010) and mitophagy (Valentin-Vega and Kastan 2012; Pizzamiglio, Focchi, and Antonucci 2020).

A mechanistic understanding of why ATM deficiency causes ataxia is still under debate, but it is far from the only DDR protein linked to ataxia, as aprataxin (APTX) (Aicardi et al. 1988), meiotic recombination 11 homolog 1 (MRE11) (Sedghi et al. 2018), nibrin (NBS1) (van der Burgt et al. 1996), senataxin (SETX) (Moreira et al. 2004), and tyrosyl-DNA phosphodiesterase 1 (TDP1) (Takashima et al. 2002) when absent or dysfunctional can cause cerebellar-related ataxia. This suggests that the neurological features of genome instability syndromes have a common underlying cause, although this idea is still to be mechanistically demonstrated (McKinnon 2009; Rass, Ahel, and West 2007).

A major factor limiting our ability to define why loss of DDR proteins, like ATM, selectively impacts the cerebellum and causes progressive ataxia is the lack of an animal model that recapitulates these neurological symptoms (Lavin 2013). Several A-T rodent models have been created over the past several years by inserting gene mutations that cause protein dysfunction (lack kinase activity) or complete deficiency (Herzog et al. 1998; Xu and Baltimore 1996; Elson et al. 1996; Spring et al. 2001; Campbell et al. 2015; Quek et al. 2016; Tal et al. 2018; Lavin 2013); a minipig was also recently reported (Beraldi et al. 2017). Unfortunately, none acquire an overt, progressive ataxia with cerebellar dysfunction and atrophy that recapitulates the human disease, even though other aspects of the disorder like thyroid cancers, infertility, and immune abnormalities do develop. It remains unclear why these prior animal models fail to display the progressive ataxic phenotype (Lavin 2013). It is possible

Perez et al. Resubmission

that species specific molecular compensations in mice provide redundancies or alternative pathways minimizing the effects of ATM deficiency in the brain (El-Brolosy and Stainier 2017). It is also possible that the shortened lifespan of prior models (Barlow et al. 1996) is too brief for the stochastic mechanisms driving cerebellar dysfunction and atrophy to accumulate and impact motor behavior. Other challenges include potentially leaky genetic manipulations that result in low levels of ATM protein or active fragments with residual kinase activity, thus limiting neuropathology (Li et al. 2011). The impact of missing such a crucial animal model has been significant, severely limiting not only experimental studies from identifying the cellular and molecular mechanisms but hampering pre-clinical development and testing of much needed therapeutics.

We test here whether increasing genotoxic stress, by placing null mutations in not just the *Atm* gene, but also the related *Aptx* gene, leads to a more representative mouse model that displays cerebellar dysfunction, atrophy, and the development of progressive ataxia. We chose to additionally knock-out *Aptx* because its deficiency causes an A-T like disorder in humans called ataxia with ocular apraxia type 1 (AOA1), but without A-T's other system defects that could increase the potential for prenatal lethality or early death (e.g., immunodeficiency and cancer predisposition) (Coutinho P 2002). Moreover, APTX is a phosphodiesterase involved in DNA reassembly after double and single stranded repair, having a function downstream of, but not directly regulated or related to ATM (Gueven et al. 2004; Schellenberg, Tumbale, and Williams 2015; Ahel et al. 2006). We therefore reasoned that deficiency of both proteins would have an additive effect on genotoxic stress capable of inducing a detectable neurological dysfunction. Our results indeed demonstrate that mice deficient in ATM and APTX develop cerebellar dysfunction, atrophy, and a progressive and profound ataxia, while deficiency in either protein alone do not. Additionally, double mutants acquired several additional characteristic symptoms of A-T, including defects in immune maturation and a high incidence of cancer (thymomas), making it the most representative model, from a phenotypic standpoint, to date.

Finally, to improve the clinical and translational relevance of this new mouse model, we inserted a point mutation (103C>T) in the *Atm* gene common to a large family North African A-T patients. This mutation

Perez et al. Resubmission

results in a premature termination codon (PTC) at what would normally be amino acid 35 resulting in loss of ATM production. As a result of this genotypic mutation, we were then able to conduct proof-of-principle experiments demonstrating the ability of so-called PTC readthrough therapeutics to restore production of the ATM protein. This includes the Small Molecule Readthrough (SMRT) compound, GJ103 we are currently developing (Du et al. 2013).

2.0 Results

2.1 Creation of a new A-T mutant mouse model expressing a clinically relevant nonsense mutation

To create a more clinically relevant mouse model of A-T we used a gateway recombination cloning and site-directed mutagenesis method to recapitulate a c.103C>T (p.R35X) mutation in the *ATM* gene found in a large population of North African A-T patients (**Fig. 1A and Methods**) (Gilad et al. 1996). The insertion of thymine in place of cytosine at this site in exon 3 results in a premature termination codon (PTC) causing nonsense mutation in the *ATM* gene. Since the c.103C>T mutation results in different PTCs in the human compared to the mouse *Atm* gene, TGA vs. TAG respectively, we created two different mice by exchanging the mouse *Atm* exon 3 with either a human or mouse exon 3 variant with the c.103C>T mutation (**Fig. 1B**). In the human variant, a 103C>T mutation of the mouse codon, where the arginine (R) encoding codon (CGA) becomes a TGA stop codon, results in a mouse we denote as *Atm*^{R35X} (officially *Atm*^{Tm1.1(103CAG)TGA)Mfgc}). In the mouse variant, the c.103C>T mutation transforms a glutamine (Q) encoding CAG codon into a TAG stop codon and is denoted *Atm*^{Q35X} (officially *Atm*^{Tm1.1(103CT)Mfgc}). The presence of the PTC results in a loss of ATM expression, either reduced by about half in the heterozygote expressing one normal mouse copy of the *Atm* gene (*Atm*^{R35X/+} or *Atm*^{Q35X/+}), or completely in the homozygote (*Atm*^{R35X/R35X} or *Atm*^{Q35X/Q35X}) (**Fig. 1C**).

Atm^{R35X/R35X}, *Aptx*^{-/-} mice were created by first crossing single mutant *Atm*^{R35X/R35X} (congenic on the C57BL/6J background) and *Aptx*^{-/-} (mixed C57BL/6J and 129 background) mice to generate double mutant heterozygote *Atm*^{R35X/+}; *Aptx*^{+/-} mice. F1-5 littermate *Atm*^{R35X/+}; *Aptx*^{+/-} mice were then crossed within litters to create sufficient numbers of the desired experimental and control genotypes to

Perez et al. Resubmission

determine how loss of different amounts of ATM and APTX affect the animal's phenotype (**Fig. 1D**). Like prior ATM deficient A-T mouse models, ATM or APTX deficiency alone did not result in mice with ataxia (**Video 1 and 2**). However, deficiency in both proteins (*Atm*^{R35X/R35X}; *Aptx*^{-/-}) results in the development of a severe and progressively ataxic phenotype (**Fig. 1E, Video 3 and 4**).

2.2 ATM deficient mice have lowered survivability and a high incidence of thymomas

We assessed the general health and development of control and experimental mice expressing different levels of ATM and APTX (**Fig. 2**). We found that *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice grew ~55% slower and reached estimated plateau weights that were ~35% less than control genotypes (log-rank, n = 21 to 40, p<0.0001; **Fig. 2A**). These differences in weight were a postnatal phenomenon, as no significant weight differences were detected just after birth (P8) across all genotypes (1-way ANOVA, n = 5 to 23, p>0.23). Adolescent double mutant mice at postnatal day 45 (P45) weighed on average 30% less in males double mutant: 14.4 ± 1.0 g (n = 13) vs. wildtype: 20.2 ± 0.5 g (n = 16), *t*-test, p<0.0001 and 25% less in females [double mutant: 12.7 ± 0.6 g (n = 17) vs. wildtype: 17.0 ± 0.2 g (n = 15), *t*-test, p<0.0001; **Fig. 1A**]. Differences across the control genotypes were observed, but they were small and not consistent across time points or sex and therefore judged to not be physiologically relevant (**Fig. 2A**). Survivability of the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice was significantly reduced compared to *Atm*^{+/-}; *Aptx*^{+/-} mice, with 53% of mice still alive at 400 days of age, compared to 97% of *Atm*^{+/-}; *Aptx*^{+/-} mice at the same time point (**Fig. 2B**). ATM deficiency alone was sufficient to reduce survivability, as compared to *Atm*^{+/-}; *Aptx*^{+/-} mice, both *Atm*^{R35X/R35X}; *Aptx*^{+/-} and *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice had significantly reduced survivability rates [42%, log-rank, $\chi^2_{(1, 56)} = 13.49$, p=0.0002 and 52%, log-rank, $\chi^2_{(1, 53)} = 19.54$, p<0.0001, respectively]. No significant difference between ATM deficient mice with partial or complete APTX deficiency was detected [log-rank, $\chi^2_{(2, 85)} = 1.01$, p=0.6]. Conversely, mice harboring at least one functional copy of the *Atm* gene had normal survivability rates, regardless of whether they expressed APTX or not [log-rank, $\chi^2_{(3, 131)} = 3.08$, p=0.4]. No significant difference between male and female mice was observed and thus data were pooled [log-rank, p>0.4 for all pairwise comparisons; **Fig. 2-figure S1B**]. Generally, a third of mice with ATM deficiency died from complications related to large thymic cancers

Perez et al. Resubmission

(thymoma) found in the thoracic cavity (**Fig. 2C**). The presence or absence of APTX did not impact cancer prevalence, and mice with at least one *Atm* transcript were cancer free up until at least P400. Overall, ATM, but not APTX deficiency had severe effects on the health and survivability of mice.

2.3 Both ATM and APTX deficiency are necessary to produce progressive motor dysfunction

The progressive development of severe ataxia is a hallmark characteristic of A-T that is recapitulated in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice but none of the other control genotypes we tested. Overall, we find motor coordination deficits emerge between 210 and 400 days after birth in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice and find no evidence of ataxia in mice with at least one copy of the *Atm* or *Aptx* gene (**Fig. 3A, B**). For the vertical pole test, *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice took twice as long to descend at P400 compared to *Atm*^{+/+}; *Aptx*^{+/+}, *Atm*^{+/+}; *Aptx*^{-/-}, *Atm*^{R35X/R35X}; *Aptx*^{+/+}, or *Atm*^{R35X/+}; *Aptx*^{-/-} mice [Male: 29.1 ± 0.9 s (n = 3) vs. 7.5 ± 0.4 s (n = 12), 12.5 ± 2.5 s (n = 9), 9.2 ± 0.9 s (n = 10), 8.6 ± 0.9 s (n = 11), 1-way ANOVA, $F_{(4, 40)} = 19.9$, $p < 0.0001$; Female: 19.0 ± 4.0 s (n = 4) vs. 7.5 ± 0.4 s (n = 12), 7.8 ± 0.4 s (n = 10), 10.5 ± 1.2 s (n = 6), 8.2 ± 0.5 s (n = 8), 1-way ANOVA, $F_{(4, 35)} = 13.9$, $p < 0.0001$]. An examination of gait indicated that *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice at P400, but not P210 need additional stabilization during ambulation, as they spend twice as much time with 3 paws, rather than the normal 2 in contact with the ground as they walk across the gait analysis platform [Male: 56.2 vs. 26.4 to 32.2 %, 1-way ANOVA, $F_{(4, 54)} = 14.3$, $p < 0.0001$; Female: 58.4 vs. 18.9 to 28.8 %, 1-way ANOVA, $F_{(3, 178)} = 95.5$, $p < 0.0001$; **Fig. 3B**]. *Atm*^{R35X/R35X}; *Aptx*^{-/-} also display a slower cadence and average speed across the platform compared to all other genotypes at P400 [cadence, Male: 9.5 vs. 13.3 to 15.9 steps/s, 1-way ANOVA, $F_{(3, 204)} = 36.8$, $p < 0.0001$; Female: 9.1 vs. 14.2 to 15.9 steps/s, 1-way ANOVA, $F_{(3, 204)} = 39.7$, $p < 0.0001$; speed, Male: 8.8 vs. 22 to 26 cm/s, 1-way ANOVA, $F_{(4, 50)} = 28.3$, $p < 0.0001$; Female: 58.4 vs. 18.9 to 28.8 cm/s, 1-way ANOVA, $F_{(3, 178)} = 39.7$, $p < 0.0001$; **Fig. 3B**; **Fig. 3-fig. S1**]. This difference in speed and cadence is unlikely due to animal size, as there are no significant differences in these parameters at earlier time points when the difference in size is significant (**Fig. 2A**). These observations across the two behavioral tests were found in both male and female mice at each of their respective time points, consistent with the lack of sex differences observed in A-T patients.

Perez et al. Resubmission

We further examined behavioral differences between the *Atm*^{R35X/R35X}, *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice using a standardized set of experimental procedures used to phenotype genetically modified mice (i.e., SHIRPA; **Fig. 3C**; **Fig. 3-fig. S1**) (Rogers et al. 1997). We first detected differences in motor function at P8, where *Atm*^{R35X/R35X}, *Aptx*^{-/-} mice took 3-4 times longer on average to right themselves compared to *Atm*^{+/+}; *Aptx*^{+/+} mice [Male: 6.4 ± 1.1 s (n = 24) vs. 1.5 ± 0.1 s (n = 23), *t*-test, *p*<0.0002; Female: 11.1 ± 1.9 s (n = 21) vs. 2.4 ± 0.3 s (n = 17), *t*-test, *p*<0.0002; **Fig. 3C bottom**]. At 30-days of age, we detected significant differences between *Atm*^{R35X/R35X}, *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice in behavioral tests that qualitatively measure body position and spontaneous activity (**Fig. 3C**). Striking differences in *Atm*^{R35X/R35X}, *Aptx*^{-/-} compared to *Atm*^{+/+}; *Aptx*^{+/+} mice were observed at P400, especially for behaviors related to movement, including locomotor activity, body position, and gait (**Fig. 3C**). The results from this battery of tests demonstrates that *Atm*^{R35X/R35X}, *Aptx*^{-/-} mice develop a severe change in behavior by P400, consistent with purely visual observations of significant motor coordination deficits in the mice up to this time point. Importantly, we do not find any significant differences between the other control genotypes, including *Atm*^{R35X/+}; *Aptx*^{-/-} mice that express at least some ATM but no APTX protein (**Fig. 3-fig. S1**).

2.4 The membrane and synaptic properties are perturbed in ATM and APTX deficient neurons of the cerebellum

Purkinje neurons (PN) are a key neuronal subtype located in the cerebellar cortex. They display considerable intrinsic excitability, firing action potentials spontaneously at rates significantly higher than most other neurons in the brain (50 to 100 Hz more in many cases). Their activity shapes cerebellar output via tonic inhibition of neurons of the cerebellar nuclei, which project to motor coordination centers in the forebrain, brainstem, and spinal cord. Dysfunction of cerebellar PNs is associated with several forms of ataxia and implicated in A-T (Hoxha et al. 2018; Cook, Fields, and Watt 2020; Shiloh 2020). We therefore examined if the electrophysiological properties of PNs in the *Atm*^{R35X/R35X}, *Aptx*^{-/-} cerebellum were abnormal.

Perez et al. Resubmission

Since PN baseline activity and responsivity to input is mediated by a baseline set of passive and active membrane properties (**Fig. 4**), we directly recorded from and compared the membrane properties of PNs in acute cerebellar slices harvested from *Atm*^{R35X/R35X}; *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice (P350 to 400). PNs recorded from *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice had significantly “tighter” membranes, displaying higher membrane input resistances (R_m) than those from *Atm*^{+/+}; *Aptx*^{+/+} mice [47.7 ± 5.6 (n = 15) vs. 30.2 ± 1.47 (n = 23) M Ω , *t*-test, *p*=0.008; **Fig. 4B**]. They also displayed a faster membrane time constant (τ) [3.6 ± 0.4 (n = 15) vs. 5.1 ± 0.3 (n = 23) ms, *t*-test, *p*=0.009; **Fig. 4B**]. These results indicate that the total membrane capacitance ($C_m = \tau/R_m$) of the *Atm*^{R35X/R35X}; *Aptx*^{-/-} PNs is significantly reduced [98.25 ± 19.23 (n = 15) vs. 175.6 ± 12.67 (n = 23) pF, *t*-test, *p*=0.0025; **Fig. 4B**]. At the cellular level, this suggests that the ATM and APTX deficient PNs have less (i.e., smaller in size) or thinner membranes than that of wildtype PNs; a result suggestive of a developmental deficit or neurodegenerative process (Dell’Orco et al. 2015). We next assessed the intrinsic excitability of PNs in *Atm*^{R35X/R35X}; *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice by examining PN action potential (AP) generation and dynamics. Significant deficits in the ability of PNs to fire continuously in response to current injection were observed in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice (**Fig. 4C**). These deficits were associated with significant perturbations to the amplitude, threshold, and area of evoked action potentials [amplitude: 66.2 ± 0.7 (n = 14) vs. 72.1 ± 1.4 (n = 13) Δ mV, *t*-test, *p* = 0.003; threshold: -55.2 ± 1.5 vs. -48.61 ± 1.9 mV, *t*-test, *p*=0.0196; area: 17.96 ± 0.6 vs. 20.63 ± 1.0 mV*ms, *t*-test, *p*=0.048; **Fig. 4C**]. Together, these experiments demonstrate significant perturbations of PN physiological properties that likely perturb their ability to function normally in the cerebellum of *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice.

We next tested whether extrinsic and/or synaptic PN properties were also impacted in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice. We first examined spontaneous mini excitatory post synaptic currents (mEPSC) generated by granule cell-to-PN synapses (i.e., parallel fiber inputs). No difference in mEPSC size was detected, indicating the function of granule cell axon terminals (i.e. parallel fibers) was relatively normal in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} cerebellum [18.92 ± 1.3 (n = 11) vs. 23.4 ± 3.3 (n = 11) pA, *t*-test, *p*=0.477; **Fig. 4D**] (Yamasaki, Hashimoto, and Kano 2006). mEPSC frequency however, was found to be significantly

Perez et al. Resubmission

increased, a phenomenon that could be attributed to either an increase in the total number of synapses, an increase in the size of the readily releasable pool of synaptic vesicles, or an increase in the probability of neurotransmitter release in PNs of *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice [18.75 ± 2.8 Hz (n = 11) vs. 11.4 ± 1.0 Hz (n = 11), *t*-test, *p*=0.047; **Fig. 4D**]. We next explored evoked synaptic release and short-term plasticity by simultaneously recording from PNs and electrically stimulating either granule cell (i.e., parallel fibers) or inferior olivary (i.e., climbing fiber) axons with a paired-pulse burst (2-pulses, 50 ms apart). The synaptic properties of parallel fibers were found to be normal, displaying no significant differences in the expected short-term facilitation (Atluri and Regehr 1996) or halfwidth and decay time constant of the evoked EPSC [PPR: 1.3 ± 0.03 (n = 10) vs. 1.4 ± 0.05 (n = 13), *t*-test, *p*=0.162; halfwidth: 3.9 ± 0.6 vs. 4.9 ± 0.4 ms, *t*-test, *p*=0.175; time constant: 3.5 ± 0.5 vs. 4.7 ± 0.4 ms, *t*-test, *p*=0.054; **Fig. 4E**]. In comparison, we found climbing fiber-to-PN synaptic responses, which normally displays pair-pulse depression (Hansel and Linden 2000), to depress at significantly greater magnitudes in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice [PPR: 0.6 ± 0.03 (n = 6) vs. 0.7 ± 0.02 (n = 9), *t*-test, *p*=0.03; **Fig. 4F**]. The overall width and decay time constant of the evoked currents were also smaller [halfwidth: 2.3 ± 0.6 (n = 6) vs. 3.0 ± 0.2 (n = 9) ms, *t*-test, *p*=0.004; time constant (fast): 1.1 ± 0.14 vs. 2.9 ± 0.4 ms, *t*-test, *p*=0.001]. While these results could be caused by a presynaptic deficit, such as reduced vesicle stores in the climbing fiber axon terminal, the fact that the initial magnitude of the EPSC is unaffected [23.5 ± 3.6 (n = 6) vs. 19.1 ± 1.9 (n = 9), *t*-test, *p*=0.3], points to a more intrinsic deficit, such as a reduced Ca²⁺ influx from the endoplasmic reticulum, which could significantly impact long term synaptic plasticity critical to cerebellar function (Hoxha et al. 2018; Kano and Watanabe 2017). Overall, the perturbations we observed here to the passive and active PN properties likely give rise to significant cerebellar dysfunction in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice.

2.5 ATM and APTX deficiency causes a progressive perturbation of PN neural activity that is associated with dendritic shrinking and overall cerebellar atrophy.

Decreased rates of spontaneous PN action potential firing, which can be indicative of PN dysfunction, have been observed in several mouse models of ataxia, including spinocerebellar ataxia (SCA) 2, 3, 5,

Perez et al. Resubmission

6, 13, 27, several models of episodic ataxia (e.g., leaner, ducky, and tottering), and autosomal-recessive spastic ataxia of the Charlevoix-Saguenay ((Hourez et al. 2011; Hansen et al. 2013; Dell'Orco, Pulst, and Shakkottai 2017; Kasumu and Bezprozvanny 2012; Liu et al. 2009; Perkins et al. 2010; Shakkottai et al. 2011a; Jayabal et al. 2016; Stoyas et al. 2020; Hurlock, McMahon, and Joho 2008; Shakkottai et al. 2009; Bosch et al. 2015; Walter et al. 2006; Alviña and Khodakhah 2010; Ady et al. 2018; Larivière et al. 2019; Cook, Fields, and Watt 2020). We therefore used this biomarker to characterize the progression of PN perturbation in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice and assess whether deficits were restricted to ATM and APTX deficient mice, consistent with the behavioral results (**Fig. 2,3**). We additionally examined whether decreased PN activity differed across the cerebellum, as anecdotal clinical pathology reports suggest degeneration may occur asymmetrically across the cerebellum, with the anterior and posterior vermis and middle cerebellar hemispheres affected the most, although no systematic analysis has been performed and the consistency of results across patients is highly variable (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979; Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967; Aguilar et al. 1968a; Paula-Barbosa et al. 1983).

Using extracellular recording methods in the acute slice, we recorded spontaneous action potentials from 3,300 PNs (**Fig. 5A**), across 188 animals, encompassing *Atm*^{R35X/R35X}; *Aptx*^{-/-} and 3 other genotypes at 4 different time points (P45, 120, 210, and 400). We visually selected “healthy” cells (see Methods), deeper in the slice, that consistently fired during the extent of the 60 second recording period. Qualitatively, tissue and cell quality did not visually differ across genotypes under DIC microscopy. Cells were sampled in a distributed fashion across the lateral, intermediate, and medial (vermis) cerebellum of each mouse to assess whether changes in PN firing activity was ubiquitous or anatomically restricted. Regions were segregated based on gross anatomical domains in the mouse defined by natural anatomical boundaries (e.g., foliation) and their general connectivity with different regions of the nervous system (e.g., forebrain, brainstem, etc.) (Voogd and Glickstein 1998).

Perez et al. Resubmission

We found that complete deficiency of both ATM and APTX, consistent with the behavioral results, was necessary to produce a significantly reduced spontaneous PN firing frequency (**Fig. 5A, B**). Although the trend of slower PN firing rates was observed across most regions of the cerebellum, some subregions appeared to be less or minimally impacted, including several areas of the lateral cerebellum, including the paraflocculus, paramedian, and crus I and II (**Fig. 5-fig. S1**). Significant age dependent changes in firing frequency were also only observed in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice (**Fig. 5B**). The most significant decline occurring between P120 and 210 [medial: 50.3 ± 2.4 Hz (n = 61) vs. 36.9 ± 2.2 Hz (n = 31), *t*-test, *p*=0.0006]. No significant difference in PN firing frequency was detected between male and female mice within each genotype, thus the data were pooled (2-way ANOVA, *p*>0.3 across all pairwise comparisons; **Fig. 5-fig. S2**). Previous studies across several mouse models of heritable ataxia, including episodic ataxia and several variants of spinocerebellar ataxia find that physiological disruption in PN firing not only changes its frequency, but also its regularity (Kasumu and Bezprozvanny 2012; Jayabal et al. 2016; Stoyas et al. 2020; Cook, Fields, and Watt 2020). We compared both the coefficient of variation (CV) and variability in adjacent intervals (CV2) between *Atm*^{R35X/R35X}; *Aptx*^{-/-} and control mice (**Fig. 5-figs. S3, S4**). No difference in these parameters across sex, age, or genotype was detected. Consistent with the behavioral results, cerebellar dysfunction was found only in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice that developed ataxia and not in mice with at least some expression of ATM or APTX.

2.5 ATM and APTX deficiency induces cerebellar atrophy

Ataxia in A-T patients is usually detected between 1 to 2-years of age and is associated with mild to no cerebellar atrophy (Tavani et al. 2003; Taylor et al. 2015). Significant structural changes are usually apparent via neuroimaging between 5 to 10-years of age (Demaerel, Kendall, and Kingsley 1992; Tavani et al. 2003). Comprehensive characterization of the progression of cerebellar atrophy is however severely lacking in large part due challenges in imaging young patients (<2-years) and the rarity of the disorder. We therefore assessed the gross developmental progression of cerebellar size and pathohistology in mice deficient in both ATM and APTX that progressively develop a severe ataxia

Perez et al. Resubmission

and perturbations in cerebellar function. Structural changes in the overall size of the cerebellum were examined in *Atm*^{R35X/R35X}; *Aptx*^{-/-} and several different control mice over 5 time points (P45, 120, 210, 400, 460; **Fig. 5C**). Cerebellar size was defined within each animal by the ratio of 2-dimensional surface area of the dorsal cerebellum to the forebrain (i.e., cerebellum area divided by forebrain area). The cerebellar size of control mice slightly increased during adolescence and early adulthood (P45-P210), was generally stable through adulthood (P210-400), and then declined slightly in older age (P400-460). In stark contrast, relative cerebellar size in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice progressively declined after P120. No cerebellar atrophy was observed in mice with at least one copy of the *Atm* gene [i.e., *Atm*^{R35X/+}; *Aptx*^{-/-}; 1-way ANOVA, $F_{(3,44)} = 1.2$, $p=0.32$; **Fig. 5C**]. To rule out the possibility that reduced cerebellar size was related to the smaller stature of *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice, we examined, but did not find a correlation between animal weight and actual cerebellar size [Pearson's correlation, $p>0.3$ for all 4 genotypes at P460, $n = 10$ to 20]. Furthermore, we found that cerebellar size did not differ between male and the on average 22% smaller female mice across genotypes at this age [2-way ANOVA, $F_{(2,153)} = 1.9$, $p=0.2$]. Therefore, cerebellar neurodegeneration in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice, which begins after P120, is correlated with ATM and APTX deficiency.

In humans, cerebellar atrophy is associated with a variety of changes in the macro-structure of the cerebellar cortex and post mortem examination has identified abnormalities in the density and distribution of granule cells (GC) and PN (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979; Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967; Aguilar et al. 1968a; Paula-Barbosa et al. 1983; Gatti and Vinters 1985). In the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice we do not qualitatively observe severe pathological changes in the anatomy of their cerebellum (**Fig. 5-fig. S5B**). To then understand the anatomical basis of the gross atrophy, we measured the width of the cortical layer of the cerebellum that contains the massive dendritic arbors of PN called the molecular layer (ML). Consistent with the temporal changes in gross cerebellar size, PN firing frequency, and behavior, ML width in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice was reduced and declined in size from P120 to P400 [P400: $120.2 \pm 2.1 \mu\text{m}$ ($n = 5$) vs. $140.2 \pm 4.8 \mu\text{m}$ ($n = 5$), Sidak

Perez et al. Resubmission

test, $p=0.0004$; **Fig. 5D**]. In contrast, the width of the GC layer did not change with age or differ significantly between *Atm*^{R35X/R35X}, *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice [P400: $135.5 \pm 2.4 \mu\text{m}$ ($n = 6$) vs. $127.5 \pm 4.3 \mu\text{m}$ ($n = 5$), Sidak test, $p=0.5$; **Fig. 5D**]. These results, along with the finding that PN density did not significantly differ between *Atm*^{R35X/R35X}, *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice [P400: 3.9 ± 0.3 ($n = 4$) vs. 4.4 ± 0.3 ($n = 5$) PNs/4000 μm^2 , Welch's test, $p=0.3$; **Fig. 5-fig. S5B**] indicate that cerebellar atrophy is largely due to a reduction in the extent of PN dendrites. At the anatomical level, we were able to qualitatively observe a few other PN abnormalities. This includes abnormally large diameter primary dendrites and axonal swellings in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice (**Fig. 5-fig. S5C**). Overall, we find a good correlation between the abnormal structural and electrophysiological properties and the progression of motor behavioral deficits.

2.6 Differential disruption of thymocyte development in ATM-deficient vs. APTX-deficient mice

Chronic sinopulmonary infections associated with immunodeficiency are one of the leading causes of death in A-T patients (Morrell, Cromartie, and Swift 1986; Bhatt and Bush 2014). Immunodeficiency is linked to deficits in the generation of B- and T-lymphocytes that have been linked to defects in the antigen receptor gene rearrangement processes during the generation of these cells in bone marrow and thymus, respectively (Staples et al. 2008). The resulting defects in mature lymphocyte numbers include decreases in CD4⁺ helper T-cells and killer CD8⁺ T-cells (Schubert, Reichenbach, and Zielen 2002). We therefore examined the percentages of T-cells in peripheral blood and of different subpopulations in the thymus of *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice using T-cell antigen receptor (TCR) and CD4/CD8 co-receptor expression.

In the peripheral blood, we observed a significant reduction in the total fraction of CD3⁺ T-cells in mice with reduced or absent ATM expression compared to wildtype mice (**Fig. 6**). This reduction was further compounded by concomitant deficiency of APTX. ATM and APTX deficiencies reduced T-cells in peripheral blood by over 65% decrease compared to wild type controls. The effect of APTX deficiency was additive to that of ATM deficiency suggesting a different mechanism of action for each of these two proteins on T-cell generation. The reduction in the percentage of T-cells in peripheral blood was mostly

Perez et al. Resubmission

associated with reduction in the CD4⁺ helper T-cell population (**Fig. 6B**). Of interest, the proportion of CD8⁺ T-cells was increased only in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice (**Fig. 6B**). Again, we observed a differential effect of ATM and APTX deficiencies as seen for the effects of these mutations on the total T-cell fraction.

Given the reduction in T-cell populations in the blood, we next assessed T-cell development in the thymus. In this organ, bone marrow-derived T-cell progenitors undergo TCR gene rearrangement followed by positive selection for MHC restriction and negative selection of autoreactive clones. The phases of thymocyte development can be followed by monitoring expression of CD4 and CD8 expression in thymocytes. The progression of this developmental program goes from double negative (CD4⁻CD8⁻) thymocytes, to double positive (CD4⁺CD8⁺) thymocytes and then to single positive (CD4⁺ or CD8⁺) thymocytes. In addition, within the double negative stage, four different subpopulations can be identified, based on expression of CD25 and CD44, known as DN1 (CD44⁺CD25⁻), DN2 (CD44⁺CD25⁺), DN3 (CD25⁺CD44⁻) and DN4 (CD44⁻CD25⁻) (Germain 2002).

Gene rearrangement during thymocyte development occurs twice, once at the double negative thymocyte stage in the CD25⁺CD44⁻ stage (Krangel 2009) and then again in double positive thymocyte stage before progressing into separate CD4⁺ and CD8⁺ single positive populations (Livák et al. 1999). ATM deficiency has been linked to defects in both bouts of rearrangement in mice (Vachio 2007, Hathcock 2013). Therefore, we compared the proportion of cells in the thymus expressing these different developmental cell surface markers in our ATM deficient and control mice (**Fig. 7**). *Atm*^{R35X/R35X}; *Aptx*^{-/-} and *Atm*^{R35X/+}; *Aptx*^{-/-}, but not *Atm*^{R35X/R35X}; *Aptx*^{+/+} mice had significantly elevated proportions of CD44⁺CD25⁻, CD44⁺CD25⁺, and CD44⁻CD25⁺ cells compared to wildtype (**Fig. 7A**). These increased proportions appear to be due in part to an impediment of CD44⁻CD25⁺ cells maturing into CD44⁻CD25⁻ double negative cells, as the fraction of cells from *Atm*^{R35X/R35X}; *Aptx*^{-/-} and *Atm*^{R35X/+}; *Aptx*^{-/-} mice is significantly lower than wildtype (**Fig. 7A**). Of interest, APTX deficiency by itself had the greatest effect on the loss of DN4 cells suggesting that APTX deficiency, rather than ATM deficiency, is

Perez et al. Resubmission

responsible for this effect. To our knowledge, this finding implicates for the first time APTX in gene rearrangement during the process of TCR β recombination.

Next, we looked at the proportions of CD4⁺CD8⁺ thymocytes compared to CD4⁺CD8⁻ and CD4⁻CD8⁺ single positive thymocytes in these four different strains. In agreement with our results in the blood and prior studies, we found that ATM-deficient mice but not control mice displayed decreased expression of CD4⁺CD8⁻ and CD4⁻CD8⁺ single positive thymocytes (**Fig. 7B**). These results support the role of ATM in TCR α/δ gene rearrangement during thymocyte development (Bredemeyer et al. 2006), a role that is independent of the role played by APTX in early thymocyte maturation.

2.7 Readthrough molecules overcome PTC to restore ATM expression

Our primary rationale for inserting a clinically relevant nonsense mutation in the *Atm* gene was to generate a mouse amenable to critical pre-clinical testing of a novel set of small molecule readthrough (SMRT) compounds. We previously demonstrated SMRT compounds recover production of ATM protein in A-T patient derived lymphoblastoid cell lines by overcoming premature termination codons (PTC) caused by nonsense mutations (Du et al. 2013). To demonstrate suitability of this new A-T animal model for SMRT compound testing we chose to directly examine their ability to restore ATM expression using an explant approach that circumvents challenges related to *in vivo* delivery (e.g., bioavailability, route of delivery, etc.). ATM expression was measured in samples from the spleen, where ATM is normally expressed at high levels, and the cerebellum, a key target tissue for the disorder. We exposed these explant tissues, harvested from homozygous *Atm*^{R35X} and *Atm*^{Q35X} mice with either a candidate SMRT compound (GJ103), or an aminoglycoside previously known to have readthrough properties (G418) for 72 hrs. and then measured ATM expression by immunoblot to assess restoration. In both types of ATM deficient mice, ATM expression was consistently restored in the spleen and cerebellum by both G418, GJ103 (**Fig. 8**). These results demonstrate that our SMRT compounds can enable readthrough of at least 2 of the 3 possible nonsense mutation causing PTCs and provide the rationale for *in vivo* efficacy testing in follow-on studies.

3.0 Discussion

By increasing genotoxic stress through the addition of a secondary hit to the DDR pathway, we generated a novel mouse model that displays the most comprehensive set of A-T symptoms of any model to date. This includes a severe and progressive ataxia associated with cerebellar atrophy and perturbations of PN properties along with a high incidence of cancer and defects in immune cell development. Together, these comorbidities encompass the three leading causes of premature death in A-T — each contributing to roughly a third. Of these, the incapacitating effect of ataxia is the most penetrant and is reported by patients and caregivers as having the greatest impact on their quality of life. For this reason, the presence of ataxia and cerebellar atrophy in this new mouse model is of great significance as it provides for the very first time a resource to not only elucidate the mechanisms of neurological dysfunction, but also a critically needed *in vivo* model to test severely needed A-T therapeutics like the readthrough compounds we describe here.

We found several similarities between the overall progression of ataxia in the *Atm*^{R35X/R35X}, *Aptx*^{-/-} mice and A-T patients. In clinical A-T, motor deficits are observable by roughly 2-years of age, when parents and doctors detect a lowered ability to transition from toddling to a smooth, reflexively coordinated gate—unfortunately, little is known about motor defects at earlier stages due to the diseases low prevalence and current lack of early diagnostic testing (Rothblum-Oviatt et al. 2016). Patients usually learn to walk without assistance and neurological symptoms tend to remain stable through the first 4 to 5 years of life (Rothblum-Oviatt et al. 2016). We found a similar early progression of motor deficits in *Atm*^{R35X/R35X}, *Aptx*^{-/-} mice, detecting mild motor deficits early at P8 (righting reflex deficit), followed by a period of relative stability, prior to onset of a progressive and severe ataxia developing after p210 that included changes in gait, startle reflex, tremor, and locomotor activity. Several important questions arise out of these findings, including whether ATM and/or APTX have a neurodevelopmental role in the cerebellum. Future studies focused on the early phase of the disorder will be critical in understanding if the cerebellum develops normally prior to dysfunction or whether developmental defects are an initial cause. We also found, similar to A-T patients, that the severity of the late developing ataxia was

Perez et al. Resubmission

variable with some mice ambulating with a clumsy, high stepping rear gate (**Video 3**) and others moving almost entirely via contortion of the rear trunk (**Fig. 1E** and **Video 4**) (Rothblum-Oviatt et al. 2016; Levy and Lang 2018; Boder and Sedgwick 1958). Overall, we found that *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice developed a visually profound and measurable progressive loss in motor coordination that was similar to that observed in A-T patients and that was rescued by expression of at least one copy of the *Atm* or *Aptx* gene.

The loss of motor coordination in A-T has been attributed to cerebellar degeneration due to its relatively selective neuropathology across the brain and its causal role in several different forms of ataxia (Hoche et al. 2012). Consistent with A-T patient neuroimaging studies (Wallis et al. 2007; Sahama et al. 2015; Sahama et al. 2014; Dineen et al. 2020; Tavani et al. 2003; Quarantelli et al. 2013), we find that cerebellar size in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice is initially normal, but progressively atrophies in line with changes in neurological function. While loss of cerebellar tissue has been considered a main cause of ataxia in humans, it is unclear from clinical data if ataxia severity is a good predictor of the extent of cerebellar degeneration found postmortem (Aguilar et al. 1968b; Crawford et al. 2006; Dineen et al. 2020). In the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice, we find clear atrophy associated with thinning of the Purkinje neuron dendrite layer that precedes the late, severe behavioral deficits. However, since the neuropathology observed in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice is not severe, our findings suggest that like several SCAs (e.g., 1 and 3), changes in cerebellar function itself, rather than profound loss of cerebellar cells, is likely sufficient to cause the ataxic phenotype (Shakkottai et al. 2011b; Lorenzetti et al. 2000; Clark et al. 1997).

Why ATM and APTX deficiency is required to generate ataxia in mice, when loss of either is sufficient to cause ataxia in humans, remains unclear. One possibility is that the rodent brain may more flexibly utilize compensatory pathways or redundant proteins while responding to the 10-20k DNA lesions that impact cells each day (Lindahl and Barnes 2000). Several forms of DNA repair exist to potentially meet this challenge, including base excision repair (BER), nucleotide excision repair (NER), as well as homologous and non-homologous end joining (HEJ and NHEJ, respectively), all of which ATM and

Perez et al. Resubmission

APT_X have been implicated in (Chou et al. 2015; Çaglayan et al. 2017; Wakasugi et al. 2014; Tumbale et al. 2018; Chatterjee and Walker 2017). Alternatively, it may be the case that deficiency in ATM or APT_X alone does not adequately impact cell health during the mouse's comparatively short lifespan, and thus eliminating both proteins is necessary to achieve sufficient accumulation of DNA damage to manifest over this time period. This possibility is strengthened by the fact that ATM and APT_X have distinct biochemical properties and functional roles in the DNA damage response, and therefore deficiency in both would be predicted to cause a broader hit to genome stability (i.e., increased genotoxic stress).

Our finding, that two genome stability pathway proteins are required to induce neurological defects in mice strongly suggests that it is the loss of ATM's role in DNA repair, rather than potential functions in oxidative stress signaling, mitophagy, or mitochondrial function that cause the cerebellar defects (Shiloh 2020). Alternatives however, cannot be completely ruled out, as APT_X, like ATM has been observed within the mitochondria of brain cells, where it is thought to support the processing of mitochondrial DNA (Meagher and Lightowlers 2014; Sykora et al. 2011). This new mouse model provides a new tool to explore these possibilities and mechanistically define how loss of ATM and APT_X ultimately causes cerebellar dysfunction.

The biophysical perturbations observed in PNs recorded from the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice are similarly found in several other mouse models of ataxia. This includes changes we observed in PN input resistance, membrane capacitance, and AP threshold and width, which have also been described in mouse models of SCA like 1, 3, and 7 (Stoyas et al. 2020; Shakkottai et al. 2011b; Dell'Orco et al. 2015). Moreover, the progressive reduction in PN action potential firing frequency we report, which correlates well with the development of ataxia in the *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice, is reported in a large number of ataxic mouse models, including SCAs 1, 2, 3, 5, 6, and 13 as well as a few episodic forms (see review (Cook, Fields, and Watt 2020)).

Given the significant overlap in PN perturbations observed across many different ataxias caused by distinct cellular defects, restoring PN AP firing frequencies has been considered as a broad-based

Perez et al. Resubmission

therapeutic approach. However, it remains unclear whether reduced PN firing is an actual causal factor of ataxia. Moreover, experimental evidence suggests changes in PN activity may in fact be a generalized response to maintain homeostasis during ongoing disease-related impairment of PN physiology (Dell'Orco et al. 2015). Thus, continued efforts across all cerebellar ataxias are needed to link the genetic, molecular, and cellular disruptions caused by disease to the specific changes in cerebellar neural signaling that ultimately generates the ataxia. Of significant importance in this effort will be defining whether disease-causing cerebellar defects commonly or differentially cause ataxia through a loss of cerebellar function (e.g., loss of coordinating signals during movement), or from a dominant negative effect (e.g., disrupting downstream neural circuits with abnormal neural output patterns). Ultimately, while a common therapeutic strategy to address cerebellar ataxias would have the greatest impact, a directed approach that addresses the distinct genetic and molecular causes of cellular dysfunction may ultimately be necessary to successfully develop an efficacious therapeutic.

The mechanistic link between deficiency in DNA stability proteins like ATM and APTX and PN dysfunction is far less obvious than it is for other types of ataxia, like the loss of voltage gated ion channels that directly shape membrane excitability in SCA6 (i.e., $Ca_{v2.1}$) (Jayabal et al. 2016). Our results suggest the effects of ATM and APTX loss on PNs is intrinsic, as we don't find changes in the presynaptic properties of granule cells, nor evidence of their cellular loss (no change in GCL thickness). Moreover, while we observed differences in short term plasticity of inferior olivary inputs in ATM and APTX deficient PNs and wildtype, these results likely point to a disruption in Ca^{2+} homeostasis potentially via reductions in Inositol 1,4,5-triphosphate receptor 1 (*Itpr1*) expression, similar to those observed in SCAs 1, 2, and 3 mouse models as well as ATM-deficient mice (Kim et al. 2020; Chen et al. 2008; Liu et al. 2009; Shakkottai et al. 2011b). While this provides a promising avenue for future examination and comparison, it is as of yet unclear, even for the SCAs, whether changes in Ca^{2+} homeostasis is the causal factor or just another symptom or even compensatory response of diseased or disturbed PNs (Dell'Orco et al. 2015).

Perez et al. Resubmission

In the immune system, ATM is implicated in the repair of DNA breaks that naturally occur during gene rearrangement of antigen receptor genes in B- and T-cell precursors, a phenomenon critical for antigen receptor (Ig and TCR) diversity of these cells. Our finding that T-cell proportions in the blood are significantly reduced is consistent with prior studies in humans and A-T knockout mice (Schubert, Reichenbach, and Zielen 2002; Hathcock et al. 2013; Chao, Yang, and Xu 2000; Barlow et al. 1996). This reduction of T-cells in the periphery likely correlates with a defect in both cellular and humoral immunity. Importantly, we found that expression of at least one copy of the ATM gene is enough to restore CD4⁺ deficits in the blood indicating that therapies able to restore at least some ATM expression would have therapeutic efficacy. Although we have not assessed B-cell development in this paper, it is likely that similar conclusions would apply to that process given their mechanistic similarities (Marshall et al. 2018).

As expected, the reduction of T-cells in peripheral blood correlated with defective thymocyte development. In the thymus, we found two main defects. One, induced primarily by APTX deficiency, manifests as a defect in the DN3 to DN4 transition coinciding with early rearrangement of TCR β locus. The other defect, primarily caused by ATM deficiency, correlates with decreased progression of double positive CD4⁺CD8⁺ to single positive cells, primarily CD4⁺ thymocytes. While the APTX finding was surprising, as its deficiency (AOA 1) is not associated with immune deficits, APTX is known to interact with TCR β gene rearrangement proteins, including XRCC4 (Clements et al. 2004). Future studies aimed at defining APTX's role in end-joining mechanisms during TCR gene rearrangement will be important, and the possibility that alternative end-joining mechanisms, like the use of microhomologies account for the lack of an immune deficit in its absence need further investigation (Bogue et al. 1997).

The survivability of *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice is considerably longer than prior A-T mouse models. In comparison, the first A-T KO mouse model reported by Barlow et al. died from thymomas usually within 2-4 months after birth (Barlow et al. 1996). The increased severity of cancer survivability in this, and many other knockout A-T mouse models is likely genetic, as the background strain harboring the mutation has been shown to have significant effects on cancer prevalence and survivability, with A/J

Perez et al. Resubmission

and C57BL/6 backgrounds having significantly increased survivability over the BALBC and 129S strains (Genik et al. 2014). The fact that our ATM deficient mice were created on a C57BL/6 background likely underlies their comparatively long lifespan. Given that the *Atm*^{R35X/R35X}; *Aptx*^{+/+} mice do not develop ataxia, it is unlikely that the early death in A-T KO mice prevents observation of an ataxic phenotype that would otherwise develop in these mice. On the other hand, it is unknown whether the C57BL/6 background confers a resilience to developing ataxia, as it does for cancer. Defining the genetic or possibly epigenetic factors that influence the severity of the disease could provide avenues for future therapeutic development.

Given the global nature of the ATM and APTX null mutation in our mouse model, we cannot entirely rule out that extra-cerebellar defects may also contribute to the severe ataxic phenotype, and thus future examination outside the cerebellum, in both the forebrain, brainstem, spinal cord, and even muscle will need to be conducted. Within the cerebellum, while we found some anatomical differences in the PN firing properties within different regions of the cerebellum, we didn't detect regional differences in ML width or PN density. However, there are challenges in using regional anatomy as a grouping factor in the cerebellum, as the physical folds of the tissue do not necessarily correlate with the boundaries of functional, molecular expression, or biophysical property domains that have been described (Apps and Hawkes 2009; Tsutsumi et al. 2015; Gao, van Beugen, and De Zeeuw 2012; Zhou et al. 2014). Experiments focused on examining the extent of cerebellar defects within these domains will be important in future studies and compared to the anecdotal reports of anatomical differences in A-T patients (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979; Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967; Aguilar et al. 1968a; Paula-Barbosa et al. 1983).

While we detect two potential stages in the progression of ataxia in the *Atm*^{R35X/R35X}; *Aptx*^{+/+} mice, the later stage of severe ataxia develops in adulthood in mice, as compared to the childhood onset in humans. This may limit its use in some neurodevelopmental based studies. Also, the interpretation of future experiments must carefully factor in the fact that this new model expresses null mutations in two

Perez et al. Resubmission

genome stability genes at the same time, a situation that has not been detected in human patients with either A-T or AOA1.

Finally, pinpointing where, when, and how ATM deficiency causes cerebellar pathology and ataxia has been a challenge as prior ATM deficient mice generally lack the characteristic features needed to causally link cellular and molecular deficits to the ataxic phenotype. Multiple promising avenues of investigation have been defined, including those focused at the neuronal level where ATM is implicated in oxidative stress signaling (Chen et al. 2003) and synaptic function (Li et al. 2009; Vail et al. 2016), as well as glial function, where recent evidence suggests glial pathology may be a leading factor in cerebellar pathology (Kaminsky et al. 2016; Campbell et al. 2016; Petersen, Rimkus, and Wassarman 2012; Weyemi et al. 2015). This novel animal model provides a new tool to test mechanistic hypotheses regarding how ATM deficiency causes cerebellar pathology and ataxia. Additionally, this model may serve most importantly as a critical preclinical tool for testing both previously proposed therapeutic candidates (Browne et al. 2004; Chen et al. 2003) and our own SMRT compounds (Du et al. 2013). The severe limitations of not having a suitable preclinical model for therapeutic testing, especially for a rare disorder like A-T and AOA1, cannot be overstated.

4.0 Materials and Methods

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (<i>Mus musculus</i>)	<i>Atm</i> ^{R35X} , <i>Atm</i> ^{Tm1.1(103 CAG)TGA} <i>Mfgc</i>	This paper	103C>T mutation, human exon replacement	Generated by Hicks laboratory. Has been backcrossed into C57b/6 9 times. Contact pmathews@lundquist.org

Perez et al. Resubmission

Strain, strain background (<i>Mus musculus</i>)	<i>Atm</i> ^{Q35X} <i>Atm</i> ^{Tm1.1'(103C)} <i>TjMfgc</i>	This paper	103C>T mutation, targeted premature termination signal in the mouse codon	Generated by Hicks laboratory. Has been backcrossed into C57b/6 9 times. Contact pmathews@lundquist.org
Strain, strain background (<i>Mus musculus</i>)	<i>Aptx</i> ^{-/-}	Ahel et al. 2006	MGI Cat# 3687171, RRID:MGI:3687171	Contact peter.mckinnon@stjude.org
Gene (<i>Mus musculus</i>)	<i>Atm</i>	MGI	MGI:107202; C030026E19 Rik; ENSMUSG0000034218	
Gene (<i>Homo Sapien</i>)	<i>ATM</i>	OMIM	OMIM: 607585 MGI: 107202 HomoloGene: 30952; ENSG00000149311	
Sequence-based reagent	<i>Atm</i> gene	Transnetix	PCR primers	F-5'- CCTTTGAG GCATAAGT TGCAACTT G-3'
Sequence-based reagent	<i>Atm</i> gene	Transnetix	PCR primers	R- 5'- GTACAGTGT ATCAGGTTA GGCATGC-3'
Chemical compound/ drugs	GJ103 salt Formula: C ₁₆ H ₁₄ N ₄ O ₃ S	TargetMol	T3448; CAS No. : 1459687-96-7	100 µM in media
Antibody	ATM (D2E2) (Rabbit-monoclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 2873, RRID:AB_206	WB(1:500) WB(1:1000)

Perez et al. Resubmission

			2659	
Antibody	GAPDH (14C10) (Rabbit- monoclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 2118, RRID:AB_561 053	WB(1:4000)
Antibody	β -Actin (D6A8) (Rabbit- monoclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 8457, RRID:AB_109 50489	WB(1:5000)
Antibody	Anti-Rabbit IgG,HRP- linked (Goat- monoclonal- polyclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 7074, RRID:AB_209 9233	WB(1:5000)
Antibody	Calbindin (D-28k) (Rabbit, polyclonal)	Swant Inc.	Swant Cat# CB 38, RRID:AB_100 00340	IF (1:1000)
Antibody	Anti-Rabbit Alexa Fluor 488 (Goat- polyclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# A- 11034, RRID:AB_257 6217	IF(1:1000)
Antibody	CD4 (GK1.5) (Rat- monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 50- 0041-82, RRID:AB_106 09337	FACS (5 ul per test)
Antibody	CD8 (53-6.7) (Rat- monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 53- 0081-82, RRID:AB_469 897	FACS (5 ul per test)

Perez et al. Resubmission

Antibody	CD3 (145-2C11) (Hamster-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 12- 0031-83, RRID:AB_465 497	FACS (5 ul per test)
Antibody	CD44 (IM7) (Rat-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 25- 0441-82, RRID:AB_469 623	FACS (5 ul per test)
Antibody	CD25 (PC61.5) (Rat-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 47- 0251-82, RRID:AB_127 2179	FACS (5 ul per test)
Other	Fluoromount -G with DAPI	Southern Biotech	Cat# 0100-20, RRID: SCR_021261	
Commercial assay or kit	BCA Protein Assay Kit	ThermoFisher Pierce	Cat# 23225	Protein Assay
Commercial assay or kit	SuperSignal West Pico Chemilumine scent Substrate	ThermoFisher Pierce	Cat# 34580	Chemilumine scent Substrate
Commercial assay or kit	Radiance plus	Azure Biosystems	Cat# AC2103	Chemilumine scent Substrate
Software, algorithm	FlowJo	https://www.flowjo.com/solutions/flowjo	RRID:SCR_0 08520	
Software, algorithm	ImageJ software	ImageJ (http://imagej.nih.gov/ij/)	RRID:SCR_0 03070	

Perez et al. Resubmission

Software, algorithm	IgorPro	http://www.wavemetrics.com/products/igorpro/igorpro.htm	RRID:SCR_000325	Version 7; Tarotools procedures
Software, algorithm	Neuroexpress	https://www.researchgate.net/project/NeuroExpress-Analysis-software-for-whole-cell-electrophysiological-data	https://www.researchgate.net/project/NeuroExpress-Analysis-software-for-whole-cell-electrophysiological-data	Version 21.1.13; Used for mEPSC analyses
Software, algorithm	GraphPad, Prism	GraphPad Prism (https://graphpad.com)	RRID:SCR_015807	Versions 8 and 9
Software, algorithm	MBF, Stereo investigator	https://www.mbfbioscience.com/stereology	RRID:SCR_017667	Version 2021
Software, algorithm	Microsoft Excel	https://www.microsoft.com/en-us/microsoft-365/excel	RRID:SCR_016137	Version 365
Software, algorithm	Catwalk XT	https://www.noldus.com/catwalk-xt	RRID:SCR_021262	

588

589 4.1 Ethics Statement

590 This study was performed in strict accordance with the recommendations in the Guide for the Care and
591 Use of Laboratory Animals of the National Institutes of Health. All the animals were handled according
592 to approved institutional animal care and use committee (IACUC) protocols at The Lundquist Institute
593 (31374-03, 31773-02) and UCLA (ARC-2007-082, ARC-2013-068). The protocol was approved by the
594 Committee on the Ethics of Animal Experiments of the Lundquist Institute (Assurance Number: D16-
595 00213). Every effort was made to minimize pain and suffering by providing support when necessary
596 and choosing ethical endpoints.

4.2 Mice

All mice were group housed and kept under a 12-h day/night cycle with food and water available *ad libitum*. Animals were housed within the general mouse house population, and not in specialized pathogen free rooms. Older animals were made available wetted food or food gel packs on the ground of the cages as ataxia developed. *Atm*^{R35X} and *Atm*^{Q35X} mice were created and provided by Dr. Hicks and colleagues at the University of Manitoba.

These mice were created to contain the c.103C>T mutation found in a large population of North African AT patients, using recombineering Gateway technology and site-directed mutagenesis. A C>T mutation at this position in the mouse *Atm* gene creates a TAG G stop codon. The same mutation in the human ATM gene produces a TGA G stop codon. In consideration of the use of these models for therapeutic interventions, we chose to create a mouse model for each of the two PTC codons (**Fig. 1A**).

A modified Gateway R3-R4-destination vector was used to pull out the desired region of the mouse *Atm* gene from a Bacterial Artificial Chromosome (BAC) and subsequently mutated to create either a TAG G stop codon at codon 35 (M00001, position 103 (C>T)) or a TGA G stop codon (M00002, position 103 (CAG>TGA), replicating the human AT PTC). The genomic alleles were then cloned into a modified version of the NorCOMM mammalian targeting vector using a 3-way Gateway Reaction (Bradley et al. 2012). The resulting targeting vectors were electroporated into C2 ES cells (C57Bl/6N, derived in A. Nagy lab, Toronto, Canada) and successfully targeted clones were identified by selection with G418 (Gertsenstein et al. 2010). Integration of the mutated targeting cassette into the *Atm* gene locus was confirmed by Southern blot, and by sequencing of PCR products to confirm the presence of the *Atm* PTC mutation, error free targeting into the *Atm* locus and error free functional components of the vector (data not shown). Positive ES clones were used for blastocyst injection to obtain the transgenic lines. The transgenic allele contained a floxed human beta actin promoter - delta TK1- Neo cassette in the intron upstream of the region containing the mutated exon. This floxed cassette was subsequently excised by crossing with a Cre driver mouse (B6.C-Tg(CMV-cre)1Cgn/J) to generate *Atm*^{R35X/+} and *Atm*^{Q35X/+} (MGI nomenclature: *Atm*^{TM1(103CAG>TGA)MFGC} and *Atm*^{TM1(103C>T)MFGC}, respectively) mouse lines

Perez et al. Resubmission

(Fig. 1A). Genotyping of the two *Atm* lines was performed by using the following primers at Tm 62°C: *Atm* gene forward (F) primer: 5'-CCTTTGAGGCATAAGTTGCAACTTG-3'; and *Atm* gene reverse (R) primer: 5'-GTACAGTGTATCAGGTTAGGCATGC-3', creating a Wild-type allele product of 151bp or targeted allele product of 241bp (**Figs. 1A, 1B**).

Atm^{R35X} and *Atm*^{Q35X} were backcrossed with C57Bl/6J mice for 9 generations (99.2% isogenic) prior to cryopreservation and subsequent rederivation using C57Bl/6J surrogate mothers. *Atm*^{R35X} and *Atm*^{Q35X} breeders were obtained from F1 sibling *Atm*^{R35X/+} and *Atm*^{Q35X/+} mice. *Atm*^{R35X/R35X} and *Atm*^{Q35X/Q35X} were both found to be fertile. *Aptx* knockout (*Aptx*^{-/-}) mice were created and provided to Dr. Mathews as embryos from Dr. McKinnon (Ahel et al. 2006), and subsequently rederived via C57Bl/6J surrogate mothers. *Aptx*^{-/-} mice are on a C57Bl/6 and 129 mixed background. *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice of various wildtype, heterozygous, and homozygous combinations were created from *Atm*^{R35X/+}; *Aptx*^{+/-} breeders generated by crossing *Atm*^{R35X/R35X} and *Aptx*^{-/-} mice. One cohort of double mutant and corresponding control mice were used in the longitudinal behavioral study for gait analyses and SHIRPA testing (**Figs. 2, 3**). Multiple additional cohorts of age matched double mutant and control mice were used for electrophysiological, immunohistological, and Vertical Pole test experiments (**Figs. 4, 7**). Immunological and protein expression experiments were carried out using mice bred from the original *Atm*^{R35X} and *Atm*^{Q35X} rederived mice (**Figs. 5, 6, and 8**).

Genotyping was performed from ear tissue samples of P8-11 mice. Real-time PCR methods conducted by Transnetyx Inc. were used to determine each animals' genotype. Animals were made identifiable via toe tattoos given at the same time as ear biopsy. Unique primers for *Atm*^{R35X} and *Atm*^{Q35X} were quantified and used to identify wildtype, hetero- and homo-zygous mice (listed above). *Aptx*^{-/-} and *Aptx*^{+/+} primers were used to assess their genotypes.

4.3 Animal Health

Animals were weighed via a digital scale at P8, 45, 120, 210, 400. Animal death was recorded as the day found dead, or on the day of euthanization when the animals reached a humane endpoint (animal unable to right itself within 60s, significant hair matting indicating lack of self-grooming, or excessive

Perez et al. Resubmission

distress as noted by the veterinary staff). Animal carcasses were immediately frozen upon death, and postmortem necropsies were carried out in batch. Probable cause of death was determined to the best of our ability in collaboration with the staff veterinarian (Dr. Catalina Guerra) by visual inspection of the internal organs. Some mice were cannibalized or accidentally disposed of by vivarium staff and were therefore labelled as “missing.” Mice with no discernable visual cause of death were labelled “indeterminable.” Mice that were found with thoracic masses near where the thymus would normally be in young mice were listed as “thymic cancer.” All other identified probable causes of death (e.g., enlarged livers, urinary blockage) were labelled “other.”

4.4 Behavior

Before performing any behavioral test, mice were acclimated to the behavioral suite for ~20-minutes. Mice were tested at varying times of the day, in line with their day cycle. A battery of behavioral tests was performed on naïve double mutant mice of the indicated genotypes at various time points depending on the behavior but in the same cohort of mice. The battery of tests included Catwalk Gait assessment (P45, 120, 210, 400) and a subset of the SmithKline-Beecham Harwell Imperial-College and Royal-London-Hospital Phenotype Assessment (SHIRPA) tests (P30 and 400). These tests were conducted by the UCLA Behavioral Core. Double mutant and control mice were additionally examined on the Vertical Pole test. All behavioral apparatus was wiped down with ethanol (70%) between each testing each subject.

Gait Analysis

We used a Noldus Catwalk Gait analysis system designed to semi-automatically measure and analyze the gait of mice during normal ambulation. Briefly, the movement of mice across a glass bottom corridor is video recorded from a ventral position. Paw prints are highlighted in the video due to light illumination across the glass walking platform. Each mouse step within a video is subsequently detected using Catwalk XT (Noldus) in a semi-automated fashion. A run for each mouse consists of 3 trials of consistent ambulation across the monitored platform. Only consistent trials are accepted, and mice may take up to 10 attempts to complete 3 compliant trials in either direction across the corridor. Compliant

Perez et al. Resubmission

trials were defined as those with movement across the platform under 5 s-long and with no more than 60% speed variation. Once placed onto the platform, mice generally ran back and forth without any need for experimenter prompting.

Vertical Pole

Mice are placed at the top of an 80 cm tall bolt with their nose faced down and hind paws as close to the top as possible. Mice are immediately released, and time started immediately upon placement. Time is stopped when the first forepaw touches the surface below the pole. A mouse's natural predilection is to immediately climb down the pole, and they are given up to 60 s to traverse the pole, otherwise they are helped off the pole. A non-completed trial is automatically given a time of 30 s, as 95% of mice that did not descend within 30 s were still on the pole at the 60 s mark.

SHIRPA

Behavioral tests were conducted by the University of California, Los Angeles Behavioral Core at P30 and P400. All parameters are scored to provide a quantitative assessment, which enables comparison of results both over time and between different laboratories. Each mouse was sequentially tested across all behaviors within ~20-min time span before moving onto the next mouse. The experimenter was blinded to animal genotype. The screen was performed as described previously (Rogers et al. 1997).

Behavioral Observation

The primary screen provides a behavioral observation profile and assessment of each animal begins by observing undisturbed behavior in a viewing jar (10 cm diameter) for 5-min. In addition to the scored behaviors of **body position**, **spontaneous activity**, **respiration rate**, and **tremor**, the observer logs any instances of bizarre or stereotyped behavior and convulsions, compulsive licking, self-destructive biting, retropulsion (walking backwards) and indications of spatial disorientation.

Arena Behavior

Perez et al. Resubmission

Thereafter, the mouse is transferred to the arena (30 cm x 50 cm) for testing of transfer arousal and observation of normal behavior. The arena is marked into a grid of 10 cm² squares to measure locomotor activity within a 30 s-period. While the mouse is active in the arena, measures of **startle response, gait, pelvic elevation, and tail elevation** are recorded.

Supine Restraint

The animal is restrained in a supine position to record autonomic behaviors. During this assessment, **grip strength, body tone, pinna reflex, corneal reflex, toe pinch, wire maneuver, and heart rate**, were evaluated.

Balance and Orientation

Finally, several measures of vestibular system function were performed. The **righting reflex, contact righting reflex**, and **negative geotaxis** tests were performed. Throughout this procedure vocalization, urination and general fear, irritability, or aggression were recorded.

Equipment Used

1. Clear Plexiglas arena (approximate internal dimensions 55 x 33 x18 cm). On the floor of the arena is a Plexiglas sheet marked with 15 squares (11 cm). A rigid horizontal wire (3 mm diameter) is secured across the rear right corner such that the animals cannot touch the sides during the wire maneuver. A grid (40 x 20 cm) with 12 mm mesh (approximate) is secured across the width of the box for measuring tail suspension and grip strength behavior.
2. A clear Plexiglas cylinder (15 x 11 cm) was used as a viewing jar.
3. One grid floor (40 x 20 cm) with 12 mm meshes on which viewing jars stand.
4. Four cylindrical stainless-steel supports (3 cm high x 2.5 cm diameter) to raise grids off the bench.
5. One square (13 cm) stainless steel plate for transfer of animals to the arena.
6. Cut lengths of 3 / 0 Mersilk held in the forceps for corneal and pinna reflex tests
7. A plastic dowel rod sharpened to a pencil point to test salivation and biting.
8. A pair of dissecting equipment forceps, curved with fine points (125 mm forceps, Philip Harris Scientific, Cat. No. D46-174), for the toe pinch.

Perez et al. Resubmission

9. A stopwatch.

10. An IHR Click box is used for testing the startle responses. The Click Box generates a brief 20 KHz tone at 90dB SPL when held 30cm above the mouse. Contact Prof. K.P. Steel, MRC Institute of Hearing Research, University Park, Nottingham NG7 2RD.

11. A ruler.

12. A 30 cm clear Plexiglas tube with an internal diameter of 2.5 cm for the contact righting reflex.

4.5 Electrophysiology

Preparation of acute cerebellar slices

Acute parasagittal slices of 300 μ m thickness were prepared from the cerebellum of experimental and control littermate mice by following published methods (Hansen et al., 2013). In brief, cerebella were quickly removed and immersed in an ice-cold extracellular solution with composition of (mM): 119 NaCl, 26 NaHCO₃, 11 glucose, 2.5 KCl, 2.5 CaCl₂, 1.3 MgCl₂ and 1 NaH₂PO₄, pH 7.4 when gassed with 5% CO₂/95% O₂. Cerebella were sectioned parasagittally using a vibratome (Leica VT-1000, Leica Biosystems, Nussloch, Germany) and initially incubated at 35°C for ~30 min, and then equilibrated and stored at room temperature until use.

Extracellular Electrophysiology

Extracellular and intracellular recordings were obtained from Purkinje neurons (PNs) in slices constantly perfused with carbogen-bubbled extracellular solution and maintained at either 37° C (extracellular) or 32° C (intracellular) $\pm$ 1° C (see above). Cells were visualized with DIC optics and a water-immersion 40x objective (NA 0.75) using a Zeiss Examiner microscope. Glass pipettes of ~3 M Ω resistance (Model P-1000, Sutter instruments, Novato, CA) were filled with either extracellular solution and positioned near PN axon hillocks in order to measure action potential-associated capacitive current transients in voltage clamp mode with the pipette potential held at 0 mV. For whole-cell patch clamp recordings, pipettes were filled with an intracellular solution (mM): 140 KMeth (CH₃KO₃S), 10 NaCl, 2 MgCl₂, 0.2 CaCl₂, 10 HEPES, 14 Phosphocreatine (tris salt), 1 EGTA, 4 Mg-ATP, 0.4 Na-GTP. 100 μ M Picrotoxin (Sigma) was added to block inhibitory GABAergic synaptic inputs. Data was acquired using

Perez et al. Resubmission

a MultiClamp 700B amplifier at 20 or 100 kHz in voltage or current clamp mode, Digidata 1440 with pClamp10 (Molecular Devices, Sunnyvale, CA) and filtered at 2 to 4 kHz. The series resistance was usually between 10 and 15 MΩ. Series resistance was compensated at 80% for short term plasticity experiments only.

For extracellular recordings, a total of 20 to 45 PNs were recorded from for each animal across all genotypes, sexes, and age groups. Recordings were distributed across both the medial-lateral and rostro-caudal axis of the cerebellum. Only cells with a “healthy” look (low contrast of cellular borders), and regular, uninterrupted firing rate were examined. During analysis, a few cells were found to have gaps in firing of greater than 2 second and these cells were eliminated from analysis, as this type of firing is associated with being “unhealthy.” Double mutant tissue did not qualitatively differ in appearance under DIC microscopy prior to recordings, nor was the number of “unhealthy” cells greater than that of other genotypes (7% vs 4 to 11% of all cells across control genotypes at P400). Spatial comparison of neural activity was obtained by recording from serial sections in the flocculus, lateral (2nd or 3rd), intermediate (6th or 7th), and medial (11th or 12th) slices. Lower number slices were used in the younger age groups (P45 and 110) to roughly match the relative positioning of recordings across age groups. 0-3 recordings were made from each lobule within each slice dependent on tissue quality and health. Each recording lasted for 1-minute. 3 to 5 mice were used for each age group and the experimenter was blinded to the genotype, age, and sex.

Intracellular recordings were obtained from PNs in either folia III or VIII of the vermis, no statistical differences in properties were observed between folia.

Analyses

Spontaneous action potential interstimulus intervals detected and analyzed using standard and custom routines in ClampFit (Molecular Device), IgorPro (Wavemetrics), and Excel (Microsoft). Specifically, action potentials were threshold detected and spiking statistics (i.e., frequency and interval length), determined using adapted IgorPro routines (Taro Tools; <https://sites.google.com/site/tarotoolsregister/>).

Perez et al. Resubmission

The coefficient of variation of the mean inter-spike interval (CV) and the median inter-spike interval ($CV2 = \frac{2}{\sqrt{2}} \frac{|ISIn+1-ISIn|}{(ISIn+1+ISIn)}$) were calculated in Excel using custom macros.

Standard membrane properties were analyzed using IgorPro. R_M was determined by averaging 3 voltage trace responses to a -5 mV step pulse from a -80 mV holding potential and measuring the resulting current deflection between 900 and 1000 ms after onset. The membrane time constant was measured by fitting a single exponential to the initial decay phase from 90 to 10% of the peak. C_M was calculated by dividing the membrane time constant by the R_M . mEPSC events were recorded over a 1-minute epoch and detected and measured using Neuroexpress (v21.1.13). Parallel and climbing fiber axons were stimulated using theta-glass electrodes (W.P.I.) and a TTL controlled stimulus isolator (ISO-Flex, A.M.P.I.). Evoked EPSC amplitudes and decay time constants (1 exp. for parallel and 2 exp. for climbing fibers) were analyzed using custom routines in IgorPro. Action potentials were examined as part of a set of 1 s current injections between -500 and 2250 pA (250 pA steps) with a holding current adjusted to maintain an ~70 mV potential. Action potential waveforms were measured using custom routines in IgorPro. Action potential threshold was defined as the first membrane voltage in which the first derivative exceeded 30 mV/ms (Zhu et al. 2006).

4.6 Examination of Cerebellar Atrophy

Cerebellar size

Immediately after brain removal from the skull, a dorsal, whole mount image was obtained. Images were then processed using Fiji (NIH). The forebrain and cerebellar sizes were assessed by outlining their 2-dimensional space and then calculating area. We normalized for possible differences in overall brain size by dividing the results of the cerebellum by forebrain size to produce a relative cerebellum to forebrain ratio. Experimenters were blind to the genotype of the animal.

Immunohistochemistry

At the respective study endpoints (P45, 120, 210, 400), male and female mice of all genotypes represented in this study were anesthetized with isoflurane and underwent transcardial perfusion with

Perez et al. Resubmission

phosphate-buffered saline followed by 4% (w/v) buffered paraformaldehyde (PFA) and then dissected to extract the brain. Images of the whole brain were taken immediately after removing the brain from the skull and the brains were then submerged in 4% PFA for 24-hours, followed by 72-hours in 30% sucrose in Tris-buffered saline (TBS) with 0.05% azide, and then cryoprotected in TBS-AF and stored at 4°C until further use. The cerebellum was separated from the forebrain and parasagittally sectioned using a sliding microtome (Microm HM 430, Thermo Scientific) set to section at 40 µm thickness. Cerebellum sections were collected in a series of six and stored in TBS-AF at 4° C until further use. For immunofluorescent visualization of Purkinje neurons, cerebellum sections of both *Atm*^{+/+}; *Aptx*^{+/+} and *Atm*^{R35X/R35X}; *Aptx*^{-/-} (n = 5 per genotype) were washed for 5 min in TBS three times, and then blocked in 15% normal goat serum at room temperature for 30 min followed by free floating incubation in rabbit anti-calbindin D-28k (1:1000) for 1 hour at room temperature on an orbital shaker, then washed for 5 min with TBS three times, followed by free floating incubation in goat anti-rabbit Alexa Fluor 488 (1:1000) for 1 h in the dark at room temperature on an orbital shaker. Following secondary antibody incubation, sections were washed for 5 min in TBS three times and stored in TBS until further use. Sections were mounted and cover-slipped with Fluoromount-G with DAPI. Slides were scanned using Stereo Investigator version 2020.1.3 64bit (MBF Bioscience) on a Zeiss Axio Imager.M2 microscope (Carl Zeiss Microscopy) using a 20x objective (NA 0.5) and images captured with a Hamamatsu ORCA Flash 4.0 LT C11440 digital camera (Hamamatsu Photonics). To quantify the number of calbindin-reactive cells in each folia in the resulting images, we randomly drew 2 lines between 300 to 500 µm long in each folia and manually counted the total number of PNs along the length within the 40 µm thickness of the tissue slice. 2D density (# of PNs/(linear length * 40 um thickness)) of the two samples per folia were then averaged for further between folia and subject comparison. Experimenter was blinded to mouse genotype.

Molecular (Calbindin) and granule cell (DAPI) layer widths were assessed by taking two width measurements at two predefined locations for each folia, roughly halfway along the long extent of each

Perez et al. Resubmission

folia. Experimenter was blinded to the mouse genotype in which slices were examined and the order of examination was interleaved.

4.7 Flow Cytometry Measurements

Flow cytometry analysis of blood and thymus cells was performed by staining with specific anti-mouse antibodies: CD4, CD8, CD3, CD44, and CD25. Briefly, whole-blood samples (50 μ l) were stained using fluorescent-labeled antibodies, then red-blood cells were lysed using BD lysing solution while live white-blood cells were stained using a viability stain. Thymi were mechanically dissociated. 1 to 2 million thymus cells were similarly stained using specific antibodies for CD4, CD8, CD44 and CD25. Analysis of immuno-stained white blood cells or thymus samples was performed using FACS ARIA III and data analyzed using FlowJo software as reported previously (Sanghez et al. 2017).

4.8 Western Blots

Protein extracts (cells/tissues) were homogenized in radioimmunoprecipitation assay (RIPA) lysis buffer (150 mM NaCl, 1% Nonidet P-40 [NP-40], 0.5% deoxycholate, 0.1% SDS, 50 mM Tris, pH 8.0) with protease inhibitors (10 μ g/ml AEBSF, 10 μ g/ml leupeptin, 5 μ g/ml pepstatin, 5 μ g/ml chymotrypsin, 10 μ g/ml aprotinin). The protein extracts were sonicated then pelleted by centrifugation at 13,000 rpm for 15 min at 4°C. BCA protein assay was used to quantify protein concentrations. Samples containing equal amounts of protein 50 to 100 μ g per lane were separated using 4 to 12% gradient TGX precast gels BioRad then transferred by TransBlot Semi-Dry BioRad system using Nitrocellulose transfer pack. Transferred blots were stained by Ponceau S stain for equal protein loading then washed and blocked with 5% nonfat dry milk in TBST for 60 min at room temp. Primary antibodies were incubated with shaking overnight at 4°C. Blots were probed for the following antibodies: ATM (D2E2) Rabbit mAb Cell Signaling, at 1:1000 dilution, β -Actin (D6A8) Rabbit mAb Cell Signaling, GAPDH (D16H11) Rabbit mAb Cell Signaling followed by the appropriate horseradish peroxidase–conjugated (HRP) secondary Anti-rabbit, Anti-mouse for 2 hours at room temperature. After multiple washes with TBST, Protein expression was detected by Radiance Plus chemiluminescence substrate using the Azure c400 and the

Perez et al. Resubmission

BioRad ChemiDoc imaging systems. Densitometric analysis of the ATM was performed using ImageJ. Experiments were performed with 2 technical and 2-3 biological replicates as indicated.

4.9 Statistical Assessment

The number of animals chosen for each group was based on a priori power analyses using GPower v3.1 based on an α size of 0.5, power of 0.8, and effect sizes estimated from preliminary data or prior studies. We used both parametric (1- and 2-way ANOVA) for normally distributed and non-parametric (Kruskal Wallace) statistical methods for interval data to test for differences between groups followed by pairwise multiple comparisons tests as indicated in the text. Outliers for immune data in Figs. 6 and 7 were excluded via the ROUT method (Q=2%). The specific analyses used for each data set is noted in each figure legend. For all figures: * $p \leq 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Data are reported as mean $\pm$ SEM and box and whisker plots indicate the minimum, first quartile, median, third quartile, and maximum data values. All figures and statistical analyses were completed using Excel (Microsoft 360) or Prism v8 and 9 (Graphpad).

5.0 Acknowledgements

We would like to thank the UCLA Behavioral Core, especially Irina Zhuravka for her efforts assaying behavioral deficits in the mice. We would also like to thank Dr. Jennifer Fogel for comments and edits to the manuscript.

6.0 Competing interests

The authors declare that no competing interests exist.

7.0 Citations

Ady, Visou, Brenda Toscano-Márquez, Moushumi Nath, Philip K Chang, Jeanette Hui, Anna Cook, François Charron, Roxanne Larivière, Bernard Brais, and R Anne McKinney. 2018. 'Altered synaptic and firing properties of cerebellar Purkinje cells in a mouse model of ARSACS', *The Journal of physiology*, 596: 4253-67.

Perez et al. Resubmission

- 875 Aguilar, M. J., S. Kamoshita, B. H. Landing, E. Boder, and R. P. Sedgwick. 1968a. 'Pathological
876 observations in ataxia-telangiectasia. A report of five cases', *Journal of Neuropathology and*
877 *Experimental Neurology*, 27: 659-76.
- 878 Aguilar, Mary Jane, Shigehiko Kamoshita, Benjamin H. Landing, Elena Boder, and Robert P. Sedgwick.
879 1968b. 'Pathological Observations in Ataxia-Telangiectasia: A Report on Five Cases*', *Journal*
880 *of Neuropathology and Experimental Neurology*, 27: 659-76.
- 881 Ahel, I., U. Rass, S. F. El-Khamisy, S. Katyal, P. M. Clements, P. J. McKinnon, K. W. Caldecott, and S.
882 C. West. 2006. 'The neurodegenerative disease protein aprataxin resolves abortive DNA
883 ligation intermediates', *Nature*, 443: 713-6.
- 884 Aicardi, J., C. Barbosa, E. Andermann, F. Andermann, R. Morcos, Q. Ghanem, Y. Fukuyama, Y.
885 Awaya, and P. Moe. 1988. 'Ataxia-ocular motor apraxia: a syndrome mimicking ataxia-
886 telangiectasia', *Annals of Neurology*, 24: 497-502.
- 887 Alviña, Karina, and Kamran Khodakhah. 2010. 'KCa channels as therapeutic targets in episodic ataxia
888 type-2', *Journal of Neuroscience*, 30: 7249-57.
- 889 Amromin, George D, Elena Boder, and Raymond Teplitz. 1979. 'Ataxia-telangiectasia with a 32 year
890 survival: a clinicopathological report', *Journal of Neuropathology and Experimental Neurology*,
891 38: 621-43.
- 892 Ando, K., J. L. Kernan, P. H. Liu, T. Sanda, E. Logette, J. Tschopp, A. T. Look, J. Wang, L. Bouchier-
893 Hayes, and S. Sidi. 2012. 'PIDD death-domain phosphorylation by ATM controls prodeath
894 versus prosurvival PIDDosome signaling', *Molecular Cell*, 47: 681-93.
- 895 Apps, R., and R. Hawkes. 2009. 'Cerebellar cortical organization: a one-map hypothesis', *Nature*
896 *Reviews: Neuroscience*, 10: 670-81.
- 897 Atluri, Pradeep P., and Wade G. Regehr. 1996. 'Determinants of the Time Course of Facilitation at the
898 Granule Cell to Purkinje Cell Synapse', *The Journal of Neuroscience*, 16: 5661-71.
- 899 Barlow, C., S. Hirotsune, R. Paylor, M. Liyanage, M. Eckhaus, F. Collins, Y. Shiloh, J. N. Crawley, T.
900 Ried, D. Tagle, and A. Wynshaw-Boris. 1996. 'Atm-deficient mice: a paradigm of ataxia
901 telangiectasia', *Cell*, 86: 159-71.
- 902 Beraldi, R., D. K. Meyerholz, A. Savinov, A. D. Kovacs, J. M. Weimer, J. A. Dykstra, R. D. Geraets, and
903 D. A. Pearce. 2017. 'Genetic ataxia telangiectasia porcine model phenocopies the multisystemic
904 features of the human disease', *Biochim Biophys Acta Mol Basis Dis*, 1863: 2862-70.
- 905 Bhatt, Jayesh M., and Andrew Bush. 2014. 'Microbiological surveillance in lung disease in ataxia
906 telangiectasia', *European Respiratory Journal*, 43: 1797-801.
- 907 Boder, Elena, and Robert P. Sedgwick. 1958. 'ATAXIA-TELANGIECTASIA', *A Familial Syndrome of*
908 *Progressive Cerebellar Ataxia, Oculocutaneous Telangiectasia and Frequent Pulmonary*
909 *Infection*, 21: 526-54.
- 910 Bogue, M. A., C. Wang, C. Zhu, and D. B. Roth. 1997. 'V(D)J recombination in Ku86-deficient mice:
911 distinct effects on coding, signal, and hybrid joint formation', *Immunity*, 7: 37-47.
- 912 Bosch, Marie K, Yarimar Carrasquillo, Joseph L Ransdell, Ajay Kanakamedala, David M Ornitz, and
913 Jeanne M Nerbonne. 2015. 'Intracellular FGF14 (iFGF14) is required for spontaneous and
914 evoked firing in cerebellar Purkinje neurons and for motor coordination and balance', *Journal of*
915 *Neuroscience*, 35: 6752-69.
- 916 Bradley, A., K. Anastassiadis, A. Ayadi, J. F. Battey, C. Bell, M. C. Birling, J. Bottomley, S. D. Brown, A.
917 Bürger, C. J. Bult, W. Bushell, F. S. Collins, C. Desaintes, B. Doe, A. Economides, J. T. Eppig,
918 R. H. Finnell, C. Fletcher, M. Fray, D. Friendewey, R. H. Friedel, F. G. Grosveld, J. Hansen, Y.
919 Hérault, G. Hicks, A. Hörlein, R. Houghton, M. Hrabé de Angelis, D. Huylebroeck, V. Iyer, P. J.
920 de Jong, J. A. Kadin, C. Kaloff, K. Kennedy, M. Koutsourakis, K. C. Lloyd, S. Marschall, J.
921 Mason, C. McKerlie, M. P. McLeod, H. von Melchner, M. Moore, A. O. Mujica, A. Nagy, M.
922 Nefedov, L. M. Nutter, G. Pavlovic, J. L. Peterson, J. Pollock, R. Ramirez-Solis, D. E. Rancourt,
923 M. Raspa, J. E. Remacle, M. Ringwald, B. Rosen, N. Rosenthal, J. Rossant, P. Ruiz Noppinger,
924 E. Ryder, J. Z. Schick, F. Schnütgen, P. Schofield, C. Seisenberger, M. Selloum, E. M.
925 Simpson, W. C. Skarnes, D. Smedley, W. L. Stanford, A. F. Stewart, K. Stone, K. Swan, H.
926 Tadepally, L. Teboul, G. P. Tocchini-Valentini, D. Valenzuela, A. P. West, K. Yamamura, Y.

Perez et al. Resubmission

- 927 Yoshinaga, and W. Wurst. 2012. 'The mammalian gene function resource: the International
928 Knockout Mouse Consortium', *Mammalian Genome*, 23: 580-6.
- 929 Bredemeyer, Andrea L, Girdhar G Sharma, Ching-Yu Huang, Beth A Helmink, Laura M Walker, Katrina
930 C Khor, Beth Nuskey, Kathleen E Sullivan, Tej K Pandita, and Craig H Bassing. 2006. 'ATM
931 stabilizes DNA double-strand-break complexes during V (D) J recombination', *Nature*, 442: 466-
932 70.
- 933 Browne, Susan E., L. Jackson Roberts, Phyllis A. Dennerly, Susan R. Doctrow, M. Flint Beal, Carrolee
934 Barlow, and Rodney L. Levine. 2004. 'Treatment with a catalytic antioxidant corrects the
935 neurobehavioral defect in ataxia-telangiectasia mice', *Free Radical Biology and Medicine*, 36:
936 938-42.
- 937 Çaglayan, Melike, Rajendra Prasad, Rachel Krasich, Matthew J. Longley, Kei Kadoda, Masataka
938 Tsuda, Hiroyuki Sasanuma, Shunichi Takeda, Keizo Tano, William C. Copeland, and Samuel H.
939 Wilson. 2017. 'Complementation of aprataxin deficiency by base excision repair enzymes in
940 mitochondrial extracts', *Nucleic Acids Research*, 45: 10079-88.
- 941 Campbell, A., B. Krupp, J. Bushman, M. Noble, C. Pröschel, and M. Mayer-Pröschel. 2015. 'A novel
942 mouse model for ataxia-telangiectasia with a N-terminal mutation displays a behavioral defect
943 and a low incidence of lymphoma but no increased oxidative burden', *Human Molecular
944 Genetics*, 24: 6331-49.
- 945 Campbell, Andrew, Jared Bushman, Joshua Munger, Mark Noble, Christoph Pröschel, and Margot
946 Mayer-Pröschel. 2016. 'Mutation of ataxia-telangiectasia mutated is associated with
947 dysfunctional glutathione homeostasis in cerebellar astroglia', *Glia*, 64: 227-39.
- 948 Chao, Connie, Eva Marie Yang, and Yang Xu. 2000. 'Rescue of Defective T Cell Development and
949 Function in *Atm*^{-/-} Mice by a Functional TCRαβ Transgene', *The Journal of Immunology*, 164:
950 345-49.
- 951 Chatterjee, Nimrat, and Graham C. Walker. 2017. 'Mechanisms of DNA damage, repair, and
952 mutagenesis', *Environmental and Molecular Mutagenesis*, 58: 235-63.
- 953 Chen, Philip, Cheng Peng, John Luff, Kevin Spring, Dianne Watters, Steven Bottle, Shigeki Furuya, and
954 Martin F. Lavin. 2003. 'Oxidative Stress Is Responsible for Deficient Survival and
955 Dendritogenesis in Purkinje Neurons from Ataxia-Telangiectasia Mutated Mutant Mice', *The
956 Journal of Neuroscience*, 23: 11453-60.
- 957 Chen, Xi, Tie-Shan Tang, Huiping Tu, Omar Nelson, Mark Pook, Robert Hammer, Nobuyuki Nukina,
958 and Ilya Bezprozvanny. 2008. 'Deranged calcium signaling and neurodegeneration in
959 spinocerebellar ataxia type 3', *Journal of Neuroscience*, 28: 12713-24.
- 960 Chou, W. C., L. Y. Hu, C. N. Hsiung, and C. Y. Shen. 2015. 'Initiation of the ATM-Chk2 DNA damage
961 response through the base excision repair pathway', *Carcinogenesis*, 36: 832-40.
- 962 Clark, H Brent, Eric N Burright, Wael S Yunis, Seth Larson, Claire Wilcox, Boyd Hartman, Antoni
963 Matilla, Huda Y Zoghbi, and Harry T Orr. 1997. 'Purkinje Cell Expression of a Mutant Allele of
964 SCA1in Transgenic Mice Leads to Disparate Effects on Motor Behaviors, Followed by a
965 Progressive Cerebellar Dysfunction and Histological Alterations', *Journal of Neuroscience*, 17:
966 7385-95.
- 967 Clements, P. M., C. Breslin, E. D. Deeks, P. J. Byrd, L. Ju, P. Bieganski, C. Brenner, M. C. Moreira,
968 A. M. Taylor, and K. W. Caldecott. 2004. 'The ataxia-oculomotor apraxia 1 gene product has a
969 role distinct from ATM and interacts with the DNA strand break repair proteins XRCC1 and
970 XRCC4', *DNA Repair (Amst)*, 3: 1493-502.
- 971 Concannon, P., and R. A. Gatti. 1997. 'Diversity of ATM gene mutations detected in patients with
972 ataxia-telangiectasia', *Human Mutation*, 10: 100-7.
- 973 Cook, Anna A., Eviatar Fields, and Alanna J. Watt. 2020. 'Losing the Beat: Contribution of Purkinje Cell
974 Firing Dysfunction to Disease, and Its Reversal', *Neuroscience*.
- 975 Coutinho P, Barbot C, Coutinho P. 2002. 'Ataxia with Oculomotor Apraxia Type 1.' in Ardinger HH
976 Adam MP, Pagon RA, et al. (ed.), *GeneReviews*® (GeneReviews® [Internet]: University of
977 Washington, Seattle).
- 978 Crawford, T. O., R. L. Skolasky, R. Fernandez, K. J. Rosquist, and H. M. Lederman. 2006. 'Survival
979 probability in ataxia telangiectasia', *Archives of Disease in Childhood*, 91: 610-11.

Perez et al. Resubmission

- De Leon, G. A., W. D. Grover, and D. S. Huff. 1976. 'Neuropathologic changes in ataxia-telangiectasia', *Neurology*, 26: 947-51.
- Dell'Orco, James M., Aaron H. Wasserman, Ravi Chopra, Melissa A. C. Ingram, Yuan-Shih Hu, Vikrant Singh, Heike Wulff, Puneet Opal, Harry T. Orr, and Vikram G. Shakkottai. 2015. 'Neuronal Atrophy Early in Degenerative Ataxia Is a Compensatory Mechanism to Regulate Membrane Excitability', *The Journal of Neuroscience*, 35: 11292-307.
- Dell'Orco, James M, Stefan M Pulst, and Vikram G Shakkottai. 2017. 'Potassium channel dysfunction underlies Purkinje neuron spiking abnormalities in spinocerebellar ataxia type 2', *Human Molecular Genetics*, 26: 3935-45.
- Demaerel, PH, BE Kendall, and D Kingsley. 1992. 'Cranial CT and MRI in diseases with DNA repair defects', *Neuroradiology*, 34: 117-21.
- Dineen, Rob A., Felix Raschke, Hannah L. McGlashan, Stefan Pszczolkowski, Lorna Hack, Andrew D. Cooper, Manish Prasad, Gabriel Chow, William P. Whitehouse, and Dorothee P. Auer. 2020. 'Multiparametric cerebellar imaging and clinical phenotype in childhood ataxia telangiectasia', *NeuroImage: Clinical*, 25: 102110.
- Du, L., M. E. Jung, R. Damoiseaux, G. Completo, F. Fike, J. M. Ku, S. Nahas, C. Piao, H. Hu, and R. A. Gatti. 2013. 'A new series of small molecular weight compounds induce read through of all three types of nonsense mutations in the ATM gene', *Molecular Therapy*, 21: 1653-60.
- El-Brolosy, Mohamed A., and Didier Y. R. Stainier. 2017. 'Genetic compensation: A phenomenon in search of mechanisms', *Plos Genetics*, 13: e1006780-e80.
- Elson, A., Y. Wang, C. J. Daugherty, C. C. Morton, F. Zhou, J. Campos-Torres, and P. Leder. 1996. 'Pleiotropic defects in ataxia-telangiectasia protein-deficient mice', *Proceedings of the National Academy of Sciences of the United States of America*, 93: 13084-9.
- Gao, Zhenyu, Boeke J van Beugen, and Chris I De Zeeuw. 2012. 'Distributed synergistic plasticity and cerebellar learning', *Nature Reviews: Neuroscience*, 13: 619-35.
- Gatti, R. A., and H. V. Vinters. 1985. 'Cerebellar pathology in ataxia-telangiectasia: the significance of basket cells', *Kroc Foundation Series*, 19: 225-32.
- Genik, P. C., H. Bielefeldt-Ohmann, X. Liu, M. D. Story, L. Ding, J. M. Bush, C. M. Fallgren, and M. M. Weil. 2014. 'Strain background determines lymphoma incidence in Atm knockout mice', *Neoplasia*, 16: 129-36.
- Germain, Ronald N. 2002. 'T-cell development and the CD4-CD8 lineage decision', *Nature Reviews Immunology*, 2: 309-22.
- Gertsenstein, Marina, Lauryl M. J. Nutter, Tammy Reid, Monica Pereira, William L. Stanford, Janet Rossant, and Andras Nagy. 2010. 'Efficient Generation of Germ Line Transmitting Chimeras from C57BL/6N ES Cells by Aggregation with Outbred Host Embryos', *PloS One*, 5: e11260.
- Gilad, Shlomit, Rami Khosravi, Dganit Shkedy, Tamar Uziel, Yael Ziv, Kinneret Savitsky, Galit Rotman, Sara Smith, Luciana Chessa, Timothy J. Jorgensen, Reli Harnik, Moshe Frydman, Ozden Sanal, Sima Portnoi, Zipora Goldwicz, N. G. J. Jaspers, Richard A. Gatti, Gilbert Lenoir, Martin F. Lavin, Kouichi Tatsumi, Rolf D. Wegner, Yosef Shiloh, and Anat Bar-Shira. 1996. 'Predominance of Null Mutations in Ataxia-Telangiectasia', *Human Molecular Genetics*, 5: 433-39.
- Gueven, N., O. J. Becherel, A. W. Kijas, P. Chen, O. Howe, J. H. Rudolph, R. Gatti, H. Date, O. Onodera, G. Taucher-Scholz, and M. F. Lavin. 2004. 'Aprataxin, a novel protein that protects against genotoxic stress', *Human Molecular Genetics*, 13: 1081-93.
- Guo, Zhi, Sergei Kozlov, Martin F. Lavin, Maria D. Person, and Tanya T. Paull. 2010. 'ATM Activation by Oxidative Stress', *Science*, 330: 517-21.
- Hansel, Christian, and David J. Linden. 2000. 'Long-Term Depression of the Cerebellar Climbing Fiber-Purkinje Neuron Synapse', *Neuron*, 26: 473-82.
- Hansen, Stephen T, Pratap Meera, Thomas S Otis, and Stefan M Pulst. 2013. 'Changes in Purkinje cell firing and gene expression precede behavioral pathology in a mouse model of SCA2', *Human Molecular Genetics*, 22: 271-83.

Perez et al. Resubmission

- Hathcock, K. S., S. Bowen, F. Livak, and R. J. Hodes. 2013. 'ATM influences the efficiency of TCRbeta rearrangement, subsequent TCRbeta-dependent T cell development, and generation of the pre-selection TCRbeta CDR3 repertoire', *PloS One*, 8: e62188.
- Herzog, Karl-Heinz, Miriam J. Chong, Manuela Kapsetaki, James I. Morgan, and Peter J. McKinnon. 1998. 'Requirement for Atm in Ionizing Radiation-Induced Cell Death in the Developing Central Nervous System', *Science*, 280: 1089-91.
- Hoche, F., K. Seidel, M. Theis, S. Vlaho, R. Schubert, S. Zielen, and M. Kieslich. 2012. 'Neurodegeneration in ataxia telangiectasia: what is new? What is evident?', *Neuropediatrics*, 43: 119-29.
- Hourez, Raphael, Laurent Servais, David Orduz, David Gall, Isabelle Millard, Alban de Kerchove d'Exaerde, Guy Cheron, Harry T Orr, Massimo Pandolfo, and Serge N Schiffmann. 2011. 'Aminopyridines correct early dysfunction and delay neurodegeneration in a mouse model of spinocerebellar ataxia type 1', *Journal of Neuroscience*, 31: 11795-807.
- Hoxha, Eriola, Ilaria Balbo, Maria Concetta Miniaci, and Filippo Tempia. 2018. 'Purkinje Cell Signaling Deficits in Animal Models of Ataxia', *Frontiers in Synaptic Neuroscience*, 10: 6-6.
- Hurlock, Edward C, Anne McMahon, and Rolf H Joho. 2008. 'Purkinje-cell-restricted restoration of Kv3.3 function restores complex spikes and rescues motor coordination in Kcnc3 mutants', *Journal of Neuroscience*, 28: 4640-48.
- Jackson, Thomas J, Gabriel Chow, Mohnish Suri, Philip Byrd, Malcolm R Taylor, and William P Whitehouse. 2016. 'Longitudinal analysis of the neurological features of ataxia-telangiectasia', *Developmental Medicine and Child Neurology*, 58: 690-97.
- Jayabal, Sriram, Hui Ho Vanessa Chang, Kathleen E Cullen, and Alanna J Watt. 2016. '4-aminopyridine reverses ataxia and cerebellar firing deficiency in a mouse model of spinocerebellar ataxia type 6', *Scientific Reports*, 6: 1-12.
- Kaminsky, Natalie, Ofer Bihari, Sivan Kanner, and Ari Barzilai. 2016. 'Connecting Malfunctioning Glial Cells and Brain Degenerative Disorders', *Genomics, Proteomics & Bioinformatics*, 14: 155-65.
- Kano, Masanobu, and Takaki Watanabe. 2017. 'Type-1 metabotropic glutamate receptor signaling in cerebellar Purkinje cells in health and disease', *F1000Research*, 6: 416-16.
- Kastan, M. B., and J. Bartek. 2004. 'Cell-cycle checkpoints and cancer', *Nature*, 432: 316-23.
- Kasumu, Adebimpe, and Ilya Bezprozvanny. 2012. 'Deranged calcium signaling in Purkinje cells and pathogenesis in spinocerebellar ataxia 2 (SCA2) and other ataxias', *The Cerebellum*, 11: 630-39.
- Kim, Jusik, Keeun Kim, Jung-Soon Mo, and Youngsoo Lee. 2020. 'Atm deficiency in the DNA polymerase β null cerebellum results in cerebellar ataxia and Itpr1 reduction associated with alteration of cytosine methylation', *Nucleic Acids Research*, 48: 3678-91.
- Krangel, Michael S. 2009. 'Mechanics of T cell receptor gene rearrangement', *Current Opinion in Immunology*, 21: 133-39.
- Larivière, Roxanne, Nicolas Sgarioto, Brenda Toscano Márquez, Rébecca Gaudet, Karine Choquet, R Anne McKinney, Alanna J Watt, and Bernard Brais. 2019. 'Sacs R272C missense homozygous mice develop an ataxia phenotype', *Molecular Brain*, 12: 1-15.
- Lavin, M. F. 2013. 'The appropriateness of the mouse model for ataxia-telangiectasia: neurological defects but no neurodegeneration', *DNA Repair (Amst)*, 12: 612-9.
- Levy, Ariel, and Anthony E. Lang. 2018. 'Ataxia-telangiectasia: A review of movement disorders, clinical features, and genotype correlations', *Movement Disorders*, 33: 1238-47.
- Li, J., Y. R. Han, M. R. Plummer, and K. Herrup. 2009. 'Cytoplasmic ATM in neurons modulates synaptic function', *Current Biology*, 19: 2091-6.
- Li, Jiali, Jianmin Chen, Harry V. Vinters, Richard A. Gatti, and Karl Herrup. 2011. 'Stable Brain ATM Message and Residual Kinase-Active ATM Protein in Ataxia-Telangiectasia', *The Journal of Neuroscience*, 31: 7568-77.
- Lindahl, T., and D. E. Barnes. 2000. 'Repair of endogenous DNA damage', *Cold Spring Harbor Symposia on Quantitative Biology*, 65: 127-33.

Perez et al. Resubmission

- 1082 Liu, Jing, Tie-Shan Tang, Huiping Tu, Omar Nelson, Emily Herndon, Duong P Huynh, Stefan M Pulst,
1083 and Ilya Bezprozvanny. 2009. 'Deranged calcium signaling and neurodegeneration in
1084 spinocerebellar ataxia type 2', *Journal of Neuroscience*, 29: 9148-62.
- 1085 Livák, Ferenc, Michelle Tourigny, David G. Schatz, and Howard T. Petrie. 1999. 'Characterization of
1086 TCR Gene Rearrangements During Adult Murine T Cell Development', *The Journal of*
1087 *Immunology*, 162: 2575-80.
- 1088 Lorenzetti, Diego, Kei Watase, Bisong Xu, Martin M Matzuk, Harry T Orr, and Huda Y Zoghbi. 2000.
1089 'Repeat instability and motor incoordination in mice with a targeted expanded CAG repeat in the
1090 Sca1 locus', *Human Molecular Genetics*, 9: 779-85.
- 1091 Marshall, Jean S., Richard Warrington, Wade Watson, and Harold L. Kim. 2018. 'An introduction to
1092 immunology and immunopathology', *Allergy, asthma, and clinical immunology : official journal of*
1093 *the Canadian Society of Allergy and Clinical Immunology*, 14: 49-49.
- 1094 Matei, I. R., C. J. Guidos, and J. S. Danska. 2006. 'ATM-dependent DNA damage surveillance in T-cell
1095 development and leukemogenesis: the DSB connection', *Immunological Reviews*, 209: 142-58.
- 1096 Matsuoka, Shuhei, Bryan A. Ballif, Agata Smogorzewska, E. Robert McDonald, Kristen E. Hurov, Ji
1097 Luo, Corey E. Bakalarski, Zhenming Zhao, Nicole Solimini, Yaniv Lerenthal, Yosef Shiloh,
1098 Steven P. Gygi, and Stephen J. Elledge. 2007. 'ATM and ATR Substrate Analysis Reveals
1099 Extensive Protein Networks Responsive to DNA Damage', *Science*, 316: 1160-66.
- 1100 McKinnon, P. J. 2009. 'DNA repair deficiency and neurological disease', *Nature Reviews:*
1101 *Neuroscience*, 10: 100-12.
- 1102 Meagher, M., and R. N. Lightowlers. 2014. 'The role of TDP1 and APTX in mitochondrial DNA repair',
1103 *Biochimie*, 100: 121-4.
- 1104 Micol, R., L. Ben Slama, F. Suarez, L. Le Mignot, J. Beauté, N. Mahlaoui, C. Dubois d'Enghien, A.
1105 Laugé, J. Hall, J. Couturier, L. Vallée, B. Delobel, F. Rivier, K. Nguyen, T. Billette de Villemeur,
1106 J. L. Stephan, P. Bordigoni, Y. Bertrand, N. Aladjidi, J. M. Pedespan, C. Thomas, I. Pellier, M.
1107 Koenig, O. Hermine, C. Picard, D. Moshous, B. Neven, F. Lanternier, S. Blanche, M. Tardieu,
1108 M. Debré, A. Fischer, and D. Stoppa-Lyonnet. 2011. 'Morbidity and mortality from ataxia-
1109 telangiectasia are associated with ATM genotype', *Journal of Allergy and Clinical Immunology*,
1110 128: 382-9.e1.
- 1111 Monaco, S., E. Nardelli, G. Moretto, T. Cavallaro, and N. Rizzuto. 1988. 'Cytoskeletal pathology in
1112 ataxia-telangiectasia', *Clinical Neuropathology*, 7: 44-6.
- 1113 Moreira, M. C., S. Klur, M. Watanabe, A. H. Németh, I. Le Ber, J. C. Moniz, C. Tranchant, P. Aubourg,
1114 M. Tazir, L. Schöls, M. Pandolfo, J. B. Schulz, J. Pouget, P. Calvas, M. Shizuka-Ikeda, M. Shoji,
1115 M. Tanaka, L. Izatt, C. E. Shaw, A. M'Zahem, E. Dunne, P. Bomont, T. Benhassine, N.
1116 Bouslam, G. Stevanin, A. Brice, J. Guimarães, P. Mendonça, C. Barbot, P. Coutinho, J.
1117 Sequeiros, A. Dürr, J. M. Warter, and M. Koenig. 2004. 'Senataxin, the ortholog of a yeast RNA
1118 helicase, is mutant in ataxia-ocular apraxia 2', *Nature Genetics*, 36: 225-7.
- 1119 Morrell, Daphne, Elizabeth Cromartie, and Michael Swift. 1986. 'Mortality and cancer incidence in 263
1120 patients with ataxia-telangiectasia', *Journal of the National Cancer Institute*, 77: 89-92.
- 1121 Paula-Barbosa, M. M., C. Ruela, M. A. Tavares, C. Pontes, A. Saraiva, and C. Cruz. 1983. 'Cerebellar
1122 cortex ultrastructure in ataxia-telangiectasia', *Annals of Neurology*, 13: 297-302.
- 1123 Perkins, Emma M, Yvonne L Clarkson, Nancy Sabatier, David M Longhurst, Christopher P Millward,
1124 Jennifer Jack, Junko Toraiwa, Mitsunori Watanabe, Jeffrey D Rothstein, and Alastair R Lyndon.
1125 2010. 'Loss of β -III spectrin leads to Purkinje cell dysfunction recapitulating the behavior and
1126 neuropathology of spinocerebellar ataxia type 5 in humans', *Journal of Neuroscience*, 30: 4857-
1127 67.
- 1128 Petersen, Andrew J., Stacey A. Rimkus, and David A. Wassarman. 2012. 'ATM kinase inhibition in glial
1129 cells activates the innate immune response and causes neurodegeneration in
1130 *Drosophila*', *Proceedings of the National Academy of Sciences*, 109: E656-E64.
- 1131 Pizzamiglio, L., E. Focchi, and F. Antonucci. 2020. 'ATM Protein Kinase: Old and New Implications in
1132 Neuronal Pathways and Brain Circuitry', *Cells*, 9.
- 1133 Quarantelli, Mario, Giuliana Giardino, Anna Prinster, Giuseppina Aloj, Barbara Carotenuto, Emilia
1134 Cirillo, Angela Marsili, Elena Salvatore, Ennio Del Giudice, and Claudio Pignata. 2013. 'Steroid

Perez et al. Resubmission

- 1135 treatment in Ataxia-Telangiectasia induces alterations of functional magnetic resonance imaging
- 1136 during pronosupination task', *European Journal of Paediatric Neurology*, 17: 135-40.
- 1137 Quek, Hazel, John Luff, KaGeen Cheung, Sergei Kozlov, Magtoug Gatei, C. Soon Lee, Mark C.
- 1138 Bellingham, Peter G. Noakes, Yi Chieh Lim, Nigel L. Barnett, Steven Dingwall, Ernst Wolvetang,
- 1139 Tomoji Mashimo, Tara L. Roberts, and Martin F. Lavin. 2016. 'A rat model of ataxia-
- 1140 telangiectasia: evidence for a neurodegenerative phenotype', *Human Molecular Genetics*, 26:
- 1141 109-23.
- 1142 Rashi-Elkeles, S., R. Elkon, N. Weizman, C. Linhart, N. Amariglio, G. Sternberg, G. Rechavi, A.
- 1143 Barzilai, R. Shamir, and Y. Shiloh. 2006. 'Parallel induction of ATM-dependent pro- and
- 1144 antiapoptotic signals in response to ionizing radiation in murine lymphoid tissue', *Oncogene*, 25:
- 1145 1584-92.
- 1146 Rass, Ulrich, Ivan Ahel, and Stephen C. West. 2007. 'Defective DNA Repair and Neurodegenerative
- 1147 Disease', *Cell*, 130: 991-1004.
- 1148 Rogers, D. C., E. M. Fisher, S. D. Brown, J. Peters, A. J. Hunter, and J. E. Martin. 1997. 'Behavioral
- 1149 and functional analysis of mouse phenotype: SHIRPA, a proposed protocol for comprehensive
- 1150 phenotype assessment', *Mammalian Genome*, 8: 711-3.
- 1151 Rothblum-Oviatt, Cynthia, Jennifer Wright, Maureen A. Lefton-Greif, Sharon A. McGrath-Morrow,
- 1152 Thomas O. Crawford, and Howard M. Lederman. 2016. 'Ataxia telangiectasia: a review',
- 1153 *Orphanet Journal of Rare Diseases*, 11: 159.
- 1154 Sahama, Ishani, Kate Sinclair, Simona Fiori, James Doecke, Kerstin Pannek, Lee Reid, Martin Lavin,
- 1155 and Stephen Rose. 2015. 'Motor pathway degeneration in young ataxia telangiectasia patients:
- 1156 A diffusion tractography study', *NeuroImage: Clinical*, 9: 206-15.
- 1157 Sahama, Ishani, Kate Sinclair, Kerstin Pannek, Martin Lavin, and Stephen Rose. 2014. 'Radiological
- 1158 imaging in ataxia telangiectasia: a review', *The Cerebellum*, 13: 521-30.
- 1159 Sandoval, N., M. Platzer, A. Rosenthal, T. Dork, R. Bendix, B. Skawran, M. Stuhmann, R. D. Wegner,
- 1160 K. Sperling, S. Banin, Y. Shiloh, A. Baumer, U. Bernthaler, H. Sennefelder, M. Brohm, B. H.
- 1161 Weber, and D. Schindler. 1999. 'Characterization of ATM gene mutations in 66 ataxia
- 1162 telangiectasia families', *Human Molecular Genetics*, 8: 69-79.
- 1163 Sanghez, Valentina, Anna Luzzi, Don Clarke, Dustin Kee, Steven Beuder, Danielle Rux, Mitsujiro
- 1164 Osawa, Joaquín Madrenas, Tsui-Fen Chou, Michael Kyba, and Michelina Iacovino. 2017.
- 1165 'Notch activation is required for downregulation of HoxA3-dependent endothelial cell phenotype
- 1166 during blood formation', *PloS One*, 12: e0186818.
- 1167 Savitsky, K., A. Bar-Shira, S. Gilad, G. Rotman, Y. Ziv, L. Vanagaite, D. A. Tagle, S. Smith, T. Uziel, S.
- 1168 Sfez, M. Ashkenazi, I. Pecker, M. Frydman, R. Harnik, S. R. Patanjali, A. Simmons, G. A.
- 1169 Clines, A. Sarti, R. A. Gatti, L. Chessa, O. Sanal, M. F. Lavin, N. G. Jaspers, A. M. Taylor, C.
- 1170 F. Arlett, T. Miki, S. M. Weissman, M. Lovett, F. S. Collins, and Y. Shiloh. 1995. 'A single ataxia
- 1171 telangiectasia gene with a product similar to PI-3 kinase', *Science*, 268: 1749-53.
- 1172 Schellenberg, M. J., P. P. Tumbale, and R. S. Williams. 2015. 'Molecular underpinnings of Aprataxin
- 1173 RNA/DNA deadenylase function and dysfunction in neurological disease', *Progress in*
- 1174 *Biophysics and Molecular Biology*, 117: 157-65.
- 1175 Schubert, R., J. Reichenbach, and S. Zielen. 2002. 'Deficiencies in CD4+ and CD8+ T cell subsets in
- 1176 ataxia telangiectasia', *Clinical and Experimental Immunology*, 129: 125-32.
- 1177 Sedghi, Maryam, Mehri Salari, Ali-Reza Moslemi, Ariana Kariminejad, Mark Davis, Hayley Goullée,
- 1178 Björn Olsson, Nigel Laing, and Homa Tajsharghi. 2018. 'Ataxia-telangiectasia-like disorder in a
- 1179 family deficient for MRE11A, caused by a MRE11 variant', *Neurology Genetics*, 4:
- 1180 e295.
- 1181 Shakkottai, Vikram G, Maria do Carmo Costa, James M Dell'Orco, Ananthakrishnan
- 1182 Sankaranarayanan, Heike Wulff, and Henry L Paulson. 2011a. 'Early changes in cerebellar
- 1183 physiology accompany motor dysfunction in the polyglutamine disease spinocerebellar ataxia
- 1184 type 3', *Journal of Neuroscience*, 31: 13002-14.
- 1185 Shakkottai, Vikram G, Maolei Xiao, Lin Xu, Michael Wong, Jeanne M Nerbonne, David M Ornitz, and
- 1186 Kelvin A Yamada. 2009. 'FGF14 regulates the intrinsic excitability of cerebellar Purkinje
- 1187 neurons', *Neurobiology of Disease*, 33: 81-88.

Perez et al. Resubmission

- Shakkottai, Vikram G., Maria do Carmo Costa, James M. Dell'Orco, Ananthakrishnan Sankaranarayanan, Heike Wulff, and Henry L. Paulson. 2011b. 'Early Changes in Cerebellar Physiology Accompany Motor Dysfunction in the Polyglutamine Disease Spinocerebellar Ataxia Type 3', *The Journal of Neuroscience*, 31: 13002-14.
- Shiloh, Yosef. 2020. 'The cerebellar degeneration in ataxia-telangiectasia: A case for genome instability', *DNA Repair*, 95: 102950.
- Shiloh, Yosef, and Yael Ziv. 2013. 'The ATM protein kinase: regulating the cellular response to genotoxic stress, and more', *Nature Reviews Molecular Cell Biology*, 14: 197-210.
- Solitare, G. B. 1968. 'Louis-Bar's syndrome (ataxia-telangiectasia). Anatomic considerations with emphasis on neuropathologic observations', *Neurology*, 18: 1180-6.
- Solitare, G. B., and V. F. Lopez. 1967. 'Louis-bar's syndrome (ataxia-telangiectasia). Neuropathologic observations', *Neurology*, 17: 23-31.
- Spring, K., S. Cross, C. Li, D. Watters, L. Ben-Senior, P. Waring, F. Ahangari, S. L. Lu, P. Chen, I. Misko, C. Paterson, G. Kay, N. I. Smorodinsky, Y. Shiloh, and M. F. Lavin. 2001. 'Atm knock-in mice harboring an in-frame deletion corresponding to the human ATM 7636del9 common mutation exhibit a variant phenotype', *Cancer Research*, 61: 4561-8.
- Staples, E. R., E. M. McDermott, A. Reiman, P. J. Byrd, S. Ritchie, A. M. Taylor, and E. G. Davies. 2008. 'Immunodeficiency in ataxia telangiectasia is correlated strongly with the presence of two null mutations in the ataxia telangiectasia mutated gene', *Clinical and Experimental Immunology*, 153: 214-20.
- Stoyas, Colleen A, David D Bushart, Pawel M Switonski, Jacqueline M Ward, Akshay Alaghata, Mi-bo Tang, Chenchu Niu, Mandheer Wadhwa, Haoran Huang, and Alex Savchenko. 2020. 'Nicotinamide pathway-dependent Sirt1 activation restores calcium homeostasis to achieve neuroprotection in spinocerebellar ataxia type 7', *Neuron*, 105: 630-44. e9.
- Strich, Sabina J. 1966. 'Pathological findings in three cases of ataxia-telangiectasia', *Journal of Neurology, Neurosurgery, and Psychiatry*, 29: 489.
- Swift, M., D. Morrell, E. Cromartie, A. R. Chamberlin, M. H. Skolnick, and D. T. Bishop. 1986. 'The incidence and gene frequency of ataxia-telangiectasia in the United States', *American Journal of Human Genetics*, 39: 573-83.
- Sykora, Peter, Deborah L. Croteau, Vilhelm A. Bohr, and David M. Wilson, 3rd. 2011. 'Aprataxin localizes to mitochondria and preserves mitochondrial function', *Proceedings of the National Academy of Sciences of the United States of America*, 108: 7437-42.
- Takashima, H., C. F. Boerkoel, J. John, G. M. Saifi, M. A. Salih, D. Armstrong, Y. Mao, F. A. Quiocho, B. B. Roa, M. Nakagawa, D. W. Stockton, and J. R. Lupski. 2002. 'Mutation of TDP1, encoding a topoisomerase I-dependent DNA damage repair enzyme, in spinocerebellar ataxia with axonal neuropathy', *Nature Genetics*, 32: 267-72.
- Tal, Efrat, Marina Alfo, Shan Zha, Ari Barzilai, Chris I. De Zeeuw, Yael Ziv, and Yosef Shiloh. 2018. 'Inactive Atm abrogates DSB repair in mouse cerebellum more than does Atm loss, without causing a neurological phenotype', *DNA Repair*, 72: 10-17.
- Tavani, F., R. A. Zimmerman, G. T. Berry, K. Sullivan, R. Gatti, and P. Bingham. 2003. 'Ataxia-telangiectasia: the pattern of cerebellar atrophy on MRI', *Neuroradiology*, 45: 315-19.
- Taylor, A. M., Z. Lam, J. I. Last, and P. J. Byrd. 2015. 'Ataxia telangiectasia: more variation at clinical and cellular levels', *Clinical Genetics*, 87: 199-208.
- Terplan, K. L., and R. F. Krauss. 1969. 'Histopathologic brain changes in association with ataxia-telangiectasia', *Neurology*, 19: 446-54.
- Tsutsumi, S., M. Yamazaki, T. Miyazaki, M. Watanabe, K. Sakimura, M. Kano, and K. Kitamura. 2015. 'Structure-function relationships between aldolase C/zebrin II expression and complex spike synchrony in the cerebellum', *Journal of Neuroscience*, 35: 843-52.
- Tumbale, Percy, Matthew J Schellenberg, Geoffrey A Mueller, Emma Fairweather, Mandy Watson, Jessica N Little, Juno Krahn, Ian Waddell, Robert E London, and R Scott Williams. 2018. 'Mechanism of APTX nicked DNA sensing and pleiotropic inactivation in neurodegenerative disease', *The EMBO Journal*, 37: e98875.

Perez et al. Resubmission

- 1240 Vacchio, Melanie S., Alexandru Olaru, Ferenc Livak, and Richard J. Hodes. 2007. 'ATM deficiency
1241 impairs thymocyte maturation because of defective resolution of T cell receptor α locus coding
1242 end breaks', *Proceedings of the National Academy of Sciences*, 104: 6323-28.
- 1243 Vail, Graham, Aifang Cheng, Yu Ray Han, Teng Zhao, Shengwang Du, Michael M. T. Loy, Karl Herrup,
1244 and Mark R. Plummer. 2016. 'ATM protein is located on presynaptic vesicles and its deficit
1245 leads to failures in synaptic plasticity', *Journal of Neurophysiology*, 116: 201-09.
- 1246 Valentin-Vega, Y. A., and M. B. Kastan. 2012. 'A new role for ATM: regulating mitochondrial function
1247 and mitophagy', *Autophagy*, 8: 840-1.
- 1248 van der Burgt, I, K H Chrzanowska, D Smeets, and C Weemaes. 1996. 'Nijmegen breakage syndrome',
1249 *Journal of Medical Genetics*, 33: 153-56.
- 1250 Verhagen, Mijke M.M., Jean-Jacques Martin, Marcel van Deuren, Chantal Ceuterick-de Groote, Corry
1251 M.R. Weemaes, Berry H.P.H. Kremer, Malcolm A.R. Taylor, Michèl A.A.P. Willemsen, and
1252 Martin Lammens. 2012. 'Neuropathology in classical and variant ataxia-telangiectasia',
1253 *Neuropathology*, 32: 234-44.
- 1254 Voogd, Jan, and Mitchell Glickstein. 1998. 'The anatomy of the cerebellum', *Trends in Neurosciences*,
1255 21: 370-75.
- 1256 Wakasugi, M., T. Sasaki, M. Matsumoto, M. Nagaoka, K. Inoue, M. Inobe, K. Horibata, K. Tanaka, and
1257 T. Matsunaga. 2014. 'Nucleotide excision repair-dependent DNA double-strand break formation
1258 and ATM signaling activation in mammalian quiescent cells', *Journal of Biological Chemistry*,
1259 289: 28730-7.
- 1260 Wallis, LI, PD Griffiths, SJ Ritchie, CAJ Romanowski, G Darwent, and ID Wilkinson. 2007. 'Proton
1261 spectroscopy and imaging at 3T in ataxia-telangiectasia', *American journal of neuroradiology*,
1262 28: 79-83.
- 1263 Walter, Joy T, Karina Alvina, Mary D Womack, Carolyn Chevez, and Kamran Khodakhah. 2006.
1264 'Decreases in the precision of Purkinje cell pacemaking cause cerebellar dysfunction and
1265 ataxia', *Nature Neuroscience*, 9: 389-97.
- 1266 Weyemi, Urbain, Christophe E. Redon, Towqir Aziz, Rohini Choudhuri, Daisuke Maeda, Palak R.
1267 Parekh, Michael Y. Bonner, Jack L. Arbiser, and William M. Bonner. 2015. 'NADPH oxidase 4 is
1268 a critical mediator in Ataxia telangiectasia disease', *Proceedings of the National Academy of
1269 Sciences*, 112: 2121-26.
- 1270 Xu, Y., and D. Baltimore. 1996. 'Dual roles of ATM in the cellular response to radiation and in cell
1271 growth control', *Genes and Development*, 10: 2401-10.
- 1272 Yamasaki, Miwako, Kouichi Hashimoto, and Masanobu Kano. 2006. 'Miniature Synaptic Events Elicited
1273 by Presynaptic Ca^{2+} Rise Are Selectively Suppressed by Cannabinoid Receptor Activation in
1274 Cerebellar Purkinje Cells', *The Journal of Neuroscience*, 26: 86-95.
- 1275 Zhou, Haibo, Zhanmin Lin, Kai Voges, Chiheng Ju, Zhenyu Gao, Laurens W. J. Bosman, Tom J. H.
1276 Ruigrok, Freek E. Hoebeek, Chris I. De Zeeuw, and Martijn Schonewille. 2014. 'Cerebellar
1277 modules operate at different frequencies', *Elife*, 3: e02536.
- 1278 Zhu, L., B. Scelfo, F. Tempia, B. Sacchetti, and P. Strata. 2006. 'Membrane excitability and fear
1279 conditioning in cerebellar Purkinje cell', *Neuroscience*, 140: 801-10.

1280

8.0 Figure Legends

Figure 1. New A-T mouse models expressing clinically related PTCs. A) The *Atm* gene locus was targeted by homologous recombination of a targeting vector containing a modified NorCOMM cassette in intron one and the corresponding A-T PTC mutation in exon 3 to create the targeted *Atm*^{R35X} and *Atm*^{Q35X} ES cell lines. Following germline transmission of these alleles in mice, the floxed NorCOMM cassette was removed by Cre excision *in vivo* to produce the final *Atm*^{R35X} and *Atm*^{Q35X} mouse lines. **B)** Genotyping of A-T mouse models. PCR agarose gel of mouse DNA shows 151 bp wildtype (+) allele band and 241 bp Cre-excised targeted allele band. **C)** ATM levels were examined using immunoblot analyses of the spleen due to its high expression density in this tissue. Exemplar blots illustrate a gene dose effect of ATM protein expression in samples harvested from wildtype (+), heterozygous (R35X/+, Q35X/+), and homozygous *Atm*^{R35X/R35X} (R35X) and *Atm*^{Q35X/Q35X} (Q35X) mice as indicated. **D)** Breeding scheme schematic for double mutant and control mice for this study. **E)** *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice develop an ataxia that at late stages results in a severe loss of motor coordination and ability to ambulate (see **Videos 1-4**). Abbreviations for panel 1: **hβA**-human beta Actin promotor; **ΔTK1**-delta TK1, inactivated Thymidine Kinase 1; **T2A**-self-cleaving peptide sequence; **Neo**-Neomycin gene; **PGKpA**-Phosphoglycerate kinase poly A tail; **loxP**-recombination elements are shown as a blue triangle; orientation of the Gateway **attB** recombination elements by an orange arrow; orientation of the genotyping **F** and **R** primers is shown by green and blue arrows respectively; and engineered PTC sites are shown in exon 3 by a red circle.

Figure 2. Health and survivability of single and double mutant mice. A) Left: The line color and symbol for each genotype is denoted and is consistent across all figures (1-8). **Right:** *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice weighed significantly less than all control genotypes as indicated by the growth curves (± 95% confidence interval; dotted lines). Growth curve (*Atm*^{R35X/R35X}; *Aptx*^{-/-} vs. controls): Male k = 0.024 vs. 0.011-0.019, Y_{max} = 21.8 vs. 32.9-41.0 g, (n = 3 to 18); Female k = 0.030 vs. 0.017-0.022, Y_{max} = 16.9 vs. 23.3-31.3, (n = 2 to 19). Sum of squares F-test run across all curves: Male F_(12, 364) = 30.5, ****p<0.0001, Female F_(12, 339) = 28.3, ****p<0.0001. **B)** ATM deficient mice, regardless of APTX

Perez et al. Resubmission

expression, displayed significantly lower survivability with ~55% of mice deceased by P400. No statistical differences between ATM deficient mice were detected. Moreover, a single wildtype copy of the *Atm* gene was sufficient to prevent premature death (no statistical difference detected between *Atm*^{R35X/+}; *Aptx*^{-/-} and *Atm*^{+/+}; *Aptx*^{+/+} mice). Log-rank (Mantel-Cox) tests across all ($\chi^2_{(6, 217)} = 48.4$, **** $p < 0.0001$), just the ATM deficient ($\chi^2_{(2, 217)} = 1.06$, $p = 0.6$), and single comparisons to wildtype (see figure) were conducted. Total number of animals indicated in panel **C**. **C**) Pie charts illustrating that ATM deficient mice displayed a high prevalence of thymomas based on postmortem necropsies. “Other” probable causes of death included enlarged livers, and obstructed kidneys. “Missing” mice were presumed dead and cannibalized by cage mates, cause of death unknown. **Figure 2-figure supplement 1, Figure 2-source data**

Figure 2-figure supplement 1. Animal weight for each time point and genotype. A) The average weights are plotted for each genotype at each of the indicated time points. Growth curves without experimental, 2-way ANOVA with age and genotype as factors. Male: $F_{(10, 226)} = 5.6$, $p < 0.0001$; Female: $F_{(10, 197)} = 7.3$, $p < 0.0001$. **B)** The survivability of each genotype of mice are plotted for male and female individually.

Figure 3. *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice develop a progressive loss in motor coordination. A) *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice take a similar amount of time to descend a vertical pole at P45, 120, and 210, but significantly longer at P400. These overall results were found to be similar for both male (left, $n = 2$ to 12) and female (right, $n = 4$ to 12) mice. **B)** Consistent with the vertical pole test, the gait of *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice measured during ambulation on a Catwalk gait analysis system was significantly different to controls by P400, but not before P210. This includes the percent of time a mouse spends with 3 vs. 1, 2, or 4 paws on the ground and the speed and cadence during each run across the platform. The effects of the two null mutations were generally similar between males (left, $n = 4$ to 21) and females (right, $n = 3$ to 18). **C)** Behavioral data for male (blue) and female (pink) *Atm*^{R35X/R35X}; *Aptx*^{-/-} (dark purple, left) and *Atm*^{+/+}; *Aptx*^{+/+} (orange, right) mice are plotted at P30 (left 2 columns) and 400 (right 2 columns). Left-right asymmetries of the horizontal bars indicate a difference

Perez et al. Resubmission

in performance between genotypes for the behavioral test listed in a column on the far left. A significant difference in the time to right during the righting reflex at P8 was observed in both Male and Female mice (bottom). **A** and **B** were examined via two-way ANOVA with age and genotype as factors followed by *postshot* Tukey's multiple comparison tests between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and each of the control genotypes. Behavioral tests in **C** were examined using a non-parametric Kruskal Wallace followed by *postdocs* Dunn's multiple comparisons tests. Symbol/color key: $Atm^{+/+}$; $Aptx^{+/+}$ (purple circle), $Atm^{+/+}$; $Aptx^{-/-}$ (blue diamond), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ (green triangle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange square), $Atm^{R35X/+}$; $Aptx^{-/-}$ (red inverted triangle) **Figure 3-figure supplement 1, Figure 3-source data**

Figure 3-figure supplement 1. $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice develop progressive ataxia. A) $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice develop an ataxia that at late stages results in a severe loss of coordination and ability to ambulate. **B)** Additional gait analyses measuring stride length and time of overlap of the hindlimbs. Examined via two-way ANOVA with age and genotype as factors followed by *postshot* Tukey's multiple comparison tests between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and each of the control genotypes. **C)** Behavioral deficits are seen only in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice across all behavioral tests and sexes. Behavioral tests were examined using a non-parametric Kruskal Wallace followed by *posthoc* Dunn's multiple comparisons tests.

Figure 4. The biophysical properties of PNs are significantly perturbed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice. A) Schematic diagram of intracellular recording from a single Purkinje neuron (PN) in an acute cerebellar tissue slice preparation used to examine their biophysical properties. **B) Left:** Voltage clamp measurements of PN neuron membrane properties were made from a 1 s, -5 mV step pulse as illustrated. **Right:** The membrane input resistance (R_m), time constant (τ), and capacitance (C_m) were perturbed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ compared to $Atm^{+/+}$; $Aptx^{+/+}$ mice. **C)** Current clamp recordings of PN action potentials (AP) after 2 nA step pulses from a -70 mV holding potential. PN action potentials recorded from $Atm^{R35X/R35X}$; $Aptx^{-/-}$ fail to maintain constant firing and summary plots show that they have lower 1st AP amplitudes, firing threshold, and area under the curve. **D) Top:** Example mEPSC traces taken from a PN under voltage clamp at a -80 mV holding potential. **Bottom:** Median frequency

Perez et al. Resubmission

and amplitude data, along with the overall probability distribution function are plotted for both *Atm*^{+/+}; *Aptx*^{+/+} (n = 11) and *Atm*^{R35X/R35X}; *Aptx*^{-/-} (n = 11) mice. The frequency, but not amplitude of PNs recorded in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice was found to be perturbed. **E and F) Left:** Example traces of evoked EPSCs recorded from PNs as a result of a 2-pulse stimulation (50 ms interval) of either parallel (**E**) or climbing (**F**) axon fibers. Traces illustrate the first (A₁) and second (A₂) amplitude (normalized) and time course of first decay (blue fitted line) of each synaptic response. **Right:** Summary plots of the paired pulse ratio. While parallel fiber paired pulse facilitation was normal in *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice, climbing fiber paired pulse depression and halfwidth was significantly perturbed compared to *Atm*^{+/+}; *Aptx*^{+/+} mice. Data in **B** were compared using an ANOVA (Kruskal-Wallis) followed by Dunn's multiple comparisons test while data in **D to F** were compared via Welch's *t*-test. Symbol/color key: *Atm*^{+/+}; *Aptx*^{+/+} (purple circle), *Atm*^{R35X/R35X}; *Aptx*^{-/-} (orange square) **Figure 4-figure supplement 1**

Figure 4-figure supplement 1. Current vs. voltage responses significantly differ between *Atm*^{+/+}; *Aptx*^{+/+} and *Atm*^{R35X/R35X}; *Aptx*^{-/-} mice. A) PN voltage responses to various current steps between -500 and 2250 pA (250 pA steps) from a -70 mV holding current in *Atm*^{+/+}; *Aptx*^{+/+} (top, purple) and *Atm*^{R35X/R35X}; *Aptx*^{-/-} (bottom, orange) mice. **(B)** I-V curves calculated from either max deflection (V_{m max}) or steady state (V_{m end}) for *Atm*^{+/+}; *Aptx*^{+/+} (purple) and *Atm*^{R35X/R35X}; *Aptx*^{-/-} (orange) mice. **(C)** Various measurements of the voltage response to -500 pA step pulse (blue box in **B**) in *Atm*^{+/+}; *Aptx*^{+/+} (purple) and *Atm*^{R35X/R35X}; *Aptx*^{-/-} (orange) mice. Significance was tested using a non-parametric Mann Whitney test.

Figure 5. Cerebellar atrophy is associated with a progressive reduction in PN neural action potential firing frequency and PN dendritic length. A) Schematic diagram of extracellular recording from a single Purkinje neuron (PN) in an acute cerebellar tissue slice preparation. Example electrophysiological traces for *Atm*^{+/+}; *Aptx*^{+/+} (purple, top) and *Atm*^{R35X/R35X}; *Aptx*^{-/-} (orange, bottom) PNs in the medial (vermis) area of the cerebellum. **B)** *Atm*^{R35X/R35X}; *Aptx*^{-/-} PN action potential firing frequency progressively decreased with age and was significantly slower in comparison to all control genotypes expressing at least one copy of the *Atm* or *Aptx* gene. [*Atm*^{+/+}; *Aptx*^{+/+} (n=52 to 59), *Atm*^{+/+}; *Aptx*^{-/-} (n =

Perez et al. Resubmission

51 to 64), $Atm^{R35X/R35X}; Aptx^{+/+}$ (n = 39 to 52), $Atm^{R35X/R35X}; Aptx^{-/-}$ (n = 24 to 71), $Atm^{R35X/+}; Aptx^{-/-}$ (n = 69)] **C)** Cartoon image of the brain highlighting the dorsal forebrain and cerebellar surface. 2-dimensional area estimates from dorsal images of the brain were used to determine the cerebellum to forebrain ratio allowing us to control for any differences in overall size of the brain. We found the cerebellum decreased in size over age in $Atm^{R35X/R35X}; Aptx^{-/-}$ (n = 5 to 10), but not control mice [$Atm^{+/+}; Aptx^{+/+}$ (n = 4 to 20), $Atm^{+/+}; Aptx^{-/-}$ (n = 4 to 12), $Atm^{R35X/R35X}; Aptx^{+/+}$ (n = 6 to 16), $Atm^{R35X/+}; Aptx^{-/-}$ (n = 6)]. **D)** Immunofluorescent images of $Atm^{+/+}; Aptx^{+/+}$ (top) and $Atm^{R35X/R35X}; Aptx^{-/-}$ (bottom) at P400 (medial cerebellar lobule VIII). Scale bar = 50 μ m **E)** The width of the molecular layer, but not the granule cell layer progressively declined over age (n = 5 to 7). Statistical significances were assessed via 2-way ANOVA with age and genotype as factors followed by *posthoc* Holm-Sidak (**B**, **C**, and **E (top)**) or Sidak (**E bottom**) pairwise multiple comparisons test. **Figure 5-figure supplements 1-5, Figure 5-source data**

Figure 5-figure supplement 1. Mean PN firing frequency across the cerebellum. Average PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400.

Figure 5-figure supplement 2. Mean PN firing frequency across genotype and sex. Average PN firing frequency for all cells recorded from male and female mice is plotted for the indicated genotype. No significant differences were observed between sex. 2-Way ANOVA with age and sex as factors, $Atm^{+/+}; Aptx^{+/+}$ ($F_{(1, 751)} = 1.15$, $p=0.3$), $Atm^{+/+}; Aptx^{-/-}$ ($F_{(1, 797)} = 1.10$, $p=0.3$), $Atm^{R35X/R35X}; Aptx^{+/+}$ ($F_{(1, 630)} = 0.17$, $p=0.7$), $Atm^{R35X/R35X}; Aptx^{-/-}$ ($F_{(1, 666)} = 1.10$, $p=0.4$), *t*-test for P400 $Atm^{R35X/+}; Aptx^{-/-}$ ($p=0.9$)

Figure 5-figure supplement 3. Coefficient of Variation of PN firing frequency across the cerebellum. Average CV of PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400. No significant differences ($p<0.5$) were detected across all areas using 2-way ANOVA with age and genotype as factors.

Figure 5-figure supplement 4. Mean variation between PN firing intervals across the cerebellum. Average CV2 of PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400.

Perez et al. Resubmission

No significant differences ($p < 0.5$) were detected across all areas using 2-way ANOVA with age and genotype as factors.

Figure 5-figure supplement 5. Histopathological effects of ATM and APTX deficiency. A) Width measurements of the molecular and granule cell layer (ML and GCL respectively) for each folia across the medial intermediate and lateral areas of the cerebellum. **B) (Top)** Summary plot indicates no significant differences in the 2D linear density of PN across each folia in $Atm^{R35X/R35X}; Aptx^{-/-}$ vs. $Atm^{+/+}; Aptx^{+/+}$. **(Bottom)** Images of parasagittal cerebellar slices (2.5X) illustrating the lack of significant gaps in PN (green, Calbindin) or granule cell layers (magenta, DAPI). Scale bar = 0.5 mm **(C)** Images of parasagittal sections from $Atm^{+/+}; Aptx^{+/+}$ (top row) and $Atm^{R35X/R35X}; Aptx^{-/-}$ (bottom rows) mice (63X). PNs (green, Calbindin) were found to have a larger diameter than those found in $Atm^{+/+}; Aptx^{+/+}$ cells (red arrows). PN axonal swellings were also observed in $Atm^{R35X/R35X}; Aptx^{-/-}$ sections (magenta arrows). Scale bar = 50 μ m

Figure 6. T-cell deficits are found in the blood of $Atm^{R35X/R35X}; Aptx^{-/-}$ mice. A) Representative flow cytometric profiles of T-cell glycoprotein marker CD3 and summary plots indicate ATM and/or APTX deficient mice have decreased proportions of CD3⁺ T-cells in the blood. **B)** Representative flow cytometric profiles of T-cell glycoprotein markers CD4 and CD8 gated on CD3⁺ cells and summary plots for CD8 and CD4 single positive cell proportions. ATM deficient mice had reduced CD4⁺ proportions compared to mice with at least one copy of the *Atm* gene. Statistical significances were assessed via 1-way ANOVA followed by *posthoc* Tukey's pairwise multiple comparisons tests. Number of animals denoted at bottom of bar. Symbol/color key: $Atm^{+/+}; Aptx^{+/+}$ (purple circle), $Atm^{R35X/R35X}; Aptx^{+/+}$ (green triangle), $Atm^{R35X/+}; Aptx^{-/-}$ (red inverted triangle), $Atm^{R35X/R35X}; Aptx^{-/-}$ (orange square) **Figure 6/7-source data**

Figure 7. ATM and APTX deficiency confer deficits in T-cell expression, but at different developmental stages. A) Representative flow cytometric profiles of T-cell glycoprotein markers CD44 and CD25 gated on CD4⁺CD8⁻ double negative (DN) cells. Summary plots show proportions of thymocytes at DN stages 1-4 (left to right). APTX deficient mice display increased proportions for DN1-

Perez et al. Resubmission

3 and decreased proportion at DN4 consistent with a deficit in ontogeny from DN3 to DN4. **B)** Representative flow cytometric profiles of T-cell glycoprotein markers CD4 and CD8 gated. ATM deficient mice display decreased proportions for CD4 and CD8 single positive cells consistent with a deficit in ontogeny from CD4⁺CD8⁺ double positive to CD4⁺ and CD8⁺ single positive fates. Statistical significances were assessed via 1-way ANOVA followed by *posthoc* Tukey's pairwise multiple comparisons tests. Number of animals denoted at bottom of bars. Symbol/color key: *Atm*^{+/+}; *Aptx*^{+/+} (purple circle), *Atm*^{R35X/R35X}; *Aptx*^{+/+} (green triangle), *Atm*^{R35X/+}; *Aptx*^{-/-} (red inverted triangle), *Atm*^{R35X/R35X}; *Aptx*^{-/-} (orange square)

Figure 8. ATM protein expression is restored after readthrough compound exposure in explant tissues from *Atm*^{R35X/R35X} and *Atm*^{Q35X/Q35X}. Spleen and cerebellar explant tissue from *Atm*^{R35X/R35X} and *Atm*^{+/+} mice were treated with vehicle, the readthrough compounds G418 (100 μM), or GJ103 (100 μM) for 72 hrs. ATM immunoblots show recovery of ATM (MW 350 kDa) production in both the spleen (n = 2) and cerebellum (n = 3). Equal loading was assessed via housekeeping genes (Actin or GAPDH) and ponceau staining. **Figure 8-source data**

11.0 Source files

Figure 2- source data. Weight, age of death, and probable cause of death

Figure 3-source data. Raw behavior data

Figure 4-source data. Electrophysiological data

Figure 5- source data. Individual average firing frequencies for each recorded cell; Individual CV for each recorded cell; Brain area

Figure 6/7- source data. Tables of FACs data

Figure 8-source data. Western blot measurements

12.0 Rich Media

Video 1. Pole test, *Atm*^{+/+} vs. *Atm*^{R35X/R35X}. *Atm*^{R35X/R35X} do not display an ataxic phenotype at P460.

Video 2. Pole test, *Aptx*^{+/+} vs. *Aptx*^{-/-}. *Aptx*^{-/-} mice do not display an ataxic phenotype at P460.

Video 3. Pole test, *Atm*^{+/+}; *Aptx*^{+/+} vs. *Atm*^{R35X/R35X}; *Aptx*^{-/-}. *Atm*^{R35X/R35X}; *Aptx*^{-/-} have considerable motor disability at P460.

Video 4. Open field, *Atm*^{+/+}; *Aptx*^{+/+} vs. *Atm*^{R35X/R35X}; *Aptx*^{-/-}. *Atm*^{R35X/R35X}; *Aptx*^{-/-} display a clear inability to ambulate in the open field at P460.

Perez et al. Resubmission

1465

1466

1467

1468

Fig. 1

Fig. 2

A

B

C

Fig. 3

A

Pole test

B

Gait Analyses

C

SHIRPA

30 Days old

400 Days old

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8