Short Title: Variation in C₄ pathways in maize growth

Authors for Contact: Jennifer J. Arp, jarp@danforthcenter.org; Doug K. Allen, doug.allen@usda.gov

Developmental Effects on Relative Use of PEPCK and NADP-ME Pathways of C₄ Photosynthesis in Maize

Jennifer J. Arp¹, Shrikaar Kambhampati¹, Kevin L. Chu¹, Somnath Koley¹, Lauren M. Jenkins^{1,2}, Todd C. Mockler¹, and Doug K. Allen^{1,2}.

(1) Donald Danforth Plant Science Center, Saint Louis, MO 63132 USA

(2) United States Department of Agriculture, Agricultural Research Service, Saint Louis, MO 63132 USA

One Sentence Summary: The proportion of the two C₄ pathways in maize plants is dependent on canopy position and not the age of the leaf.

Author Contributions: J.A. and D.K.A. conceived the research plans; S.Ka., S.Ko., K.C., L.J., and J.A. performed labeling and metabolite quantitation experiments; J.A. designed the experiments and analyzed the data; J.A. wrote the article with contributions of all the authors; D.K.A. supervised and completed the writing. D.K.A. agrees to serve as the author responsible for contact and ensures communication.

Responsibilities of the Author for Contact: It is the responsibility of the author for contact to ensure that all scientists who have contributed substantially to the conception, design or execution of the work described in the manuscript are included as authors, in accordance with the guidelines from the Committee on Publication Ethics (COPE) (<http://publicationethics.org/resources/guidelines>). It is the responsibility of the author for contact also to ensure that all authors agree to the list of authors and the identified contributions of those authors.

Funding Sources: Funding for the research was provided by the National Science Foundation Plant Genome Research Program (PGRP) postdoctoral fellowship (NSF-1812235) to J.A., and PGRP Award IOS-1829365, the United States Department of Agriculture National Institute of Food and Agriculture (2017-67013-26156; 2016-67013-24585), the Office of Biological and Environmental Research in the DOE Office of Science (DE-SC0008769), and the National Institutes of Health (U01 CA235508). Support was also provided by United States Department of Agriculture Agricultural Research Service. Support for the acquisition of the 6500 QTRAP LC-MS/MS was provided by the National Science Foundation (DBI #1427621).

Abstract

C₄ photosynthesis is an adaptive photosynthetic pathway which concentrates CO₂ around Rubisco in specialized bundle sheath cells to reduce photorespiration. Historically, the pathway has been characterized into three different subtypes based on the decarboxylase involved, although recent work has provided evidence that some plants can use multiple decarboxylases, with maize in particular using both the NADP-malic enzyme (NADP-ME) pathway and phosphoenolpyruvate carboxykinase (PEPCK) pathway. Parallel C₄ pathways could be advantageous in balancing energy and reducing equivalents between bundle sheath and mesophyll cells, in decreasing the size of the metabolite gradients between cells and may better accommodate changing environmental conditions or source to sink demands on growth. The enzyme activity of C₄ decarboxylases can fluctuate with different stages of leaf development, but it remains unclear if the pathway flexibility is an innate aspect of leaf development or an adaptation to the leaf microenvironment that is regulated by the plant. In this study, variation in the two C₄ pathways in maize were characterized at nine plant ages throughout the life cycle. Two positions in the canopy were examined for variation in physiology, gene expression, metabolite concentration, and enzyme activity, with particular interest in asparagine as a potential regulator of C₄ decarboxylase activity. Variation in C₄ and C₃ metabolism was observed for both leaf age and canopy position, reflecting the ability of C₄ pathways to adapt to changing microenvironments.

Introduction

C₄ photosynthesis is a beneficial adaptation to environmental conditions that enables plants to more effectively assimilate carbon dioxide relative to C₃ plants, resulting in some of the most productive crops on a biomass basis. C₄ photosynthesis has evolved over 60 times in plants to overcome the inefficiencies of Rubisco by minimizing the oxidation reaction and photorespiration (Sage et al., 2011). The pathway has historically been categorized into three subtypes based on the decarboxylase—NAD-malic enzyme (NAD-ME), NADP-malic enzyme (NADP-ME) or phosphoenolpyruvate carboxykinase (PEPCK)—but additional evidence has identified plants that can use a combination of decarboxylases in C₄ photosynthesis (Chapman and Hatch, 1981; Furbank, 2011; Wang et al., 2014). Several advantages of using a combination of C₄ pathways have been proposed. By dividing the transfer of metabolites between the mesophyll and bundle sheath into a combination of amino and organic acids, smaller gradients of each are sufficient to drive the transfer between cell types (Pick et al., 2011; Stitt and Zhu, 2014; Wang et al., 2014). The two pathways distribute energy and reducing equivalents differently between the two cell types which could be complementary and provide faster response to shifting light environments (Furbank, 2011; Stitt and Zhu, 2014). The ratio of NADP-ME to PEPCK pathway flux may not be fixed within species, across plant development, or under changing environment. Literature reports in maize indicate PEPCK flux is 10-25% of total C₄ flux (Chapman and Hatch, 1981; Weissmann et al., 2016; Arrivault et al., 2017), whereas in the C₄ dicot *Flaveria bidentis*, 50% of assimilated carbon comes through the PEPCK pathway (Meister et al., 1996). More broadly, enzyme activities for key C₄ enzymes in NADP-ME-type species also indicate extensive variation in the ratio of aspartate to malate translocated in C₄ plants (Kanai and Edwards, 1999). The ratio of malate and aspartate can also fluctuate in response to nitrogen limitation (Khamis et al., 1992). Though multiple studies have observed differences in pathway flux, the factors that elicit particular C₄ photosynthetic subtype use remain to be firmly established and could aid efforts to sustainably increase crop productivity.

The role for dual C₄ pathways has been considered over plant development in *Cleome gyandra* (Sommer et al., 2012). In this dicot species, decarboxylase activity varied with leaf age along the canopy using the NAD-malic enzyme pathway and a supplemental PEPCK pathway. In younger leaves, the activity of NAD-ME was twice that of PEPCK; however, in older leaves, PEPCK activity increased and NAD-ME activity decreased, leading to nearly double PEPCK activity compared to NAD-ME while the total amount of decarboxylase activity remained nearly constant. Though the study was intended to characterize the establishment of C₄

photosynthesis across development, the findings established within-plant differences and indicated the ratio of pathway use is pliable within species.

Canopy position and the resulting light environment of the leaf has an effect on the rate of photosynthesis. In the maize canopy, all new leaves form in full sunlight at the top of the canopy and become progressively shaded by new growth, requiring leaves to adapt to the new environment after they are fully formed. In the maize canopy, the gradients of light and age generate a pattern of increasing photosynthetic capacity at the top of the canopy, and less activity at the bottom of the canopy from older, self-shaded leaves (Ubierna et al., 2013; Niinemets, 2016; Pons, 2016; Collison et al., 2020). Up to 50% of maize photosynthesis occurs in these shaded parts of the canopy (Baker et al., 1988), and leaves in the lower canopy are capable of high rates of photosynthesis when not exposed to large degrees of self-shading (Collison et al., 2020). In addition to differences in overall rates of photosynthesis, foundational work on light regulation in maize has shown that the C_4 enzymes are more light-controlled than the C_3 enzymes (Sugiyama et al., 1984; Ward and Woolhouse, 1986). However, at low light, the C_3 pathways are expected to be more downregulated than C_4 as the plant shifts N away from Rubisco and toward light harvesting (Boardman, 1977; Björkman, 1981; Hikosaka and Terashima, 1995; Evans and Poorter, 2001; Walters, 2005; Tazoe et al., 2006; Pengelly et al., 2010). Moreover, species utilizing different C_4 subtypes have different degrees of shade acclimation, with C_4 grasses which use the NADP-ME subtype, such as *Z. mays*, able to acclimate to shade more readily than those using NAD-ME or PEP-CK subtypes (Sonawane et al., 2018).

The interaction between C_4 pathways and plant nitrogen status is well-documented (Khamis et al., 1992; Taub and Lerdau, 2000; Ghannoum et al., 2005; Pinto et al., 2014; Pinto et al., 2016) but mechanistically is not completely understood. The PEPCK pathway shuttles carbon using the amino acid aspartate which provides a potential regulatory link between C_4 photosynthesis and the nitrogen status of the plant. Aspartate can be rapidly converted to asparagine that is associated with important developmental cues during the maize life cycle (Seebauer et al., 2004) and transports carbon and nitrogen between leaves and developing kernels (Cañas et al., 2010). Asparagine is believed to play a role in signaling seed storage protein deposition (Hernández-Sebastià et al., 2005; Pandurangan et al., 2012) and, importantly, appears to be a regulator of PEPCK activity in seeds. When asparagine was added exogenously to grape seeds, PEPCK activity was increased 100-fold (Walker et al., 1997). This relationship is likely

important *in vivo* because PEPCK proteins are developmentally regulated in tomato and grape seeds (Walker et al., 1999; Bahrami et al., 2001), and PEPCK activity coincides with peak amino acid metabolism and storage protein deposition, possibly through asparaginase activity (Walker et al., 1999). Despite known anaplerotic functional relationships between asparagine and PEPCK, it remains unclear if asparagine could be important for the regulation of the C₄ PEPCK gene during development and aging of the maize leaf.

In this study, we grew maize plants at ten day intervals to sample leaves from the top and middle of the canopy at nine plant ages to determine the effect of canopy position and plant age on the proportion of C₄ flux through PEPCK relative to NADP-ME. Parts of the canopy with the greatest fluctuations in light environment, including lower leaves, would be expected to use proportionally more mixed C₄ pathways to allow leaf cells to maintain an energetic homeostasis. Additionally, plant age might affect the C₄ pathways such that older plants would use proportionately more equal C₄ pathways to maintain moderate levels of photosynthesis as leaf N is remobilized to the grain during senescence. One possible regulator of PEPCK variation based on work in other species is asparagine, so asparagine concentration across development was compared with C₄ pathway activity to identify potential regulatory relationships. Overall, we found the plant modifies the proportion of the two C₄ photosynthetic pathways in response to the developmental and environmental microenvironment of the leaf throughout the growing season.

Results

Distinguishing maize plant developmental progression from canopy influence

To understand variation in the C₄ pathways over the course of the plant life cycle and within the canopy, maize plants (*var.* W22) were grown in ten-day intervals for 100 days in the greenhouse from February to July 2019. Plants were sampled on July 2-3, 2019, for the nine developmental times, representing plants 15 days after sowing to 100 days after sowing (DAS; Figure 1). Measurements were taken from the top collared leaf ("top leaf") and leaf 13 ("subtending leaf"), which is the leaf subtending the ear in W22. All measurements were taken from the middle of the leaf, 20 cm from the leaf tip. The top leaf represents the leaf receiving the highest amount of sunlight and is the youngest expanded leaf on the maize plant. The subtending leaf provides the most nutrients to support ear growth (Subedi and Ma, 2005). Measurements were not taken from leaves within the whorl nor from leaves which had fully senesced. In total, these leaf

samples represented the span of the maize life cycle and two functionally distinct parts of the leaf canopy.

Photosynthetic CO₂ assimilation decreased with plant age

Overall photosynthetic rate has been shown to be variable during plant growth and in the canopy (Dwyer and Stewart, 1986; Chen et al., 2016; Niinemets, 2016), so we measured net CO₂ assimilation (A_{net}) on the plants at the nine growth stages. A_{net} was measured at both ambient (350 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and high (1500 $\mu\text{mol m}^{-2} \text{s}^{-1}$) light levels in the LI-6800 chamber. During plant growth A_{net} consistently decreased with age in the plant for both top and subtending leaves at the ambient and high light levels (Figure 2A). Leaf position did not have a significant effect on A_{net} ($p=0.90$, ANOVA type II). The interaction between light and plant age was significant ($p=5.96\text{e-}6$, ANOVA): in young plants, A_{net} increased substantially in the high light conditions compared to 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$; however, the response to high light was dampened in older plants.

Chlorophyll content and A_{net} are correlated with plant age

A_{net} was modestly correlated with chlorophyll concentration in the leaf (Pearson's $R = 0.39$) as an overall indicator of photosynthetic capacity. Total chlorophyll concentration increased in the top leaf from 0.6 mg chlorophyll g^{-1} FW to 1.1 mg g^{-1} FW between 15 DAS and 60 DAS before decreasing back to 0.6 mg g^{-1} FW at 80DAS, the last sampling time for the top leaf before complete senescence (Figure 2B). Leaf position in the canopy did not have a significant effect on total chlorophyll concentration ($p=0.12$, ANOVA type II); however, plant age had a significant effect ($p=0.004$). Chlorophyll concentration decreased steadily in the subtending leaf, likely as a consequence of the initiation of leaf senescence (Figure 2B). Throughout the growth cycle, the nitrogen status of the leaf and the light environment are expected to vary and can affect the chlorophyll a to b ratio (Hikosaka and Terashima, 1995). Chlorophyll a to b ratio was not correlated with the age of the plant or canopy position ($p=0.14$ and $p=0.73$, ANOVA type II); however, the interaction effect between leaf position and plant age was significant ($p=0.001$, ANOVA type II; Figure 2C), and the ratio drops slightly in the top leaves at 70 and 80 days after sowing, reflecting the beginning of senescence.

Ratio of C₄ decarboxylases changed with canopy position and plant age

The responses of the two C₄ pathway decarboxylases were investigated using transcriptional analysis since the enzymes involved in the C₄ pathway are at least partially regulated at this level (Pick et al., 2011). Using the gene expression atlas resource for maize (Stelpflug et al., 2016), gene expression levels were probed for the two decarboxylase enzymes in maize leaves over developmental age. Both PEPCK1 (GRMZM2G001696) and NADP-ME (NADP-ME3, GRMZM2G085019) expression varied with growth stage, beginning with the highest expression in the youngest leaves and decreasing with leaf age. NADP-ME3 expression decreased quickly after the V9 growth stage; its expression was approximately 2-fold higher than PEPCK1 until the V9 growth stage, where expression became approximately equal for the two decarboxylases (Figure 3A). Based on the transcriptomic evaluation, a comprehensive experiment was designed to assess the C₄ subtypes through the lifetime of leaves.

The expression of NADP-ME3 and PEPCK1 in the plants grown in the greenhouse was assessed by quantitative real-time PCR to confirm the trend indicated by the maize atlas. Similar to the pattern from the maize expression atlas, both C₄ decarboxylase genes decreased in relative expression as the leaf aged (Figure 3B). For both genes, expression was highest in the youngest leaf tissues and decreased continually in older leaves; the trend held for both the leaf at the top of the canopy and the leaf subtending the ear (Figure 3B). On average, NADP-ME3 was expressed at a level 2.4 times higher than PEPCK1. Unlike in the gene expression atlas, PEPCK1 expression was less divergent from NADP-ME3 expression in the youngest leaves, and more divergent in older leaves. Leaf position significantly impacted the expression of both NADP-ME3 and PEPCK1 expression, but not the ratio between the two genes.

Decarboxylase activity was also measured using enzyme activity assays. Maize plants exhibited variation in both the amount of enzyme activity and the ratio of NADP-ME to PEPCK activity depending on both canopy position of the leaf and growth stage of individual leaves (ANOVA activity ~ leaf * plant age, $p < 0.05$). Both enzyme activities in the top leaf were not significantly different with plant age. Canopy position had a large effect, with leaves in the top of the canopy exhibiting 35.7% more PEPCK and 62.2% more NADP-ME activity in the four growth stages where both leaves were measured (Figure 4). In the top of the canopy, NADP-ME activity was 2-fold higher than PEPCK on average; however, in the subtending leaves, the ratio decreased to 1.5-fold higher, indicating a decreased relative role for NADP-ME in that part of the canopy. Thus, in the top leaves, 66% of decarboxylase activity came from NADP-ME and 34% from

PEPCK, while in the subtending leaves, the proportion shifted to 60% from NADP-ME and 40% from PEPCK.

Pool sizes of C₄ and C₃ metabolites

Malate and aspartate are the transfer acids that move CO₂ from the mesophyll to the bundle sheath. Concentrations of both metabolites were measured in the two leaves for nine growth stages with LC-MS/MS. In the 15-day old plants, the ratio of malate to aspartate was 4.6:1, similar to reported values (Chapman and Hatch, 1981; Weissmann et al., 2016; Arrivault et al., 2017) and with expected pool sizes: 6.5±1.4 μmol malate g⁻¹ FW and 1.4±0.6 μmol aspartate g⁻¹ FW (Khamis et al., 1992; Lohaus et al., 1998; Szecowka et al., 2013; Arrivault et al., 2017). Both plant age and canopy position had significant effects on the pool sizes of malate and aspartate (ANOVA, type II, p<0.0001; Figure 5). In the top leaf, the pool size of malate decreased from 6.5±1.4 μmol g⁻¹ FW at 15 days to 2.2±0.8 μmol g⁻¹ FW for 30-80 DAS, while the aspartate pool size was largely unchanged with time. Thus, the ratio between the malate and aspartate pools decreased to 1.8±0.4 in the top leaf from 30 to 80 DAS (Figure 5). In the subtending leaf, the concentration of malate was generally greater than in the top leaf and both pool sizes decreased with plant age (Figure 5). In contrast, metabolites involved in the Calvin Benson Cycle were significantly affected by plant age, but only GAP/DHAP and E4P were significantly different between parts of the canopy, both having larger pools in the subtending leaf (Supplemental Figure 1).

One hypothesis for why plants might use the NADP-ME and PEPCK pathway with similar ratios in subtending leaves and older leaves is to maintain smaller pools of the five transfer acids, rather than large pools of malate and pyruvate needed for the NADP-ME pathway alone. Pyruvate, PEP, and aspartate all had strong age effects, decreasing with leaf age, while alanine increased in the top leaf with plant age, and the malate pool size did not change with plant age. Because the malate pool is mostly inactive (discussed below), its pool was excluded from a total transfer acid pool comparison. This combined pool was slightly larger in the top leaf and smaller in the bottom leaf (p=0.055, ANOVA type II), and decreased with leaf age (p=1.56e-5, ANOVA type II)

¹³CO₂ labeling in C₃ and C₄ metabolites

Differences in pool sizes for each growth stage may be the consequence of cumulative changes in metabolism over the life of the plant; however, isotopic labeling provides a snapshot of active metabolism. $^{13}\text{CO}_2$ was provided to the leaf of 15-day or 55-day old plants, and metabolites were measured for label incorporation and absolute pool size. These two ages were chosen to represent plants at the beginning of the vegetative and reproductive growth stages, as well as contrasting C_4 metabolite pool sizes. Plants were labeled for up to 5 minutes, and the average label incorporation in C_4 shuttle metabolites and several central carbon metabolism metabolites were quantified (Figure 6A, Supplemental Figure 1). Leaves from 15-day old plants incorporated ^{13}C label faster than the 55-day old plants, reaching an average labeling amount of 47-55% for measured Calvin Benson Cycle intermediates (i.e. phosphoglyceric acid (PGA), fructose biphosphate (FBP), glyceraldehyde 3-phosphate (GAP) and dihydroxyacetone phosphate (DHAP)) at 15 days compared to 12-27% for 55-day old plants in the top or subtending leaf, in agreement with the net CO_2 assimilation data (Figure 1). The C_4 intermediates were less enriched at both ages: 15-day old plants contained pyruvate that was 20% average labeled, 24.6% labeled phosphoenolpyruvate (PEP), 35% labeled aspartate, and 8.5% labeled malate. In the 55-day old plants, C_4 intermediates were approximately half the average label of the 15-day old plants after 5 minutes of exposure to $^{13}\text{CO}_2$ (Supplemental Figure 1).

Malate and aspartate predominantly incorporate label from $^{13}\text{CO}_2$ in the C-4 position, which is subsequently decarboxylated in the bundle sheath causing loss of the label (Chapman and Hatch, 1981). Label incorporation in the C1-C3 positions of malate and aspartate comes from downstream PEP and pyruvate labeling as a result of Calvin Benson Cycle activity. The malate pool also accumulates little ^{13}C label due to a large proportion of the malate pool not participating in photosynthetic metabolism, resulting in a large proportion considered an inactive pool during stable isotope labeling (Szecowka et al., 2013; Ma et al., 2014; Allen, 2016; Arrivault et al., 2017). The inactive pool in the 15-day and 55-day plants were similar, with approximately 80% of malate unlabeled after 5 minutes (Figure 6B) and only 10% average labeling (Supplemental Figure 1), though the final asymptotic value where M_0 levels off is incompletely defined at the five min time point (Arrivault et al., 2017; AuBuchon-Elder et al., 2020). Looking more deeply at the distribution of isotopologues, malate and aspartate labeling matched expectations based on the C_4 pathways (Figure 6B). The m+1 isotopologue for aspartate increased rapidly from 0 to 60 seconds to account for 50% of all aspartate in 15-day old plants and 20-30% in subtending and top leaves of 55-day old plants. Malate m+1 isotopologues

accounted for 5-10% of all malate. Very little (<5%) label accumulated in the C1-C3 positions of malate and aspartate during the first minute of labeling, also consistent with C₄ topology.

The metabolism of the NADP-ME and PEPCK pathways was calculated in a cross-comparable way in the three leaves (15-day top, 55-day top, 55-day subtending) using the method described in Arrivault et al (Arrivault et al., 2017) to calculate n-atom equivalents in the C-4 position of malate and aspartate. N-atom equivalent measurements for malate and aspartate use the number of labeled carbons, the metabolite concentration, and the assumption that all malate or aspartate molecules with at least one ¹³C contain label in the C-4 position to quantify labeling from a molar basis (see Supplemental Table 1). From this calculation, malate and aspartate labeling were directly compared. During the first 60 seconds of labeling, the C-4 position of malate labeled at twice the rate of aspartate in the 15-day old plants and 4.3 and 2.6 times faster in the 55-day old top leaf and subtending leaf, respectively (Figure 6B), reflecting 66.5% of C₄ activity going through the NADP-ME pathway and 33.5% through PEPCK and indicating a larger role for aspartate than previously identified, where aspartate only accounted for 10-25% of the activity of the C₄ shuttle (Chapman and Hatch, 1981; Weissmann et al., 2016; Arrivault et al., 2017). In the 55-day old plants, the rate of labeling in the C-4 position of malate in the subtending leaf was approximately one-half the top leaf, while aspartate labeling was consistent throughout the canopy, resulting in 18.8% of C₄ activity through PEPCK in the top leaf, and 27.9% in the subtending leaf (Figure 6C).

Asparagine concentrations in the plant

Asparagine is an important storage and transport form of nitrogen in maize plants (Lohaus et al., 1998; Lea and Azevedo, 2007) and has been shown to specifically induce PEPCK activity in grape seeds (Walker et al., 1999). Asparagine can be synthesized from aspartate via asparagine synthetase and converted back to aspartate through L-asparaginase; however, the relationship between the C₄ PEPCK and asparagine concentration is not known.

Asparagine concentration was measured using LC-MS/MS. The asparagine concentration in the leaf varied from 0.15 μmol mg⁻¹ to 6.6 μmol mg⁻¹ and did not have a clear pattern with leaf age or canopy position in this experiment (Figure 7). Concentrations were more variable than other metabolites (Supplemental Figure 1), resulting in a decreased ability to observe any trends in the data. As such, asparagine was very weakly correlated with PEPCK activity (Pearson's R = 0.17) and somewhat correlated with PEPCK expression (Pearson's R = 0.32) in the data.

PEPCK activity was more closely correlated with its substrate, aspartate ($R=0.74$), although it was not correlated with PEPCK expression ($R=0.07$). Aspartate and asparagine correlations were not correlated ($R = 0.01$). No correlation was observed between asparagine concentration and PEPCK expression or activity, predominantly because asparagine concentrations were highly variable within samples, but similar in all plant ages and both canopy positions. To fully elucidate this relationship in maize leaves, a more targeted approach would be necessary, focusing on asparagine concentrations in bundle sheath cells. Moreover, asparagine concentration fluctuates diurnally, with its highest concentrations at night (Harmer et al., 2018; Kambhampati et al., 2018), so the relationship between asparagine concentration and PEPCK activity in the leaf may also have a diurnal component which was not assessed in these experiments.

Discussion

The activities of C_4 subtype pathways vary during maize growth; with the greatest difference between leaves at the top and middle of the canopy, rather than with plant age. PEPCK activity was consistent between the top leaf and subtending leaf; however, NADP-ME activity decreased in the older subtending leaf compared to the top leaf, leading to a decreased overall rate of decarboxylation and changing the ratio of the decarboxylases between the two canopy positions. Similarly, the PEPCK pathway metabolite aspartate labeled at the same rate in the top and subtending leaf of 55-day old plants and correlated with PEPCK activity, while malate labeling decreased by three-fold in the subtending leaf compared to the top leaf. Decreased malate labeling in the subtending leaf was surprising considering the subtending leaves from all plant growth stages had larger malate pools than top leaves. These results indicate a large, inactive pool of malate in the subtending leaf, in addition to the photosynthetically active pool of malate in plant leaves (Szecowka et al., 2013; Ma et al., 2014; Weissmann et al., 2016; Arrivault et al., 2017). The large malate pool may serve to buffer the fluctuating light environment within the canopy or could function as a carbon reserve for remobilization to the grain.

Fluctuating environmental conditions, in particular with regard to light, are hypothesized to be a reason that plants might use a combination of C_4 pathways in parallel (Furbank, 2011; Bellasio and Griffiths, 2014a; Stitt and Zhu, 2014; Wang et al., 2014). The more stable, typically unshaded light environment at the top of the canopy may not require the same degree of metabolic flexibility afforded by parallel pathways and benefits from the efficiency of the NADP-ME pathway with CO_2 decarboxylation directly in the chloroplast (Wang et al., 2014). The subtending leaves receive less light in a closed canopy but can have high intensity sun flecks

under field conditions and thus may benefit through shared C_4 pathway operation (Bellasio and Griffiths, 2014a; Stitt and Zhu, 2014; Wang et al., 2014). Our experiments suggest the balanced activities occur though a reduction in NADP-ME activity and decreased total C_4 shuttling in lower parts of the canopy (Figure 4, Figure 6), which did not result in a lower carbon assimilation rate or reduced Calvin Benson Cycle metabolite labeling. Carbon assimilation rates were however significantly affected by the age of the plant (Figure 2, Supplemental Figure 1). Subtending leaves maintained the same level of CBC metabolism as leaves at the top of the canopy, likely as a result of even light distribution through the canopy due to growth in a greenhouse rather than a dense field canopy (Collison et al., 2020).

This study was designed to address differences in C_4 metabolism based on leaf age and canopy position, such that the same two leaves were compared between plants of different ages. There were few differences in C_4 metabolism between plants of different ages (i.e. top leaf on plants), but significant differences in C_4 pathways between leaves in different positions on the same plant (i.e. microenvironment). These results bring up important possibilities of regulation by light that differ between leaves on the same plant. Light may play a crucial role in regulating the C_4 decarboxylases between canopy positions. The relationship between light and C_4 pathway activity has been studied in the context of different species with different C_4 subtypes (Ubierna et al., 2013; Sonawane et al., 2018). Less consideration has been given to the plasticity of the C_4 pathways within a species. NADP-ME is strongly light regulated by light quality (Casati et al., 1998) and quantity (Hatch and Kagawa, 1976; Bellasio and Griffiths, 2014b), and by time of day, through phosphorylation, with its activity peaking at two hours after dawn and decreasing slowly throughout the day (Bovdilova et al., 2019). Leaves positioned lower in the canopy may not receive the quality and quantity of light to activate NADP-ME to the levels observed at the top of the canopy (Figure 4, Figure 6). In contrast, PEPCK activity may be unresponsive to light (Wingler et al., 1999) or potentially negatively regulated by light (Chao et al., 2014); though it should be noted that the experiments reported here were performed in the greenhouse where plant leaves are not shaded to the extent they would be in the field. PEPCK expression levels are more clearly responsive to other environmental conditions such as N supply which could have effects in leaves subtending the ear (Delgado-Alvarado et al., 2007; Penfield et al., 2012). N distribution in the canopy also regulates the level of photosynthesis in herbaceous canopies (Niinemets, 2016).

Conclusions

The top and middle of the canopy and different ages of the plant experience changes in C_3 and C_4 cycle metabolism. Variation in the relative proportion of the two C_4 pathways was strongest between leaves at the top and bottom of the canopy, while differences in overall photosynthetic rate were predominantly caused by the age of the plant. C_4 characteristics changed as the leaf aged, although they did not shift relative to each other, instead generally decreasing overall to coordinate with decreased C_3 metabolism. At the top of the canopy, the plant predominantly uses NADP-ME and shifts to more equal use of the two C_4 pathways lower in the canopy, as the microenvironment in the lower canopy is more variable and the leaves shift to supporting growth of sink tissues throughout the plant. Plasticity within the plant life cycle and within the canopy is a potential advantage of the parallel C_4 pathways, allowing fine-tuning of the C_4 pathways to optimize for specific growth conditions of the leaf.

Methods

Plant Growth

Plants were grown in the greenhouse in Saint Louis, MO from February to July 2019. Four maize plants (genotype W22) were planted at ten day intervals for 90 days. Plants were self-pollinated upon flowering. Plants were grown in Berger 35, 7% bark medium in 2.5 gallon pots. Plants were grown with 14-hour day length, 10 hour nights using supplemental light to extend the day length and when available sunlight was below $600 \mu\text{mol m}^{-2} \text{s}^{-1}$. Plant growth temperature was 28°C day, 22°C night and a minimum 40% relative humidity.

Plants were sampled at 100 days after initiation of the experiment. Plants were 10, 20, 30, 40, 50, 60, 70, 80, or 90 days old at the time of sampling. Leaf tissue was collected from the topmost leaf and the leaf subtending the ear (leaf 13 in W22), when available. For the two oldest sets of plants (80 or 90 days after sowing), the topmost leaf on each plant was fully senesced and not sampled. For the three youngest sets of plants (10, 20, or 30 days after sowing), the subtending leaf had not emerged from the whorl and was not sampled. All measurements were taken from 10cm leaf segments beginning 20cm from the tip of the leaf.

Gas Exchange Measurements

Net CO₂ assimilation was measured using gas exchange measurements on a LI-6800 Photosynthesis System (Li-Cor Inc., Lincoln, Nebraska). Net CO₂ assimilation was measured at steady state at 400ppm CO₂ and 350 followed by $1500 \mu\text{mol m}^{-2} \text{s}^{-1}$ light. The leaf was clamped into the instrument head using a 6cm aperture, temperature was controlled at 25°C, and the gas flow rate was $500 \mu\text{mol s}^{-1}$.

Gene Expression of PEPCK and NADP-ME

Leaf tissue was sampled from a 10cm segment starting 20cm from the leaf tip, from either the leaf at the top of the canopy or from leaf number 13, which is the leaf below the ear in W22 (subtending leaf). RNA was extracted from 50mg of leaf tissue using Trizol following the manufacturer instructions (Invitrogen 15596026). Residual genomic DNA was removed using DNaseI (Invitrogen TURBO DNA-free, AM1907), and first strand cDNA synthesis was performed using Invitrogen Superscript II (18064022). Gene expression for PEPCK1 (Zm00004b001002) and NADP-ME3 (Zm00004b015828) were quantified via qRT-PCR using a Roche Light Cycler 480 II using Roche SYBR green I (Roche 04707516001). The delta cT method was used to quantify relative gene expression of PEPCK1 and NADP-ME3 over time.

NAC26 (Zm00004b022707) was used as a housekeeping gene using published primers (CCGCCGTCAACAGGGGAAATCTG, GTAGCACGCCCAAGACCAACAG; (Lin et al., 2014) Genome-wide identification of housekeeping genes in maize). Primers for PEPCK1 were forward: CCCGATCAACACCTGGACG and reverse: GACGCACCCATGACAATACC; primers for NADP-ME3 were forward: GAGTCAGGGCCGTTCAATCT and reverse: ACAGAGTACCATCCGCGTTG.

Enzyme Activity

PEPCK activity was measured using the method of Walker et al (Walker et al., 1999) detailed on protocols.org (Osorio et al., 2014). Briefly, crude protein was extracted from ~100mg of fresh leaf tissue in a buffer containing 0.5M bicine-KOH (pH 9.0), 0.2M KCl, 3mM EDTA, 5% (w/v) PEG-4000, 25mM DTT and 0.4% bovine serum albumin. The extract was centrifuged for 20 minutes and the supernatant was added to a buffer containing 0.5M bicine-KOH (pH 9.0), 3mM EDTA, 55% w/v PEG-4000, and 25mM DTT. The sample was incubated for 10 minutes on ice, centrifuged at 13,000 x g at 4°C for 20 minutes. The supernatant was discarded and the pellet was resuspended in 10mM bicine-KOH (pH 9.0) with 25mM DTT. Activity was measured in the carboxylation direction by coupling the reaction with malate dehydrogenase and following the oxidation of NADH at 340nm using Molecular Devices SpectraMax M2 spectrophotometer. Total protein was measured using the protein extract for PEPCK activity using Bradford reagent (Millipore Sigma; Cat: B6916) and commercial bovine serum albumin standards (Thermo Fisher 23208). NADP-malic enzyme activity was measured using the method described in Osorio et al (Osorio et al., 2014) from Detarsio et al (Detarsio et al., 2003), using 50mg of fresh leaf tissue. Enzyme activity was measured by following the reduction of NADP⁺ at 340nm for 5 minutes.

Compositional Analysis

Chlorophyll content was measured using the method of Arnon et al (Arnon, 1949). Amino acid, sugar and sugar phosphate content was measured using the method described in Czajka et al, (Czajka et al., 2020). Briefly, metabolites were extracted from 100mg fresh weight of ground leaf tissue using 3:7 (v/v) methanol:chloroform solution incubated for two hours on a rotator at 4°C. After two hours, 0.5 mL of ddH₂O was added, the solution was centrifuged and the supernatant (aqueous phase) was centrifuged in 3KDa filters, frozen, lyophilized, and resuspended in 50uL 1:1 methanol:ddH₂O. Metabolite quantities were measured using LC-MS using a Shimadzu Prominence-xR UFLC system and a SCIEX hybrid triple quadrupole-linear ion trap MS equipped with Turbo V^{TM} electrospray ionization source for separation and detection of

metabolites, and samples were injected into an InfinityLab Poroshell 120 HILIC-Z (2.1 x 100 mm, 2.7 µm, Agilent Technologies) column.

Stable isotope labeling using $^{13}\text{CO}_2$

Isotopic labeling was performed on plants from two of the stages, 15 days and 55 days after sowing. For 15-day old plants, the top collared leaf was sampled, and for 55-day old plants, both the top collared leaf and the subtending leaf were used. The 15-day old plants were selected to represent the growth stage commonly characterized for C_4 pathway labeling (Chapman and Hatch, 1981; Weissmann et al., 2016; Arrivault et al., 2017), where young maize leaves use the NADP-ME pathway for 75-90% of metabolism, relying on the PEPCK pathway as a minor contributor. The 55-day sample was selected to represent plants that have generated all of their leaves, reached the end of vegetative growth, and are approaching the developmental shift at anthesis. Labeling in central carbon metabolites was measured using $^{13}\text{CO}_2$ provided to the leaf with a hand-held clamp for 0, 20, 60, or 300 seconds to generate a time course of label incorporation. Leaves were freeze clamped at the end of each time interval, flash frozen in liquid nitrogen, and stored at -80°C until further processing. Metabolites were extracted in chloroform:methanol as above, and metabolite labeling and pool size were measured by LC-MS/MS on a Qtrap6500 (Chu et al, in prep; Czajka et al., 2020)). Samples were corrected for natural abundance of ^{13}C using IsoCorrector (Heinrich et al., 2018).

Statistical Analysis

ANOVA comparisons were performed using two-way ANOVAs with factors canopy position (leaf) and plant age, with the Anova function in the car package in R (Fox and Weisberg, 2018) to use a type II ANOVA to be able to compare factors with unequal sample sizes.

Acknowledgements

The authors would like to thank the Donald Danforth Plant Science Center Plant Growth Facility and Donald Danforth Plant Science Center Proteomics and Mass Spectrometry Facility. Thank you to Emma Smith, Emily Frankenreiter, and Genesis Hudson for assistance with performing tissue sampling.

Figures

Figure 1: Experimental Design: Plants were grown at ten day intervals in the greenhouse to generate a gradient of maize development which could be sampled on the same day. Nine individual time points were used and captured maize development from the V3 stage until physiological maturity. The leaf at the top of the canopy for each time point was sampled until the top leaf senesced at 90 days after sowing. Also sampled was leaf 13, which subtends the ear in genotype W22. Leaf 13 emerged from the whorl at 50 days after sowing.

Figure 2: Measures of overall photosynthesis: A.) Net CO₂ assimilation (A) was measured at two light levels using a LI-COR 6800 gas exchange instrument. B.) Total chlorophyll was measured using a spectrometric assay. C.) Chlorophyll a to b ratio. All data are means $\pm$ sd of 3-4 replicate plants.

Figure 3: Gene expression differences: A.) Expression of NADP-ME3 (Zm00004b015828) and PEPCK1 (Zm00004b001002) in the maize gene expression atlas (Stelpflug et al, 2016) decreased with plant age, and the ratio of NADP-ME3 (gray) to PEPCK1 (blue) decreased in older plants. B.) Relative gene expression for PEPCK1 and NADP-ME3 was quantified using qPCR. Data are means $\pm$ sd of 3-4 replicate plants.

Figure 4: Enzyme activity assays: *In vitro* activities of NADP-ME (gray) and PEPCK (blue) from whole leaf segments of maize leaves at nine growth stages from the top or subtending leaf. Data are means $\pm$ sd of 3-4 replicate plants.

Figure 5: C₄ transfer acid quantification: Malate and aspartate pools were quantified using an LC-MS approach and normalized by sample fresh weight. Malate pools (black) were larger than aspartate pools (blue) for all measured time points. Data are means $\pm$ sd of 3-4 replicate plants.

Figure 6: C₄ transfer acid labeling: A.) Quantitation of malate and aspartate pools using LC-MS. Malate (gray); aspartate (blue). B.) Time course of labeling for malate and aspartate. Mass isotopologs (m_n) represent malate or aspartate molecules that have incorporated n molecules of ¹³C. C.) Moles of ¹³C in the C-4 position of malate (gray) and aspartate (blue) using the method of (Arrivault et al., 2017).

Figure 7: Asparagine: asparagine concentration during plant growth measured by LC-MS normalized by sample fresh weight. Data are means $\pm$ sd of 3-4 replicate plants.

Supplemental Data

Supplemental Table 1: Calculations to determine degree of labeling in the C-4 position of malate and aspartate.

Supplemental Figure 1: A.) Metabolite quantitation of C₄ intermediates. PEP: phospho*eno*/pyruvate. B.) Metabolite quantitation of Calvin Benson Cycle intermediates which were able to be quantified. TP: Triose phosphate (i.e. GAP and DHAP: Glyceraldehyde 3-phosphate and Dihydroxyacetone phosphate), FBP: Fructose 1,6-bisphosphate; E4P: Erythrose 4-phosphate; S7P: Sedoheptulose 7-phosphate; R5P: Ribose 5-phosphate.

Supplemental Figure 2: Average labeling in C₃ and C₄ metabolites during the five minute time course. PEP: phospho*eno*/pyruvate, PGA: phosphoglyceric acid, FBP: fructose 1,6-bisphosphate, TP: triose phosphate (i.e. GAP and DHAP: glyceraldehyde 3-phosphate and

553 dihydroxyacetone phosphate), UDPG: uridine diphosphate glucose, 2OG: 2-oxoglutarate, ASN:
 554 asparagine, GLN: glutamine, Glu: glutamate.
 555
 556 Supplemental Figure 3: Isotopologue distribution graphs for labeled metabolites during the five
 557 minute time course. PEP: phospho*eno*lpyruvate, PGA: phosphoglyceric acid, FBP: fructose 1,6-
 558 biphosphate, TP: triose phosphate (i.e. GAP and DHAP: glyceraldehyde 3-phosphate and
 559 dihydroxyacetone phosphate), UDPG: uridine diphosphate glucose, 2OG: 2-oxoglutarate, ASN:
 560 asparagine, GLN: glutamine, Glu: glutamate.

References

- Allen DK (2016) Quantifying plant phenotypes with isotopic labeling & metabolic flux analysis. *Curr Opin Biotechnol* 37: 45–52
- Arnon DI (1949) COPPER ENZYMES IN ISOLATED CHLOROPLASTS. POLYPHENOLOXIDASE IN BETA VULGARIS. *Plant Physiol* 24: 1–15
- Arrivault S, Obata T, Szecówka M, Mengin V, Guenther M, Hoehne M, Fernie AR, Stitt M (2017) Metabolite pools and carbon flow during C4 photosynthesis in maize: 13CO2 labeling kinetics and cell type fractionation. *J Exp Bot* 68: 283–298
- AuBuchon-Elder T, Coneva V, Goad DM, Jenkins LM, Yu Y, Allen DK, Kellogg EA (2020) Sterile Spikelets Contribute to Yield in Sorghum and Related Grasses. *Plant Cell* 32: 3500–3518
- Bahrami AR, Chen ZH, Walker RP, Leegood RC, Gray JE (2001) Ripening-related occurrence of phosphoenolpyruvate carboxykinase in tomato fruit. *Plant Mol Biol* 47: 499–506
- Baker NR, Long SP, Ort DR (1988) Photosynthesis and temperature, with particular reference to effects on quantum yield. *Symp Soc Exp Biol* 42: 347–375
- Bellasio C, Griffiths H (2014a) The Operation of Two Decarboxylases, Transamination, and Partitioning of C4 Metabolic Processes between Mesophyll and Bundle Sheath Cells Allows Light Capture To Be Balanced for the Maize C4 Pathway. *Plant Physiol* 164: 466–480
- Bellasio C, Griffiths H (2014b) Acclimation of C4 metabolism to low light in mature maize leaves could limit energetic losses during progressive shading in a crop canopy. *J Exp Bot* 65: 3725–3736
- Björkman O (1981) Responses to different quantum flux densities. *Physiological plant ecology I*
- Boardman NK (1977) Comparative Photosynthesis of Sun and Shade Plants. *Annu Rev Plant Physiol* 28: 355–377
- Bovdilova A, Alexandre BM, Höppner A, Luís IM, Alvarez CE, Bickel D, Gohlke H, Decker C, Nagel-Steger L, Alseekh S, et al (2019) Posttranslational Modification of the NADP-Malic Enzyme Involved in C4 Photosynthesis Modulates the Enzymatic Activity during the Day. *Plant Cell* 31: 2525–2539
- Cañas RA, Quilleré I, Lea PJ, Hirel B (2010) Analysis of amino acid metabolism in the ear of maize mutants deficient in two cytosolic glutamine synthetase isoenzymes highlights the importance of asparagine for nitrogen translocation within sink organs. *Plant Biotechnol J* 8: 966–978
- Casati P, Drincovich MF, Andreo CS, Donahue R, Edwards GE (1998) UV-B, red and far-red light regulate induction of the C4 isoform of NADP-malic enzyme in etiolated maize seedlings. *Funct Plant Biol* 25: 701–708
- Chao Q, Liu X-Y, Mei Y-C, Gao Z-F, Chen Y-B, Qian C-R, Hao Y-B, Wang B-C (2014) Light-regulated phosphorylation of maize phosphoenolpyruvate carboxykinase plays a vital role in its activity. *Plant Mol Biol* 85: 95–105
- Chapman KSR, Hatch MD (1981) Aspartate Decarboxylation in Bundle Sheath Cells of Zea

599 mays and Its Possible Contribution to C3 Photosynthesis. *Funct Plant Biol* 8: 237–248

600 Chen Y, Wu D, Mu X, Xiao C, Chen F, Yuan L, Mi G (2016) Vertical Distribution of
601 Photosynthetic Nitrogen Use Efficiency and Its Response to Nitrogen in Field-Grown Maize.
602 *Crop Sci* 56: 397–407

603 Collison RF, Raven EC, Pignon CP, Long SP (2020) Light, Not Age, Underlies the
604 Maladaptation of Maize and *Miscanthus* Photosynthesis to Self-Shading. *Front Plant Sci*
605 11: 783

606 Czajka JJ, Kambhampati S, Tang YJ, Wang Y, Allen DK (2020) Application of Stable Isotope
607 Tracing to Elucidate Metabolic Dynamics During *Yarrowia lipolytica* α -Ionone Fermentation.
608 *iScience* 23: 100854

609 Delgado-Alvarado A, Walker RP, Leegood RC (2007) Phosphoenolpyruvate carboxykinase in
610 developing pea seeds is associated with tissues involved in solute transport and is
611 nitrogen-responsive. *Plant Cell Environ* 30: 225–235

612 Detarsio E, Wheeler MCG, Bermúdez VAC, Andreo CS, Drincovich MF (2003) Maize C4 NADP-
613 malic enzyme: Expression in *Escherichia coli* and characterization of site-directed mutants
614 at the putative nucleotide-binding sites. *J Biol Chem* 278: 13757–13764

615 Dwyer LM, Stewart DW (1986) Effect of leaf age and position on net photosynthetic rates in
616 maize (*Zea mays* L.). *Agric For Meteorol* 37: 29–46

617 Evans JR, Poorter H (2001) Photosynthetic acclimation of plants to growth irradiance: the
618 relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain.
619 *Plant Cell Environ* 24: 755–767

620 Fox J, Weisberg S (2018) *An R Companion to Applied Regression*. SAGE Publications

621 Furbank RT (2011) Evolution of the C(4) photosynthetic mechanism: are there really three C(4)
622 acid decarboxylation types? *J Exp Bot* 62: 3103–3108

623 Ghannoum O, Evans JR, Chow WS, Andrews TJ, Conroy JP, von Caemmerer S (2005) Faster
624 Rubisco is the key to superior nitrogen-use efficiency in NADP-malic enzyme relative to
625 NAD-malic enzyme C4 grasses. *Plant Physiol* 137: 638–650

626 Harmer SL, Covington MF, Bläsing O, Stitt M (2018) Circadian regulation of global gene
627 expression and metabolism. *Annual Plant Reviews online* 133–165

628 Hatch MD, Kagawa T (1976) Photosynthetic activities of isolated bundle sheath cells in relation
629 to differing mechanisms of C-4 pathway photosynthesis. *Arch Biochem Biophys* 175: 39–53

630 Heinrich P, Kohler C, Ellmann L, Kuerner P, Spang R, Oefner PJ, Dettmer K (2018) Correcting
631 for natural isotope abundance and tracer impurity in MS-, MS/MS- and high-resolution-
632 multiple-tracer-data from stable isotope labeling experiments with IsoCorrectoR. *Sci Rep* 8:
633 17910

634 Hernández-Sebastià C, Marsolais F, Saravitz C, Israel D, Dewey RE, Huber SC (2005) Free
635 amino acid profiles suggest a possible role for asparagine in the control of storage-product
636 accumulation in developing seeds of low-and high-protein soybean lines. *J Exp Bot* 56:
637 1951–1963

638 Hikosaka K, Terashima I (1995) A model of the acclimation of photosynthesis in the leaves of
639 C3 plants to sun and shade with respect to nitrogen use. *Plant Cell Environ* 18: 605–618

640 Kambhampati S, Ajewole E, Marsolais F (2018) Advances in Asparagine Metabolism. *In* FM
641 Cánovas, U Lüttge, R Matyssek, eds, *Progress in Botany* Vol. 79. Springer International
642 Publishing, Cham, pp 49–74

643 Kanai R, Edwards GE (1999) The biochemistry of C4 photosynthesis. *C4 plant biology* 49: 87

644 Khamis S, Lamaze T, Farineau J (1992) Effect of nitrate limitation on the photosynthetically
645 active pools of aspartate and malate in maize, a NADP malic enzyme C 4 plant. *Physiol*
646 *Plant* 85: 223–229

647 Lea PJ, Azevedo RA (2007) Nitrogen use efficiency. 2. Amino acid metabolism. *Ann Appl Biol*
648 151: 269–275

649 Lin F, Jiang L, Liu Y, Lv Y, Dai H, Zhao H (2014) Genome-wide identification of housekeeping
650 genes in maize. *Plant Mol Biol* 86: 543–554

651 Lohaus G, Büker M, Hußmann M, Soave C, Heldt H-W (1998) Transport of amino acids with
652 special emphasis on the synthesis and transport of asparagine in the Illinois Low Protein
653 and Illinois High Protein strains of maize. *Planta* 205: 181–188

654 Ma F, Jazmin LJ, Young JD, Allen DK (2014) Isotopically nonstationary ¹³C flux analysis of
655 changes in *Arabidopsis thaliana* leaf metabolism due to high light acclimation. *Proceedings*
656 *of the National Academy of Sciences* 111: 16967–16972

657 Meister M, Agostino A, Hatch MD (1996) The roles of malate and aspartate in C4 photosynthetic
658 metabolism of *Flaveria bidentis* (L.). *Planta* 199: 262–269

659 Niinemets Ü (2016) Leaf age dependent changes in within-canopy variation in leaf functional
660 traits: a meta-analysis. *J Plant Res* 129: 313–338

661 Osorio S, Vallarino JG, Szecowka M, Ufaz S, Tzin V, Angelovici R, Galili G, Fernie AR (2014)
662 Extraction and Measurement the Activities of Cytosolic Phosphoenolpyruvate
663 Carboxykinase (PEPCK) and Plastidic NADP-dependent Malic Enzyme (ME) on Tomato
664 (*Solanum lycopersicum*). *Bio-protocol* 4: e1122–e1122

665 Pandurangan S, Pajak A, Molnar SJ, Cober ER, Dhaubhadel S, Hernández-Sebastià C, Kaiser
666 WM, Nelson RL, Huber SC, Marsolais F (2012) Relationship between asparagine
667 metabolism and protein concentration in soybean seed. *J Exp Bot* 63: 3173–3184

668 Penfield S, Clements S, Bailey KJ, Gilday AD, Leegood RC, Gray JE, Graham IA (2012)
669 Expression and manipulation of phosphoenolpyruvate carboxykinase 1 identifies a role for
670 malate metabolism in stomatal closure. *Plant J* 69: 679–688

671 Pengelly JJJ, Sirault XRR, Tazoe Y, Evans JR, Furbank RT, von Caemmerer S (2010) Growth
672 of the C4 dicot *Flaveria bidentis*: photosynthetic acclimation to low light through shifts in leaf
673 anatomy and biochemistry. *J Exp Bot* 61: 4109–4122

674 Pick TR, Bräutigam A, Schlüter U, Denton AK, Colmsee C, Scholz U, Fahnenstich H,
675 Pieruschka R, Rascher U, Sonnewald U, et al (2011) Systems analysis of a maize leaf
676 developmental gradient redefines the current C4 model and provides candidates for

677 regulation. *Plant Cell* 23: 4208–4220

678 Pinto H, Powell JR, Sharwood RE, Tissue DT, Ghannoum O (2016) Variations in nitrogen use
679 efficiency reflect the biochemical subtype while variations in water use efficiency reflect the
680 evolutionary lineage of C4 grasses at inter-glacial CO₂. *Plant Cell Environ* 39: 514–526

681 Pinto H, Sharwood RE, Tissue DT, Ghannoum O (2014) Photosynthesis of C3, C3–C4, and C4
682 grasses at glacial CO₂. *J Exp Bot* 65: 3669–3681

683 Pons TL (2016) Regulation of Leaf Traits in Canopy Gradients. *In* K Hikosaka, Ü Niinemets,
684 NPR Anten, eds, *Canopy Photosynthesis: From Basics to Applications*. Springer
685 Netherlands, Dordrecht, pp 143–168

686 Sage RF, Christin P-A, Edwards EJ (2011) The C(4) plant lineages of planet Earth. *J Exp Bot*
687 62: 3155–3169

688 Seebauer JR, Moose SP, Fabbri BJ, Crossland LD, Below FE (2004) Amino acid metabolism in
689 maize earshoots. Implications for assimilate preconditioning and nitrogen signaling. *Plant*
690 *Physiol* 136: 4326–4334

691 Sommer M, Bräutigam A, Weber APM (2012) The dicotyledonous NAD malic enzyme C4 plant
692 *Cleome gynandra* displays age-dependent plasticity of C4 decarboxylation biochemistry.
693 *Plant Biol* 14: 621–629

694 Sonawane BV, Sharwood RE, Whitney S, Ghannoum O (2018) Shade compromises the
695 photosynthetic efficiency of NADP-ME less than that of PEP-CK and NAD-ME C4 grasses.
696 *J Exp Bot* 69: 3053–3068

697 Stelpflug SC, Sekhon RS, Vaillancourt B, Hirsch CN, Buell CR, de Leon N, Kaeppler SM (2016)
698 An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to
699 Explore Root Development. *Plant Genome*. doi: 10.3835/plantgenome2015.04.0025

700 Stitt M, Zhu X-G (2014) The large pools of metabolites involved in intercellular metabolite
701 shuttles in C4 photosynthesis provide enormous flexibility and robustness in a fluctuating
702 light environment. *Plant Cell Environ* 37: 1985–1988

703 Subedi KD, Ma BL (2005) Ear Position, Leaf Area, and Contribution of Individual Leaves to
704 Grain Yield in Conventional and Leafy Maize Hybrids. *Crop Sci* 45: 2246–2257

705 Sugiyama T, Mizuno M, Hayashi M (1984) Partitioning of nitrogen among ribulose-1, 5-
706 biphosphate carboxylase/oxygenase, phosphoenolpyruvate carboxylase, and pyruvate
707 orthophosphate dikinase as related to biomass productivity in maize seedlings. *Plant*
708 *Physiol* 75: 665–669

709 Szecowka M, Heise R, Tohge T, Nunes-Nesi A, Vosloh D, Huege J, Feil R, Lunn J, Nikoloski Z,
710 Stitt M, et al (2013) Metabolic fluxes in an illuminated *Arabidopsis* rosette. *Plant Cell* 25:
711 694–714

712 Taub DR, Lerdau MT (2000) Relationship between leaf nitrogen and photosynthetic rate for
713 three NAD-ME and three NADP-ME C4 grasses. *Am J Bot* 87: 412–417

714 Tazoe Y, Noguchi K, Terashima I (2006) Effects of growth light and nitrogen nutrition on the
715 organization of the photosynthetic apparatus in leaves of a C4 plant, *Amaranthus cruentus*.

716 Plant Cell Environ 29: 691–700

717 Ubierna N, Sun W, Kramer DM, Cousins AB (2013) The efficiency of C4 photosynthesis under
718 low light conditions in *Zea mays*, *Miscanthus x giganteus* and *Flaveria bidentis*. Plant Cell
719 Environ 36: 365–381

720 Walker RP, Acheson RM, Técsi LI, Leegood RC (1997) Phosphoenolpyruvate Carboxykinase in
721 C4 Plants: Its Role and Regulation. Funct Plant Biol 24: 459–468

722 Walker RP, Chen ZH, Tecsli LI, Famiani F, Lea PJ, Leegood RC (1999) Phosphoenolpyruvate
723 carboxykinase plays a role in interactions of carbon and nitrogen metabolism during grape
724 seed development. Planta 210: 9–18

725 Walters RG (2005) Towards an understanding of photosynthetic acclimation. J Exp Bot 56:
726 435–447

727 Wang Y, Bräutigam A, Weber APM, Zhu X-G (2014) Three distinct biochemical subtypes of C4
728 photosynthesis? A modelling analysis. J Exp Bot 65: 3567–3578

729 Ward DA, Woolhouse HW (1986) Comparative effects of light during growth on the
730 photosynthetic properties of NADP-ME type C4 grasses from open and shaded habitats. II.
731 Photosynthetic enzyme activities and metabolism. Plant Cell Environ 9: 271–277

732 Weissmann S, Ma F, Furuyama K, Gierse J, Berg H, Shao Y, Taniguchi M, Allen DK, Brutnell
733 TP (2016) Interactions of C4 Subtype Metabolic Activities and Transport in Maize Are
734 Revealed through the Characterization of DCT2 Mutants. Plant Cell 28: 466–484

735 Wingler A, Walker RP, Chen ZH, Leegood RC (1999) Phosphoenolpyruvate carboxykinase is
736 involved in the decarboxylation of aspartate in the bundle sheath of maize. Plant Physiol
737 120: 539–546

Figure 1: **Experimental Design:** Plants were grown at ten day intervals in the greenhouse to generate a gradient of maize development which could be sampled on the same day. Nine individual time points were used and captured maize development from the V3 stage until physiological maturity. The leaf at the top of the canopy for each time point was sampled until the top leaf senesced at 90 days after sowing. Also sampled was leaf 13, which subtends the ear in genotype W22. Leaf 13 emerged from the whorl at 50 days after sowing.

Figure 2: Measures of overall photosynthesis: A.) Net CO₂ assimilation (A) was measured at two light levels using a LI-COR 6800 gas exchange instrument. B.) Total chlorophyll was measured using a spectrometric assay. C.) Chlorophyll a to b ratio. All data are means $\pm$ standard deviation of 3-4 replicate plants.

Figure 3: Gene expression differences: A.) Expression of NADP-ME3 (Zm00004b015828) and PEPCK1 (Zm00004b001002) in the maize gene expression atlas (Stelpflug et al, 2016) decreased with plant age, and the ratio of NADP-ME3 (gray) to PEPCK1 (blue) decreased in older plants. B.) Relative gene expression for PEPCK1 and NADP-ME3 was quantified using qPCR. Data are means $\pm$ sd of 3-4 replicate plants.

Figure 4: **Enzyme activity assays:** *In vitro* activities of NADP-ME (gray) and PEPCK (blue) from whole leaf segments of maize leaves at nine growth stages from the top or subtending leaf. Data are means $\pm$ sd of 3-4 replicate plants. U: μ mol NADH oxidation (PEPCK) or NADP⁺ reduction (NADP-ME) per minute.

Figure 5: **C₄ transfer acid quantification:** Malate and aspartate pools were quantified using an LC-MS approach and normalized by sample fresh weight. Malate pools (black) were larger than aspartate pools (blue) for all measured time points. Data are means $\pm$ sd of 3-4 replicate plants.

Figure 6: **C₄ transfer acid labeling**: A.) Quantitation of malate and aspartate pools using LC-MS. Malate (gray); aspartate (blue). B.) Time course of labeling for malate and aspartate. Mass isotopologs (m_n) represent malate or aspartate molecules that have incorporated n molecules of ^{13}C . C.) Moles of ^{13}C in the C-4 position of malate (gray) and aspartate (blue) using the method of Arrivault et al, ([Arrivault et al., 2017](https://doi.org/10.1002/ps.5000)).

Figure 7: Asparagine concentration during plant growth measured by LC-MS normalized by sample fresh weight. Data are means $\pm$ sd of 3-4 replicate plants.

Parsed Citations

Allen DK (2016) Quantifying plant phenotypes with isotopic labeling & metabolic flux analysis. *Curr Opin Biotechnol* 37: 45–52

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Arnon DI (1949) COPPER ENZYMES IN ISOLATED CHLOROPLASTS. POLYPHENOLOXIDASE IN BETA VULGARIS. *Plant Physiol* 24: 1–15

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Arrivault S, Obata T, Szczówka M, Mengin V, Guenther M, Hoehne M, Fernie AR, Stitt M (2017) Metabolite pools and carbon flow during C4 photosynthesis in maize: 13CO2 labeling kinetics and cell type fractionation. *J Exp Bot* 68: 283–298

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

AuBuchon-Elder T, Coneva V, Goad DM, Jenkins LM, Yu Y, Allen DK, Kellogg EA (2020) Sterile Spikelets Contribute to Yield in Sorghum and Related Grasses. *Plant Cell* 32: 3500–3518

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bahrani AR, Chen ZH, Walker RP, Leegood RC, Gray JE (2001) Ripening-related occurrence of phosphoenolpyruvate carboxykinase in tomato fruit. *Plant Mol Biol* 47: 499–506

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Baker NR, Long SP, Ort DR (1988) Photosynthesis and temperature, with particular reference to effects on quantum yield. *Symp Soc Exp Biol* 42: 347–375

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bellasio C, Griffiths H (2014a) The Operation of Two Decarboxylases, Transamination, and Partitioning of C4 Metabolic Processes between Mesophyll and Bundle Sheath Cells Allows Light Capture To Be Balanced for the Maize C4 Pathway. *Plant Physiol* 164: 466–480

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bellasio C, Griffiths H (2014b) Acclimation of C4 metabolism to low light in mature maize leaves could limit energetic losses during progressive shading in a crop canopy. *J Exp Bot* 65: 3725–3736

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Björkman O (1981) Responses to different quantum flux densities. *Physiological plant ecology I*

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Boardman NK (1977) Comparative Photosynthesis of Sun and Shade Plants. *Annu Rev Plant Physiol* 28: 355–377

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bovdilova A, Alexandre BM, Höppner A, Luís IM, Alvarez CE, Bickel D, Gohlke H, Decker C, Nagel-Steger L, Aseekh S, et al (2019) Posttranslational Modification of the NADP-Malic Enzyme Involved in C4 Photosynthesis Modulates the Enzymatic Activity during the Day. *Plant Cell* 31: 2525–2539

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cañas RA, Quilleré I, Lea PJ, Hirel B (2010) Analysis of amino acid metabolism in the ear of maize mutants deficient in two cytosolic glutamine synthetase isoenzymes highlights the importance of asparagine for nitrogen translocation within sink organs. *Plant Biotechnol J* 8: 966–978

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Casati P, Drincovich MF, Andreo CS, Donahue R, Edwards GE (1998) UV-B, red and far-red light regulate induction of the C4 isoform of NADP-malic enzyme in etiolated maize seedlings. *Funct Plant Biol* 25: 701–708

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chao Q, Liu X-Y, Mei Y-C, Gao Z-F, Chen Y-B, Qian C-R, Hao Y-B, Wang B-C (2014) Light-regulated phosphorylation of maize phosphoenolpyruvate carboxykinase plays a vital role in its activity. *Plant Mol Biol* 85: 95–105

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chapman KSR, Hatch MD (1981) Aspartate Decarboxylation in Bundle Sheath Cells of Zea mays and Its Possible Contribution to C3 Photosynthesis. *Funct Plant Biol* 8: 237–248

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen Y, Wu D, Mu X, Xiao C, Chen F, Yuan L, Mi G (2016) Vertical Distribution of Photosynthetic Nitrogen Use Efficiency and Its Response to Nitrogen in Field-Grown Maize. *Crop Sci* 56: 397–407

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Collison RF, Raven EC, Pignon CP, Long SP (2020) Light, Not Age, Underlies the Maladaptation of Maize and Miscanthus Photosynthesis to Self-Shading. *Front Plant Sci* 11: 783

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Czajka JJ, Kambhampati S, Tang YJ, Wang Y, Allen DK (2020) Application of Stable Isotope Tracing to Elucidate Metabolic Dynamics During Yarrowia lipolytica α -Ionone Fermentation. *iScience* 23: 100854

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

- Delgado-Alvarado A, Walker RP, Leegood RC (2007) Phosphoenolpyruvate carboxykinase in developing pea seeds is associated with tissues involved in solute transport and is nitrogen-responsive. Plant Cell Environ 30: 225–235**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Detarsio E, Wheeler MCG, Bermúdez VAC, Andreo CS, Drincovich MF (2003) Maize C4 NADP-malic enzyme: Expression in escherichia coli and characterization of site-directed mutants at the putative nucleotide-binding sites. J Biol Chem 278: 13757–13764**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dwyer LM, Stewart DW (1986) Effect of leaf age and position on net photosynthetic rates in maize (Zea Mays L.). Agric For Meteorol 37: 29–46**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Evans JR, Poorter H (2001) Photosynthetic acclimation of plants to growth irradiance: the relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain. Plant Cell Environ 24: 755–767**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fox J, Weisberg S (2018) An R Companion to Applied Regression. SAGE Publications**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Furbank RT (2011) Evolution of the C(4) photosynthetic mechanism: are there really three C(4) acid decarboxylation types? J Exp Bot 62: 3103–3108**
- Ghannoum O, Evans JR, Chow WS, Andrews TJ, Conroy JP, von Caemmerer S (2005) Faster Rubisco is the key to superior nitrogen-use efficiency in NADP-malic enzyme relative to NAD-malic enzyme C4 grasses. Plant Physiol 137: 638–650**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Harmer SL, Covington MF, Bläsing O, Stitt M (2018) Circadian regulation of global gene expression and metabolism. Annual Plant Reviews online 133–165**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hatch MD, Kagawa T (1976) Photosynthetic activities of isolated bundle sheath cells in relation to differing mechanisms of C-4 pathway photosynthesis. Arch Biochem Biophys 175: 39–53**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Heinrich P, Kohler C, Ellmann L, Kuerner P, Spang R, Oefner PJ, Dettmer K (2018) Correcting for natural isotope abundance and tracer impurity in MS-, MS/MS- and high-resolution-multiple-tracer-data from stable isotope labeling experiments with IsoCorrectorR. Sci Rep 8: 17910**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hernández-Sebastià C, Marsolais F, Saravitz C, Israel D, Dewey RE, Huber SC (2005) Free amino acid profiles suggest a possible role for asparagine in the control of storage-product accumulation in developing seeds of low-and high-protein soybean lines. J Exp Bot 56: 1951–1963**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hikosaka K, Terashima I (1995) A model of the acclimation of photosynthesis in the leaves of C3 plants to sun and shade with respect to nitrogen use. Plant Cell Environ 18: 605–618**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kambhampati S, Ajewole E, Marsolais F (2018) Advances in Asparagine Metabolism. In FM Cánovas, U Lüttge, R Matyssek, eds, Progress in Botany Vol. 79. Springer International Publishing, Cham, pp 49–74**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kanai R, Edwards GE (1999) The biochemistry of C4 photosynthesis. C4 plant biology 49: 87**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Khamis S, Lamaze T, Farineau J (1992) Effect of nitrate limitation on the photosynthetically active pools of aspartate and malate in maize, a NADP malic enzyme C 4 plant. Physiol Plant 85: 223–229**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Lea PJ, Azevedo RA (2007) Nitrogen use efficiency. 2. Amino acid metabolism. Ann Appl Biol 151: 269–275**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Lin F, Jiang L, Liu Y, Lv Y, Dai H, Zhao H (2014) Genome-wide identification of housekeeping genes in maize. Plant Mol Biol 86: 543–554**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Lohaus G, Bükler M, Hußmann M, Soave C, Heldt H-W (1998) Transport of amino acids with special emphasis on the synthesis and transport of asparagine in the Illinois Low Protein and Illinois High Protein strains of maize. Planta 205: 181–188**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Ma F, Jazmin LJ, Young JD, Allen DK (2014) Isotopically nonstationary ¹³C flux analysis of changes in Arabidopsis thaliana leaf metabolism due to high light acclimation. Proceedings of the National Academy of Sciences 111: 16967–16972**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Meister M, Agostino A, Hatch MD (1996) The roles of malate and aspartate in C4 photosynthetic metabolism of *Flaveria bidentis* (L.). *Planta* 199: 262–269

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Niinemets Ü (2016) Leaf age dependent changes in within-canopy variation in leaf functional traits: a meta-analysis. *J Plant Res* 129: 313–338

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Osorio S, Vallarino JG, Szczewka M, Ufaz S, Tzin V, Angelovici R, Galili G, Fernie AR (2014) Extraction and Measurement the Activities of Cytosolic Phosphoenolpyruvate Carboxykinase (PEPCK) and Plastidic NADP-dependent Malic Enzyme (ME) on Tomato (*Solanum lycopersicum*). *Bio-protocol* 4: e1122–e1122

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pandurangan S, Pajak A, Molnar SJ, Cober ER, Dhaubhadel S, Hernández-Sebastià C, Kaiser WM, Nelson RL, Huber SC, Marsolais F (2012) Relationship between asparagine metabolism and protein concentration in soybean seed. *J Exp Bot* 63: 3173–3184

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Penfield S, Clements S, Bailey KJ, Gilday AD, Leegood RC, Gray JE, Graham IA (2012) Expression and manipulation of phosphoenolpyruvate carboxykinase 1 identifies a role for malate metabolism in stomatal closure. *Plant J* 69: 679–688

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pengelly JLL, Sirault XRR, Tazoe Y, Evans JR, Furbank RT, von Caemmerer S (2010) Growth of the C4 dicot *Flaveria bidentis*: photosynthetic acclimation to low light through shifts in leaf anatomy and biochemistry. *J Exp Bot* 61: 4109–4122

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pick TR, Bräutigam A, Schlüter U, Denton AK, Colmsee C, Scholz U, Fahnenstich H, Pieruschka R, Rascher U, Sonnewald U, et al (2011) Systems analysis of a maize leaf developmental gradient redefines the current C4 model and provides candidates for regulation. *Plant Cell* 23: 4208–4220

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pinto H, Powell JR, Sharwood RE, Tissue DT, Ghannoum O (2016) Variations in nitrogen use efficiency reflect the biochemical subtype while variations in water use efficiency reflect the evolutionary lineage of C4 grasses at inter-glacial CO2. *Plant Cell Environ* 39: 514–526

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pinto H, Sharwood RE, Tissue DT, Ghannoum O (2014) Photosynthesis of C3, C3–C4, and C4 grasses at glacial CO2. *J Exp Bot* 65: 3669–3681

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pons TL (2016) Regulation of Leaf Traits in Canopy Gradients. In K Hikosaka, Ü Niinemets, NPR Anten, eds, *Canopy Photosynthesis: From Basics to Applications*. Springer Netherlands, Dordrecht, pp 143–168

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sage RF, Christin P-A, Edwards EJ (2011) The C(4) plant lineages of planet Earth. *J Exp Bot* 62: 3155–3169

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Seebauer JR, Moose SP, Fabbri BJ, Crossland LD, Below FE (2004) Amino acid metabolism in maize earshoots. Implications for assimilate preconditioning and nitrogen signaling. *Plant Physiol* 136: 4326–4334

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sommer M, Bräutigam A, Weber APM (2012) The dicotyledonous NAD malic enzyme C4 plant *Cleome gynandra* displays age-dependent plasticity of C4 decarboxylation biochemistry. *Plant Biol* 14: 621–629

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sonawane BV, Sharwood RE, Whitney S, Ghannoum O (2018) Shade compromises the photosynthetic efficiency of NADP-ME less than that of PEP-CK and NAD-ME C4 grasses. *J Exp Bot* 69: 3053–3068

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stelpflug SC, Sekhon RS, Vaillancourt B, Hirsch CN, Buell CR, de Leon N, Kaeppler SM (2016) An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to Explore Root Development. *Plant Genome*. doi: 10.3835/plantgenome2015.04.0025

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stitt M, Zhu X-G (2014) The large pools of metabolites involved in intercellular metabolite shuttles in C4 photosynthesis provide enormous flexibility and robustness in a fluctuating light environment. *Plant Cell Environ* 37: 1985–1988

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Subedi KD, Ma BL (2005) Ear Position, Leaf Area, and Contribution of Individual Leaves to Grain Yield in Conventional and Leafy Maize Hybrids. *Crop Sci* 45: 2246–2257

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sugiyama T, Mizuno M, Hayashi M (1984) Partitioning of nitrogen among ribulose-1, 5-bisphosphate carboxylase/oxygenase, phosphoenolpyruvate carboxylase, and pyruvate orthophosphate dikinase as related to biomass productivity in maize seedlings. *Plant Physiol* 75: 665–669

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Szeczowka M, Heise R, Tohge T, Nunes-Nesi A, Vosloh D, Huege J, Feil R, Lunn J, Nikoloski Z, Stitt M, et al (2013) Metabolic fluxes in an illuminated Arabidopsis rosette. Plant Cell 25: 694–714

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Taub DR, Lerdau MT (2000) Relationship between leaf nitrogen and photosynthetic rate for three NAD-ME and three NADP-ME C4 grasses. Am J Bot 87: 412–417

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tazoe Y, Noguchi K, Terashima I (2006) Effects of growth light and nitrogen nutrition on the organization of the photosynthetic apparatus in leaves of a C4 plant, Amaranthus cruentus. Plant Cell Environ 29: 691–700

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ubierna N, Sun W, Kramer DM, Cousins AB (2013) The efficiency of C4 photosynthesis under low light conditions in Zea mays, Miscanthus x giganteus and Flaveria bidentis. Plant Cell Environ 36: 365–381

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Walker RP, Acheson RM, Técsi LI, Leegood RC (1997) Phosphoenolpyruvate Carboxykinase in C4 Plants: Its Role and Regulation. Funct Plant Biol 24: 459–468

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Walker RP, Chen ZH, Tecsli LI, Famiani F, Lea PJ, Leegood RC (1999) Phosphoenolpyruvate carboxykinase plays a role in interactions of carbon and nitrogen metabolism during grape seed development. Planta 210: 9–18

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Walters RG (2005) Towards an understanding of photosynthetic acclimation. J Exp Bot 56: 435–447

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang Y, Bräutigam A, Weber APM, Zhu X-G (2014) Three distinct biochemical subtypes of C4 photosynthesis? A modelling analysis. J Exp Bot 65: 3567–3578

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ward DA, Woolhouse HW (1986) Comparative effects of light during growth on the photosynthetic properties of NADP-ME type C4 grasses from open and shaded habitats. II. Photosynthetic enzyme activities and metabolism. Plant Cell Environ 9: 271–277

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weissmann S, Ma F, Furuyama K, Gierse J, Berg H, Shao Y, Taniguchi M, Allen DK, Brutnell TP (2016) Interactions of C4 Subtype Metabolic Activities and Transport in Maize Are Revealed through the Characterization of DCT2 Mutants. Plant Cell 28: 466–484

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wingler A, Walker RP, Chen ZH, Leegood RC (1999) Phosphoenolpyruvate carboxykinase is involved in the decarboxylation of aspartate in the bundle sheath of maize. Plant Physiol 120: 539–546

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)