1 **Title:** Insulin-dependent and -independent dynamics of insulin receptor trafficking in
2 muscle cells

3 **Authors:** Haoning Howard Cen¹, Jason Rogalski², Libin Abraham³, Michael R. Gold³,
4 Leonard J. Foster², James D. Johnson^{1*}

5

6 **Affiliation:**

7 1 Department of Cellular and Physiological Sciences, Life Sciences Institute, University of
8 British Columbia, Vancouver, BC. Canada.

9 2 Department of Biochemistry & Molecular Biology, and Michael Smith Laboratories,
10 University of British Columbia, Vancouver, BC, Canada

11 3 Department of Microbiology and Immunology, Life Life Sciences Institute, University of
12 British Columbia, Vancouver, BC. Canada

13

14 **Running title:** Muscle cell insulin receptor trafficking and interactions

15 **Keywords:** insulin, insulin receptor, caveolin, clathrin, endocytosis

16

17 ***Address correspondence to:** James D. Johnson, Ph.D., Professor. Diabetes Research
18 Group, Dept. of Cellular and Physiological Sciences & Dept. of Surgery, University of
19 British Columbia, 5358 Life Sciences Building, 2350 Health Sciences Mall, Vancouver,
20 BC, Canada, V6T 1Z3

21 E-mail: James.D.Johnson@ubc.ca; Fax: (604) 822-2316

22 Twitter: @JimJohnsonSci

23

Acknowledgments: Spinning disk confocal and STED microscopy were performed in the LSI IMAGING facility of the Life Sciences Institute of the University of British Columbia using infrastructure funded by the Canadian Foundation of Innovation and BC Knowledge Development Fund as well as a Strategic Investment Fund (Faculty of Medicine, University of British Columbia). We thank Dr. Ivan Robert Nabi and Dr. Guang Gao of the LSI IMAGING facility for guidance on imaging experiments. Mass Spectrometry used here was supported by the Canada Foundation for Innovation and Genome Canada (264PRO). We thank Yuanyuan Amanda Yang for guidance on Python and R coding and Yi Han Xia for assistance in R coding. The schematics in Figure 7 is created with BioRender.com.

Contributions: HHC designed the study, performed all experiments, analyzed data, and wrote the manuscript. JR performed mass spectrometry, analyzed the output data, and edited the manuscript. LA and MG contributed to the MATLAB scripts and the analysis of single particle tracking and edited the manuscript. LF contributed to the design of the mass-spectrometry experiments and edited the manuscript. JDJ designed the study, supervised the research, and edited the manuscript. JDJ is the ultimate guarantor of this work.

Funding: This study was funded by a Canadian Institutes of Health Research Operating Grant (P-133692) to JDJ.

Conflict of interest: The authors declare that they have no conflicts of interest with the contents of this article.

50

51 **Abstract**

52 Insulin resistance contributes to type 2 diabetes and can be driven by
53 hyperinsulinemia. Insulin receptor (INSR) internalization and cell-surface dynamics at
54 rest and during insulin exposure are incompletely understood in muscle cells. Using
55 surfacing labeling and live-cell imaging, we observed robust basal internalization of INSR
56 in C2C12 myoblasts, without a robust effect of added insulin. Mass-spectrometry analysis
57 of INSR-binding proteins identified potential molecular mechanisms associated with
58 internalization. We confirmed known interactors, including IGF1R, but also identified
59 underappreciated INSR-binding factors such as ANXA2. Protein-protein interaction
60 network mapping suggested links between INSR and caveolin-mediated endocytosis.
61 INSR interacted with both caveolin and clathrin heavy chain (CLTC) in mouse skeletal
62 muscle and C2C12 myoblasts. Whole cell 2D super-resolution imaging revealed that high
63 levels of insulin (20 nM) increased INSR colocalization with CAV1 but decreased its
64 colocalization with CLTC. Single particle tracking confirmed the colocalization of
65 cell-surface INSR with both over-expressed CAV1-mRFP and CLTC-mRFP. INSR tracks
66 that colocalized with CAV1 exhibited longer radii and lifetimes, regardless of insulin
67 exposure, compared to non-colocalized tracks, whereas insulin further increased the
68 lifetime of INSR/CLTC colocalized tracks. Overall, these data suggest that muscle cells
69 utilize both CAV1 and CLTC-dependent pathways for INSR dynamics and internalization.

70

71 **Introduction**

72 The insulin receptor (INSR) is one of the most studied membrane proteins due to its
73 important roles in insulin resistance and diabetes. However, many fundamental aspects
74 of INSR biology remain unclear, including the mechanisms and kinetics of its
75 internalization. The INSR can be internalized via endocytosis^{1,2} and endosomal INSR
76 localization contributes to the phosphorylation of specific downstream targets when

compared with INSR at the cell surface^{3,4}. Moreover, ligand-receptor complex dissociation in acidified endosomes permits insulin degradation while the receptor can be sorted to recycling or degradation pathways^{1,2}. INSR internalization is reduced in monocytes^{5,6} and adipocytes⁷ obtained from patients with insulin resistance and type 2 diabetes, suggesting the importance of this process in disease pathogenesis.

Tissue-specific insulin resistance has been implicated in the development of diabetes^{8,9}, but it is less clear whether there are tissue-specific INSR internalization kinetics and/or mechanisms. Interestingly, adipocytes internalize insulin at a lower rate in the presence of high insulin concentrations, as compared to low insulin conditions¹⁰, while hepatocytes are the opposite¹¹. The consensus view is that INSR is endocytosed via a clathrin-coated pit mechanism, based on evidence from a variety of cell types, including human and rat hepatocytes, fibroblasts, and lymphocytes¹¹⁻¹⁴. However, other studies have shown that caveolin-rich membrane lipid rafts, caveolae, are critical for INSR internalization in other cell types, including adipocytes^{15,16}. In pancreatic β -cells, we showed that caveolin-1 (CAV1) mediates INSR internalization, bypassing clathrin positive compartments⁴. In muscle cells, previous studies have shown that clathrin-mediated endocytosis is involved in INSR internalization, but the role of the caveolin-mediated pathway has not been investigated¹⁷. The INSR internalization kinetics and routes are less well characterized in muscle cells.

In the present study, we employed inter-domain tagged INSR with fluorescent proteins⁴ and SNAP-tag¹⁸, surface biotinylation assays, as well as super-resolution and TIRF imaging, to explore the dynamics of INSR internalization and the effects of ambient insulin. We also used proteomics to identify INSR interacting proteins and defined the relationship between INSR and caveolin- or clathrin-mediated internalization pathways. Together, our data show that INSR exhibited a high degree of insulin-independent internalization in C2C12 myoblasts.

Results

Effects of insulin on INSR internalization

First, we characterized INSR internalization in C2C12 myoblasts under basal conditions and in the context of added insulin. A high rate of basal INSR internalization was observed in high-glucose serum-free DMEM medium when compared with PBS. INSR internalization over 5 min was not significantly stimulated by addition of insulin to the DMEM medium (Fig. 1A). Therefore, INSR has a high rate of constitutive internalization in this cell type, which is enhanced by the nutrients in medium, but that is largely independent of acute exposure to insulin across a range of physiological doses. These observations in myoblasts are consistent with our previous measurements in differentiated C2C12 myotubes¹⁹.

Qualitative analysis of INSR kinetics in the context of insulin doses

We also visualized INSR internalization at single-cell level using inter-domain-tagged INSR, which we have previously found more faithfully recapitulates the sub-cellular localization of INSR compared to terminal fusion proteins⁴. C2C12 myoblast cells with stable overexpression of SNAP-tagged INSR isoform A (INSR-A-SNAP) were labeled using a cell non-permeable SNAP-Surface Alexa Fluor 488 dye (Fig. 1B). Live-cell time-lapse imaging of surface-labeled INSR-A-SNAP by spinning disk confocal microscopy revealed INSR internalization in the context of 0, 0.2, and 20 nM insulin over 15 min (Fig. 1B, Video S1). Consistent with the surface biotinylation assay results in Figure 1A, we did not observe a clear difference in INSR internalization in the presence of insulin in these cells. Internalized INSR vesicles exerted dynamic movement and interaction, which was more evident between 20-30 min (Fig. 1C-F, Video S2). INSR-containing vesicles exhibited many apparent fusion and fission events regardless of insulin concentrations. Some smaller vesicles merged into a larger vesicle that remained in the field undergoing limited and apparently random motion (one example

is shown in Fig. 1F), suggestive of a fusion event. A caveat associated with 2D imaging and limited Z resolution is that the observed INSR vesicles might not all be on the same optical plane. Our experiments using surface labeling of SNAP-tagged INSR permitted the visualization of the dynamic behavior of internalized INSR vesicles. These qualitative observations motivated the more quantitative studies below.

Quantitative analysis of INSR kinetics in the context of insulin

Lateral movement of tyrosine kinase receptors has been linked to their functional state. Ligand binding of some monomeric receptors, such as EGFR, promotes their dimerization and aggregation and decreases their lateral mobility²⁰. While INSRs are thought to already exist as dimers in plasma membrane, aggregation of INSR by antibodies can stimulate INSR signaling²⁰, and INSR may need to localize to membrane lipid rafts for normal signaling^{21,22}. Therefore, we thought to quantitatively assessed whether insulin affects the lateral movement of INSR in or near the plasma membrane using time-lapse TIRF microscopy of C2C12 myoblast transiently expressing inter-domain-tagged INSR-A-EGFP. We imaged on the same cells at 0, 6, or 11 min in the presence of 2 nM insulin, as well as a control group without insulin to control for other variables such as photobleaching (Fig. 2A). Single particle tracking was conducted on the INSR-A-EGFP (Fig. 2B), and we calculated their diffusion coefficient (Fig. 2C,D) and track radius (Fig. 2E,F). Insulin did not affect the track radius or diffusion coefficient at any time points. Together with the qualitative data reported above, these observations suggest that the majority of INSR mobility is independent of acute insulin stimulation in this cell model.

Novel INSR interactors identified by immunoprecipitation and mass spectrometry

To explore the molecular mechanisms associated with INSR internalization, we conducted co-immunoprecipitation followed by mass-spectrometry to identify INSR

interactors. We also used INSR knockout C2C12 myoblasts as a negative control and an INSR antibody that does not cross-react with Insulin-like growth factor 1 receptor (IGF1R)²³ to identify INSR interactors with higher specificity than any previously reported study. After filtering the detected proteins against the INSR knockout cell control lysates, we identified INSR and five high-confidence INSR-binding proteins (Fig. 3A). The immunoprecipitated abundance of these detected interactors was not significantly changed by the addition of 2 nM insulin (Fig. 4A). INSR is well established to form heterodimers with IGF1R^{24,25}, and its immunoprecipitation represented a type of internal positive control for this study. The other detected proteins are not well-known INSR interactors.

To discover whether any of the identified INSR-binding proteins might be connected with endocytosis machinery, protein-protein interaction (PPI) network was constructed between these co-immunoprecipitated proteins and all the proteins under the “Endocytosis” pathway in Kyoto Encyclopedia of Genes and Genomes (KEGG). We manually extracted a subnetwork that connected all our identified proteins, except for HIST1H1B which did not connect with any proteins in this network (Fig. 3B). In the resultant PPI network, annexin A2 (ANXA2) is connected with CAV1 and CAV3 (Fig. 3B), and it has been reported to localize in caveolae²⁶. Ribosomal protein SA (RPSA) is a laminin receptor reported to localize in plasma membrane lipid rafts²⁷. Heat shock protein B1 (HSPB1; a.k.a. HSP27) is exclusively found in insulin-responsive tissues and has been implicated in insulin sensitivity and INSR/IGF1R signalling²⁸⁻³⁰. The PPI network and cellular localization of these interactors pointed to the possibility that INSR may also associate with caveolins and/or related proteins in myoblasts.

Insulin-dependent interaction between INSR and CAV3 or CLTC *in vivo*

We sought to confirm that INSR could interact with both caveolin and clathrin in skeletal muscle *in vivo*. Skeletal muscle was collected 5 or 10 min after intraperitoneal

injection of insulin (1.5 U/kg) or PBS, timepoints sufficient to measure acute insulin signalling via both AKT and ERK signalling pathways (Fig. 3C). Acute insulin administration did not change the amount of INSR that could be immunoprecipitated from skeletal muscle, nor did it alter the association between INSR and CLTC (Fig. 3D). However, 10 min after insulin injection, there was a significantly increased interaction between INSR and caveolin, specifically, the mature skeletal muscle-specific isoform CAV3^{31,32} (Fig. 3D). These data show that acute insulin signaling results in an increased association between INSR and CAV3 *in vivo*, but does not define the cellular context for this interaction.

Super-resolution imaging of INSR, CAV1, and CLTC interactions in the presence of insulin

We next used super-resolution STED imaging to examine colocalization between INSR and clathrin (CLTC) or caveolin (C2C12 myoblasts only express CAV1 before differentiation³⁰). C2C12 myoblast cells with stable overexpression of INSR-A-SNAP were labeled by cell-nonpermeable SNAP substrate, to image only newly internalized INSR. Cells were incubated with 0, 0.2 or 20 nM insulin for 30 min, and then fixed. Endogenous CLTC and CAV1 were then labeled using immunofluorescent antibody staining (Fig. 4A). The colocalization between INSR-SNAP, CAV1, and CLTC was accessed by Object Pearson Coefficient (Fig. 4B,C). The degree of colocalization between INSR and CAV1 was similar in the presence of 0 or 0.2 nM insulin but was higher under 20 nM insulin (Fig. 4B). However, INSR and CLTC colocalized more at 0.2 nM insulin, but less at 20 nM insulin, compared to 0 nM insulin (Fig. 4B). CAV1 and CLTC do not have known biological interactions; therefore, their low level of colocalization served as an internal negative control and, indeed, it did not change under different insulin levels (Fig. 4B). Using the same data, the relative colocalizations between INSR and CAV1 or CLTC were also considered at different insulin concentrations, further

illustrating how high insulin drives INSR to colocalize with CAV1 (Fig. 4C). Together, these data revealed the insulin-dependent interaction of INSR with both CAV1 in vivo and in vitro, alongside the constitutive interaction with CLTC.

Effects of CAV1 and CLTC on INSR mobility

To better understand the interplay between INSR, CAV1, and CLTC, we co-expressed INSR-B-TagBFP with CAV1-mRFP or CLTC-mRFP in C2C12 myoblasts and imaged them by TIRF time-lapse microscopy. Colocalization between INSR and CAV1 or CLTC were observed (Fig. 5A,B). Some colocalized puncta moved or disappeared together with closely associated intensity, referred to here as colocalized tracks (examples in Fig. 5C,D). Tracks were considered as colocalized when the distance between the centers of particle signals was less than 2 pixels for more than 4 continuous frames. These observations suggested that INSR interacted with CAV1 and CLTC on the cell surface and might be endocytosed through both caveolin- and clathrin-mediated pathways.

We also conducted single particle tracking on these INSR puncta to define the effects of CAV1 or CLTC on INSR movement in the plasma membrane. Similar to our observations above (Fig. 2), insulin did not alter the diffusion coefficient or track radius of INSR. Cells with CAV1 co-expression had lower diffusion coefficients and track radius of INSR compared to cells with CLTC co-expression, regardless of ambient insulin exposure (Fig. 5E-H). CLTC-expressing cells under 0 nM insulin had shorter INSR lifetimes than other groups (Fig. 5I,J), which might suggest a more rapid internalization. These data suggest that CAV1 may restrain INSR movement, relative to CLTC.

Finally, we performed an analysis within the same cells, comparing INSR tracks that were colocalized or not colocalized with either CAV1 or CLTC tracks. Within cells, INSR spent more time associated with CAV1, and for longer periods, compared with CLTC (Fig. 6A,B). There were no clear effects of time on the association between INSR and

CAV1 or CLTC across different moments of their lifetimes together (Fig. 6A,B). Within the same cells, we compared the colocalized and the non-colocalized INSR tracks, for both CAV1 and CLTC, and found no significant differences in mean diffusion coefficients (Fig. 6C,D). The track radius of INSR tracks colocalized with CAV1 was larger than non-colocalized tracks regardless of insulin (Fig. 6E). The track radius of INSR tracks colocalized with CLTC was larger than non-colocalized tracks, but only in the no insulin condition (Fig. 6F). The lifetimes of INSR tracks colocalized with CAV1 were longer than non-colocalized tracks regardless of insulin (Fig. 6G). INSR tracks colocalized with CLTC also had longer lifetimes than non-colocalized tracks, and insulin further increased the lifetime of colocalized tracks (Fig. 6H). Overall, our live-cell imaging of INSR confirmed its separate interactions with either CAV1 or CLTC on the cell surface.

Discussion

The goal of this study was to explore the INSR trafficking dynamics and interactions in a myoblast cell line. We demonstrated that INSR was constitutively internalized both in the absence and presence of insulin. We used a high-stringency proteomics approach to identify specific INSR binding proteins and PPI network modeling to predict a role for caveolin in INSR internalization. Biochemical experiments, super-resolution imaging, and TIRF imaging suggested a role for CAV1 in INSR internalization and cell-surface trafficking (Fig. 7).

The constitutive internalization of INSR has been described previously in lymphocytes and shown not to require ligand binding and kinase activity, although insulin-stimulated autophosphorylation of INSR could accelerate internalization¹⁴. In differentiated C2C12 myotubes, INSR phosphorylation levels were constitutively high, with little additional activation by insulin across the physiological range¹⁹, consistent with the lack of stimulation of INSR internalization by acute insulin treatment in this study. Surface labeling by SNAP-tag enabled clear visualization of internalized INSR-containing

vesicles, permitting the study of INSR endosomal sorting and traffic. The lateral movement of INSR in or near the plasma membrane was also not significantly affected by insulin, consistent with a previous study that reported unchanged INSR diffusion coefficient with or without insulin in HEK293T, HepG2, and human IR-overexpressing Rat-1 cells³³. CAV1 may have a role in decreasing the mobility of INSR. Indeed, caveolae have restricted lateral movement in the plasma membrane due to the connection to actin cytoskeleton through Filamin-A³⁴, so the CAV1-dependent decrease in INSR mobility might reflect the interaction of INSR with caveolae. Similarly, the accumulation of ganglioside GM3 in TNF α -induced insulin resistance state was associated with the elimination of INSR from the caveolae and increased INSR mobility on the cell surface²¹. Additional studies will be required to fully define the processes that control INSR movement within the plasma membrane.

The possibility of more than one internalization pathway for INSR has been understudied. The current study supports the concept that INSR utilizes both clathrin- and caveolin-mediated endocytosis. Another study confirmed the involvement of the clathrin-mediated pathway in myoblast by identifying LMBD1 protein as the cargo-specific adaptor of INSR in clathrin-mediated endocytosis¹⁷. The combination of dual clathrin and caveolin endocytosis routes has been demonstrated for IGFR³⁵, which has a similar structure and function as INSR and can form heterodimers with INSR³⁶. In hepatocytes, it has also been suggested that INSRs are internalized in two functionally separate pathways based on the much more rapid recycling of receptors internalized by high concentrations of insulin (17.2 nM) versus low insulin (0.17 nM), and treatments known to inhibit coated pit formation blocked internalization of the insulin-INSR complex at low concentrations of insulin but allowed internalization in response to high concentrations of insulin¹¹. Most existing studies on INSR internalization looked at one pathway at a time. The mechanisms of coordinated multiple internalization routes, implied by the insulin-dependent clathrin and caveolin interactions we observed in muscle cells, might

direct INSR to different destinations (e.g. recycle or degradation)³⁷ and should be comprehensively characterized in more cell types to assess potential roles in tissue-specific insulin resistance.

We used stringent proteomics and INSR CRISPR knockout control cells to identify known, novel, and under-appreciated interactors of INSR. As expected, we identified IGF1R, an established dimerization partner of INSR. We also demonstrated the specific binding of HSBP1/HSP27, which is involved in tyrosine kinase signalling pathways downstream of p38 MAPK, ERK, and MAPKAPK5³⁸ and has been previously linked to insulin sensitivity²⁸⁻³⁰. We demonstrated an interaction between INSR and RPSA, which has been previously co-immunoprecipitated with INSR and insulin receptor substrates in 3T3L1 preadipocytes³⁹. In other cell types, RPSA drives ERK signalling and cytoskeletal organization, with nuclear functions affecting cell growth, survival, migration, and protein synthesis²⁷. ANXA2 is not a well-known interactor of INSR, but one study reported ANXA2 interaction with INSR and IGF1R⁴⁰. Other studies found that INSR signalling can tyrosine-phosphorylate^{41,42} and SUMOylate⁴³ ANXA2, which mediates insulin-stimulated actin rearrangement. Interestingly, INSR internalization seemed to be necessary for ANXA2 phosphorylation because mutant INSR that was incapable of internalization but normal for tyrosine kinase signalling failed to phosphorylate ANXA2⁴¹. On the other hand, ANXA2 has a strong direct connection with epidermal growth factor receptor (EGFR) in the PPI network which belongs to the same receptor tyrosine kinase superfamily as INSR⁴⁴. ANXA2 knockdown halted EGFR endocytosis beyond the early endosome, which was accompanied by enhanced EGFR and AKT phosphorylation⁴⁵. These data support the possibility that ANXA2 may have a profound role in INSR trafficking and signalling that is worth investigating. Both *Anxa2* and *Rspa* mRNAs were upregulated by prolonged insulin in C2C12 myotubes⁸. HIST1H1B belongs to the family of linker histones that is important in chromatin organization⁴⁶, but the meaning of the INSR interaction requires further investigation. This proteomic analysis provides clear evidence for a

limited number of proteins that are likely to be direct interactors with INSR in myoblast cells, and demonstrates that the majority of these interactors were not significantly altered by insulin exposure.

There are caveats and unexplained observations in the current study. Although STED superresolution microscopy might be able to reach 20-nm resolution, primary and secondary antibody labeling might add ~30-nm distance and reduce the accuracy⁴⁷. TIRF microscopy has an axial resolution of ~100 nm and a defraction-limited lateral resolution (~200 – 300 nm)⁴⁸. Therefore, the colocalization analyses supports the co-occurrence but may not confirm the functional interaction of the proteins. Nonetheless, the insulin-dependent colocalization observed in STED microscopy and the track colocalization over multiple frames observed in TIRF microscopy are evidences that the proteins had functional interaction instead of random colocalization. The CAV1- or CLTC-colocalized INSR tracks had longer lifetime than non-colocalized tracks, which conflicted with the assumption that CAV1 or CLTC internalized INSR, but instead, they seemed to retain INSR on the membrane. There are multiple possible explanations; for example, overexpressed CAV1 might stabilize caveolae⁴⁹, or CLTC could form stable flat lattices, termed clathrin plaques⁵⁰, which might have novel involvement in INSR internalization. We did not perform loss-of-function studies to confirm the roles of caveolin and clathrin in INSR endocytosis. Further research is needed to elucidate the roles of specific INSR internalization routes on insulin actions. Inhibition of clathrin-mediated endocytosis of INSR by knocking down LMBD1 in myoblasts increased activation of the INSR and AKT¹⁷. We also recently discovered that reducing CAV1-dependent INSR internalization blocked the activation of ERK, but not AKT, signalling in pancreatic beta-cells⁴. Cholesterol extraction by methyl- β -cyclodextrin, which is often used as a caveolae-specific inhibitor but also affects clathrin-coated pit formation⁵¹, inhibited the phosphorylation of IRS1, SHC, and AKT, but not ERK or INSR phosphorylation, in adipocytes^{16,52}. In contrast, inhibition of both clathrin- and caveolin-mediated endocytosis

by the dynamin K44A dominant-negative mutant⁵³ reduced the phosphorylation of SHC and ERK phosphorylation, but not IRS1 or AKT phosphorylation, in rat hepatoma H4IIE cells¹². Therefore, INSR internalization enables selective activation of downstream signalling pathways, and in a tissue-specific manner.

In summary, this study provides important information on the internalization of INSR in myoblast cells and defines the stringent INSR interactome. We also provide evidence using a variety of methods that INSR is internalized via both clathrin- and caveolin-dependent pathways in this cell model. The current study can serve as a stepping stone for more detailed research on how multiple internalization routes can provide cell-type customized regulations on INSR internalization.

Materials and Methods

Cell culture

The C2C12 mouse myoblast (ATCC cell line provided by Dr. Brain Rodrigues, University of British Columbia, Vancouver, Canada) was maintained in Dulbecco's modified Eagle's medium (DMEM, Gibco) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco), and 1% penicillin-streptomycin (100 µg/ml; Gibco). All downstream experiments were conducted after 6-hour serum starvation in serum-free medium (DMEM supplemented with 1% penicillin-streptomycin), except as otherwise indicated.

Western blot analyses

C2C12 myotubes or mice skeletal muscle (gastrocnemius) tissues were sonicated in RIPA buffer (50 mM β-glycerol phosphate, 10 mM HEPES, 1% Triton X-100, 70 mM NaCl, 2 mM EGTA, 1 mM Na₃VO₄, and 1 mM NaF) supplemented with complete mini protease inhibitor cocktail (Roche, Laval, QC). Lysates were incubated in Blue Loading Buffer containing 50 mM DTT (Cell Signaling, Danvers, MA) at 95°C for 5 min and

resolved by SDS-PAGE. Proteins were then transferred to PVDF membranes (BioRad, CA) and probed with antibodies against p-ERK1/2 (Thr202/Tyr204) (1:1000, Cat. #4370), ERK1/2 (1:1000, Cat. #4695), p-AKT (Ser473) (1:1000, Cat. #9271), p-AKT (Thr308) (1:1000, Cat. #9275), AKT (1:1000, Cat. #9272), INSR- β subunit (1:1000, Cat. #3020S), p-INSR β (Tyr1150/1151) (1:1000, Cat. #3024), FOXO1 (1:1000, Cat. #2880), p-FOXO1 (Thr24) (1:1000, Cat. #9464), all from Cell Signalling (CST), and β -tubulin (1:2000, Cat. #T0198, Sigma). The signals were detected by secondary HRP-conjugated antibodies (anti-mouse Cat. #7076, anti-rabbit Cat. #7074; Cell Signaling) and Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific). Protein band intensities were quantified with Image Studio Lite software (LI-COR).

Surface Protein Biotinylation Assay

Biotinylation of surface proteins was performed as previously described¹⁹. In brief, cells were incubated with cell-impermeable EZ-Link-NHS-SS-biotin (300 μ g/ml in PBS; Pierce) at 37°C for 2 min and washed with ice-cold 50 mM Tris-buffered saline (TBS) to remove excess biotin. For detecting internalized proteins, cells were washed with PBS and incubated in PBS or serum-free medium supplemented with 0, 0.2, 2, or 20 nM insulin at 37°C to stimulate INSR internalization. After certain time periods, cells were incubated with glutathione solution (50 mM glutathione, 75 mM NaCl, 1 mM EDTA, 1% BSA, 75 mM NaOH) to strip the remaining surface biotin lysed in complete RIPA buffer. Biotinylated internalized proteins in lysates were captured by NeutrAvidin beads (Pierce) and eluted by incubating in Blue Loading Buffer (50 mM DTT, Cell Signaling, Danvers, MA) containing 50 mM DTT at 95°C for 5 min. Internalized INSR in eluent and total INSR in lysates were detected in Western blot analysis.

Plasmids, SNAP-tag cloning and lentivirus infection

CAV1-mRFP and CLTC-mRFP plasmids were generous gifts from Dr. Ivan Robert Nabi, Vancouver, BC, Canada. INSR-A-TagBFP and INSR-B-TagBFP plasmids are previously described⁴.

PCR amplification of DNA fragments was performed using Q5 High-Fidelity DNA Polymerase (Cat. #M0491, New England Biolabs, Ipswich, MA, USA) according to the manufacturer's instructions. Prior ligation with the target vector, all PCR products were subcloned into pJET1.2 vector using the CloneJET PCR Cloning Kit (Cat. #K1232, Thermo Scientific, Waltham, MA, USA) according to the manufacturer's instructions. Subsequently, the PCR products were excised from the pJET1.2 vector, gel purified and ligated with the digested target vector using T4 DNA ligase. To generate INSR-A-EGFP and INSR-B-EGFP plasmids, eGFP was PCR amplified using primers 5'-CGT ACG ATC GAA TTC CCC GTG AGC AAG GGC GAG GAG CTG-3' (forward) and 5'- ACC GGT ATC TGA TCC GGA CTT GTA CAG CTC GTC CAT GCC GA-3' (reverse) and inserted into pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP (previously described⁴) to replace TagRFP by EGFP using BsiWI and AgeI restriction sites.

To generate pLenti-INSR-A-SNAPf and pLenti-INSR-B-SNAPf lentiviral plasmids, SNAP tag was firstly PCR amplified from pSNAPf Vector (Cat. #N9183S, New England Biolabs, Ipswich, MA, USA) using primers 5'-CGT ACG ATC GAA TTC CCC ATG GAC AAA GAC TGC GAA ATG AAG CGC ACC ACC-3' (forward) and 5'- ACC GGT ATC TGA TCC GGA ACC CAG CCC AGG CTT GCC CAG-3' (reverse) and inserted into pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP to replace TagRFP by SNAP-tag using BsiWI and AgeI restriction sites. Subsequently, INSR-A-SNAP and INSR-B-SNAP ORFs were PCR amplified from pcDNA3.1(-)-INSR-A/B-SNAPf plasmids using primers 5'-GAA CCC ACT GCT TAC TGG CT-3' (forward) and 5'-GGT GGT TGT ACA CTA GAA GGC ACA GTC GAG GC -3' (reverse) and inserted into pLenti-C-Myc-DDK-IRES-GFP vector to replace the Myc tag, Flag tag, IRES, and

TurboGFP using NheI and BsrGI restriction sites. The PCR program for this long INSR-SNAPf fragments was optimized to be as follows: initial denaturation 98°C 3min; 30 cycles of 98°C 10 s, 71°C 30 s, 72°C 4min; final extension 72°C 2min; hold 4°C.

SNAP-tag labeling and spinning disk confocal imaging of C2C12 myoblast

Cell non-permeable SNAP-Surface Alexa Fluor 488 substrate (Cat. #S9129S) was dissolved and diluted to 5 µM concentration in serum-free medium containing 0.5% BSA according to the manufacture's instruction. Next, stable C2C12 myoblasts cell line expressing INSR-A-SNAP seeded on glass-bottom dishes (Cat. #P35G-1.5-14-C, MatTek) were incubated in the substrate for 30 min on ice and washed 3 times with serum-free medium (0.5% BSA).

C2C12 dish was then set up on the Spinning disk microscope, applied 37°C FuoroBrite DMEM (Cat. #A1896701, Gibco) serum-free medium containing 0, 0.2 or 20 nM human insulin (Cat. #I9278, SigmaAldrich), and immediately started imaging in a 37°C chamber using a 100x/1.45 oil objective and QuantEM 512SC Photometrics camera on a spinning disk confocal system based on Zeiss Axiovert 200M microscope (LSI Imaging Core, UBC). Time-lapse imaging was conducted at 30 s intervals between 0 – 20 min or at 5 s intervals between 20-30 min.

Immunofluorescence staining

Stable C2C12 cell line seeded on glass-bottom dishes were labeled with cell non-permeable SNAP-Surface Alexa Fluor 488 substrate, incubated with 0, 0.2 or 20 nM human insulin for 30 min, fixed with 4% (w/v) paraformaldehyde (Sigma-Aldrich) for 20 minutes at room temperature, and washed 3 times with PBS. After permeabilized in 0.1% (v/v) Triton X-100 (Sigma-Aldrich) for 15 minutes, primary antibodies targeting Caveolin1 (1:400, Rabbit mAb, Cat. #3267, Cell Signaling) and CLTC heavy chain (1:200, Goat pAb, Cat. #sc-6579, Santa Cruz) were incubated overnight at 4°C.

Coverslips were then washed in PBS, blocked in Dako Protein Block (Cat #X0909, Agilent), and stained by secondary antibodies (1:1000, Cy3 Donkey anti-Rabbit, Cat. #711-165-152, Jackson ImmunoResearch; 1:500, Alexa Fluor 633 Donkey anti-Goat, Cat. # A21082, Invitrogen) diluted in Dako Antibody Diluent (Cat. # S0809, Agilent) for 2 hours at room temperature. After washing in PBS, coverslips were mounted with Prolong Diamond Antifade Mountant (Cat. #P36961, Invitrogen).

STED microscopy and image analysis

Stimulated emission depletion (STED) microscopy was performed with the 100×/1.4 Oil HC PL APO CS2 STED White objective of a Leica TCS SP8 3× STED microscope (Leica, Wetzlar, Germany) equipped with a white light laser, HyD detectors, and Leica Application Suite X (LAS X) software (LSI Imaging Core, Life Sciences Institute, University of British Columbia). Time-gated fluorescence detection was used to further improve lateral resolution. INSR-SNAP labeled by Alexa Fluor 488 was excited at 488 nm and depleted using the 592-nm depletion laser. CAV1 labeled by Cy3 was excited at 555 nm and depleted using the 592-nm depletion laser. CLTC labeled by Alex Fluor 633 was excited at 633 nm and depleted using the 660-nm depletion laser. Sequential acquisition in the order of AF633/Cy3/AF488 was used to avoid cross-talk.

Huygens Professional software (Scientific Volume Imaging, Hilversum, the Netherlands) was used to deconvolve STED images and calculate the Object Pearson correlation coefficient for INSR-SNAP, CAV1, and CLTC.

TIRF microscopy and single particle tracking analysis

Total internal reflection fluorescence (TIRF) microscopy was performed on a Zeiss Axiovert 200M with a 100x Alpha-PlanFluar NA 1.45 oil objective (Zeiss, Germany) and a TIRF laser angle modifier (Zeiss). Cells were kept at 37°C in a temperature-controlled incubation chamber (Harvard Apparatus, Holliston, MA). Images were acquired with a

CoolSNAP HQ2 CCD camera (Photometrics, Tucson, AZ). 405 nm and 561 nm solid-state diode laser system was used to excite TagBFP and mRFP tagged proteins.

The ImageJ distribution Fiji was used to process microscope images⁵⁴. The local background of TIRF images was subtracted using the rolling ball method (20-pixel radius) before single particle tracking.

Single particles (INSR, CAV1, or CLTC) were detected and tracked using Icy bioimaging analysis software version 1.9.8.1⁵⁵. Particles were detected by the UnDecimated Wavelet Transform Detector plugin that was set to detect bright spots over a dark background, exhibit 70% sensitivity to spots <3 pixels, detect particle size between 2 and 3000 pixels. Particles were tracked using the Multiple Hypothesis Tracking method built into the Spot Tracking plugin that was set to recognize particles that exhibited either diffusive or directed motion^{56,57}. Track radius (maximum displacement of the particle track) and lifetime of the particles were exported from the Motion Profile Processor of the Spot Tracking plugin. Diffusion coefficients were calculated in MATLAB as described previously but without corrections for positional errors and blurring⁵⁸. Colocalization of tracks was determined by Track Processor Interaction Analysis (distance threshold 2 pixels) in the Spot Tracking plugin. INSR tracks that colocalized with CAV1 or CLTC tracks for more than 4 continuous frames are determined to be colocalized tracks. The exported time and duration of colocalization were organized in Python and plotted in R studio. Diffusion coefficients, track radius, and the lifetime of tracks were plotted in R studio. Specifically, SuperPlots were used to plot the mean value of the cells and individual tracks of each cell⁵⁹. Code is available at GitHub deposit https://github.com/hcen/INSR_tracking.

Skeletal muscle co-immunoprecipitation

Mice received 1.5U/kg insulin or PBS through intraperitoneal injection. Skeletal muscle (gastrocnemius) was collected after 5 or 10 min of injection. INSR was

immunoprecipitated by incubating 500 µg of protein lysate with 2 µg INSR antibodies (Rabbit pAb, Cat. #sc-711, Santa Cruz) at 4°C overnight. Subsequently, the solution was incubated with PureProteome™ Protein G Magnetic Beads (Cat. # LSKMAGG10, Millipore, Burlington, MA, USA) and washed according to the manufacturer's instructions. INSR protein complexes were eluted from the beads by incubation in Blue Loading Buffer (50 mM DTT, Cell Signaling) at 95°C for 5 min and were analyzed by standard western blot procedures. INSR, CAV1, and CLTC were detected by the following Cell Signaling primary antibodies: INSRβ (1:1000, Mouse mAb, Cat. #3020S), CLTC (1:1000, Rabbit mAb, Cat. #4796), Caveolin1 (1:1000, Rabbit mAb, Cat. #3267), and HRP-conjugated VeriBlot secondary antibodies (1:200, anti-mouse Cat. #ab131368, anti-rabbit ab131366, Abcam) that do not bind to IgG heavy and light chains to reduce background noise.

Immunoprecipitation-Mass Spectrometry

C2C12 myoblasts were serum-starved for 6 hours and were given 0 or 2 nM insulin for 15 min before collecting their lysates in RIPA buffer. INSRα antibody (NNC0276-3000, a gift from NovoNordisk compound sharing program) that does not cross-react with IGF1R²³ was covalently conjugated to Dynabeads M-270 Epoxy beads (Cat. #14301, Invitrogen) to immunoprecipitate INSR according to the manufacture's protocol. INSR protein complexes were eluted from the beads by incubation in Blue Loading Buffer (50 mM DTT, Cell Signaling) at 95°C for 5 min.

Eluted co-immunoprecipitation samples were run into 10% SDS PAGE gel for clean-up procedure, fixed, stained, and destained in water. Each of the samples was excised, reduced, alkylated, and digested according to standard in-gel digest protocols⁶⁰. Extracted peptide samples were then cleaned up via STAGE-tip purification, as previously described⁶¹.

Samples were reconstituted in 2% ACN, 0.5% formic acid, and the peptides were analyzed using a quadrupole – time of flight mass spectrometer (Impact II; Bruker

Daltonics) on-line coupled to an EasyLC 1000 HPLC (ThermoFisher Scientific) using a Captive spray nanospray ionization source (Bruker Daltonics) including Aurora Series Gen2 (CSI) analytical column, (25cm x 75µm 1.6µm FSC C18, with Gen2 nanoZero and CSI fitting; Ion Opticks, Parkville, Victoria, Australia), and a µ-Precolumn, 300 µm ID x 5 mm, C18 PepMap, 5 µm, 100 Å, Thermo Scientific, Waltham, Massachusetts, United States). Samples were reconstituted in (0.1% aqueous formic acid and 2 % acetonitrile in water) and 1ug of sample was loaded, injected in triplicates. A standard 90 min reversed-phase separation was employed, using water:acetonitrile gradients.

LC and MS acquisition parameters were standard for peptide sequencing. Acquired data were then searched by MaxQuant (V1.6.7.) against the Uniprot protein database for Mouse C2C12 cell line, and LFQ intensities extracted and normalized using the MaxLFQ algorithm, with 20ppm and 30ppm mass accuracies for precursor and product ion masses, respectively, and a 1% false discovery rate cut-off. Differential expression was determined by comparing the mean protein intensities of the triplicate injections between the groups. Significance ($p < 0.05$) was calculated via t-test using default parameters in Perseus (v1.6.14.0).

Statistics

Data were presented as mean $\pm$ SEM in addition to the individual data points. A significance level of $p < 0.05$ was used throughout. Student's t-tests were used for comparisons between two groups. Comparisons between more than two groups were performed using multiple comparison p-values were adjusted using the Tukey method. Comparisons between multiple groups and two categorical variables were performed using Two-way ANOVA followed by Tukey post hoc test. Statistical analysis above was performed using GraphPad Prism (Version 9.0.2). Two-sided Kolmogorov-Smirnov Tests were used for comparing cumulative probability, which was performed with `ks.test()` function in R.

561

562 References

- 563 1. Knutson VP. Cellular trafficking and processing of the insulin receptor. *FASEB J.*
564 1991;5(8):2130-2138.
- 565 2. Kahn CR, Crettaz M. Insulin receptors and the molecular mechanism of insulin
566 action. *Diabetes Metab Rev.* 1985;1(1-2):5-32.
- 567 3. Bevan AP, Burgess JW, Drake PG, Shaver A, Bergeron JJ, Posner BI. Selective
568 activation of the rat hepatic endosomal insulin receptor kinase. Role for the
569 endosome in insulin signaling. *The Journal of biological chemistry.*
570 1995;270(18):10784-10791.
- 571 4. Boothe T, Lim GE, Cen H, et al. Inter-domain tagging implicates caveolin-1 in
572 insulin receptor trafficking and Erk signaling bias in pancreatic beta-cells. *Mol*
573 *Metab.* 2016;5(5):366-378.
- 574 5. Trischitta V, Gullo D, Squatrito S, Pezzino V, Goldfine ID, Vigneri R. Insulin
575 Internalization into Monocytes Is Decreased in Patients with Type II Diabetes
576 Mellitus. 1986;62:522-528.
- 577 6. Trischitta V, Brunetti A, Chiavetta A, Benzi L, Papa V, Vigneri R. Defects in
578 insulin–receptor internalization and processing in monocytes of obese subjects
579 and obese NIDDM patients. 1989;38:1579–1584.
- 580 7. Jochen AL, Berhanu P, Olefsky JM. Insulin internalization and degradation in
581 adipocytes from normal and type II diabetic subjects. *J Clin Endocrinol Metab.*
582 1986;62(2):268-274.
- 583 8. Rask-Madsen C, Kahn CR. Tissue-specific insulin signaling, metabolic syndrome,
584 and cardiovascular disease. *Arterioscler Thromb Vasc Biol.*
585 2012;32(9):2052-2059.
- 586 9. Guo S. Insulin signaling, resistance, and the metabolic syndrome: insights from
587 mouse models into disease mechanisms. *The Journal of endocrinology.*
588 2014;220(2):T1-T23.
- 589 10. Jochen A, Hays J, Lee M. Kinetics of insulin internalization and processing in
590 adipocytes: effects of insulin concentration. *J Cell Physiol.* 1989;141(3):527-534.
- 591 11. McClain DA, Olefsky JM. Evidence for two independent pathways of
592 insulin-receptor internalization in hepatocytes and hepatoma cells. *Diabetes.*
593 1988;37(6):806-815.
- 594 12. Ceresa BP, Kao AW, Santeler SR, Pessin JE. Inhibition of clathrin-mediated
595 endocytosis selectively attenuates specific insulin receptor signal transduction
596 pathways. *Mol Cell Biol.* 1998;18(7):3862-3870.
- 597 13. Carpentier JL, Van Obberghen E, Gorden P, Orci L. Surface redistribution of
598 125I-insulin in cultured human lymphocytes. *J Cell Biol.* 1981;91(1):17-25.

- 599 14. Paccaud JP, Siddle K, Carpentier JL. Internalization of the human insulin receptor.
600 The insulin-independent pathway. *The Journal of biological chemistry*.
601 1992;267(18):13101-13106.
- 602 15. Bickel PE. Lipid rafts and insulin signaling. *Am J Physiol Endocrinol Metab*.
603 2002;282(1):E1-E10.
- 604 16. Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in
605 insulin signaling and diabetes. *Am J Physiol Endocrinol Metab*.
606 2003;285(6):E1151-1160.
- 607 17. Tseng LT, Lin CL, Tzen KY, Chang SC, Chang MF. LMBD1 protein serves as a
608 specific adaptor for insulin receptor internalization. *The Journal of biological*
609 *chemistry*. 2013;288(45):32424-32432.
- 610 18. Gautier A, Juillerat A, Heinis C, et al. An engineered protein tag for multiprotein
611 labeling in living cells. *Chem Biol*. 2008;15(2):128-136.
- 612 19. Cen H, Botezelli JD, Johnson JD. Modulation of Insr and insulin receptor signaling
613 by hyperinsulinemia in vitro and in vivo. *Biorxiv*. 2021.
- 614 20. Jans DA. Lateral Mobility of Polypeptide Hormone Receptors and GTP-Binding
615 Proteins. In: *The Mobile Receptor Hypothesis: The Role of Membrane Receptor*
616 *Lateral Movement in Signal Transduction*. Boston, MA: Springer US; 1997:83-115.
- 617 21. Kabayama K, Sato T, Saito K, et al. Dissociation of the insulin receptor and
618 caveolin-1 complex by ganglioside GM3 in the state of insulin resistance. *Proc*
619 *Natl Acad Sci U S A*. 2007;104(34):13678-13683.
- 620 22. Ikonen E, Vainio S. Lipid microdomains and insulin resistance: is there a
621 connection? *Sci STKE*. 2005;2005(268):pe3.
- 622 23. Ørstrup LH, Slaaby R, Rasch MG, et al. Cross-species reactive monoclonal
623 antibodies against the extracellular domains of the insulin receptor and IGF1
624 receptor. *J Immunol Methods*. 2019;465:20-26.
- 625 24. Benyoucef S, Surinya KH, Hadaschik D, Siddle K. Characterization of insulin/IGF
626 hybrid receptors: contributions of the insulin receptor L2 and Fn1 domains and the
627 alternatively spliced exon 11 sequence to ligand binding and receptor activation.
628 *Biochem J*. 2007;403(3):603-613.
- 629 25. Moxham CP, Duronio V, Jacobs S. Insulin-like growth factor I receptor
630 beta-subunit heterogeneity. Evidence for hybrid tetramers composed of
631 insulin-like growth factor I and insulin receptor heterodimers. *The Journal of*
632 *biological chemistry*. 1989;264(22):13238-13244.
- 633 26. Shu L, Lee L, Chang Y, et al. Caveolar structure and protein sorting are
634 maintained in NIH 3T3 cells independent of glycosphingolipid depletion. *Arch*
635 *Biochem Biophys*. 2000;373(1):83-90.
- 636 27. DiGiacomo V, Meruelo D. Looking into laminin receptor: critical discussion
637 regarding the non-integrin 37/67-kDa laminin receptor/RPSA protein. *Biol Rev*
638 *Camb Philos Soc*. 2016;91(2):288-310.

- 639 28. Zoubeidi A, Zardan A, Wiedmann RM, et al. Hsp27 promotes insulin-like growth
640 factor-I survival signaling in prostate cancer via p90Rsk-dependent
641 phosphorylation and inactivation of BAD. *Cancer Res.* 2010;70(6):2307-2317.
- 642 29. Muranova LK, Shatov VM, Bukach OV, Gusev NB. Cardio-Vascular Heat Shock
643 Protein (cvHsp, HspB7), an Unusual Representative of Small Heat Shock Protein
644 Family. *Biochemistry (Mosc).* 2021;86(Suppl 1):S1-S11.
- 645 30. Yuan H, Wang T, Niu Y, Liu X, Fu L. AMP-activated protein kinase-mediated
646 expression of heat shock protein beta 1 enhanced insulin sensitivity in the skeletal
647 muscle. *FEBS Lett.* 2017;591(1):97-108.
- 648 31. Chidlow JH, Sessa WC. Caveolae, caveolins, and cavins: complex control of
649 cellular signalling and inflammation. *Cardiovasc Res.* 2010;86(2):219-225.
- 650 32. Capozza F, Cohen AW, Cheung MW, et al. Muscle-specific interaction of caveolin
651 isoforms: differential complex formation between caveolins in fibroblastic vs.
652 muscle cells. *Am J Physiol Cell Physiol.* 2005;288(3):C677-691.
- 653 33. Chang M, Kwon M, Kim S, et al. Aptamer-based single-molecule imaging of
654 insulin receptors in living cells. *J Biomed Opt.* 2014;19(5):051204.
- 655 34. Hubert M, Larsson E, Lundmark R. Keeping in touch with the membrane; protein-
656 and lipid-mediated confinement of caveolae to the cell surface. *Biochem Soc*
657 *Trans.* 2020;48(1):155-163.
- 658 35. Martins AS, Ordóñez JL, Amaral AT, et al. IGF1R signaling in Ewing sarcoma is
659 shaped by clathrin-/caveolin-dependent endocytosis. *PLoS One.*
660 2011;6(5):e19846.
- 661 36. Pandini G, Frasca F, Mineo R, Sciacca L, Vigneri R, Belfiore A. Insulin/insulin-like
662 growth factor I hybrid receptors have different biological characteristics depending
663 on the insulin receptor isoform involved. *The Journal of biological chemistry.*
664 2002;277(42):39684-39695.
- 665 37. Tomas A, Futter CE, Eden ER. EGF receptor trafficking: consequences for
666 signaling and cancer. *Trends Cell Biol.* 2014;24(1):26-34.
- 667 38. Kammanadiminti SJ, Chadee K. Suppression of NF-kappaB activation by
668 *Entamoeba histolytica* in intestinal epithelial cells is mediated by heat shock
669 protein 27. *The Journal of biological chemistry.* 2006;281(36):26112-26120.
- 670 39. Ku HC, Chang HH, Liu HC, et al. Green tea (-)-epigallocatechin gallate inhibits
671 insulin stimulation of 3T3-L1 preadipocyte mitogenesis via the 67-kDa laminin
672 receptor pathway. *Am J Physiol Cell Physiol.* 2009;297(1):C121-132.
- 673 40. Zhao WQ, Chen GH, Chen H, et al. Secretion of Annexin II via activation of insulin
674 receptor and insulin-like growth factor receptor. *The Journal of biological*
675 *chemistry.* 2003;278(6):4205-4215.
- 676 41. Biener Y, Feinstein R, Mayak M, Kaburagi Y, Kadowaki T, Zick Y. Annexin II is a
677 novel player in insulin signal transduction. Possible association between annexin
678 II phosphorylation and insulin receptor internalization. *The Journal of biological*
679 *chemistry.* 1996;271(46):29489-29496.

680 42. Rescher U, Ludwig C, Konietzko V, Kharitononkov A, Gerke V. Tyrosine
681 phosphorylation of annexin A2 regulates Rho-mediated actin rearrangement and
682 cell adhesion. *J Cell Sci.* 2008;121(Pt 13):2177-2185.

683 43. Caron D, Boutchueng-Djidjou M, Tanguay RM, Faure RL. Annexin A2 is
684 SUMOylated on its N-terminal domain: regulation by insulin. *FEBS Lett.*
685 2015;589(9):985-991.

686 44. Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell.*
687 2010;141(7):1117-1134.

688 45. de Graauw M, Cao L, Winkel L, et al. Annexin A2 depletion delays EGFR
689 endocytic trafficking via cofilin activation and enhances EGFR signaling and
690 metastasis formation. *Oncogene.* 2014;33(20):2610-2619.

691 46. Behrends M, Engmann O. Linker histone H1.5 is an underestimated factor in
692 differentiation and carcinogenesis. *Environ Epigenet.* 2020;6(1):dvaa013.

693 47. Vicidomini G, Bianchini P, Diaspro A. STED super-resolved microscopy. *Nat*
694 *Methods.* 2018;15(3):173-182.

695 48. Fish KN. Total internal reflection fluorescence (TIRF) microscopy. *Curr Protoc*
696 *Cytom.* 2009;Chapter 12:Unit12.18.

697 49. Le PU, Guay G, Altschuler Y, Nabi IR. Caveolin-1 is a negative regulator of
698 caveolae-mediated endocytosis to the endoplasmic reticulum. *Journal of*
699 *Biological Chemistry.* 2002;277(5):3371-3379.

700 50. Humphries AC, Way M. The non-canonical roles of clathrin and actin in pathogen
701 internalization, egress and spread. *Nat Rev Microbiol.* 2013;11(8):551-560.

702 51. Rodal SK, Skretting G, Garred O, Vilhardt F, van Deurs B, Sandvig K. Extraction
703 of cholesterol with methyl-beta-cyclodextrin perturbs formation of clathrin-coated
704 endocytic vesicles. *Mol Biol Cell.* 1999;10(4):961-974.

705 52. Parpal S, Karlsson M, Thorn H, Strålfors P. Cholesterol depletion disrupts
706 caveolae and insulin receptor signaling for metabolic control via insulin receptor
707 substrate-1, but not for mitogen-activated protein kinase control. *The Journal of*
708 *biological chemistry.* 2001;276(13):9670-9678.

709 53. Hussain KM, Leong KL, Ng MM, Chu JJ. The essential role of clathrin-mediated
710 endocytosis in the infectious entry of human enterovirus 71. *The Journal of*
711 *biological chemistry.* 2011;286(1):309-321.

712 54. Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for
713 biological-image analysis. *Nat Methods.* 2012;9(7):676-682.

714 55. de Chaumont F, Dallongeville S, Chenouard N, et al. Icy: an open bioimage
715 informatics platform for extended reproducible research. *Nat Methods.*
716 2012;9(7):690-696.

717 56. Chenouard N, Bloch I, Olivo-Marin JC. Multiple hypothesis tracking for cluttered
718 biological image sequences. *IEEE Trans Pattern Anal Mach Intell.*
719 2013;35(11):2736-3750.

57. Chenouard N, Smal I, de Chaumont F, et al. Objective comparison of particle tracking methods. *Nat Methods*. 2014;11(3):281-289.
58. Abraham L, Lu HY, Falcão RC, et al. Limitations of Qdot labelling compared to directly-conjugated probes for single particle tracking of B cell receptor mobility. *Sci Rep*. 2017;7(1):11379.
59. Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating reproducibility and variability in cell biology. *J Cell Biol*. 2020;219(6).
60. Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal Chem*. 1996;68(5):850-858.
61. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat Protoc*. 2007;2(8):1896-1906.

Figure legends

Figure 1. Biochemical and live-cell imaging characterization of the effects of insulin on INSR.

(A) Internalized to total INSR ratio was quantified using surface biotinylation in undifferentiated C2C12 myoblasts were incubated in PBS (no insulin) or serum-free DMEM containing 0, 0.2, 2, or 20 nM insulin for 15 min. (B) Western blot showing SNAP-tagged INSR expressed from lentiviral vector in comparison to wild-type INSR or in C2C12 myoblasts. (C,D). Qualitative snapshots of surface INSR-A-SNAP labeled by Surface Alexa Flour 488 cell non-permeable dye in 0, 0.2 or 20 nM insulin conditions were imaged in live undifferentiated C2C12 myoblasts from (C) 0 - 15 min or (D) after 20 min using a spinning disk confocal microscope. Time-lapse images of INSR vesicle interactions starting from the selected subregions (white squares) of snapshots in are shown in the *insets*. (n=3 cells)

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

(A) Experimental design of TIRF microscopy of same cells at 3 timepoints with 2 nM

insulin or no insulin. **(B)** Representative TIRF image of INSR-A-EGFP puncta, binary image of detected INSR spots, and INSR tracks. **(C)** Diffusion coefficient of INSR-A-EGFP tracks in control or insulin group. Data is plotted as SuperPlot with bigger dots showing mean values of the cells and smaller dots showing individual INSR tracks of the cells. **(D)** The cumulative probability shows the distribution of diffusion coefficient in **(C)**. **(E)** Track radius of INSR-A-EGFP tracks in control or insulin group. **(F)** The cumulative probability shows the distribution of diffusion coefficient in **(E)**. (n=3 cells in control group, n=9 cells in insulin group)

Figure 3. Identification of INSR interactors.

(A) Proteins that are detected in INSR immunoprecipitation mass-spectrometry, and significantly different between wild-type and INSR knockout groups. Proteins in green area were specific interactors that had little detection in knockout cells. **(C)** AKT and ERK phosphorylation in. **(C)** Western blot of mice skeletal muscle lysates after insulin injection. Phospho-AKT (Ser473) to total AKT ratio or phospho-ERK1/2 to total ERK ratio were quantified. (n=6) **(D)** Co-immunoprecipitation of skeletal muscle INSR after PBS or insulin injection. Interacting CLTC to INSR ratio or CAV3 to INSR ratio were quantified. (n=7-10)

Figure 4. Colocalization and interaction between INSR, CAV1 and CLTC under different insulin concentrations.

(A) Representative STED microscopy images of C2C12 myoblasts expressing INSR-A-SNAP (surface labeled) that were fixed after stimulated with 0 nM, 0.2 nM, or 20 nM insulin for 30 min and stained for CAV1 and CLTC (Scale bar = 5 μ m). **(B,C)** Colocalizations between INSR-A-SNAP, CAV1, and CLTC were quantified by Object Pearson Coefficient. Data are plotted to show differences between insulin concentrations **(B)** or between protein pairs **(C)**. (n=7-9 images, 1-4 cells per image. *p<0.05, Tukey's

multiple comparison after 2-ANOVA. Box represents median and 25th to 75th percentiles.)

Figure 5. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

(A,B) TIRF microscopy showed colocalization between INSR-B-TagBFP and **(A)** CAV1-mRFP or **(B)** CLTC-mRFP. **(C,D)** The associated fluorescent intensity of representative INSR-B-TagBFP and colocalized **(C)** CAV1-mRFP or **(D)** CLTC-mRFP. **(E)** Diffusion coefficient of INSR-B-TagBFP tracks in cells expressing CAV1-mRFP under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin (n=2 cells, 324 tracks) or expressing CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2 nM insulin (n=2 cells, 146 tracks). Data is plotted as SuperPlot, with bigger dots showing mean values of the cells and smaller dots showing individual INSR tracks of the cells. The cumulative probability shows the distribution of the diffusion coefficient. **(F)** Track radius of INSR-B-TagBFP tracks in cells expressing CAV1-mRFP under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin (n=2 cells, 324 tracks) or expressing CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2 nM insulin (n=2 cells, 146 tracks). The cumulative probability shows the distribution of the track radius. **(G)** Track radius of INSR-B-TagBFP tracks in cells expressing CAV1-mRFP under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin (n=2 cells, 324 tracks) or expressing CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2 nM insulin (n=2 cells, 146 tracks). The cumulative probability shows the distribution of the diffusion coefficient. (# effect of co-expression, 2-ANOVA.) Experiments shown were conducted in Ringer's buffer.

Figure 6. Analysis of INSR tracks colocalized or not colocalized with CAV1 or CLTC.

(A,B) TIRF microscopy showed colocalization between INSR-B-TagBFP and **(A)** CAV1-mRFP or **(B)** CLTC-mRFP membrane. **(A)** INSR tracks that colocalized with CAV1 under 0 or 2 nM insulin (referred to as CAV1.Y.0nM or CAV1.Y.2nM below). **(B)** INSR

805 tracks that colocalized with CLTC under 0 or 2 nM insulin (referred to as CLTC.Y.0nM or
806 CLTC.Y.2nM below). **(C)** Diffusion coefficient of INSR-B-TagBFP tracks that did not
807 colocalize with CAV1-mRFP tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin
808 (CAV1.N.2nM) or colocalized with CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM)
809 or 2 nM insulin (CAV1.Y.2nM). The cumulative probability shows the distribution of
810 diffusion coefficient. **(D)** Diffusion coefficient of INSR-B-TagBFP tracks that did not
811 colocalize with CLTC-mRFP tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin
812 (CLTC.N.2nM) or colocalized with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM)
813 or 2 nM insulin (CLTC.Y.2nM). The cumulative probability shows the distribution of
814 diffusion coefficient. **(E)** Track radius of INSR-B-TagBFP tracks that did not colocalize
815 with CAV1-mRFP tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM)
816 or colocalized with CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin
817 (CAV1.Y.2nM). The cumulative probability shows the distribution of track radius **(F)** Track
818 radius of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP tracks under 0
819 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized with CLTC-mRFP
820 tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM). The cumulative
821 probability shows the distribution of track radius. **(G)** Lifetime of INSR-B-TagBFP tracks
822 that did not colocalize with CAV1-mRFP tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM
823 insulin (CAV1.N.2nM) or colocalized with CAV1-mRFP tracks under 0 nM insulin
824 (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM). The cumulative probability shows the
825 distribution of lifetime. **(H)** Lifetime of INSR-B-TagBFP tracks that did not colocalize with
826 CLTC-mRFP tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or
827 colocalized with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin
828 (CLTC.Y.2nM). The cumulative probability showing the distribution of lifetime.
829 (CAV1.N.0nM, n=2 cells, 125 tracks; CAV1.N.2nM, n=2 cells, 96 tracks; CAV1.Y.0nM,
830 n=2 cells, 235 tracks; CAV1.Y.2nM, n=2 cells, 228 tracks; CLTC.N.0nM, n= 4 cells, 597
831 tracks; CLTC.N.2nM, n= 2 cells, 91 tracks; CLTC.Y.0nM, n= 4 cells, 320 tracks;

CLTC.Y.2nM, n= 2 cells, 55 tracks. # effect of colocalization, \$ effect of insulin, % interaction of the 2 variables, 2-ANOVA.) Experiments shown were conducted in Ringer's buffer.

Figure 7. Summary of the effects of CAV1 and CLTC on INSR tracks. Higher insulin (20 nM) promoted INSR and CAV1 colocalization, while lower insulin (0.2 nM) promoted INSR and CLTC colocalization. INSR tracks had lower diffusion coefficient and track radius in CAV1-overexpressing cells than CLTC-overexpressing cells. Within the same cells, INSR tracks colocalized with CAV1 had longer lifetime and larger track radius than non-colocalized tracks. INSR tracks colocalized with CLTC had longer lifetime than non-colocalized tracks, which is further increased by insulin.

Figure 1. Biochemical and live-cell imaging characterization of the effects of insulin on INSR.

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

A

B

C

D

Figure 3. Identification of INSR interactors.

Figure 4. Colocalization and interaction between INSR, CAV1 and CLTC under different insulin concentrations.

Figure 5. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

Figure 6. Analysis of INSR tracks colocalized or not colocalized with CAV1 or CLTC.

Figure 7. Summary of the effects of CAV1 and CLTC on INSR tracks.