Pain and the Emotional Brain: Affective Rather than Cognitive Processes Drive the Cortical Encoding of Pain

Astrid Mayr¹, Anne Stankewitz¹, Stephanie Irving¹, Viktor Witkovsky³, Enrico Schulz^{1,2,4}

¹ Department of Neurology, Ludwig-Maximilians-Universität München, Munich, Germany

² Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK

³ Department of Theoretical Methods, Institute of Measurement Science, Slovak Academy of Sciences, Bratislava, Slovak Republic

⁴ Department of Medical Psychology, Ludwig-Maximilians-Universität München, Munich, Germany

Corresponding author:

Enrico Schulz

Ludwig-Maximilians-Universität München

Neurologische Klinik und Poliklinik

A: Fraunhoferstr. 20, 82152 Martinsried, Germany

O: +49 89 4400 74826

E: es@pain.sc

ABSTRACT

The experience of pain has been dissociated into two interwoven aspects: a sensory-discriminative aspect assessed in ratings of pain intensity and an affective-motivational aspect assessed in ratings of unpleasantness. In a pain attenuation paradigm, participants were asked to evaluate applied cold pain. The majority of the trials showed a distinct rating: Some trials were rated higher for unpleasantness, others were rated higher for intensity. Using linear mixed effect models on single trials, we related the variable difference between unpleasantness and intensity ratings to functional MRI data. The direct comparison revealed a stronger relationship between cortical data and pain ratings for unpleasantness. No region showed a stronger effect for pain intensity. The present study underlines the importance of the emotional-affective aspects of pain-related cortical processes in the brain. These findings reflect the biological function of the pain system to prevent harm and to preserve the physical integrity of the body.

INTRODUCTION

The experience of pain has been dissociated into sensory-discriminative and affective-motivational aspects ¹, which are assessed as subjective ratings of pain intensity and pain unpleasantness ²⁻⁴. Both descriptors of pain share a conceptual core variance and are therefore often highly correlated: very intense pain is associated with higher levels of unpleasantness ^{5,6}. The experience of pain unpleasantness is considered to exhibit a particularly strong influence on well-being ^{7,8}, as it has been found to be tightly connected to pain catastrophising ⁷.

Studies exploring the cortical underpinnings of pain unpleasantness are scarce and non-specific as they implicitly include the shared variance with pain intensity ^{9,10}. The conceptual and empirical proximity of both aspects has made a differential investigation of pain intensity and pain unpleasantness challenging. A series of early PET studies attempted to independently modulate pain intensity and pain unpleasantness using hypnosis. For pain unpleasantness, the group reported a specific modulation in the anterior cingulate cortex (ACC) ^{6,11}. This region was also active to a lesser extent in the low-unpleasant condition, but was not found in any of the conditions of the intensity modulation. This absence of activity in this ACC region in the pain intensity study, as well as the lack of a direct comparison with pain unpleasantness, leaves some questions regarding distinct cortical processing of pain intensity and pain affect ¹².

Nonetheless, despite a high correlation between both subjective pain aspects, it is indeed possible to dissociate the cortical underpinnings of pain intensity and pain unpleasantness in a within-subject design without the need for any modulatory intervention. To this end, we aimed to investigate the brain regions that differentially process pain intensity and pain unpleasantness. As a major advantage, both aspects are explored in relation to the very same underlying cortical data; each trial has been evaluated for either descriptor of the subjective experience of pain. The majority of the trials indeed differed and enabled us to elucidate which aspect of pain perception is more deeply rooted in the human brain.

RESULTS

Behavioural data

The single-trial pain ratings varied substantially across trials; we recorded an average of 34 (± 14) for pain intensity and of 31 (± 17) for pain unpleasantness. Both aspects of pain perception were highly correlated ($r=0.86$, $p<0.01$, Figure 1). Nonetheless, the majority of the trials exhibited a distinct rating for intensity and unpleasantness (75%, Figure 1).

Figure 1. The left side (A) shows the correlation between pain intensity [I] and pain unpleasantness [U]. The correlation coefficient is $r=0.86$. Nevertheless, the ratings for both aspects were identical only for a minority of the trials. The histogram (B) shows the difference between pain intensity ratings and pain unpleasantness ratings; most trials (75%) exhibited different ratings. For some trials, pain intensity was rated higher (negative Δ), for others pain unpleasantness was rated higher (positive Δ).

Imaging data

The results indicate a stronger positive relationship for *pain unpleasantness* in regions that are known to be involved in pain processing: the bilateral anterior and posterior insula, postcentral regions, regions in the frontal cortex, the paracingulate regions, and the nucleus ruber. Additionally, we found results for the negative gradient, indicating a more positive relationship for *pain intensity* for the retrosplenial cortex, the precuneus, the posterior cingulate cortex, the occipital cortex, the amygdala, and the middle frontal gyrus. However, these negative results occur exclusively in brain regions that are suppressed in response to pain. Therefore, they reflect a stronger negative relationship between pain ratings and brain activity for unpleasantness (Figure 2).

Please note that the differences are in a gradual fashion; due to the fact that both aspects of pain evaluation are highly correlated, they are considered to be commonly processed in the same cortical regions, but to a different extent. All trials on intensity and unpleasantness are relying on the same cortical responses.

Figure 2. Several cortical regions reflect the gradual differences between pain intensity and pain unpleasantness. Warm colours indicate regions that show a more positive relationship for pain unpleasantness; cold colours indicate brain regions that show a stronger negative

relationship for pain unpleasantness. In order to better separate the different regions we increased the statistical threshold beyond the significance level ($t > 4.9$) for the positive effects.

max t-value	x	y	z	region
-8.45	1	-69	25	Retrosplenial Cortex, Precuneus, Posterior Cingulate Cortex
12.4	42	13	12	Insular Cortex, Inferior Frontal Cortex, Striatum
12.5	4	16	48	Superior Frontal Gyrus, Juxtapositional Gyrus, Paracingulate Gyrus
12.3	-32	13	3	Insular Cortex, Striatum
10.8	-33	-50	46	Superior Parietal Lobe
10.6	40	-44	46	Superior Parietal Lobe
-8.34	26	31	43	Middle Frontal Gyrus
10.2	-24	-5	56	Superior Frontal Gyrus
9.73	38	47	20	Frontal Pole
7.08	-1	-22	28	Posterior Cingulate Cortex
9.1	29	-1	55	Middle Frontal Gyrus
9.37	-37	-23	56	Postcentral Gyrus
-5.35	4	59	-5	Frontal Pole
9.65	-47	3	34	Precentral Gyrus
-5.87	58	-55	4	Middle Temporal Gyrus
10.2	8	-54	-10	Cerebellum
-6.74	26	-5	-18	Amygdala
8.79	-42	28	30	Middle Frontal Gyrus
6.86	-9	-22	-7	Brainstem, ncl. ruber

Table 1. Differences in the encoding of pain intensity and pain unpleasantness. Positive effects indicate a stronger positive relationship for unpleasantness; negative relationships indicate a stronger negative relationship for unpleasantness (all $p < 0.01$, PALM corrected).

DISCUSSION

In the present study, we were investigating the cortical underpinnings of pain intensity and pain unpleasantness. Despite the fact that each shares the same fundamental core, we took advantage of the sufficient variability, which allowed us to investigate cortical functions related to the subjective experience of pain in a within-subject design at single-trial level. Overall, we revealed a stronger relationship for the pain-related cortical

processes regarding the affective aspect compared to the sensory-discriminative aspect, suggesting that the emotional-affective aspect of pain is more deeply rooted in the human brain.

Regions that increased activity in response to pain had a more positive effect for unpleasantness, and regions that exhibited decreased activity showed a more negative effect for unpleasantness. Consequently, unpleasantness ratings may reflect the more direct and intuitive processing of emotions in the human brain rather than the more complex and “minded” evaluation of pain intensity. Due to the methodological differences of previous work and their inherent weaknesses, the present study is not comparable to previous imaging studies^{11,13}. However, our results underline other research indicating the preferential processing of pain-related emotions in the human brain^{14,15}. It has been suggested that the feeling of unpleasantness belongs to the initial part of the noxious sensation and occurs before any conscious and cognitive processing of the sensation¹⁶. Therefore, the stronger effect of pain unpleasantness may better reflect the biological function of the pain system to prevent harm and to preserve the physical integrity of the body¹⁷.

Indeed, there is a large framework of research that supports the notion that emotional processes are tightly associated with pain processing in the brain. Higher pain affect has been found to increase autonomic responses, such as skin conductance^{18,19}, which in turn were found to encode pain-related cortical responses^{20,21}. Potentially emotion-related autonomic responses were found to be more tightly related to cortical activity than ratings of pain intensity²⁰.

Our study complements previous findings by Zeidan and colleagues who shed light on the individual abilities to modulate pain intensity and pain unpleasantness through meditation¹³. The better performing participants were able to make use of specific subunits of the pain modulation system. The participants that exhibited a better performance in the attenuation of pain unpleasantness, modulated the activity of the orbitofrontal cortex¹³. This region that has been suggested to serve as a major hub of emotional modulation²². In a similar vein, the hub that modulates the ability to attenuate pain intensity through meditation could be located in the anterior insula¹³. Depending on the pain attenuation strategy, we would assume the existence of further cortical hubs that are differently utilised by individual subjects and which alter pain intensity and pain unpleasantness^{23,24}.

Taken together, the present study shows that pain unpleasantness is more tightly related to cortical processing than the cognitive evaluation of pain intensity. These findings generalise to all brain regions that exhibit pain-related responses, irrespective of whether they increase or decrease their activity. Consequently, studies utilising the evaluation of painful stimulation are suggested to assess pain unpleasantness rather than pain intensity. These suggestions may also be expanded to investigations of endogenous pain in chronic patients. Future studies need to shed further light on either aspect of pain evaluation under specific psychological conditions, and we hypothesise temporal differences in the cascade of pain-related cortical processes: the cortical processes of pain unpleasantness evaluation are processed faster and may occur before the cognitive and “minded” cortical processes of pain intensity evaluation.

ACKNOWLEDGEMENTS

This project was funded by the “Deutsche Forschungsgemeinschaft” (DFG, SCHU 2879/1-2”).

DECLARATION OF INTERESTS

All authors declare no conflict of interest.

DATA AVAILABILITY

The data will be made available via the Open Science Framework (<https://osf.io/tbc2u/>).

CODE AVAILABILITY

No custom code has been developed for this study.

MATERIALS AND METHODS

Subjects. 20 healthy subjects (16 female/4 male, mean age 27 ± 5 years) were included in the study. All subjects gave written informed consent. The study was approved by the Medical Sciences Interdivisional Research Ethics Committee of the University of Oxford and conducted in conformity with the Declaration of Helsinki.

The experimental procedure has been described in detail in our previous publications^{23,24}. The experiment consisted of four conditions across four separate blocks, where each block comprised 12 trials from the same condition. In all conditions and trials, the subjects received cold pain stimuli on the dorsum of their left hand delivered by a thermode (Pathway II; Medoc Ltd, Israel). The first condition of the experiment was always the unmodulated pain condition. The subsequent three conditions were counterbalanced blocks of pain attenuation: (A) an attentional shift, (B) an imaginal strategy, and (C) a non-imaginal reinterpretation. After each trial, the subjects were prompted to rate pain intensity and pain unpleasantness within 10 s. The scale ranged between 0 and 100 in steps of 5 points. The endpoints of the scale were determined as no pain (0) and the maximum pain the subjects were willing to tolerate (100). There were a total of 48 trials during fMRI recording. The 40 s of painful stimulation were preceded by a rest period of 10 s at 38 °C thermode temperature.

Supplementary Figure 1: Schematic time course of a single trial. After 10 s of rest, painful stimulation was applied for 40 s. The pain was rated regarding intensity ([I], 10 s) and unpleasantness ([U], 10 s).

Data Acquisition. Imaging data were acquired on a 7 tesla Siemens MRI scanner using parallel image acquisition (GRAPPA, factor = 2). In order to cover the whole brain, each of the 1768 functional echo-planar imaging (EPI) volumes comprised 34 axial slices of 2 mm thickness and 2 × 2 mm in-plane resolution with a 1 mm gap between slices. The repetition time (TR) was 1.96 s, the echo time (TE) was 25 ms (flip angle 90°), the field of view (FOV) was 220 × 220 mm, and the matrix size was 110 × 110 pixels. A T1-weighted structural image (isotropic 1 mm³ voxel) was acquired for the registration of the EPI to the MNI (Montreal Neurological Institute) template.

Imaging analysis. The data were pre-processed and analysed with FSL²⁵ and Matlab (version R2020a, The Mathworks, USA). The preprocessing consisted of brain extraction, high-pass filtering with a frequency cutoff of 1/90 Hz, a correction for head motion during scanning, spatial normalisation to the MNI template, and spatial smoothing (6 mm FWHM). The data were further semi-automatically cleaned of artefacts with MELODIC²⁶. Beta coefficients that represent the magnitude of cortical activity for each trial were computed in FEAT²⁴.

Statistical analysis. Using Linear Mixed Effects models^{27,28}, we aimed to differentiate the relationship between the two aspects of pain ratings and cortical activity. Across all trials we created a vector representing the single trial difference between pain intensity and pain unpleasantness (UI_gradient). Differences between pain intensity and pain unpleasantness occurred in 75% of trials. For the intensity trials we created a new vector by subtracting the ratings from the mean between intensity and unpleasantness for the trial. We used the analogue procedure for the unpleasantness trials:

$$(1) UI_gradient = [Unpleas-Intens/2+Unpleas/2; Intens-Intens/2+Unpleas/2]$$

The statistical model is expressed in Wilkinson notation²⁹; the included fixed effect of interest (UI_gradient:fmri) describes the magnitudes of the population common intercept and the population common slopes for the relationship between cortical data and the single trial differences in pain perception. The added random effect (i.e. 1|subject) models the specific intercept differences for each subject (e.g. subject-specific differences in fMRI beta coefficients):

$$(2) pain \sim fmri + UI_gradient:fmri + (1|subject)$$

Analysis was restricted to pain-related brain regions; we created a 3D mask consisting of the group result of the analysis in FEAT contrasting the pure pain trials with the baseline period (t>2). To correct for multiple comparisons we applied a randomisation approach. Behavioural data were shuffled and the entire analysis was repeated 5000 times within the boundaries of the 3D brain mask. The highest absolute t-values of each repetition across the whole volume were extracted. This procedure resulted in right-skewed distribution for each condition. Based on these distributions, the statistical thresholds were determined using the “palm_datapval” function publicly available in PALM³⁰.

REFERENCES

- 1 Melzack R, Casey KL. Sensory, motivational, and central control determinants of pain: a new conceptual model in pain. In: Kenshalo DR (ed). . Springfield, Ill., 1968, pp 423–443.
- 2 Kunz M, Lautenbacher S, LeBlanc N, Rainville P. Are both the sensory and the affective dimensions of pain encoded in the face? *Pain* 2012; **153**: 350–358.
- 3 Rainville P, Carrier B, Hofbauer RK, Bushnell CM, Duncan GH. Dissociation of sensory and affective dimensions of pain using hypnotic modulation. *Pain* 1999; **82**: 159–171.
- 4 Perlman DM, Salomons TV, Davidson RJ, Lutz A. Differential effects on pain intensity and unpleasantness of two meditation practices. *Emotion* 2010; **10**: 65–71.
- 5 Coghill RC, Sang CN, Maisog JM, Iadarola MJ. Pain intensity processing within the human brain: a bilateral, distributed mechanism. *J Neurophysiol* 1999; **82**: 1934–1943.
- 6 Hofbauer RK, Rainville P, Duncan GH, Bushnell MC. Cortical representation of the sensory dimension of pain. *J Neurophysiol* 2001; **86**: 402–411.
- 7 Gagnon-Dolbec A, Fortier M, Cormier S. Pain intensity and pain unpleasantness in triathletes: A study examining their associations with pain catastrophizing and pain expectations. *Psychol Sport Exerc* 2021; : 101928.
- 8 Shriver A. The unpleasantness of pain for humans and other animals. *Philosophy of Pain*. 2018; : 147–162.
- 9 Zunhammer M, Geis S, Busch V, Eichhammer P, Greenlee MW. Pain modulation by intranasal oxytocin and emotional picture viewing — a randomized double-blind fMRI study. *Scientific Reports*. 2016; **6**. doi:10.1038/srep31606.
- 10 Schreckenberger M, Siessmeier T, Viertmann A, Landvogt C, Buchholz H-GG, Rolke R *et al*. The unpleasantness of tonic pain is encoded by the insular cortex. *Neurology* 2005; **64**: 1175–1183.
- 11 Rainville P, Duncan GH, Price DD, Carrier B, Bushnell MC. Pain affect encoded in human anterior cingulate but not somatosensory cortex. *Science* 1997; **277**: 968–971.
- 12 Poldrack RA, Fletcher PC, Henson RN, Worsley KJ, Brett M, Nichols TE. Guidelines for reporting an fMRI study. *Neuroimage* 2008; **40**: 409–414.
- 13 Zeidan F, Martucci KT, Kraft R, Gordon NS, McHaffie JG, Coghill RC. Brain mechanisms supporting the modulation of pain by mindfulness meditation. *J Neurosci* 2011; **31**: 5540–5548.
- 14 Zhou F, Li J, Zhao W, Xu L, Zheng X, Fu M *et al*. Empathic pain evoked by sensory and emotional-communicative cues share common and process-specific neural representations. *Elife* 2020; **9**. doi:10.7554/eLife.56929.
- 15 Baliki MN, Chialvo DR, Geha PY, Levy RM, Harden RN, Parrish TB *et al*. Chronic pain and the emotional brain: specific brain activity associated with spontaneous fluctuations of intensity of chronic back pain. *J Neurosci* 2006; **26**: 12165–12173.
- 16 Kong J, White NS, Kwong KK, Vangel MG, Rosman IS, Gracely RH *et al*. Using fMRI to dissociate sensory encoding from cognitive evaluation of heat pain intensity. *Hum Brain Mapp* 2006; **27**: 715–721.
- 17 Wiech K, Tracey I. Pain, decisions, and actions: a motivational perspective. *Front Neurosci* 2013; **7**: 46.
- 18 Loggia ML, Juneau M, Bushnell CM. Autonomic responses to heat pain: Heart rate, skin conductance, and their relation to verbal ratings and stimulus intensity. *Pain* 2011; **152**: 592–598.
- 19 Geuter S, Gamer M, Onat S, Büchel C. Parametric trial-by-trial prediction of pain by easily available physiological measures. *Pain* 2014; **155**: 994–1001.
- 20 Nickel MM, May ES, Tiemann L, Postorino M, Ta Dinh S, Ploner M. Autonomic responses to tonic pain are more closely related to stimulus intensity than to pain intensity. *Pain* 2017; **158**: 2129–2136.
- 21 Mobascher A, Brinkmeyer J, Warbrick T, Musso F, Wittsack HJ, Stoermer R *et al*. Fluctuations in electrodermal activity reveal variations in single trial brain responses to painful laser stimuli--a

- fMRI/EEG study. *Neuroimage* 2009; **44**: 1081–1092.
- 22 Rolls ET. The orbitofrontal cortex and emotion in health and disease, including depression. *Neuropsychologia* 2019; **128**: 14–43.
- 23 Schulz E, Stankewitz A, Winkler AM, Irving S, Witkovský V, Tracey I. Ultra-high-field imaging reveals increased whole brain connectivity underpins cognitive strategies that attenuate pain. *Elife* 2020; **9**. doi:10.7554/eLife.55028.
- 24 Schulz E, Stankewitz A, Witkovský V, Winkler AM, Tracey I. Strategy-dependent modulation of cortical pain circuits for the attenuation of pain. *Cortex*. 2019; **113**: 255–266.
- 25 Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. *Neuroimage* 2012; **62**: 782–790.
- 26 Griffanti L, Salimi-Khorshidi G, Beckmann CF, Auerbach EJ, Douaud G, Sexton CE *et al*. ICA-based artefact removal and accelerated fMRI acquisition for improved resting state network imaging. *Neuroimage* 2014; **95**: 232–247.
- 27 Witkovský V. Estimation, testing, and prediction regions of the fixed and random effects by solving the Henderson's mixed model equations. *Measurement science review* 2012; **12**: 234–248.
- 28 Michail G, Dresel C, Witkovský V, Stankewitz A, Schulz E. Neuronal Oscillations in Various Frequency Bands Differ between Pain and Touch. *Front Hum Neurosci* 2016; **10**: 182.
- 29 Wilkinson GN, Rogers CE. Symbolic Description of Factorial Models for Analysis of Variance. *J R Stat Soc Ser C Appl Stat* 1973; **22**: 392–399.
- 30 Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation inference for the general linear model. *Neuroimage* 2014; **92**: 381–397.