Cerebrospinal fluids from healthy pregnant women does not harbor a detectable microbial community

Yongyong Kang^{1,3,11#}, Xinchao Ji^{2#}, Li Guo^{3,4,5#}, Han Xia^{3,6#}, Xiaofei Yang^{1,4,7}, Zhen Xie⁸, Xiaodan Shi⁹, Rui Wu⁸, Dongyun Feng⁹, Chen Wang¹⁰, Min Chen¹⁰, Wenliang Zhang⁴, Hong Wei⁴, Yuanlin Guan⁶, Kai Ye^{1,3,4,5,11*}, Gang Zhao^{8,9*}

¹Genome Institute, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, China

²Department of Neurology, Xi'an No. 3 Hospital, The Affiliated Hospital of Northwest University, Xi'an 710061, China

³School of Automation Science and Engineering, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an 710049, China

⁴MOE Key Laboratory for Intelligent Networks & Network Security, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an 710049, China

⁵School of Life Science and Technology, Xi'an Jiaotong University, Xi'an 710049 China

⁶Hugobiotech Co., Ltd., Beijing 100000, China

⁷School of Computer Science and Technology, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an 710049, China

⁸School of Medicine, Northwest University, Xi'an 710069, China

⁹Department of Neurology, Xijing Hospital, The Fourth Military Medical University, Xi'an 710032, China

¹⁰Department of Anesthesiology, Xijing Hospital, The Fourth Military Medical University, Xi'an, 710032, China

¹¹Center for Mathematical Medical, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, China

* Corresponding author: kaiye@xjtu.edu.cn; zhaogang@nwu.edu.cn

# These authors contributed equally to this work

ABSTRACT Cerebrospinal fluids circulating human central nervous system have long been considered aseptic in healthy individuals, because normally the blood-brain barrier protects against microbial invasions. However, this dogma has been questioned by several reports that microbes were identified in human brains, raising the question whether a microbial community is present in cerebrospinal fluids of healthy individuals without neurological diseases. Here, we collected and analyzed metagenomic and metatranscriptomic sequencing data of cerebrospinal fluid specimens from a cohort of 23 pregnant women aged between 23 and 40 and one-to-one matched contamination controls. From data analysis of 116 specimens of eight different types, we detected 619 nonredundant microbial taxa which were dominated by bacteria (75%) and viruses (24%). In cerebrospinal fluids metagenomic samples, a total of 76 redundant species were detected including four (one nonredundant) eukaryota taxa, eleven (four nonredundant) bacteria, and 61 (21 nonredundant) viruses that were mostly bacteriophages. Metagenomic data analysis found no significant difference between cerebrospinal fluid specimens and negative controls in terms of microbial species diversity. In addition, no active or viable microbiome were present in the cerebrospinal fluid samples after subtracting microbes detected in contamination controls. In conclusion, we found no strong evidence that colonized microbial community exist in the cerebrospinal fluids of healthy individuals.

IMPORTANCE Microbiome are prevalent throughout human bodies with profound health implications. However, it remains unclear whether a microbiome is present and active in human cerebrospinal fluids that are long considered aseptic given the blood-

53 brain barrier. Here, we applied unbiased metagenomic and metatranscriptomic
 54 sequencing to detect microbiome in cerebrospinal fluids collected from a cohort of 23
 55 pregnant women with matched controls. By analyzing 116 specimens of eight types, no
 56 strong evidence was found to support a presence of colonized microbiome in the
 57 cerebrospinal fluids. Our findings have profound implications to human immunity
 58 against neurological infections and disorders, providing a guide for disease diagnostics,
 59 prevention and therapeutics in clinical settings.

60

61 **Keywords:** cerebrospinal fluids, mNGS, metagenomics, pathogen, microbiome

INTRODUCTION

First defined by Joshua Lederberg in 2001(1), human microbiome has since been discovered at almost every part of human bodies such as gut, oral, skin, bladder, vagina, lungs(2-8). It has profound impact on human health, being associated with a broad range of human diseases including cancers, diabetes, schizophrenia and autoimmune diseases etc.(9-12). However, due to difficulties in the identification and traceability of contaminations, it remains controversial whether there are colonized microbial community at some sites such as placenta, blood and amniotic fluid (13-17), although recently both experimental and analytical methods have improved in sensitivity and accuracy regarding microbiome discovery.

Cerebrospinal fluid (CSF) circulating the human central nervous system (CNS) has long been considered sterile given that the blood-brain barrier is thought to effectively protect against microbial invasions. However, this traditional knowledge of microbe-free CSF has been challenged in recent years with several reports of microbes detected in human brains and CSF. For example, a bacterial pathogen *Porphyromonas gingivalis* was identified in brain regions including cerebral cortex and hippocampus in patients with Alzheimer's disease(18). In addition, a number of DNA viruses in CSF were identified from a subjects of virome(19). It remains elusive whether these reports are evidence of a common microbiome in human CSF and CNS, or simply sporadic and accidental events.

Given the debate over the existence of any microbial community in CSF and the importance of understanding microbial infection in human central nervous systems, we

performed microbiome analysis to characterize bacteria, archaea, eukaryota and viruses of CSF from a cohort of 23 donors without neurological disorders, as well as one-to-one matched positive controls (oral and skin) and negative controls (normal saline). DNA/RNA extraction buffers and sterile swab were also collected as controls. In total, 116 specimens of eight types were used in this study. Considering the limitations of 16S ribosomal RNA based approach to achieve a consistent result in species or strain level(20-23), we choose unbiased metagenomic next-generation sequencing (mNGS) for rapidly detecting all genetic materials of microbiome at species resolution and use metatranscriptomic sequencing to assess the physiological states of microbial communities in CSF(24, 25). mNGS as a promising approach, its clinical diagnostic performance in infectious diseases has been widely adopted in the medical community by multi-center studies(26-28).

In our data analysis, we found no significant difference between CSF specimens and negative controls in microbial species diversity. In all CSF samples, no active or viable microbiome was present after subtracting microbial taxa detected in CSF by those detected in contamination controls. Taken together, no strong evidence was found in our study supporting that colonized microbiome exists in the cerebrospinal fluids.

RESULTS

Metagenomic sequencing of cerebrospinal fluids in healthy individuals

To investigate whether there is microbiome in CSF, we collected and analyzed microbiome of CSF samples from 23 pregnant women aged 23–40 years who underwent intraspinal anesthesia before the caesarean section via lumbar puncture,

coupled with normal saline collected with syringe as negative controls. For each subject, oral and skin microbiomes were also collected and analyzed as positive controls (Figure 1a). All samples were then subjected to DNA extraction and metagenomic shotgun sequencing and analysis. Finally, to validate whether the microbiome, if any detected in CSF, is physiologically active, metatranscriptome profilings for 12 of the pregnant women CSF samples were performed. After quality control (QC) with KneadData(29) (v0.7.2) for metagenomic and metatranscriptomic sequencing data, MetaPhlAn(30) (latest version 3), a state-of-the-art taxonomic classification tool based on unique clade-specific marker genes, was used to detect microbes in each sample.

In total, we detected 619 nonredundant microbial taxa in 116 specimens using metagenomic and metatranscriptomic sequencing and analysis (Supplementary Table S1). These microbes detected in all samples were dominated by bacteria (75%) and viruses (24%). Overall, skin, oral and swab samples had the most abundant microbiome of all samples with 393, 199 and 137 nonredundant microbial taxa, respectively. This came as no surprise because skins and orals are well known to harbor a plethora of microbes. By contrast, the number of microbial taxa detected in CSF_DNA (26), negative controls (49) and extraction buffers (27) were relatively fewer (Figure 1b).

We then compared the taxa detected in different specimen types, finding that there was little overlap among all samples. Skin, oral and sterile swab had a large amount of unique microbial taxa among all sample types, with 243, 129 and 36 taxa found only in these samples, respectively (Figure 1b). Although a large variation in the number of microbes detected was observed for skins, orals and swabs, a much smaller variation

was found for CSF_DNA, CSF metatranscriptomic (CSF_RNA), negative controls and extraction buffers (DNA/RNA buffer) (Figure 1b). The oral samples were rich in *Streptococcus*, *Veillonella*, *Neisseria*, *Rothia* and *Prevotella*, while the skin samples were rich in *Cutibacterium*, *Staphylococcus*, *Micrococcus*, *Malassezia*, consistent with many previous studies(4, 6, 31) (Supplementary Figure 1). Our successful detection of known microbiome for orals and skins provided a proof-of-concept of NGS-based metagenomic sequencing method, laying a solid foundation for our exploration of CSF microbiome using such a method.

We next focused on examining the microbial species detected for each CSF_DNA specimen. In CSF samples, a total of 76 redundant species including 11 (4 nonredundant) bacteria, 61 (21 nonredundant) viruses and four (one nonredundant) eukaryota taxa were detected (Figure 1c). Most of these viruses were bacteriophages. The relative abundance of microbes suggested the species “*Cyprinid herpesvirus 3*” are the predominant species in 19 of 23 CSF_DNA samples (Figure 1d). *Cutibacterium acnes* in species level appeared in 5 specimens. Additionally, 100%, 26%, 22%, and 22% of all CSF_DNA samples contain *Cyprinid herpesvirus 3*, *Human alphaherpesvirus 2*, *Enterobacteria phage mEp460* and *Dasheen mosaic virus*, respectively. However, “*Cyprinid herpesvirus 3*” detected in all CSF_DNA samples were also found in all negative and skin samples, suggesting a likely external source of these microbes during the CSF sampling procedure.

The microbiome signature of cerebrospinal fluids and negative controls is similar

149 Since microbial species were identified in both CSF_DNA and negative controls, it is
150 likely that microbial cells and/or DNA present in negative controls may have been
151 introduced into CSF during the sampling process. Similarly, the possibility of skin
152 microbiome being introduced into CSF during lumbar puncture could not be ruled out,
153 despite the application of skin surface sterilization. Therefore, we asked how similar in
154 general the microbiome signature is for different sample types by comparing the
155 microbial species detected in these samples. We first performed Non-metric
156 Multidimensional Scaling (NMDS) analysis and principal coordinates analysis (PCoA),
157 and then characterized the beta diversity of CSF and other specimen types using Bray-
158 Curtis distances, a metric commonly used to evaluate microbiome difference among
159 samples supported by Wilcoxon statistical significance. NMDS, PCoA (Supplementary
160 Figure 2), and beta-diversity analysis revealed an overall clear separation of microbial
161 communities for each sample type, except that microbiome in CSF_DNA specimens
162 overlapped partially with negative controls (Figure 2a). Statistical analysis suggested
163 beta-diversity between CSF_DNA and other sample types was significantly different
164 from CSF_DNA self-comparison. However, there was no significant difference
165 between CSF_DNA self beta-diversity and CSF_DNA-negative beta diversity (Wilcox
166 test: $p=0.59$) (Figure 2b). In addition, the low diversity suggested the microbial
167 communities in CSF_DNA and negative controls were highly similar. In fact, shared
168 microbial taxa between CSF_DNA and negative control accounted for 42% and 22%
169 of CSF_DNA and negative control, respectively. By contrast, 58% microbial taxa in
170 CSF-DNA were detected in skin samples, whereas only 4% of skin microbes were

found in CSF-DNA specimens. On one hand, these results indicated the microbial cells or DNA detected in CSF samples may partly have come from negative controls during sample collection. On the other hand, the high beta-diversity between skin and CSF specimens implied that the skin surface sterilization before lumbar punctures effectively prevented the contamination of CSF samples with most, if not all, skin microbes.

No microbiome is present in the CSF after subtracting microbes from controls

With the detected microbiome in CSF samples, we questioned whether these microbes were truly CSF inhabitants or simply brought in from external sources such as skins, negative controls and DNA extraction buffer. To verify whether the CSF contains de facto colonized microbial communities, we subtracted the microbes collectively detected in negative control samples and DNA extraction buffer samples from microbes of each CSF_DNA sample, a method commonly used and previously described by human microbiome study(17). After subtraction, 12 CSF samples contained no microbe, whereas the other 11 CSF samples contain a total of 14 microbial taxa including 11 viruses, 2 bacteria and 1 eukaryota. Since an introduction of microbes from skins could not be completely ruled out, we further checked whether the 14 taxa were present in skins as well and found that 6 of the 14 taxa were also found in the matching skin microbiome. This left eight microbial taxa after subtraction as potential CSF inhabiting microbes, including five viruses (“*Bovine alphaherpesvirus 1*”, “*Escherichia virus V5*”, “*Klebsiella virus KP27*”, “*Macaca mulatta polyomavirus 1*”, “*Trichoplusia_ni_single_nucleopolyhedrovirus*”), two bacteria (*Hydrogenophilus thermoluteolus*, *Tepidimonas fonticaldi*), and one eukaryota (*Aspergillus turcosus*).

193 The detection of microbes using the metagenomic approach offers a glimpse of
 194 microorganisms present in certain niches. However, it remains uncertain whether these
 195 microbes are live or dead, as DNA from dead cells are also detectable by mNGS.
 196 Therefore, we further evaluated the physiological activities of the potential CSF-
 197 inhabiting microbes using metatranscriptomic sequencing, because the number of
 198 microbes detected by both in metagenomic and metatranscriptome would indicate
 199 active microbes may be present in CSF samples. CSF transcriptomics revealed
 200 transcripts for several microbial taxa including “*Equine infectious anemia virus*” and
 201 “*Cyprinid herpesvirus 3*” appearing in all samples, and *Escherichia coli* and *Dasheen*
 202 *mosaic virus* appearing in eleven and nine samples, respectively. We then asked, for the
 203 eleven CSF-DNA samples with microbes left after subtraction by negative controls
 204 and DNA extraction buffers, whether these microbes have detected in
 205 metatranscriptomic data. The result showed that only “*human alphaherpesvirus 1*” had
 206 signals from both CSF genomics and transcriptomics. However, “*human*
 207 *alphaherpesvirus 1*” also appeared in the skin, suggesting no active microbiome was
 208 detectable in CSF after removing this species potentially originated from skins.
 209 Although metagenomic analysis detected the one *Aspergillus turcosus* species from four
 210 individuals (Figure 3), no transcripts of *Aspergillus turcosus* were detected in
 211 metatranscriptomic, suggesting a lack of living cell activity. *Aspergillus turcosus* is well
 212 known as opportunistic human pathogens and can cause infections in individuals of
 213 compromised immune systems(32). How this fungal species (cells or DNA) reaches the
 214 CSF of the five healthy individuals is unknown, but it shows CSF, though without an

active microbiome, might not be entirely free of opportunistic fungi which could potentially cause infections in central nervous systems when host immune system is compromised. Taken together, our study found no strong evidence supporting actively transcribed microbiome in the CSF.

DISCUSSION

Hereby, CSF samples from a cohort of 23 healthy individuals without neurological disease with a matched set of controls were collected for microbiome detection using culture-independent approach by a whole genome shotgun sequencing. The metagenomic data analysis indicated that there was no significant difference between CSF specimens and negative controls in beta diversity of detected microbes. In addition, no clear signal of active microbiome in the CSF samples was found by comparing CSF and contamination controls. Except *Aspergillus turcosus* appeared in four samples, no microbiome was present in more than two CSF samples after being subtracted by microbes in negative controls and DNA extraction buffer.

Compared with bacteria, more viruses were detected in CSF specimens. These viruses are mainly bacteriophages, most of which are also present in negative and skin samples. Although bacteriophages in the CSF have been reported, clear evidence for regular colonization of CSF by these viruses is lacking (19). Whether these viruses appear accidentally or colonized in CSF needs further investigation.

In our results, four CSF samples contained *Aspergillus* DNA, but no *Aspergillus* nucleic acid was detected in the corresponding RNA samples, suggesting that these *Aspergillus* DNA fragments may have come from contaminations. Due to the high sensitivity of

mNGS, it can also detect trace amount of nucleic acid fragments released from dead microorganisms present in human periphery blood or tissues, experiment reagents and consumables. Furthermore, when using puncture to collect CSF specimen, tissues such as skin, muscle, blood vessels are potential sources of contamination. Except for strict disinfection measures before operation, constructing a database of colonizing microorganisms of these healthy tissues will enable subsequent bioinformatics analysis to filter out noise signals and reduce the false positive rate.

Highly sensitive mNGS represents a powerful tool for detecting microbiome at species resolution, especially for microbiome studies in specimens of low-abundance biomass, such as CSF. Currently, mNGS has become an important auxiliary method for clinical pathogenic diagnosis and treatment of infectious diseases. Main challenges of studying this issue have been an overall lack of CSF samples from healthy human subjects and the technically sound sampling as well as data analysis methods based on different reference databases and taxonomic strategies.

We focused on determining whether a CSF microbiome is present in healthy individuals without neurological disorders, a long-disputed issue in scientific and clinical research field. Our data analysis demonstrated that the microbiome of CSF was indistinguishable from contamination controls. It is intriguing but remains unclear whether a microbiome is present in CSF of patients diagnosed with diseases such as Alzheimer's disease, multiple sclerosis, Parkinson's disease and what roles the CSF microbiome plays in the development of these disorders.

In conclusion, using metagenomic combined with metatranscriptomic deep sequencing,

we found microbiome profile in CSF samples was indistinguishable from that in contamination controls. Our data indicated that by current approaches there was no evidence to support the existence of a CSF microbiome in the populations without known neurological disorders. Such findings shall have great implications to human health especially neurological disorders and infections, providing a guide for disease diagnostics, prevention and therapeutics in clinical settings.

MATERIALS AND METHODS

Subjects

Twenty-three donors were recruited from the Xijing Hospital of the Fourth Military Medical University. All subjects were enrolled from obstetrics department in which the pregnant woman aged 23–40 years need intraspinal anesthesia before the caesarean section. Subjects who have suffered from central nervous system infection disease (eg, meningitis, encephalitis) or any systemic infection disease and autoimmune disease (eg, hepatitis, tuberculosis, systemic lupus erythematosus, rheumatism) and have received antibiotics treatment in the past six months prior to sample collection were excluded. We also excluded subjects with a history of hypertension, diabetes, heart disease, cancer and neurological disease (eg, Alzheimer’s disease, Parkinson’s disease, multiple sclerosis, epilepsy).

Sample collection

Lumbar puncture was performed in the 23 subjects enrolled in this study and the CSF samples were collected in a 4ml centrifugal tube with syringe and then stored in a –80 °C freezer for metagenomics analysis. Twelve CSF samples were randomly

selected from 23 pregnant women for metatranscriptome studies and RNA protection reagent was added to the CSF immediately after collection. Then, the samples were centrifuged and the pellets stored at -80°C for metatranscriptomic sequencing. Meanwhile, normal saline was collected with syringes for environmental controls (negative control). Furthermore, oral and skin samples were selected from 23 enrolled subjects as one-to-one matched positive control. For skin positive controls: The back skin of $5\times 5\text{ cm}^2$ areas around the puncture site (L3-L4 intervertebral space) were swabbed with a sterile cotton swab before skin clean with povidone iodine. To maximize microbial load, no bathing was permitted within 24 hours of sample collection. For oral positive controls, all subjects were forbidden to eat and drink six hours before operation. The surfaces of tongue, buccal fold, hard palate, soft palate, tooth, gingiva and saliva were swabbed with sterile swab. Unused sterile swabs were collected for negative controls (“sterile swab”). Details of Matching information between samples are described in Supplementary Table S2.

DNA extraction and purification

DNA was isolated using the QIAamp DNA Mini Kit (Qiagen 51304) according to the manufacturer’s instructions. 1) DNA extraction from swabs: Swab tips were cut and placed in a 2 ml microcentrifuge tube and then 400 μl PBS were added. Next, 20 μl of proteinase K and 400 μl of buffer AL were added, vortexed for 10 s, and the solution was incubated for 15 min at 56°C . And then added 400 μl ethanol (100%) and mixed again by vortexing. Lastly, DNA purification was performed with buffer AW1 and AW2 using QIAamp Mini spin column, followed by elution with 35 μl of buffer EB. 2) DNA

extraction from CSF and normal saline controls: 200 µl sample was added into the microcentrifuge tube, and then added 20 µl of proteinase K and 200 µl of buffer AL respectively, vortexed for 10 s, and the solution was incubated for 15 min at 56 °C. Next, added 200 µl ethanol (100%) and mixed again by vortexing. DNA purification was performed as described above.

Metagenomics library construction

For preparation of metagenomics libraries, the QIAseq FX DNA Library Kit (Qiagen; 180715) was used. The construction involved five steps: 1) Fragmentation and End-repair: to generate 200–300 bp fragments, 32.5µl purified DNA were fragmented by incubation with FX buffer 5µl, FX enhancer 2.5µl and 10 µl FX enzyme mix at cycling program: 4 °C 1minute → 32 °C 12minutes → 65 °C 30minutes → 4 °C hold. 2) Adapter ligation: 5 µl of adaptor, 20 µl of ligation buffer, 10 µl of DNA ligase and 15 µl of nuclease-free water were added and incubated for 15 minutes at 20 °C to initiate adapter ligation. Adapter ligation cleanup was performed immediately, 3) Adapter ligation cleanup: 80 µl of resuspended AMPure® XP beads (0.8×) were added to each ligated sample and mix well by pipetting. Next, the mixture was incubated for 5 minutes at room temperature and then the beads were pelleted on a magnetic stand (Invitrogen) for 2 minutes. The supernatant was discarded and the pellet was washed twice with 200 µl of 80% ethanol, then the beads were eluted with 52.5 µl of buffer EB. Subsequently, 50 µl of supernatant was transferred into a new 1.5 ml microcentrifuge tube and a second purification was performed with 50 µl (1×) AMPure® XP beads. The final, 23.5 µl of purified DNA sample was obtained. 4) Amplification of library DNA: 25 µl of HiFi

PCR Master Mix, 1.5 µl of Primer Mix and 23.5 µl of library DNA were added in PCR tube. PCR enrichment was performed under the cycle conditions: 2 minutes at 94 °C, 12 × (20 s at 98 °C, 30 s at 60 °C, 30 s at 72 °C), and 1 minute at 72 °C. The final, to obtain libraries, the PCR products were purified with AMPure XP beads as described above.

RNA extraction and purification

Total RNA was extracted using the RNeasy Mini Kit (Qiagen; 74104) according to the manufacturer's instructions. The pellet of each sample which has been treated with RNA protection reagent as described above, was resuspended in 100µl TE buffer containing lysozyme, and Proteinase K was added into the mixture, then incubated for 10 minutes at room temperature. 350µl of buffer RLT was added and vortexed vigorously. The final, RNA isolation and purification was performed with buffer AW1 and RPE respectively using RNeasy Mini spin column, followed by elution with RNase-free water.

RNA library preparation for metatranscriptomics sequencing

For construction of RNA libraries, the QIAseq FX Single Cell RNA Library Kit (Qiagen; 180733) was used. The construction involved five steps: 1) Genomic DNA (gDNA) removal: 8 µl of purified RNA and 3 µl of NA denaturation buffer were added into a sterile PCR tube and incubated for 3 minutes at 95 °C. To remove genomic DNA, 2 µl of gDNA wipeout buffer was added and incubated for 10 minutes at 42 °C. 2) Reverse transcription: 4 µl of RT/Polymerase buffer, 1 µl of random primer, 1 µl of Oligo dT primer and 1 µl of Quantiscript RT enzyme mix were added in each sample and reverse

transcription was carried out for 60 minutes at 42 °C. 3) Ligation: 8 µl of ligase buffer and 2 µl of ligase mix were added into the RT reaction and incubated at 24°C for 30 minutes. 4) Whole transcriptome amplification: 1 µl of REPLI-g SensiPhi DNA Polymerase and 29µl of reaction buffer were used for Multiple Displacement Amplification (MDA), then incubate at 30°C for 2 h. The final, an approximate length of 2000–70,000 bp amplified cDNA was produced. 5) Enzymatic Fragmentation: The amplified cDNA was diluted 1:3 in H₂O sc, 10 µl of the diluted DNA and FX Enzyme Mix were used to obtain 300 bp library fragment with reaction conditions: 4 ° C 1minute→32 ° C 15minutes→65 ° C 30minutes→4 ° C hold. 6) Adapter ligation: 5 µl of adapter and 45 µl of ligation master mix were added into each sample and incubated at 20° C for 15 minutes. Subsequently, the adapter ligation cleanup was performed with AMPure XP beads as described above. The final, purified libraries were obtained ready for sequencing without further PCR amplification.

Next generation equencing

Shotgun sequencing was performed on Illumina Hiseq platform for all samples (paired end library of 150-bp and 150-bp read length). Approximately, 25 Gb and 5 Gb of raw paired-end reads were obtained per sample in the CSF genomics samples and negative samples, respectively.

Data quality control

To reduce the impact of host reads on the results, we need to remove human reads involved in the raw sequencing data before bioinformatics analysis. KneadData(29) (v0.7.2), a widely used tool, is designed to perform quality control on metagenomic and

metatranscriptomic sequencing data, especially for microbiome experiments. All reads were filtered using KneadData with the following trimmomatic options: ILLUMINACLIP: TruSeq3-PE-2.fa:2:30:10:8:true, SLIDINGWINDOW:4:20, MINLEN:50 and bowtie2 options: --very-sensitive, --dovetail. The proportion of human reads in CSF genomics samples is up to 92%.

Detecting potential microbiome

MetaPhlAn(30) (v 3.0.1) is a computational tool for profiling the composition of microbial communities (bacteria, archaea, viruses and eukaryotes) from shotgun sequencing data. Based on ~1.1M unique clade-specific marker genes identified from ~100,000 reference genomes, MetaPhlAn can profile unambiguous taxonomic assignments and accurate estimation of organismal relative abundance in species-level resolution. Classifying the reads to marker genes database, MetaPhlAn outputs a file containing detected microbes and relative abundance in different level. MetaPhlAn ran with custom parameters: --add_viruses --input_type fastq --read_min_len 50. It's worth noting that MetaPhlAn (previous version 2) was the only bioinformatics tool with 0% false positive relative abundance and the best diversity estimate(33).

β -diversity and phylogenetic analysis

Using R (version 3.6.3) with R studio environment, β -diversity (between-sample diversity) was estimated by Bray-Curtis dissimilarity in vegan package. All figures are plotted using R.

Supplementary files

Supplementary Table S1. Microbes detected in different samples.

Supplementary Figure S1. Top 10 genus in oral and skin samples, respectively. Microbial community structures of 23 Oral (figure S1a) and Skin (figure S1b) samples shown in a stacked barplot that summarizes the relative abundance of different genus detected.

Supplementary Figure S2. Principal Coordinates Analysis (PCoA) analysis of microbial species detected from different sample types.

Supplementary Table S2. Samples label and matching information.

Availability of data and materials

The clean sequence data reported in this paper have been deposited in the Genome Sequence Archive in BIG Data Center(34, 35), Chinese Academy of Sciences, under accession number [PRJCA004977XXXXX](https://bigd.big.ac.cn/bioproject/XXXXX) that are publicly accessible at [https://bigd.big.ac.cn/bioprojectXXXXX](https://bigd.big.ac.cn/bioproject/XXXXX).

Acknowledgements

We thank Peng Jia, Tingjie Wang, Ningxin Dang, Honghui Shen and Tun Xu for helpful discussions regarding data analysis and Jing Hai for administrative and technical support. We thank the High-Performance Computing Cluster of the First Affiliated Hospital of Xi'an Jiaotong University for data processing.

This study was supported by the National Key R&D Program of China (Grant Nos. 2018YFC0910400, 2017YFC0907500, and 2016 YFC0904501), the National Natural Science Foundation of China (Grant Nos. 31671372, 61702406, 31701739, and 31970317), the National Science and Technology Major Project of China (Grant No.

2018ZX10302205), as well as the General Financial Grant from the China Postdoctoral Science Foundation (Grant Nos. 2017M623178 and 2017M623188).

Ethics Statement

This study was approved by the Ethics Committee of the Xijing Hospital of the Fourth Military Medical University. All procedures were conducted in accordance with the approved guidelines. All patients read and signed the consent form before sample collection.

Competing interests

The authors declare that they have no competing interests.

References

1. Lederberg J, McCray A. 2001. 'Ome Sweet 'Omics--A Genealogical Treasury of Words. *The Scientist* 15:8-8.
2. Gilbert JA, Blaser MJ, Caporaso JG, Jansson JK, Lynch SV, Knight R. 2018. Current understanding of the human microbiome. *Nature Medicine* 24:392-400.
3. Clemente JC, Ursell LK, Parfrey LW, Knight R. 2012. The impact of the gut microbiota on human health: an integrative view. *Cell* 148:1258-1270.
4. Nasidze I, Li J, Quinque D, Tang K, Stoneking M. 2009. Global diversity in the human salivary microbiome. *Genome research* 19:636-643.
5. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket CO. 2011. Vaginal microbiome of reproductive-age women. *Proceedings of the National Academy of Sciences* 108:4680-4687.
6. Grice EA, Segre JA. 2011. The skin microbiome. *Nature reviews microbiology* 9:244-253.
7. Beck JM, Young VB, Huffnagle GB. 2012. The microbiome of the lung. *Translational Research* 160:258-266.
8. Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, Chinwalla AT, Creasy HH, Earl AM, FitzGerald MG, Fulton RS. 2012. Structure, function and diversity of the healthy human microbiome. *nature* 486:207.
9. Iida N, Dzutsev A, Stewart CA, Smith L, Bouladoux N, Weingarten RA, Molina DA, Salcedo R, Back T, Cramer S. 2013. Commensal bacteria control cancer response to therapy by modulating the tumor microenvironment. *Science* 342:967-970.
10. Moreno-Indias I, Cardona F, Tinahones FJ, Queipo-Ortuño MI. 2014. Impact of the gut microbiota on the development of obesity and type 2 diabetes mellitus. *Frontiers in microbiology* 5:190.
11. Zhu F, Ju Y, Wang W, Wang Q, Guo R, Ma Q, Sun Q, Fan Y, Xie Y, Yang Z, Jie Z, Zhao B, Xiao L, Yang L, Zhang T, Feng J, Guo L, He X, Chen Y, Chen C, Gao C, Xu X, Yang H, Wang J, Dang Y, Madsen L, Brix S, Kristiansen K, Jia H, Ma X. 2020. Metagenome-wide association of gut microbiome features for schizophrenia. *Nature Communications* 11:1612.
12. Morgan XC, Tickle TL, Sokol H, Gevers D, Devaney KL, Ward DV, Reyes JA, Shah SA, LeLeiko N, Snapper SB. 2012. Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. *Genome biology* 13:R79.
13. de Goffau MC, Lager S, Sovio U, Gaccioli F, Cook E, Peacock SJ, Parkhill J, Charnock-Jones DS, Smith GCS. 2019. Human placenta has no microbiome but can contain potential pathogens. *Nature* 572:329-334.
14. Lauder AP, Roche AM, Sherrill-Mix S, Bailey A, Laughlin AL, Bittinger K, Leite R, Elovitz MA, Parry S, Bushman FD. 2016. Comparison of placenta samples with contamination controls does not provide evidence for a distinct placenta microbiota. *Microbiome* 4:29.
15. Castillo DJ, Rifkin RF, Cowan DA, Potgieter M. 2019. The Healthy Human Blood Microbiome: Fact or Fiction? *Frontiers in Cellular and Infection Microbiology* 9.
16. Whittle E, Leonard MO, Harrison R, Gant TW, Tonge DP. 2019. Multi-Method Characterization of the Human Circulating Microbiome. *Frontiers in Microbiology* 9.
17. Lim ES, Rodriguez C, Holtz LR. 2018. Amniotic fluid from healthy term pregnancies does not harbor a detectable microbial community. *Microbiome* 6:1-8.
18. Dominy SS, Lynch C, Ermini F, Benedyk M, Marczyk A, Konradi A, Nguyen M, Haditsch U, Raha

- 465 D, Griffin C. 2019. Porphyromonas gingivalis in Alzheimer's disease brains: Evidence for
466 disease causation and treatment with small-molecule inhibitors. Science advances
467 5:eaau3333.
- 468 19. Ghose C, Ly M, Schwanemann LK, Shin JH, Atabakhsh K, Barr JJ, Little M, Schooley RT, Pride
469 DT. 2019. The virome of cerebrospinal fluid: viruses where we once thought there were none.
470 Frontiers in microbiology 10:2061.
- 471 20. Engelbrektson A, Kunin V, Wrighton KC, Zvenigorodsky N, Chen F, Ochman H, Hugenholtz P.
472 2010. Experimental factors affecting PCR-based estimates of microbial species richness and
473 evenness. The ISME journal 4:642-647.
- 474 21. Klappenbach JA, Saxman PR, Cole JR, Schmidt TM. 2001. rrndb: the Ribosomal RNA Operon
475 Copy Number Database. Nucleic acids research 29:181-184.
- 476 22. Claesson MJ, Wang Q, O'Sullivan O, Greene-Diniz R, Cole JR, Ross RP, O'Toole PW. 2010.
477 Comparison of two next-generation sequencing technologies for resolving highly complex
478 microbiota composition using tandem variable 16S rRNA gene regions. Nucleic acids research
479 38:e200-e200.
- 480 23. Gevers D, Cohan FM, Lawrence JG, Spratt BG, Coenye T, Feil EJ, Stackebrandt E, Van de Peer Y,
481 Vandamme P, Thompson FL. 2005. Re-evaluating prokaryotic species. Nature Reviews
482 Microbiology 3:733.
- 483 24. Bikel S, Valdez-Lara A, Cornejo-Granados F, Rico K, Canizales-Quinteros S, Soberón X, Del
484 Pozo-Yauner L, Ochoa-Leyva A. 2015. Combining metagenomics, metatranscriptomics and
485 viromics to explore novel microbial interactions: towards a systems-level understanding of
486 human microbiome. Computational and Structural Biotechnology Journal 13:390-401.
- 487 25. Shakya M, Lo C-C, Chain PSG. 2019. Advances and Challenges in Metatranscriptomic Analysis.
488 Frontiers in Genetics 10.
- 489 26. Ramachandran PS, Wilson MR. 2020. Metagenomics for neurological infections — expanding
490 our imagination. Nature Reviews Neurology 16:547-556.
- 491 27. Chiu CY, Miller SA. 2019. Clinical metagenomics. Nature Reviews Genetics 20:341-355.
- 492 28. Wilson MR, Sample HA, Zorn KC, Arevalo S, Yu G, Neuhaus J, Federman S, Stryke D, Briggs B,
493 Langelier C, Berger A, Douglas V, Josephson SA, Chow FC, Fulton BD, DeRisi JL, Gelfand JM,
494 Naccache SN, Bender J, Dien Bard J, Murkey J, Carlson M, Vespa PM, Vijayan T, Allyn PR,
495 Campeau S, Humphries RM, Klausner JD, Ganzon CD, Memar F, Ocampo NA, Zimmermann LL,
496 Cohen SH, Polage CR, DeBiasi RL, Haller B, Dallas R, Maron G, Hayden R, Messacar K,
497 Dominguez SR, Miller S, Chiu CY. 2019. Clinical Metagenomic Sequencing for Diagnosis of
498 Meningitis and Encephalitis. New England Journal of Medicine 380:2327-2340.
- 499 29. Anonymous. KneadData software: <http://huttenhower.sph.harvard.edu/kneaddata>.
- 500 30. Segata N, Waldron L, Ballarini A, Narasimhan V, Jousson O, Huttenhower C. 2012.
501 Metagenomic microbial community profiling using unique clade-specific marker genes.
502 Nature Methods 9:811-814.
- 503 31. Chen YE, Fischbach MA, Belkaid Y. 2018. Skin microbiota–host interactions. Nature 553:427-
504 436.
- 505 32. Wilson MR, O'Donovan BD, Gelfand JM, Sample HA, Chow FC, Betjemann JP, Shah MP, Richie
506 MB, Gorman MP, Hajj-Ali RA, Calabrese LH, Zorn KC, Chow ED, Greenlee JE, Blum JH, Green G,
507 Khan LM, Banerji D, Langelier C, Bryson-Cahn C, Harrington W, Lingappa JR, Shanbhag NM,
508 Green AJ, Brew BJ, Soldatos A, Strnad L, Doernberg SB, Jay CA, Douglas V, Josephson SA,

509 DeRisi JL. 2018. Chronic Meningitis Investigated via Metagenomic Next-Generation
510 Sequencing. *JAMA Neurology* 75:947-955.

511 33. Amos GCA, Logan A, Anwar S, Fritzsche M, Mate R, Bleazard T, Rijpkema S. 2020. Developing
512 standards for the microbiome field. *Microbiome* 8:98.

513 34. Wang Y, Song F, Zhu J, Zhang S, Yang Y, Chen T, Tang B, Dong L, Ding N, Zhang Q, Bai Z, Dong X,
514 Chen H, Sun M, Zhai S, Sun Y, Yu L, Lan L, Xiao J, Fang X, Lei H, Zhang Z, Zhao W. 2017. GSA:
515 Genome Sequence Archive*. *Genomics, Proteomics & Bioinformatics* 15:14-18.

516 35. Members NGDC, Partners. 2019. Database Resources of the National Genomics Data Center
517 in 2020. *Nucleic Acids Research* 48:D24-D33.

518

Figure 1: Microbial community structure in human CSF of 23 healthy individuals. (a) Metagenomic experimental design in this study: CSF and matched control samples (positive controls: oral and skin; negative controls: saline solution) collected from 23 pregnant women along with DNA and RNA extraction buffers (number indicates replicates) and were sequenced for metagenomic and metatranscriptomic analysis (see Methods). (b) An overview of microbes detected in each sample type. The number of microbes detected in each sample, and shared species between different samples were shown in the upset plot, with the dots representing intersections among sample types, and the bars representing the number of microbes for each sample type (horizontal bars) and ones shared for each intersection type (vertical bars). The inset shows a box plot summarizing distributions of the number of species detected for different sample types. (c) Circle barplot summarizing the the number of microbial species in each CSF_DNA sample, categorized into three major types: eukaryota, virus and bacteria. (d) Microbial community structures of 23 CSF_DNA samples shown in a stacked barplot that summarizes the relative abundance of different species of microbes detected for each CSF_DNA sample.

Figure 2. Microbiome similarity among sample types. (a) NMDS (Non-metric Multidimensional Scaling) analysis of microbial species detected from different sample types. Shapes and colors represent sample types. (b) Boxplot summarizing the beta diversity within CSF_DNA and between CSF_DNA and other specimens using Bray-Cruits dissimilarity. Statistical significance was assessed by Wilcoxon test whose significance level is indicated with asterisks (***: $P < 0.001$).

Figure 3. Microbes remained in the cerebrospinal fluids. Subtracting the microbes appeared in the negative control and DNA extraction buffer, 14 species (6 species labeled with star appeared in skin samples) remained in CSF genomic samples.

Supplementary Figure S1: Top 10 genus in oral and skin samples, respectively.

Microbial community structures of 23 Oral (figure S1a) and Skin (figure S1b) samples shown in a stacked barplot that summarizes the relative abundance of different genus detected.

550 **Supplementary Figure S2:** Principal Coordinates Analysis (PCoA) analysis of
 551 microbial species detected from different sample types.

552
 553 PCoA (Principal Coordinates Analysis) analysis of microbial species detected from different sample
 554 types. Shapes and colors represent sample types.
 555

556 **Supplementary Table S2: Sample labels and matching information.**

Table S1: Samples label and matching information							
CSF_DNA	Negative	Skin	Oral	CSF_RNA	Swab	RNA_buffer	DNA_buffer
1	1	1	1	1	1	1	1
2	2	2	2		2	2	2
3	3	3	3		3	3	3
4	4	4	4	4	4		
5	5	5	5		5		
6	6	6	6		6		
7	7	7	7				
8	8	8	8	8			
9	9	9	9	9			
10	10	10	10	10			
11	11	11	11				
12	12	12	12				
13	13	13	13				
14	14	14	14	14			
15	15	15	15	15			
16	16	16	16	16			
17	17	17	17				
18	18	18	18	18			
19	19	19	19	19			
20	20	20	20				
21	21	21	21	21			
22	22	22	22				
23	23	23	23	23			
CSF_DNA: Cerebrospinal fluids metagenomic							
CSF_RNA: Cerebrospinal fluids metatranscriptomic							

557