Knockout of *Anopheles stephensi* immune gene LRIM1 by CRISPR-Cas9 reveals its unexpected role in reproduction and vector competence

Ehud Inbar¹, Abraham G. Eappen¹, Robert T. Alford², William Reid^{1*}, Robert A. Harrell², Maryam

Hosseini¹, Sumana Chakravarty¹, Tao Li¹, B. Kim Lee Sim¹, Peter F. Billingsley¹ and Stephen L. Hoffman^{1&}

¹Sanaria Inc. Rockville, MD 20850. ²Insect Transformation Facility, Institute for Bioscience and

Biotechnology Research, University of Maryland, Rockville, MD 20850.

*Current address: Dept. Veterinary Pathobiology, University of Missouri, Columbia MO 65211.

&Correspondence and requests for materials should be addressed to Stephen L. Hoffman

Abstract

PfSPZ Vaccine against malaria is composed of *Plasmodium falciparum* (Pf) sporozoites (SPZ) manufactured using aseptically reared *Anopheles stephensi* mosquitoes. Immune response genes of *Anopheles* mosquitoes such as Leucin-Rich protein (LRIM1), inhibit *Plasmodium* SPZ development (sporogony) in mosquitoes by supporting melanization and phagocytosis of ookinetes. With the aim of increasing PfSPZ infection intensities, we generated an *A. stephensi* LRIM1 knockout line, *Δaslr1*, by embryonic genome editing using CRISPR-Cas9. *Δaslr1* mosquitoes had a significantly increased midgut bacterial load and an altered microbiome composition, including elimination of commensal acetic acid bacteria. The alterations in the microbiome caused increased mosquito mortality and unexpectedly, significantly reduced sporogony. The survival rate of *Δaslr1* and their ability to support PfSPZ development, were

1 partially restored by antibiotic treatment of the mosquitoes, and fully restored to baseline
 2 when *Δaslim1* mosquitoes were produced aseptically. Deletion of LRIM1 also affected
 3 reproductive capacity: oviposition, fecundity and male fertility were significantly
 4 compromised. Attenuation in fecundity was not associated with the altered microbiome. This
 5 work demonstrates that LRIM1's regulation of the microbiome has a major impact on vector
 6 competence and longevity of *A. stephensi*. Additionally, LRIM1 deletion identified an
 7 unexpected role for this gene in fecundity and reduction of sperm transfer by males.

8

Introduction

Sanaria® PfSPZ vaccines, composed of aseptic, purified, cryopreserved *Plasmodium falciparum* (Pf) sporozoites (SPZ) are produced in aseptically reared *Anopheles stephensi* mosquitoes [1-5].

The cost of each dose of vaccine would be reduced significantly if more PfSPZ could be produced per mosquito. When Pf sexual stage parasites are ingested by *Anopheles* spp.

mosquitoes, the parasites mate then transform to ookinetes which penetrate the mosquito midgut and form oocysts, which during sporogony give rise to PfSPZ. After migrating to and

invading the mosquito salivary glands, PfSPZ are inoculated into humans when the mosquitoes feed (reviewed by [6, 7]). One way to increase PfSPZ production in mosquitoes would be to

reduce innate immune responses in the mosquito midgut [8-11] that are thought to inhibit parasite development.

Anopheles spp. mosquitoes possess innate immune systems that regulate microbial infections, including *Plasmodium* [10, 12-15]. Invasion of the midgut epithelium by *Plasmodium* ookinetes

activates a complement-like cascade initiated by a C3-homolog, thioester-containing

glycoprotein 1 (TEP1), that circulates in the mosquito hemolymph [8]. Upon ookinete invasion

of the midgut epithelium, TEP1 is cleaved by proteolysis. Cleaved TEP1 forms a complex with

leucine-rich repeat (LRR) proteins, LRIM1 and APCL1, which is crucial for the stability of the

cleaved form of TEP1 while circulating in the hemolymph [9, 11]. The complex binds to the

ookinete surface and labels it for melanization and phagocytosis [8, 10, 16, 17]. Knock-down of

TEP1 by RNA interference (RNAi) resulted in a 5-fold increase in *P. berghei* (Pb) oocysts in the

midguts of susceptible *A. gambiae* and elimination of melanization in a *P. berghei* refractory

A. gambiae line, L3-5 [8]. Likewise, knock down of LRIM1 or APCL1 by RNAi led to an ~50-fold

increase in oocysts in the midguts of susceptible *A. gambiae* and elimination of melanized parasites in a refractory *A. gambiae* [11].

Based on these findings, we hypothesized that deletion of *A. stephensi* immune genes would increase PfSPZ infection intensities. We have previously knocked down LRIM1 by RNAi using the UAS-GAL4 system in which LRIM1 dsRNA was endogenously expressed [18]. The knock-down resulted in 4-13-fold increase in midgut oocysts and 2-10-fold increase in salivary gland PfSPZ compared to WT [18]. Here, we used CRISPR-Cas9 to generate a stable LRIM1 knockout line ($\Delta aslrim1$) of *A. stephensi*. The LRIM1 knockout line was dramatically more susceptible than the wild type (WT) to bacterial infection, but not more susceptible to PfSPZ infection. Our embryonic genome editing had an unexpected off-target effect, profoundly reducing the reproductive capacity of the mosquitoes.

Results

Generation of LRIM1 knockout line in Anopheles stephensi, using CRISPR-Cas9

Multiple short guide (sg) RNAs (Table S1) were used to target the first and the second exons on the 5' end of *A. stephensi* LRIM1 (*aslrim1*) gene (Vector base - ASTE000814). The sgRNAs were mixed with recombinant CRISPR-associated protein 9 (Cas9) and injected into 549 *A. stephensi* embryos (Table S2). Nineteen out of the 549 eggs hatched (3.5%), from which, only 4 females and 6 males had developed to G₀ adults. The adults were sorted, and backcrossed *en masse* to WT males and females as appropriate. Females were provided with a blood meal and allowed to lay eggs. Twenty G₁ egg pools were collected of which, 8 groups of larvae were tested by

PCR, using primers flanking the expected editing position (see primers 1 and 2 in figure S1A and primers table S3). Sequence analysis of the PCR products revealed a 13 bp deletion in the 5' of exon 2 (nucleotides 942-954), in one group of larvae out of the eight tested (table S2). The adults from the positive group were sorted and outcrossed to WT and after the blood feeding, the females were set to lay eggs individually. Prior to egg laying, a leg was removed from each female, DNA was extracted from each leg and PCR used to screen for females carrying the deletion. The G₂ adults derived from the positive females were then in-crossed and individual females were allowed to lay eggs. Females were collected after egg laying and tested by PCR-sequencing, and G₃ eggs were collected from positive females. Groups of G₃ larvae, derived from each of the positive G₂ females, were tested by PCR with primers designed to detect alleles that were homozygous or heterozygous for the deletion (primers 3 and 4 in Figure S1A). Groups of larvae with the highest frequencies of the deletion alleles were continued on and the cycle repeated itself until a homozygous female was found in G₈. The G₉ larvae from that female were all homozygous, demonstrating that the null deletion was fixed in the mosquito line. PCR and sequence analysis on individual mosquitoes from G₁₂ indicated that the deletion allele remained stable in the population in later generations (Figures S1B and S1C). Importantly, no evidence for the WT LRIM1 allele was seen in any of the mosquitoes tested, indicating the establishment of a stable, homozygous LRIM1 knockout line. This line will be referred to as *Δaslrin1* hereafter.

To confirm that the deletion resulted in silenced gene transcription, we performed qPCR on cDNA made from mRNA from different life stages of WT and *Δaslrin1* mosquitoes using two different reactions. In the first reaction the primers were targeting a region on the transcript

that did not include the 13 bp deletion; therefore, this reaction should have amplified the fragment from both the WT and the *Δaslr1* lines (Figure S2A, primers 5 and 6). In the second reaction, the 3' end of the reverse primer was anchored in the deletion, which should not have resulted in amplification in the *Δaslr1* line (primers 7 and 8). Approximately 30% decrease in the total abundance of LRIM1 transcript was observed in all tested life stages of *Δaslr1* when qPCR was done with the first reaction (primers 5 and 6), suggesting that while the transcript was still present, its abundance was somehow affected by the deletion. As expected, there was no amplification of *Δaslr1* using the second qPCR reaction (primers 7 and 8) in any of the different life stages, indicating that LRIM1 is not expressed in the *Δaslr1* line. The functional relationship between LRIM1 and TEP1 [17] prompted us to check whether transcription of TEP1 was affected by the 13 bp deletion in LRIM1 and/or its potential loss of function. No significant change was observed in TEP1 transcription TEP1 in all life stages of *Δaslr1* compared to WT (Fig S2A).

The deletion of 13bp from the gene was predicted to cause a frame shift in the amino acid sequence of the protein at positions 57 to 62, which would potentially result in a stop codon, leaving only a small portion (62 amino acids) of the N-terminus of the protein. A polyclonal antiserum was raised against a 20 amino acid peptide in the middle of the LRIM1 protein, starting in position 84 which should not have reacted with protein in *Δaslr1*. Western blot analysis of hemolymph extracted from both WT and *Δaslr1* demonstrated clearly that LRIM1 protein was not present in the hemolymph of the *Δaslr1* line (Figure S2B).

LRIM1 controls the midgut microflora and thereby longevity of A. stephensi

We investigated the effect of LRIM1 deletion on microbial populations in the mosquito midgut.

First, we assessed the general bacterial loads in the mosquitoes using qPCR. DNA was

extracted from pools of 10 mosquitoes, both WT and *Δaslr1*, (5 pools from each line) and

bacterial loads were analyzed by real-time PCR using general 16S rDNA primers, targeting

variable region 4 [19] (Figure 1A and see primers 13-14 table S3 for primers). Mosquitoes were

washed in ethanol and twice in sterile PBS to prevent inclusion of bacteria from the mosquito

surface in the extractions. The total bacterial load in non-fed *Δaslr1* was 52 ± 18 -fold higher

than in WT. Similar results were observed when using a culture-based approach when colony

forming units (CFU) in *Δaslr1* midguts were approximately 200-fold higher than the CFU in

WT midguts (Figure S4E). In the WT mosquitoes, blood feeding led to an increase in total

bacterial loads by 21 ± 8 -fold. Uptake of blood increased total bacterial load in *Δaslr1* by 2.6-

fold. However, the variability in the bacterial quantity in this line was high and the results

between non blood fed and blood fed were not statistically significant.

To evaluate the effect of the LRIM1 deletion on the microbial community composition in the

mosquitoes, the DNA pools from both lines were assessed by 16S ribosomal RNA (rRNA) gene

amplicon sequencing [20], and bacteria identified to genus level. Alpha diversity indices

(calculated at a sequencing depth of 12,000 sequences/sample) indicated significantly lower

microbial richness in *Δaslr1* relative to WT (Figure 1B), but this phenomenon did not

manifest itself in significantly different Shannon index values due to higher evenness in

Δaslr1 samples (Figure S3A,B). Microbial community structure was significantly different between groups, as measured using analysis of similarity (ANOSIM) on Bray-Curtis similarity values (Figure S3C and S3D). The microbial communities in *Δaslr1* mosquitoes were significantly different as compared to WT, and within-group *Δaslr1* microbial communities were also more similar to each other than within-group WT communities (Figure S3D). Using DEApp software (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5291987/>) DESeq2 analysis showed that 16 taxa were significantly different in abundance between groups, with 15 of them lower in *Δaslr1* relative to WT (Figure 1C, Table 1). The most reduced bacteria in *Δaslr1* mosquitoes were from the genera *Asaia* and *Tanticharoenia*, as well as an unidentified genus of *Acetobacteraceae*. Interestingly, *Acetobacteraceae* are Gram negative bacteria that oxidize sugars to acetic acid during fermentation, some of which are known to be insect commensals, specifically of *Anopheles* mosquitoes [21-23]. A heatmap was generated of the 10 most abundant genera in WT mosquitoes and relative abundance compared with *Δaslr1*. Proteobacterial genera (*Enterobacter*, *Klebsiella*, *Serratia*) and a genus of *Flavobacteriales* (*Elizabethkingia*) were variably abundant in all animals, but were not significantly different between groups (Figure 1C). Again, the absence of the acetic acid bacteria can be clearly observed. Overall, deletion of LRIM1 led to significant alterations in the mosquito midgut microbiome.

We noticed high mortality of the *Δaslr1* adults and hypothesized that this resulted from the high numbers of bacteria. We therefore monitored the mortality of WT and *Δaslr1* mosquitoes with and without addition of penicillin-streptomycin (PS) in their sucrose meals. Providing 1% but not 0.5% PS eliminated internal bacterial populations from the WT and

Δaslr1 mosquitoes [24, 25] and did not compromise their survival and fecundity (Figure S4). With PS treatment, the longevity of *Δaslr1* mosquitoes increased while there was no effect on WT (Figure 2A). The mosquitoes were blood fed three days after collection. At that time point, over 90% of the WT mosquitoes were still alive whether they fed on regular sucrose or sucrose with PS. The survival of the *Δaslr1* was only $69 \pm 7\%$ on the day of blood feeding. The survival of the *Δaslr1* was improved ($81 \pm 4\%$) when PS was added to the sucrose meal. A big difference in the mortality was observed 6 days after the collection of the mosquitoes (3 days post bloodmeal), when only $20 \pm 3\%$ of the *Δaslr1* were still alive compared to $84 \pm 10\%$ of the WT ($P < 0.05$). The survival in *Δaslr1* was partially rescued by feeding on PS, as $54 \pm 10\%$ of the mosquitoes were still alive at this point. A dramatic drop in the survival of *Δaslr1* was observed 8- and 14-days post collection, when only $8 \pm 5\%$ and $6 \pm 2\%$ of the mosquitoes had survived respectively, compared to $80 \pm 9\%$ and $42 \pm 7\%$ survival of WT at these time points, respectively ($P < 0.05$). Again, a significant, rescue of the survival in *Δaslr1* was observed 8-and 14- days after collection, with the addition of PS to the sucrose meal. The results suggest a profound role for LRIM1 in determining the longevity of the mosquitoes via controlling their internal microflora.

LRIM1 is indirectly important for development of the Pf in non-aseptic A. stephensi

The compliment-like cascade, and specifically LRIM1, are considered pivotal in the interactions between *Plasmodium* parasites and mosquito vectors. To assess the effect of the LRIM1 knockout on growth and development of Pf, we infected WT and *Δaslr1* mosquitoes, grown

1 under normal (non-aseptic) and aseptic conditions, with Pf (Figure 2). Two-three day old WT
2 and *Δaslr1* adult mosquitoes were maintained non-aseptically on 15% sucrose ± 1% PS for
3 three days, prior to blood feeding. Seven days post blood feeding, midguts were dissected and
4 oocyst infection intensities were assessed (Figure 2B). In WT mosquitoes, the oocyst intensity
5 was 7.9 geometric mean (GM) oocysts/midgut (95% confidence interval (CI) = 7.8-17.3), with a
6 prevalence of 90%. Only 6 *Δaslr1* female adults survived to 7 days post feeding and
7 surprisingly, none were infected. Addition of PS to the sugar meal of the WT mosquitoes
8 significantly increased the infection to 50.5 oocysts/midgut (95% CI = 57-105.3) and prevalence
9 remained unchanged at 90% (Figure 2B). PS partially rescued the infection in *Δaslr1* with an
10 intensity of 3.2 (95% CI = 2.3-10.6) oocysts/midgut and prevalence of 60%. Due to the high
11 mortality and the modest rescue by PS in *Δaslr1* mosquitoes, we repeated the experiment
12 starting with more mosquitoes and treating them with PS for their entire adult life, including
13 throughout Pf development (Figure 2C). Oocyst infections were significantly lower in *Δaslr1*
14 (GM = 2 oocysts/midgut, 95% CI= 2.5-9.2) versus WT (GM = 50.8 oocysts/midgut, 95% CI =
15 52.3-104.8) ($p < 0.0001$). Unlike the previous observation (Figure 2B), feeding on PS by WT
16 mosquitoes reduced infection intensity slightly (GM = 32.2 oocysts/midgut, 95% CI = 26.4-61.4)
17 and prevalence remained high (100%). Addition of PS to the sugar meal of *Δaslr1*
18 mosquitoes again partially recovered the infections (GM = 8.6 oocysts/midgut, 95% CI = 10.5-
19 31.7, prevalence = 73%). In contrast, under aseptic conditions, the infection intensities
20 between WT (GM = 29.6 oocysts/midgut, 95% CI = 27.3-56) and *Δaslr1* (GM = 29.9
21 oocysts/midgut, 95% CI = 28.9-51.4) were almost identical and the prevalence was 94% for
22 both lines (Figure 2D). A small, non-significant difference was observed in intensities of PfSPZ

infections; 35,824 PfSPZ/mosquito (95% CI = 41085-87442) in the WT versus 11,590 PfSPZ/mosquito (95% CI = 25469-56433) in *Δaslr1* (Figure 2E). The infection prevalence was 100% and 95% in WT and *Δaslr1*, respectively. Overall, under normal growth conditions, oocyst infection intensities in *Δaslr1* were significantly lower than in WT mosquitoes. This decrease was partially rescued by addition of PS to the sugar meal, and fully rescued when the mosquitoes were grown aseptically.

A role for LRIM1 in mosquito fecundity and reproduction

As there was a profound attenuation in the reproductive capacity of *Δaslr1* compared to their WT counterparts, we examined fecundity (egg production and egg hatching rate) in females and fertility in males. WT and *Δaslr1* mosquitoes were provided with 15% sucrose ± 1% PS from the day of adult emergence. Females were provided with a bloodmeal 1 week post emergence and engorged females were separated into cages. Three days post bloodmeal, females from WT, WT+PS, *Δaslr1* and *Δaslr1*+PS, were placed individually into *Drosophila* tubes and allowed to oviposit. The number of females that laid eggs was determined and the eggs were counted in 34-35 tubes collected randomly from each group (Table 2). The majority of WT females had laid eggs, 92% and 95% of the WT and WT+PS females, respectively while only 61% and 55% of the *Δaslr1* and *Δaslr1*+PS, respectively, had oviposited. The number of eggs laid by individual *Δaslr1* females was reduced significantly to 55.7 (95% CI = 47.2-65.8) and 60.9 (95% CI = 54.9-67.6) in the absence and presence of PS, respectively, from WT 111.1 (95% CI = 104.2-126.0) and 97.6 (95% CI = 92.3-114.1) in the absence or presence of PS,

1 respectively (Figure 3A, Table 2). Deletion of LRIM1 also resulted in significant decrease in
2 hatching rates (Figure 3B and Table 2). In WT mosquitoes, the mean egg hatching rate 52.9%
3 (95% CI = 37.5-74.5) and 72.4% (95% CI = 66.7-78.6) in WT and WT + PS, respectively. In
4 *Δaslr1*, egg hatching was significantly lower ($p < 0.0001$) at 22.8% (95% CI = 13.5-38.7) and
5 25% (95% CI = 17.5-36) for *Δaslr1* and *Δaslr1* + PS, respectively. Altogether, deletion of
6 LRIM1 resulted in a significant reduction in fecundity in terms of oviposition rate, number of
7 laid eggs per female and egg hatching rate. None of these were rescued by the addition of PS
8 to the sucrose meal, suggesting that low fecundity in *Δaslr1* was not associated with the
9 increased bacterial loads. To determine whether the low number of eggs was due to a
10 reduction in blood intake by the *Δaslr1* mosquitoes, females were randomly collected
11 immediately after the bloodmeal and the volume of bloodmeal was determined). The
12 geometric mean volume of bloodmeal taken by WT mosquitoes was 4.6 μL (95% CI = 4.2 - 5.0 μL)
13 while in *Δaslr1* females it was only 2.7 μL (95% CI = 2.0 - 3.6 μL) μL of blood (1.7-fold less,
14 $P < 0.0001$) (Figure 3C). *Anopheles* mosquitoes concentrate host blood cells and proteins,
15 simultaneously excreting excess salts and water, in a process called prediuresis, which is
16 exemplified by the release of large blood-colored droplets during feeding [26, 27]. During
17 rearing of the mosquitoes, we noticed a dramatic reduction in prediuresis products in *Δaslr1*
18 mosquitoes compared to WT. To demonstrate this difference, we transferred 50 WT and
19 *Δaslr1* females each to a paper 473 mL paper container and placed a round filter paper at
20 the bottom of the container. After feeding, more prediuresis products were seen on filter
21 papers from the WT containers compared to filter papers from the *Δaslr1* containers (Figure
22 3D), indicating that bloodmeal processing is severely disrupted by the LRIM1 deletion. Total

blood meal protein was measured in both lines to determine if the difference in prediuresis was also manifested in lower acquisition of proteins from the bloodmeal (Figure 3D). The geometric mean of blood meal protein contents/ midgut in *Δaslr1* was 553 μg protein/midgut (95% CI = 451.8 -676.1 μg protein/midgut) compared to 706 μg protein/midgut (95% CI = 604.0 – 826.5 μg protein/midgut) in the WT. The reduction in blood intake and/or in prediuresis in *Δaslr1* did translate to a significant difference in midgut protein contents.

LRIM1 has a role in male fertility

The significant decrease in oviposition and hatching rate in *Δaslr1* mosquitoes could be due to reduced fertility of *Δaslr1* males. To assess whether the deletion of LRIM1 had an effect on male fertility, WT female pupae were sorted and crossed with either WT (W-W) or *Δaslr1* (W-L) males. Mosquitoes were reared to adults, females were provided with a bloodmeal, and the number of eggs per female and the proportion of fecund females were determined 1-day post egg laying. The hatch rate was determined 3-days post egg laying. The females that did not lay eggs were dissected and examined microscopically for the presence of eggs in the ovaries and sperm in the spermatheca (Table 3). Bloodmeal size was the same in W-W and W-L females (data not shown), confirming that blood intake was not a factor in oviposition outcome or the number of eggs. In the W-W, 60.3% of the females laid eggs compared to only 14.3% in W-L, consistent with the reduced percentage of egg laying females observed previously in *Δaslr1* (Table 2). The mean fecundity in W-L was moderately but significantly ($P<0.05$) lower than that of W-W cross (89.1 ± 10.5 eggs/female and 115.3 ± 5.2 eggs/female,

respectively) (Figure 4A). An equal proportion of females that did not lay eggs in W-W and W-L, had eggs in the ovaries (47.1% and 46.7%, respectively). Sperms were detected from the spermathecae of 52.9% of females that did not lay eggs in W-W and only 22.2% in W-L. These data suggest that *Δaslr1* males have a reduced capacity to inseminate females or that the number of sperm that they deposit in the females is below the detection level. However, *Δaslr1* and WT males had similar numbers of sperm in their testes (Figure 4B).

LRIM1 is expressed in the hemolymph of *A. stephensi* and *A. gambiae* (See figure S2B and [11]), and in the midgut of *A. gambiae*, 24-48 hours following an infected bloodmeal [28]. Due to the effect of LRIM1 deletion on both fecundity and male fertility, we looked for LRIM1 expression in the reproductive organs of the mosquitoes. RNA was extracted from pools of male reproductive organs (MRO - testis, male Accessory glands, vas deferens and ejaculatory duct) and ovaries, from 30 sugar-fed 4-6-day old male and female mosquitoes, respectively. The LRIM1 mRNA abundance in those tissues relative to whole mosquitoes was assessed using real-time PCR with LRIM1 specific primers (primers 7 and 8 table S3). LRIM1 mRNA was significantly more abundant in the male reproductive organs compared to whole mosquitoes. Conversely, we did not detect LRIM1 mRNA in the ovaries (Figure 4C). Western blot analysis indicated expression of LRIM1 protein on male reproductive organs from WT but not *Δlr1* males.

Discussion

We generated LRIM1 knock out (*Δaslrin1*) *A. stephensi* to reduce the innate immune responses in the midgut against Pf ookinetes and oocysts, with the aim of improving PfSPZ numbers for manufacturing PfSPZ products. Using CRISPR-Cas9, we were able to generate a deletion in the LRIM1 coding region that completely prevented expression of LRIM1 protein. Deletion of the LRIM1 gene had a profound impact on the quantity and diversity of the mosquito midgut microbiome, increasing total bacterial load and reducing midgut microflora diversity and richness. The *Δaslrin1* mosquitoes were colonized predominantly by known mosquito commensals such as the proteobacteria *Enterobacter*, *Klebsiella*, *Serratia* and *Flavobacteriales*, as well as by *Elizabethkingia* but other classes such as the acetic acid bacteria were lost from the midgut. The removal of the acetic acid bacteria and specifically the *Asaia* genus, is of a particular interest because it is an important commensal bacterium that is highly abundant in *Aedes* and *Anopheles* species, particularly, *A. stephensi* [21, 29-31]. The *A. stephensi* abdomen is colonized predominantly by different *Asaia* species which account for 41%, 25% and 20% of the total population in the gut, salivary glands and female reproductive system, respectively [29], and 58% of the bacterial population in the male reproductive system. The changes in the mosquito microbiome and the bacterial overgrowth in *Δaslrin1* mosquitoes significantly reduced their survival, indicating the pivotal but probably indirect role of LRIM1 in mosquito longevity.

Contrary to our expectations, PfSPZ infection intensities did not increase in *Δaslrin1*. Under normal rearing conditions, there was a dramatic reduction in PfSPZ intensities which was partially rescued by treating the mosquitoes with antibiotics, and completely rescued when mosquitoes were reared under aseptic conditions. LRIM1 appears crucial for the development of Pf in non-aseptic *A. stephensi*, most likely by regulation of the mosquito microflora. Therefore, establishing infection and completion of the sporogonic cycle by the parasite, depends heavily on the interactions between the parasite and the internal microflora. The concept of parasite-microbiota interactions inside the mosquito is not new and the antiparasitic effect of bacteria has been demonstrated for a variety of *Anopheles* and *Plasmodium* species [32]. Most of these studies suggest that microbiota-related parasite killing is achieved mainly by the stimulation of the mosquito immune response by the bacteria [33-36]. However, the results of our work, specifically the overwhelming increase in bacterial loads in *Δaslrin1* mosquitoes, strongly suggest that even with a complete immune response, competition for nutrients and/or reduced fitness due to increased bacterial load also plays a major part in determining the ability of the parasite to develop optimally in the mosquito. Whether the anti-parasitic effect of the microbiota is through stimulation of the immune response or simply by competition for resources or reduced mosquito fitness, LRIM1 is important in controlling the microbiome and thereby providing the parasite with an environment in which it can flourish.

The decrease in PfSPZ infection intensity following the LRIM1 deletion was contrary to what had been reported by us and by others, showing an increase in either *P. berghei* (Pb) and Pf

oocyst or PfSPZ numbers upon the knock-down of LRIM1 by RNAi [11, 17, 18, 37]. The differences are likely due to the different genetic manipulation approaches that were taken. Unlike in gene knockouts, RNAi does not lead to a complete elimination of the targeted proteins. In line with this, reduced expression through RNAi of a fibrinogen-related protein 1 (FREP1) resulted in only partial protein depletion and subsequently, partial reduction in midgut oocyst infections (50%) [38]. In contrast, complete knockout of FREP1 protein by CRISPR-Cas9 resulted in a much stronger suppression of the infection (~80%) [37, 39]. Moreover, the knock-down of LRIM1 in *A. stephensi* by RNAi resulted in significant increase in Pf infection intensity [18]. However, in that work, at least 40% of LRIM1 transcript was still present at all life stages tested. Importantly, the knock-down did not affect the bacterial load in the transgenic mosquitoes and this is likely to be the main reason for the differences between the knock-down of LRIM1 and its complete elimination by CRISPR-Cas9, in the present work. LRIM1 dsRNA was injected into 1-2-day old adult *A. gambiae*, which were infected a few days later [11, 17], probably too short a time frame for the excessive bacterial burdens to establish compared to the present work, where the LRIM1 deletion was permanent, affecting mosquitoes right from the early stages of development. In summary, while we cannot rule out that LRIM1 and the complement-like system interacts directly with the parasites as suggested previously, the comparable infections in aseptically reared WT and $\Delta aslr1$ mosquitoes suggest that the effect of LRIM1 on Pf infection is indirect and mediated by the mosquito microbiota.

Deletion of LRIM1 significantly reduced the fecundity and reproduction of the *Δaslrin1* mosquitoes which could not be rescued by addition of antibiotics to the mosquito sucrose meal, suggesting that the role of LRIM1 in these processes was not mediated by the microbiota. This reduction in fecundity maybe explained at least in part by the reduction in blood intake and in prediuresis by *Δlrin1* females. A direct role of LRIM1 in oogenesis is yet to be defined even though there is some evidence for an association between oogenesis and the complement-like system. Vitellogenin (Vg) is a nutrient transporter, essential for delivering digested bloodmeal peptides to the maturing mosquito oocytes [40, 41]. Disruption of Vg resulted in defects in egg development and in parallel, significant reduction in TEP1-mediated ookinete killing [42]. MosGILT, a mosquito saliva protein, is also involved in this process and its knock-down led to a profound impairment in ovarian development that was coupled with significant reduction in TEP1-mediated parasite killing [43]. In another study, deletion of FREP1, a *Plasmodium* agonist, in *A. gambiae*, led to a significant reduction in infection intensity by both *P. berghei* and *P. falciparum* but also had a profound fitness cost in which blood ingestion, fecundity and egg hatching all decreased significantly [39], supporting the premise of association between parasite infection and reproduction capacity.

In *Anopheles*, oviposition depends heavily on male's ability to inseminate females and fertilize the eggs [44, 45]. Thus, the reduction in egg laying might also be explained by the effect of LRIM1 deletion on the fertility of the males and their ability to fertilize the eggs. Despite the fact that LRIM1 is expressed in the male reproductive organs, sperm counts in *Δaslrin1* males were comparable to those in WT. Therefore, the role of LRIM1 in male fertility is not based on

the ability of the males to produce sperm but rather on the quality of the sperm, the seminal fluids, and/or on the ability of the males to inseminate the females. The involvement of the complement-like system in determining the quality of the sperm was demonstrated previously [46]; in *A. gambiae*, TEP1 binds to the surface of damaged sperm in the testes, labeling them for removal and thereby allowing for a high rate of healthy sperm production. In this context, the supporting role of LRIM1 in stabilizing the active form of TEP1 [9, 17] may also be relevant in maintaining sperm quality.

It is interesting to note that acetic acid bacteria and specifically *Asaia*, that were removed from the microflora in $\Delta aslrim1$ mosquitoes, typically populate the reproductive organs of male and female *A. gambiae* and *A. stephensi* [21, 23, 29]. Moreover, in these mosquitoes, *Asaia* is transmitted horizontally from males to females through mating and vertically from the female to the eggs [29, 47]. We do not have evidence for an association between the elimination of bacteria species such as *Asaia* and the low fecundity in the $\Delta aslrim1$, but it is possible that the regulation of internal microbiota by LRIM1 allows important commensal bacteria such as *Asaia* to successfully colonize the reproductive organs, and in the absence of LRIM1, such regulation may be disrupted to see overgrowth of other species.

The unexpected reduction in mosquito fecundity following LRIM1 deletion was independent of the role of LRIM1 in controlling the midgut microbiota, inferring that LRIM1 has an additional, unidentified role in the reproductive capacity of the mosquitoes. This demonstrates the power of precise CRISPR-Cas9 genome editing to reveal novel functions of targeted genes while raising serious concerns about our ability to accurately attribute specific biological functions to a gene. Such off-target effects are important considerations for embryonic gene

1 editing by CRISPR-Cas9 in other species, including humans, where creation of adverse
2 phenotypes might not be manifested until later in life [48-50].

3 Our goal was to increase PfSPZ yields in *A. stephensi* and thereby improve PfSPZ production by
4 deleting the gene encoding the *A. stephensi* immune deficiency protein, LRIM1. The resultant
5 mutant line, *Δaslr1*, was significantly more susceptible to bacteria but not to Pf and
6 moreover, was severely compromised in reproductive capacity. Therefore, this line of
7 mosquitoes will not be useful for the manufacturing of PfSPZ vaccines. We are currently
8 testing the possibility of improving PfSPZ yields by deleting other innate immune responses
9 genes.

10

11

Table 1. Relative abundance of bacterial genera, *Δaslr1* vs wild-type.

Taxon (Genus-Level)	Base Mean	Log2 Fold Change	Padj value
Asaia	2581.84	-14.56	6.38E-21
Tanticharoenia	166.62	-12.24	1.15E-09
Caulobacter	6.64	-7.58	5.90E-03
Aquabacterium	3.89	-6.82	1.57E-02
Niabella	2.73	-6.24	3.58E-02
Enterococcus	1.96	-6.09	3.87E-02
Cellvibrio	10.35	-6.08	4.24E-04
Acetobacteraceae	2100.01	-4.91	1.90E-09
Variovorax	21.36	-4.51	5.93E-04
Deinococcus	6.64	-4.28	8.09E-03
Leptothrix	7.61	-3.99	7.73E-03
Sphingobacteriales; env.OPS 17	15.76	-3.27	2.54E-02
Sphingomonas	12.97	-3.02	5.90E-03
Chloroplast	8.09	-2.99	3.32E-02

Unassigned	82.2	-1.99	3.40E-02
Acetobacter	2.62	4.17	4.25E-02

1

2

1

Table 2. Fecundity in wild-type and Δ aslr1 *Anopheles stephensi* grown on 15% sucrose with or without pen strep

Group	Number of females	Proportion of females ovipositing (%)	Geometric mean number of eggs laid per female (95% Confidence interval)	Geometric mean % Hatching (95% Confidence interval)
WT	34	95.2	111.1 (104.2-126)	52.9 (37.5-74.5)
WT + PS	35	84.9	97.6 (92.3-114.1)	72.4 (66.7-78.6)
Δ aslr1	34	61.4	55.7 (53.1-68.7)	22.8 (13.5-38.7)
Δ aslr1 + PS	35	55.4	60.9 (57.1-70.3)	25 (17.5-36)

4

5

Table 3. Egg production in WT females and oviposition following a cross with Δ aslr1 or WT males

Cross	#Live males ²	#Live females		% Females laying eggs	% Females with eggs in the ovaries [*]	% Females inseminated [*]
		Feeding ¹	Egg laying ²			
♂WT X ♀WT	63	43	27	60.3	47.1	52.9
♂ Δ aslr1 X ♀WT	70	33	12	14.3	46.7	22.2

* Percentage is calculated from the number of females that did not lay eggs. ¹ The numbers were

determined at the day of blood feeding. ² The numbers were determined at the day of egg laying.

9

10

11

12

13

Figure legends

Figure 1. Effect of LRIM1 deletion on the mosquito microflora. (A) Real-time PCR

quantification of the bacterial population densities in the mosquitoes. The reaction used 16S rDNA primers, 515F and 806R (primers 13 and 14, table S3), targeting the bacterial V4 region of the SSU rDNA [51]. The PCR was done on DNA from pools of 10 females from each mosquito line (WT and *Δaslr1*) before and after blood feeding. The results represent the mean of 5 replicates ± SD. Mosquito S7 rDNA was used as housekeeping gene. (B) Alpha diversity richness analysis (to genus) in WT and *Δaslr1* indicating higher microbial richness in WT relative *Δaslr1* (Mann-Whitney U, P=0.047). (C) Heatmap of the 10 most abundant microbial taxa across mosquito lines. Depth of sequencing ranged from 12827 to 65,296 sequences/sample (mean=38,682; median=31,785).

Figure 2. Effect of deletion of LRIM1 on mosquito longevity and *Plasmodium falciparum*

infection intensity. (A) Survival of WT and *Δaslr1* *A. stephensi* over a period of 14 days. Each point represents the mean ± SD of survival percentage in three different cages. * p≤0.05. (B-C) Oocyst numbers in non-aseptic mosquitoes. Oocysts were counted by microscopy 7 days post bloodmeal for WT and *Δaslr1* mosquitoes grown in non-aseptic conditions and fed on 15% sucrose with or without 1% penicillin/streptomycin (PS). The PS was added to the sugar meal for a short period (3- days prior to the bloodmeal) (B) or, throughout the entire adult life of the mosquitoes (C). Each point represents the oocyst number in a single midgut. (B) n=22, 6, 20

and 20 for WT, *Δaslr1*, WT + PS and *Δaslr1* + PS, respectively. (C) n=20, 19, 18 and 22 for WT, *Δaslr1*, WT + PS and *Δaslr1* + PS, respectively. The results were analyzed in non-parametric, Kruskal-Wallis and Dunn's multiple comparisons tests. (D) The number of oocysts/midgut in WT and *Δaslr1* mosquitoes grown under aseptic conditions, n=18 and 17 for WT and *Δaslr1*, respectively. The results were analyzed by the non-parametric, Mann Whitney test. (B-D) The results are expressed as a geometric means $\pm$ SD. Since some values were zero, a value of 1 was added to the entire dataset to allow calculation of the geometric means. (E) Number of sporozoites/mosquito in mosquitoes grown in aseptic conditions. The numbers of sporozoites were determined by dissecting the salivary glands of the mosquitoes and counting sporozoites from each mosquito by microscopy. The results are expressed as the geometric mean $\pm$ SD (n=20 for both groups). The results for panel E were analyzed by the non-parametric, Mann Whitney test. For all panels, NS p>0.05, * p≤0.05, ** p≤0.01 *** p≤0.001, ****p≤0.0001.

Figure 3. Fecundity in WT and *Δaslr1* mosquitoes. (A) Number of eggs/ females. The number of eggs was determined by counting the eggs laid in single *Drosophila* tubes. The results are expressed as the geometric mean of the number of eggs/females $\pm$ SD; n=34 for WT and 35 for the other 3 treatments. (B) Percentage of eggs hatching. The hatching was determined by counting the number hatching larvae in the single *Drosophila* tubes by microscopy, 2 days after egg laying. The percentage is of the total number of eggs laid in that tube (panel A). The results are reported as the geometric mean of the percent hatching $\pm$ SD; n=34 for WT and 35 for the other 3 treatments. Since some values were zero, a value of 1 was

added to the entire dataset to allow calculation of the geometric means. (A and B) The data were analyzed by Kruskal-Wallis and Dunn's multiple comparisons tests. **** $p \leq 0.0001$. (C) The bloodmeal volumes in WT and *Δaslr1* mosquitoes immediately after blood feeding. The results represent the geometric mean of bloodmeal volumes $\pm$ SD; $n=20$. (D) Remnants of blood on the filter paper after bloodmeal of 50 females, as an indication of prediuresis. (E) Bloodmeal protein contents in midguts of WT and *Δaslr1* mosquitoes, immediately after bloodmeal. The protein contents were determined using Lowry protein assay. The results represent the geometric mean of midgut protein contents in micrograms $\pm$ SD; $n=20$. (C and E) The results were analyzed using non-parametric, Mann Whitney test. ** $P \leq 0.01$, **** $P \leq 0.0001$.

Figure 4. Effect of deletion of LRIM1 on fertility in male mosquitoes. (A) Number of eggs laid in single *Drosophila* tubes by WT females, mated with WT (W-W) or *Δaslr1* (W-L) males. The results are the mean number $\pm$ SD ($n=38$ and 10 for W-W and W-L, respectively). Results were analyzed by unpaired T-test. (B) The number of sperms/males was determined by dissecting the male testes and accessory glands from each male and counting the number of sperms in hemocytometer under the microscope. The results are presented as the geometric mean $\pm$ SD ($n= 40$ and 31 for WT and *Δaslr1* respectively). The data were analyzed using non-parametric, Mann Whitney test. Since some values were zero, a value of 1 was added to the entire dataset to allow calculation of the geometric means. (C) Relative abundance of LRIM1 mRNA in ovaries (OVA) and male reproductive organs (MRO) of WT mosquitoes was determined by real-time PCR using primers specific to LRIM1 (primers 5 and 6, Table S3). The abundance was normalized to the mosquito S7 rDNA gene (Primers 9 and 10, table S3). The

results in panel C represent two independent experiments with similar results ** $P \leq 0.01$. (D) Western blot analysis done on pools of male reproductive organs (MRO) from 30 WT and *Δaslr1* male mosquitoes. The proteins were transferred to PVDF membrane and reacted with 1:250 rabbit polyclonal anti LRIM1, generated in this work. Betta actin (1:1000) was used as a loading control.

Figure S1. Generation of *Δaslr1* mosquito line. (A) PCR strategy for detection of indels in mosquitoes during the CRISPR procedure. Primers 1 and 2 flank the expected deletion site (marked in red) and are used to amplify both the WT and the deletion alleles. The 3' end of primer 3 (13 bp, marked in red) is anchored in the deletion and thus the primer should anneal only to the WT allele. One base pair of the 3' end of primer 4 is anchored upstream to the deletion while the other 17 bp are anchored downstream for the deletion and thus PCR with this primer should only amplify the deletion allele. (B) Alignment showing Homozygous deletion of 13 nucleotides in LRIM1 gene in 10 randomly collected *Δaslr1* mosquitoes from G₁₂. The alignment was done on sequences generated from PCR done with primers 1 and 2. (C) Diagnostic PCR amplifying the WT allele (upper panel, primers 1 and 3) and the deletion allele (lower panel, primers 1 and 4) in 2 WT and 10 randomly collected G₁₂ *Δaslr1* mosquitoes.

Figure S2. Conformation of LRIM1 knockout at the RNA and protein levels. (A) Upper panel - Scheme of the real-time PCR strategy. The nucleotide position indicated refers to the position of the primers on the LRIM1 gene (ASTE000814). PCR I was done with Primers 5 and 6 (Table S3) and targets a region downstream to the deletion and therefore should amplify both the WT and the Deletion alleles. PCR II is done with primers 7 and 8 and is aimed to amplify only the WT allele as the 3' end of primer 5 is anchored in the deletion. Lower panel- Real-time PCR

done with the above primer sets. Results are the mean mRNA abundance $\pm$ SD of 4 different replicate RNA samples from larvae, pupae and female and male adults. Each sample is a pool of 10 individuals from each of the life stages. The results are indicated as the LRIM1 or TEP1 mRNA abundance relative to the housekeeping, S7 rDNA gene (Primers 9 and 10, table S3). (B) Western blot analysis: Hemolymph from 7 female mosquitoes were loaded in each well in the gel and transferred to PVDF membranes. Membranes were reacted with anti 1:250 LRIM1 antiserum generated in this work. The expected molecular weight of LRIM1 protein is 58.3 kDa. For loading control, membrane was reacted with 1:500 AsTEP1 antiserum (ABBIOTECH # 250881).

Figure S3. Alpha diversity and analysis of similarity in WT and *Δaslr1* mosquitoes

(A-B) Comparison of alpha diversity. (A) Shannon index (log base e) of microbial communities in WT and *Δaslr1* mosquitoes. (B) Microbial community evenness. The differences in both measures were not statistically significant (MWU, $P=0.4034$ and MWU, $P=1$ in A and B, respectively). Alpha diversity indices were calculated on rarefied datasets (12,000 sequences/sample). (C) Metric Multidimensional Scaling (mMDS) plot of Mosquito-associated microbial communities. Analysis was performed at the taxonomic level of genus. Data was $\log(x+1)$ transformed, and Bray-Curtis similarity was calculated for all pairwise comparisons. (D) Bray-Curtis similarity values were represented in two dimensions, and 2D stress was 0.08.

Analysis of similarity (ANOSIM) indicated that microbial communities between the two mosquito lines were significantly different ($R=0.524$, $p=0.008$).

Figure S4. The effect of Pen Strep addition to the sucrose meal on midgut bacterial loads, survival and fecundity. (A) Survival of WT mosquitoes maintained on 0%, 0.5% and 1% (v/v) of Penicillin-Streptomycin (PS) solution (500 U/mL), diluted in 15% sucrose. The survival Percentage is the number of live mosquitoes in each time point relative to the number of mosquitoes put in the cage on day zero (30). (B) Mean $\pm$ SD of the number of eggs per females grown on different PS concentrations. The mosquitoes were provided with bloodmeal 3 days after transferring them to cages and individual mosquitoes were put in *Drosophila* tubes for oviposition 4 days after bloodmeal. Eggs were counted in each tube. (C) The number of larvae in each *Drosophila* tube was determined 1-2 days post oviposition. The results show the % of larvae out of the total number of eggs laid in that particular tube. (B, C) $n=4$, 5 and 5 for 0%, 0.5% and 1% PS, respectively. (D) Colony forming units (CFU) in individual guts of WT mosquitoes in different PS concentrations. Each gut was diluted 10 and 100 times in sterile PBS and the CFU for each gut is the mean between the two dilutions. The results show the mean CFU for different guts ($n=4$, 3 and 3 for 0%, 0.5% and 1%, respectively). (E) Colony forming units (CFU) in individual gut of WT and $\Delta aslrim1$ mosquitoes grown with and without 1%PS ($P \leq 0.05$). Each gut was diluted 10, 100 and 1000 times in sterile PBS and the CFU for each gut is the mean between the three dilutions. The results show the mean $\pm$ SD CFU for different guts ($n=4$).

1

2

3

4 **Materials and Methods**

5 *Mosquito and parasite growth and infection*

6 The mosquitoes used in this study were *Anopheles stephensi* SDA500, reared at the University
 7 of Maryland Insect Transformation Facility (ITF) at the Institute for Bioscience and
 8 Biotechnology Research (IBBR), using standard conditions (28°C and 75% humidity). For aseptic
 9 rearing of mosquitoes, egg laying was induced and eggs were transferred to Sanaria for aseptic
 10 production[1]. For infection of mosquitoes, Pf NF54 strain were grown in blood culture for 18-
 11 20 days for the development of gametocytes. Mosquitoes were then fed through a membrane
 12 with infected blood (5 million gametocytes/mL), as established previously [1, 52, 53].

13 *Mosquito rearing, CRISPR mix and Injection into A. stephensi embryos*

14 CRISPR mix was done based on the protocol of Kistler et al. [25]. Briefly, short-guide (sg) RNAs
 15 were generated by PCR amplification of the guides following by in vitro transcription using the
 16 Ambion Megascript T7 kit (#AM1334) according to the manufacturer instructions. The RNAs
 17 were purified using Megaclear Transcription Clean-Up kit (Thermo Fisher #AM1908). Multiple
 18 sgRNAs (40 ng/μL) were mixed with 300ng of recombinant CAS9 (PNA bio - CP01). CRISPR
 19 injection mix was injected by standard methods [54, 55] into preblastoderm *A. stephensi*
 20 embryos. Embryos were injected between 40 min (TS) and 60 min (TF) post start of embryo

collection, with T0 being the time mosquitoes were added to egg chamber, Ts being average start time of injection, and TF being average time of completion for the last embryo of an injection round. All injections were done at the University of Maryland ITF at IBBR.

DNA and RNA assessments.

Adults or larvae of *A. stephensi* mosquitoes were homogenized using blue pestles and genomic (g)DNA was extracted from the homogenates using Qiagen DNeasy Blood & Tissue Kit (#69506). RNA was extracted from mosquito homogenates using Qiagen RNeasy mini kit (#74106) according to the manufacturer instructions. PCR reactions were done using MyTaq red DNA polymerase (bioline USA #BIO-21108). Complimentary (c)DNAs were synthesized using High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher # 4368814) on extracted RNAs. Real-time PCR reactions were done on cDNAs or gDNAs (for assessment of bacterial loads) using SensiFAST™ Real-Time PCR Kits (bioline # BIO-82005).

Production of LRIM1 antibody and western blot analysis.

Poly-clonal α LRIM1 was produced by injecting a short peptide (KLH conjugated), corresponding to amino acids 437-456 of *A. stephensi* LRIM1 protein, into rabbits (done by Envigo Bioproducts). For western blots, mosquito hemolymph was extracted as described previously [11] and male reproductive organs (testes, male accessory glands, *vasa deferentia* and ejaculatory ducts) were each pooled from 30 mosquitoes and lysed in RIPA lysis and extraction buffer (Thermo Scientific #89900) according to the manufacturer instructions. Samples were run on

SDS-PAGE gel and probed with the polyclonal α LRIM1 antiserum at 1:250 dilution, then a 1:5000 secondary anti rabbit (AP conjugated).

Survival, fecundity bloodmeal volume and protein content.

To estimate mosquitoes' survival, a known number of mosquitoes were separated to a clean gallon container. Dead mosquitoes were removed from the containers at the indicated time points and counted. The number of dead mosquitoes was subtracted from the total number of mosquitoes. To determine egg laying, single blood-fed mosquitoes were removed from the cages and placed in a single *Drosophila* tube containing water and sealed with a cotton ball. Mosquitoes were allowed to lay eggs for 1-2 days in a humidified 28°C room before being removed from the tubes. The eggs were counted under dissection microscope. The *Drosophila* tubes were incubated for an additional 1-2 days in the same conditions and the larvae were counted under dissection microscope. The percentage of larvae hatching was determined per the total number of eggs in the tube. For determination of bloodmeal, midguts were dissected immediately after blood feeding (2 cycles of 20 minutes per cage) and homogenized in 100 microliter PBS. A standard curve was made and hemoglobin was determined in spectrophotometer at a wavelength of 412 nM, as was done previously [56]. Similarly, bloodmeal protein contents were determined using the Lowry protein assay kit (Thermo # 23240) on midguts dissected immediately after blood feeding.

Midgut bacterial loads

To assess the effect of adding antibiotics to the sucrose meal on the midgut bacterial loads, adult mosquitoes were fed with 15% sucrose with or without different concentrations of Penicillin-streptomycin from the time of emergence. Midguts were dissected and homogenized in 100 microliters of sterile PBS. Serial dilutions of guts homogenates were then plated on LB-agar plates and colonies were counted to determine colony forming units (CFU).

Microbial diversity analyses

Genomic DNA was PCR amplified with primers 515F and 926R [57], targeting the V4 and V5 variable regions of the microbial small subunit ribosomal RNA gene using a two-stage “targeted amplicon sequencing (TAS)” protocol as described previously [20]. Primers were modified to include linker sequences at the 5’ ends (i.e., so-called “common sequences” or CS1 and CS2 on forward and reverse primers, respectively). First stage PCR amplifications were performed in 10 microliter reactions in 96-well plates, using the MyTaq HS 2X master mix (BioLine, Taunton, MA, USA). PCR conditions were 95°C for 5 minutes, followed by 28 cycles of 95°C for 30”, 50°C for 60” and 72°C for 90”. Subsequently, a second PCR amplification was performed in 10 microliter reactions in 96-well plates. A master mix for the entire plate was made using the MyTaq HS 2X master mix. Each well received a separate primer pair with a unique 10-base barcode, obtained from the Access Array Barcode Library for Illumina (Fluidigm, South San Francisco, CA; Item# 100-4876). These AccessArray primers contained the CS1 and CS2 linkers at the 3’ ends of the oligonucleotides. Cycling conditions were as follows: 95°C for 5 minutes, followed by 8 cycles of 95°C for 30”, 60°C for 30” and 72°C for 30”. A final, 7-minute elongation step was performed at 72°C. Samples were pooled in equal volume using an EpMotion5075 liquid handling robot (Eppendorf, Hamburg, Germany). The pooled library

was purified using an AMPure XP cleanup protocol (0.6X, vol/vol; Agencourt, Beckmann-Coulter) to remove fragments smaller than 300 bp. The pooled libraries, with a 20% phiX spike-in, were loaded onto an Illumina MiniSeq mid-output flow cell (2x150 paired-end reads). Based on the distribution of reads per barcode, the amplicons (before purification) were re-pooled to generate a more balanced distribution of reads. The re-pooled library was purified using AMPure XP cleanup, as described above. The re-pooled libraries, with a 15% phiX spike-in, were loaded onto a MiSeq v3 flow cell, and sequenced using an Illumina MiSeq sequencer. Fluidigm sequencing primers, targeting the CS1 and CS2 linker regions, were used to initiate sequencing. De-multiplexing of reads was performed on instrument. Library preparation, pooling, sequencing was performed at the Genome Research Core, Research Resources Center (RRC), University of Illinois at Chicago (UIC).

References

1. Hoffman SL, Billingsley PF, James E, Richman A, Loyevsky M, Li T, et al. Development of a metabolically active, non-replicating sporozoite vaccine to prevent *Plasmodium falciparum* malaria. *Hum Vaccin*. 2010;6(1):97-106. doi: 10.4161/hv.6.1.10396. PubMed PMID: 19946222.
2. Luke TC, Hoffman SL. Rationale and plans for developing a non-replicating, metabolically active, radiation-attenuated *Plasmodium falciparum* sporozoite vaccine. *J Exp Biol*. 2003;206(Pt 21):3803-8. doi: 10.1242/jeb.00644. PubMed PMID: 14506215.
3. Mordmuller B, Surat G, Lagler H, Chakravarty S, Ishizuka AS, Lalremruata A, et al. Sterile protection against human malaria by chemoattenuated PfSPZ vaccine. *Nature*. 2017;542(7642):445-9. doi: 10.1038/nature21060. PubMed PMID: 28199305.
4. Sissoko MS, Healy SA, Katile A, Omaswa F, Zaidi I, Gabriel EE, et al. Safety and efficacy of PfSPZ Vaccine against *Plasmodium falciparum* via direct venous inoculation in healthy malaria-exposed adults in Mali: a randomised, double-blind phase 1 trial. *Lancet Infect Dis*. 2017;17(5):498-509. doi: 10.1016/S1473-3099(17)30104-4. PubMed PMID: 28216244; PubMed Central PMCID: PMC6803168.
5. Seder RA, Chang LJ, Enama ME, Zephir KL, Sarwar UN, Gordon IJ, et al. Protection against malaria by intravenous immunization with a nonreplicating sporozoite vaccine. *Science*. 2013;341(6152):1359-65. doi: 10.1126/science.1241800. PubMed PMID: 23929949.
6. Simonetti AB. The biology of malarial parasite in the mosquito--a review. *Mem Inst Oswaldo Cruz*. 1996;91(5):519-41. doi: 10.1590/s0074-02761996000500001. PubMed PMID: 9137738.
7. Abraham EG, Jacobs-Lorena M. Mosquito midgut barriers to malaria parasite development. *Insect Biochem Mol Biol*. 2004;34(7):667-71. doi: 10.1016/j.ibmb.2004.03.019. PubMed PMID: 15242707.
8. Blandin S, Shiao SH, Moita LF, Janse CJ, Waters AP, Kafatos FC, et al. Complement-like protein TEP1 is a determinant of vectorial capacity in the malaria vector *Anopheles gambiae*. *Cell*. 2004;116(5):661-70. doi: 10.1016/s0092-8674(04)00173-4. PubMed PMID: 15006349.
9. Fraiture M, Baxter RH, Steinert S, Chelliah Y, Frolet C, Quispe-Tintaya W, et al. Two mosquito LRR proteins function as complement control factors in the TEP1-mediated killing of *Plasmodium*. *Cell Host Microbe*. 2009;5(3):273-84. doi: 10.1016/j.chom.2009.01.005. PubMed PMID: 19286136.
10. Moita LF, Wang-Sattler R, Michel K, Zimmermann T, Blandin S, Levashina EA, et al. In vivo identification of novel regulators and conserved pathways of phagocytosis in *A. gambiae*. *Immunity*. 2005;23(1):65-73. doi: 10.1016/j.immuni.2005.05.006. PubMed PMID: 16039580.
11. Povelones M, Waterhouse RM, Kafatos FC, Christophides GK. Leucine-rich repeat protein complex activates mosquito complement in defense against *Plasmodium* parasites. *Science*. 2009;324(5924):258-61. doi: 10.1126/science.1171400. PubMed PMID: 19264986; PubMed Central PMCID: PMC6803168.
12. Abraham EG, Pinto SB, Ghosh A, Vanlandingham DL, Budd A, Higgs S, et al. An immune-responsive serpin, SRPN6, mediates mosquito defense against malaria parasites. *Proc Natl Acad Sci U S A*

- A. 2005;102(45):16327-32. doi: 10.1073/pnas.0508335102. PubMed PMID: 16260729; PubMed Central PMCID: PMCPMC1283470.
13. Christophides GK, Vlachou D, Kafatos FC. Comparative and functional genomics of the innate immune system in the malaria vector *Anopheles gambiae*. *Immunol Rev.* 2004;198:127-48. PubMed PMID: 15199960.
14. Meister S, Kanzok SM, Zheng XL, Luna C, Li TR, Hoa NT, et al. Immune signaling pathways regulating bacterial and malaria parasite infection of the mosquito *Anopheles gambiae*. *Proc Natl Acad Sci U S A.* 2005;102(32):11420-5. doi: 10.1073/pnas.0504950102. PubMed PMID: 16076953; PubMed Central PMCID: PMCPMC1183586.
15. Michel K, Budd A, Pinto S, Gibson TJ, Kafatos FC. *Anopheles gambiae* SRPN2 facilitates midgut invasion by the malaria parasite *Plasmodium berghei*. *EMBO Rep.* 2005;6(9):891-7. doi: 10.1038/sj.embor.7400478. PubMed PMID: 16113656; PubMed Central PMCID: PMCPMC1369158.
16. Collins FH, Sakai RK, Vernick KD, Paskewitz S, Seeley DC, Miller LH, et al. Genetic selection of a *Plasmodium*-refractory strain of the malaria vector *Anopheles gambiae*. *Science.* 1986;234(4776):607-10. doi: 10.1126/science.3532325. PubMed PMID: 3532325.
17. Povelones M, Upton LM, Sala KA, Christophides GK. Structure-function analysis of the *Anopheles gambiae* LRIM1/APL1C complex and its interaction with complement C3-like protein TEP1. *PLoS Pathog.* 2011;7(4):e1002023. doi: 10.1371/journal.ppat.1002023. PubMed PMID: 21533217; PubMed Central PMCID: PMCPMC3077365.
18. Peter F. Billingsley KIG, Abraham Eappen, Robert Harrell, Robert Alford, Tao Li, Sumana Chakravarty, B. Kim Lee Sim, Stephen L. Hoffman, David A. O'Brochta. Transient knockdown of *Anopheles stephensi* LRIM1 using RNAi increases *Plasmodium falciparum* sporozoite salivary gland infections. *Malaria Journal.* 2021;In Press. doi: 10.1186/s12936-021-03818-8.
19. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci U S A.* 2011;108 Suppl 1:4516-22. doi: 10.1073/pnas.1000080107. PubMed PMID: 20534432; PubMed Central PMCID: PMCPMC3063599.
20. Naqib A, Poggi S, Wang W, Hyde M, Kunstman K, Green SJ. Making and Sequencing Heavily Multiplexed, High-Throughput 16S Ribosomal RNA Gene Amplicon Libraries Using a Flexible, Two-Stage PCR Protocol. *Methods Mol Biol.* 2018;1783:149-69. doi: 10.1007/978-1-4939-7834-2_7. PubMed PMID: 29767361.
21. Favia G, Ricci I, Marzorati M, Negri I, Alma A, Sacchi L, et al. Bacteria of the genus *Asaia*: a potential paratransgenic weapon against malaria. *Adv Exp Med Biol.* 2008;627:49-59. doi: 10.1007/978-0-387-78225-6_4. PubMed PMID: 18510013.
22. Raspor P, Goranovic D. Biotechnological applications of acetic acid bacteria. *Crit Rev Biotechnol.* 2008;28(2):101-24. doi: 10.1080/07388550802046749. PubMed PMID: 18568850.
23. Crotti E, Rizzi A, Chouaia B, Ricci I, Favia G, Alma A, et al. Acetic acid bacteria, newly emerging symbionts of insects. *Appl Environ Microbiol.* 2010;76(21):6963-70. doi: 10.1128/AEM.01336-10. PubMed PMID: 20851977; PubMed Central PMCID: PMCPMC2976266.
24. Moreno M, Tong-Rios C, Orjuela-Sanchez P, Carrasco-Escobar G, Campo B, Gamboa D, et al. Continuous Supply of *Plasmodium vivax* Sporozoites from Colonized *Anopheles darlingi* in the Peruvian Amazon. *ACS Infect Dis.* 2018;4(4):541-8. doi: 10.1021/acsinfecdis.7b00195. PubMed PMID: 29465219; PubMed Central PMCID: PMCPMC5902790.
25. Toure AM, Mackey AJ, Wang ZX, Beier JC. Bactericidal effects of sugar-fed antibiotics on resident midgut bacteria of newly emerged anopheline mosquitoes (Diptera: Culicidae). *J Med Entomol.* 2000;37(2):246-9. doi: 10.1093/jmedent/37.2.246. PubMed PMID: 10730495.

26. Billingsley PF, Hecker H. Blood digestion in the mosquito, *Anopheles stephensi* Liston (Diptera: Culicidae): activity and distribution of trypsin, aminopeptidase, and alpha-glucosidase in the midgut. *J Med Entomol.* 1991;28(6):865-71. doi: 10.1093/jmedent/28.6.865. PubMed PMID: 1770523.
27. Briegel H, Rezzonico L. Concentration of host blood protein during feeding by anopheline mosquitoes (Diptera: Culicidae). *J Med Entomol.* 1985;22(6):612-8. doi: 10.1093/jmedent/22.6.612. PubMed PMID: 4078846.
28. Osta MA, Christophides GK, Kafatos FC. Effects of mosquito genes on *Plasmodium* development. *Science.* 2004;303(5666):2030-2. doi: 10.1126/science.1091789. PubMed PMID: 15044804.
29. Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, Marzorati M, et al. Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc Natl Acad Sci U S A.* 2007;104(21):9047-51. doi: 10.1073/pnas.0610451104. PubMed PMID: 17502606; PubMed Central PMCID: PMCPMC1885625.
30. Chouaia B, Rossi P, Montagna M, Ricci I, Crotti E, Damiani C, et al. Molecular evidence for multiple infections as revealed by typing of *Asaia* bacterial symbionts of four mosquito species. *Appl Environ Microbiol.* 2010;76(22):7444-50. doi: 10.1128/AEM.01747-10. PubMed PMID: 20851960; PubMed Central PMCID: PMCPMC2976182.
31. Dong Y, Taylor HE, Dimopoulos G. AgDscam, a hypervariable immunoglobulin domain-containing receptor of the *Anopheles gambiae* innate immune system. *PLoS Biol.* 2006;4(7):e229. doi: 10.1371/journal.pbio.0040229. PubMed PMID: 16774454; PubMed Central PMCID: PMCPMC1479700.
32. Romoli O, Gendrin M. The tripartite interactions between the mosquito, its microbiota and *Plasmodium*. *Parasit Vectors.* 2018;11(1):200. doi: 10.1186/s13071-018-2784-x. PubMed PMID: 29558973; PubMed Central PMCID: PMCPMC5861617.
33. Bahia AC, Dong Y, Blumberg BJ, Mlambo G, Tripathi A, BenMarzouk-Hidalgo OJ, et al. Exploring *Anopheles* gut bacteria for *Plasmodium* blocking activity. *Environ Microbiol.* 2014;16(9):2980-94. doi: 10.1111/1462-2920.12381. PubMed PMID: 24428613; PubMed Central PMCID: PMCPMC4099322.
34. Gendrin M, Rodgers FH, Yerbanga RS, Ouedraogo JB, Basanez MG, Cohuet A, et al. Antibiotics in ingested human blood affect the mosquito microbiota and capacity to transmit malaria. *Nat Commun.* 2015;6:5921. doi: 10.1038/ncomms6921. PubMed PMID: 25562286; PubMed Central PMCID: PMCPMC4338536.
35. Meister S, Agianian B, Turlure F, Relogio A, Morlais I, Kafatos FC, et al. *Anopheles gambiae* PGRPLC-mediated defense against bacteria modulates infections with malaria parasites. *PLoS Pathog.* 2009;5(8):e1000542. doi: 10.1371/journal.ppat.1000542. PubMed PMID: 19662170; PubMed Central PMCID: PMCPMC2715215.
36. Dong Y, Manfredini F, Dimopoulos G. Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog.* 2009;5(5):e1000423. doi: 10.1371/journal.ppat.1000423. PubMed PMID: 19424427; PubMed Central PMCID: PMCPMC2673032.
37. George KI. GENETIC MANIPULATION OF ANOPHELES STEPHENSI IMMUNITY TO INCREASE PLASMODIUM FALCIPARUM SALIVARY GLAND SPOOROZOITE INFECTION LEVELS. Dissertation submitted to the Faculty of the Graduate School of the University of Maryland, College Park, in partial fulfillment of the requirements for the degree of Doctor of Philosophy. 2016.
38. Zhang G, Niu G, Franca CM, Dong Y, Wang X, Butler NS, et al. *Anopheles* Midgut FREP1 Mediates *Plasmodium* Invasion. *J Biol Chem.* 2015;290(27):16490-501. doi: 10.1074/jbc.M114.623165. PubMed PMID: 25991725; PubMed Central PMCID: PMCPMC4505404.

39. Dong Y, Simoes ML, Marois E, Dimopoulos G. CRISPR/Cas9 -mediated gene knockout of *Anopheles gambiae* FREP1 suppresses malaria parasite infection. *PLoS Pathog.* 2018;14(3):e1006898. doi: 10.1371/journal.ppat.1006898. PubMed PMID: 29518156; PubMed Central PMCID: PMC5843335.
40. Raikhel AS, Dhadialla TS. Accumulation of yolk proteins in insect oocytes. *Annu Rev Entomol.* 1992;37:217-51. doi: 10.1146/annurev.en.37.010192.001245. PubMed PMID: 1311540.
41. Tufail M, Takeda M. Molecular characteristics of insect vitellogenins. *J Insect Physiol.* 2008;54(12):1447-58. doi: 10.1016/j.jinsphys.2008.08.007. PubMed PMID: 18789336.
42. Rono MK, Whitten MM, Oulad-Abdelghani M, Levashina EA, Marois E. The major yolk protein vitellogenin interferes with the anti-plasmodium response in the malaria mosquito *Anopheles gambiae*. *PLoS Biol.* 2010;8(7):e1000434. doi: 10.1371/journal.pbio.1000434. PubMed PMID: 20652016; PubMed Central PMCID: PMC2907290.
43. Yang J, Schleicher TR, Dong Y, Park HB, Lan J, Cresswell P, et al. Disruption of *mosGILT* in *Anopheles gambiae* impairs ovarian development and Plasmodium infection. *J Exp Med.* 2019. doi: 10.1084/jem.20190682. PubMed PMID: 31658986.
44. Chambers GM, Klowden MJ. Age of *Anopheles gambiae* Giles male mosquitoes at time of mating influences female oviposition. *J Vector Ecol.* 2001;26(2):196-201. PubMed PMID: 11813657.
45. Voordouw MJ, Koella JC. Genetic variation of male reproductive success in a laboratory population of *Anopheles gambiae*. *Malar J.* 2007;6:99. doi: 10.1186/1475-2875-6-99. PubMed PMID: 17663767; PubMed Central PMCID: PMC1971063.
46. Pompon J, Levashina EA. A New Role of the Mosquito Complement-like Cascade in Male Fertility in *Anopheles gambiae*. *PLoS Biol.* 2015;13(9):e1002255. doi: 10.1371/journal.pbio.1002255. PubMed PMID: 26394016; PubMed Central PMCID: PMC4579081.
47. Damiani C, Ricci I, Crotti E, Rossi P, Rizzi A, Scuppa P, et al. Paternal transmission of symbiotic bacteria in malaria vectors. *Curr Biol.* 2008;18(23):R1087-8. doi: 10.1016/j.cub.2008.10.040. PubMed PMID: 19081038.
48. Cho GY, Schaefer KA, Bassuk AG, Tsang SH, Mahajan VB. Crispr Genome Surgery in the Retina in Light of Off-Targeting. *Retina.* 2018;38(8):1443-55. doi: 10.1097/IAE.0000000000002197. PubMed PMID: 29746416; PubMed Central PMCID: PMC6054556.
49. Greely HT. CRISPR'd babies: human germline genome editing in the 'He Jiankui affair'. *J Law Biosci.* 2019;6(1):111-83. doi: 10.1093/jlb/lsz010. PubMed PMID: 31666967; PubMed Central PMCID: PMC6813942.
50. Skryabin BV, Kummerfeld DM, Gubar L, Seeger B, Kaiser H, Stegemann A, et al. Pervasive head-to-tail insertions of DNA templates mask desired CRISPR-Cas9-mediated genome editing events. *Sci Adv.* 2020;6(7):eaax2941. doi: 10.1126/sciadv.aax2941. PubMed PMID: 32095517; PubMed Central PMCID: PMC7015686.
51. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol.* 2016;18(5):1403-14. doi: 10.1111/1462-2920.13023. PubMed PMID: 26271760.
52. Ponnudurai T, Lensen AH, Van Gemert GJ, Bensink MP, Bolmer M, Meuwissen JH. Infectivity of cultured *Plasmodium falciparum* gametocytes to mosquitoes. *Parasitology.* 1989;98 Pt 2:165-73. doi: 10.1017/s0031182000062065. PubMed PMID: 2668861.
53. Li T, Eappen AG, Richman AM, Billingsley PF, Abebe Y, Li M, et al. Robust, reproducible, industrialized, standard membrane feeding assay for assessing the transmission blocking activity of vaccines and drugs against *Plasmodium falciparum*. *Malar J.* 2015;14:150. doi: 10.1186/s12936-015-0665-8. PubMed PMID: 25890243; PubMed Central PMCID: PMC4491417.
54. Abraham EG, Donnelly-Doman M, Fujioka H, Ghosh A, Moreira L, Jacobs-Lorena M. Driving midgut-specific expression and secretion of a foreign protein in transgenic mosquitoes with AgAper1

regulatory elements. *Insect Mol Biol.* 2005;14(3):271-9. doi: 10.1111/j.1365-2583.2004.00557.x.
PubMed PMID: 15926896.
55. Catteruccia F, Nolan T, Loukeris TG, Blass C, Savakis C, Kafatos FC, et al. Stable germline transformation of the malaria mosquito *Anopheles stephensi*. *Nature.* 2000;405(6789):959-62. doi: 10.1038/35016096. PubMed PMID: 10879538.
56. Feldmann AM, Billingsley PF, Savelkoul E. Bloodmeal digestion by strains of *Anopheles stephensi* liston (Diptera: Culicidae) of differing susceptibility to *Plasmodium falciparum*. *Parasitology.* 1990;101 Pt 2:193-200. doi: 10.1017/s003118200006323x. PubMed PMID: 2263414.
57. Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, et al. Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys. *mSystems.* 2016;1(1). doi: 10.1128/mSystems.00009-15. PubMed PMID: 27822518; PubMed Central PMCID: PMC5069754.

Acknowledgments

We thank the following Sanaria staff for their support: Stephen Matheny for mosquito rearing, Asha Patil, Yonas Abebe, Christiane Tran, Lauren Smith and Alicia Robinson for maintenance of *P. falciparum* parasites and infection of mosquitoes, and Urvashi Rai, Alyssa Arnheim, Michelle Ryan and Tanya Samuel for dissection of mosquito salivary glands and assessment of sporozoite infections. This work was supported by NIAID SBIR grant 5R44AI131806-03.

Author Contributions

E.I., A.G.E., R.T.A., W.R. R.A.H., M.H., S.C. and T.L. Performed the work and contributed to data collection; E.I., A.G.E., P.F.B and S.L.H. designed the study and wrote the manuscript; B.K.L.S. P.F.B., A.G.E. and S.L.H. supervised the project. All authors discussed the results and commented on the manuscript.

Competing interests

E.I., A.G.E., M.H., T.L., S.C., P.F.B., B.K.L.S., and S.L.H. are salaried employees of Sanaria Inc., the developer and owner of PfSPZ Vaccine. In addition, S.L.H. and B.K.L.S. have a financial interest in Sanaria Inc. No other authors have a competing interest.

Fig 1

Fig 2

A

B

C

D

E

Fig 3

Fig 4

Fig S1

A

B

C

Fig S2

Fig S3

Fig S4