Music-Based Intervention Connects Auditory and Reward Systems

Milena Aiello Quinci¹, Alexander Belden¹, Valerie Goutama¹, Dayang Gong¹, Suzanne Hanser², Nancy J. Donovan³, Maiya Geddes³, Psyche Loui^{1*}

¹ Northeastern University

² Berklee College of Music

³ Brigham and Women's Hospital and Harvard Medical School

* Corresponding author: p.loui@northeastern.edu

Abstract

Listening to pleasurable music is known to engage the brain's reward system, but little is known about how this engagement develops over time. Here we show for the first time that brain network connectivity can change longitudinally as a result of a personalized receptive music-based intervention (MBI) in cognitively unimpaired older adults. Using a combination of whole-brain regression, seed-based connectivity analysis, and representational similarity analysis (RSA), we compared fMRI responses during a simple music listening task in older adults before and after an eight-week personalized music listening program. Participants rated self-selected and researcher-selected musical excerpts on liking and familiarity. Parametric effects of liking, familiarity, and selection showed significant activation of auditory, reward, default mode, and sensorimotor areas both pre- and post-intervention. Seed-based connectivity comparing pre- and post-intervention showed a significant increase in functional connectivity between auditory regions and the medial prefrontal cortex (mPFC), and this auditory-mPFC connectivity was modulated by participant liking and familiarity ratings. RSA showed significant representations of selection and novelty in auditory regions at both time-points, and an increase in striatal representation of musical stimuli following intervention. Taken together, results show a sensitivity of auditory, reward, default, and sensorimotor regions to individual differences in music familiarity and liking, as well as a shift in brain network dynamics following the personalized MBI. Results show how regular music listening can provide an auditory channel towards the mPFC, thus offering a potential neural mechanism supporting healthy brain aging.

Keywords

Music, aging, dementia, longitudinal, intervention, fMRI

Listening to music engages a network of auditory areas, including the Heschl's gyrus (HG), superior temporal gyrus (STG), and superior temporal sulcus (STS) (1,2). When listening to pleasurable music, areas in this auditory network are functionally connected with areas in the dopaminergic reward system, specifically medial prefrontal cortex (mPFC) as well as dorsal and ventral striatum (3,4,5,6). The reward system is involved in motivated behavior (7) and is active during the processing of biologically relevant stimuli such as food, as well as cues that are tightly linked to biological stimuli, such as money (8). Reward system areas are also involved in the learning of associations between stimuli and their subjective value (4,9,10,11,12). Despite their known coactivity, little is understood about how connectivity between auditory and reward areas change over time, either naturally or as a result of training-dependent plasticity. Understanding time-dependent changes in auditory-reward connectivity will inform studies on the evolutionary bases and social roles of music (13) as well as applied studies on music, health, and well-being, with especially widespread implications for music therapy.

In addition to activating the reward system, the experience of music is fundamentally intertwined with the listener's knowledge, autobiographical memories, and understanding of the intentions of the composer and the performer, through its structure and performance (14,15,16). These rich experiences are supported by activity in the default mode network (DMN), which encompasses the mPFC (which is also part of the reward network), along with the posterior cingulate cortex and temporoparietal junction or angular gyrus. The DMN is involved during self-generated thought (17) and is disrupted in activity and connectivity in multiple neuropsychiatric disorders and conditions such as depression and learned helplessness (18).

Agency in the selection of music, or the ability to choose the music that one listens to, may provide the listener with a psychological platform to link the consequences of one's musical choices to perceptual predictive processes, giving rise to a relatively active mode of listening (19) that has implications for the health-relevance of music and the use of music in therapy (20). Listening to self-selected music reduces anxiety and improves task performance and enjoyment, compared to music selected by an experimenter (21). While previous fMRI studies suggest that the reward network and DMN are likely involved (22), there is little direct evidence for the involvement of these networks in self-selection of music listening per se. Since the mPFC is both within the DMN and in the reward network, testing how self-selected music activates these disparate networks may inform neuroscientific understanding about how the reward and default networks interact in the brain.

Here we present a task fMRI study of music listening before and after a music-based intervention (MBI) in cognitively unimpaired, community-dwelling older adults. MBIs are therapeutic strategies facilitated by music, and can be classified as either receptive if they are primarily focused on music listening, or interactive if they involve active engagement in music making activities (23). Receptive MBIs have been shown to reduce depressive symptoms in individuals with clinical depression (24,25) and to improve quality of life and emotional well-being, while decreasing anxiety for people with dementia (26). Relatedly, an eight-week intervention on mindfulness-based stress reduction has shown changes in reward-related fMRI activity and connectivity (27). Thus, we developed an eight-week receptive MBI program, based on previous research with older adults (28), and collected neuropsychological and neuroimaging

measures both prior to and after participation in the intervention. Through this project, we aim to investigate the effects of music liking, familiarity, and self-selection on reward processing before and after intervention, through a combination of whole-brain regression, seed-based analysis, and representational similarity analysis (RSA).

<insert Figure 1 here>

Our predictions (preregistered [here: https://osf.io/qsevr/](https://osf.io/qsevr/)) are threefold: first, we predict that brain responses to musical stimuli will be sensitive to differences in liking, familiarity, and self-selection. Specifically, from a whole-brain analysis, we expect higher brain activity when listening to musical stimuli that are well-liked, familiar, and self-selected. Second, we predict that the activity and connectivity within and between auditory and reward areas, as observed in region-of-interest (ROI) analyses, will be sensitive to differences in liking, familiarity, and self-selection. Third, we expect some experience-dependent changes in how the brain responds to music over time, to be observed by comparing activity and connectivity in auditory and reward regions before and after MBI.

Results

Whole-Brain Univariate Effects of Liking, Familiarity, and Self-Selection

Our first hypothesis, i.e. that brain responses to musical stimuli would be sensitive to differences in liking, familiarity, and self-selection, was tested using parametric contrasts on liking and familiarity ratings, and a contrast between self-selected and other-selected music. Whole-brain effects for each liking rating, each familiarity rating, and each stimulus type can be found in <https://neurovault.org/collections/TCPSFBMF/>, and in Supplementary Materials Figure S1-S3.

Parametric Effect of Liking

Table 1 and Figure 2A show the whole-brain corrected univariate, second-level parametric effects of liking ratings pre-intervention and post-intervention at the $p < .05$ voxelwise and clusterwise FDR-corrected level. Pre-intervention, the parametric effect of liking was observed in auditory regions including bilateral STG, STS, and MTG, DMN regions including mPFC, PCC, and temporoparietal junction (TPJ)/inferior parietal lobule (IPL), reward regions including mPFC, dorsal striatum (caudate), ventral striatum (nucleus accumbens NAcc, putamen), and amygdala, and motor regions including premotor cortex, SMA, precentral gyrus (PCG), superior frontal gyrus, and frontal operculum. Post-intervention, the parametric effect of liking was observed in auditory areas including bilateral STG, STS, and MTG, in DMN areas including mPFC, PCC, and TPJ/IPL, and in ventral striatum. Pre-intervention and post-intervention effects of liking did not differ significantly at the whole-brain FDR-corrected levels.

Parametric Effect of Familiarity

Table 2 and Figure 2B show the whole-brain corrected univariate, second-level parametric effects of familiarity ratings pre- and post-intervention, again at the $p < .05$ voxelwise and

clusterwise FDR-corrected level. Pre-intervention, the parametric effect of familiarity was observed in a large cluster spanning the auditory areas including bilateral STG, STS, and MTG, reward areas including dorsal and ventral striatum, DMN areas including the TPJ/IPL and PCC, and sensorimotor areas including the precentral and postcentral gyri and SMA. Post-intervention, the parametric effect of familiarity was observed bilaterally in auditory regions including STG, STS, MTG, and the PCC. Pre-intervention and post-intervention effects of familiarity did not differ significantly at the FDR-corrected levels.

Effect of Self vs. Other-Selection

Table 3 and Figure 2C show the whole-brain corrected univariate, second-level contrast of self-selected vs. other-selected music listening pre- and post-intervention, again at the $p < .05$ voxelwise and clusterwise FDR-corrected level. Pre-intervention, self-selected music, compared to other-selected music, elicited greater activation in the bilateral STG/HG, orbitofrontal cortex, dorsal and ventral striatum, mPFC, PCC, TPJ/SPL, and PCG. Post-intervention, self-selected compared to other-selected music elicited greater activation in the STG, MTG, mPFC, ventral striatum, PCC, TPJ/SPL, and parahippocampal gyrus. Pre-intervention and post-intervention effects of selection did not differ significantly at the FDR-corrected levels.

<insert Figure 2 here>

Seed-Based Connectivity: Post-intervention > Pre-intervention

Whole-brain analyses of cortical activity during music listening showed that while classic auditory areas in the superior temporal lobe were active both pre- and post-intervention, music that received the highest liking and familiarity ratings elicited significant activity in the mPFC post-intervention, but not pre-intervention (Figure 3A; all conditions are shown in Supplementary Materials Figure S1-S3). This additional involvement of mPFC with auditory areas following the intervention suggests that intervention increased connectivity between these areas. To test our hypothesis of experience-dependent changes in activity and connectivity of auditory and reward regions, we conducted seed-based connectivity analyses, using auditory regions as the seed ROI and comparing its functional connectivity patterns before and after MBI. Figure 3B shows seed-based functional connectivity from the seed ROIs of the auditory network, including HG, STG, STS, and MTG (29). A between-sessions contrast (Post-intervention > Pre-intervention) on functional connectivity maps seeded from the auditory network showed a significant positive cluster in the mPFC at the $p < .05$ FDR-corrected level, indicating higher functional connectivity post-intervention than pre-intervention in this region. Thus, auditory areas increased in their connectivity to the mPFC after the intervention. Since this specific location within mPFC was the only region that increased its functional connectivity with the auditory network, we extracted the beta series from this significant mPFC cluster for further ROI-based analyses.

ROI analysis

To test our hypothesis that auditory and reward regions would be sensitive in their activity and connectivity to the self-selectedness of music, we extracted beta values from the auditory

network (29) and the mPFC from the reward network. Figure 3E shows between-subject correlations between auditory and mPFC areas, pre- and post-intervention for the four levels of liking ratings. While auditory and mPFC activity were positively correlated across all conditions, the correlation was highest ($r = 0.75$) for musical stimuli that received a “Love” rating. Furthermore, the between-subjects correlation between auditory and mPFC regions were higher post-intervention than pre-intervention, suggesting an increase in auditory-reward connectivity over time.

To address the functional connectivity of auditory and reward regions over the time-course of listening to music, we extracted the time series of beta values ($TR = .475$ s) from the significant mPFC cluster identified in the auditory seed-based connectivity analysis above, and compared them across conditions and across sessions. While there was no clear difference between the conditions in pre-intervention beta series, a clear pattern emerged between conditions in the post-intervention beta series, such that at the midpoint of the 20-second-long musical stimuli, beta values were highest for self-selected, highly familiar, and well-liked stimuli (Figure 3CD). In other words, auditory-reward connectivity increased over time especially for well-liked, familiar, and self-selected musical excerpts, but did not increase over time for well-known vs. novel pieces of music.

<insert Figure 3 here>

Representational Similarity Analysis

A representational similarity analysis (methods outlined in Figure 4A) was conducted to compare how similarly the distinct ROIs and networks of interest (listed in Section 2.7.5) represented psychological characteristics of the musical stimuli, such as their novelty (well-known vs. novel stimuli), their selection (self- vs other-selected), and their liking and familiarity ratings, before and after intervention. Permutation testing comparisons of model RDMs to first-level fMRI-derived RDMs (Figure 4B) indicated a significant ($p < 0.05$, T-max corrected) representation of novelty in the auditory network pre-intervention, but not post-intervention. In addition, there was significant representation of novelty in the left and right anterior and posterior STG, as well as representation of stimulus selection in the right posterior STG, and representation of liking in the anterior salience network at pre-intervention. Post-intervention, there were significant representations of stimulus selection in the left and right anterior and posterior STG and the basal ganglia, significant representations of novelty in bilateral posterior STG and left anterior STG, and significant representations of familiarity in both the basal ganglia and the language networks. Comparing pre-intervention and post-intervention fMRI-derived RDMs showed a significant effect of session in the representation of stimulus selection in the right executive control network, with a significant increase in relatedness post-intervention.

Multidimensional scaling visualisations of second-level RDMs (Figure 4C) indicate a tight clustering of auditory (green) ROIs at both time points. These regions assort most closely with the model of novelty at both time points, and reward associated regions (pink) show a closer assortment with these regions post-intervention. A close relationship between models of liking, familiarity, and selection was also observed, and this relationship became even closer post-

intervention. These models were most highly similar to bilateral SMA and the anterior salience network at pre-intervention, and they were most similar to anterior salience and language networks at post-intervention. Post-intervention representations of dorsal DMN and mPFC were also more closely related to bilateral nucleus accumbens as compared to pre-intervention. Overall, neural representations of musical stimuli were mostly similar before and after intervention, except that the reward network became more similar to the auditory network after intervention (Figure 4C), and the right executive control network showed increased differentiation between self-selected and other-selected stimuli, resulting in higher representational similarity with the stimulus selection model after intervention (Figure 4B, right panel).

<insert Figure 4 here>

Discussion

We present a first longitudinal fMRI study of a receptive music-based intervention on auditory-reward activity and connectivity in older adults. Musical excerpts that were self-selected, well-liked, and highly familiar were the most effective at driving the activity of the auditory, DMN, and auditory-motor systems. Functional connectivity of the auditory system and the mPFC increased over the course of the intervention and became more selective for self-selected and well-liked music. Results have implications for the design and implementation of music-based interventions for special populations who have disrupted DMN or reward systems, including Alzheimer's disease and depression, respectively.

Univariate patterns of brain activity showed many similarities pre- and post-intervention and included auditory and DMN areas. Auditory-motor activity (SMA, pre-SMA, cingulate) significantly increased with liking before but not after the intervention. The parametric contrast of familiarity showed activity in the salience network (cingulate and anterior insula) before but not after intervention. While stimuli that are familiar and well-liked both engage the DMN as well as the auditory network, self-selected music is most effective at engaging many wider swaths of regions throughout the brain, with activity extending into sensorimotor areas (PCG, postcentral gyri, SMA), executive control areas (middle frontal gyri), and areas in the salience network (cingulate cortex, insula). These effects of liking and familiarity are largely consistent with previous reports, but this is the first study that parametrically tested different levels of liking and familiarity in the same study. The effect of familiarity on mental representations of music has been shown on activity of the auditory network identified here (30,31,32), with effects on emotion (31) and mental representation of music (32). A prior systematic review found that music familiarity tended to elicit a motor pattern of activation (31). While this is consistent with our findings pre-intervention, we found no significant activity in the auditory-motor areas such as the premotor areas, precentral gyrus, and SMA, after intervention. As we used the same stimuli for pre-intervention and post-intervention scans, this reduction in motor activity may be because the unfamiliar stimuli became more familiar by the time the post-intervention session took place. However, the effects of familiarity persisted post-intervention in the auditory network and the DMN. The persistent auditory activity post-intervention may reflect that the auditory system was

still able to form stronger predictions for familiar music than for unfamiliar music following the intervention. The persistent DMN activity may reflect mind-wandering, or stimulus-independent thought (17,33), including autobiographical memory and associations that may be elicited by music that is well-known to the listener.

The parametric effects of liking are very similar to those of familiarity, but the effects of liking are more focused on auditory, DMN (mPFC, PCC, TPJ/IPL), and motor (premotor, SMA, pre-SMA, PCG) areas pre-intervention, while again only the auditory and DMN regions were active post-intervention. Comparing the effects of liking against effects of familiarity, the TPJ/IPL and mPFC are two regions that persisted both pre and post-intervention for liking but not for familiarity. This distinction between liking and familiarity is important: the mPFC is part of both the DMN and the reward system, and it is consistently active during pleasurable experiences including but not limited to music (34). The fact that it is active during the liking contrast and not the familiarity contrast, adds further support to mounting evidence that the mPFC has a unique role within the DMN, as a region that is common to the DMN and the dopaminergic system.

The TPJ/IPL is frequently active during theory of mind tasks (35). In addition to being part of the DMN (33), the TPJ and the mPFC have together been posited as a theory-of-mind network (36). This role of TPJ/IPL in thinking about other minds fits well with the idea that musical experiences are fundamentally intertwined with the listener's understanding of the intentions of the composer and the performer. It has been posited that "the best music" should have rich structure that can be inferred from its surface, and that the subjective value of music arises from an alliance between "compositional grammar" and the listener's "listening grammar" (37). While the relationship between the meaning of music and its subjective value is complex and abstract, the role of a theory-of-mind network may add to the claim that comprehensibility — here, the understanding of other minds — is necessary if not sufficient for subjective value in music. Here, we see that while the TPJ/IPL and mPFC responded selectively to well-liked music, they showed even greater activity in the self-selected vs. other-selected contrast. The fact that self-selected music was more effective at engaging the theory-of-mind network, compared to music that was retroactively rated as well-liked, again points to a role of agency in determining the subjective value of music.

A strength of the present study is that we obtained fMRI data using a fast TR with many different kinds of musical stimuli, including some that were self-selected by the participant, and others selected by the researcher to span well-known musical selections as well as novel stimuli. The quality and quantity of data enabled more time-sensitive analyses of specific ROI data, including representational similarity analysis to capture the relative distance between specific ROIs and functional networks, and between ROIs and experimentally-defined variables such as liking, familiarity, novelty, and self-selection. Representational similarity analysis confirmed that liking and familiarity for music are very similarly represented in the brain, and that both are strongly predicted by agency, or the effect of self-selection. Interestingly, the variables of liking, familiarity, and self-selection were more closely represented post-intervention than pre-intervention. This is consistent with the univariate analyses, in showing that effects of liking, familiarity, and self-selection are more similar to each other post-intervention than pre-intervention. Another important observation from RSA is that the reward network became more

similar (closer in the MDS plot) to the auditory network after intervention. Novelty was represented differently from liking, familiarity, and self-selection. Relatedly, the right executive control network (RECN) was the only network that showed a significant change in representation of stimulus features, specifically of the feature of selection. This is reflected in the increased dissimilarity in the RECN's representation of the self-selected and novel stimuli post-intervention (Supplementary Materials Figure S6). In other words, the RECN was more representative of stimulus selection after intervention, in contrast to other networks such as the DMN. This pattern of increased dissimilarity between RECN and DMN that emerged after intervention converges with the well-replicated finding of anticorrelated activity between DMN and ECN at rest and during most cognitive tasks (38).

Seed-based connectivity showed an increase in functional connectivity from auditory regions to mPFC after intervention. This increase in connectivity is modulated by liking and familiarity, as well as by self-selection. This is borne out by between-subjects correlations as well as within-subjects beta time-series comparisons: Highest auditory-mPFC connectivity was observed across subjects during liked and loved stimuli, especially after intervention. The beta time-series data showed an increasing differentiation by liking, familiarity, and self-selection after the intervention. In contrast, stimuli that were chosen to be well-known to the participants did not differ from novel stimuli in auditory-mPFC functional connectivity. Thus, the music-based intervention increased auditory-reward connectivity, but especially for some individuals and for some listening conditions. This is an important finding, as it is the first longitudinal demonstration of change in auditory-mPFC connectivity. Auditory-reward connectivity has been shown previously during the experience of pleasure when listening to music in task fMRI, PET, (5,6,39), resting state fMRI, and white matter connectivity as identified by DTI (40,41,42). Auditory-reward connectivity has also been modulated pharmacologically (3) and with transcranial magnetic brain stimulation (43). However, this is the first study to show a natural (i.e. as opposed to experimentally-induced) longitudinal change in auditory-reward connectivity after a receptive music-based intervention. By mindfully listening to music over the course of weeks, we have shown that participants can change their auditory-reward connectivity. As the reward system is important for many kinds of motivated behavior, these results have profound implications for the design of lifestyle interventions, such as music listening, that may affect motivated behavior through its underlying brain mechanisms.

The mPFC is a crucial hub across multiple networks in the brain, due to its membership in the dopaminergic reward network, in the DMN, and in coding for subjective value and a sense of ownership or self-referential processing (44,45). Rather than disentangling between these theoretical contributions to mPFC activity, the finding that music selectively engages mPFC may highlight the role that music has in all of these psychological functions. Our findings support the idea that music provides an auditory channel towards reward centers including the mPFC. This auditory channel towards the reward system is posited to be the evolutionary basis for the role of music for social bonding (13).

From a clinical standpoint, the mPFC and PCC are most sensitive to changes in functional connectivity in Alzheimer's disease (46). mPFC activity and connectivity are also disrupted in psychiatric diseases, including depression and schizophrenia (18). Thus, these results suggest

that MBI could help mitigate aberrant network changes in depression and AD, by modulating activity and connectivity of the mPFC. While these results suggest that MBI is effective in changing auditory-reward connectivity, one caveat is that the present longitudinal study is not a randomized controlled trial; thus the results could be explained by time-dependent changes that are extraneous to music listening. However, given that the older adult population sampled here is unlikely to have spontaneously increasing brain connectivity in these areas without a focused intervention, and given that auditory-reward connectivity is a relatively well-established set of regional hypotheses from previous research (6,34,47), we expect that the observed changes here point to the longitudinal effects of music listening. Nevertheless, in future studies we plan to define a control intervention that helps isolate the roles of specific musical features on changes in auditory-reward activity and connectivity.

More generally, the findings of this study suggest that participants should choose the music to be applied in a given intervention or music therapy treatment program, in order to leverage the effects of agency on engagement of brain functions. Allowing individuals to self-select music will also necessitate collaborations between neuroscientists and music therapists to implement MBIs in ways that respect the cultural dependence and personal salience of musical experiences (48). By demonstrating the ability of music to engage the connection between auditory and reward systems in healthy older adults, our findings shed light on potential therapeutic effects of music listening on healthy aging.

Materials and Methods

Participants

Twenty-two older adults met inclusion criteria to participate. Three of them dropped out during the first session for various reasons related to the MRI portion of the study, and three were lost-to-follow-up after the second session as a result of the COVID-19 pandemic, resulting in a final sample of 16 older adults (8 males and 8 females) between the ages of 54 and 89 ($M=66.38$, $SD=8.74$). Recruitment took place through craigslist.org and through coauthors (MG and ND) at Brigham and Women's Hospital. Participants were included if they 1) were at least 50 years old, 2) had normal hearing, and 3) passed MRI screening. Participants were excluded if they 1) changed medications within 6 weeks of screening; 2) had a history of psychotic or schizophrenic episodes, major neurological diagnosis, or other medical condition that might impair cognition; 3) had a history of chemotherapy within the past 10 years; or 4) experienced serious physical trauma or were diagnosed with a serious chronic health condition requiring medical treatment and monitoring within 3 months of screening. Participants were compensated for their time. This study was approved by the Northeastern University Institutional Review Board.

Procedure

Prior to their enrollment, participants completed a pre-screening by telephone. Participants were screened for the inclusion and exclusion criteria and completed the Telephone Interview for Cognitive Status (TICS; 49) with a score of $\geq 31/41$ considered cognitively normal and an MRI pre-entry screening form. They were also asked to provide the researcher with the names of six

musical pieces they enjoy listening to. Participants who passed pre-screening were invited to the lab for a pre-intervention session, which included a battery of neuropsychological tests and behavioral measures, an MRI, and a blood draw. The neuropsychological tests were administered by a trained researcher, and the behavioral measures were completed through an online survey. Later that week, participants were invited back to the lab to meet with a music therapist for an assessment of their musical preferences and creation of two personalized playlists for the MBI (see section 2.4). At the end of the 8-week intervention period (12 weeks for the 2 participants enrolled during COVID-19; see Section 2.3), participants returned to the lab for a post-MBI session to complete the same battery of neuropsychological tests and behavioral measures administered during the first session, as well as a post-intervention MRI and blood draw. See Figure 1 for a timeline of the procedure. For the present manuscript, we report data from the task fMRI portion of the study; data from the neuropsychological battery and the behavioral measures will be reported separately.

Special Considerations due to COVID-19

Data collection for this study began in July of 2019, and had been completed per protocol on 10 participants before the United States entered a COVID-19-related lockdown. Two participants included in the final sample were actively undergoing MBI when the lockdown began in March 2020. For those two participants, the music intervention was extended to 12 weeks rather than the typical 8 weeks, to complete the intervention as intended. Resumption of Research Plan was developed and approved in consultation with the Institutional Review Board (IRB) of Northeastern University. Removing those two participants did not alter the overall pattern of results, thus they were analyzed along with the rest of the sample. The remaining 4 participants in the final sample were enrolled during the COVID-19 pandemic, and underwent an 8-week MBI and performed both pre- and post-intervention testing in compliance with the COVID-specific Resumption of Research plan.

For our COVID-specific Resumption of Research plan: consenting and neuropsychological testing were done via a HIPAA-compliant version of Zoom to minimize the amount of time participants spent in the lab for in-person data collection. Participants were called one day before their scheduled session, screened using a COVID-19 pre-screener based on the COVID-19 CDC guidelines, and were provided with the lab's IRB-approved COVID-19-compliant guidelines. If participants answered no to all of COVID-19 pre-screener questions, they were invited to come in for their session the next day. After arriving for their session, participants were screened again using the same COVID-19 pre-screener and their temperatures were taken. If participants answered no to all of the questions and had a temperature reading of less than 100.4° F, then they were invited into the lab. Participants and researchers wore full personal protective equipment (PPE: surgical masks, face shields, lab coats, and surgical gloves) and were socially distanced as much as possible during the onsite portions of the study, and the lab was disinfected before and after each session. In the MRI, when alone in the scanner, participants were allowed the option of taking their masks off. Recent fMRI studies suggest that wearing a mask during the scan does not significantly affect task activation (50).

Music-Based Intervention

Prior to the start of the MBI, a Premium YouTube Music account was created for each participant for music listening. At the start of the MBI, participants met with a music therapist who familiarized them with the YouTube Music platform and assessed their music preferences. An energizing playlist and a relaxing playlist, of approximately 25 pieces of music each, were compiled, beginning with the six pieces selected during pre-screening. The music therapist instructed participants to listen to their choice of selections from one or both of their playlists for one hour daily, over the course of the next 8 weeks. They were asked to pay attention to how the music affected their moods and emotions, while noting any memories, images, or associations with the music. After each focused listening experience, they recorded these reflections and self-observations in a journal. Once a week, the music therapist called the participant to discuss their compliance and responses to the daily listening, and to help them add or remove pieces from their playlists. Prior to COVID-19, the start of the MBI occurred in person; however, during COVID-19, participants met with the music therapist over Zoom.

fMRI Task

The fMRI task consisted of 24 trials altogether. In each trial, participants were first presented with a musical stimulus (lasting 20 seconds), then they were given the task of rating how familiar they found the music to be (familiarity rating lasted 2 seconds), and how much they liked the music (liking rating also lasted 2 seconds). Musical stimuli for the MRI task consisted of 24 different audio excerpts, each of which was 20 seconds in duration. Each audio stimulus was from one of the following three categories: participant self-selected music (6/24 stimuli), other-selected (researcher-selected) music including well-known excerpts spanning multiple musical genres (42; 10/24 stimuli) and novel music spanning multiple genres (8/24 stimuli). A list of the researcher-selected musical selections is given in Supplementary Materials Table S1. Stimuli were presented in a randomized order, and participants made ratings of familiarity and liking on the scales of 1 to 4: for familiarity: 1=very unfamiliar, 2=unfamiliar, 3=familiar, 4=very familiar; for liking: 1=hate, 2=neutral, 3=like, 4=love. Participants made these ratings by pressing a corresponding button on a button-box (Cambridge Research Systems) inside the scanner. Participants wore MR-compatible over-the-ear headphones (Cambridge Research Systems) over musician-grade silicone ear plugs during MRI data acquisition. The spatial mapping between buttons and the numerical categories of ratings were counterbalanced between subjects to reduce any systematic association between familiarity or liking and the motor activity resulting from making responses. This fMRI task was completed before and after the MBI.

fMRI Data Acquisition

Images were acquired using a Siemens Magnetom 3T MR scanner with a 64-channel head coil at Northeastern University. For task fMRI data, continuous acquisition was used for 1440 volumes with a fast TR of 475 ms, for a total acquisition time of 11.4 minutes. Forty-eight axial slices (slice thickness = 3 mm, anterior to posterior, z volume = 14.4 mm) were acquired as echo-planar imaging (EPI) functional volumes covering the whole brain (TR = 475 ms, TE = 30 ms, flip angle = 60°, FOV = 240mm, voxel size = 3 x 3 x 3 mm³).

T1 images were also acquired using a MPRAGE sequence, with one T1 image acquired every 2400 ms, for a total task time of approximately 7 minutes. Sagittal slices (0.8 mm thick, anterior to posterior) were acquired covering the whole brain (TR = 2400 ms, TE = 2.55 ms, flip angle = 8°, FOV= 256, voxel size = 0.8 x 0.8 x 0.8 mm³). Resting state and diffusion tensor imaging data were also collected, but as they are outside the scope of the present study, they will be reported in another manuscript.

Data Analysis

Preprocessing

Task fMRI data was preprocessed using the Statistical Parametric Mapping 12 (SPM12) software (51) with the CONN Toolbox (52). Preprocessing steps included functional realignment and unwarp, functional centering, functional slice time correction, functional outlier detection using the artifact detection tool, functional direct segmentation and normalization to MNI template, structural centering, structural segmentation and normalization to MNI template, and functional smoothing to an 8mm gaussian kernel (53). Denoising steps for fMRI data included white matter and cerebrospinal fluid confound correction (54), and bandpass filtering to 0.008–0.09 Hz.

Univariate Whole-Brain Analysis

Response data from the fMRI task were imported into R Studio, and onset values were extracted for each trial, including its liking rating (hate, neutral, like, love), familiarity rating (very unfamiliar, unfamiliar, familiar, very familiar), and music condition (self-selected, other-selected well-known, other-selected novel).

First- and second-level analyses were completed in SPM12. For each participant, data were converted from 4D to 3D images, resulting in 1440 scans. The model was specified using the following criteria: interscan interval = 0.475 seconds, microtime resolution = 16, microtime onset = 8, duration = 42. Only data from the time while the participant was listening to the musical excerpt (rather than when the participant was making the familiarity and liking ratings) were included in this model. At the first-level, the following contrasts were extracted both pre- and post-intervention: the parametric effect of liking (linear contrast with all liking ratings), the parametric effect of familiarity (linear contrast with all familiarity ratings), and self-selected > other-selected (a combination of common researcher-selected and novel music. The resulting first-level contrasts were then analyzed using a one-sample t-test across all participants at the second level. Whole-brain results were rendered to a standard MNI brain. Results from the second-level analyses were statistically corrected using a cluster and voxel threshold of $p < 0.05$ (FDR-corrected) through CONN Toolbox (52). Data were also visualized through Conn.

Seed-Based Connectivity Analysis

Seed-based connectivity analysis was performed using the CONN toolbox (52) to determine changes in functional connectivity after participation in MBI. Seed ROIs consisted of Auditory

and Reward networks defined by previous work in our lab (29,47), with the effect of session used as a between-conditions contrast (post-intervention > pre-intervention). Significant clusters from these comparisons were then processed as new ROI masks, and beta time-series for these ROIs were extracted and compared trial-by-trial across the four liking and familiarity ratings for each subject.

Representational Similarity Analysis

Representational dissimilarity matrix (RDM) definition and comparisons were performed in MATLAB (Mathworks, R2020b) using tools from the RSA toolbox (55). RDMs used in this study are a combination of fMRI-derived RDMs and model RDMs. Selection and Novelty RDMs were generated by assigning each stimulus either a 1 or a 0 depending on if it was one of the six participant-selected stimuli or one of the 16 well-known stimuli (including participant-selected and researcher selected excerpts), respectively, and then subtracting each pair of stimulus values to form a binary difference matrix. The model RDM of Familiarity was defined as the difference in participant familiarity ratings between each pair of stimuli, and similarly the model of Liking used subject liking ratings.

fMRI-derived RDMs were generated using ROI-specific beta timeseries information for each stimulus. Each fMRI-derived RDM was defined as one minus the correlation coefficient of each pair of stimuli's average beta timeseries for a given ROI. Regions of interest included in this analysis consist of the 14 networks defined by the Stanford FIND lab atlas (56), as well as 12 hypothesis-driven ROIs, including ROIs for bilateral anterior and posterior STG, bilateral HG, NAcc, and supplementary motor area (SMA), the mPFC, and the Posterior Cingulate (PCC) derived from CONN Default and Networks Atlas (www.nitrc.org/projects/conn, RRID:SCR_009550). Finally, we included auditory and reward networks as defined in a previous study (29).

First-level, subject specific RDMs were generated for each ROI at both pre-intervention and post-intervention timepoints, and then compared to model RDMs. ROI to model comparisons were achieved through two independent analysis pipelines. In the first analysis pipeline, each first-level fMRI-derived RDM was compared to models of Selection, Novelty, Liking, and Familiarity through 10,000-fold bootstrap resampling (55,57) to generate relatedness values for each subject between each ROI-Model pair. These values then underwent 50,000 fold permutation T-max testing in order to determine significant ROI Model relationships at each session, and to determine any significant differences between sessions. The second analysis pipeline averaged first-level RDMs in order to form second-level, subject averaged RDMs. These second-level RDMs were then visualized using second order multidimensional scaling (55).

Acknowledgments

Supported by the Grammy Foundation, the Kim and Glen Campbell Foundation, NSF-STTR #2014870, NSF-CAREER award #1945436 to PL. We thank all our participants, NUBIC, and Juliet Davidow and the Northeastern University MRI Users Group for helpful comments.

References

1. Koelsch, S., Fritz, T., Schulze, K., Alsop, D., & Schlaug, G. (2005). Adults and children processing music: An fMRI study. *NeuroImage*, 25(4), 1068–1076.
2. Ellis, R. J., Bruijn, B., Norton, A. C., Winner, E., & Schlaug, G. (2013). Training-mediated leftward asymmetries during music processing: A cross-sectional and longitudinal fMRI analysis. *NeuroImage*, 75, 97–107.
3. Ferreri, L., Mas-Herrero, E., Zatorre, R. J., Ripollés, P., Gomez-Andres, A., Alicart, H., Olivé, G., Marco-Pallarés, J., Antonijoan, R. M., Valle, M., Riba, J., & Rodriguez-Fornells, A. (2019). Dopamine modulates the reward experiences elicited by music. *Proceedings of the National Academy of Sciences of the United States of America*, 116(9), 3793–3798.
4. Gold, B. P., Mas-Herrero, E., Zeighami, Y., Benovoy, M., Dagher, A., & Zatorre, R. J. (2019a). Musical reward prediction errors engage the nucleus accumbens and motivate learning. *Proceedings of the National Academy of Sciences of the United States of America*, 116(8), 3310–3315.
5. Salimpoor, V. N., Benovoy, M., Larcher, K., Dagher, A., & Zatorre, R. J. (2011). Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. *Nature Neuroscience*, 14(2), 257–264.
6. Salimpoor, V. N., Van Den Bosch, I., Kovacevic, N., McIntosh, A. R., Dagher, A., & Zatorre, R. J. (2013). Interactions between the nucleus accumbens and auditory cortices predict music reward value. *Science*, 340(6129), 216–219.
7. Olds, J., & Milner, P. (1954). Positive reinforcement produced by electrical stimulation of septal area and other regions of rat brain. *Journal of Comparative and Physiological Psychology*, 47(6), 419–427.
8. Knutson, B., Westdorp, A., Kaiser, E., & Hommer, D. (2000). fMRI visualization of brain activity during a monetary incentive delay task. *NeuroImage*, 12(1), 20–27.
9. Bartra, O., McGuire, J. T., & Kable, J. W. (2013). The valuation system: A coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *NeuroImage*, 76, 412–427. <https://doi.org/10.1016/j.neuroimage.2013.02.063>
10. Clithero, J. A., & Rangel, A. (2014). Informatic parcellation of the network involved in the computation of subjective value. *Social Cognitive and Affective Neuroscience*, 9(9), 1289–1302. <https://doi.org/10.1093/scan/nst106>

11. Gold, B. P., Pearce, M. T., Mas-Herrero, E., Dagher, A., & Zatorre, R. J. (2019b). Predictability and uncertainty in the pleasure of music: a reward for learning?. *Journal of Neuroscience*, 39(47), 9397-9409.
12. Cheung, V. K., Harrison, P. M., Meyer, L., Pearce, M. T., Haynes, J. D., & Koelsch, S. (2019). Uncertainty and surprise jointly predict musical pleasure and amygdala, hippocampus, and auditory cortex activity. *Current Biology*, 29(23), 4084-4092.
13. Savage, P. E., Loui, P., Tarr, B., Schachner, A., Glowacki, L., Mithen, S., & Fitch, W. T. (2020). Music as a coevolved system for social bonding. *Behavioral and Brain Sciences*, 1-36.
14. Janata, P. (2009). The neural architecture of music-evoked autobiographical memories. *Cerebral Cortex*, 19(11), 2579–2594.
15. Aydogan, G., Flaig, N., Ravi, S. N., Large, E. W., McClure, S. M., & Margulis, E. H. (2018). Overcoming Bias: Cognitive Control Reduces Susceptibility to Framing Effects in Evaluating Musical Performance. *Scientific Reports*, 8(1), 6229.
16. Alluri, V., Toiviainen, P., Burunat, I., Kliuchko, M., Vuust, P., & Brattico, E. (2017). Connectivity patterns during music listening: Evidence for action-based processing in musicians. *Human Brain Mapping*, 38(6), 2955–2970.
17. Andrews-Hanna, J. R. (2012). The brain's default network and its adaptive role in internal mentation. *The Neuroscientist*, 18(3), 251-270.
18. Whitfield-Gabrieli, S., & Ford, J. M. (2012). Default mode network activity and connectivity in psychopathology. *Annual review of clinical psychology*, 8, 49-76.
19. Koelsch, S., Vuust, P., & Friston, K. (2019). Predictive Processes and the Peculiar Case of Music. *Trends in Cognitive Sciences*, 23(1), 63–77.
20. Ruud, E. (1997). Music and the quality of life. *Nordic Journal of Music Therapy*, 6(2), 86–97.
21. Cassidy, G., & MacDonald, R. (2009). The effects of music choice on task performance: A study of the impact of self-selected and experimenter-selected music on driving game performance and experience. *Musicae Scientiae*, 13(2), 357–386.
22. Taruffi, L., Pehrs, C., Skouras, S., & Koelsch, S. (2017). Effects of sad and happy music on mind-wandering and the Default Mode Network. *Scientific Reports*, 7(1), 1–10.
23. Hanser, S. B. (2018). The new music therapist's handbook (3rd Ed.). Boston: Berklee Press.

24. Maratos, A., Crawford, M. J., & Procter, S. (2011). Music therapy for depression: It seems to work, but how?. *British Journal of Psychiatry*, 199(2), 92–93.
25. Maratos, A., Gold, C., Wang, X., & Crawford, M. (2008). Music therapy for depression. *Cochrane database of systematic reviews*, (1).
26. van der Steen, J. T., Smaling, H. J. A., van der Wouden, J. C., Bruinsma, M. S., Scholten, R. J. P. M., & Vink, A. C. (2018). Music-based therapeutic interventions for people with dementia. *Cochrane Database of Systematic Reviews*, (7).
27. Gotink, R. A., Meijboom, R., Vernooij, M. W., Smits, M., & Hunink, M. G. M. (2016). 8-week Mindfulness Based Stress Reduction induces brain changes similar to traditional long-term meditation practice – A systematic review. *Brain and Cognition*, 108, 32–41.
28. Hanser, S. B., & Thompson, L. W. (1994). Effects of a music therapy strategy on depressed older adults. *Journal of Gerontology*, 49(6), P265-P269.
29. Wang, D., Belden, A., Hanser, S. B., Geddes, M. R., & Loui, P. (2020). Resting-state connectivity of auditory and reward systems in Alzheimer’s Disease and Mild Cognitive Impairment. *Frontiers in Human Neuroscience*, 14, 1–10.
30. Freitas, C., Manzato, E., Burini, A., Taylor, M. J., Lerch, J. P., & Anagnostou, E. (2018). Neural correlates of familiarity in music listening: A systematic review and a neuroimaging meta-analysis. *Frontiers in Neuroscience*, 12, 1–14.
31. Pereira, C. S., Teixeira, J., Figueiredo, P., Xavier, J., Castro, S. L., & Brattico, E. (2011). Music and Emotions in the Brain: Familiarity Matters. *PLoS ONE*, 6(11).
32. Sternin, A., McGarry, L. M., Owen, A. M., & Grah, J. A. (2021). The Effect of Familiarity on Neural Representations of Music and Language. *Journal of Cognitive Neuroscience*, 1–17.
33. Christoff, K., Irving, Z. C., Fox, K. C. R., Spreng, R. N., & Andrews-Hanna, J. R. (2016). Mind-wandering as spontaneous thought: A dynamic framework. *Nature Reviews Neuroscience*, 17(11), 718–731.
34. Mas-Herrero, E., Maini, L., Sescousse, G., & Zatorre, R. J. (2021). Common and distinct neural correlates of music and food-induced pleasure: A coordinate-based meta-analysis of neuroimaging studies. *Neuroscience and Biobehavioral Reviews*, 123, 61–71. \
35. Saxe, R., & Kanwisher, N. (2003). People thinking about thinking people: the role of the temporo-parietal junction in “theory of mind”. *Neuroimage*, 19(4), 1835-1842.

36. Bedny, M., Pascual-Leone, A., & Saxe, R. R. (2009). Growing up blind does not change the neural bases of Theory of Mind. *Proceedings of the National Academy of Sciences*, 106(27), 11312-11317.
37. Lerdahl, F. (1992). Cognitive Constraints on Compositional Systems. *Contemporary Music Review*, 6(2), 97-121.
38. Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences*, 102(27), 9673-9678.
39. Blood, A. J., & Zatorre, R. J. (2001). Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proceedings of the National Academy of Sciences of the United States of America*, 98(20), 11818–11823.
40. Loui, P., Patterson, S., Sachs, M. E., Leung, Y., Zeng, T., & Przysinda, E. (2017). White matter correlates of musical anhedonia: Implications for evolution of music. *Frontiers in Psychology*, 8, 1664.
41. Martínez-Molina, N., Mas-Herrero, E., Rodríguez-Fornells, A., Zatorre, R. J., & Marco-Pallarés, J. (2019). White matter microstructure reflects individual differences in music reward sensitivity. *Journal of Neuroscience*, 39(25), 5018-5027.
42. Sachs, M. E., Ellis, R. J., Schlaug, G., & Loui, P. (2016). Brain connectivity reflects human aesthetic responses to music. *Social cognitive and affective neuroscience*, 11(6), 884-891.
43. Mas-Herrero, E., Dagher, A., & Zatorre, R. J. (2018). Modulating musical reward sensitivity up and down with transcranial magnetic stimulation. *Nature human behaviour*, 2(1), 27-32.
44. Kim, K., & Johnson, M. K. (2014). Extended self: spontaneous activation of medial prefrontal cortex by objects that are 'mine'. *Social cognitive and affective neuroscience*, 9(7), 1006-1012.
45. Nicolle, A., Klein-Flügge, M. C., Hunt, L. T., Vlaev, I., Dolan, R. J., & Behrens, T. E. (2012). An agent independent axis for executed and modeled choice in medial prefrontal cortex. *Neuron*, 75(6), 1114-1121.
46. Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: Anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*, 1124, 1–38.
47. Belfi, A. M., & Loui, P. (2020). Musical anhedonia and rewards of music listening: Current advances and a proposed model. *Annals of the New York Academy of Sciences*, 1464(1), 99–114.

48. McAuley, J. D., Wong, P. C. M., Mamidipaka, A., Phillips, N., & Margulis, E. H. (2021). Do you hear what I hear? Perceived narrative constitutes a semantic dimension for music. *Cognition*, 212, 104712.
49. Brandt, J., Spencer, M., & Folstein, M. (1988). The telephone interview for cognitive status. *Neuropsychiatry Neuropsychol Behav Neurol*, 1(2), 111–117.
50. Law, C. S. W., Lan, P. S., & Glover, G. H. (2021). Effect of wearing a face mask on fMRI BOLD contrast. *NeuroImage*, 229, 1–8.
51. Penny, W. D., Friston, K. J., Ashburner, J. T., Kiebel, S. J., & Nichols, T. E. (Eds.). (2011). *Statistical parametric mapping: the analysis of functional brain images*. Elsevier.
52. Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connectivity*, 2(3), 125–141.
53. Friston, K. J., Ashburner, J., Frith, C. D., Poline, J. B., Heather, J. D., & Frackowiak, R. S. (1995). Spatial registration and normalization of images. *Human brain mapping*, 3(3), 165–189.
54. Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage*, 37(1), 90–101.
55. Nili, H., Wingfield, C., Walther, A., Su, L., Marslen-Wilson, W., & Kriegeskorte, N. (2014). A toolbox for representational similarity analysis. *PLoS computational biology*, 10(4), e1003553.
56. Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD (2012). Decoding subject-driven cognitive states with whole-brain connectivity patterns. *Cereb Cortex*.
57. Kriegeskorte, N., Mur, M., & Bandettini, P. A. (2008). Representational similarity analysis-connecting the branches of systems neuroscience. *Frontiers in systems neuroscience*, 2, 4.

Tables

Table 1 | Parametric Effect of Liking - Anatomical labels based on CONN Toolbox, MNI coordinates, Voxel and Cluster Threshold: $p < 0.05$ (FDR corrected)

Region(s)/Clusters	X	Y	Z	Cluster Size
Pre-Intervention				
STG/MTG	-57	-36	15	465
Insular Cortex				
STG/MTG	51	-3	-15	431
PCC	-3	-36	33	233
TPJ/IPL	-42	-69	27	98
Superior Frontal Gyrus	-27	33	45	81
SMA	3	3	6	72
PCG	-48	-3	42	69
	-36	15	18	27
mPFC	0	60	0	45
Cerebellum	-33	-69	-27	45
	48	-3	36	26
Amygdala*	-3	-15	-15	21
Putamen*	24	9	3	16
Post-Intervention				
PCC	-9	-45	30	349
STG/MTG	-57	-24	-6	252
	57	-18	-12	192
TPJ/IPL	-45	-69	33	58
Cerebellum	12	-69	-18	26
mPFC	-9	57	9	24
Amygdala*	12	3	-15	12

Voxel Threshold: p FDR corrected < 0.05 ; Cluster Threshold: p FDR < 0.05

*Voxel Threshold: p FDR corrected < 0.05 ; Cluster Threshold: p uncorrected < 0.05

Table 2 | Parametric Effect of Familiarity - Anatomical labels based on CONN Toolbox, MNI coordinates, Voxel and Cluster Threshold: $p < 0.05$ (FDR corrected)

Region(s)/Clusters	X	Y	Z	Cluster Size
Pre-Intervention				
PCC				
TPJ/IPL				
STG/MTG				
Cerebellum				
PCG				
Postcentral Gyrus				
SMA				
Thalamus				
Hippocampus	3	-60	24	12793
Amygdala				
Caudate				
Putamen				
Parahippocampal Gyrus				
Insular Cortex				
Pallidum				
Post-Intervention				
STG/MTG	51	-18	-6	200
	-60	-15	-9	123
PCC	-3	-54	12	111
Supramarginal Gyrus	-51	-42	18	67
Cerebellum	12	-63	-18	40
Cingulate	-6	-42	33	24
Cerebellum	6	-51	-51	22
Voxel Threshold: p FDR corrected <0.05 ; Cluster Threshold: p uncorrected <0.05				

Table 3 | Self-Selected > Other-Selected - Anatomical labels based on CONN Toolbox, MNI coordinates, Voxel and Cluster Threshold: $p < 0.05$ (FDR corrected)

Region(s)/Clusters	X	Y	Z	Cluster Size
Pre-Intervention				
STG/HG				
mPFC				
PCC				
TPJ/IPL				
PCG				
Thalamus				
Insular Cortex				
Hippocampus	57	0	-9	17621
Amygdala				
Parahippocampal Gyrus				
Caudate				
Putamen				
Pallidum				
Nucleus Accumbens				
Orbitofrontal Cortex				
Putamen	-24	27	-15	147
Caudate				
Post-Intervention				
STG/MTG	-42	-42	15	812
	57	6	-9	392
PCC	-9	-54	18	540
Cerebellum	6	-51	-51	70
	15	-66	-18	68
	-12	-69	-18	39
mPFC	-9	54	-9	58
Parahippocampal Gyrus	-30	-21	-9	26
Hippocampus				
Thalamus*	-9	0	3	17

Voxel Threshold: p FDR corrected < 0.05 ; Cluster Threshold: p FDR corrected < 0.05

*Voxel Threshold: p FDR corrected < 0.05 ; Cluster Threshold: p uncorrected < 0.05

Figures

Figure 1. Study timeline and fMRI experiment design.

Figure 2. Univariate whole-brain analyses. (A) Parametric effects of liking pre- and post-intervention. Auditory (STG and HG), reward (dorsal and ventral striatum), and DMN (mPFC and PCC) regions showed activity that varied parametrically with liking ratings. Motor areas also showed activity that varied parametrically with liking ratings, but only pre-intervention. **(B) Parametric effects of familiarity pre- and post-intervention.** Auditory, reward, and DMN regions showed activity that also varied parametrically with familiarity ratings. The parametric effect of familiarity showed a much larger cluster pre-intervention than post-intervention, spanning motor areas as well as auditory and DMN areas. **(C) Self-selected > Other-selected music, pre- and post-intervention.** Pre-intervention, greater widespread engagement was observed in auditory, reward, and DMN regions for self-selected music compared to other-selected music. Post-intervention, greater engagement was observed in auditory, reward, and DMN regions for the same contrast. All images are results of second-level analyses showing significant clusters at the $p < .05$ FDR-corrected level.

Figure 3. Seed-based connectivity from auditory ROIs. (A) Cortical regions active during musical stimuli that received the highest liking rating (“Love”) pre- and post-intervention, showing significant activity in mPFC post-intervention but not pre-intervention. Voxelwise and clusterwise FDR-corrected $p < 0.05$. (B) The mPFC showed a significant effect of session (post-intervention > pre-intervention) in auditory network seed-based functional connectivity (voxel height $p < 0.001$, uncorrected; cluster size $p < 0.05$, FDR corrected). (C-D) Time-series data of ROIs extracted from the positive contrast in (B), representing post > pre-intervention increase in functional connectivity. (C) Beta series for different liking and familiarity ratings pre- and post-intervention, showing an increase in beta post-intervention, representing increased functional

connectivity between auditory regions and mPFC, for loved and very familiar music. (D) Beta series for self-selected and other-selected, and for well-known and novel music listening, showing an increase in auditory-mPFC functional connectivity after intervention for self-selected music but not for other-selected music. This increase in functional connectivity was not observed when comparing well-known vs. novel music listening. (E) Correlations between auditory and mPFC activity in beta values for the four levels of liking ratings before and after intervention. Each data point is one subject. Comparing across the panels from “Hate” (lowest liking rating) to “Love” (highest liking rating) shows more correlated auditory and mPFC activity during high liking ratings compared to low liking ratings, with the between-subject correlation being highest after the intervention for stimuli that received higher ratings.

Figure 4: Representational Similarity Analysis. (A) Schematic of RSA processing pipeline. Subject specific Ratings and task fMRI data are used to generate first-level representational dissimilarity matrices. First-Level RDMs are then used in 10,000-fold bootstrap resampling comparisons of models to functional RDMs for each subject. Resulting relatedness values are used in 50,000-fold t-max permutation testing (B) to determine significant (yellow) model-brain relationships at pre-intervention and post-intervention, as well as any significant effect of session ($p < 0.05$). First-Level RDMs are then averaged across subjects to generate second-level RDMs, which are used to produce second order multidimensional scaling visualizations of RDM relatedness (C).

Supplementary Materials

Table S1 | Researcher-Selected Song List

Title	Artist/Composer	Music Selection
Adagio for Strings	Barber	Well-Known
Symphony No. 2, Finale	Mahler	Well-Known
Cavatina Op. 130	Beethoven	Well-Known
John Wayne Gacy Jr.	Sufjan Stevens	Well-Known
Hurt	Johnny Cash	Well-Known
Sound of Silence	Simon and Garfunkel	Well-Known
Reckoner	Radiohead	Well-Known
Nocturne in C Sharp Minor	Chopin	Well-Known
Nessun Dorma	Andrea Bocelli	Well-Known
Symphony No. 6, movement 1	Tchaikovsky	Well-Known
Bohlen-Pierce Pan Flute Folk Tune	Arturo Grolimund	Novel
Hoquetus II	Johannes Kretz	Novel
Reminiscences	Steven Yi	Novel
Beyond The Horizon	Georg Hajdu	Novel
When the Moon Casts Shadows	Hubert Ho	Novel
Simmer	Hubert Ho	Novel
Vertigo Temporum	Hubert Ho	Novel
Manual Labor	Hubert Ho	Novel

Figure S1. Univariate whole-brain analyses for different levels of liking. (A) Love. Auditory and DMN regions showed activity both pre- and post-intervention, with the TPJ showing pre-intervention activity only. Motor areas showed deactivation both pre- and post-intervention. **(B) Like.** Auditory regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. **(C) Neutral.** Auditory regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. **(D) Hate.** Auditory

regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. All images are results of second-level analyses showing significant clusters at the $p < .05$ FDR-corrected level.

Figure S2. Univariate whole-brain analyses for different levels of familiarity. (A) Very Familiar. Auditory and DMN regions showed activity both pre- and post-intervention, with the TPJ showing pre-intervention activity only. Motor areas showed deactivation both pre- and post-intervention. **(B) Familiar.** Auditory regions showed activity both pre- and post-intervention, post-intervention to a greater extent. Motor regions showed deactivation only for post-intervention. **(C) Very Unfamiliar.** Auditory regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. **(D) Unfamiliar.** Auditory regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. All images are results of second-level analyses showing significant clusters at the $p < .05$ FDR-corrected level.

Figure S3. Univariate whole-brain analyses for music selection. (A) Self-Selected.

Auditory and DMN regions showed activity both pre- and post-intervention. Motor areas showed deactivation both pre- and post-intervention. **(B) Well-Known.** Auditory regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. **(C) Novel.** Auditory regions showed activity both pre- and post-intervention. Regions spanning the motor and DMN areas showed deactivation both pre- and post-intervention. All images are results of second-level analyses showing significant clusters at the $p < .05$ FDR-corrected level.

Figure S4: Second Level Model RDMs. (A) Time invariant binary RDMs representing stimulus selection (top) and novelty (bottom). (B) subject-averaged RDMs representing stimulus liking (top) and familiarity (bottom) ratings at the pre-intervention timepoint. (C) subject-averaged RDMs representing stimulus liking (top) and familiarity (bottom) ratings at the post-intervention timepoint.

Pre-Intervention

Figure S5: Pre-intervention fMRI-Derived RDMs. Participant-averaged RDMs representing differences in cortical responses to stimuli at the pre-intervention timepoint.

Post-Intervention

Figure S6: Post-intervention fMRI-Derived RDMs. Participant-averaged RDMs representing differences in cortical responses to stimuli at the post-intervention timepoint.

Figure S7: Bootstrap Comparisons of Selection RDM. results of 10,000-fold bootstrap comparisons of selection RDM to second level fMRI-derived RDMs at pre-intervention (left) and post-intervention (right) time points. Asterisk (*) represents a significant correlation between selection and a given fMRI-derived RDM ($p < 0.05$, FWE corrected), and Black bars represent a significant difference between fMRI-derived RDMs ($p < 0.05$, FWE corrected)

Figure S8: Bootstrap Comparisons of Novelty RDM. results of 10,000-fold bootstrap comparisons of novelty RDM to second level fMRI-derived RDMs at pre-intervention (left) and post-intervention (right) time points. Asterisk (*) represents a significant correlation between novelty and a given fMRI-derived RDM ($p < 0.05$, FWE corrected).

Figure S9: Bootstrap Comparisons of Liking RDM. results of 10,000-fold bootstrap comparisons of second-level liking RDM to second level fMRI-derived RDMs at pre-intervention (left) and post-intervention (right) time points. Asterisk (*) represents a significant correlation between liking and a given fMRI-derived RDM ($p < 0.05$, FWE corrected).