Title: Multivalency enhances the specificity of Fc-cytokine fusions

Authors: Brian Orcutt-Jahns¹, Peter C. Emmel¹, Eli M. Snyder¹, Cori Posner², Scott M. Carlson², Aaron S. Meyer^{1,3,4,5,*}

Affiliations:

¹Department of Bioengineering, University of California, Los Angeles, United States of America

²Visterra, Inc., Waltham, MA, United States of America

³Department of Bioinformatics, University of California, Los Angeles, United States of America

⁴Jonsson Comprehensive Cancer Center, University of California, Los Angeles, United States of America

⁵Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of

California, Los Angeles, United States of America

*Corresponding author. Email: ameyer@asmlab.org

Classification: Research report

Keywords: Cytokines, IL-2, IL-15, IL-7, Immunity, Multivalency

Author Emails/Contact Information:

Brian Orcutt-Jahns: orcuttjahnsbrian@gmail.com

Peter C. Emmel: petercemmel@gmail.com

Eli M. Snyder: emsnyder@g.ucla.edu

Cori Posner: Cori.posner@gmail.com

Scott M. Carlson: smcarlson@gmail.com

Aaron S. Meyer: ameyer@ucla.edu

Abstract: The common γ -chain receptor cytokines are promising immune therapies due to their central role in coordinating the proliferation and activity of various immune cell populations. One of these cytokines, interleukin (IL)-2, has potential as a therapy in autoimmunity but is limited in effectiveness by its modest specificity toward regulatory T cells (T_{reg} s). Therapeutic ligands are often made dimeric as antibody Fc fusions to confer desirable pharmacokinetic benefits, with under-explored consequences on signaling. Here, we systematically profiled the signaling responses to a panel of wild type and mutein IL-2 molecules in various Fc fusion configurations. We used a tensor-structured dimensionality reduction scheme to decompose the responses of each cell population to each ligand over a range of time points and cytokine concentrations. We found that dimeric muteins are uniquely specific for T_{reg} s at intermediate ligand concentrations. We then compared signaling response across all treatments to a simple, two-step multivalent binding model. Our model was able to predict cellular responses with high accuracy. Bivalent Fc fusions display enhanced specificity and potency for T_{reg} s through avidity effects toward IL-2R α . We then utilize our model to identify the potential benefits conferred by valency engineering as an additional mechanism for cytokines with optimized therapeutic benefits. In total, these findings represent a comprehensive analysis of how ligand properties, and their consequent effects on surface receptor-ligand interactions, translate to selective activation of immune cell populations. It also identifies a new route toward engineering even more selective therapeutic cytokines.

Significance Statement: Signaling in off-target immune cells has hindered the effectiveness of IL-2 as an immunotherapeutic. We show that bivalent IL-2 muteins exhibit more regulatory T cell-selective signaling than monovalent forms. This altered selectivity is explained by altered surface receptor-ligand binding kinetics and can be quantitatively predicted using a multivalent binding model. Finally, our model shows that even more selective IL-2 therapies may be developed by designing cytokines in higher valency formats, revealing valency as an unexplored mechanism for engineering specific IL-2 responses.

Main Text:

INTRODUCTION

Cytokines that bind to the common γ -chain (γ_c) receptor, including interleukin (IL)-2, 4, 7, 9, 15, and 21, are a critical hub in modulating both innate and adaptive immune responses (1). The cytokine family operates through a common theme of binding private receptors for each ligand before engaging the common γ_c receptor to induce signaling. A prominent phenotypic outcome of γ_c receptor signaling is lymphoproliferation, and so the cytokines are often observed to be an endogenous or exogenous mechanism for altering the balance of immune cell types. This phenotype is observed most extremely from loss-of-function or reduced activity mutations in γ_c which subvert T and NK cell maturation (2). Disruptive mutations in private receptors can lead to more selective reductions in cell types such as regulatory T cells (T_{reg} s) with IL-2R α or T cells with IL-7R α (1). Conversely, activating mutations in these receptors, such as IL-7R α , promote cancers such as B and T cell leukemias (3).

The importance of these cytokines to immune homeostasis and challenges in altering their signaling toward specific therapeutic goals have inspired a variety of engineered forms. Perhaps the most common approach has been to alter the receptor affinities of IL-2 to weaken its interaction with IL-2R α , IL-2R β , or both receptors. IL-2R α confers T_{reg} s with greater sensitivity

toward IL-2, and so IL-2R α affinity tunes the relative amount of signaling toward regulatory versus effector populations, while IL-2R β modulates the overall signaling potency (4). In most cases, the wild-type cytokine or mutein is fused to an IgG antibody to take advantage of FcRn-mediated recycling for extended half-life. The antibody has also been employed in a more active role by binding to IL-2 to influence its availability to each receptor in so-called immunocytokines (5). Fc fusion has taken many forms such as conjugation to the cytokine in an N-terminal or C-terminal orientation, including one or two cytokines per IgG, and including or excluding Fc effector functions (6). Notably, bivalent cytokine fusions have been shown to be more potent in T_{reg}-targeted engineered therapies, perhaps through pharmacokinetic or other means (7). The potential design space for these molecules quickly becomes experimentally intractable without consistent design principles.

To address this challenge, here we systematically evaluate the signaling specificity effects of engineered cytokine alterations, including affinity-altering mutations and Fc-fusion formats. We explore three hypotheses for the observed effect of bivalency in Fc-cytokine fusions. We find that the effect of bivalency can be fully explained by altered binding selectivity toward cell based on their abundances. The signaling specificity of all muteins and Fc-formats match well with a multivalent binding model, both between cell types and across cell-to-cell variation within a cell type. We propose that cytokine valency is an unexplored axis for further enhancing selective signaling responses and that many opportunities for using multivalency engineering exist within the γ_c cytokine family.

RESULTS

Multiple potential mechanisms of bivalency effects

Bivalent γ_c cytokines have been shown to display altered potency and cell-type-selective signaling profiles, both *in vitro* and *in vivo* (8). This alteration has led to both increases and reductions in the mutein's propensity for immune suppression; these variable outcomes may occur due to the differing affinities of the IL-2 mutein employed in the Fc fusions (7, 8). The mechanism by which multivalency induces altered signaling compared to monovalent ligands in the γ_c signaling pathways has not been systematically explored. The possible origins of the differences between monovalent and bivalent cytokine signaling profiles are varied, and each would influence the use of this alteration toward therapeutic ends.

As a first possibility, multivalent γ_c ligands may induce signaling through altered surface-receptor interactions (Fig. 1a). While monovalent ligands can only bind and ultimately transduce signal through a single receptor complex, multivalent ligands can potentially engage two or more complexes simultaneously. This change in signaling activity ultimately leads to an altered functional affinity, referred to as avidity. Among other effects, avidity leads to varied binding depending on the abundance of target receptors (9). In fact, this has been used with IL-2 to deliver cytotoxic cargo more selectively to T_{reg}S (10).

A second possible effect of multivalency is assembly of fundamentally distinct signaling complexes (Fig. 1b). Multivalent ligands, by binding multiple receptor complexes, can increase local receptor concentration leading to increases in downstream signal transduction (11). Clustering multiple IL-2R β / γ_c /IL-2 complexes may also help promote more potent or distinct forms of signaling compared to a single complex achievable with monovalent IL-2. Composition

of the active receptor complex is critical to the types of signaling responses observed, exemplified by the distinct signaling of “synthekines” that activate combinations of cytokine receptors (12).

Finally, multivalent ligands may have distinct responses due to pharmacokinetic (PK) or cell trafficking changes (Fig. 1c). IL-2 is rapidly cleared primarily through receptor-mediated endocytosis after Fc fusion, meaning that PK is closely tied to its activity (13). Nanoparticles and ligands of higher valency are endocytosed through altered mechanisms, ultimately leading to changes in intracellular degradation rates (14). The PK and *in vivo* half-lives of γ_c cytokines are also modulated by their valency, with effects, in turn, on PK (8). Changes in PK are especially important to selective expansion of T_{reg} s as this seems to be promoted most prominently by sustained signaling, over transient boluses of IL-2 (15). Overall, bivalency can have a multitude effects on the use of IL-2 therapeutically, in what experimental systems it will be observed, and how it might be further exploited, depending upon the mechanism of its effects.

Bivalent Fc-cytokine fusions have distinct cell specificity but shared dynamics

To explore the determinants of IL-2 response in a systematic manner, we stimulated peripheral blood mononuclear cells (PBMCs) collected from a single donor with 13 IL-2 muteins with varying receptor affinities in both monomeric and dimeric Fc-fusion formats (Table S1). Stimulated cells were collected at four time points using 12 treatment concentrations. The PBMCs were then stained for canonical cell type markers and pSTAT5, a read-out of signaling response, allowing us to separate response by cell type. Four different cell types (T_{reg} , T_{helper} , CD8+, NK) were gated and quantified (Fig. S1a-d). T_{reg} s, and T_{helper} s were further dissected according to their IL-2R α abundances into low, average, and high expression subpopulations by isolating sub-populations using three evenly logarithmically spaced bins (Fig. S1j).

Exploring how dynamic responses vary across responding cell types and ligand treatments is challenging due to the multi-dimensional space of potential origins of variations. Restricting ones’ view to a single time point, cell type, or ligand concentration provides only a slice of the picture (6, 16). To provide a more comprehensive view of response, we organized our profiling experiments into a four-dimensional tensor, wherein the position along each dimension represented the ligand used, concentration, treatment duration, or cell type in the profiling. We then factored this data using non-negative canonical polyadic (CP) decomposition to derive factors summarizing the influence of each dimension (Fig. 2a). Three components explained roughly 90% of the variance within the dataset (Fig. 2b).

Factorization separated distinct response profiles into separate components, and the effect of each dimension into separate factors. For instance, component 1 almost exclusively represented responses to wild-type cytokines (Fig. 2c), which were the only ligands which were not Fc-conjugated, showing a response primarily at high concentrations (Fig. 2d), with broad specificity (Fig. 2e) and a signaling profile that peaks at 30 minutes and then more rapidly decreases (Fig. 2f). An alternative way to interpret the factorization results is to compare profiles within a single dimension. For example, component 1 led to a less sustained profile of signaling response as compared to the other signaling patterns (Fig. 2f).

Remarkably, components 2 and 3 cleanly separated ligands conjugated in bivalent or monovalent forms respectively (Fig. 2c). In fact, ligand valency was represented more prominently than

differences in receptor affinity between muteins. Component 2 had uniquely high T_{reg} specificity (Fig. 2e) and was most represented at intermediate concentrations (Fig. 2d). Component 2 was also highly correlated with IL-2R α abundance in subsets of T_{reg} and T_{helper} cells, suggesting that bivalent molecule's specificity for T_{reg} s is mediated by their higher abundance of IL-2R α . Component 3 had a broad cell response (Fig. 2e) and increased monotonically with concentration (Fig. 2d). Despite these strong differences in specificities both components had nearly identical time dynamics (Fig. 2f). In total, these results indicated ligand valency as a critical determinant of IL-2 specificity, and that while Fc fusion subtly affected the specificity of response, monovalent and bivalent ligands had identical dynamics.

Differences among IL-2 responses are explained by a simple multivalent binding model

With the indication that T_{reg} specificity is enhanced by multivalency without changes to signaling dynamics, we sought to evaluate how well cell surface binding on its own could predict response. We employed a two-step, equilibrium, multivalent binding model to predict cellular response to IL-2 muteins by assuming that signaling response was proportional to the amount of active receptor-ligand complexes (17). See methods for specifics about how binding predictions were used to predict pSTAT5 response. We fit this model to our signaling profiling experiments and evaluated its concordance with the data. The only non-scaling fitting parameter, K_x^* , had an optimum at 1.2×10^{-11} #/cell, consistent with that seen for other receptor families (18–20). Overall, we observed remarkable consistency between predicted and observed response ($R^2 = 0.85$; Fig. 3b). This high accuracy was maintained within subsets of the data including individual cell types and ligands (Fig. 3c–d). To verify our model's predictions were dependent on the valency of the molecule, we again fit the model enforcing that all ligands were monovalent. This significantly reduced model accuracy for many of the bivalent ligands (Fig. 3e), confirming that multivalent binding was critical in modeling signaling response.

To ensure that our model was not simply capturing a trend towards higher signaling with increasing concentration, we examined our model's accuracy within specific cytokine concentrations (Fig. 3f). Our model was unable to predict response at the lowest concentrations as these did not stimulate signaling and were dominated by experimental noise but increased in accuracy at concentrations where any signaling response was observed. Finally, we examined how our model's accuracy varied with time by fitting the model to each time point individually. As expected, given that longer treatments likely involve various compensatory signaling such as the degradation or increased transcription of IL-2 receptor subunits (21, 22), we most accurately predicted initial responses (30 mins) with a slight decrease in accuracy over longer timescales (Fig. 3g). In total, multivalent cell surface binding showed quantitative agreement with the pattern of cell-type-specific responses to IL-2 muteins, supporting that the specificity enhancement of bivalency is derived from cell surface binding avidity effects.

Multivalency provides a general strategy for further enhanced signaling selectivity

Given that a simple binding model accurately predicted cell-type-specific responses to IL-2 and that bivalent, Fc-fused IL-2 muteins have favorable specificity properties, we sought to computationally explore to what extent multivalency might be a generally useful strategy. While monovalent ligand binding scales linearly with receptor abundance, multivalent ligands bind nonlinearly depending upon receptor abundance (23). Thus, multivalent ligands should be able to selectively target cells with uniquely high expression of certain γ_c family receptors.

Valency enhancements are only apparent with coordinated changes in receptor-ligand binding affinities (24). Therefore, we optimized the receptor affinities of simulated ligands while varying valency. We first designed IL-2 muteins of varying valency to obtain optimal T_{reg} specificity (Fig. 4a). As expected, ligand valency increased achievable selectivity past what was achievable using the monovalent cytokine format at any receptor affinity. Higher valency required reduced IL-2R α affinities (Fig. 4b).

We then explored whether IL-2 muteins lacking IL-2R α binding could selectively target NK cells, based on their uniquely high expression of IL-2R β , with similar results; IL-2 muteins of higher valency were predicted to be increasingly selective for activation of NK cells, so long as IL-2R β / γ_c affinity was coordinately decreased (Fig. 4c,d). Finally, we sought to explore T_{helper} -selective muteins of IL-7, as they express high amounts of IL-7R α (Fig. S1i). We again found that ligands of higher valency should achieve higher degrees of selectivity for these cells, but that the benefits of valency were less than the targeting of T_{reg} s or NK cells using IL-2 mutants because CD8 $^+$ T cells have similar IL-7R α amounts (Fig. 4e). These benefits were again found to be contingent on decreasing affinity of the IL-7 muteins for IL-7R α at higher valency (Fig. 4f). In total, these results show that valency has unexplored potential for designing cytokines with enhanced therapeutic efficacy and reduced toxicity. They also show the potential benefit of the modeling framework described above in guiding therapeutic development.

DISCUSSION

Here, we have systematically explored how ligand properties determine signaling response across 13 ligands including wild-type IL-2, and both monovalent and bivalent Fc fusion IL-2 variants. A tensor-based dimensionality reduction technique identified the patterns of changing response with ligand properties, revealing that multivalent cytokines have unique specificity but identical dynamics (Fig. 2). The mechanism responsible for this unique specificity was then explored using a multivalent binding model, which able to reproduce cell-type-specific responses to IL-2 muteins with high accuracy, indicating that specificity is derived from surface binding avidity effects (Fig. 3). Through this model, we found that cytokines of higher valency offer even greater cell type selectivity given corresponding affinity adjustments, which should translate to therapies of improved potency and reduced toxicity (Fig. 4).

The design of cell-type-selective ligands is complicated by the complex signaling processes by which they induce cellular signaling and ultimately response, which leads to a combinatorial explosion of potential ligand design objectives (25). This work serves to demonstrate the importance of the relatively unexplored axis of ligand development represented by the design of multivalent ligands. Multivalent ligands have several documented effects, including altered signal transduction (11, 26), binding avidity, and pharmacokinetics (27) or intracellular trafficking (28). While valency has been extensively explored as a means to introduce binding selectivity based on receptor density (9, 29), how this effect interacts with the presence of multiple receptor species and cell signaling is surely more complex. The ability of the binding model described in this work to accurately predict immune cell-type-specific response indicates that the cell-type-specific signaling profiles of these cytokines can still be understood as principally arising through receptor avidity effects at the cell surface (Fig. 3).

Our work here may be used to guide the design of IL-2 muteins with high selectivity for T_{reg} s, an important criterion in the design of IL-2 based treatments for autoimmune diseases (30, 31). By

optimizing the predicted selectivity of high-valency ligands, we showed that multivalency may be exploited to design more effective IL-2 based therapeutics for use in the clinical setting, where IL-2 based therapies have traditionally struggled (Fig. 4a,b) (32). Combined with the superior *in vivo* half-life conferred by Fc-conjugated IL-2 muteins, these multivalent therapeutics could potentially be used in an out-patient setting and require less frequent dosing (33). Furthermore, the superior selectivity offered by engineered multivalent ligands will likely further increase their *in vivo* half-lives, due to a reduction in receptor-mediated clearance by off-target populations. We also demonstrated the potential benefit which multivalency may confer in the selective activation of NK cells, which could lead to similarly improved anti-cancer treatments (34). Our approach here may also be applied to engineer selectivity into other signaling pathways characterized by cell type pleiotropy, such as IL-4/IL-13 or TNF systems (35, 36).

While our approach does effectively capture cell type specific responses to IL-2 and IL-2 muteins and can be readily applied to other well-studied signaling families, several challenges in its implementation and broad translation to therapeutic development remain. Our model can be used to generate guidelines for the modulation of receptor-ligand interactions in conjunction with valency engineering to design more selective ligands, but also predicts that there are specific receptor affinities required to achieve these benefits (Fig. 4). While a host of methods have been developed and employed to enhance or ablate cytokine affinity for receptor subunits (37), precisely tuning ligand affinities remains a complicated protein engineering challenge. Furthermore, we have not yet experimentally validated whether the advantages conferred by higher valency ligands *in vitro* translate to *in vivo* settings. For example, a research group found that while a bivalent IL-2 molecule conjugated to a diphtheria-based toxin showed an enhanced ability to target and eliminate T_{reg} s in an *in vitro* setting (38), its enhanced selectivity failed to be replicated in *in vivo* trials and did not eliminate T_{reg} s any more selectively than its monovalent counterpart (39). However, the increased efficacy and reduced toxicity of a bivalent EGFR-based therapy over its monovalent counterpart in an *in vivo* setting was demonstrated by the same group, suggesting that the specificity conferred by multivalent ligands can indeed transfer to clinical settings (40). This discrepancy between *in vivo* and *in vitro* selectivity indicates that further experimental modeling is undoubtedly necessary to explore the *in vivo* benefits of multivalent IL-2 therapies.

MATERIALS AND METHODS

Modeling

Binding model

Model was formulated as described in Tan *et al* (24). The monomer composition of a ligand complex was represented by a vector $\theta = (\theta_1, \theta_2, \dots, \theta_{N_L})$, where each θ_i was the number of monomer ligand type i on that complex. Let C_θ be the proportion of the θ complexes in all ligand complexes, and Θ be the set of all possible θ 's. We have $\sum_{\theta \in \Theta} C_\theta = 1$.

The binding between a ligand complex and a cell expressing several types of receptors can be represented by a series of q_{ij} . The relationship between q_{ij} 's and θ_i is given by $\theta_i = q_{i0} + q_{i1} + \dots + q_{iN_R}$. Let the vector $\mathbf{q}_i = (q_{i0}, q_{i1}, \dots, q_{iN_R})$, and the corresponding θ of a binding configuration $\mathbf{q}$ be $\theta(\mathbf{q})$. For all i in $\{1, 2, \dots, N_L\}$, we define $\psi_{ij} = R_{eq,j} K_{a,ij} K_x^*$ where $j =$

275 $\{1, 2, \dots, N_R\}$ and $\psi_{i0} = 1$. The relative number of complexes bound to a cell with configuration
276 $\mathbf{q}$ at equilibrium is

$$277 \quad v_{\mathbf{q},eq} = \frac{L_0 C_{\theta}(\mathbf{q})}{K_x^*} \prod_{i=1}^{N_L} \prod_{j=0}^{N_R} \psi_{ij}^{q_{ij}} \prod_{i=1}^{N_L} \binom{\theta_i}{q_i}.$$

278 Then we can calculate the relative amount of bound receptor n as

$$279 \quad R_{\text{bound},n} = \frac{L_0}{K_x^*} \sum_{\theta \in \Theta} C_{\theta} \left[\sum_{i=1}^{N_L} \frac{\psi_{in} \theta_i}{\sum_{j=0}^{N_R} \psi_{ij}} \right] \prod_{i=1}^{N_L} \left(\sum_{j=0}^{N_R} \psi_{ij} \right)^{\theta_i}.$$

280 By $R_{\text{tot},n} = R_{\text{eq},n} + R_{\text{bound},n}$, we can solve $R_{\text{eq},n}$ numerically for each type of receptor.

281 Application of multivalent binding model to IL-2 signaling pathway

282 Each IL-2 molecule was allowed to bind to one free IL-2R α and one IL-2R β/γ_c receptor. Initial
283 IL-2-receptor association proceeds with the known kinetics of monomeric ligand-receptor
284 interaction (Table S1). Subsequent ligand-receptor binding interactions then proceed with an
285 association constant proportional to available receptor abundance and affinity multiplied by the
286 scaling constant, K_x^* , as described above. To predict pSTAT5 response to IL-2 stimulation, we
287 assumed that pSTAT5 is proportional to the amount of IL-2-bound IL-2R β/γ_c , as complexes
288 which contain these species actively signal through the JAK/STAT pathway. Scaling factors
289 converting from predicted active signaling species to pSTAT5 abundance were fit to
290 experimental data on a per-experiment and cell type basis. A single K_x^* value was fit for all
291 experiments and cell types.

292 Tensor Factorization

293 Before decomposition, the signaling response data was background subtracted and variance
294 scaled across each cell population. Tensor decomposition was performed using the Python
295 package TensorLy (41), using non-negative canonical polyadic decomposition.

296 Experimental Methods

297 All experimental methods were performed as described in Farhat *et al.* (42)

298 Receptor abundance quantitation

299 Cryopreserved PBMCs (ATCC, PCS-800-011, lot#81115172) were thawed to room temperature
300 and slowly diluted with 9 mL pre-warmed RPMI-1640 medium (Gibco, 11875-093)
301 supplemented with 10% fetal bovine serum (FBS, Seradigm, 1500-500, lot#322B15). Media was
302 removed, and cells washed once more with 10 mL warm RPMI-1640 + 10% FBS. Cells were
303 brought to 1.5×10^6 cells/mL, distributed at 250,000 cells per well in a 96-well V-bottom plate,
304 and allowed to recover 2 hrs at 37°C in an incubator at 5% CO₂. Cells were then washed twice
305 with PBS + 0.1% BSA (PBSA, Gibco, 15260-037, Lot#2000843) and suspended in 50 μ L PBSA
306 + 10% FBS for 10 min on ice to reduce background binding to IgG.

Antibodies were diluted in PBSA + 10% FBS and cells were stained for 1 hr at 4°C in darkness with a gating panel (Panel 1, Panel 2, Panel 3, or Panel 4) and one anti-receptor antibody, or an equal concentration of matched isotype/fluorochrome control antibody. Stain for CD25 was included in Panel 1 when CD122, CD132, CD127, or CD215 was being measured (CD25 is used to separate T_{reg}S from other CD4⁺ T cells).

Compensation beads (Simply Cellular Compensation Standard, Bangs Labs, 550, lot#12970) and quantitation standards (Quantum Simply Cellular anti-Mouse IgG or anti-Rat IgG, Bangs Labs, 815, Lot#13895, 817, Lot#13294) were prepared for compensation and standard curve. One well was prepared for each fluorophore with 2 µL antibody in 50 µL PBSA and the corresponding beads. Bead standards were incubated for 1 hr at room temperature in the dark.

Both beads and cells were washed twice with PBSA. Cells were suspended in 120 µL per well PBSA, and beads to 50 µL, and analyzed using an IntelliCyt iQue Screener PLUS with VBR configuration (Sartorius) with a sip time of 35 and 30 secs for cells and beads, respectively. Antibody number was calculated from fluorescence intensity by subtracting isotype control values from matched receptor stains and calibrated using the two lowest binding quantitation standards. T_{reg} cells could not be gated in the absence of CD25, so CD4⁺ T cells were used as the isotype control to measure CD25 in T_{reg} populations. Cells were gated as shown in Fig. S1. Measurements were performed using four independent staining procedures over two days. Separately, the analysis was performed with anti-receptor antibodies at 3x normal concentration to verify that receptor binding was saturated. Replicates were summarized by geometric mean.

pSTAT5 Measurement of IL-2 and -15 Signaling in PBMCs

Human PBMCs were thawed, distributed across a 96-well plate, and allowed to recover as described above. IL-2 (R&D Systems, 202-IL-010) or IL-15 (R&D Systems, 247-ILB-025) were diluted in RPMI-1640 without FBS and added to the indicated concentrations. To measure pSTAT5, media was removed, and cells fixed in 100 µL of 10% formalin (Fisher Scientific, SF100-4) for 15 mins at room temperature. Formalin was removed, cells were placed on ice, and cells were gently suspended in 50 µL of cold methanol (-30°C). Cells were stored overnight at -30°C. Cells were then washed twice with PBSA, split into two identical plates, and stained 1 hr at room temperature in darkness using antibody panels 4 and 5 with 50 µL per well. Cells were suspended in 100 µL PBSA per well, and beads to 50 µL, and analyzed on an IntelliCyt iQue Screener PLUS with VBR configuration (Sartorius) using a sip time of 35 seconds and beads 30 seconds. Compensation was performed as above. Populations were gated as shown in Fig. S1, and the median pSTAT5 level extracted for each population in each well.

Recombinant proteins

IL-2/Fc fusion proteins were expressed using the Expi293 expression system according to manufacturer instructions (Thermo Scientific). Proteins were as human IgG1 Fc fused at the N- or C-terminus to human IL-2 through a (G4S)₄ linker. C-terminal fusions omitted the C-terminal lysine residue of human IgG1. The AviTag sequence GLNDIFEAQKIEWHE was included on whichever terminus did not contain IL-2. Fc mutations to prevent dimerization were introduced into the Fc sequence (43). Proteins were purified using MabSelect resin (GE Healthcare). Proteins were biotinylated using BirA enzyme (BPS Biosciences) according to manufacturer instructions, and extensively buffer-exchanged into phosphate buffered saline (PBS) using

Amicon 10 kDa spin concentrators (EMD Millipore). The sequence of IL-2R β / γ Fc heterodimer was based on a reported active heterodimeric molecule (patent application US20150218260A1), with the addition of (G4S)₂ linker between the Fc and each receptor ectodomain. The protein was expressed in the Expi293 system and purified on MabSelect resin as above. IL2-R α ectodomain was produced with C-terminal 6xHis tag and purified on Nickel-NTA spin columns (Qiagen) according to manufacturer instructions.

Octet binding assays

Binding affinity was measured on an Octet RED384 (ForteBio). Briefly, biotinylated monomeric IL-2/Fc fusion proteins were uniformly loaded to Streptavidin biosensors (ForteBio) at roughly 10% of saturation point and equilibrated for 10 minutes in PBS + 0.1% bovine serum albumin (BSA). Association time was up to 40 minutes in IL-2R β / γ titrated in 2x steps from 400 nM to 6.25 nM, or IL-2R α from 25 nM to 20 pM, followed by dissociation in PBS + 0.1% BSA. A zero-concentration control sensor was included in each measurement and used as a reference signal. Assays were performed in quadruplicate across two days. Binding to IL-2R α did not fit to a simple binding model so equilibrium binding was used to determine the K_D within each assay. Binding to IL-2R β / γ fit a 1:1 binding model so on-rate (k_{on}), off-rate (k_{off}) and K_D were determined by fitting to the entire binding curve. Kinetic parameters and K_D were calculated for each assay by averaging all concentrations with detectable binding signal (typically 12.5 nM and above).

Statistical analysis

The number of replicates performed for each experimental measurement, and the values of confidence intervals are described in corresponding figure captions. N is used to describe the number of times a particular experiment was performed. All flow cytometry experiments performed on hPBMCs were conducted using separate experimental replicates on cells gathered from a single donor. To quantify population-level flow cytometry measurements for both signaling and receptor quantitation experiments, the mean fluorescent intensity (MFI) of a gated population was measured. Compensation to removed fluorescent spectral overlap was performed for each experimental measurement. Subtraction of either negative controls or cells treated with isotype antibodies was performed on signaling and receptor quantitation data respectively to remove background signal. Cells which were measured to display fluorescent intensities above 1,000,000 were excluded from analysis during signaling experiments. Pearson correlation coefficients (R^2) values were used to describe model accuracy when predicting signaling response to IL-2 and IL-2 muteins. The K_x^* parameter was fit using least-squares fitting and the Broyden–Fletcher–Goldfarb–Shanno minimization algorithm as implemented in SciPy.

References

1. W. J. Leonard, J.-X. Lin, J. J. O'Shea, The γc Family of Cytokines: Basic Biology to Therapeutic Ramifications, *Immunity* **50**, 832–850 (2019).
2. M. Noguchi, H. Yi, H. M. Rosenblatt, A. H. Filipovich, S. Adelstein, W. S. Modi, O. W. McBride, W. J. Leonard, Interleukin-2 receptor gamma chain mutation results in X-linked severe combined immunodeficiency in humans., *Cell* **73**, 147–57 (1993).

3. P. P. Zenatti, D. Ribeiro, W. Li, L. Zuurbier, M. C. Silva, M. Paganin, J. Tritapoe, J. A. Hixon, A. B. Silveira, B. A. Cardoso, L. M. Sarmiento, N. Correia, M. L. Toribio, J. Kobarg, M. Horstmann, R. Pieters, S. R. Brandalise, A. A. Ferrando, J. P. Meijerink, S. K. Durum, J. A. Yunes, J. T. Barata, Oncogenic IL7R gain-of-function mutations in childhood T-cell acute lymphoblastic leukemia, *Nature Genetics* **43**, 932–939 (2011).
4. S. Mitra, A. M. Ring, S. Amarnath, J. B. Spangler, P. Li, W. Ju, S. Fischer, J. Oh, R. Spolski, K. Weiskopf, H. Kohrt, J. E. Foley, S. Rajagopalan, E. O. Long, D. H. Fowler, T. A. Waldmann, K. C. Garcia, W. J. Leonard, Interleukin-2 activity can be fine-tuned with engineered receptor signaling clamps., *Immunity* **42**, 826–38 (2015).
5. J. B. Spangler, J. Tomala, V. C. Luca, K. M. Jude, S. Dong, A. M. Ring, P. Votavova, M. Pepper, M. Kovar, K. C. Garcia, Antibodies to Interleukin-2 Elicit Selective T Cell Subset Potentiation through Distinct Conformational Mechanisms., *Immunity* **42**, 815–25 (2015).
6. A. M. Farhat, A. C. Weiner, C. Posner, Z. S. Kim, B. Orcutt-Jahns, S. M. Carlson, A. S. Meyer, Modeling Cell-Specific Dynamics and Regulation of the Common Gamma Chain Cytokines, *Cold Spring Harbor Laboratory* (2020), doi:10.1101/778894.
7. L. Khoryati, M. N. Pham, M. Sherve, S. Kumari, K. Cook, J. Pearson, M. Bogdani, D. J. Campbell, M. A. Gavin, An IL-2 mutein engineered to promote expansion of regulatory T cells arrests ongoing autoimmunity in mice, *Science Immunology* **5**, eaba5264 (2020).
8. C. J. M. Bell, Y. Sun, U. M. Nowak, J. Clark, S. Howlett, M. L. Pekalski, X. Yang, O. Ast, I. Waldhauer, A. Freimoser-Grundschober, E. Moessner, P. Umana, C. Klein, R. J. Hosse, L. S. Wicker, L. B. Peterson, Sustained in vivo signaling by long-lived IL-2 induces prolonged increases of regulatory T cells., *J Autoimmun* **56**, 66–80 (2014).
9. C. M. Csizmar, J. R. Petersburg, T. J. Perry, L. Rozumalski, B. J. Hackel, C. R. Wagner, Multivalent Ligand Binding to Cell Membrane Antigens: Defining the Interplay of Affinity, Valency, and Expression Density., *J Am Chem Soc* **141**, 251–261 (2018).
10. J. S. Peraino, H. Zhang, P. V. Rajasekera, M. Wei, J. C. Madsen, D. H. Sachs, C. A. Huang, Z. Wang, Diphtheria toxin-based bivalent human IL-2 fusion toxin with improved efficacy for targeting human CD25(+) cells., *J Immunol Methods* **405**, 57–66 (2014).
11. L. L. Kiessling, J. E. Gestwicki, L. E. Strong, Synthetic Multivalent Ligands as Probes of Signal Transduction, *Angewandte Chemie International Edition* **45**, 2348–2368 (2006).
12. I. Moraga, J. B. Spangler, J. L. Mendoza, M. Gakovic, T. S. Wehrman, P. Krutzik, K. C. Garcia, Synthekines are surrogate cytokine and growth factor agonists that compel signaling through non-natural receptor dimers, *eLife* **6**, e22882 (2017).
13. S. Gillies, Y. Lan, B. Brunkhorst, W.-K. Wong, Y. Li, K.-M. Lo, Bi-functional cytokine fusion proteins for gene therapy and antibody-targeted treatment of cancer, *Cancer Immunology, Immunotherapy* **51**, 449–460 (2002).

14. S. J. York, L. S. Arneson, W. T. Gregory, N. M. Dahms, S. Kornfeld, The rate of internalization of the mannose 6-phosphate/insulin-like growth factor II receptor is enhanced by multivalent ligand binding., *J Biol Chem* **274**, 1164–71 (1999).
15. R. Varma, K. Liu, K. Avery, R. Rashid, S. Schubert, I. Leung, C. Bonzon, L. Bogaert, J. Desjarlais, M. J. Bennett, Regulatory T Cell Selective IL-2-Fc Fusion Proteins for the Treatment of Autoimmune Diseases, *Blood* **132**, 3709–3709 (2018).
16. M. Murphy, S. D. Taylor, A. S. Meyer, Structured decomposition improves systems serology prediction and interpretation, *Cold Spring Harbor Laboratory* (2021), doi:10.1101/2021.01.03.425138.
17. Z. C. Tan, A. S. Meyer, A general model of multivalent binding with ligands of heterotypic subunits and multiple surface receptors, *Cold Spring Harbor Laboratory* (2021), doi:10.1101/2021.03.10.434776.
18. R. A. Robinett, N. Guan, A. Lux, M. Biburger, F. Nimmerjahn, A. S. Meyer, Dissecting FcγR Regulation through a Multivalent Binding Model., *Cell Syst* **7**, 41–48.e5 (2018).
19. A. S. Perelson, C. DeLisi, *Receptor clustering on a cell surface. I. theory of receptor cross-linking by ligands bearing two chemically identical functional groups* (Elsevier, 1980; <https://www.sciencedirect.com/science/article/abs/pii/0025556480900176>).
20. J. D. Stone, J. R. Cochran, L. J. Stern, T-cell activation by soluble MHC oligomers can be described by a two-parameter binding model., *Biophys J* **81**, 2547–57 (2001).
21. R. F. Pass, C. Hutto, R. Ricks, G. A. Cloud, Increased rate of cytomegalovirus infection among parents of children attending day-care centers., *N Engl J Med* **314**, 1414–8 (1986).
22. L. T. Yamamoto, H. N. Wigder, D. J. Fligner, M. Rauen, R. A. Dershewitz, Relationship of bacteremia to antipyretic therapy in febrile children., *Pediatr Emerg Care* **3**, 223–7 (1987).
23. Y. Mazor, K. F. Sachsenmeier, C. Yang, A. Hansen, J. Filderman, K. Mulgrew, H. Wu, W. F. Dall'Acqua, Enhanced tumor-targeting selectivity by modulating bispecific antibody binding affinity and format valence, *Scientific Reports* **7**, 40098 (2017).
24. Z. C. Tan, B. Orcutt-Jahns, A. S. Meyer, A quantitative view of strategies to engineer cell-selective ligand binding, *Cold Spring Harbor Laboratory* (2020), doi:10.1101/2020.11.25.398974.
25. Y. E. Antebi, J. M. Linton, H. Klumpe, B. Bintu, M. Gong, C. Su, R. McCardell, M. B. Elowitz, Combinatorial Signal Perception in the BMP Pathway, *Cell* **170**, 1184–1196.e24 (2017).
26. I. Moraga, J.B. Spangler, J.L. Mendoza, M. Gakovic, T.S. Wehrman, P. Krutzik, K.C. Garcia, SyntheKines are surrogate cytokine and growth factor agonists that compel signaling through non-natural receptor dimers, *eLife* (2017).
27. Á. M. Cuesta, N. Sainz-Pastor, J. Bonet, B. Oliva, L. Álvarez-Vallina, Multivalent antibodies: when design surpasses evolution, *Trends in Biotechnology* **28**, 355–362 (2010).

28. S. J. York, L. S. Arneson, W. T. Gregory, N. M. Dahms, S. Kornfeld, The Rate of Internalization of the Mannose 6-Phosphate/Insulin-like Growth Factor II Receptor Is Enhanced by Multivalent Ligand Binding, *Journal of Biological Chemistry* **274**, 1164–1171 (1999).
29. C. B. Carlson, P. Mowery, R. M. Owen, E. C. Dykhuizen, L. L. Kiessling, Selective Tumor Cell Targeting Using Low-Affinity, Multivalent Interactions, *ACS Chemical Biology* **2**, 119–127 (2007).
30. L. B. Peterson, C. J. M. Bell, S. K. Howlett, M. L. Pekalski, K. Brady, H. Hinton, D. Sauter, J. A. Todd, P. Umana, O. Ast, I. Waldhauer, A. Freimoser-Grundschober, E. Moessner, C. Klein, R. J. Hosse, L. S. Wicker, A long-lived IL-2 mutein that selectively activates and expands regulatory T cells as a therapy for autoimmune disease, *Journal of Autoimmunity* **95**, 1–14 (2018).
31. C. Ye, D. Brand, S. G. Zheng, Targeting IL-2: an unexpected effect in treating immunological diseases, *Signal Transduction and Targeted Therapy* **3**, 2 (2018).
32. J. P. Siegel, R. K. Puri, Interleukin-2 toxicity., *Journal of Clinical Oncology* **9**, 694–704 (1991).
33. A. Carr, S. Emery, A. Lloyd, J. Hoy, R. Garsia, M. French, G. Stewart, G. Fyfe, D. A. Cooper, Australian IL-2 Study Group, Outpatient Continuous Intravenous Interleukin-2 or Subcutaneous, Polyethylene Glycol-Modified Interleukin-2 in Human Immunodeficiency Virus-Infected Patients: A Randomized, Controlled, Multicenter Study, *The Journal of Infectious Diseases* **178**, 992–999 (1998).
34. D.-A. Silva, S. Yu, U. Y. Ulge, J. B. Spangler, K. M. Jude, C. Labão-Almeida, L. R. Ali, A. Quijano-Rubio, M. Ruterbusch, I. Leung, T. Biary, S. J. Crowley, E. Marcos, C. D. Walkey, B. D. Weitzner, F. Pardo-Avila, J. Castellanos, L. Carter, L. Stewart, S. R. Riddell, M. Pepper, G. J. L. Bernardes, M. Dougan, K. C. Garcia, D. Baker, De novo design of potent and selective mimics of IL-2 and IL-15, *Nature* **565**, 186–191 (2019).
35. A. E. Kelly-Welch, Interleukin-4 and Interleukin-13 Signaling Connections Maps, *Science* **300**, 1527–1528 (2003).
36. G. A. Efimov, A. A. Kruglov, Z. V. Khlopchatnikova, F. N. Rozov, V. V. Mokhonov, S. Rose-John, J. Scheller, S. Gordon, M. Stacey, M. S. Drutskaya, S. V. Tillib, S. A. Nedospasov, Cell-type–restricted anti-cytokine therapy: TNF inhibition from one pathogenic source, *Proceedings of the National Academy of Sciences* **113**, 3006–3011 (2016).
37. J. B. Spangler, I. Moraga, J. L. Mendoza, K. C. Garcia, Insights into Cytokine–Receptor Interactions from Cytokine Engineering, *Annual Review of Immunology* **33**, 139–167 (2015).
38. J. S. Peraino, H. Zhang, P. V. Rajasekera, M. Wei, J. C. Madsen, D. H. Sachs, C. A. Huang, Z. Wang, Diphtheria toxin-based bivalent human IL-2 fusion toxin with improved efficacy for targeting human CD25+ cells, *Journal of Immunological Methods* **405**, 57–66 (2014).

39. Z. Wang, Q. Zheng, H. Zhang, R. T. Bronson, J. C. Madsen, D. H. Sachs, C. A. Huang, Z. Wang, Ontak-like human IL-2 fusion toxin, *Journal of Immunological Methods* **448**, 51–58 (2017).
40. Z. Qi, Y. Qiu, Z. Wang, H. Zhang, L. Lu, Y. Liu, D. Mathes, E. A. Pomfret, D. Gao, S. Lu, Z. Wang, A novel diphtheria toxin-based bivalent human EGF fusion toxin for treatment of head and neck squamous cell carcinoma, *Molecular Oncology* **15**, 1054–1068 (2021).
41. J. Kossai, Y. Panagakis, A. Anandkumar, M. Pantic, *TensorLy: Tensor Learning in Python* (arXiv, 2018; <https://arxiv.org/abs/1610.09555>).
42. A. M. Farhat, A. C. Weiner, C. Posner, Z. S. Kim, B. Orcutt-Jahns, S. M. Carlson, A. S. Meyer, Modeling cell-specific dynamics and regulation of the common gamma chain cytokines, *Cell Reports* **35**, 109044 (2021).
43. T. Ishino, M. Wang, L. Mosyak, A. Tam, W. Duan, K. Svenson, A. Joyce, D. M. O’Hara, L. Lin, W. S. Somers, R. Kriz, Engineering a Monomeric Fc Domain Modality by N-Glycosylation for the Half-life Extension of Biotherapeutics, *Journal of Biological Chemistry* **288**, 16529–16537 (2013).

Acknowledgements:

Funding: This work was supported by a research agreement with Visterra, Inc.

Author contributions statement: A.S.M. and S.M.C. conceived of the study. C.P. and S.M.C. performed the PBMC experiments and engineered the IL-2 fusion proteins. A.S.M, B.O.J., E.M.S., and P.C.E. performed the computational analysis. All authors helped to design experiments and/or analyze the data. All authors contributed to writing the paper.

Competing interests: S.M.C. and C.P. are employees of Visterra Inc. A.S.M. has filed an invention disclosure on the use of multivalent cytokines to enhance cell type selective responses.

Data Sharing Plan: All analysis was implemented in Python v3.9 and can be found at <https://github.com/meyer-lab/gc-valent>, release 1.0, along with all experimental data.

Figures

Fig. 1. Possible effects of bivalent Fc-cytokine formats. (A) Multivalent ligands are known to bind to surface receptors with altered functional affinities that vary as a function of the number of receptors displayed by the cell. (B) Cartoon illustrating how ligand valency may alter signal transduction dynamics. Multivalent binding has been shown to be critical in some signaling pathways via the local aggregation of active signaling complexes. (C) Multivalently bound receptors have been shown to be degraded intracellularly at a rate differing from monovalently bound receptors and have also been shown to possess altered *in vivo* pharmacokinetics.

Fig. 2. pSTAT5 response varies in a cell type- and treatment-specific manner. (A) Schematic representation of non-negative canonical polyadic (CP) decomposition. Experimental pSTAT5 measurements are arranged in a tensor according to the duration of treatment, ligand used, cytokine concentration, and cell type. CP decomposition then helps to visualize this space. **(B)** Percent variance reconstructed (R2X) versus the number of components used. **(C)** Component values for each IL-2 form. **(D)** Component values representing the effect of IL-2 concentration. **(E)** Component values representing cell type specificity. **(F)** Component values for the effect of treatment duration.

Fig. 3. Responses are predicted by a simple multivalent binding model. All accuracies are calculated as a Pearson's correlation R^2 score for experimental cytokine responses at the 30 minute and one hour time points. **(A)** A simplified cartoon of the model. Initial association of multivalent ligands proceeds according to monovalent affinity, and subsequent binding events proceed with that affinity scaled by the K_x^* parameter. **(B)** Experimentally measured vs. predicted pSTAT5 responses for all cell types, ligands, and concentrations. Each point represents a single experimental measurement ($N=1$). **(C,D)** Model's accuracy subset by cell type **(C)** and ligand **(D)** for all mono- and bivalent IL-2 muteins. **(E)** Model's accuracy when predicting the responses of all cell types to bivalent ligands either correctly as dimers, or as IL-2 monomers. **(F,G)** Model's accuracy subset by concentration **(F)** for all ligands and time **(G)** for all ligands, concentrations, and cell types.

Fig. 4. Multivalency with coordinate affinity adjustments can enhance selectivity. Affinities were allowed to vary across K_D s of 10 pM–1 μ M while K_x^* was fixed at its fitting optimum. All optimizations were performed using a concentration of 1 nM. Selectivity was calculated as the ratio of predicted pSTAT5 in target cells to the mean pSTAT5 predicted in off-target cells. **(A,C,E)** Signaling response of T_{reg} , NK cells, and T_{helper} cells predicted for ligand of optimal selectivity at different valencies. Response predictions were normalized to each population's response for the monovalent case. Selectivity for T_{regs} and NK cells were derived from IL-2 muteins, and selectivity for $T_{helpers}$ was calculated using IL-7 muteins. **(B,D,F)** Optimal receptor-ligand dissociation constants for each ligand optimized for selectivity. Mutein affinity for IL-2R α and IL-2R β/γ_c was allowed to vary for IL-2 muteins, and affinity for IL-7R α was allowed to vary for IL-7 muteins.