A PKA Inhibitor Motif within Smoothed Controls Hedgehog Signal Transduction

John T. Happ^{1,8}, Corvin D. Arveseth^{1,8}, Jessica Bruystens^{2,8}, Daniela Bertinetti^{3,8}, Isaac B. Nelson^{1,8}, Cristina Olivieri⁴, Danielle S. Hedeem¹, Ju-Fen Zhu¹, Jacob L. Capener¹, Jan Wilfried Bröckel³, Lily Vu⁵, C.C. King^{2,7}, Victor L. Ruiz-Perez⁶, Gianluigi Veglia⁴, Friedrich W. Herberg³, Susan S. Taylor^{2,7,*}, Benjamin R. Myers^{1,*}

¹Department of Oncological Sciences, Department of Biochemistry, and Department of Bioengineering, University of Utah School of Medicine, Salt Lake City, UT, USA

²Department of Pharmacology, University of California, San Diego, La Jolla, CA, USA

³Institute for Biology, Department of Biochemistry, University of Kassel, Kassel, Germany

⁴Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, Minneapolis, MN, USA

⁵Department of Neurobiology, University of California, San Diego, La Jolla, CA, USA

⁶Instituto de Investigaciones Biomédicas “Alberto Sols,” Consejo Superior de Investigaciones Científicas (CSIC)—Universidad Autónoma de Madrid (UAM), and CIBER de Enfermedades Raras (CIBERER), Instituto de Salud Carlos III (ISCIII), Madrid, Spain

⁷Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA, USA

⁸These authors contributed equally to this work

*Correspondence: staylor@ucsd.edu or benjamin.myers@hci.utah.edu

ABSTRACT

The Hedgehog (Hh) cascade is central to development, tissue homeostasis, and cancer. A pivotal step in Hh signal transduction is the activation of GLI transcription factors by the atypical G protein-coupled receptor (GPCR) Smoothened (SMO). How SMO activates GLI has remained unclear for decades. Here we show that SMO employs a decoy substrate sequence to physically block the active site of the PKA catalytic subunit (PKA-C) and extinguish its enzymatic activity. As a result, GLI is released from phosphorylation-induced inhibition. Using a combination of *in vitro*, cellular, and organismal models, we demonstrate that interfering with SMO / PKA pseudosubstrate interactions prevents Hh signal transduction. The mechanism we uncovered echoes one utilized by the Wnt cascade, revealing an unexpected similarity in how these two essential developmental and cancer pathways signal intracellularly. More broadly, our findings define a new mode of GPCR-PKA communication that may be harnessed by a range of membrane receptors and kinases.

INTRODUCTION

The Hh signaling cascade is fundamental to embryogenesis, controlling the development of nearly every vertebrate organ¹⁻⁴. Insufficient Hh pathway activity underlies birth defects affecting the nervous, cardiovascular, and musculoskeletal systems⁵⁻⁷. On the other hand, Hh pathway overactivation drives several common cancers, including basal cell carcinoma of the skin (the most common cancer in North America) and medulloblastoma (the most common pediatric brain tumor)^{8,9}. The Hh pathway utilizes an unusual signal transduction mechanism involving layers of repressive interactions¹⁻⁴. In the pathway “off” state, PKA-C phosphorylates GLI, stimulating its proteolysis into a truncated transcriptional repressor that inhibits target gene expression^{4,10}. In the pathway “on” state, Hh ligands bind to and inactivate the 12-transmembrane sterol transporter PATCHED1 (PTCH1), which releases SMO from PTCH1-mediated inhibition⁴. This process allows SMO to access its endogenous sterol ligands and undergo an activating conformational change¹¹⁻¹⁴. Once activated, SMO blocks PKA-C-mediated phosphorylation of GLI¹⁵⁻¹⁸, a key step that likely occurs within the tiny cell-surface compartment formed by the primary cilium¹⁹⁻²¹. Consequently, GLI is activated and can control the expression of proliferative or differentiative genes¹⁰. SMO regulation of PKA-C is thus a critical event in transducing Hh signals from the cell surface to the nucleus. However, the underlying mechanism has remained mysterious for decades¹⁻⁴.

GPCRs classically inhibit PKA-C via well-characterized signaling cascades involving heterotrimeric G protein-mediated effects on cAMP which promote formation of inactive PKA holoenzymes^{22,23}. In contrast, we recently found that SMO prevents PKA-C from phosphorylating substrates via a noncanonical mechanism. SMO directly interacts with PKA-C subunits, recruiting them to the membrane and thereby restricting their access to soluble GLI proteins. SMO / PKA-C interactions are triggered by GPCR kinases 2 and 3 (GRK2/3), which recognize the SMO active state and phosphorylate the cytoplasmic tail (C-tail) of SMO to promote PKA-C binding. Based on these observations, we proposed that active, phosphorylated SMO binds to and sequesters PKA-C within the cilium, which prevents phosphorylation of GLI and thereby promotes GLI activation²⁴.

Here we uncover a critical and unexpected component of the SMO / PKA-C regulatory mechanism in which SMO physically blocks PKA-C enzymatic activity. We show that the SMO proximal C-tail (pCT) acts as a decoy PKA-C substrate that binds to and occludes the kinase active site, thereby preventing phosphorylation of PKA-C targets. SMO is, to our knowledge, the first example of a GPCR that functions as a direct PKA-C inhibitor. However, this decoy substrate mechanism appears to apply more generally to transmembrane receptors and kinases in other signaling pathways.

RESULTS

SMO binds and inhibits PKA-C as a pseudosubstrate

The principal regulators of PKA-C activity within cells are PKA regulatory (PKA-R) subunits²⁵ and the heat-stable protein kinase inhibitor (PKI) proteins²⁶. PKI proteins and type I (PKA-RI)

regulatory subunits operate as pseudosubstrates that bind within the PKA-C active site but cannot undergo phosphorylation. As a result, the enzyme's phosphoryl transfer and substrate turnover cycle is interrupted²⁵⁻²⁸. PKA-R / PKA-C holoenzymes dissociate upon cAMP binding to PKA-R, releasing catalytically active PKA-C²⁵. In contrast, PKI proteins interact with PKA-C independently of cAMP levels²⁶. Despite their divergent regulatory influences, PKI and PKA-RI engage the PKA-C active site cleft using similar sequence motifs^{25,28}.

Inspection of the mouse SMO sequence revealed a region within the pCT (residues 615-638) that bears the hallmarks of a PKA-C pseudosubstrate (Fig. 1a). First, in contrast to canonical PKA-C substrates, which contain a serine or threonine at a canonical phosphorylation site (P-site)²⁹, SMO possesses a non-phosphorylatable residue, alanine, at this position. Second, SMO contains arginines at the P-2 and P-3 positions that are essential in other pseudosubstrates (PKI and PKA-RI) for binding the PKA-C active site³⁰⁻³³. Third, SMO contains a hydrophobic residue, isoleucine, at P+1 and a bulkier aromatic residue, tryptophan, at P-13, both of which contribute to high-affinity PKI interactions with PKA-C³⁴⁻³⁶. Finally, SMO harbors a predicted α -helical sequence N-terminal to the pseudosubstrate region (Fig. 1a), resembling a domain in PKI that is required for high-affinity interactions with PKA-C^{37,38}. Based on these observations and the studies described below, we refer to SMO residues 615-638 as the "SMO PKI motif".

We hypothesized that SMO utilizes its PKI motif as a pseudosubstrate to bind and inhibit PKA-C, thereby activating GLI. Consistent with this hypothesis, residues 615-637 of mouse SMO are essential for communication with GLI³⁹, although the structure, function, and interacting partner(s) of this SMO region are all undefined. Furthermore, the P, P+1, P-2, P-3, and P-13 residues critical for PKA-C pseudosubstrate function in *bona fide* PKI proteins are highly conserved among SMO orthologs (Extended Data Fig. 1), suggestive of a central role in SMO functionality.

We measured the affinity of the SMO PKI motif for PKA-C α , the best-studied and most ubiquitously expressed PKA-C isoform²⁷, using fluorescence polarization assays. A fluorescently labeled peptide encompassing the SMO PKI motif (standard SMO peptide, Fig. 1a, red) bound saturably to purified human (K_D = 823 nM, Fig. 1b (top)) or mouse (K_D = 911 nM, Extended Data Fig. 2a (left)) PKA-C α . These interactions showed pseudosubstrate characteristics: they were strongly ATP-dependent⁴⁰ (EC_{50} for ATP = 19.2 μ M (human) or 0.59 μ M (mouse), Fig. 1b (bottom) and Extended Data Fig. 2a (right)) and were blocked by alanine substitution of the P-2 / P-3 arginines and the P-13 tryptophan, hereafter referred to as the "WRR mutation" (Fig. 1c, Extended Data Fig. 2b). Along similar lines, direct surface plasmon resonance (SPR) binding studies revealed that a recombinant protein encompassing the entire 93 amino-acid SMO pCT bound specifically to PKA-C α (Fig. 1d). These interactions displayed fast on- and off-rates with transient kinetics, and steady-state analysis of the binding data revealed a K_D of 835 nM (Extended Data Fig. 2c,d). Consistent with our peptide binding studies, PKA-C interactions with the SMO pCT depended strictly on ATP and $MgCl_2$ (Extended Data Fig. 2e), and only minimal PKA-C binding was observed for the WRR mutant (Fig. 1e, Extended Data Fig. 2c,d). Interestingly, the above SMO PKI binding assays revealed weaker interactions with PKA-C than observed for conventional pseudosubstrates (K_D = 1.1 nM for PKI α (5-24) peptide⁴¹). We discuss the biochemical basis and biological significance for these affinity differences below.

We next studied the interaction between PKA-C and the SMO PKI motif at a structural level using nuclear magnetic resonance (NMR) spectroscopy. Binding of pseudosubstrate sequences from either SMO or PKI α induced similar overall chemical shift perturbations throughout the PKA-C kinase core (Fig. 2a, Extended Data Fig. 2f). SMO PKI peptides also blocked PKA-C enzymatic activity *in vitro*, as assessed via a PKA-C substrate phosphorylation assay⁴², with an extended length peptide (Fig. 1a) inhibiting PKA-C more profoundly (Fig. 2b). In keeping with this trend, the recombinant SMO pCT (Extended Data Fig. 2g,h) required 5-10-fold lower concentrations to efficiently block PKA-C activity compared to SMO PKI peptides (24-38 amino acids in length) (IC₅₀ = 10.9 μ M for pCT vs. 50-125 μ M for SMO PKI peptides, Fig. 2c). Consistent with a pseudosubstrate mode of inhibition, introducing the WRR mutation into the SMO pCT restored PKA-C activity (Fig. 2d). These data indicate that the SMO PKI motif is sufficient to inhibit PKA-C substrate phosphorylation, with additional sequences in the pCT enhancing the efficiency of inhibition.

The SMO PKI motif is required for Hh signal transduction

We next asked whether the SMO PKI motif contributes to PKA-C binding and inhibition in living systems. Our initial studies involved a simplified HEK293 cell model for SMO / PKA-C regulation and employed truncated SMO constructs (either SMO657 or SMO674, with the number indicating the C-terminal-most residue) that exhibit improved expression and biochemical stability compared to full-length SMO²⁴. We used bioluminescence resonance energy transfer (BRET)^{43,44} to detect interactions between nanoluciferase (nanoluc)-tagged SMO and YFP-tagged PKA-C constructs in a cellular environment²⁴. In this assay, PKA-C exhibited strong, specific BRET with wild-type SMO657, but not with SMO657 harboring the WRR mutation (Fig. 3a). Coimmunoprecipitation assays confirmed these results (Extended data Fig. 3a). Similarly, the WRR mutation prevented SMO674 from colocalizing with PKA-C at the membrane²⁴, as determined by live-cell confocal microscopy (Fig. 3b).

We next assessed the impact of the SMO PKI motif on PKA-C activity in cells using a model PKA-C substrate, the cyclic AMP response element binding protein (CREB) transcription factor⁴⁵, and a CREB transcriptional reporter assay. This experimental paradigm entails overexpression of PKA-C at levels exceeding those of endogenous PKA-R, thereby minimizing potentially confounding contributions from heterotrimeric G protein- and cAMP-containing cascades²⁴. Under these conditions, wild type SMO674 blocked PKA-C-mediated CREB reporter activation while SMO674 harboring the WRR mutation failed to do so (Fig. 3c). We observed similar effects with serine substitution of the P-site alanine (A635S, Fig. 3c and Extended data Fig. 3a), which converts pseudosubstrates to substrates and thereby promotes their dissociation from the PKA-C active site^{33,46}.

We extended the above findings to more physiological cellular models of Hh signaling in primary cilia, using SMO constructs with full-length unmodified C-termini. We studied SMO / PKA-C colocalization in cilia using inner medullary collecting duct (IMCD3) cells, a robustly ciliated kidney cell line that is used extensively in studies of ciliary Hh signal transduction^{24,47-49}. Upon Hh

pathway activation, SMO accumulates in IMCD3 primary cilia⁴⁷⁻⁴⁹, where it colocalizes with PKA-C²⁴. In contrast, PKA-C displayed dramatically reduced colocalization in cilia with SMO containing the WRR mutation (Extended Data Fig. 3b). The WRR mutation also prevented interaction of SMO with PKA-C (expressed at near-endogenous levels²⁴) in these ciliated cells (Fig. 3d).

To capture the entire process of Hh signal transduction from the cell surface to the nucleus, we used mouse embryonic fibroblasts (MEFs) and a GLI transcriptional reporter assay^{11,50,51}. This model strictly requires SMO, PKA-C, GLI, and an intact cilium^{50,52,53}. In *Smo*^{-/-} MEFs, transfection of wild-type SMO enabled strong GLI transcriptional responses to Hh ligands, whereas the WRR and A635S mutants were almost completely devoid of activity (Fig. 3e).

Mutations in the SMO PKI motif specifically affect binding to and inhibition of PKA-C, based on the following control studies. First, the expression levels and electrophoretic mobilities of these mutants were similar to those of their wild-type counterparts (Extended Data Fig. 3a and 4a,b,d), and the mutants localized correctly to primary cilia (Extended Data Fig. 3b, 4c). In addition, the SMO mutations only minimally affected BRET with nanobody 2 (NbSmo2, Fig. 3a), which selectively binds the active state of SMO via the cytoplasmic face of its seven-transmembrane (7TM) domain²⁴. Finally, the mutants underwent normal SMO activity-dependent phosphorylation by GRK2/3 kinases^{24,54} (Extended Data Fig. 4d). These data rule out possible nonspecific effects of the mutations on SMO expression, trafficking, conformational activation, or ability to serve as a GRK2/3 substrate.

To examine the role of the SMO PKI motif in Hh signal transduction *in vivo*, we studied the specification of slow muscle cell types in zebrafish embryos, a widely utilized model of morphogenetic Hh signal transduction during vertebrate development^{24,55,56}. While expression of wild-type SMO restored correct muscle specification to *smo*^{-/-} zebrafish, the WRR or A635S mutants did not (Fig. 3f, Extended Data Table 2).

Taken together, our studies demonstrate a requirement for PKA-C pseudosubstrate interactions in SMO inhibition of PKA-C, and activation of GLI, during Hh signal transduction in both cultured cells and organisms.

An avidity-based mechanism for SMO inhibition of PKA-C

PKA-C interactions with the SMO PKI motif, while essential for Hh signal transduction (Fig. 3), appear weaker ($K_D = 823\text{-}911$ nM for SMO PKI peptides, Fig. 1b, Extended Data Fig. 2a,d) than those with a canonical PKA-C pseudosubstrate, ($K_D = 1.1$ nM for PKI α (5-24) peptide⁴¹). SMO / PKA-C interactions, however, are also influenced by sequences outside the PKI motif, indicating that they are more favorable *in vivo* than the *in vitro* measurements suggest. Several lines of evidence support this proposal. First, SMO mutations known to reduce PKA-C interactions *in vivo*, including deletion of a predicted amphipathic helix spanning residues 570-581 or alanine substitution of GRK2/3 phosphorylation sites²⁴, map to non-PKI regions of the pCT (summarized in Fig. 4a). Second, SMO pCT interactions with PKA-C are enhanced by the distal SMO C-tail (dCT, residues 658-793)²⁴, demonstrating that this region harbors additional PKA-C binding

determinants. Accordingly, in HEK293 cells, PKA-C interactions with SMO containing its full-length C-tail are only partially blocked by the WRR mutation (Fig. 4b), in contrast to the near-complete loss of interaction with SMO lacking the dCT (Fig. 3a). Finally, a tiled SMO C-tail peptide microarray identified several PKA-C-binding sequences in addition to the PKI motif, including one in the dCT with PKI-like attributes (Fig. 4c, Extended Data Fig. 5).

Taken together, our data support an avidity-based mechanism for SMO / PKA-C binding. A set of ancillary interactions cooperate to stabilize the core binding events between the SMO PKI motif and the PKA-C active site. This avidity-based mechanism may provide the essential link between SMO activation and disruption of PKA-C phosphoryl transfer during Hh signal transduction.

DISCUSSION

SMO inhibition of PKA-C is a central aspect of Hh signal transduction in development and disease. Instead of obeying the existing paradigms for GPCR-PKA signaling, SMO enjoys a private signaling pathway whereby it directly binds PKA-C as a pseudosubstrate. This both restricts access of PKA-C to soluble targets and extinguishes its enzymatic activity. A PKI motif encoded by the SMO cytoplasmic domain is central to this mechanism in two ways. First, the PKI motif sequesters PKA-C away from GLI by promoting SMO / PKA-C interactions, and thus recruitment of PKA-C to the membrane. Second, the PKI motif occludes the PKA-C active site, interrupting catalysis. By utilizing this two-pronged strategy, SMO can efficiently block PKA-C-mediated phosphorylation, activate GLI, and promote transcription of Hh pathway target genes.

Based on our findings, we propose a revised model for Hh signal transduction downstream of SMO (Fig. 4d). In the pathway “off” state, SMO is in an inactive conformation^{11,51,57,58}. As a result, SMO cannot be phosphorylated efficiently by GRK2/3^{24,54}, and exists at low levels in the ciliary membrane⁵⁹⁻⁶¹. We expect interactions between PKA-C and the SMO PKI motif to be unfavorable under these conditions due to their distinct subcellular localizations and relatively weak affinity. In the pathway “on” state, Hh-mediated inactivation of PTCH1 enables SMO to bind its endogenous sterol ligands and adopt an active conformation^{11-13,51}. Consequently, SMO undergoes phosphorylation by GRK2/3 and accumulates in the cilium, bringing SMO in close proximity to ciliary PKA-C. This allows SMO to bind PKA-C via its PKI motif, which both sequesters PKA-C at the ciliary membrane²⁴ and blocks PKA-C-mediated catalysis. These interactions are facilitated by other PKA-C-binding sequences in the SMO pCT and dCT. We expect the interactions are also enhanced by the membrane itself, which restricts diffusion of proteins to two dimensions and can thereby dramatically increase effective concentrations relative to soluble compartments⁶². Additionally, SMO activation could conceivably decrease ciliary cAMP via coupling to inhibitory ($G_{\alpha_{i/o/z}}$) G protein cascades^{51,63,64} or by triggering the ciliary exit of GPR161, a constitutively active GPCR that couples to stimulatory (G_{α_s}) G proteins^{47,49,65}. These processes, while not absolutely required for Hh signal transduction⁶⁵⁻⁷⁰, may reduce levels of free (non-PKA-R-bound) PKA-C in cilia⁷¹, and thereby aid the SMO PKI motif in binding and inhibiting the ciliary pool of PKA-C. Thus, activation of SMO prevents PKA-C substrate phosphorylation via a host of auxiliary interactions and regulatory influences that assist the essential pseudosubstrate action of the SMO PKI motif.

Our study helps explain a crucial but enigmatic aspect of Hh signal transduction, namely how PKA-C is inhibited, and GLI activated, only when the Hh pathway is in an “on” state. Indeed, our findings are consistent with a number of prior observations, such as that inactivation of GRK2/3 decreases SMO / PKA-C interactions²⁴, blocks SMO / PKA-C colocalization at the membrane²⁴, and abolishes Hh signal transduction^{56,72,73}. Based on our model, these effects can be explained by GRK2/3 inhibition weakening the SMO / PKA-C interaction below a critical threshold such that the PKI motif cannot efficiently engage the enzyme’s active site. Furthermore, our findings suggest that SMO / PKA-C interactions are appropriately tuned: If the SMO PKI motif were to bind PKA-C with high affinity, the Hh pathway could misfire, as PKA-C would be inhibited (and GLI dephosphorylated) even when SMO is inactive. Conversely, reducing the affinity of PKA-C for the SMO PKI motif, as occurs with the SMO WRR or A635S mutations, prevents the SMO active state from inhibiting PKA-C and thereby hinders GLI transcription. Future structural studies of full-length SMO / PKA-C complexes will provide a more detailed understanding of how the SMO pCT engages PKA-C, and may also reveal precisely how GRK2/3 facilitates these interactions.

The mechanism described above represents, to our knowledge, an unprecedented mode of GPCR-kinase communication. It is, however, remarkably similar to a strategy used by another critical developmental and cancer pathway: the Wnt cascade⁷⁴⁻⁷⁷ (Extended Data Fig. 6). During the initial steps of Wnt signal transduction at the plasma membrane, the cytoplasmic tail of LDL receptor-related proteins 5/6 (LRP5/6) serves as a pseudosubstrate inhibitor of GSK-3 β kinases. This blocks phosphorylation and degradation of the transcriptional coactivator β -catenin⁷⁸⁻⁸¹. Moreover, the LRP5/6 tail binds GSK-3 β with low affinity, but Wnt pathway activation enhances this interaction through phosphorylation of the LRP5/6 tail as well as recruitment of GSK-3 β -containing protein complexes to the membrane⁷⁸⁻⁸⁰. Thus, in both the Hh and Wnt cascades, a phosphorylated receptor tail sequesters and competitively inhibits a kinase via an avidity-based mechanism. It was unexpected that these two pathways, which utilize distinct molecular components and rely on disparate subcellular environments (cilium vs. plasma membrane)^{21,82}, nevertheless share a common mechanism for intracellular signal transduction. Why such a mechanism evolved to carry out central steps in these pathways is presently unclear. One possibility is that it serves to spatially restrict the kinase-inhibiting effects of upstream receptors and thereby avoid pleiotropic effects of inhibiting these kinases globally. This may help to ensure signaling specificity and reduce untoward crosstalk with other pathways^{75,78,81}. It is also possible that such a mechanism is particularly well-suited to the embryogenesis- and homeostasis-related functions of Hh and Wnt signaling. In any event, the use of a similar strategy during Hh and Wnt signal transduction hints at a more general applicability to a range of transmembrane receptors and kinases. Given the ubiquity of these proteins in metazoan physiology and disease^{22,23,28}, and their prevalence as therapeutic targets^{83,84}, this represents an exciting area for future study.

METHODS:

Cell culture and zebrafish husbandry. HEK293FT cells, IMCD3 Flp-in cells, *Smo*^{-/-} MEFs, and HEK293-Freestyle cells were grown as previously described²⁴. Stably transfected HEK293 Flp-in T-rex cells were constructed and maintained as previously described¹¹. IMCD3 stable line

expressing SMO(WRR)-IRES-PKACmNG produced as previously described, others (PKAC-mNG and β 2ARNb80-mNG) already described. Zebrafish were maintained as previously described²⁴.

Antibodies, small molecules, and other reagents. SAG21k was a gift from P. Beachy. Cmpd101 was obtained from Hello Bio. Rabbit anti-GFP (which also detects YFP) was obtained from Thermo Fisher Scientific (A11122). Mouse anti-Arl13b was obtained from Antibodies Inc (75-287). Alexa 647-conjugated M1 FLAG antibody and M1 FLAG affinity resin were prepared in-house. Control or ShhN conditioned medium was prepared from stably transfected HEK293 cells as previously described²⁴. Dual-luciferase assay was obtained from Promega. Coelenterazine h was obtained from NanoLight Technology (301–500). Furimazine was obtained from AOBIOUS (AOB36539). For transfection studies, TransIT 2020 and TransIT 293 were obtained from Mirus Bio.

DNA constructs. For expression and purification of the SMO pCT in *E. coli*, the ShhN gene was excised from ShhN / pHTSHP⁵¹ and replaced with residues 565-657 of mouse SMO (⁵⁶⁵KRIKK...⁶⁵⁷PEEQAN⁶⁵⁷), downstream of the MBP-His₈-SUMO tags in the vector pHTSHP, or for GST-constructs in the vector pGexKG. GST-PKI construct in pGexKG was previously described⁸⁵. The resulting construct was termed SMO pCT(565-657). The SMO655-nanoluc, SMO674, and SMO-nanoluc constructs (all in pVLAD6, with N-terminal hemagglutinin signal sequence and FLAG tag) were previously described²⁴. Full-length SMO (in pGEN, with C-terminal myc tag) was previously described⁸⁶. Mouse PKA-C α -YFP, PKA-Rl α -YFP, and NbSmo2-YFP constructs in pVLAD6 were previously described²⁴. Barrestin1-YFP in pCDNA3.1-zeo was previously described²⁴. To construct IMCD3 Flp-in stable lines coexpressing FLAG-tagged mutant SMO with mNeonGreen-tagged PKA-C α (using an IRES element), the previously described construct SMO-IRES-PKAC α mNG / pEF5-FRT²⁴ was modified to introduce the WRR mutation into SMO. To construct stable Flp-in HEK293 cell lines coexpressing wild-type or mutant SMO674 (which includes an N-terminal hemagglutinin signal sequence and FLAG tag) along with GFP-tagged PKAC α , we cloned a SMO674-IRES-PKAC α GFP cassette into the pCDNA5-FRT-TO vector. Untagged human and mouse PKAC α constructs in the vector pRSETb were previously described⁸⁵. All mutant DNA sequences were prepared via PCR-based mutagenesis, cloned into their respective vector backbones via Gibson assembly, and verified by Sanger sequencing.

Peptide synthesis. SMO PKI peptides (standard = SADVSSAWAQHVTKMVARRGAILP; extended = SADVSSAWAQHVTKMVARRGAILPQDVSVPVATPVPP) were prepared via standard solid-phase synthesis by either GenScript or Elim Bio, purified via reversed-phase HPLC to $\geq 85\%$ purity, and the sequence / molecular mass confirmed by mass spectrometry. For fluorescence polarization assays, a FAM fluorophore and 3xPEG linker were added to the N-terminus of the standard SMO PKI peptide during synthesis. PKI α (5-24) Peptide (TTYADFIASGRTGRRNAIHD) was synthesized as described for the SMO-PKI peptides by GeneCust (Boynes, France).

Fluorescence polarization assays. Human or mouse PKA-C α subunits were purified as previously described⁸⁷. In addition, the protein obtained from the S200 was further purified via cation exchange chromatography. For this, the S200 peak fractions were pooled and dialyzed

overnight into MonoS buffer (20 mM KH_2PO_4 pH 6.5, 5 mM DTT) before loading onto a MonoS cation exchange column. PKA-C α was eluted with a gradient of 0-300mM KCl in MonoS buffer. Binding of a SMO peptide containing the PKI motif was investigated by fluorescence polarization (FP). Assay buffer for all FP experiments consisted of 50 mM MOPS pH 7, 35 mM NaCl, 10 mM MgCl_2 , 1 mM DTT and 0.005% Triton X-100 with the addition of 1 mM ATP in the PKA-C titrations. Experiments were performed by adding 50 μl of the titration component (PKA-C or ATP) to 150 μl of FAM-SMO containing solution. For SMO/PKA-C binding experiments, a two-fold dilution series from 16 μM to 0 μM of either mouse or human PKA-C α was added to 40 nM FAM-SMO. To assess the ATP dependence of SMO binding to PKA-C, a titration with varying the ATP concentration from 0 to 1 mM (human PKA-C α) or 12.8 μM (mouse PKA-C α) was used with keeping PKA-C at 3 μM and FAM-SMO at 25 nM. Readings were taken with a Tecan Genios plate reader using black flat bottom 96-well Costar plates. Each experiment was carried out in at least triplicate with FAM readings at 485 nm excitation and 535 nm emission.

Peptide arrays. Peptide arrays were synthesized with a MultiPep Flexible Parallel Peptide Synthesizer (Intavis Bioanalytical Instruments, Germany). The blots were probed with mouse or human PKA C α protein, purified as described above for “Fluorescence polarization assays”. Unless noted otherwise all steps were carried out at room temperature. The blots were initially soaked with 100% ethanol for 5 minutes followed by 5 times 5-minute washes with water. All subsequent washes were carried out for 5 minutes 5 times with TTBS. The membranes were washed with TTBS, blocked with 5% milk in TTBS for 1 hour and washed again with TTBS. They were then soaked with TTBS containing 2 μg PKA-C α , 5% milk, 10 mM MgCl_2 and 1mM ATP overnight at 4° C. The next day the blots were washed, blocked with 5% milk in TTBS and washed again before incubation overnight at 4° C with TTBS containing 5% milk and a primary PKA-C antibody (generated in-house previously). The next day the blots were washed, then incubated with an HRP-conjugated anti-rabbit antibody (PrometheusTM) in 5% milk TTBS for one hour followed by a final round of washes with TTBS. The membranes were finally covered with SuperSignal West Pico PLUS Chemiluminescent Substrate for detection of HRP (Thermo Scientific # 34580) and imaged with a ChemiDoc MP Imaging System from BIO RAD. A representative array image (representative of two separate trials) is shown; we excluded from analysis any spots that were not observed consistently across replicates.

Surface plasmon resonance (SPR). Human PKA-C α was overexpressed in *E. coli* BL21(DE3) cells after induction with 0.4 mM IPTG for 16 h at RT using the expression vector pRSETb-hPKAC α and then purified by affinity chromatography using an IP20-resin as described earlier⁸⁸. GST-PKI, GST-SMO pCT wt and WRR mutant were overexpressed in *E. coli* BL21(DE3) for 16 hr at room temperature, and the GST fusion proteins were purified using Protino glutathione agarose 4B according to the manufacturer's instruction (Macherey-Nagel). SPR interaction studies were performed according to previous studies^{46,85}. Briefly, the interaction studies were performed in running buffer (20 mM MOPS, pH 7.0, 150 mM NaCl, 50 μM EDTA, 0.005% P20 surfactant) at 25 °C using a Biacore 3000 instrument (GE Healthcare). For measurements involving ATP/ MgCl_2 , the buffer was supplemented with 1 mM ATP and 10 mM MgCl_2 . Polyclonal anti-GST antibody (Carl Roth, 3998.1) was covalently immobilized to all four flow cells of a CM5

sensorchip (GE Healthcare) to a level of 8,000 response units (RU) via standard NHS/EDC amine coupling. Each measurement cycle started with the sequential capture of 60–130 RU of GST-PKI and 500–1,000 RU of GST-SMO-RLG wt and GST-SMO-RLG WRR mutant on separate flow cell (flow rate 10 μ L/min). Interaction analysis was then initiated by the injection of increasing concentrations of human PKA C α (156 nM – 10 μ M) at a flow rate of 30 μ L/min for 120 or 240 s (association) followed by 120 or 240 s dissociation with buffer without PKA C α . Nonspecific binding was removed by subtracting SPR signals from a blank flow cell (without GST-protein) and additional buffer blank runs without PKA C α (double referencing) employing BIAevaluation Software 4.1.1 (Cytiva, Marlborough, MA, USA). After each cycle, the sensorchip was regenerated by injecting up to 5 times with 10 mM glycine, pH 1.9 or 2.2, to remove the GST-fusion proteins from the antibody surfaces until the baseline level was reached. Steady-state analysis was performed with GraphPadPrism with a one-site binding (hyperbola) model as previously described⁸⁵.

NMR spectroscopy. Recombinant human PKA-C α (PRKACA, uniprot P17612) was expressed and purified as reported^{89,90}. Briefly, transformed *E.coli* BL21 (DE3) pLysS (Agilent) cells were cultured in deuterated (²H) M9 minimal medium supplemented with ¹⁵NH₄Cl. Protein overexpression was initiated by adding 0.4 mM of isopropyl β -D-1-thiogalactopyranoside (IPTG) and carried out for 12 hours at 20 °C. The collected cell pellet was then resuspended in 50 mM Tris-HCl pH 8.0, 30 mM KH₂PO₄, 200 mM NaCl, 5 mM β -mercaptoethanol, 0.15 mg/ mL lysozyme, 200 μ M ATP, DnaseI, and protease inhibitor (Sigma) and pass through a French press (2 times). The cell resuspension was then cleared by centrifugation (18,000 rpm for 45 minutes), and the supernatant was incubated overnight with Ni²⁺-NTA resin (Thermo Fisher). His-tagged PKA-C α was eluted using 50 mM Tris-HCl pH 8.0, 30 mM KH₂PO₄, 100 mM NaCl, 5 mM β -mercaptoethanol, 1 mM phenylmethylsulfonyl fluoride (PMSF) supplemented with 200 mM of imidazole. The His-tag was removed during an overnight dialysis step performed in 20 mM KH₂PO₄ (pH 6.5), 25 mM KCl, 5 mM β -mercaptoethanol, 0.1 mM PMSF, using stoichiometric quantities of recombinant tobacco etch (TEV) protease. Finally, cationic exchange chromatography was performed to separate the three different isoforms of PKA-C α , representing the three different phosphorylation states of the kinase, using a linear gradient of KCl in 20 mM KH₂PO₄ at a pH of 6.5 (HiTrap Q-SP column, GE Healthcare Life Science). All the NMR experiments were performed using PKA-C α isoform II, corresponding to phosphorylation at S10, T197, and S338 residues⁹¹. The purity of the protein preparation was tested using SDS-PAGE electrophoresis and Mass spectrometry analysis (purity >97%).

A portion of the lyophilized extended SMO PKI peptide (see above) was resuspended in 20 mM KH₂PO₄ (pH 6.5), 90 mM KCl, 10 mM MgCl₂, 10 mM DTT, 1 mM NaN₃, to a final concentration of 4 mM. The NMR experiments were performed on 100 μ M uniformly ²H, ¹⁵N-labeled PKA-C α sample in 20 mM KH₂PO₄, 90 mM KCl, 10 mM MgCl₂, 10 mM DTT, 1 mM NaN₃ at pH of 6.5 and saturated with 12 mM of a non-hydrolyzable ATP analog (ATP γ N). Modified [¹H-¹⁵N]-TROSY-HSQC spectra were acquired on a Bruker Advance III spectrometer operating at a proton frequency of 850 MHz, equipped with a TCI cryoprobe, at an acquisition temperature of 300K. First, a spectrum of ATP γ N-saturated PKA-C α complex (PKA-C α /ATP γ N) was recorded with 2048

(proton) and 128 (nitrogen) complex points. Stoichiometric amounts of SMO PKI peptide (hereafter referred to as SMO-PKI) were then added to the PKA-C α /ATP γ N complex until saturation (1:0, 1:1, 1:2, and 1:4 SMO-PKI:PKA-C/ATP γ N molar ratio), with a concentration of SMO-PKI ranging from 0.1 to 0.4 mM. All the spectra were processed using NMRPipe⁹², and visualized using NMRFAM-SPARKY software⁹³. Combined chemical shift perturbation (CSPs) were calculated using the ¹H and the ¹⁵N chemical shift derived from the PKA-C α /ATP γ N complex and the 1:4 SMO-PKI:PKA-C/ATP γ N complex. The CSPs was calculated using the following equation⁹⁴:

$$\Delta\delta = \delta_{SMO} - \delta_{ATP\gamma N} = \sqrt{(\Delta\delta H)^2 + (0.154 \times \Delta\delta N)^2}$$

The PKI α (5-24) CSPs were originally published earlier⁹⁰ and are included here for reference.

Purification of SMO pCT for PKA-C activity assays. MBP-His₈-SUMO-tagged SMO pCT (SMO pCT(565-657) / pHTSHP) was transformed into BL21 (DE3) *E. coli* and grown to OD₆₀₀ = ~0.5-1.0 in 1 L terrific broth (TB) with ampicillin at 37°C in 2800ml baffled Fernbach flasks until. IPTG was then added to 0.4mM and the temperature lowered to 18°C for 18 hours. Cells were harvested by centrifugation, snap-frozen in liquid nitrogen, and stored at -80°C. Cell pellets were resuspended in 20ml binding buffer (50mM Tris pH 8.0, 300mM NaCl, 10mM imidazole, protease inhibitors (Pierce)) at 4°C with stirring. Lysozyme was added at 1mg/ml as well as benzonase (10,000X, Sigma), and samples were lysed by sonication. Lysates were clarified by centrifugation at 40,000 x g, 30 min, 4°C. Supernatant was run by gravity over a column of 6ml NiNTA affinity resin (Qiagen) pre-equilibrated in binding buffer. The column was washed twice with 5 column volumes of wash buffer (50mM Tris pH 8.0, 300mM NaCl, 25mM imidazole) and eluted with four successive 1 column volumes of elution buffer (50mM Tris pH 8.0, 300mM NaCl, 250mM imidazole). Protein content of fractions was estimated by Quickstart Bradford assay and purity determined by SDS-PAGE. Protein-rich fractions were pooled and cleaved by the addition of 1.4mg Ulp1 enzyme (prepared in-house) and 1mM DTT for 1 hour at room temperature. After incubation, the samples were dialyzed overnight against dialysis buffer (50mM HEPES pH 8.0, 150mM NaCl, 7mM β -mercapthoethanol. The completed digestion reaction was applied to NiNTA resin, washed twice with 3 column volumes of dialysis buffer, and eluted with 3 column volumes of elution buffer. The flow-through and wash fractions were pooled, centrifugated at 20,000 x g, 5 min, 4°C to pellet any insoluble material, and further purified by loading onto a 1ml HiTrap HP SP cation exchange column pre-equilibrated in low-salt buffer (50mM HEPES pH8.0, 150mM NaCl). Column was washed extensively with low salt buffer and protein was eluted with a gradient to high-salt buffer (50mM HEPES pH 8.0, 1.2M NaCl). Fractions containing the cleaved SMO pCT were pooled, concentrated, and injected onto a Superdex 200 (10/300) gel filtration column equilibrated in gel filtration buffer (50 mM HEPES pH 7.5, 300 mM NaCl). The peak fractions were collected and analyzed by SDS-PAGE. Fractions containing intact pCT were pooled, snap-frozen in liquid nitrogen, and stored at -80°C. For the experiments in Fig. 2d, wild-type or mutant pCT proteins were purified by NiNTA affinity as described above and used directly in PKA-C activity assays.

In vitro PKA-C activity assays. For *in vitro* PKA-C activity assays the recombinant mouse PKA catalytic subunit (C α) was overexpressed in *E. coli* BL21(DE3) cells after induction with 0.4 mM IPTG for 14 hr at room temperature using the expression vector pRSETb / mPKAC α and then

purified by affinity chromatography using an IP20-resin as described earlier⁸⁸. PKA catalytic activity was assayed using a coupled spectrophotometric assay as described previously⁴². Briefly, the reaction mixture contained 100 mM MOPS (pH 7), 10 mM MgCl₂, 100 μM ATP, 1 mM phosphoenolpyruvate, 15 U/mL lactate dehydrogenase, 70 U/mL pyruvate kinase, 200 mM reduced nicotinamide adenine dinucleotide, 5 mM β-mercaptoethanol with 15-30 nM Cα and 260 μM Kemptide (LRRASLG; GeneCust) as a substrate. Formation of PKA-C:SMO complex (concentration of SMO peptide or recombinant SMO pCT protein are indicated in the figures) was carried out for 3 minutes at room temperature in the assay mixture. The apparent IC₅₀ for SMO pCT were determined by fitting the concentration-dependent activity to a sigmoid dose-response model. All data were plotted as means of at least two independent experiments measured in duplicate each with standard deviation (SD).

MALS. Purified SMO pCT was concentrated to 5 mg/ml and analyzed via SEC-MALS using a Superdex 75 gel chromatography column (GE Healthcare) equilibrated in gel filtration buffer (50 mM HEPES pH 7.5, 300 mM NaCl) with an in-line DAWN MALS detector (Wyatt Technology.)

BRET. BRET in HEK293 or IMCD3 cells was performed as previously described²⁴. Briefly, HEK293 or IMCD3 cells were transiently transfected with nanoluc-tagged SMO (0.3 μg) along with YFP-tagged βarrestin1, PKA-Cα, or PKA-Rlα plasmids. Typically 0.1 μg of each BRET acceptor plasmid was used, with the following two exceptions: (1) Fig. 4b which examined a range of PKA-Cα DNA amounts as indicated in the figure legend; (2) Fig 3d which used 0.3 μg of PKA-Cα-YFP (corresponding to a 0.33-2.68% increase in the size of the endogenous PKA-C pool, as determined by quantitative immunoblotting²⁴) or PKA-Rlα-YFP (which displays no BRET with SMO even though it expresses at substantially higher levels than does PKA-Cα-YFP²⁴); Cells were replated in poly-D-lysine coated white opaque 96-well plates, loaded with 5 μM coelenterazine h (HEK293) or 10 μM furimazine (IMCD3), and analyzed for BRET on a Tecan Spark multimode plate reader. The background signal from cells expressing nanoluc-tagged SMO without BRET acceptor was subtracted from all measurements. For all BRET assays, data represent mean ± SEM from triplicate wells, and data are representative of at least two independent experiments.

Flow cytometry. Cell surface expression of wild-type and mutant SMO constructs was analyzed via flow cytometry as previously described²⁴. Briefly, HEK293-Freestyle cells were infected with BacMam viruses encoding the indicated wild-type or mutant SMO674 constructs. 1-2 days later, live cells were stained with Alexa 647-conjugated M1 anti-FLAG antibody (1:1000, 5 minutes) and analyzed on a Cytotflex flow cytometer (Beckman Coulter).

CREB reporter assays. CREB transcriptional reporter assays were performed as previously described²⁴. Briefly, HEK293FT cells were transfected in 24-well plates with a 30:1 mixture of CRE-Firefly reporter (pGL4.29[luc2p/CRE/Hygro]) and constitutively expressing SV40-Renilla plasmids (20%(w/w)), PKA-Cα (0.625%(w/w)), along with the indicated wild-type or mutant SMO674 plasmids (24%(w/w)). A GFP expression plasmid was used to bring the total amount of DNA in each well to 250 ng. Two days later, reporter activity was measured via dual luciferase assay. Reporter activity is expressed as a ratio of Firefly/Renilla (relative luciferase units (RLU))

For all CREB assays, data represent mean $\pm$ SEM from triplicate wells, and data are representative of at least two independent experiments.

Coimmunoprecipitation assay. SMO / PKA-C coimmunoprecipitation was performed as previously described²⁴, with minor modifications. Briefly, 3 ml HEK293-Freestyle cells were infected with BacMam viruses encoding PKA-C α -YFP the indicated wild-type or mutant SMO674 constructs and solubilized in low-salt solubilization buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 0.1 mM TCEP, 0.5% GDN, 1 mM CaCl₂•6H₂O, protease inhibitor tablet) to prepare a whole-cell lysate. A portion of the lysate was reserved for SDS-PAGE analysis, and the remainder was incubated with FLAG affinity resin (10 μ l settled resin per condition). After a one-hour incubation, resin was washed three times in low-salt wash buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 0.1 mM TCEP, 0.05% GDN, 1 mM CaCl₂•6H₂O), and protein eluted in 40 μ l of the same buffer supplemented with 5 mM EDTA and 0.2 mg/ml FLAG peptide. Proteins were separated by SDS-PAGE on a 4-20% Stain-Free TGX gel (BioRad), and total protein in lysate or eluate was visualized via Stain Free imaging. To detect PKA-C α -YFP, inputs and eluates were transferred to PVDF membranes and processed for immunoblotting with anti-GFP antibodies as previously described²⁴. In-gel Pro-Q Diamond assay to detect phosphoprotein was performed according to the manufacturer's instructions, as previously described²⁴.

HEK293 imaging. Imaging and quantification of SMO / PKA-C colocalization in HEK293 cells were performed as previously described²⁴, with minor modifications. Briefly, HEK293 Flp-in T-rex stable cell lines coexpressing GFP-tagged PKA-C α with either wild-type or mutant FLAG-tagged SMO674 were plated onto μ -slide 8-well glass chamberslides (ibidi), and treated overnight with 1 μ g/ml doxycycline (to induce SMO expression) and 1 μ M SAG21k (to induce SMO activation). Live cells were subsequently stained for 5 min with an Alexa647-conjugated M1 anti-FLAG antibody (1:2000) followed by washing in HBSS, mounting, and visualization. Images were collected on a Leica SP8 laser scanning confocal microscope, using a 40x water immersion lens. All images were acquired with identical zoom / exposure / gain settings and processed identically in Fiji. SMO / PKA-C colocalization was determined by measuring the background-subtracted PKA-C (green) fluorescence at the membrane ($\text{GFP}_{\text{membrane}} - \text{GFP}_{\text{cytoplasm}}$) and the background-subtracted SMO (red) fluorescence at the membrane ($\text{Alexa647}_{\text{membrane}} - \text{Alexa647}_{\text{cytoplasm}}$), using four independent line-scans across the membrane in Fiji. The background-subtracted green fluorescence divided by the background-subtracted red fluorescence is referred to as "SMO / PKA-C colocalization" and reported in arbitrary units (AU).

IMCD3 imaging. Imaging and quantification of SMO ciliary accumulation and SMO / PKA-C colocalization in IMCD3 cells were performed as previously described²⁴. Briefly, to assess ciliary accumulation of wild-type or mutant SMO, IMCD3 cells were transiently transfected on coverslips with the indicated myc-tagged wild-type or mutant SMO / pGEN constructs, grown to confluency, fixed, and permeabilized. Coverslips were then stained with anti-myc (SMO) and anti-Arl13b (cilia) antibodies, along with DAPI to mark the nucleus. For quantitative assessment of SMO / PKA-C colocalization in live IMCD3 cilia, cell lines coexpressing mNeonGreen-tagged PKA-C α with either wild-type or mutant FLAG-tagged SMO were plated onto μ -slide 8-well glass chamberslides (ibidi), grown to confluency, and treated overnight in low-serum medium with 1 μ M SAG21k (to

induce SMO activation and ciliary accumulation.) A previously described cell line coexpressing mNeonGreen-tagged Nb β 2AR80 with wild-type FLAG-tagged SMO²⁴ was used as a negative control. Live cells were subsequently stained for 5-10 min with an Alexa647-conjugated M1 anti-FLAG antibody (1:1000) and Hoechst counterstain, followed by washing in HBSS, mounting, and visualization. Cells were imaged on a Leica SP8 laser scanning confocal using a 40x water immersion lens. Three-dimensional reconstructions of Z-stacks were performed in Fiji using the 3D Viewer plugin. Quantification of SMO (red) and PKA-C/Nb β 2AR80 (green) staining was performed using CiliaQ⁹⁵ and reported as a ratio of the green fluorescence in each cilium, normalized to the red fluorescence ("Colocalization with SMO in cilia (AU)"), as previously described²⁴. All images were acquired with identical zoom / exposure / gain settings.

GLI reporter assays. GLI transcriptional reporter assays were performed as previously described²⁴. Briefly, *Smo*^{-/-} MEFs were transfected with a 30:1 mixture of 8xGli-Firefly and SV40-Renilla plasmids (50% (w/w)), along with the indicated full-length wild-type or mutant SMO constructs (in a pGEN vector, with C-terminal myc tag) (2%(w/w)), and GFP to adjust the amount of DNA to 250 ng/well. Cells were cultured to confluency, shifted to 0.5% FBS-containing medium, and treated with control or ShhN conditioned medium (1:20 dilution) for 2 days. Reporter activity was then measured via dual luciferase assay. For all GLI assays, data represent mean $\pm$ SEM from triplicate wells, and data are representative of at least two independent experiments.

Zebrafish embryological studies. Zebrafish mRNA injection, fixing, mounting, and staining of muscle cell markers were all performed as previously described²⁴, except that a far-red secondary antibody (donkey anti-rabbit 649, Jackson Laboratory, 1:500) was used instead of a red secondary antibody during immunohistochemistry.

Software. PDB files were viewed using Pymol. Graphs, curve fitting, and statistical analysis was performed in GraphPad Prism. Alignments were generated using CLUSTAL Omega. Graphics were generated using BioRender or Adobe Illustrator.

ACKNOWLEDGMENTS:

We thank J. Zalatan for making us aware of the parallels between SMO / PKA-C regulation in the Hh pathway and LRP / GSK-3 β regulation in the Wnt pathway. We thank S. Lusk and K. Kwan for providing *smo*-null zebrafish (*smo*^{hi1640}), and D. Klatt Shaw and D. Grunwald for sharing advice and reagents regarding zebrafish immunohistochemistry. We thank J. Müller and S. Kasten for excellent technical assistance. We thank the Johnson Foundation Structural Biology and Biophysics Core at the Perelman School of Medicine (Philadelphia, PA) for performing SEC-MALS analyses. We thank D. Julius, K. Basham, A. Manglik, M. He, and S. Nakielnny for providing feedback on this manuscript. B.R.M. acknowledges support from the 5 for the Fight Foundation and a Cancer Center Support Grant Pilot Project Fund from the Huntsman Cancer Institute. This work was supported by the funding line Future (PhosMOrg) of University of Kassel (F.W.H.), and NIH grants R01GM100310-08 (G.V.), 1R35GM130389 (S.S.T.), 1R03TR002947 (S.S.T.) and 1R35GM133672 (B.R.M.).

AUTHOR CONTRIBUTIONS:

J.T.H. designed, executed, and interpreted CREB and GLI reporter assays. C.D.A. designed, executed, and interpreted HEK293 BRET assays. J.B. designed, executed, and interpreted fluorescence polarization studies and peptide array studies. D.B. designed, executed, and interpreted *in vitro* PKA-C activity assays and SPR studies (the latter with assistance from J.W.B.). I.B.N. developed SMO pCT purification approaches and purified this domain for *in vitro* PKA-C activity assays. C.O. designed, executed, and interpreted NMR studies. D.S.H. designed, executed, and interpreted zebrafish embryology studies. J-F.Z. designed, executed, and interpreted co-immunoprecipitation and IMCD3 BRET assays. J.L.C. designed, executed, and interpreted HEK293 confocal imaging studies. L.V. performed initial fluorescence polarization studies. C.C.K. collaborated with J.B. to develop SMO peptide arrays. V.L.R-P. provided advice and guidance on mutagenesis experiments to disrupt SMO / PKA-C interactions. S.S.T. and B.R.M. conceived the project. G.V., F.W.H., S.S.T., and B.R.M. interpreted data and provided overall project supervision. B.R.M. performed IMCD3 ciliary imaging studies and wrote the manuscript with assistance from J.T.H., C.D.A., and I.B.N.

COMPETING FINANCIAL INTERESTS:

The authors declare no competing financial interests.

Correspondence and requests for materials should be addressed to S.S.T. or B.R.M.

FIGURES

Fig. 1: SMO binds PKA-C as a pseudosubstrate. **a**, CLUSTAL alignment of the mouse SMO pCT with the PKIα pseudosubstrate region. Additional PKA-C pseudosubstrate and substrate sequences are provided for comparison^{27,96,97}. P-site is yellow; other key conserved residues are green. Spiral cartoon above alignment indicates predicted SMO helical region. Standard (615-638) and extended (615-652) SMO peptides used for *in vitro* assays are colored red or black, respectively. Inset, structure of PKA-Cα bound to PKIα(5-24) (PDB: 3FJQ), with ATP colored orange and key PKI residues colored as described above. **b**, Top, fluorescence polarization assay employing FAM-labeled SMO peptide, 1 mM ATP, and varying concentrations of human PKA-Cα. Bottom, the same assay except with 3 μM PKA-Cα and varying concentrations of ATP. **c**, Overlay of purified mouse PKA-Cα onto an array of SMO peptides containing the indicated substitutions in the P-13, P-3, and P-2 positions. **d**, SPR sensorgram for binding of GST-tagged wild-type SMO pCT to a series of PKA-Cα concentrations ranging from 0.156 μM to 10 μM. Buffer contained 1 mM ATP and 10 mM MgCl₂. **e**, As **d**, but with SMO pCT harboring the WRR mutation.

Fig. 2: SMO is a pseudosubstrate inhibitor of PKA-C. **a**, Mapping of amide backbone chemical shift perturbations (CSP, δ) for $[^1\text{H}, ^{15}\text{N}]$ -labeled PKA-Ca bound to nucleotide (ATPyN) and either an extended SMO PKI peptide (black) or a control PKI α (5-24) peptide⁹⁰ (blue), calculated relative to ATPyN-bound PKA-Ca without peptide. Key functional domains of PKA-C are highlighted along the X-axis. Magenta spheres indicate PKA-Ca residues (R144, D166, R194, and F297) that show a signal for PKI α (5-24) but not SMO. Right, CSP values were mapped onto the PKA-Ca structure (PDB: 4WB5) and displayed as a heatmap. **b**, Spectrophotometric assay of PKA-Ca substrate phosphorylation, in the presence of standard (red) or extended (coral) SMO peptides (see Fig. 1a) or a control PKI α (5-24) peptide (grey). **c**, Concentration dependent inhibition of PKA-Ca with recombinant SMO pCT. **d**, As **b**, but comparing wild-type vs. WRR mutant versions of the recombinant SMO pCT. Inhibition in **b-d** is calculated relative to a control without SMO peptide (Ctrl.). See Extended Data Table 1 for statistical analysis.

Fig. 3: The SMO PKI motif is required for Hh signal transduction. **a**, Top, schematic diagram of truncated SMO expression constructs. Bottom, BRET analysis of SMO / PKA-C interactions in HEK293 cells expressing nanoluc-tagged wild-type (wt) SMO657 or SMO657 harboring the WRR mutation (WRR), along with YFP-tagged PKA-Ca. SMO566 (which lacks the C-tail) serves as a negative control donor. YFP-tagged β arrestin1 (which exhibits minimal binding to SMO) and NbSmo2 (which binds the intracellular surface of the SMO 7TM domain) serve as negative and positive control acceptors, respectively²⁴. **b**, Left, confocal images of live HEK293 cells coexpressing GFP-tagged PKA-Ca with FLAG-tagged wild-type or mutant SMO674. Cells were treated with SMO agonist (SAG21k). Scale bar = 10 μ m. Right, quantification of SMO / PKA-C colocalization (n=34-48 cells per condition). **c**, CREB transcriptional reporter assay, reflecting PKA-C mediated substrate phosphorylation, in HEK293 cells transfected with PKA-Ca and the indicated SMO674 constructs. **d**, BRET analysis of SMO / PKA-C interactions in IMCD3 cells expressing nanoluc-tagged wild-type or mutant SMO, along with low levels of PKA-Ca-YFP. PKA-R1a-YFP serves as a negative control. Under these conditions, PKA-Ca-YFP is expressed at substantially lower levels than PKA-R1a-YFP²⁴. **e**, GLI transcriptional reporter assay in *Smo*^{-/-} MEFs transfected with a GFP negative control (Ctrl.), or the indicated wild-type (wt) or mutant SMO constructs. Cells were treated with conditioned medium containing the N-terminal signaling domain of Sonic hedgehog (ShhN), or control, non-ShhN-containing conditioned medium (Ctrl). **f**, Wild-type or *smo*^{-/-} zebrafish injected with the indicated mRNA constructs were stained for Prox1 (magenta) or Engrailed (En, green) to mark muscle fiber nuclei. n=12 (uninjected), n=41 (SMO wt), n= 47 (SMO WRR). See Extended Data Table 1 for statistical analysis.

Fig. 4: An avidity-based mechanism for SMO inhibition of PKA-C. **a**, Annotated sequence of the mouse SMO pCT. PKI motif is indicated in red, along with GRK2/3 phosphorylation sites (orange) and residues 570-581 (aquamarine box), previously shown to influence SMO / PKA-C interactions and Hh signal transduction^{24,61}. **b**, BRET analysis of SMO / PKA-C interactions in HEK293 cells transfected with full-length wild-type SMO (wt, navy) WRR mutant (purple), or C-terminally truncated (SMO566, white) versions of SMO as donor, and the indicated DNA amounts of PKA-Cα as acceptor. Nonspecific signal is indicated by the negative control BRET acceptor

710 β arrestin1 (dashed line). See Extended Data Table 1 for statistical analysis. **c**, Representative
 711 image of a tiled array of 18mer peptides covering the complete C-tail of human SMO, probed with
 712 PKA-C α as in Fig. 1c. Peptides that bind are boxed and their sequences indicated. **d**, Proposed
 713 model for Hh signal transduction, as described in "Discussion".
 714

		P ₁₃	P ₃	P ₂	P ₁	
Mus_musculus	AGLAFDLNEPS-----ADVSSAWAQHV-TKMVARRGAILPQDVS-VTPVATPVPP---EE					654
Homo_sapiens	AGLAFDLNEPS-----ADVSSAWAQHV-TKMVARRGAILPQDIS-VTPVATPVPP---EE					650
Danio_rerio	AGINFDLNEPS-----IEMSSAWAQHV-TKMVARRGAILPQDIS-VTPTGTPIPP--PEE					630
Xenopus_laevis	AGLNFDMEPS-----ADMSSAWAQHV-TKMVARRGAILPQDVS-VTPVATPVPP---EE					623
Gallus_gallus	AGLAFDINEPS-----ADVSSAWAQHV-TKMVARRGAILPQDVS-VTPVATPVPP---EE					517
Drosophila_melanogaster	VGLNFDVNDLNS-SETNDISSTWAAYL-PQCVKRRMALTGAATGNSSSH-GPRKNSLDSE					671
Ciona_intestinalis	LGMNFDLHVS-----QEMSSSWVRNV-PNMVKRRGMLPMEQPHDNVE-----					611
Lytechinus_variegatus	IGMKLDLPPSSVVGDDPTSSSSWGNNVPVRMLARRGAAYPITLGNSPRATPDSS--DSG					641
Petromyzon_marinus	VGMDLNLDDSG-----S-DSSSWINHV-TKMVARRGAILP-----					581
Novocrania_anomala	LGMKFDLNSVT----SQDMSSAWMEAV-PRLVRRRGGMTHPTAGTLRRYSDS-----DI					621

Extended Data Fig.1: Sequence alignment of SMO PKI motif. Extended alignment of a portion of the pCT from the indicated SMO orthologs, with key PKI motif residues colored as in Fig. 1a.

Extended Data Fig. 2: Further binding and peptide array studies, SPR sensorgrams, NMR spectra, and SMO pCT purification strategy. **a**, Fluorescence polarization assays using mouse PKA-Cα, performed as in Fig. 1b. **b**, Peptide array, performed as in Fig. 1c, but with individual residues in the human SMO PKI motif mutated to alanine. **c**, SPR sensorgram for 5 μM PKA-Cα binding to GST-tagged wild-type (blue) or WRR mutant (purple) SMO pCT, or a PKIα positive control (red), in the presence of ATP and MgCl₂. **d**, Steady-state analysis of binding interactions

between human PKA-C α and a recombinant wild-type (wt, blue) or mutant (WRR, magenta) SMO pCT, as assessed by SPR. **e**, SPR sensorgram, performed as in **c**, but with ATP and MgCl₂ omitted from the buffer. PKA-C α was present at 2.5 μ M. Note that although removal of ATP and MgCl₂ does not completely eliminate steady-state binding to the PKI α positive control, it dramatically accelerates the dissociation rate, as expected. **f**, [¹H, ¹⁵N] heteronuclear single quantum coherence (HSQC) spectra used to calculate CSP values. Spectra were acquired from PKA-C α / ATP γ N complexed with a SMO peptide at 1:0 (red), 1:1 (yellow), 1:2 (green), or 1:4 (purple) molar ratios. CSP values were calculated from the spectrum corresponding to a 1:4 molar ratio and plotted in Fig. 2a. **g**, Purification of SMO pCT domain from *E. coli*. Following size exclusion chromatography (left), the purified protein was analyzed by SDS-PAGE (right). The SMO pCT elutes as two peaks, the earlier of which is the intact pCT and the later of which corresponds to a truncated fragment (data not shown). Fractions containing the intact pCT (red box) were pooled and used for subsequent experiments. **h**, Multi-angle light scattering coupled with size exclusion chromatography (SEC-MALS) was used to determine the protein oligomeric state for the pooled fractions indicated in **g**. The average molecular mass was calculated as $M_w = 11.25 \pm 2.1$ kDa, close to the predicted molecular mass for a monomer (10.1 kDa).

Extended Data Fig. 3: Coimmunoprecipitation studies and ciliary colocalization studies to assess SMO / PKA-C interactions. **a**, Coimmunoprecipitation of PKA-C α -YFP with the indicated FLAG-tagged wild-type or mutant SMO constructs was assessed using FLAG chromatography from lysates of transfected HEK293 cells. **b**, Left, Colocalization of FLAG-tagged wild-type or mutant SMO674 (magenta) with mNeonGreen-tagged PKA-C α (green) in ciliated IMCD3 cells stably expressing both constructs and treated with a SMO agonist, SAG21k, to induce SMO ciliary localization. Cilia are marked by the SMO stain. mNeonGreen-tagged Nb β 2AR80 (which does not bind SMO²⁴) serves as a negative control. 3D reconstructions from Z-stacks of confocal live-

752 cell images are shown. Right, quantification of microscopy studies (n=142-244 cilia per condition).
753 See Extended Data Table 1 for statistical analysis.
754

Extended Data Fig. 4: Controls for SMO / PKA-C binding, colocalization, and signaling studies. **a**, Expression levels of SMO constructs in Fig. 3a, assessed by whole-cell nanoluc measurements. **b**, Surface levels of N-terminally FLAG-tagged wild-type or mutant SMO674 constructs were quantified via expression in HEK293 cells followed by FLAG staining and flow cytometry. Mock-infected cells stained with FLAG antibody (red) serve as a negative control. **c**, Ciliary localization in IMCD3 cells of myc-tagged wild-type or mutant SMO proteins (magenta). Cilia were visualized with Arl13b antibody (green). Scale bar = 5 μ m. **d**, GRK2/3-dependent phosphorylation of FLAG-tagged wild-type or mutant SMO674 constructs was determined via expression in HEK293 cells treated with or without the GRK2/3 inhibitor cmpd101, followed by FLAG purification. Levels of total and phosphorylated SMO were assessed by Stain Free imaging and ProQ Diamond fluorescence, respectively. SMO566, which is not phosphorylated by GRK2/3 (as it does not contain the C-tail and therefore lacks all previously mapped physiological GRK2/3 phosphorylation sites²⁴), serves as a negative control.

b Human SMO C-tail

535 WTKATLLIWRRTWCRLTGQSDDEPKRIKKSKMIAKAFSKRHELLQNPGQELSFMSM
HTVSHDGPVAGLAFDLNEPSADVSSAWAQHVTKMVARRGAILPQDISVTPVATPVPPE
EQANLWLVEAEISPELQKRLGRKKRRKRKKEVCPLAPPELHPPAPAPSTIPRLPQL
PRQKCLVAAGAWGAGDSCRQGAWTLVSNPFCPEPSPPQDPFLPSAPAPVAWAHGRROG
LGPIHSRTNLMDELMADSSDF 786

Extended Data Fig. 5: Complete data set from SMO C-tail peptide array studies. **a**, The same SMO tiled peptide array from Fig. 4c, but including the sequences of all positive hits in each array cluster. **b**, Complete human SMO C-tail sequence used to create the peptide array. In **a,b**, the SMO PKI motif identified in the pCT is indicated in red. Key residues in this PKI motif, along with ones in the candidate PKI motif in the dCT, are colored as in Fig. 1a.

Extended Data Fig. 6: Similarity between signal transduction mechanisms in the Hh and Wnt pathways. Schematic diagram of transmembrane signal transduction in the Hh (left) and Wnt (right) pathways. During Hh signal transduction, active SMO is phosphorylated on its cytoplasmic tail by GRK2/3, triggering membrane sequestration and inhibition of PKA-C, and ultimately stabilization and activation of GLI. During Wnt signal transduction, active LRP5/6 is phosphorylated on its cytoplasmic tail by glycogen synthase kinase (GSK)-3β and casein kinase (CK)-1α, triggering membrane sequestration and inhibition of GSK-3β, and ultimately stabilization and activation of β-catenin⁷⁴⁻⁷⁷. Note that this is a simplified and highly schematized diagram and is not intended to be comprehensive; many other components of both pathways (for example, the destruction complex in which GSK-3β and β-catenin reside) are omitted in order to highlight mechanistic similarities between the underlying transmembrane signaling mechanisms.

797 **Extended Data Table 1.** Tests of statistical significance for all figures.

Figure	Condition 1	Condition 2	Significant?	p-value
2b	multiple comparison	multiple comparison	yes	<0.001 ^a
2d	None	wt	yes	<0.0001 ^b
2d	WRR	wt	yes	<0.0001 ^b
2d	None	WRR	no	n.s. ^b
3a	SMO657 wt: β arrestin1	SMO657 wt: NbSmo2	yes	0.00053
3a	SMO657 wt: β arrestin1	SMO657 wt: PKA-C	yes	0.00438
3a	SMO657 wt: PKA-C	SMO657 WRR: PKA-C	yes	0.001553
3a	SMO657 wt: PKA-C	SMO566 wt: PKA-C	yes	0.003181
3a	SMO657 WRR: β arrestin1	SMO657 WRR: PKA-C	no	0.051139
3a	SMO657 WRR: β arrestin1	SMO657 WRR: NbSmo2	yes	0.000165
3a	SMO566 wt: β arrestin1	SMO566 wt: PKA-C	no	0.22268
3a	SMO566 wt: β arrestin1	SMO566 wt: NbSmo2	yes	0.00162
3b	SMO674: wt	SMO674: WRR	yes	<0.0001
3c	None	PKA-C	yes	0.000723
3c	PKA-C	PKA-C + SMO674 wt	yes	0.000995
3c	PKA-C	PKA-C + SMO674 WRR	no	0.267104
3c	PKA-C	PKA-C + SMO674 A635S	no	0.544481
3c	PKA-C + SMO674 wt	PKA-C + SMO674 WRR	yes	0.004314
3c	PKA-C + SMO674 wt	PKA-C + SMO674 A635S	yes	0.000513
3d	SMO wt: PKA-R	SMO wt: PKA-C	yes	0.012831
3d	SMO wt: PKA-C	SMO WRR: PKA-C	yes	0.01315
3d	SMO WRR: PKA-R	SMO WRR: PKA-C	no	0.113776
3e	Ctrl.: Ctrl	Neg.: ShhN	no	0.63116
3e	SMO wt: Ctrl	SMO wt: ShhN	yes	0.010704
3e	SMO wt: ShhN	SMO WRR: ShhN	yes	0.010346
3e	SMO wt: ShhN	SMO A635S: ShhN	yes	0.010945
4b	β arrestin1: SMO WRR	PKA-C (0.04 μ g): SMO WRR	no	0.885586
4b	β arrestin1: SMO WRR	PKA-C (0.08 μ g): SMO WRR	yes	0.001529
4b	β arrestin1: SMO WRR	PKA-C (0.12 μ g): SMO WRR	yes	0.000368
4b	β arrestin1: SMO WRR	PKA-C (0.16 μ g): SMO WRR	yes	0.000109
4b	β arrestin1: SMO WRR	PKA-C (0.20 μ g): SMO WRR	yes	0.000085

4b	PKA-C (0.04 µg): SMO wt	PKA-C (0.04 µg): SMO WRR	yes	0.004444
4b	PKA-C (0.08 µg): SMO wt	PKA-C (0.08 µg): SMO WRR	yes	0.000502
4b	PKA-C (0.12 µg): SMO wt	PKA-C (0.12 µg): SMO WRR	yes	0.001341
4b	PKA-C (0.16 µg): SMO wt	PKA-C (0.16 µg): SMO WRR	yes	0.004344
4b	PKA-C (0.20 µg): SMO wt	PKA-C (0.20 µg): SMO WRR	yes	0.000029
4b	PKA-C (0.04 µg): SMO566	PKA-C (0.04 µg): SMO WRR	yes	0.00368
4b	PKA-C (0.08 µg): SMO566	PKA-C (0.08 µg): SMO WRR	no	0.180776
4b	PKA-C (0.12 µg): SMO566	PKA-C (0.12 µg): SMO WRR	yes	0.014786
4b	PKA-C (0.16 µg): SMO566	PKA-C (0.16 µg): SMO WRR	yes	0.006063
4b	PKA-C (0.20 µg): SMO566	PKA-C (0.20 µg): SMO WRR	yes	0.001566
ED 3b	SMO wt + PKA-C	SMO WRR + PKA-C	yes	<0.0001
ED 4a	SMO657	SMO657 WRR	no	0.057394

^a To check for significant differences between PKA-C lacking peptide (control) and each peptide concentration a one-way ANOVA with Tukey's multiple comparison was performed. All comparisons show $p \leq 0.0001$ (****) significance except the comparison between 125µM standard vs extended peptide ($p \leq 0.001$, ***).

^b To check for significant differences between PKA lacking peptide (control) and the indicated recombinant pCT fragment a one-way ANOVA with Tukey's multiple comparison was performed (**** $\triangleq p \leq 0.0001$, ns $\triangleq$ not significant).

Extended Data Table 2 Quantification of phenotypes for zebrafish embryogenesis studies. The # and % of animals exhibiting U-shaped somites (indicative of a failure in Hh signaling during somitogenesis^{55,98-100}) are indicated. Note that in both the WRR and A635S mutant conditions, close to 25% of animals exhibited U-shaped somites, consistent with the Mendelian inheritance of a null *smo* allele from the initial heterozygous incross.

mRNA injected	# Animals from <i>smo</i> (+/-) incross	# Animals with U-shaped somites	% Animals with U-shaped somites
SMO wt	154	2	1.3%
SMO WRR	226	61	27.0%
SMO A635S	70	19	27.1%

REFERENCES:

- 1 Briscoe, J. & Therond, P. P. The mechanisms of Hedgehog signalling and its roles in development and disease. *Nat Rev Mol Cell Biol* **14**, 416-429, doi:10.1038/nrm3598 (2013).
- 2 Ingham, P. W. & McMahon, A. P. Hedgehog signaling in animal development: paradigms and principles. *Genes Dev* **15**, 3059-3087, doi:10.1101/gad.938601 (2001).
- 3 Ingham, P. W., Nakano, Y. & Seger, C. Mechanisms and functions of Hedgehog signalling across the metazoa. *Nat Rev Genet* **12**, 393-406, doi:10.1038/nrg2984 (2011).
- 4 Kong, J. H., Siebold, C. & Rohatgi, R. Biochemical mechanisms of vertebrate hedgehog signaling. *Development* **146**, doi:10.1242/dev.166892 (2019).
- 5 Muenke, M. & Beachy, P. A. Genetics of ventral forebrain development and holoprosencephaly. *Curr Opin Genet Dev* **10**, 262-269, doi:10.1016/s0959-437x(00)00084-8 (2000).
- 6 Digilio, M. C. *et al.* Atrioventricular canal defect and genetic syndromes: The unifying role of sonic hedgehog. *Clin Genet* **95**, 268-276, doi:10.1111/cge.13375 (2019).
- 7 Anderson, E., Peluso, S., Lettice, L. A. & Hill, R. E. Human limb abnormalities caused by disruption of hedgehog signaling. *Trends Genet* **28**, 364-373, doi:10.1016/j.tig.2012.03.012 (2012).
- 8 Pak, E. & Segal, R. A. Hedgehog Signal Transduction: Key Players, Oncogenic Drivers, and Cancer Therapy. *Dev Cell* **38**, 333-344, doi:10.1016/j.devcel.2016.07.026 (2016).
- 9 Wu, F., Zhang, Y., Sun, B., McMahon, A. P. & Wang, Y. Hedgehog Signaling: From Basic Biology to Cancer Therapy. *Cell Chem Biol* **24**, 252-280, doi:10.1016/j.chembiol.2017.02.010 (2017).
- 10 Hui, C. C. & Angers, S. Gli proteins in development and disease. *Annu Rev Cell Dev Biol* **27**, 513-537, doi:10.1146/annurev-cellbio-092910-154048 (2011).
- 11 Deshpande, I. *et al.* Smoothed stimulation by membrane sterols drives Hedgehog pathway activity. *Nature* **571**, 284-288, doi:10.1038/s41586-019-1355-4 (2019).
- 12 Huang, P. *et al.* Structural Basis of Smoothed Activation in Hedgehog Signaling. *Cell* **174**, 312-324 e316, doi:10.1016/j.cell.2018.04.029 (2018).
- 13 Byrne, E. F. X. *et al.* Structural basis of Smoothed regulation by its extracellular domains. *Nature* **535**, 517-522, doi:10.1038/nature18934 (2016).
- 14 Qi, X., Friedberg, L., De Bose-Boyd, R., Long, T. & Li, X. Sterols in an intramolecular channel of Smoothed mediate Hedgehog signaling. *Nat Chem Biol* **16**, 1368-1375, doi:10.1038/s41589-020-0646-2 (2020).
- 15 Wang, B., Fallon, J. F. & Beachy, P. A. Hedgehog-regulated processing of Gli3 produces an anterior/posterior repressor gradient in the developing vertebrate limb. *Cell* **100**, 423-434, doi:10.1016/s0092-8674(00)80678-9 (2000).
- 16 Humke, E. W., Dorn, K. V., Milenkovic, L., Scott, M. P. & Rohatgi, R. The output of Hedgehog signaling is controlled by the dynamic association between Suppressor of Fused and the Gli proteins. *Genes Dev* **24**, 670-682, doi:10.1101/gad.1902910 (2010).
- 17 Niewiadomski, P. *et al.* Gli protein activity is controlled by multisite phosphorylation in vertebrate Hedgehog signaling. *Cell Rep* **6**, 168-181, doi:10.1016/j.celrep.2013.12.003 (2014).
- 18 Li, J. *et al.* PKA-mediated Gli2 and Gli3 phosphorylation is inhibited by Hedgehog signaling in cilia and reduced in Talpid3 mutant. *Dev Biol* **429**, 147-157, doi:10.1016/j.ydbio.2017.06.035 (2017).
- 19 Nachury, M. V. & Mick, D. U. Establishing and regulating the composition of cilia for signal transduction. *Nat Rev Mol Cell Biol* **20**, 389-405, doi:10.1038/s41580-019-0116-4 (2019).

- 20 Reiter, J. F. & Leroux, M. R. Genes and molecular pathways underpinning ciliopathies. *Nat Rev Mol Cell Biol* **18**, 533-547, doi:10.1038/nrm.2017.60 (2017).
- 21 Gigante, E. D. & Caspary, T. Signaling in the primary cilium through the lens of the Hedgehog pathway. *Wiley Interdiscip Rev Dev Biol* **9**, e377, doi:10.1002/wdev.377 (2020).
- 22 Lefkowitz, R. J. The superfamily of heptahelical receptors. *Nat Cell Biol* **2**, E133-136, doi:10.1038/35017152 (2000).
- 23 Pierce, K. L., Premont, R. T. & Lefkowitz, R. J. Seven-transmembrane receptors. *Nat Rev Mol Cell Biol* **3**, 639-650, doi:10.1038/nrm908 (2002).
- 24 Arveseth, C. D. *et al.* Smoothed transduces hedgehog signals via activity-dependent sequestration of PKA catalytic subunits. *PLoS Biol* **19**, e3001191, doi:10.1371/journal.pbio.3001191 (2021).
- 25 Taylor, S. S., Ilouz, R., Zhang, P. & Kornev, A. P. Assembly of allosteric macromolecular switches: lessons from PKA. *Nat Rev Mol Cell Biol* **13**, 646-658, doi:10.1038/nrm3432 (2012).
- 26 Dalton, G. D. & Dewey, W. L. Protein kinase inhibitor peptide (PKI): a family of endogenous neuropeptides that modulate neuronal cAMP-dependent protein kinase function. *Neuropeptides* **40**, 23-34, doi:10.1016/j.npep.2005.10.002 (2006).
- 27 Soberg, K. & Skalhogg, B. S. The Molecular Basis for Specificity at the Level of the Protein Kinase a Catalytic Subunit. *Front Endocrinol (Lausanne)* **9**, 538, doi:10.3389/fendo.2018.00538 (2018).
- 28 Taylor, S. S., Zhang, P., Steichen, J. M., Keshwani, M. M. & Kornev, A. P. PKA: lessons learned after twenty years. *Biochim Biophys Acta* **1834**, 1271-1278, doi:10.1016/j.bbapap.2013.03.007 (2013).
- 29 Kemp, B. E. & Pearson, R. B. Protein kinase recognition sequence motifs. *Trends Biochem Sci* **15**, 342-346, doi:10.1016/0968-0004(90)90073-k (1990).
- 30 Scott, J. D., Fischer, E. H., Demaille, J. G. & Krebs, E. G. Identification of an inhibitory region of the heat-stable protein inhibitor of the cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A* **82**, 4379-4383, doi:10.1073/pnas.82.13.4379 (1985).
- 31 Cheng, H. C., van Patten, S. M., Smith, A. J. & Walsh, D. A. An active twenty-amino-acid-residue peptide derived from the inhibitor protein of the cyclic AMP-dependent protein kinase. *Biochem J* **231**, 655-661, doi:10.1042/bj2310655 (1985).
- 32 Corbin, J. D. *et al.* Studies on the properties and mode of action of the purified regulatory subunit of bovine heart adenosine 3':5'-monophosphate-dependent protein kinase. *J Biol Chem* **253**, 3997-4003 (1978).
- 33 Scott, J. D., Glaccum, M. B., Fischer, E. H. & Krebs, E. G. Primary-structure requirements for inhibition by the heat-stable inhibitor of the cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A* **83**, 1613-1616, doi:10.1073/pnas.83.6.1613 (1986).
- 34 Glass, D. B., Cheng, H. C., Mende-Mueller, L., Reed, J. & Walsh, D. A. Primary structural determinants essential for potent inhibition of cAMP-dependent protein kinase by inhibitory peptides corresponding to the active portion of the heat-stable inhibitor protein. *J Biol Chem* **264**, 8802-8810 (1989).
- 35 Knighton, D. R. *et al.* Structure of a peptide inhibitor bound to the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 414-420, doi:10.1126/science.1862343 (1991).
- 36 Knighton, D. R. *et al.* Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 407-414, doi:10.1126/science.1862342 (1991).
- 37 Johnson, D. A., Akamine, P., Radzio-Andzelm, E., Madhusudan, M. & Taylor, S. S. Dynamics of cAMP-dependent protein kinase. *Chem Rev* **101**, 2243-2270, doi:10.1021/cr000226k (2001).

- 38 Olivieri, C. *et al.* Multi-state recognition pathway of the intrinsically disordered protein kinase inhibitor by protein kinase A. *Elife* **9**, doi:10.7554/eLife.55607 (2020).
- 39 Kim, J. *et al.* The role of ciliary trafficking in Hedgehog receptor signaling. *Sci Signal* **8**, ra55, doi:10.1126/scisignal.aaa5622 (2015).
- 40 Whitehouse, S. & Walsh, D. A. Mg X ATP2-dependent interaction of the inhibitor protein of the cAMP-dependent protein kinase with the catalytic subunit. *J Biol Chem* **258**, 3682-3692 (1983).
- 41 Knape, M. J. *et al.* Divalent metal ions control activity and inhibition of protein kinases. *Metallomics* **9**, 1576-1584, doi:10.1039/c7mt00204a (2017).
- 42 Cook, P. F., Neville, M. E., Jr., Vrana, K. E., Hartl, F. T. & Roskoski, R., Jr. Adenosine cyclic 3',5'-monophosphate dependent protein kinase: kinetic mechanism for the bovine skeletal muscle catalytic subunit. *Biochemistry* **21**, 5794-5799, doi:10.1021/bi00266a011 (1982).
- 43 Marullo, S. & Bouvier, M. Resonance energy transfer approaches in molecular pharmacology and beyond. *Trends Pharmacol Sci* **28**, 362-365, doi:10.1016/j.tips.2007.06.007 (2007).
- 44 Machleidt, T. *et al.* NanoBRET--A Novel BRET Platform for the Analysis of Protein-Protein Interactions. *ACS Chem Biol* **10**, 1797-1804, doi:10.1021/acschembio.5b00143 (2015).
- 45 Shaywitz, A. J. & Greenberg, M. E. CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu Rev Biochem* **68**, 821-861, doi:10.1146/annurev.biochem.68.1.821 (1999).
- 46 Knape, M. J. *et al.* Divalent Metal Ions Mg(2)(+) and Ca(2)(+) Have Distinct Effects on Protein Kinase A Activity and Regulation. *ACS Chem Biol* **10**, 2303-2315, doi:10.1021/acschembio.5b00271 (2015).
- 47 Mick, D. U. *et al.* Proteomics of Primary Cilia by Proximity Labeling. *Dev Cell* **35**, 497-512, doi:10.1016/j.devcel.2015.10.015 (2015).
- 48 Desai, P. B., Stuck, M. W., Lv, B. & Pazour, G. J. Ubiquitin links smoothened to intraflagellar transport to regulate Hedgehog signaling. *J Cell Biol* **219**, doi:10.1083/jcb.201912104 (2020).
- 49 Pal, K. *et al.* Smoothened determines beta-arrestin-mediated removal of the G protein-coupled receptor Gpr161 from the primary cilium. *J Cell Biol* **212**, 861-875, doi:10.1083/jcb.201506132 (2016).
- 50 Varjosalo, M., Li, S. P. & Taipale, J. Divergence of hedgehog signal transduction mechanism between Drosophila and mammals. *Dev Cell* **10**, 177-186, doi:10.1016/j.devcel.2005.12.014 (2006).
- 51 Myers, B. R., Neahring, L., Zhang, Y., Roberts, K. J. & Beachy, P. A. Rapid, direct activity assays for Smoothened reveal Hedgehog pathway regulation by membrane cholesterol and extracellular sodium. *Proc Natl Acad Sci U S A* **114**, E11141-E11150, doi:10.1073/pnas.1717891115 (2017).
- 52 Tuson, M., He, M. & Anderson, K. V. Protein kinase A acts at the basal body of the primary cilium to prevent Gli2 activation and ventralization of the mouse neural tube. *Development* **138**, 4921-4930, doi:10.1242/dev.070805 (2011).
- 53 Lipinski, R. J., Bijlsma, M. F., Gipp, J. J., Podhaizer, D. J. & Bushman, W. Establishment and characterization of immortalized Gli-null mouse embryonic fibroblast cell lines. *BMC Cell Biol* **9**, 49, doi:10.1186/1471-2121-9-49 (2008).
- 54 Chen, W. *et al.* Activity-dependent internalization of smoothened mediated by beta-arrestin 2 and GRK2. *Science* **306**, 2257-2260, doi:10.1126/science.1104135 (2004).
- 55 Wolff, C., Roy, S. & Ingham, P. W. Multiple muscle cell identities induced by distinct levels and timing of hedgehog activity in the zebrafish embryo. *Curr Biol* **13**, 1169-1181, doi:10.1016/s0960-9822(03)00461-5 (2003).

969 56 Zhao, Z. *et al.* An essential role for Grk2 in Hedgehog signalling downstream of
970 Smoothed. *EMBO Rep* **17**, 739-752, doi:10.15252/embr.201541532 (2016).
971 57 Taipale, J. *et al.* Effects of oncogenic mutations in Smoothed and Patched can be
972 reversed by cyclopamine. *Nature* **406**, 1005-1009, doi:10.1038/35023008 (2000).
973 58 Rohatgi, R., Milenkovic, L., Corcoran, R. B. & Scott, M. P. Hedgehog signal transduction
974 by Smoothed: pharmacologic evidence for a 2-step activation process. *Proc Natl Acad Sci U S A* **106**, 3196-3201, doi:10.1073/pnas.0813373106 (2009).
975 59 Corbit, K. C. *et al.* Vertebrate Smoothed functions at the primary cilium. *Nature* **437**,
976 1018-1021, doi:10.1038/nature04117 (2005).
977 60 Rohatgi, R., Milenkovic, L. & Scott, M. P. Patched1 regulates hedgehog signaling at the
978 primary cilium. *Science* **317**, 372-376, doi:10.1126/science.1139740 (2007).
979 61 Kim, J., Kato, M. & Beachy, P. A. Gli2 trafficking links Hedgehog-dependent activation of
980 Smoothed in the primary cilium to transcriptional activation in the nucleus. *Proc Natl Acad Sci U S A* **106**, 21666-21671, doi:10.1073/pnas.0912180106 (2009).
981 62 Yogurtcu, O. N. & Johnson, M. E. Cytosolic proteins can exploit membrane localization
982 to trigger functional assembly. *PLoS Comput Biol* **14**, e1006031, doi:10.1371/journal.pcbi.1006031 (2018).
983 63 Riobo, N. A., Saucy, B., Dilizio, C. & Manning, D. R. Activation of heterotrimeric G
984 proteins by Smoothed. *Proc Natl Acad Sci U S A* **103**, 12607-12612, doi:10.1073/pnas.0600880103 (2006).
985 64 Shen, F., Cheng, L., Douglas, A. E., Riobo, N. A. & Manning, D. R. Smoothed is a
986 fully competent activator of the heterotrimeric G protein G(i). *Mol Pharmacol* **83**, 691-
987 697, doi:10.1124/mol.112.082511 (2013).
988 65 Mukhopadhyay, S. *et al.* The ciliary G-protein-coupled receptor Gpr161 negatively
989 regulates the Sonic hedgehog pathway via cAMP signaling. *Cell* **152**, 210-223, doi:10.1016/j.cell.2012.12.026 (2013).
990 66 Hwang, S. H. *et al.* The G protein-coupled receptor Gpr161 regulates forelimb formation,
991 limb patterning and skeletal morphogenesis in a primary cilium-dependent manner.
992 *Development* **145**, doi:10.1242/dev.154054 (2018).
993 67 Shimada, I. S. *et al.* Basal Suppression of the Sonic Hedgehog Pathway by the G-
994 Protein-Coupled Receptor Gpr161 Restricts Medulloblastoma Pathogenesis. *Cell Rep*
995 **22**, 1169-1184, doi:10.1016/j.celrep.2018.01.018 (2018).
996 68 Regard, J. B. *et al.* Activation of Hedgehog signaling by loss of GNAS causes
997 heterotopic ossification. *Nat Med* **19**, 1505-1512, doi:10.1038/nm.3314 (2013).
998 69 Moore, B. S. *et al.* Cilia have high cAMP levels that are inhibited by Sonic Hedgehog-
999 regulated calcium dynamics. *Proc Natl Acad Sci U S A* **113**, 13069-13074, doi:10.1073/pnas.1602393113 (2016).
1000 70 Low, W. C. *et al.* The decoupling of Smoothed from Galphai proteins has little effect
1001 on Gli3 protein processing and Hedgehog-regulated chick neural tube patterning. *Dev Biol* **321**, 188-196, doi:10.1016/j.ydbio.2008.06.014 (2008).
1002 71 Tschaikner, P., Enzler, F., Torres-Quesada, O., Aanstad, P. & Stefan, E. Hedgehog and
1003 Gpr161: Regulating cAMP Signaling in the Primary Cilium. *Cells* **9**, doi:10.3390/cells9010118 (2020).
1004 72 Pusapati, G. V. *et al.* G protein-coupled receptors control the sensitivity of cells to the
1005 morphogen Sonic Hedgehog. *Sci Signal* **11**, doi:10.1126/scisignal.aao5749 (2018).
1006 73 Meloni, A. R. *et al.* Smoothed signal transduction is promoted by G protein-coupled
1007 receptor kinase 2. *Mol Cell Biol* **26**, 7550-7560, doi:10.1128/MCB.00546-06 (2006).
1008 74 MacDonald, B. T., Tamai, K. & He, X. Wnt/beta-catenin signaling: components,
1009 mechanisms, and diseases. *Dev Cell* **17**, 9-26, doi:10.1016/j.devcel.2009.06.016 (2009).
1010 75 Metcalfe, C. & Bienz, M. Inhibition of GSK3 by Wnt signalling--two contrasting models. *J Cell Sci* **124**, 3537-3544, doi:10.1242/jcs.091991 (2011).
1011

1020 76 Nusse, R. & Clevers, H. Wnt/beta-Catenin Signaling, Disease, and Emerging
1021 Therapeutic Modalities. *Cell* **169**, 985-999, doi:10.1016/j.cell.2017.05.016 (2017).
1022 77 Steinhart, Z. & Angers, S. Wnt signaling in development and tissue homeostasis.
1023 *Development* **145**, doi:10.1242/dev.146589 (2018).
1024 78 Piao, S. *et al.* Direct inhibition of GSK3beta by the phosphorylated cytoplasmic domain
1025 of LRP6 in Wnt/beta-catenin signaling. *PLoS One* **3**, e4046,
1026 doi:10.1371/journal.pone.0004046 (2008).
1027 79 Cselenyi, C. S. *et al.* LRP6 transduces a canonical Wnt signal independently of Axin
1028 degradation by inhibiting GSK3's phosphorylation of beta-catenin. *Proc Natl Acad Sci U*
1029 *S A* **105**, 8032-8037, doi:10.1073/pnas.0803025105 (2008).
1030 80 Wu, G., Huang, H., Garcia Abreu, J. & He, X. Inhibition of GSK3 phosphorylation of
1031 beta-catenin via phosphorylated PPPSPXS motifs of Wnt coreceptor LRP6. *PLoS One*
1032 **4**, e4926, doi:10.1371/journal.pone.0004926 (2009).
1033 81 Stamos, J. L., Chu, M. L., Enos, M. D., Shah, N. & Weis, W. I. Structural basis of GSK-3
1034 inhibition by N-terminal phosphorylation and by the Wnt receptor LRP6. *Elife* **3**, e01998,
1035 doi:10.7554/eLife.01998 (2014).
1036 82 Ocbina, P. J., Tuson, M. & Anderson, K. V. Primary cilia are not required for normal
1037 canonical Wnt signaling in the mouse embryo. *PLoS One* **4**, e6839,
1038 doi:10.1371/journal.pone.0006839 (2009).
1039 83 Hauser, A. S., Attwood, M. M., Rask-Andersen, M., Schioth, H. B. & Gloriam, D. E.
1040 Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug*
1041 *Discov* **16**, 829-842, doi:10.1038/nrd.2017.178 (2017).
1042 84 Oprea, T. I. *et al.* Unexplored therapeutic opportunities in the human genome. *Nat Rev*
1043 *Drug Discov* **17**, 317-332, doi:10.1038/nrd.2018.14 (2018).
1044 85 Zimmermann, B., Schweinsberg, S., Drewianka, S. & Herberg, F. W. Effect of metal ions
1045 on high-affinity binding of pseudosubstrate inhibitors to PKA. *Biochem J* **413**, 93-101,
1046 doi:10.1042/BJ20071665 (2008).
1047 86 Myers, B. R. *et al.* Hedgehog pathway modulation by multiple lipid binding sites on the
1048 smoothened effector of signal response. *Dev Cell* **26**, 346-357,
1049 doi:10.1016/j.devcel.2013.07.015 (2013).
1050 87 Lu, T. W. *et al.* Two PKA RIalpha holoenzyme states define ATP as an isoform-specific
1051 orthosteric inhibitor that competes with the allosteric activator, cAMP. *Proc Natl Acad Sci*
1052 *U S A* **116**, 16347-16356, doi:10.1073/pnas.1906036116 (2019).
1053 88 Olsen, S. R. & Uhler, M. D. Affinity purification of the C alpha and C beta isoforms of the
1054 catalytic subunit of cAMP-dependent protein kinase. *J Biol Chem* **264**, 18662-18666
1055 (1989).
1056 89 Walker, C. *et al.* Cushing's syndrome driver mutation disrupts protein kinase A allosteric
1057 network, altering both regulation and substrate specificity. *Sci Adv* **5**, eaaw9298,
1058 doi:10.1126/sciadv.aaw9298 (2019).
1059 90 Olivieri, C. *et al.* Defective internal allosteric network imparts dysfunctional
1060 ATP/substrate-binding cooperativity in oncogenic chimera of protein kinase A. *Commun*
1061 *Biol* **4**, 321, doi:10.1038/s42003-021-01819-6 (2021).
1062 91 Walsh, D. A. & Ashby, C. D. Protein kinases: aspects of their regulation and diversity.
1063 *Recent Prog Horm Res* **29**, 329-359, doi:10.1016/b978-0-12-571129-6.50012-9 (1973).
1064 92 Delaglio, F. *et al.* NMRPipe: a multidimensional spectral processing system based on
1065 UNIX pipes. *J Biomol NMR* **6**, 277-293, doi:10.1007/BF00197809 (1995).
1066 93 Lee, W., Tonelli, M. & Markley, J. L. NMRFAM-SPARKY: enhanced software for
1067 biomolecular NMR spectroscopy. *Bioinformatics* **31**, 1325-1327,
1068 doi:10.1093/bioinformatics/btu830 (2015).
1069 94 Williamson, M. P. Using chemical shift perturbation to characterise ligand binding. *Prog*
1070 *Nucl Magn Reson Spectrosc* **73**, 1-16, doi:10.1016/j.pnmrs.2013.02.001 (2013).

1071 95 Hansen, J. N., Rassmann, S., Stuvén, B., Jurisch-Yaksi, N. & Wachten, D. CiliaQ: a
1072 simple, open-source software for automated quantification of ciliary morphology and
1073 fluorescence in 2D, 3D, and 4D images. *Eur Phys J E Soft Matter* **44**, 18,
1074 doi:10.1140/epje/s10189-021-00031-y (2021).
1075 96 Eccles, R. L. *et al.* Bimodal antagonism of PKA signalling by ARHGAP36. *Nat Commun*
1076 **7**, 12963, doi:10.1038/ncomms12963 (2016).
1077 97 Han, P., Sonati, P., Rubin, C. & Michaeli, T. PDE7A1, a cAMP-specific
1078 phosphodiesterase, inhibits cAMP-dependent protein kinase by a direct interaction with
1079 C. *J Biol Chem* **281**, 15050-15057, doi:10.1074/jbc.M601333200 (2006).
1080 98 van Eeden, F. J. *et al.* Mutations affecting somite formation and patterning in the
1081 zebrafish, *Danio rerio*. *Development* **123**, 153-164 (1996).
1082 99 Stickney, H. L., Barresi, M. J. & Devoto, S. H. Somite development in zebrafish. *Dev Dyn*
1083 **219**, 287-303, doi:10.1002/1097-0177(2000)9999:9999<::AID-DVDY1065>3.0.CO;2-A
1084 (2000).
1085 100 Chen, W., Burgess, S. & Hopkins, N. Analysis of the zebrafish smoothened mutant
1086 reveals conserved and divergent functions of hedgehog activity. *Development* **128**,
1087 2385-2396 (2001).
1088