Lung adenocarcinoma cell-derived exosomal miR-328 promote osteoclastogenesis by targeting Nrp2

Chengcheng Zhang^{12#}, Jingru Qin^{1#}, Lu Yang^{1#}, Zhiyao Zhu¹, Xinle Qian¹, Jia Yang¹, Wan Su¹, Fengying Wang², Wenlian Chen², Haibin Deng^{1*} & Zhongqi Wang^{1*}

¹ Department of Medical Oncology, Longhua Hospital affiliated to Shanghai University of Traditional Chinese Medicine, 725 Wanpingnan Road, Shanghai 200032, China.

² Cancer Institute, Longhua Hospital affiliated to Shanghai University of Traditional Chinese Medicine, 725 Wanpingnan Road, Shanghai 200032, China.

Corresponding Author:

Haibin Deng, MD, Department of Medical Oncology, Longhua Hospital affiliated to Shanghai University of Traditional Chinese Medicine, 725 Wanpingnan Road, Shanghai 200032, China. (Tel: 008613661578200, E-mail: 13661578200@139.com)

Zhongqi Wang, MD, Department of Medical Oncology, Longhua Hospital affiliated to Shanghai University of Traditional Chinese Medicine, 725 Wanpingnan Road, Shanghai 200032, China. (Tel: 008618917763093, E-mail: aledx@sina.com)

#These authors contributed equally to this work, and both should be considered first author.

Abstract

Bone metastasis of lung cancer and detailed mechanisms are still elusive, and the roles of exosomes derived from lung adenocarcinoma cells in this process have attracted much attention. In this study, we found that lung adenocarcinoma cell-derived exosomes (LCC-Exos) promoted osteogenesis and bone resorption *in vitro*. Furthermore, LCC-Exos target bone *in vivo* and promoted bone resorption *in vivo*. Mechanistically, LCC-Exosomal miR-328 promoted bone resorption by targeting Nrp2 and LCC-Exos^{miR-328 Inhibitors} inhibited bone resorption *in vivo*. Thus, LCC-Exosomal miR-328 promote osteoclastogenesis by targeting Nrp2 and LCC-Exos^{miR-328 Inhibitors} may serve as a potential nanomedicine for the treatment of bone metastasis.

Introduction

Bone metastasis is the second most frequent site of lung cancer metastasis¹ and most lung cancer bone metastasis can be classified into osteolytic bone metastasis², which can cause pain, pathological, spinal instability, spinal cord compression and hypercalcemia and result in a decline in the quality of life^{3,4}. However, lung cancer cells can not directly lead to osteolytic metastasis by proteolytic enzymes. Instead, lung cancer cells derived circulating miRNA and exosomes played important roles by activating osteoclasts^{5,6}. Thus, we focus on the impact of lung cancer derived exosomes on osteoclast differentiation.

Exosomes are small cup or sphere-shaped vesicles, and their diameters primarily ranged from 80 nm to 150 nm, that contains mRNAs, miRNAs⁷, lipids⁸ and proteins⁹. It has been reported that the role of exosomes in cellular-communication by the transfer of proteins¹⁰, bioactive lipids and miRNAs¹¹. Lung cancer derived exosomes have been attracted attention for its role in bone metastasis and prognosis¹². Furthermore, exosomes may be potentially used as biomarkers, nanoparticle drugs and carriers of antitumor drugs. Recently, numerous investigations indicated that exosomal miRNAs played important roles in tumor growth, invasion and metastasis¹³. Therefore, finding the key exosomal miRNAs in the bone metastasis could be beneficial for treatment of bone metastasis and exosomal miRNAs may serve as a potential therapeutic target for bone metastasis¹⁴.

In this study, we found that A549 cells derived exosomes is an important extracellular vesicle in osteoclast differentiation and in bone metastasis. Furthermore, we found that A549-Exos promoted bone resorption in vitro and in vivo. We furthermore explored its mechanism and found that A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2. Therefore, we generated A549-Exos loaded in miR-328 inhibitors and found that bone resorption was prevented by injecting A549-Exos^{miR-328 Inhibitors}. Therefore, we focused on the roles of A549 cells derived exosomes in bone resorption in this study.

Results

Characteristics of the isolated A549-Exos.

To investigate the role of lung adenocarcinoma cell in osteoclastogenesis, we first determined whether lung adenocarcinoma cell could affect the differentiation of osteoclasts, RAW264.7 cells were co-cultured with A549 cells in a transwell system. The differentiation of osteoclasts was enhanced when co-culturing with A549 cells (Figure 1A). Then, we performed TEM, Dynamic light scattering analysis and Western Blot analysis to characterize the exosomes (Exos) derived from A549. A549-Exos presented a cup or sphere-shaped morphology and their diameters primarily ranged from 80 nm to 150 nm (Figure 1B, C). Western Blot analysis showed that A549-Exos expressed exosomal marker proteins including CD9 and TSG101 (Figure 1D). Thus, these data suggested that Osteoclastic differentiation was enhanced when co-culturing with A549 cells and A549-Exos showed characteristic features of exosomes.

A549-Exos promoted osteoclast differentiation and bone resorption in vitro.

To investigate the role of A549-Exos in osteoclastogenesis in vitro. A549-Exos were labeled with lipophilic dye Dio and assessed whether A549-Exos could be transported to Raw264.7 cells. The fluorescence microscope revealed that A549-Exos could be successfully absorbed by Raw264.7 cells (Figure 2A). Then, we treated Raw264.7 cells with A549-Exos or an equal volume of Phosphate Buffer Saline (PBS). osteoclastic differentiation was upregulated as the concentration of exosomes derived from A549 cells increased (Figure 2B, C). Furthermore, we found that mRNA expression of Ctsk and Traf-6 were upregulated by A549-Exos compared to its' negative control at day 7 (Figure 2D, E). Taken together, these data suggested that A549-Exos promoted osteoclast differentiation in vitro.

A549-Exos target bone in vivo.

To determine the bio-distribution of A549-Exos in vivo. The tissue distribution of DID-labeled A549-Exos (A549Exos-DID) was examined by fluorescence molecular tomography (FMT) imaging system. A549Exos-DID was injected into the tail vein of 8-week-old male mice for 3 h, 6 h or 12 h. 3 hours after intravenous injection, the

major fluorescence signals were detected in the liver and lungs and the signals can be detected in the limbs. After 6 h, the major fluorescence signals were still accumulated in the liver and lung. Significantly, fluorescence signals in the limbs were enhanced, but still weaker than those in the liver and lung. After 12 hours, fluorescence signals still can be detected in the limbs, and fluorescence signals faded from liver and lungs. The fluorescence signals can be rarely detected in the kidney. Taken together, A549-Exos could be targeted for bone tissues in vivo (Figure 3A).

A549-Exos promoted bone resorption in vivo.

To investigate the role of A549-Exos in vivo, we injected A549-Exos or an equal volume of PBS into tail vein of 8-week-old male mice third per week for 12 weeks. Microcomputed tomography (micro-CT) showed significantly lower trabecular bone volume, number, thickness, and higher trabecular separation in those mice injected with A549-Exos compared to their controls (Figure 4A, B).

A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2.

Since exosomes may concentrate large numbers of miRNAs, which can regulate gene expression by binding 3'-untranslated regions. MiR-328 has been reported to promote osteoclast differentiation¹⁶. Subsequently, we examined the expression levels of miR-328 in the exosomes, and qRT-PCR showed that A549 cells derived exosomes had significantly higher expression of miR-328 in comparison with TC-1 cells derived exosomes (Figure 5A).

To investigate osteoclastogenesis regulated by miR-328 in vitro. RAW264.7 cells were transfected with miR-328 mimics or inhibitors. The miR-328 expression levels were significantly upregulated by miR-328 mimic transfection and markedly inhibited by miR-328 inhibitor transfection (Figure 5B). Furthermore, overexpression of miR-328 suppress endogenous expression of OPG protein level (Figure 5C). Consistent with the results of OPG protein expression, we found that miR-328 mimics enhanced TRAP staining, whereas miR-328 inhibitors weakened TRAP staining (Figure 5D). Overall, these data showed that osteoclast differentiation was promoted by miR-328 in vitro.

To investigate target mRNAs of miR-328 in osteoclast differentiation process, we have checked the miRNA-binding site in Targetscan and found that miR-328 binding site on Nrp2 3'-UTR. Previously, Nrp2 deficiency has been reported to promote osteoclast differentiation and reduced osteoblast numbers¹⁷. Thus, luciferase reporter constructs containing the WT or mutated predicted miRNA-binding sites of Nrp2 were constructed. Then, RAW264.7 cells were transfected with WT-Nrp2-3'-UTR or MUT-Nrp2-3'-UTR with miR-328 mimics or negative controls and the luciferase activity of the Nrp2 reporter gene was inhibited by miR-328 mimics and the Nrp2 reporter gene was promoted by miR-328 inhibitors, whereas MUT-Nrp2-3'-UTR prevented this inhibition (Figure 5F). In conclusion, these data showed that miR-328 directly targets Nrp2 and inhibited Nrp2 expression in osteoclast cells.

A549-Exos^{miR-328 Inhibitors} inhibited bone resorption *in vivo*.

To investigate the therapeutic potential of miR-328 on bone metastasis, A549-Exos^{miR-328 Inhibitors} was generated. A549 cells transfected with miR-328 inhibitors derived exosomes were labeled with lipophilic dye Dio and assessed whether A549-Exos could carry miR-328 inhibitors and be transported to Raw264.7 cells. The fluorescence microscope revealed that A549-Exos^{miR-328 Inhibitors} could be successfully absorbed by Raw264.7 cells (Figure 6A). Then, we injected A549-Exos or A549-Exos^{miR-328 Inhibitors} into tail vein of 8-week-old male mice third per week for 12 weeks. Mice treated with A549-Exos^{miR-328 Inhibitors} showed increased trabecular bone volume, number, and thickness, and decreased trabecular separation relative to A549-Exos group, which showed that A549-Exos^{miR-328 Inhibitors} could alleviate bone loss caused by the bone metastasis (Figure 6B, C). In conclusion, these data suggested that intravenous injection of A549-Exos^{miR-328 Inhibitors} inhibited bone resorption *in vivo* and alleviate the osteopenia caused by bone metastasis.

Discussion

In this study, we found that osteoclast differentiation was enhanced when co-culturing with A549 cells. A549-Exos promote osteoclast differentiation in vitro. Furthermore, A549-Exos could be targeted for bone tissues and promoted bone resorption in vivo. Moreover, we found that A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2. In addition, A549-Exos^{miR-328 Inhibitors} treatment in vivo ameliorated

bone loss and prevented pathological fractures.

Previously, several cancer cells derived exosomes have been reported to be home to the bone microenvironment and cause metastatic lesions¹⁸. As murine prostate cancer line TRAMP-C1 inhibits the differentiation of the monocytic cell line RAW264.7 into osteoclasts¹⁹, non-small cell lung cancer (NSCLC) cells secrete exosomes promoted osteoclast differentiation of murine RAW264.7 cells by activation of EGFR phosphorylation²⁰, exosomes released by multiple myeloma cells increase the viability and migration of osteoclast precursors, through the increasing of CXCR4 expression²¹ and extracellular vesicles isolated from PC3 culture medium are internalized into osteoclast precursors and osteoblasts²². In this study, A549 cell-derived exosomal miR-328 is the regulator of bone resorption and A549-Exos^{miR-328 Inhibitors} could ameliorate bone loss. Thus, A549 cells derived exosomes could load small molecule drugs and target bone tissues. Since a set of tumor cells derived exosomes could target bone tissues, we hypothesized that these exosomes can also be used as targeted drug carriers for the corresponding primary tumors. Our work might provoke interesting future works regarding the relationship between bone metastasis and tumor cells derived exosomes leading to new potential therapeutic targets.

A set of exosomal miRNAs has been reported to be correlated with bone formation or bone resorption²³. For example, Exosomal miR-141-3p regulates osteoblast activity to promote the osteoblastic metastasis of prostate cancer²⁴, overexpression of hsa-miR-940 in tumor exosomes and its ability to ultimately lead to extensive osteoblastic lesions in the resulting tumor²⁵ and mi-192 cargo within exosome like vesicle transfer influences metastatic bone colonization²⁶. These miRNAs may also

serve as therapeutic potential targets for the treatment of bone metastases from lung cancers, which requires further investigation.

Methods and materials

Cell culture

A549 cell line was purchased from the American Type Culture Collection (ATCC) and cultured in Dulbecco's modified Eagle's medium (DMEM) (Gibco, USA), supplemented with 10% fetal bovine serum (FBS) (Gibco, USA). The Raw264.7 cell line was cultured in DMEM (Gibco) supplemented with 10% FBS. We cultured cells at 37 °C in a 5% CO₂ humidified incubator.

Exosome isolation and characterization

A549 cells were cultured in fresh complete medium containing EVs-free FBS (Thermo, USA) for 48 h. Then, we collected the conditioned medium and centrifuged them at 1500 g for 15 minutes at 4°C. After filtering through a 0.22 mm filter (Millipore, Billerica, USA). The supernatant was ultracentrifuged twice at 110 000 g for 1 hour at 4°C. Then, we resuspended the pellets in a right amount of PBS and stored A549-Exos at –80°C for later use. We observed the morphology of A549-Exos by transmission electron microscope (TEM). The size distribution of A549-Exos was determined by using Nanosizer™ instrument (Malvern Instruments, Malvern, UK). The expression of exosomal surface markers (CD9, HSP70) was analyzed by western blot (WB).

Western blot (WB) analysis

Cells or A549-Exos were lysed in a RIPA lysis buffer supplemented with protease inhibitor cocktail (CST). Then, we quantified total protein concentration by using the BCA protein assay kit (Thermo, USA). Western blotting was performed as previously described¹⁵. Primary antibodies were used in this study as follows: CD9 (CST, USA), HSP70 (CST, USA), GAPDH (CST, USA) and OPG (CST, USA). All the secondary antibodies (1:5000) were acquired from Abcam.

In vitro Osteoclastogenesis assays of RAW264.7 cells

We seeded RAW264.7 cells on 48-well plates ((1×10^4)) in D-MEM (Gibco, USA) supplemented with 10% FBS (Gibco, USA). M-CSF (30ng/ml) and RANKL (100ng/ml) were used to promote osteoclast maturation for 7 days. Then, we

performed TARP-staining according to general protocols. We considered TRAP positive cells (more than 5 nucleus) considered as mature osteoclasts.

Luciferase reporter assay

We constructed the Nrp2 3'-UTR luciferase reporter construct by amplifying the mouse Nrp2 mRNA 3'-UTR sequence (including the predicted miR-328 binding site) by PCR. Then, the PCR products were purified and cloned into the XbaI site of the pGL3-promoter vector (Promega) immediately downstream of the stop codon of luciferase. The Nrp2 mutations were prepared by using a Quick-Change Site-Directed Mutagenesis Kit (Stratagene) to get MUT-pGL3-Nrp2, which is confirmed by sequencing. RAW264.7 cells were transfected with either WT or mutant pGL3 construct and 40 ng of pRL-TKRenilla-luciferase plasmid and miR-328 mimics or inhibitors for 48 h. Luciferase activities were measured as previously described¹⁵.

Real-time PCR (RT-PCR)

Total RNA of RAW264.7 cells or A549-Exos were isolated by TRIzol reagent (Life Technologies, USA) and synthesized complementary DNAs (cDNAs). Then, Quantitative reverse transcriptase PCR (qRT-PCR) was performed as previously described¹⁵. The primer sequences used for qRT-PCR were as follows: miR-328: forward, 5'-CGG GCC TGG CCC TCT CTG CC-3'; reverse, 5'-CAG CCA CAA AAG AGC ACA AT-3'; Ctsk: forward, 5'-CTC GGC GTT TAA TTT GGG AGA-3'; reverse, 5'-TCG AGA GGG AGG TAT TCT GAG T-3'; TRAF-6: forward, 5'-AAG GTG GTG GCG TTA TAC TGC-3'; reverse, 5'-CTG GCA CAG CGG ATG TGA G-3' and GAPDH: forward, 5'-AAT GGA TTT GGA CGC ATT GGT-3'; reverse, 5'-TTT GCA CTG GTA CGT GTT GAT-3';

Animal studies

All animal experiments were conducted in accordance with the National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals, with the approval of Longhua hospital (No.LHERAW-19038). C57BL/6 mice (6-8 weeks old) were obtained from the Animal Center of Longhua hospital. The mice were housed in barrier housing conditions in the Animal Center of Longhua hospital. For delivery of A549-Exos or A549-Exos^{miR-328 Inhibitors}, 100ng/ml in 1 ml PBS was injected into tail

vein once every 3 days for 4 weeks. The mice were sacrificed at 4 weeks after the treatment.

Micro-CT analysis

femur samples dissected from mice were scanned and analyzed through micro-CT (Quantum GX, PE, USA). We performed the micro-CT scans under the same conditions: voltage 100 kV, current 80 μ A, spatial resolution 12 μ m, scanning 500 continuous sections. The data was collected and analyzed automatically through computer software (Skyscan) to analyze the number of trabecular bones (Tb.N), trabecular bone thickness (Tb.Th), trabecular bone space (Tb.Sp), bone volume fraction (BV/TV) and three-dimensional reconstruction.

Exosomes labeling and tracking in vivo.

All animal experiments were conducted in accordance with the National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals, with the approval of Longhua hospital (No.LHERAW-19038). To visualize the target organ and the metabolism process of exosomes, we labelled A549-Exos according to the instructions of the manufacturer (Thermo, USA). First, we incubated 100ug of exosomes with Vybrant DID (1:1000 in PBS) in the dark for 30 mins, and labelled exosomes were washed in PBS, with centrifugation at $100,000 \times g$ for 60 min. Then, the labelled exosomes were injected into the tail vein of mice (8 weeks old, n=3). The mice were harvested at 3, 6, 12 hours after injection. The mean intensities for DID signals were quantified from four random visual fields for each section.

Statistical analysis

All data are shown as means $\pm$ standard deviation (SD) from at least three independent experiments. Comparisons were performed using t test and one-way ANOVA as appropriate. $P < 0.05$ was considered statistically significant.

Acknowledgements

This study was supported in part by National Natural Science Foundation of China (No. 81904129), Innovation Program of Shanghai Municipal Education Commission (No. 2017-01-07-00-10-E00064), Shanghai Hospital Development Center (SHDC2020CR4050), Shanghai Medical Innovation Research Project of "Science and Technology Innovation Action Plan" (No.20Y21902300), Science and Technology Innovation Project of Longhua Hospital (NO.LYTD-33).

Competing interests: The authors indicate no potential conflicts of interest.

Author contributions:

Conceived and designed the review: HBD and ZQW. Collected the data: ZYZ and XLQ. Analyzed the data: JY and WS. Supervised the study: FYW and WLC. Wrote the paper: CCZ, JRQ and LY.

References

1. Fornetti, J., Welm, A.L. & Stewart, S.A. Understanding the Bone in Cancer Metastasis. *J Bone Miner Res* **33**, 2099-2113 (2018).
2. Zhang, L. & Gong, Z. Clinical Characteristics and Prognostic Factors in Bone Metastases from Lung Cancer. *Med Sci Monit* **23**, 4087-4094 (2017).
3. Altorki, N.K., *et al.* The lung microenvironment: an important regulator of tumour growth and metastasis. *Nat Rev Cancer* **19**, 9-31 (2019).
4. Wu, S., Pan, Y., Mao, Y., Chen, Y. & He, Y. Current progress and mechanisms of bone metastasis in lung cancer: a narrative review. *Transl Lung Cancer Res* **10**, 439-451 (2021).
5. Yin, J.J., Pollock, C.B. & Kelly, K. Mechanisms of cancer metastasis to the bone. *Cell Res* **15**, 57-62 (2005).
6. Xu, S., *et al.* Serum microRNA-139-5p is downregulated in lung cancer patients with lytic bone metastasis. *Oncol Rep* **39**, 2376-2384 (2018).
7. Hu, Y., *et al.* Exosomes from human umbilical cord blood accelerate cutaneous wound healing through miR-21-3p-mediated promotion of angiogenesis and fibroblast function. *Theranostics* **8**, 169-184 (2018).
8. Record, M., Silvente-Poirot, S., Poirot, M. & Wakelam, M.J.O. Extracellular vesicles: lipids as key components of their biogenesis and functions. *J Lipid Res* **59**, 1316-1324 (2018).
9. Chen, C.Y., *et al.* Exosomal DMBT1 from human urine-derived stem cells facilitates diabetic wound repair by promoting angiogenesis. *Theranostics* **8**, 1607-1623 (2018).
10. Zeelenberg, I.S., *et al.* Targeting tumor antigens to secreted membrane vesicles in vivo induces efficient antitumor immune responses. *Cancer Res* **68**, 1228-1235 (2008).
11. Kumar, S. & Reddy, P.H. Are circulating microRNAs peripheral biomarkers for Alzheimer's

- disease? *Biochim Biophys Acta* **1862**, 1617-1627 (2016).
12. Li, F.X., *et al.* Role of tumor-derived exosomes in bone metastasis. *Oncol Lett* **18**, 3935-3945 (2019).
13. Dai, J., *et al.* Exosomes: key players in cancer and potential therapeutic strategy. *Signal Transduct Target Ther* **5**, 145 (2020).
14. Zhang, J. & Wu, J. The Potential Roles of Exosomal miR-214 in Bone Metastasis of Lung Adenocarcinoma. *Front Oncol* **10**, 611054 (2020).
15. Wang, R., *et al.* miR-143 promotes angiogenesis and osteoblast differentiation by targeting HDAC7. *Cell Death Dis* **11**, 179 (2020).
16. Chen, J., *et al.* Identification of suitable reference gene and biomarkers of serum miRNAs for osteoporosis. *Sci Rep* **6**, 36347 (2016).
17. Verlinden, L., *et al.* Nrp2 deficiency leads to trabecular bone loss and is accompanied by enhanced osteoclast and reduced osteoblast numbers. *Bone* **55**, 465-475 (2013).
18. Rossi, M., Battafarano, G., D'Agostini, M. & Del Fattore, A. The Role of Extracellular Vesicles in Bone Metastasis. *Int J Mol Sci* **19**(2018).
19. Karlsson, T., Lundholm, M., Widmark, A. & Persson, E. Tumor Cell-Derived Exosomes from the Prostate Cancer Cell Line TRAMP-C1 Impair Osteoclast Formation and Differentiation. *PLoS One* **11**, e0166284 (2016).
20. Taverna, S., *et al.* Amphiregulin contained in NSCLC-exosomes induces osteoclast differentiation through the activation of EGFR pathway. *Sci Rep* **7**, 3170 (2017).
21. Raimondi, L., *et al.* Involvement of multiple myeloma cell-derived exosomes in osteoclast differentiation. *Oncotarget* **6**, 13772-13789 (2015).
22. Inder, K.L., *et al.* Cavin-1/PTRF alters prostate cancer cell-derived extracellular vesicle content and internalization to attenuate extracellular vesicle-mediated osteoclastogenesis and osteoblast proliferation. *J Extracell Vesicles* **3**(2014).
23. Luga, V., *et al.* Exosomes mediate stromal mobilization of autocrine Wnt-PCP signaling in breast cancer cell migration. *Cell* **151**, 1542-1556 (2012).
24. Ye, Y., *et al.* Exosomal miR-141-3p regulates osteoblast activity to promote the osteoblastic metastasis of prostate cancer. *Oncotarget* **8**, 94834-94849 (2017).
25. Hashimoto, K., *et al.* Cancer-secreted hsa-miR-940 induces an osteoblastic phenotype in the bone metastatic microenvironment via targeting ARHGAP1 and FAM134A. *Proc Natl Acad Sci U S A* **115**, 2204-2209 (2018).
26. Valencia, K., *et al.* miRNA cargo within exosome-like vesicle transfer influences metastatic bone colonization. *Mol Oncol* **8**, 689-703 (2014).

Figure legends

Figure 1 Characteristics of the isolated A549-Exos. (A) Representative images of TRAP staining osteoclasts. Scale bar: 100 μ m. (B) Representative TEM image of A549-Exos (scale bar = 200 nm). (C) Representative NTA results showing the size distribution of A549-Exos. (D) Detection of A549-Exos surface markers (CD9 and HSP70) by Western Blot analysis.

Figure 2 A549-Exos promoted osteoclast differentiation and bone resorption in vitro. (A) Fluorescence microscopy analysis of RAW264.7 cells treated with DiO-labeled A549-Exos for 2h. (B) Representative images of TRAP staining of mature osteoclasts treated with different concentrations of A549-Exos. Scale bar: 100 μ m. (C) Quantitative analysis of positive TRAP staining osteoclasts in Figure 2B (**P<0.01). (D) Relative levels of Ctsk (left) and TRAF-6 (right) mRNA expression in Figure 2B (**P<0.01).

Figure 3. A549-Exos target bone in vivo. (A) Representative FMT images of the near-infrared fluorescence signals in organs isolated from mice administered with PBS, A549-Exos for different periods of time (3 hours, 6 hours and 12 hours).

Figure 4. A549-Exos promoted bone resorption in vivo. (A) Representative micro-CT images in femora from A549-Exos mice and negative control group (n = 8 per group). Scale bar: 100 μm. (B) Quantitative analysis of trabecular microarchitecture from A549-Exos group and negative control group. BV/TV, trabecular bone volume per tissue volume; Tb. Th, trabecular thickness; Tb. Sp, trabecular separation; Tb. N, trabecular number (n=8 per group) (*P<0.05).

Figure 5. A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2. (A) miR-328 relative expressions in N.C. and LCC-Exos. (B) miR-328 levels in N.C., miR-328 mimics, I.N.C. and miR-328 inhibitors were measured by qRT-PCR. (C) Western blot (WB) analysis of OPG and GAPDH in RAW264.7 cells transfected with miR-328 mimics and miR-328 or their respective negative controls. (D) Representative images of TRAP staining of mature osteoclasts treated with N.C., miR-328 mimics, I.N.C. and miR-328 inhibitors. (E) Nrp2 might be the target of miR-328. (F) Raw264.7 were transfected with a luciferase reporter carrying WT or MUT-Nrp2 3'-UTR in the miR-328 mimics (F left) and miR-328 (F right) or their respective negative controls. * $p < 0.05$, ** $p < 0.01$. Data are representative of at least three independent experiments.

Figure 6. A549-Exos^{miR-328} Inhibitors inhibited bone resorption *in vivo*. (A) Fluorescence microscopy analysis of RAW264.7 cells treated with DiO-labeled A549-Exos transfected with miR-328 inhibitors for 2h (B) Representative micro-CT images in femora from A549-Exos group and A549-Exos^{miR-328} Inhibitors group (n = 6 per group). (C) Quantitative analysis of trabecular microarchitecture from A549-Exos group and A549-Exos^{miR-328} Inhibitors group (n=8 per group) (*P<0.05).