1 **Cytokinin oxidase/dehydrogenase family genes exhibit functional divergence and overlap in rice growth**
2 **and development**

3 Chenyu Rong¹, Yuexin Liu¹, Zhongyuan Chang¹, Ziyu Liu¹, Yanfeng Ding^{1,2,3} and Chengqiang Ding^{1,2,3,*}

4 ¹College of Agriculture, Nanjing Agricultural University, Nanjing 210095, People's Republic of China

5 ²Key Laboratory of Crop Physiology Ecology and Production Management, Ministry of Agriculture, Nanjing

6 210095, People's Republic of China

7 ³Collaborative Innovation Center for Modern Crop Production co-sponsored by Province and Ministry, Nanjing

8 210095, People's Republic of China

9 *Correspondence: Chengqiang Ding, email: dingcq@njau.edu.cn.

10 **Running title:** The important functions of OsCKX family genes in rice

11 **Highlight:** The *osckx4 osckx9* double mutant had significantly more tillers, whereas the *osckx1 osckx2* double
12 mutant had the opposite phenotypic change.

13

14 ABSTRACT

15 Cytokinin has important functions during plant growth and development; hence, many researchers have
 16 extensively studied cytokinin biosynthesis and degradation. Cytokinin oxidase/dehydrogenases (CKXs) are a
 17 group of enzymes that regulate oxidative cleavage to maintain cytokinin homeostasis. In rice, 11 *OsCKX* genes
 18 have been identified to date; however, most of their functions remain unknown. Here, we comprehensively
 19 analyzed the expression patterns of *OsCKX* genes and their functions. Using CRISPR/Cas9 technology, we
 20 constructed nine *osckx* mutants to determine the *OsCKX* function in rice development. The results revealed that
 21 the single *osckx* and higher-order *osckx4 osckx9* mutant lines showed functional overlap and
 22 subfunctionalization. Notably, we found that the *osckx1 osckx2* and *osckx4 osckx9* double mutants displayed
 23 contrasting phenotypic changes in tiller number, culm diameter, and panicle size as compared to the wild-type.
 24 Moreover, we identified several genes that were significantly expressed in *osckx4* and *osckx9* single and double
 25 mutant plants. Many differentially expressed genes were found to be associated with auxin, cytokinin, and
 26 nitrogen pathways. Therefore, our findings provide new insights into the functions of *OsCKX* genes in rice
 27 growth, which may be used as a foundation for future, studies aimed at improving rice yield and initiating green
 28 production.

29 **Key words:** Cytokinin, expression pattern, *OsCKX*, panicle, phenotype, rice, tiller

30

31

32 INTRODUCTION

33 Cytokinin controls plant growth and development by promoting plant cell division, growth, and differentiation.
 34 Specifically, it is responsible for stem enlargement and differentiation, organ development and structure,
 35 nutrition absorption, senescence, and stress response. Cytokinin biosynthesis starts from the ADP, ATP, or tRNA
 36 catalyzed by different adenosine phosphate isopentenyltransferases and activated by LONELY GUY enzymes
 37 (Kurakawa *et al.*, 2007; Osugi *et al.*, 2017). Previous studies reported that in rice, an overexpression of
 38 adenosine phosphate isopentenyltransferase inhibited root formation (Sakamoto *et al.*, 2006), while *lonely guy*
 39 mutants developed smaller panicles with reduced branches and floral organ defects (Kurakawa *et al.*, 2007).

40 After performing its function, cytokinin is inactivated by certain enzymes such as cytokinin
 41 oxidase/dehydrogenases (CKXs) that induce cytokinin oxidative cleavage, which is an irreversible process.
 42 Cytokinin can also be inactivated by combining with sugars, including *O*-glucosyltransferase (reversible process)
 43 and *N*-glucosyltransferase (partly irreversible process) as the storage form of cytokinin (Hosek *et al.*, 2020;
 44 Kudo *et al.*, 2012). However, this inactivation process occurs at different stages of plant development in various
 45 tissues and employs several enzymes with similar functions, most of which remain unknown. CKXs are
 46 essential for maintaining cytokinin homeostasis during plant growth and development. Therefore, extensive
 47 research on CKXs has been conducted and is still ongoing. Each plant species has a group of *CKX* genes, for
 48 example, *Arabidopsis* has seven, rice has 11, and maize has 13. Most CKXs in maize and *Arabidopsis* have
 49 putative N-terminal secretory peptides that assist in locating the endoplasmic reticulum (ER) (Niemann *et al.*,
 50 2018; Zalabak *et al.*, 2016). Previous studies have discovered several functions of CKXs in *Arabidopsis* and
 51 other crops. In *Arabidopsis*, the overexpression of *AtCKX1* and *AtCKX2* resulted in smaller shoots, larger roots,
 52 and smaller shoot apical meristems (SAMs), while *AtCKX7* overexpression resulted in shorter primary roots
 53 (Kollmer *et al.*, 2014; Werner *et al.*, 2001). Furthermore, *ckx3 ckx5* mutants developed larger SAMs and more
 54 siliques (Bartrina *et al.*, 2011). On the other hand, sextuple *ckx3 ckx5* mutants of oilseed rape showed larger and

55 more active inflorescence meristems (Schwarz *et al.*, 2020). In barley, *HvCKX1* downregulation resulted in more
56 spikes, more grains per spike, and a higher 1,000-grain weight (Zalewski *et al.*, 2010). *TaCKX6-D1* was found to
57 be associated with grain filling and grain size in wheat (Zhang *et al.*, 2012).

58 In rice, *OsCKX2/Gn1a* was the first *CKX* gene identified. Rice varieties with low *OsCKX2* expression
59 yielded more grains per panicle, while *OsCKX2* downregulation or *osckx2* mutation produced more tillers, more
60 grains per panicle, and heavier grains (Ashikari *et al.*, 2005; Yeh *et al.*, 2015). In contrast, *OsCKX4*
61 overexpression reduced the tiller number, grain number per panicle, grain weight, and plant height, while it
62 increased the number of roots (Gao *et al.*, 2014). On the other hand, *osckx9* mutants developed more tillers and
63 smaller panicles, while *OsCKX9* overexpression mutants displayed more tillers, shorter culms, smaller panicles,
64 and lower setting rates (Duan *et al.*, 2019). *OsCKX11* was found to regulate leaf senescence and grain number,
65 while *osckx11* mutation resulted in the increased number of spikelets per panicle and tillers (Zhang *et al.*, 2020).
66 The expression of *CKX* genes is regulated by several transcription factors. For example, *OsCKX2* expression is
67 controlled by zinc finger transcription factor DROUGHT AND SALT TOLERANCE and VIN3-LIKE 2 (Li *et*
68 *al.*, 2013; Yang *et al.*, 2018), while the rice NAC family transcription factor OsNAC2 regulates *OsCKX4*
69 expression (Gao *et al.*, 2014; Mao *et al.*, 2020).

70 *OsCKX* genes play an important role in the crosstalk between cytokinin and other hormones by regulating
71 cytokinin content in plant tissues, maintaining plant hormone homeostasis, and controlling plant growth and
72 development. Most *OsCKX* genes are upregulated by exogenous cytokinins, such as *trans*-zeatin (tZ),
73 *N*⁶-(Δ^2 -isopentenyl) adenine (iP) and 6-benzylaminopurine (Duan *et al.*, 2019). *OsCKX4* can also be induced by
74 auxin, which plays a role in regulating the root system (Gao *et al.*, 2014). The auxin response factor OsARF25
75 and the type-B response regulators ORR2 and ORR3 regulate the expression of *OsCKX4* downstream of auxin
76 and cytokinin activation, respectively. However, *OsCKX4* overexpression can downregulate the expression of
77 the auxin-biosynthesis gene *OsYUCCA1* and auxin-transport genes *PIN FORMED1b* (*OsPIN1b*) and *OsPIN2* in

the roots, showing a negative feedback between auxin and cytokinin. *OsCKX9* acts downstream of strigolactones (SLs) and plays a key role in the crosstalk between cytokinin and SLs. The SL-induced activation of *OsCKX9* relies on D53, which acts as a repressor of SL signaling and consequently functions in the SL-induced regulation of tiller development (Duan *et al.*, 2019). On the other hand, *OsCKX11* has antagonistic roles between the cytokinin and abscisic acid (ABA) pathways during leaf senescence. In *osckx11* mutants, the downregulation of ABA biosynthesis genes and the upregulation of ABA degradation genes result in the reduction of ABA content in flag leaves and subsequently regulate leaf senescence, which shows the relationship between cytokinin and ABA. Thus, these findings demonstrate that *OsCKX* genes act as bridges between cytokinin and other plant hormones (Zhang *et al.*, 2020).

OsCKX genes also respond to environmental changes and stressors. For example, high-concentration nitrogen treatment significantly increased *OsCKX* expression not only in the nodes and leaves, but also in young panicles (Ding *et al.*, 2014; Xu *et al.*, 2015), which also affected the tiller number and spikelets per panicle. Furthermore, *OsCKX1* and *OsCKX4* were found to be upregulated in nitrogen- and inorganic phosphate-starved roots, potentially affecting root system formation and assisting in the absorption of more nutrients (Shin *et al.*, 2018). *OsCKX4* also plays a role in zinc absorption (Gao *et al.*, 2019), while *OsCKX2* downregulation facilitates the adaptation to saline stress (Joshi *et al.*, 2018). However, it is unknown whether *OsCKX* genes play other roles during rice growth and development. In rice breeding, *OsCKX2* has been artificially selected from wild rice since ancient times and can be divided into alleles from different geographic distributions (Ashikari *et al.*, 2005; Wang *et al.*, 2015). Since most plants with downregulated *CKX* expression exhibit better phenotypes, researchers believed that *CKX* is a potential target to improve yield or to initiate another “Green Revolution” (Ashikari *et al.*, 2005; Chen *et al.*, 2020; Jameson and Song, 2020). Numerous studies have analyzed the structures of and genetic relationships between *OsCKXs*. However, the expression patterns that may determine their functions during plant growth, especially of *OsCKXs* with similar amino acid sequences, are yet to be

elucidated.

In this study, we investigated the genetic relationships and expression patterns of *OsCKX* genes using RNA sequencing (RNA-seq). We evaluated the phenotypes of *osckx* single and double mutants produced using CRISPR/Cas9 technology to determine the function of different *OsCKX*s. We also analyzed the transcriptomes of the roots and shoot bases from Nipponbare (NIP), *osckx4*, *osckx9*, and *osckx4/9* mutants to identify the genes downstream of *OsCKX4* and *OsCKX9*. Our findings may provide new insights regarding the potential application of *OsCKX* genes for improving agricultural traits.

MATERIALS AND METHODS

Plant materials and growth conditions

Oryza sativa L. ssp. *japonica* (cultivars: NIP and ZH11) was the WT rice material chosen for this study. The mutants were comprised of *osckx1/2*, *osckx3*, *osckx4*, *osckx5*, *osckx9*, and *osckx4/9* from NIP and *osckx1*, *osckx2*, *osckx7*, *osckx8*, *osckx9*, and *osckx11* from ZH11. The target for *osckx1/2* matches *OsCKX1* and *OsCKX2* completely, and has a 1-bp mismatch for *OsCKX11*. The sequencing results showed that the *osckx1/2-19* line had a 1-bp insertion in *OsCKX1* and *OsCKX2*, but no change was observed in *OsCKX11* (Supplemental Figure S2).

The rice plants were grown in a field in Danyang, Jiangsu Province, China (31.907° N 119.466° E). We used 300 kg/hm² nitrogen, 150 kg/hm² P₂O₅, and 240 kg/hm² K₂O as base fertilizers for the field experiment. For the pot experiment, 4.28 g of urea was added to each pot, while 1.92 g of KH₂PO₄ and 1.49 g KCl were used during the whole growth period. For the hydroponic experiment, the nutrient solution was composed of 2 mM KNO₃, 2 mM NH₄Cl, 0.32 mM NaH₂PO₄·2H₂O, 0.011 mM MnCl₂·4H₂O, 0.0185 mM H₃BO₃, 0.0006 mM Na₂MoO₄·2H₂O, 0.0014 mM ZnSO₄·7H₂O, 0.0016 mM CuSO₄·5H₂O, 0.3 mM MgCl₂·6H₂O, 0.66 mM CaCl₂, and 0.0448 mM Fe(II)-EDTA (pH = 5.5–6). The plants

were grown in climate chambers under long-day conditions (15 h light at 28 °C and 9 h dark at 24 °C) with a relative humidity of ~70%, and the treatment lasted 20 days.

Plasmid construction for genetic transformation

To construct the mutants for research by using CRISPR/Cas9 technology, as previously described, each *OsCKX* gene was assigned one to two single guide RNA oligo targets by using CRISPR/Cas9 technology as previously described (Mao *et al.*, 2013). The primers used for vector construction and genotyping are shown in Supplemental Table S1.

To construct *pOsCKX::GUS*, an upstream fragment of the *OsCKX*-encoding region (>3 kb) was amplified by PCR, and the resulting amplicon was excised with the corresponding restriction endonuclease and ligated into the *pCAMBIA1300::GUS* vector (Wang *et al.*, 2020).

Analysis of GUS activity

The GUS reporter activity was assayed by histochemical staining using GUS Staining Kit (FCNCS, <https://www.fcncs.com>). Various tissues were collected from *pOsCKX::GUS* transgenic plants at each developmental stage. After dipping in the GUS staining solution, the samples were taken out and then incubated in the staining solution for 6–48 h at 37 °C in the dark. The samples were destained thrice with 70% ethanol in a water bath for 5 min. Images were taken using SZX16 microscope (Olympus, Tokyo, Japan; <https://www.olympus-lifescience.com/en/>).

Sampling, RNA extraction and gene expression analysis

We collected roots from 2 cm under the tip and shoot bases (~5-mm segments from the first nodes) containing the SAMs, axillary buds, young leaves, and tiller nodes from the WT, *osckx4*, *osckx9*, and *osckx4/9* plants. The samples were stored at –80 °C until further use.

Total RNA was extracted using an E.Z.N.A.[®] Plant RNA Kit (Omega Bio-tek Inc., Norcross, GA, USA; <https://www.omegabiotek.com/>) and subjected to reverse-transcription using a PrimeScript[™] RT Reagent Kit

(Takara Biotechnology, Tokyo, Japan). Quantitative reverse-transcription PCR was performed on an ABI PRISM 7300 Real-Time PCR System (Applied Biosystems, Thermo Fisher Scientific, Waltham, MA, USA; <https://www.thermofisher.com/>) with SYBR[®] Premix Ex Taq[™] (Takara), following the manufacturer's instructions. Relative expression analysis was performed using the actin gene as internal control. The primers used are listed in Supplemental Table S1.

Sequence alignment and phylogenetic analysis

The amino acid sequence alignment was performed using MUSCLE (Madeira *et al.*, 2019), and phylogenetic trees were generated using a neighbor-joining method and 1,000 bootstrap iterations in MEGA X (Kumar *et al.*, 2018).

RNA-Seq and data analysis

Before library preparation, oligo (dT)-attached magnetic beads (Invitrogen, Cat. No. 61006) were used to purify the mRNA. An MGIEasy RNA Library preparation kit was used for library construction from purified mRNA according to the manufacturer's instruction (BGI, Cat. No. 1000006383). The quality of the constructed library was checked and sequenced after passing. High-throughput sequencing was paired-end sequenced on the BGISEQ-500 platform (BGI-Shenzhen, China). The obtained reads were processed and analyzed, and genes with Q values of < 0.05 and fold change values of > 1 were considered to be significantly differentially expressed. Based on the list of DEGs, we created and modified Venn diagrams in Microsoft Excel. Furthermore, we performed KEGG enrichment analysis ($Q < 0.05$) (Kanehisa, 2019), and generated a bubble chart using R.

ACCESSION NUMBERS

Sequence data from this article can be found in Rice Annotation Project database under the following accession numbers: *Actin* (Os03g0718100), *OsCKX1* (Os01g0187600), *OsCKX2* (Os01g0197700), *OsCKX3* (Os10g0483500), *OsCKX4* (Os01g0940000), *OsCKX5* (Os01g0775400), *OsCKX7* (Os02g0220100), *OsCKX8*

170 (Os04g0523500), *OsCKX9* (Os05g0374200), *OsCKX11* (Os08g0460600), *OsMADS25* (Os04g0304400),
 171 *OsMADS57* (Os02g0731200), *OsD17* (Os04g0550600), *OsPIN2* (Os06g0660200), *OsWOX11* (Os07g0684900),
 172 *OsRR4* (Os01g0952500), *OsTBI* (Os03g0706500).

RESULTS

OsCKX family genes display different expression patterns

Similar to the findings of previous studies, the *OsCKX* phylogenetic tree consisted of four major clades (Figure 1A). *OsCKX1*, which is commonly expressed in the roots, flowers, and grains, and *OsCKX2* were grouped together in the first clade. The β -glucuronidase (GUS) staining results revealed that *OsCKX1* had high expression in the shoot base and top of axillary buds, but very low expression in the leaf blade (Figure 1B). On the other hand, *OsCKX2*, usually expressed at high levels in the leaf sheath and inflorescence, was found to be highly expressed in the lateral root primordium, leaf blade, shoot base, and inflorescence. *OsCKX6*, *OsCKX7*, and *OsCKX10* were grouped together in the second clade and showed very low expression in all tissues.

The third clade was composed of *OsCKX4*, *OsCKX5*, and *OsCKX9*. *OsCKX4*, which is generally strongly expressed in the vegetative organs, displayed extremely high expression in the roots. *OsCKX9* was generally expressed at low levels in all tissues, showed special expression in the leaf blade, and whole axillary buds. *OsCKX5*, usually highly expressed in all tissues, also exhibited high expression in the roots and leaves.

The fourth clade consisted of *OsCKX3*, *OsCKX8*, and *OsCKX11*. *OsCKX3*, which is expected to have the highest expression in the stem and young panicles, was found to have high expression in the shoot base and young panicle. *OsCKX8*, commonly expressed at lower levels than *OsCKX3* in all vegetative organs, was highly expressed in the shoot base, flag leaf primordia, and inflorescence. Furthermore, *OsCKX11*, generally expressed at a higher level than *OsCKX3* and *OsCKX8* in all tissues, specifically in the reproductive-stage root and inflorescence, displaying high expression in the root, shoot base, and young inflorescence.

Cytokinins have many functions during plant growth and development in different organs, and the biodegradation of cytokinins in different periods or tissues can help plants accurately regulate their growth and development. In rice, we found all of eight of the *OsCKX* genes expressed at the shoot base and inflorescence meristem at different stages and positions. However, each *OsCKX* gene was expressed in a distinct pattern. As

described above, certain *OsCKX* genes were highly expressed in certain organs, and we found some tissues with certain *OsCKX* gene expressed at low levels, such as *OsCKX1* and *OsCKX3*, which were expressed at very low levels in the leaves, and *OsCKX2*, *OsCKX3*, and *OsCKX11*, which were expressed at very low levels in the flag leaf primordia. Additionally, we also determined the expression patterns of *OsCKX* genes in axillary buds: *OsCKX1* was expressed in the top, *OsCKX2*, *OsCKX3*, *OsCKX4*, *OsCKX5* were expressed in the base, and *OsCKX9* was expressed throughout the buds. These distinct expression patterns stimulated our interest in their functions.

***OsCKX* single mutants exhibit different phenotypes**

RNA-Seq analysis and GUS staining assay provided a clear view of the expression levels of *OsCKXs*; however, functional characterization of these genes is still lacking. Previous studies suggested that the knockdown of specific *OsCKXs* might help improve rice production by influencing important agronomic traits, such as tillering, development of organs facilitating nutrition, and panicle phenotype. To confirm this hypothesis, we analyzed the phenotypes of nine *OsCKX* single mutants (except for *OsCKX6* and *OsCKX10*) created using the CRISPR/Cas9 system (Supplemental Figures S1 and S2; Supplemental Table S1).

Since leaf size affects the photosynthetic capacity of rice, we investigated this and discovered that the top three leaves in the *osckx1*, *osckx2*, *osckx8*, and *osckx11* mutants were distinctly longer and wider compared to those of Zhonghua 11 (Supplemental Figures S3–S5; Supplemental Table S2). The last three leaves of the *osckx9* mutant were also distinctly longer, yet no change in their width was observed compared to Zhonghua 11. The leaf length and width of the other mutants did not significantly change (Supplemental Figures S4, A–D, and S5). On the other hand, culm diameter and plant height are closely related to the lodging resistance of rice. We found that the diameters of the basal internodes in *osckx1*, *osckx2*, *osckx8*, and *osckx11* mutants were significantly wider than those in the wild-type (WT), while the *osckx3*, *osckx4*, and *osckx5* mutants had thinner basal internodes (Supplemental Figure S4, E and F). Furthermore, the *osckx3*, *osckx4*, and *osckx5* mutants

showed significantly decreased plant height compared to others (Supplemental Figure S4, G and H).

The panicle number is determined by the tiller number per plant and has a significant effect on rice yield.

In the field and pot experiments, the *osckx4* and *osckx9* mutants developed more tillers than the WT plants, which is consistent with the results of previous studies (Supplemental Figures S3B and S4, I and J; Supplemental Tables S2 and S3). Although the mutants produced more panicles per plant, the number of grains per panicle decreased, resulting in no significant change in yield (Supplemental Figure S6, A and B; Supplemental Table S2). In contrast, the *osckx2* mutant showed significantly reduced tiller number in both the 2019 and 2020 field and pot experiments (Supplemental Tables S2 and S3). On the other hand, no significant differences were observed in the tiller numbers of the other *osckx* mutants. Additionally, there was no significant change in the yield per plant of the *osckx* mutants, except for *osckx1* and *osckx11-1*, which had significantly improved production. In contrast, *osckx3-1*, *osckx3-26*, and *osckx4-6* had significantly low yields (Supplemental Figures S6, A and B). Despite this, we observed several agronomic traits that were improved in the mutants. For example, the *osckx1*, *osckx2*, *osckx7*, and *osckx8* mutants presented heavier grain weight, while the *osckx11* mutant developed more spikelets per panicle (Supplemental Figure S6, E–H).

The *osckx1 osckx2* double mutants have significantly reduced tiller numbers

Since the phenotypes of *osckx1* and *osckx2* plants were similar, we created the new double mutant *osckx1/2*. In the T1 generation, we obtained enough plants of the *osckx1/2-19* line, which has 1-bp insertions in *OsCKX1* and *OsCKX2* (Figure 2A; Supplemental Figure S2). The *osckx1/2-19* mutant showed significantly reduced tiller number per plant (Figure 2B), consequently leading to lower panicle number per plant (Figure 2C). Notably, the *osckx1/2-19* mutants have large panicle sizes and more spikelets per panicle because the number of primary branches increased while the number of secondary branches remained the same (Figure 2, D–G). Furthermore, the 1,000-grain weight and grain length were significantly increased in *osckx1/2-19* compared with those in NIP (Figure 2H–J). In contrast, the grain width and thickness did not show significant changes between *osckx1/2-19*

and NIP (Figure 2, I, K, and L). However, the total number of panicles and setting rate were lower in *osckx1/2-19* compared to those in NIP (Figure 2M), which resulted in a lower yield with no significant differences (Figure 2N). Overall, the *osckx1/2-19* plants showed better plant architecture than the WT. In *osckx1/2-19* plants, the flag leaf became wider (Figure 2, O and P) and the plants had thick basal internodes, while the plant height did not noticeably change (Figure 2, Q and R). Considered together, these observations suggest that the double knockout of *OsCKX1* and *OsCKX2* may have improved the lodging resistance and optimized plant architecture in rice.

Double knockout of *OsCKX4* and *OsCKX9* significantly promotes tillering

Since the *osckx4* and *osckx9* single mutants developed more tillers, we created two new single mutants, namely *osckx4-6* and *osckx9-1*, and an *osckx4/9* double mutant in NIP to verify if these will also produce more tillers (Supplemental Figure S7, A and B). As expected, we discovered that the *osckx4/9* double mutant had an extremely higher tiller number compared to NIP and the two single mutants (Figure 3, A–D). In contrast, panicle size was significantly reduced in the *osckx4/9* mutant (Figure 3E). The *osckx9-1* mutant had a high panicle number per plant, but the number was markedly higher in the *osckx4/9* mutant (Figure 3F). However, the number of spikelets per panicle of *osckx4* was significantly decreased, while that of *osckx4/9* was even lower (Figure 3G). The decreased number of spikelets per panicle of *osckx4-6* was due to the low number of secondary branches per panicle, while the low number of spikelets per panicle of *osckx4/9* was caused by the reduced number of both the primary and secondary branches (Figure 3, E, H, and I). On the other hand, seed setting rates were similar in the single and double mutants, and were significantly lower than those in WT (Figure 3J). Despite the increase in grain length, the *osckx4-6* mutant presented decreased 1,000-grain weight due to reduced grain thickness (Figure 3K; Supplemental Figure S7, C–G). The 1,000-grain weight of *osckx4/9* was lower than that of the *osckx4-6* mutant due to reduced grain width and thickness (Figure 3I; Supplemental Figure S7). As a result of these changes, the yield of *osckx4/9* was significantly decreased (Figure 3L). Moreover, the root length,

diameter of basal internode, plant height, and flag leaf length and width were further decreased in the double mutant compared to those in the two single mutants (Figure 3, A, M and N; Supplemental Figure S7, H–J).

We also conducted a hydroponic experiment to investigate the root phenotypes of the mutants (Figure 4A). We verified that *OsCKX4* and *OsCKX9* have redundant roles in regulating tillering, plant height, and root length (Figure 4 B and C). The *osckx4/9* mutant not only had shorter roots, but also fewer crown roots and smaller root diameter (Supplemental Figure S8, A and B). The length and width of the 8th leaf were also decreased in *osckx4-6* and *osckx4/9* compared to those in NIP (Figure 4, D and E). These results indicate that the reduction in the leaf size of *osckx4/9* started at the vegetative stage. Overall, these results suggest that *OsCKX4* and *OsCKX9* have a functional overlap during rice growth and development.

RNA-seq analysis reveals the molecular functions of *OsCKX4* and *OsCKX9*

To elucidate the functions of *OsCKX4* and *OsCKX9*, we sampled the roots and shoot bases from NIP, *osckx4*, *osckx9*, and *osckx4/9* and performed RNA-seq analysis. Results revealed that *osckx4* had 297 differentially expressed genes (DEGs) in the shoot bases and 415 DEGs in the roots. There were 500 DEGs in the shoot bases and 342 DEGs in the roots of the *osckx9* mutant. The *osckx4/9* double mutant had the highest number of DEGs, with 806 DEGs in the shoot base and 4,556 DEGs in the roots (Figure 5A). To verify the accuracy and reproducibility of the RNA-seq results, we randomly selected seven previously studied genes associated with the phenotypes observed in this study for quantitative reverse-transcription PCR analysis. The expression profiles of these genes were found to corroborate the RNA-seq results (Supplemental Figure S9).

However, although there were numerous DEGs between the WT and mutants, there were 71 DEGs in the roots and 90 DEGs in the shoot bases that overlapped between the *osckx4* vs. WT and *osckx9* vs. WT comparisons and occupied only a small fraction of the total DEGs (Figure 5B). These included some genes that regulate the absorption of silicon (*Lsi1*, *Lsi2*), zinc (*OsZIP16*), and iron (*OsIRO2*) (Fig 5C). *OsZIP9*, *OsTCP19*, *OsHKT1.1*, and *OsNPF5.5* were upregulated in the shoots of *osckx4*, while *OsLBD37*, *OsARF19*, *OsMYB61*,

and *MT1a* were upregulated in the shoots of *osckx9*. In the roots, the downregulation of *OsAAP5*, *OsAAP8*, and *OsNLP6* and the upregulation of certain genes influenced the iron and phosphorus absorption in *osckx4*. We also observed the downregulation of *OsPUP1*, *OsPUP5*, *OsRR10*, and *OsNAAT4* in *osckx9*. These DEGs were also revealed to be associated with the different pathways controlled by *OsCKX4* and *OsCKX9* (Figure 5C). However, the functions of most DEGs remain unknown (Supplemental Dataset 1). The difference in the number of DEGs identified may imply that *OsCKX4* and *OsCKX9* have different functions. Moreover, 87.1% and 78.2% of DEGs in the roots and shoot bases, respectively, between the *osckx4/9* and WT were identified only in the double mutants, which may have been caused by the duplicate effect of the functional loss of *OsCKX4* and *OsCKX9* (Figure 5B).

Since we observed that *osckx4/9* exhibited more severe phenotypes and possessed more DEGs, we analyzed the DEGs identified in the roots and shoot bases between the WT and *osckx4/9* plants by using Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis (Figure 5D). In both tissues, the top enriched KEGG pathways included ‘glycolysis/gluconeogenesis,’ ‘biosynthesis of amino acids,’ and those related to carbon and nitrogen metabolism. Since most of the DEGs were not functionally annotated, the relationship between the DEGs and the enriched pathways and their influence on the growth and development of the mutant plants were not fully understood. However, we identified several genes that may be associated with the phenotypes observed in this study.

***OsCKX4* and *OsCKX9* regulate the metabolism and transport of hormones**

First, we identified genes related to gibberellin metabolism. We found that the expression of *OsKSI* and *OsKS2* was significantly downregulated in the roots of *osckx4/9* (Figure 6A; Supplemental Dataset 1). *OsKSI* and *OsKS2* are enzymes involved in gibberellin biosynthesis. The expression of *OsKS2* was downregulated in the roots of the *osckx9* mutant, while no changes were observed in the *osckx4* mutant. Other genes for GA biosynthesis and degradation, such as *GA20ox* and *GA2ox*, were markedly downregulated in the roots of the

osckx4/9 mutant. Some GA receptors were also downregulated in the roots of the *osckx4/9* mutant. The changes in the expression of gibberellin-related genes may explain the significantly reduced plant height of *osckx4/9* double mutants.

Further, we examined the genes associated with SLs to investigate the cause of the increased tiller numbers in the single and double mutants. We discovered that *OsDI7* was downregulated 1.20-fold, which was not significant in the roots of *osckx4/9* (Figure 6B; Supplemental Dataset 1), this may have been the cause of the increased tiller number and another reason for reduced plant height of *osckx4/9*. However, *OsDI10* was upregulated 1.59-fold in the shoot base of *osckx4/9*, and *OsDI7* was upregulated 1.38-fold in the shoot base of *osckx9*, neither of which reflect significant differences. This may have been caused by feedback regulation (Arite *et al.*, 2007). Furthermore, the SL-related transcription factors *OsTBI* and *OsMADS57* were upregulated 1.28-fold and 1.03-fold, respectively, in the shoot bases of *osckx4*, neither of which were significant changes. Finally, *OsTBI* and *OsMADS57* were downregulated 1.71-fold and 2.29-fold in the roots of *osckx4/9*, and *OsMADS57* was downregulated 1.24-fold in the roots of *osckx9*, but their functions in roots remain unknown.

Additionally, investigation of several differentially expressed auxin-related genes revealed that auxin efflux transporters *OsPIN2* and *OsPIN8* were upregulated in the shoot bases and the roots, respectively, of the mutants. We found that the auxin primary response genes *OsGH3-1*, *OsGH3-2*, *OsIAA9*, and *OsSAUR12* were upregulated in the shoot bases of the *osckx4/9* and single mutants. Additionally, *OsGH3-2*, nine *OsIAA* genes (*OsIAA1*, 4, 9, 12, 14, 15, 19, 23, 24), and five *OsSAUR* genes (*OsSAUR6*, 20, 27, 28, 33) were downregulated in the roots of *osckx4/9*; however, *OsIAA16* was upregulated in the roots of *osckx4/9*.

Furthermore, the functional loss of *OsCKX4* and *OsCKX9* also affected the expression of genes associated with cytokinin metabolism and signal transduction. The cytokinin-related genes *OsCKX2*, *OsCKX5*, *OsRR4*, *OsRR6*, *OsRR7*, *OsRR9*, and *OsRR10* were downregulated in the roots of *osckx4/9*, while *OsCRL4* was upregulated in the shoot bases of *osckx4/9* (Supplemental Dataset 1). These results may have been caused by the

high cytokinin contents in the double mutants.

***OsCKX4* and *OsCKX9* are associated with nitrogen absorption**

Many genes related to nitrogen absorption and assimilation, such as *OsNRT1*, *OsNRT2.2*, *OsNRT2.3*, *OsAMT1.3*, *OsNAR2.1*, and *OsNIR2*, were also discovered to be significantly upregulated in the roots of the *osckx4/9* mutant (Figure 7; Supplemental Dataset 1). In contrast, we observed the downregulation of *OsGS1;3* in the roots and that of *OsAS1* in the shoot bases of *osckx4/9*. Most of the genes related to nitrogen absorption and assimilation also showed similar trends in *osckx4*, while some did not show significant differences compared to NIP. However, *OsAMT3.2* and *OsGS1;3* were upregulated in the shoot bases of *osckx9*. The changes in the expression of these genes were significantly different from those of the *osckx4/9* double mutants. Several nitrate and peptide transporter (*OsNPF*) family genes were also found to be differentially expressed.

DISCUSSION

Functional differences and redundancy between *OsCKX* genes

To systematically determine the functions of *OsCKX* genes, we analyzed their expression patterns and investigated the phenotypes of *osckx* single and double mutants. Our study revealed the roles of most *OsCKX* genes in the growth and development of rice. After two years of field experiments, we discovered that *osckx4* and *osckx9* developed more tillers at the vegetative stage, while *osckx2* had fewer tillers. At the reproductive stage, the top three leaves were longer and wider and the basal internodes were thicker in *osckx1*, *osckx2*, *osckx8*, and *osckx11* (Zhonghua11 [ZH11] background), while wider flag leaves and thicker basal internodes were observed in *osckx1/2* (NIP background) (Figure 8). The *osckx9*, *osckx3*, *osckx4*, and *osckx5* mutants showed little changes in flag leaf length and width compared to their background, while *osckx3*, *osckx4*, and *osckx5* developed thinner basal internodes. For yield-related phenotypes, *osckx9* developed more panicles while *osckx9* and *osckx4* had fewer spikelets per panicle. Moreover, *osckx1*, *osckx2*, and *osckx8* displayed increased

1,000-grain weight, while *osckx11* showed reduced 1,000-grain weight.

Previous studies have described the functions of *OsCKX2*, *OsCKX4*, *OsCKX9*, and *OsCKX11*. First, researchers elucidated the function of *OsCKX2* after observing that null mutants or plants with downregulation of *OsCKX2* developed more spikelets per panicle (Ashikari *et al.*, 2005), more tillers, and larger grains (Yeh *et al.*, 2015). In the present study, we observed very little increase in the number of spikelets per panicle during the two-year field experiment, with the observed increase being lower than that observed by the other researchers. However, we found that the *osckx2* mutant of ZH11 showed significantly reduced tiller number within the first year of the field experiment. Another mutant, *osckx1/2* of NIP, also showed a reduced tiller number, which supports the hypothesis that fewer tillers may be related to thicker culms. Furthermore, both the *osckx2* mutant of ZH and *osckx1/2* mutant of NIP presented increased 1,000-grain weight. The discrepancies between our results and previous findings regarding panicle size may be attributed to the different cultivated varieties or environmental impact.

Furthermore, previous reports stated that *OsCKX4* overexpression can cause the development of a larger root system, but result in shorter plant height, fewer tillers, and smaller panicles (Gao *et al.*, 2014). For example, *OsCKX4-RNAi* or *osckx4* mutants showed fewer crown roots and shorter root lengths (Gao *et al.*, 2014; Mao *et al.*, 2020). Since *OsCKX4* is commonly expressed in the root, previous studies were more focused on investigating the relationships between *OsCKX4* and the root system instead of the shoot system. Here, we discovered that *osckx4* mutants produced more tillers, especially at the early stages, but has fewer spikelets per panicle. Additionally, *osckx9* mutants have been reported to develop more tillers, fewer spikelets per panicle, and shorter plant height (Duan *et al.*, 2019). Since the null mutations of *OsCKX4* and *OsCKX9* can affect the tiller number, we constructed *osckx4/9* mutants, which distinctly showed more tillers, shorter plant height, and smaller panicles. The increase in the tiller number of *osckx4/9* double mutants was greater than observed in *osckx4* and *osckx9* single mutants. Furthermore, the decrease in plant height and number of spikelets per panicle

of *osckx4/9* double mutants was also higher than observed in *osckx4* and *osckx9* single mutants. These results suggest the partial functional overlap of *OsCKX4* and *OsCKX9* during rice growth and development. Moreover, *osckx11* mutants previously displayed more tillers, larger panicles, and lighter 1,000-grain weight (Zhang *et al.*, 2020); these were also observed in our study. In general, we found that each *OsCKX* gene had its own function during rice growth and development as well as their own distinct characteristics, and each *osckx* single or double mutant had a distinct phenotype, but why these different phenotypes happen need to be explored in depth.

Potential causes of the functional differences and redundancy in *OsCKX* genes

The main function of CKX enzymes is the biodegradation of cytokinins. Previous studies have reported that CKXs catalyze different kinds of cytokinins. In maize, the ZmCKX enzyme activity has been systematically analyzed, revealing that ZmCKX1 and ZmCKX12 specially targeted iP, tZ, and *cis*-zeatin (cZ). Furthermore, ZmCKX5 was partial to iP and isopentenyladenosine 5'-monophosphate, while ZmCKX2, ZmCKX3, ZmCKX4a, and ZmCKX4b preferred the biodegradation of iP-9-glucoside (iP9G) and cZ. Finally, ZmCKX8, ZmCKX9, and ZmCKX10 showed preference for tZ and cZ (Zalabak *et al.*, 2014). In *Arabidopsis*, AtCKX1 showed the highest preference for tZ phosphates, iP phosphates, and iP9G, while AtCKX3 preferred iP phosphates and iPR, and AtCKX7 preferred iP9G and tZ (Kowalska *et al.*, 2010). However, little is known about the preference of OsCKX enzymes, in vitro experiment shows OsCKX9 that is partial to tZ and OsCKX11 that specially targets tZ and cZ (Duan *et al.*, 2019; Zhang *et al.*, 2020). OsCKX11 has the same substrate as its homolog protein, ZmCKX10, while other OsCKX enzyme substrates may need to be inferred from endogenous hormone content data. From the hormone content detected in the *OsCKX4-overexpression* line, it was found that the content of iP and tZ cytokinin showed no obvious change compared to WT, while DHZ and the combined type of iP and tZ showed a significant reduction (Gao *et al.*, 2014). Therefore, the distinctive functions of OsCKXs may be caused by their different target substrates.

CKX proteins are mainly localized in the ER because most contain an N-terminal signal peptide sequence

(Niemann *et al.*, 2018; Zalabak *et al.*, 2016). In the ER, cytokinins combine with their receptors and initiate signaling transduction, while most ER-localized CKXs regulate cytokinin concentrations to precisely adjust cytokinin signals (Niemann *et al.*, 2018; Romanov and Schmulling, 2021). Previous studies revealed that OsCKX4 and OsCKX11 are localized in the cytosol, while OsCKX9 is localized in the cytosol and nuclei (Duan *et al.*, 2019; Gao *et al.*, 2014; Zhang *et al.*, 2020). These three CKXs potentially regulate the cytokinin balance in other cellular organelles. However, the locations of the remaining OsCKXs remain unknown. Hence, we hypothesize that the subcellular localization of OsCKXs may also influence their functional differentiation.

In this study, the *OsCKX* genes exhibited different expression patterns. For example, *OsCKX1* and *OsCKX2* were expressed in the flower and grains, but *OsCKX1* showed lower expressed in the inflorescence meristem, which is the same as the previous study (Ashikari *et al.*, 2005). As *OsCKX4* and *OsCKX9* showed similar phenotypes in tiller number, the expression pattern showed a considerable difference. *OsCKX4* has a high expression highly in nutrition organs, yet the expression of *OsCKX9* in these organs is very low, and the different expression can get illation from previous studies (Duan *et al.*, 2019; Gao *et al.*, 2014). In inflorescence meristem, the two genes showed low expression level, which is also the same as previous studies by RT-southern blot (Ashikari *et al.*, 2005). In summary, the differences in the nature of catalytic substrates, subcellular localization, and expression in tissues may contribute to the different functions observed in *OsCKX* family genes.

Significantly altered transcriptome of the *osckx4 osckx9* mutant

Although *OsCKX4* and *OsCKX9* have been extensively studied, we observed new phenotypes in the single mutants, which have not been reported previously. Interestingly, these two genes were found to exhibit redundant functions during rice growth and development. Hence, we first investigated the genes associated with plant height and then identified the DEGs related to gibberellin metabolism between the NIP and mutants. *OsKSI* and *OsKS2*, which encode enzymes that catalyze the second step of the GA biosynthesis pathway (Ji *et*

al., 2014; Tezuka *et al.*, 2015), were significantly downregulated in the roots of *osckx4/9*. On the other hand, possibly due to the limited substrate, the expression of several *GA20ox* and *GA2ox* genes for gibberellin biosynthesis and biodegradation, respectively, were downregulated in *osckx4/9*. We hypothesize that the disrupted biosynthesis of gibberellin may have caused the reduction in *osckx4/9* plant height. However, the mechanism by which *OsCKX4* and *OsCKX9* regulate the expression of *OsKSI* and *OsKS2* is unknown.

Since the double mutants exhibited semi-dwarf phenotypes with more tillers, we believed that the expression of SL-related genes may have changed in the double mutants. However, we did not observe any changes in the expression of SL-signaling genes but found that *OsD17* was downregulated in the roots of *osckx4/9* mutants. This contradicts a previous report that stated *d17* mutants developed dwarf plants and more tillers (Zou *et al.*, 2005). On the other hand, the SL-related transcript factors *OsTBI* and *OsMADS57* were only downregulated in the roots. Furthermore, *OsTBI* was upregulated in the shoot bases of *osckx4*. Previous studies on *tb1* mutants and *OsMADS57* overexpression showed that these plants developed more tillers than the WT (Guo *et al.*, 2013; Takeda *et al.*, 2003). Since SL directly activates *OsCKX9* to regulate shoot architecture in rice, we also determined whether *OsCKX4* and *OsCKX9* can regulate the tiller number phenotype through the SL pathway by analyzing the expression of SL-related genes between WT and mutants. Based on these findings, cytokinin may be located downstream of the SL pathway during rice tillering.

During the outgrowth of tiller buds, the auxin at the tip of the axillary bud is transported by auxin efflux carriers. In *Arabidopsis*, cytokinin response factors regulate *PIN* expression (Simaskova *et al.*, 2015; Waldie and Leyser, 2018). Here, we found that *OsPIN2* was upregulated in the shoot bases of *osckx4* and *osckx4/9*. In rice, *OsPIN2* transports auxins from the shoot to the root–shoot junction, and overexpression of *OsPIN2* results in the increased number of tillers (Chen *et al.*, 2012). Moreover, in *OsCKX4-overexpression* line, the content of IAA is significantly reduced in root and the expression of *OsPIN1b* and *OsPIN2* showed significantly lower (Gao *et al.*, 2014), the connection between cytokinin and auxin transporters is needed for further research.

We also found that several type-A transcription factors, including *OsRR4*, *OsRR6*, *OsRR7*, *OsRR9*, and *OsRR10*, showed lower expression in the roots of *osckx4/9*. Some were downregulated in single mutants. However, these results were contradictory, since type-A RR genes are primary response genes for cytokinin signaling, while *OsRR4*, *OsRR9*, and *OsRR10* were previously found to be upregulated in *OsCKX4*-overexpression lines (Gao *et al.*, 2014). Since *OsRR4*, *OsRR9*, and *OsRR10* share similar amino acid sequences (Tsai *et al.*, 2012; Wang *et al.*, 2019) and *osrr9 osrr10* mutant lines developed fewer spikelets per panicle and higher tolerance to saline stress, we hypothesized that *OsRR4*, *OsRR9*, and *OsRR10* may have important functions during the cytokinin-induced inhibition of root growth. In the present study, we observed the downregulation of *OsWOX11* (Zhao *et al.*, 2009) and *OsMADS25* (Yu *et al.*, 2015) in the roots of *osckx4* and *osckx4/9*, which may explain why *osckx4* and *osckx4/9* plants have smaller root systems. Furthermore, we discovered that *OsCKX4* and *OsCKX9* influence several genes related to nitrogen absorption and utilization, providing new potential targets for improving nitrogen usage efficiency in rice. However, the mechanisms underlying this process are unknown; hence, further research is still required.

Interestingly, in addition to the genes regulated by both *OsCKX4* and *OsCKX9*, we also identified several genes that are regulated by *OsCKX4* and *OsCKX9* individually. In summary, the similar amino acid sequences but different expression patterns and different subcellular localization of *OsCKX4* and *OsCKX9* enable them to functionally overlap in the regulation of some pathways, while also individually regulating specific pathways.

In conclusion, Our systematic analysis of 11 genes in the *OsCKX* family reveals their complex expression patterns in rice. A total of nine *OsCKX* mutants were produced using CRISPR/Cas9 technology. By examining the phenotypes of the mutants throughout the rice growth period, we determined the functions of specific *OsCKX* genes in plant development. In specific, we discovered that *OsCKX4* and *OsCKX9* inhibited tillering, while *OsCKX1* and *OsCKX2* promoted tillering. Considered together, our findings establish a community resource for fully elucidating the function of *OsCKXs*, providing new insights that may be used for future

472 studies to improve rice yield and initiate green production.

473

474 **Supplemental data**

475 The following supplemental materials are available.

476 **Supplemental Figure S1.** Gene structures and mutation details of *OsCKXs*.

477 **Supplemental Figure S2.** Chromatograms of the *osckx* mutant lines.

478 **Supplemental Figure S3.** The *osckx* mutants of Zhonghua11 from the 2020 field experiment.

479 **Supplemental Figure S4.** Phenotypic characterization of the vegetative organs in *osckx* mutants from the 2020
480 field experiment.

481 **Supplemental Figure S5.** Phenotypic characterization of the second and third top leaves in *osckx* mutants from
482 the 2020 field experiment.

483 **Supplemental Figure S6.** Phenotypic characterization of yield-related phenotypes in *osckx* mutants from the
484 2020 field experiment.

485 **Supplemental Figure S7.** Mutation details of *OsCKX9*, chromatograms of the *osckx9* and *osckx4/9* mutant lines,
486 and phenotypic characterization of the flag leaves and seeds in wild-type, *osckx4*, *osckx9*, and *osckx4/9* plants.

487 **Supplemental Figure S8.** Phenotypic characterization of the leaf and root systems in wild-type and *osckx4/9*
488 plants.

489 **Supplemental Figure S9.** Relative expression levels and FPKM values of *D17*, *RR4*, *WOX11*, *PIN2*, *TB1*,
490 *OsMADS25*, and *OsMADS57* in the roots and shoot bases (BP) of Nipponbare (NIP), *osckx4*, *osckx9*, and
491 *osckx4/9*.

492 **Supplemental Table S1.** Information of primers used in the study.

493 **Supplemental Table S2.** Characterization of the vegetative organs and yield-related phenotypes in *osckx*
494 mutants from the 2019 field experiment.

495 **Supplemental Table S3.** Tiller numbers in *osckx* mutants from the 2019 field experiment.

496 **Supplemental Dataset 1.** RNA sequencing data.

Funding:

This work was supported by the National Natural Science Foundation of China (Grant number 31872855 and 31971842).

Data availability

All data supporting the findings of this study are available within the paper and within its supplementary data published online.

Acknowledgments

We thank Jiankang Zhu and Caixia Gao for providing the vectors of the CRISPR-Cas9 system. We also thank Biogle and Biorun genome editing center for producing transgenic rice.

Author contributions

C.D. and Y.D. conceived the original screening and research plans; C.R., Y.L., Z.C., Z.L., and C.D. performed the experiments and analyzed the data; C.R. and C.D. wrote the article with contributions of all the authors; C.D. agrees to serve as the author responsible for contact and ensures communication.

Conflict of interest

The authors declare no competing interests in relation to this work.

Reference

- Ashikari M, Sakakibara H, Lin S, Yamamoto T, Takashi T, Nishimura A, Angeles ER, Qian Q, Kitano H, Matsuoka M.** 2005. Cytokinin oxidase regulates rice grain production. *Science* **309**, 741-745.
- Bartrina I, Otto E, Strnad M, Werner T, Schmulling T.** 2011. Cytokinin regulates the activity of reproductive meristems, flower organ size, ovule formation, and thus seed yield in *Arabidopsis thaliana*. *Plant Cell* **23**, 69-80.
- Chen L, Zhao J, Song J, Jameson PE.** 2020. Cytokinin dehydrogenase: a genetic target for yield improvement in wheat. *Plant Biotechnol J* **18**, 614-630.
- Chen Y, Fan X, Song W, Zhang Y, Xu G.** 2012. Over-expression of OsPIN2 leads to increased tiller numbers, angle and shorter plant height through suppression of OsLAZY1. *Plant Biotechnol J* **10**, 139-149.
- Ding C, You J, Chen L, Wang S, Ding Y.** 2014. Nitrogen fertilizer increases spikelet number per panicle by enhancing cytokinin synthesis in rice. *Plant Cell Rep* **33**, 363-371.
- Duan J, Yu H, Yuan K, Liao Z, Meng X, Jing Y, Liu G, Chu J, Li J.** 2019. Strigolactone promotes cytokinin degradation through transcriptional activation of CYTOKININ OXIDASE/DEHYDROGENASE 9 in rice. *Proc Natl Acad Sci U S A* **116**, 14319-14324.
- Gao S, Fang J, Xu F, Wang W, Sun X, Chu J, Cai B, Feng Y, Chu C.** 2014. CYTOKININ OXIDASE/DEHYDROGENASE4 Integrates Cytokinin and Auxin Signaling to Control Rice Crown Root Formation. *Plant Physiol* **165**, 1035-1046.
- Gao S, Xiao Y, Xu F, Gao X, Cao S, Zhang F, Wang G, Sanders D, Chu C.** 2019. Cytokinin-dependent regulatory module underlies the maintenance of zinc nutrition in rice. *New Phytol* **224**, 202-215.
- Guo S, Xu Y, Liu H, Mao Z, Zhang C, Ma Y, Zhang Q, Meng Z, Chong K.** 2013. The interaction between OsMADS57 and OsTB1 modulates rice tillering via DWARF14. *Nat Commun* **4**, 1566.
- Hosek P, Hoyerova K, Kiran NS, Dobrev PI, Zahajska L, Filepova R, Motyka V, Muller K, Kaminek M.** 2020. Distinct metabolism of N-glucosides of isopentenyladenine and trans-zeatin determines cytokinin

metabolic spectrum in Arabidopsis. *New Phytol* **225**, 2423-2438.

Jameson PE, Song J. 2020. Will cytokinins underpin the second 'Green Revolution'? *J Exp Bot* **71**, 6872-6875.

Ji SH, Gururani MA, Lee JW, Ahn BO, Chun SC. 2014. Isolation and characterisation of a dwarf rice mutant exhibiting defective gibberellins biosynthesis. *Plant Biol (Stuttg)* **16**, 428-439.

Joshi R, Sahoo KK, Tripathi AK, Kumar R, Gupta BK, Pareek A, Singla-Pareek SL. 2018. Knockdown of an inflorescence meristem-specific cytokinin oxidase - OsCKX2 in rice reduces yield penalty under salinity stress condition. *Plant Cell Environ* **41**, 936-946.

Kanehisa M. 2019. Toward understanding the origin and evolution of cellular organisms. *Protein Science* **28**, 1947-1951.

Kollmer I, Novak O, Strnad M, Schmulling T, Werner T. 2014. Overexpression of the cytosolic cytokinin oxidase/dehydrogenase (CKX7) from Arabidopsis causes specific changes in root growth and xylem differentiation. *Plant J* **78**, 359-371.

Kowalska M, Galuszka P, Frebortova J, Sebela M, Beres T, Hluska T, Smehilova M, Bilyeu KD, Frebort I. 2010. Vacuolar and cytosolic cytokinin dehydrogenases of Arabidopsis thaliana: heterologous expression, purification and properties. *Phytochemistry* **71**, 1970-1978.

Kudo T, Makita N, Kojima M, Tokunaga H, Sakakibara H. 2012. Cytokinin activity of cis-zeatin and phenotypic alterations induced by overexpression of putative cis-Zeatin-O-glucosyltransferase in rice. *Plant Physiol* **160**, 319-331.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549.

Kurakawa T, Ueda N, Maekawa M, Kobayashi K, Kojima M, Nagato Y, Sakakibara H, Kyoizuka J. 2007. Direct control of shoot meristem activity by a cytokinin-activating enzyme. *Nature* **445**, 652-655.

Li S, Zhao B, Yuan D, Duan M, Qian Q, Tang L, Wang B, Liu X, Zhang J, Wang J, Sun J, Liu Z, Feng YQ,

Yuan L, Li C. 2013. Rice zinc finger protein DST enhances grain production through controlling Gn1a/OsCKX2 expression. *Proc Natl Acad Sci U S A* **110**, 3167-3172.

Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN, Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res* **47**, W636-W641.

Mao C, He J, Liu L, Deng Q, Yao X, Liu C, Qiao Y, Li P, Ming F. 2020. OsNAC2 integrates auxin and cytokinin pathways to modulate rice root development. *Plant Biotechnol J* **18**, 429-442.

Mao Y, Zhang H, Xu N, Zhang B, Gou F, Zhu JK. 2013. Application of the CRISPR-Cas system for efficient genome engineering in plants. *Mol Plant* **6**, 2008-2011.

Niemann MCE, Weber H, Hluska T, Leonte G, Anderson SM, Novak O, Senes A, Werner T. 2018. The Cytokinin Oxidase/Dehydrogenase CKX1 Is a Membrane-Bound Protein Requiring Homooligomerization in the Endoplasmic Reticulum for Its Cellular Activity. *Plant Physiol* **176**, 2024-2039.

Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H. 2017. Systemic transport of trans-zeatin and its precursor have differing roles in Arabidopsis shoots. *Nat Plants* **3**, 17112.

Romanov GA, Schmulling T. 2021. Opening Doors for Cytokinin Trafficking at the ER Membrane. *Trends Plant Sci* **26**, 305-308.

Sakamoto T, Sakakibara H, Kojima M, Yamamoto Y, Nagasaki H, Inukai Y, Sato Y, Matsuoka M. 2006. Ectopic expression of KNOTTED1-like homeobox protein induces expression of cytokinin biosynthesis genes in rice. *Plant Physiol* **142**, 54-62.

Schwarz I, Scheirlinck MT, Otto E, Bartrina I, Schmidt RC, Schmulling T. 2020. Cytokinin regulates the activity of the inflorescence meristem and components of seed yield in oilseed rape. *J Exp Bot* **71**, 7146-7159.

Shin SY, Jeong JS, Lim JY, Kim T, Park JH, Kim JK, Shin C. 2018. Transcriptomic analyses of rice (*Oryza sativa*) genes and non-coding RNAs under nitrogen starvation using multiple omics technologies. *BMC*

Genomics **19**, 532.

Simaskova M, O'Brien JA, Khan M, Van Noorden G, Otvos K, Vieten A, De Clercq I, Van Haperen JMA,

Cuesta C, Hoyerova K, Vanneste S, Marhavy P, Wabnik K, Van Breusegem F, Nowack M, Murphy A,

Friml J, Weijers D, Beeckman T, Benkova E. 2015. Cytokinin response factors regulate PIN-FORMED auxin transporters. *Nat Commun* **6**, 8717.

Takeda T, Suwa Y, Suzuki M, Kitano H, Ueguchi-Tanaka M, Ashikari M, Matsuoka M, Ueguchi C. 2003.

The OsTB1 gene negatively regulates lateral branching in rice. *Plant J* **33**, 513-520.

Tezuka D, Ito A, Mitsuhashi W, Toyomasu T, Imai R. 2015. The rice ent-KAURENE SYNTHASE LIKE 2 encodes a functional ent-beyerene synthase. *Biochem Biophys Res Commun* **460**, 766-771.

Tsai YC, Weir NR, Hill K, Zhang W, Kim HJ, Shiu SH, Schaller GE, Kieber JJ. 2012. Characterization of genes involved in cytokinin signaling and metabolism from rice. *Plant Physiol* **158**, 1666-1684.

Waldie T, Leyser O. 2018. Cytokinin Targets Auxin Transport to Promote Shoot Branching. *Plant Physiol* **177**, 803-818.

Wang J, Xu H, Li N, Fan F, Wang L, Zhu Y, Li S. 2015. Artificial Selection of Gn1a Plays an Important role in Improving Rice Yields Across Different Ecological Regions. *Rice (N Y)* **8**, 37.

Wang WC, Lin TC, Kieber J, Tsai YC. 2019. Response Regulators 9 and 10 Negatively Regulate Salinity Tolerance in Rice. *Plant Cell Physiol* **60**, 2549-2563.

Wang Y, Lu Y, Guo Z, Ding Y, Ding C. 2020. RICE CENTRORADIALIS 1, a TFL1-like Gene, Responses to Drought Stress and Regulates Rice Flowering Transition. *Rice (N Y)* **13**, 70.

Werner T, Motyka V, Strnad M, Schmulling T. 2001. Regulation of plant growth by cytokinin. *Proc Natl Acad Sci U S A* **98**, 10487-10492.

Xu J, Zha M, Li Y, Ding Y, Chen L, Ding C, Wang S. 2015. The interaction between nitrogen availability and auxin, cytokinin, and strigolactone in the control of shoot branching in rice (*Oryza sativa* L.). *Plant Cell Rep* **34**,

1647-1662.

Yang J, Cho LH, Yoon J, Yoon H, Wai AH, Hong WJ, Han M, Sakakibara H, Liang W, Jung KH, Jeon JS,

Koh HJ, Zhang D, An G. 2019. Chromatin interacting factor OsVIL2 increases biomass and rice grain yield.

Plant Biotechnol J **17**, 178-187.

Yeh SY, Chen HW, Ng CY, Lin CY, Tseng TH, Li WH, Ku MS. 2015. Down-Regulation of Cytokinin

Oxidase 2 Expression Increases Tiller Number and Improves Rice Yield. Rice (N Y) **8**, 36.

Yu C, Liu Y, Zhang A, Su S, Yan A, Huang L, Ali I, Liu Y, Forde BG, Gan Y. 2015. MADS-box

transcription factor OsMADS25 regulates root development through affection of nitrate accumulation in rice.

PLoS One **10**, e0135196.

Zalabak D, Galuszka P, Mrizova K, Podlesakova K, Gu R, Frebortova J. 2014. Biochemical

characterization of the maize cytokinin dehydrogenase family and cytokinin profiling in developing maize

plantlets in relation to the expression of cytokinin dehydrogenase genes. Plant Physiol Biochem **74**, 283-293.

Zalabak D, Johnova P, Plihal O, Senkova K, Samajova O, Jiskrova E, Novak O, Jackson D, Mohanty A,

Galuszka P. 2016. Maize cytokinin dehydrogenase isozymes are localized predominantly to the vacuoles. Plant

Physiol Biochem **104**, 114-124.

Zalewski W, Galuszka P, Gasparis S, Orczyk W, Nadolska-Orczyk A. 2010. Silencing of the HvCKX1 gene

decreases the cytokinin oxidase/dehydrogenase level in barley and leads to higher plant productivity. J Exp Bot

61, 1839-1851.

Zhang L, Zhao YL, Gao LF, Zhao GY, Zhou RH, Zhang BS, Jia JZ. 2012. TaCKX6-D1, the ortholog of rice

OsCKX2, is associated with grain weight in hexaploid wheat. New Phytol **195**, 574-584.

Zhang W, Peng KX, Cui FB, Wang DL, Zhao JZ, Zhang YJ, Yu NN, Wang YY, Zeng DL, Wang YH,

Cheng ZK, Zhang KW. 2021. Cytokinin oxidase/dehydrogenase OsCKX11 coordinates source and sink

relationship in rice by simultaneous regulation of leaf senescence and grain number. Plant Biotechnol J **19**,

335-350.

Zhao Y, Hu Y, Dai M, Huang L, Zhou DX. 2009. The WUSCHEL-related homeobox gene WOX11 is required to activate shoot-borne crown root development in rice. *Plant Cell* **21**, 736-748.

Zou J, Chen Z, Zhang S, Zhang W, Jiang G, Zhao X, Zhai W, Pan X, Zhu L. 2005. Characterizations and fine mapping of a mutant gene for high tillering and dwarf in rice (*Oryza sativa* L.). *Planta* **222**, 604-612.

FIGURE LEGENDS

Figure 1. Genetic relationship and expression patterns of *OsCKX* genes based on RNA sequencing data and GUS staining assay.

(A) Genetic relationship and expression patterns of *OsCKX* genes investigated in the leaf blade at the vegetative stage (LBV), leaf sheath at the vegetative stage (LSV), root at the vegetative stage (RV), leaf blade at the reproductive stage (LBR), leaf sheath at the reproductive stage (LSR), root at the reproductive stage (RR), stem (S), inflorescence meristem (IM), flower (F), and grain (G). The expression pattern was based on $\log_2$ (Fragments Per Kilobase of exon model per Million mapped fragments (FPKM) + 1) values. (B) Histochemical GUS staining of *pOsCKX::GUS* transgenic plants at various developmental stages. The scale bar in B was 1 mm.

Figure 2. Genetic and phenotypic characterization between wild-type (WT) and *osckx1/2-19* mutant plants.

(A) Mutation details of the coding sequences in the *OsCKX1* and *OsCKX2* of *osckx1/2-19* mutant. Both *OsCKX1* and *OsCKX2* had functional loss due to the 1-bp insertion caused by frameshift mutation. The solid yellow, blue, and red boxes represent the exons, untranslated regions, and target sequences, respectively. The introns are shown as solid lines, while protospacer adjacent motif (PAM) sequences are shown in red color and underlined with red. (B) Number of tillers at the vegetative stage 70 days after sowing. (C) Panicles at the reproductive stage (n = 20 for Nipponbare [NIP], n = 10 for *osckx1/2-19*). (D) Panicle phenotypes of NIP and *osckx1/2* plants. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). (E–H) Measurement of the spikelets per panicle (E), primary branches per panicle (F), secondary branches per panicle (G), 1,000-grain weight (H) (n > 20 for E–G, n = 5 for H) and *osckx1/2-19* (n > 10 for E–G, n = 5 for H) plants. I, Grain phenotypes of NIP and *osckx1/2* plants. The image was digitally extracted and scaled for comparison

(scale bar = 1 cm). J-R, Measurement of the grain length (J), grain width (K), and grain thickness (L), seed setting rate (M), yield per plant (N), length of flag leaves (O), width of flag leaves (P), basal internode diameter (Q), and plant height (R) of NIP ($n > 50$ for J-L, $n > 20$ for M-Q) and *osckx1/2-19* ($n > 50$ for J-L, $n > 10$ for M-Q) plants. The P-values indicate the significant differences between NIP and *osckx1/2-19* determined by Student's *t*-test.

Figure 3. Phenotypic characterization between wild-type (WT) and *osckx4*, *osckx9*, and *osckx4/9* mutant plants from the field experiment.

(A) Phenotypic features of Nipponbare (NIP), *osckx4*, *osckx9* and *osckx4/9* seedlings at the vegetative stage 46 days after sowing. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). (B, C) Number of tillers at the vegetative stage 46 days after sowing (B) and reproductive stage 70 days after sowing (C) ($n > 20$ for NIP, *osckx4-6*, and *osckx9-1*, $n = 10$ for *osckx4/9*). (D, E) Habits (D) and panicle phenotypes (E) of NIP, *osckx4-6*, *osckx9-1*, and *osckx4/9* plants at the mature stage. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). (F-N) Measurement of the panicle number (F), spikelets per panicle (G), primary branches per panicle (H), secondary branches per panicle (I), seed setting rate (J), 1,000-grain weight (K), yield per plant (L), diameter of basal internode (M), and plant height (N) of NIP, *osckx4-6*, *osckx9-1* ($n > 20$ for F-H and J-N, $n = 5$ for I), and *osckx4/9* ($n > 10$ for F-J and L-N, $n = 5$ for K). Different capital letters represent significant differences ($P < 0.01$) determined by one-way ANOVA and shortest significant ranges (SSR) test.

Figure 4. Phenotypic characterization between wild-type (WT) and *osckx4*, *osckx9*, and *osckx4/9* mutant plants from the hydroponic experiment.

(A) Phenotypic features of Nipponbare (NIP), *osckx4*, *osckx9*, and *osckx4/9* seedlings 21 days after transferring to hydroponic solutions. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). (B-E) Measurement of the shoot and root lengths (B), tiller numbers (C), length of 8th leaf (D), and width of 8th leaf (E)

of NIP, *osckx4-6*, *osckx9-1*, and *osckx4/9* seedlings (n = 24 each). Different capital letters represent significant differences ($P < 0.01$) determined by one-way ANOVA and SSR test.

Figure 5. RNA-seq analysis of the roots and shoot bases (BP) between wild-type (WT) and *osckx4*, *osckx9*, and *osckx4/9* mutant plants.

(A) The number of differentially expressed genes (DEGs) identified in the roots and shoot bases of the following pairwise comparisons: Nipponbare (NIP) vs. *osckx4*, NIP vs. *osckx9*, and NIP vs. *osckx4/9*. The changes in gene expression levels were calculated using the $\log_2$ fold change and Q values from three biological replicates. (B) Overlapping DEGs in the roots and shoot bases (BP) of NIP vs. *osckx4*, NIP vs. *osckx9*, and NIP vs. *osckx4/9*. (C) DEGs in *osckx4* and *osckx9* specifically and in common in the roots and shoots, genes in magenta are upregulated; genes in sky blue are downregulated. (D) Top enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways of the DEGs identified in the roots and shoot bases of NIP vs. *osckx4/9*.

Figure 6. *OsCKX4* and *OsCKX9* are associated with several signal transduction pathways.

(A) Differentially expressed genes (DEGs) related to gibberellin metabolism and signaling transduction. (B) DEGs related to strigolactone biosynthesis and signaling transduction. * $Q < 0.05$, ** $Q < 0.01$.

Figure 7. *OsCKX4* and *OsCKX9* associated with nitrogen absorption.

Differentially expressed genes related to nitrogen absorption, metabolism, and transportation. * $Q < 0.05$, ** $Q < 0.01$.

Figure 8. Patterns of *OsCKX* genes in rice.

Figure 1. Genetic relationship and expression patterns of *OsCKX* genes based on RNA sequencing data and *GUS* staining assay.

(A) Genetic relationship and expression patterns of *OsCKX* genes investigated in the leaf blade at the vegetative stage (LBV), leaf sheath at the vegetative stage (LSV), root at the vegetative stage (RV), leaf blade at the reproductive stage (LBR), leaf sheath at the reproductive stage (LSR), root at the reproductive stage (RR), stem (S), inflorescence meristem (IM), flower (F), and grain (G). The expression pattern was based on log2 Fragments Per Kilobase of exon model per Million mapped fragments (FPKM) values. **(B)** Histochemical *GUS* staining of *pOsCKX::GUS* transgenic plants at various developmental stages. The scale bar in B was 1 mm.

Figure 2. Genetic and phenotypic characterization between wild-type (WT) and *osckx1/2-19* mutant plants.

(A) Mutation details of the coding sequences in the *OsCKX1* and *OsCKX2* of *osckx1/2-19* mutant. Both *OsCKX1* and *OsCKX2* had functional loss due to the 1-bp insertion caused by frameshift mutation. The solid yellow, blue, and red boxes represent the exons, untranslated regions, and target sequences, respectively. The introns are shown as solid lines, while protospacer adjacent motif (PAM) sequences are shown in red color and underlined with red. **(B)** Number of tillers at the vegetative stage 70 days after sowing. **(C)** Panicles at the reproductive stage (n = 20 for Nipponbare [NIP], n = 10 for *osckx1/2-19*). **(D)** Panicle phenotypes of NIP and *osckx1/2-19* plants. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). **(E–H)** Measurement of the spikelets per panicle (E), primary branches per panicle (F), secondary branches per panicle (G), 1,000-grain weight (H) (n > 20 for E–G, n = 5 for H) and *osckx1/2-19* (n > 10 for E–G, n = 5 for H) plants. **I**, Grain phenotypes of NIP and *osckx1/2-19* plants. The image was digitally extracted and scaled for comparison (scale bar = 1 cm). **J–R**, Measurement of the grain length (J), grain width (K), and grain thickness (L), seed setting rate (M), yield per plant (N), length of flag leaves (O), width of flag leaves (P), basal internode diameter (Q), and plant height (R) of NIP (n > 50 for J–L, n > 20 for M–Q) and *osckx1/2-19* (n > 50 for J–L, n > 10 for M–Q) plants. The P-values indicate the significant differences

Figure 3. Phenotypic characterization between wild-type (WT) and *osckx4*, *osckx9*, and *osckx4/9* mutant plants from the field experiment.

(A) Phenotypic features of Nipponbare (NIP), *osckx4*, *osckx9* and *osckx4/9* seedlings at the vegetative stage 46 days after sowing. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). **(B, C)** Number of tillers at the vegetative stage 46 days after sowing (B) and reproductive stage 70 days after sowing (C) ($n > 20$ for NIP, *osckx4-6*, and *osckx9-1*, $n = 10$ for *osckx4/9*). **(D, E)** Habits (D) and panicle phenotypes (E) of NIP, *osckx4-6*, *osckx9-1*, and *osckx4/9* plants at the mature stage. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). **(F–N)** Measurement of the panicle number (F), spikelets per panicle (G), primary branches per panicle (H), secondary branches per panicle (I), seed setting rate (J), 1,000-grain weight (K), yield per plant (L), diameter of basal internode (M), and plant height (N) of NIP, *osckx4-6*, *osckx9-1* ($n > 20$ for F–H and J–N, $n = 5$ for I), and *osckx4/9* ($n > 10$ for F–J and L–N, $n = 5$ for K). Different capital letters represent significant differences ($P < 0.01$) determined by one-way ANOVA and shortest significant ranges (SSR) test.

Figure 4. Phenotypic characterization between wild-type (WT) and *osckx4*, *osckx9*, and *osckx4/9* mutant plants from the hydroponic experiment. **(A)** Phenotypic features of Nipponbare (NIP), *osckx4-6*, *osckx9-1*, and *osckx4/9* seedlings 21 days after transferring to hydroponic solutions. The image was digitally extracted and scaled for comparison (scale bar = 10 cm). **(B–E)** Measurement of the shoot and root lengths (B), tiller numbers (C), length of 8th leaf (D), and width of 8th leaf (E) of NIP, *osckx4-6*, *osckx9-1*, and *osckx4/9* seedlings (n = 24 each). Different capital letters represent significant differences ($P < 0.01$) determined by one-way ANOVA and SSR test.

Figure 5. RNA-seq analysis of the roots and shoot bases (BP) between wild-type (WT) and *osckx4*, *osckx9*, and *osckx4/9* mutant plants.

(A) The number of differentially expressed genes (DEGs) identified in the roots and shoot bases of the following pairwise comparisons: Nipponbare (NIP) vs. *osckx4*, NIP vs. *osckx9*, and NIP vs. *osckx4/9*. The changes in gene expression levels were calculated using the $\log_2$ fold change and Q values from three biological replicates. **(B)** Overlapping DEGs in the roots and shoot bases (BP) of NIP vs. *osckx4*, NIP vs. *osckx9*, and NIP vs. *osckx4/9*. **(C)** DEGs in *osckx4* and *osckx9* specifically and in common in the roots and shoots, genes in magenta are upregulated; genes in sky blue are downregulated. **(D)** Top enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways of the DEGs identified in the roots and shoot bases of NIP vs. *osckx4/9*.

Figure 7. OsCKX4 and OsCKX9 associated with nitrogen absorption.

Differentially expressed genes related to nitrogen absorption, metabolism, and transportation.

* Q < 0.05, ** Q < 0.01.

Figure 8. Patterns of *OsCKX* genes in rice.