mtDNA replication in dysfunctional mitochondria promotes deleterious heteroplasmy via the UPR^{mt}

Authors:

Qiyuan Yang¹, Pengpeng Liu¹, Nadine S. Anderson¹, Tomer Shpilka¹, YunGuang Du¹, Nandhitha Uma Naresh¹, Kevin Luk¹, Josh Lavelle¹, Rilee D. Zeinert^{2,3}, Peter Chien^{2,3}, Scot A. Wolfe¹, Cole M. Haynes^{1*}

Affiliations:

¹Department of Molecular, Cell and Cancer Biology, University of Massachusetts Medical School, Worcester, MA 01605, USA.

²Department of Biochemistry and Molecular Biology, University of Massachusetts Amherst, MA 01002, USA

³Molecular and Cellular Biology Graduate Program, University of Massachusetts Amherst, MA 01002, USA

Lead Contact

*Correspondence to: cole.haynes@umassmed.edu

Key words: ATFS-1; LONP-1; heteroplasmy; mitochondrial DNA (mtDNA).

Highlights:

1. OXPHOS dysfunction increases mtDNA content via ATFS-1 and POLG.
2. The bZIP protein ATFS-1 accumulates in dysfunctional mitochondria and binds mtDNA.
3. In heteroplasmic worms, ATFS-1 and POLG binding is enriched on mutant mtDNAs relative to wildtype mtDNAs.
4. The enriched POLG/mtDNA interaction is regulated by degradation of ATFS-1 via the mtDNA-bound protease LONP-1.
5. Inhibition of LONP-1 in *C. elegans* and human cells improves heteroplasmy and mitochondrial function.

ABSTRACT

The accumulation of deleterious mitochondrial genomes (Δ mtDNAs) underlies inherited mitochondrial diseases and contributes to the aging-associated decline in mitochondrial function. In heteroplasmic cells, oxidative phosphorylation (OXPHOS) function declines as the population of Δ mtDNAs increase relative to wildtype mtDNAs. In response to mitochondrial perturbations, the bZIP protein ATFS-1 induces a transcription program to promote the recovery of mitochondrial function. Paradoxically, ATFS-1 is also required to maintain Δ mtDNAs in heteroplasmic worms. However, the mechanism(s) by which ATFS-1 promotes Δ mtDNA accumulation relative to wildtype mtDNAs is unclear. Here, we show that mitochondrial-localized ATFS-1 binds almost exclusively to Δ mtDNAs in heteroplasmic worms. Moreover, we demonstrate that mitochondrial ATFS-1 promotes the preferential binding of the mtDNA replicative polymerase (POLG) to Δ mtDNAs. Interestingly, inhibition of the mtDNA-bound protease LONP-1 increased ATFS-1 and POLG binding to wildtype mtDNAs. Furthermore, LONP-1 inhibition in *C. elegans* and human cybrid cells improved the heteroplasmy ratio and restored OXPHOS function. Our findings suggest that ATFS-1 promotes mtDNA replication by recruiting POLG to mtDNA in a manner that is antagonized by LONP-1. We speculate that this mechanism promotes the repair and expansion of the mitochondrial network by synchronizing mtDNA replication with UPR^{mt} activation driven by nuclear ATFS-1 activity. However, this repair mechanism cannot resolve OXPHOS defects in mitochondria harboring Δ mtDNAs, resulting in an accumulation of ATFS-1 in dysfunctional mitochondria and constitutive replication of Δ mtDNAs.

INTRODUCTION

Mitochondria provide numerous metabolic functions including being the site of energy production via oxidative phosphorylation (OXPHOS). Most of the ~1200 proteins comprising the mitochondrial proteome are encoded by nuclear genes and must be imported into each mitochondrion following synthesis on cytosolic ribosomes¹. However, 13 essential OXPHOS components alongside the tRNAs and rRNAs required for their synthesis are encoded by mitochondrial genomes (mtDNAs), which reside in the mitochondrial matrix. Each mitochondrion harbors at least one mtDNA, and most cells harbor 100s-1,000s of mtDNAs.

Mitochondrial dysfunction accumulates as cells age and is accelerated in multiple age-associated diseases, including Alzheimer's Diseases and Parkinson's Disease. However, a variety of mitochondrial syndromes or diseases are caused by inherited mutations that impair OXPHOS function. The disease causing mutations can occur in genes required for OXPHOS encoded by either the nuclear genome, or mtDNAs, which acquire mutations at a significantly higher rate than the nuclear genome². A number of single nucleotide variants and deletions are associated with inherited mitochondrial diseases and syndromes affecting ~1:4000 individuals³. Because of the high number of mtDNAs per cell, a single mutant mtDNA has little impact. To cause the OXPHOS dysfunction that underlies mitochondrial diseases, the mutant mtDNA must accumulate to ~60% of the total cellular mtDNAs. The mixture of mutant mtDNAs and wildtype mtDNAs is known as heteroplasmy. Several studies using mitochondrial-targeted nucleases that specifically cleave mutant mtDNAs have indicated that a relatively modest reduction in the percentage of Δ mtDNAs is sufficient to improve mitochondrial function in

heteroplasmic cells and mouse tissues⁴⁻⁶.

The initial mtDNA mutation or deletion likely occurs because an error in mtDNA replication⁷. Two mechanisms are thought to contribute to the “clonal expansion” of the Δ mtDNA. In dividing cells, random, or non-selective, genetic drift can disproportionately increase either genome. An alternative model suggests that large mtDNA deletions allow for quicker replication simply because these genomes are smaller and thus replicate more rapidly^{7, 8}. Consistent with these models, mutations that reduce the function of the replicative mtDNA polymerase POLG cause a preferential depletion of mutant mtDNAs and improve heteroplasmy^{6,9}. However, the underlying mechanism(s) that drive the clonal amplification of Δ mtDNAs and maintain them at a high enough percentage to cause OXPHOS defects in mitochondrial diseases¹⁰⁻¹³, aging¹⁴ and Parkinson’s Disease¹⁵ remain unresolved.

Previously, the bZIP protein ATFS-1 was shown to be required to maintain deleterious heteroplasmy in *C. elegans*^{9, 16}. ATFS-1 harbors both a mitochondrial targeting sequence (MTS) and a nuclear localization sequence (NLS) (Fig. 1a) and regulates a transcriptional program known as the mitochondrial unfolded protein response (UPR^{mt})¹⁷. Under basal conditions, the majority of ATFS-1 is imported into mitochondria, where it is degraded by the protease LONP-1. Mitochondrial dysfunction reduces mitochondrial import capacity, resulting in a percentage of ATFS-1 trafficking to the nucleus, where it activates a transcriptional program that re-establishes mitochondrial homeostasis and promotes mitochondrial biogenesis¹⁸. Importantly, both nuclear and mitochondrial accumulation of ATFS-1 are required for development during mitochondrial dysfunction^{9, 19}. However, the function of ATFS-1 within mitochondria is unclear.

Here, we report that the maintenance of deleterious heteroplasmy requires the accumulation of ATFS-1 within mitochondria. We demonstrate that in a heteroplasmic *C. elegans* strain, ATFS-1 binds predominantly to Δ mtDNAs. Moreover, the replicative polymerase POLG also binds predominantly to Δ mtDNAs, and that this interaction depends on ATFS-1. Lastly, we demonstrate that the mitochondrial protease LONP-1, which degrades ATFS-1 in functional mitochondria^{17, 19}, is required to establish the preferential interaction between ATFS-1 and Δ mtDNAs in heteroplasmic worms. Our findings in *C. elegans* are conserved in cultured human cells as we found that inhibition of LONP1 by siRNA or the drug CDDO²⁰, improves heteroplasmy and rescues OXPHOS function in heteroplasmic cells harboring either a mutant mtDNA with a large deletion or with a mutant mtDNA harboring a single base pair substitution.

RESULTS

OXPHOS perturbations increase mtDNA quantity

OXPHOS proteins are encoded by genes located within the mitochondria and the nucleus. The heteroplasmic *C. elegans* strain *uaDf5* harbors approximately 40% wildtype mtDNAs and 60% Δ mtDNAs that lack four essential OXPHOS protein coding genes (Fig. 1b)²¹. As shown previously, *uaDf5* worms have impaired respiration²², constitutive UPR^{mt} activation, as determined by increased expression of the *hsp-6_{pr}::gfp* UPR^{mt} reporter^{22, 23} (Fig. 1c), and an increase in total mtDNAs (Fig. 1d)²¹.

To examine the impact of OXPHOS perturbation on mtDNA content, we impaired several OXPHOS components in wildtype homoplasmic worms. As expected, worms raised on *cco-1*(RNAi) (complex IV) or *cyc-1*(RNAi) (cytochrome c) had increased *hsp-6_{pr}::gfp* activation, as did strains with loss of function mutations *isp-1(qm150)* (complex III) or *clk-1(qm30)* (ubiquinone biosynthesis) (Figs. 1e,f). Intriguingly, each of the OXPHOS perturbations resulted in increased mtDNA content as determined by qPCR (Figs. 1g,h and Supplementary Fig. 1a), which is consistent with previous reports^{24, 25}. Importantly, the increase in mtDNAs caused by OXPHOS perturbation was impaired in *atfs-1(null)* worms that lack the entire *atfs-1* open reading frame²⁶ (Fig. 1g) as well as worms raised on *atfs-1*(RNAi) (Fig. 1h). Similarly, the increase in total mtDNAs in heteroplasmic worms was also reduced when raised on *atfs-1*(RNAi) (Fig. 1d). Combined, these findings indicate that the increased mtDNA content in both homoplasmic and heteroplasmic worms caused by OXPHOS perturbation requires *atfs-1*.

Increased POLG-mtDNA binding during OXPHOS perturbation requires ATFS-1

We next sought to gain insight into the mechanism by which mtDNAs are increased during OXPHOS perturbation. We previously found that ATFS-1 accumulates within mitochondria when mitochondrial function is perturbed by inhibiting the essential mitochondrial protease SPG-7 or by raising worms in the presence of ethidium bromide, which impairs mtDNA replication^{17, 27}. Intriguingly, ChIP-sequencing indicated that during mitochondrial dysfunction, ATFS-1 binds mtDNAs at a single site within the non-coding region (NCR)²⁸ (Fig. 2a), which in mammals contains sequence elements that regulate mtDNA replication²⁹. Here, we found that ATFS-1 also accumulates in the mitochondrial fraction of worms raised on *cco-1*(RNAi) (Fig. 2b). Furthermore, ATFS-1 ChIP followed by qPCR indicated that ATFS-1 also interacts with mtDNA when worms are raised on *cco-1*(RNAi), which was impaired in *atfs-1*(null) worms (Fig. 2c). Combined these data suggest that ATFS-1 accumulates within mitochondria and interacts with mtDNAs when OXPHOS is impaired.

To further explore the relationship between ATFS-1 and mtDNA accumulation during OXPHOS perturbation, we generated POLG antibodies which detected a ~120 KD band that co-fractionated with the OXPHOS protein NDUF53 (Supplementary Fig. 1b) and was depleted by *polg*(RNAi) (Supplementary Fig. 1c). Similar to ATFS-1, POLG protein levels increased within mitochondria when raised on *cco-1*(RNAi) (Fig. 2d), and POLG interacted with more mtDNAs (Fig. 2e). To determine if the increased POLG-mtDNA interaction during OXPHOS perturbation required *atfs-1*, we also performed POLG ChIP-mtDNA in wildtype and *atfs-1*(null) worms raised on *cco-1*(RNAi). Interestingly, the increased POLG-mtDNA interaction in wildtype worms raised on *cco-*

1(RNAi) was impaired in *atfs-1(null)* worms, suggesting that ATFS-1 is required for increased POLG-mtDNA binding during OXPHOS dysfunction (Fig. 2e).

Our data indicate that ATFS-1 is required for POLG to bind mtDNA during OXPHOS perturbation. We next sought to examine if the increased POLG-mtDNA binding required nuclear-localized ATFS-1. To circumvent the potentially confounding effects of the induction of *polg* mRNA by ATFS-1 during mitochondrial stress^{9, 17}, we first generated the *atfs-1^{nuc(-)}* strain in which the NLS within ATFS-1 was impaired via genome editing (Supplementary Figs. 2a,b). Importantly, the *atfs-1^{nuc(-)}* strain was unable to induce *hsp-6* mRNA when raised on *spg-7*(RNAi), consistent with impaired nuclear function (Supplementary Fig. 2c). Furthermore, we introduced the *atfs-1^{nuc(-)}* mutation into *atfs-1(et18)* worms, which causes constitutive induction of the UPR^{mt} reporter *hsp-6_{pr}::gfp* and *hsp-6* mRNA due to a mutation that perturbs the MTS of ATFS-1³⁰. The *atfs-1^{nuc(-)}* mutation in *atfs-1(et18)* worms suppressed activation of the UPR^{mt} reporter *hsp-6_{pr}::gfp*, and *hsp-6* mRNA (Supplementary Figs. 2e-g). Induction of *hsp-6* and *polg* mRNA when raised on *cco-1*(RNAi) was also impaired in *atfs-1^{nuc(-)}* worms consistent with impaired nuclear transcription (Supplementary Figs. 2g,h). Furthermore, unlike in wildtype worms, POLG protein levels were not increased in *atfs-1^{nuc(-)}* worms indicating that *atfs-1^{nuc(-)}* worms were unable to regulate nuclear transcription during OXPHOS perturbation (Supplementary Figs. 2h,i). Importantly, ATFS-1^{nuc(-)} protein accumulated within mitochondria to similar levels as wildtype ATFS-1 upon LONP-1 inhibition (Supplementary Fig. 2j) indicating the protein was expressed and processed similarly to wildtype ATFS-1. Lastly, ATFS-1^{nuc(-)} bound similar amounts of mtDNA as wildtype ATFS-1 when raised on *cco-1*(RNAi) as determined by ChIP (Figs. 2c,f).

To determine if the nuclear function of ATFS-1 is required for POLG to bind mtDNAs during OXPHOS perturbation, we examined the POLG-mtDNA interaction in *atfs-1(null)* and *atfs-1^{nuc(-)}* worms raised on control or *cco-1*(RNAi). Strikingly, significantly more POLG was bound to mtDNA in *atfs-1^{nuc(-)}* worms relative to *atfs-1(null)* worms (Fig. 2g), suggesting that the *atfs-1*-dependent increase in POLG-mtDNA binding during OXPHOS perturbation does not require the nuclear activity of ATFS-1. mtDNA content also increased in *atfs-1^{nuc(-)}* worms upon OXPHOS perturbation caused by *cco-1*(RNAi) (Fig. 2h) as well as in *clk-1(qm30)* worms (Fig. 2i), similar to ATFS-1 wildtype worms. Combined, these results suggest that the accumulation of ATFS-1 within mitochondria during OXPHOS perturbation is sufficient to increase POLG-mtDNA binding and mtDNA content.

Degradation of ATFS-1 by LONP-1 within mitochondria impairs mtDNA propagation

We next sought to determine the events leading to the accumulation of ATFS-1 within mitochondria when OXPHOS is perturbed. During normal growth, the ATP-dependent protease LONP-1 degrades the majority of ATFS-1 once the bZIP protein has been imported into the mitochondrial matrix¹⁷. LONP-1 is an ATP-dependent protease known to recognize and degrade mitochondrial proteins damaged by reactive oxygen species³¹. LONP-1 has also been shown to interact with mtDNA in diverse species (Fig. 1a)^{32, 33}, and regulate mtDNA replication^{34, 35}.

To further examine the interaction between LONP-1 and mtDNA in *C. elegans*, we generated a strain in which the C-terminus of LONP-1 was tagged with the FLAG epitope via genome editing (Fig. 3a and Supplementary Fig. 3a). Introduction of the FLAG epitope

did not impair worm development or cause UPR^{mt} activation, suggesting it did not adversely affect LONP-1 function (Supplementary Figs. 3b,c). Consistent with findings in other species^{32, 33}, LONP-1 interacted with mtDNA in *C. elegans* as determined by LONP-1^{FLAG} ChIP-mtDNA qPCR (Fig. 3b). We next examined where in mtDNA LONP-1^{FLAG} binds in wildtype homoplasmic worms. LONP-1^{FLAG} ChIP-seq indicated that the protease was enriched at several G-rich sites throughout mtDNA (Fig. 3c), but was especially enriched within the NCR (Fig. 3d). Interestingly, the strongest LONP-1^{FLAG} peak within the NCR overlapped with the ATFS-1 binding site (Fig. 3d and Supplementary Fig. 3d) suggesting a potential role in mediating the ATFS-1-mtDNA interaction.

We next examined the impact of *lonp-1*(RNAi) on mtDNA accumulation in homoplasmic wildtype worms. When raised on *lonp-1*(RNAi), the binding of ATFS-1 to mtDNA increased by ~6-fold (Fig. 3e) which correlated with an increase in mtDNA content (Fig. 3f). Importantly, exposure to *lonp-1*(RNAi) also increased the binding of POLG to mtDNA (Fig. 3g). Moreover, LONP-1 inhibition increased mtDNA content in *atfs-1^{nuc(-)}* worms (Supplementary Fig. 3e), but not in *atfs-1(null)* worms (Fig. 3f). In short, these findings support a role for mitochondrial-localized ATFS-1 in promoting mtDNA replication which is impaired by LONP-1-dependent degradation (Fig. 3h). While it remains to be determined how LONP-1 mediates ATFS-1 accumulation, we propose that 1) the decrease of ATP within perturbed mitochondria or 2) the increased accumulation of misfolded proteins that are the primary substrates of the protease³⁶ may impair LONP-1 function and allow ATFS-1 to accumulate within dysfunctional mitochondria.

Maintenance of deleterious heteroplasmy does not require ATFS-1 nuclear activity

We and the Patel Lab previously found that ATFS-1 is required to maintain Δ mtDNAs^{9, 16} in a heteroplasmic worm strain²¹. Here, we crossed the *atfs-1(null)* allele into *uaDf5* heteroplasmic worms. Consistent with previous results, Δ mtDNAs were severely depleted in heteroplasmic *atfs-1(null)* worms (Fig. 4a). The loss of Δ mtDNAs in the absence of *atfs-1* could either be due to increased mitophagy of mitochondria harboring Δ mtDNAs^{16, 37}, or decreased replication of Δ mtDNAs. To examine the role of mitophagy, we generated heteroplasmic strains lacking the mitophagy component Parkin (PDR-1 in worms) (Supplementary Fig. 4a). As expected, *pdr-1*-deficient worms had increased Δ mtDNAs relative to wildtype worms consistent with mitophagy limiting the accumulation of Δ mtDNAs^{22, 23}. However, *atfs-1(null);pdr-1(tm598)* worms had a significant reduction in Δ mtDNAs relative to *pdr-1(tm598)* worms, indicating that ATFS-1 promotes heteroplasmy via a mechanism independent of mitophagy.

As OXPHOS perturbation increased mtDNA content in homoplasmic wildtype worms via ATFS-1, we hypothesized that OXPHOS dysfunction may contribute to the increased mtDNA content in heteroplasmic worms (Fig. 1d). We previously found that heteroplasmic worms consumed less oxygen than wildtype worms⁹. To further evaluate mitochondrial function, we examined mitochondrial membrane potential by staining with tetramethylrhodamine ethyl ester (TMRE). As expected, TMRE staining was decreased in heteroplasmic worms relative to wildtype worms, but stronger than in *cco-1(RNAi)*- or *spg-7(RNAi)*-treated worms consistent with intermediate OXPHOS function (Figs. 4b,c and Supplementary Figs. 4b,c). Importantly, ATFS-1 also accumulated within the mitochondrial fraction of heteroplasmic worms, as in worms raised on *spg-7(RNAi)* or *cco-1(RNAi)* (Fig. 4d, also see Fig. 2b). Combined, these results indicate that ATFS-1

accumulates within dysfunctional mitochondria caused by impairment of either a nuclear-encoded OXPHOS component, a nuclear-encoded mitochondrial protease, or in worms harboring Δ mtDNAs.

We next examined if maintenance of Δ mtDNAs required the nuclear activity of ATFS-1 by crossing the *atfs-1^{nuc(-)}* allele into the heteroplasmic background. Consistent with impaired nuclear activity, *hsp-6_{pr}::gfp* was not increased in heteroplasmic *atfs-1^{nuc(-)}* worms (Supplementary Figs. 4d,e). Impressively, unlike *atfs-1(null)* worms, *atfs-1^{nuc(-)}* worms were able to maintain Δ mtDNAs to nearly the same level as wildtype *atfs-1* worms (Supplementary Fig. 4f) indicating the role of ATFS-1 in promoting heteroplasmy is largely independent of the nuclear activity. Consistent with the accumulation of ATFS-1 in dysfunctional mitochondria (Figs. 2b and 4d), ATFS-1 bound more total mtDNAs in heteroplasmic worms than in wildtype homoplasmic worms as determined by ChIP-mtDNA (Fig. 4e). Similarly, POLG also interacted with more total mtDNAs in heteroplasmic worms than in wildtype homoplasmic worms suggesting increased mtDNA replication in heteroplasmic worms (Supplementary Fig. 4g). Combined, these findings suggest a role for mitochondrial-localized ATFS-1 in maintaining deleterious mtDNA heteroplasmy.

In heteroplasmic worms, ATFS-1 and POLG primarily bind to Δ mtDNAs

To gain insight into the mechanism by which ATFS-1 maintains heteroplasmy, we again examined the ATFS-1-mtDNA interaction. Because wildtype mtDNAs and Δ mtDNAs both harbor the ATFS-1 binding site, we sought to determine if ATFS-1 differentially interacted with each genome. The interaction between ATFS-1 and wildtype

mtDNAs or Δ mtDNAs was evaluated via qPCR or 3D digital PCR following ATFS-1 ChIP (Fig. 4f)²². As before, qPCR of mtDNA from heteroplasmic whole worm lysate indicated the strain harbored ~60% Δ mtDNAs and ~40% wildtype mtDNAs (Fig 1b), (Supplementary Figs. 5a,b). qPCR following ATFS-1 ChIP indicated that of the mtDNAs that interacted with ATFS-1, 90% were Δ mtDNAs and 10% were wildtype mtDNAs, indicating that ATFS-1 is significantly enriched on Δ mtDNAs (Fig. 4g and Supplementary Figs. 5c,d).

Previously, we found that inhibition of POLG caused depletion of Δ mtDNAs in heteroplasmic worms relative to wildtype mtDNAs⁹, which is similar to findings in a *Drosophila* model of deleterious mtDNA heteroplasmy³⁸. Both findings suggest increased replication of Δ mtDNAs in heteroplasmic cells. To further explore the relationship between ATFS-1 and mtDNA replication, we performed POLG ChIP-mtDNA. Interestingly, ChIP-mtDNA using the POLG antibody indicated that the replicative polymerase also interacted with ~90% Δ mtDNAs and 10% wildtype mtDNAs (Fig. 4h and Supplementary Figs. 5e,f), similarly to ATFS-1 (Fig. 4g). As a control, we generated antibodies to the mtDNA packaging protein HMG-5 (TFAM in mammals) (Supplementary Figs. 5g,h), which interacts with mtDNAs independent of replication³⁹. In contrast to ATFS-1 and POLG, the percentage of Δ mtDNAs bound to HMG-5 reflected the percentage within the whole worm lysate (Fig. 4i). Combined, these data indicate that ATFS-1 and a component of the replisome are enriched on Δ mtDNAs relative to wildtype mtDNAs in heteroplasmic worms, consistent with the mutant mtDNA having a replicative advantage.

The protease LONP-1 is required to maintain heteroplasmy

Because LONP-1 binds mtDNA and its proteolytic activity prevents ATFS-1 accumulation in functional mitochondria¹⁷ (See [Figs. 3d,e](#)), we examined the role of LONP-1 in heteroplasmy maintenance. We generated antibodies to *C. elegans* LONP-1 that recognized a ~130 KD band that was reduced when raised on *lonp-1*(RNAi) ([Supplementary Fig. 6a](#)). Via ChIP-qPCR, we found that LONP-1 binds ~60 fold more mtDNAs than ATFS-1 in wildtype worms ([Supplementary Fig. 6b](#)). In heteroplasmic worms, LONP-1 bound similar percentages of wildtype and Δ mtDNAs suggesting that unlike ATFS-1 and POLG-1, LONP-1 interacts with mtDNAs independent of mitochondrial dysfunction ([Fig. 5a and Supplementary Fig. 6c](#)). Combined, these data suggest that LONP-1 is constitutively bound to mtDNAs and heteroplasmy is not maintained by uneven mtDNA binding by the protease.

Unlike ATFS-1 and POLG, LONP-1 interacts equally with both wildtype mtDNAs and Δ mtDNAs ([Fig. 5a and Supplementary Fig. 6c](#)), suggesting heteroplasmy is not maintained by uneven mtDNA binding by the protease. We next examined the effect of inhibiting LONP-1 on heteroplasmy. Strikingly, *lonp-1* inhibition via RNAi increased wildtype mtDNAs ([Fig. 5b](#)) and reduced Δ mtDNAs, which improved the heteroplasmy ratio from 59% Δ mtDNAs to 25% ([Fig. 5c](#)). Similar results were obtained in *atfs-1^{nuc(-)}* worms upon LONP-1 inhibition ([Supplementary Fig. 6d](#)). Importantly, *lonp-1*(RNAi) did not reduce the brood size of heteroplasmic worms ([Supplementary Fig. 6e](#)), suggesting that *lonp-1*(RNAi) did not select against embryos with dysfunctional mitochondria.

We next examined the interaction between ATFS-1 and mtDNAs in heteroplasmic worms upon *lonp-1* inhibition. Interestingly, when raised on *lonp-1*(RNAi), the percentage of Δ mtDNAs and wildtype mtDNAs bound to ATFS-1 nearly reflected the heteroplasmy

percentage within the whole worm lysate (Fig. 5d). Thus, the mtDNA-bound protease is required to establish the enriched interaction between ATFS-1 and Δ mtDNAs. In addition to increasing the percentage of wildtype mtDNAs bound by ATFS-1, *lonp-1*(RNAi) also increased the percentage of POLG that interacted with wildtype mtDNAs (Fig. 5e, also see Fig. 4h). However, *lonp-1*(RNAi) did not alter the percentage of HMG-5/TFAM bound to Δ mtDNAs, consistent with HMG-5 interacting with all mtDNAs independent of replication (Supplementary Fig. 6f). In support of this interpretation, when heteroplasmic worms raised on *cco-1*(RNAi) or in *clk-1* mutant strain, wildtype mtDNAs were specifically increased (Supplementary Figs. 6g,h). Combined, these findings suggest that LONP-1-mediated proteolysis antagonizes the ability of mitochondrial ATFS-1 to stimulate mtDNA replication. LONP-1 activity may be compromised within mitochondrial compartments harboring Δ mtDNAs, leading to ATFS-1 accumulation and mtDNA replication. We propose that globally inhibiting LONP-1 promotes ATFS-1-mediated mtDNA replication throughout the cell, not just in compartments enriched in Δ mtDNAs, consequently leading to a recovery of wildtype mtDNAs and a depletion of Δ mtDNAs.

LONP1 inhibition improves heteroplasmy and OXPHOS function in cybrid cells

Lastly, we examined if the role of LONP1 in maintaining Δ mtDNAs is conserved in mammals by examining two human heteroplasmic cybrid cell lines, which harbor a combination of wildtype mtDNA and Δ mtDNAs associated with mitochondrial disease⁴⁰. While cybrid cells are often used as models of deleterious mtDNA heteroplasmy, it is important to note that cybrid cells are cancer cells in which patient-derived heteroplasmic mtDNAs were introduced by cell fusion. Thus, these lines may not replicate all aspects of

deleterious heteroplasmy observed in affected patient tissues.

One cybrid line harbors a single nucleotide transition (COXI G6930A) that introduces a premature stop codon in the cytochrome c oxidase subunit I gene, which was isolated from a patient with a multisystem mitochondrial disorder (Fig. 6a)⁴¹. We also examined a cybrid line harboring a 4977 base pair deletion known as the “common deletion” which removes multiple OXPHOS genes and is associated with Kearns-Sayre Syndrome (KSS), progressive external ophthalmoplegia, cancer and aging (Fig. 6a)⁴²⁻⁴⁵. We first examined the impact of LONP1 siRNA on heteroplasmy in the KSS cybrid line. Similar to inhibition of *C. elegans* LONP-1, inhibition of human LONP1 by siRNA for 4 days (Fig. 6b), resulted in a 1.5-fold increase of wildtype mtDNAs (Fig. 6c) while KSS mtDNAs were decreased ~2 fold (Fig. 6d), resulting in a shift in the heteroplasmy ratio from 57.5% to 25.6% (Fig. 6d). This result suggests that the role of LONP1 in promoting heteroplasmy is conserved in mammals.

To further investigate the impact of LONP1 protease activity on heteroplasmy, we used the LONP1 inhibitor CDDO (2-cyano-3,12-dioxo-oleana-1,9(11)-dien-28-oic acid), also known as Bardoxolone (Fig. 6e)^{20,46}. As determined by deep sequencing, the CoxI G6930A cybrid line initially harbored ~90% G6930A mutant mtDNAs and ~10% wildtype mtDNAs (Figs. 6a,f). Impressively, incubation with 0.1 μ M or 0.25 μ M CDDO for 3 weeks resulted in depletion of the CoxI G6930A mtDNA from 86% to 68% and 72%, respectively, with a concomitant increase in wildtype mtDNAs (Fig. 6f). Furthermore, continuous incubation with 0.1 μ M or 0.25 μ M CDDO for 18.5 weeks further decreased the heteroplasmic ratios from ~90% to ~47% and 62%, respectively (Figs. 6f and Supplementary Fig. 7a). Similar results were obtained when the KSS cybrid cells were

incubated with CDDO. As determined by qPCR, KSS cells initially harbored ~50% Δ mtDNAs. Incubation with 0.1 μ M or 0.25 μ M CDDO for 4 weeks and 13 weeks depleted Δ mtDNAs to 19.5%, 20.3 %, 18.6% and 10.7%, respectively (Fig. 6g). Importantly, neither 0.1 μ M nor 0.25 μ M CDDO affected viability or mitochondria respiration of the KSS, Coxl G6930A or 143b wildtype cell lines (Fig. 6h and Supplementary Figs. 7b,c) suggesting that the improved heteroplasmy ratio was not due to selection of cells with low levels of mutant mtDNAs.

Last, we determined the impact of the CDDO-dependent shifts in heteroplasmy on OXPHOS. Mitochondrial respiratory function was measured throughout the time course (Figs. 7a-d). Impressively, incubation of KSS cells with 0.1 μ M or 0.25 μ M CDDO for 4 or 13 weeks resulted in significant increases in basal respiration suggesting improved OXPHOS (Supplementary Fig. 7d). Furthermore, the improved heteroplasmy caused by CDDO in Coxl G6930A mtDNA also resulted in increased basal respiration and maximal respiratory capacity (Figs. 7b-d). For example, 3 weeks exposure to 0.1 μ M CDDO improved basal oxygen consumption ~2-fold, while exposure for 18.5 weeks improved basal oxygen consumption over 3-fold (Figs. 7b,c). Taken together, these findings suggest that inhibition of LONP1 improves deleterious heteroplasmy and recovers mitochondrial respiration. Notably, these phenotypes are independent of mtDNA length, as maintenance of mutant mtDNAs with either a large deletion or a single base pair substitution require LONP1 function.

Discussion

The underlying mechanisms that govern heteroplasmy dynamics are largely unknown, however it has been proposed that mutant genomes have a selective advantage that promotes their accumulation. Previously, we and others reported that heteroplasmy depends on the transcription factor ATFS-1^{9, 16}. In wildtype homoplasmic worms raised under normal conditions, ATFS-1 is imported into mitochondria and degraded by the protease LONP-1¹⁷. However, during mitochondrial dysfunction, a fraction of ATFS-1 accumulates within mitochondria and the nucleus. Moreover, we also reported that ATFS-1 binds to mtDNA within dysfunctional mitochondria¹⁹. Here, we report that mitochondrial-localized ATFS-1 promotes the accumulation of mtDNAs upon OXPHOS perturbation. Importantly, ATFS-1 and POLG binding to mtDNA was increased upon OXPHOS perturbation.

Deleterious heteroplasmy also causes OXPHOS dysfunction⁴⁷ and increased mtDNA content which also required *atfs-1*. Furthermore, binding of ATFS-1 and POLG was also increased in heteroplasmic worms. Strikingly, ATFS-1 and POLG were both enriched on Δ mtDNAs by ~9:1 (Figs. 4g and 4h). As the ATFS-1 binding site is present in both wildtype and Δ mtDNAs, we propose that ATFS-1 accumulates within dysfunctional compartments, which in heteroplasmic worms harbor Δ mtDNAs. In support of this model, the nuclear activity of ATFS-1 is not required for the increased mtDNAs caused by *cco-1*(RNAi) in homoplasmic worms, or for maintaining Δ mtDNAs in heteroplasmic worms. Furthermore, nuclear ATFS-1 is not required for the increased POLG-mtDNA binding during OXPHOS perturbation in homoplasmic worms. Collectively, these results suggest that mitochondrial accumulation of ATFS-1 promotes recruitment of the mitochondrial replisome to mtDNA during OXPHOS dysfunction. It is important to note that additional

regulatory mechanisms of mtDNA replication are likely involved during normal development as homoplasmic *atfs-1(null)* worms have only a slight reduction in mtDNAs (Fig. 3f).

We find that degradation of ATFS-1 by the mtDNA-bound protease LONP-1 is required to maintain heteroplasmy. In homoplasmic worms, knockdown of LONP-1 increases total mtDNA levels in an ATFS-1-dependent manner and increases POLG binding to mtDNA (Figs. 3f,g), consistent with a role for LONP-1 in regulating mtDNA replication^{35, 48, 49}. In heteroplasmic worms, knockdown of LONP-1 also increases wildtype mtDNAs, but concomitantly reduces Δ mtDNA levels, improving heteroplasmy. Moreover, knockdown of LONP-1 abolishes the enriched binding of ATFS-1 and POLG to Δ mtDNAs (Figs. 5d,e). These results support a model wherein LONP-1 antagonizes the ability of ATFS-1 to promote mtDNA replication in healthy mitochondria. Because LONP-1 is an ATP-dependent protease that degrades proteins damaged by reactive oxygen species³¹, mitochondrial dysfunction may impede degradation of ATFS-1 by LONP-1, resulting in recruitment of POLG to mtDNA. We propose that this mechanism may serve to coordinate mtDNA replication with expansion of the mitochondrial network during cell growth or recovery from mitochondrial dysfunction. However, if compartmental dysfunction is caused by an enrichment of Δ mtDNAs, they are inadvertently, but preferentially, replicated. Our data suggests that inhibiting LONP1 throughout the mitochondrial network negates this preferential replication, leading to a reduction in the heteroplasmic ratio and recovery of mitochondrial function.

There are currently no FDA-approved treatments for diseases caused by mutant mtDNAs. However, the TORC1 inhibitor rapamycin has been shown to improve

heteroplasmy in cybrid cells⁵⁰. The findings indicate that inhibition of TORC1 in cybrid results in increased autophagy of dysfunctional mitochondria. It is perhaps interesting to note that inhibition of several TORC1 components also inhibit ATFS-1 function and *atfs-1*-dependent mitochondrial biogenesis in *C. elegans*¹⁸ suggesting TORC1 may function upstream of ATFS-1 in maintaining heteroplasmy. However, the role of TORC1 in regulating the function of mitochondrial-localized ATFS-1 remains to be determined. Here, we report that inhibition of LONP1 through siRNA-mediated knockdown or the small-molecule inhibitor CDDO reduces Δ mtDNA abundance in cybrid cells harboring patient-derived mutant mtDNAs with either a large deletion or a single base pair substitution. Moreover, this decrease in Δ mtDNA levels was accompanied by improved mitochondrial respiration in cybrid cells treated with CDDO. Our data suggests that LONP1 inhibition may represent a therapeutic strategy for diseases caused by mutant mtDNAs.

REFERENCES

1. Baker, B.M. & Haynes, C.M. Mitochondrial protein quality control during biogenesis and aging. *Trends in biochemical sciences* **36**, 254-261 (2011).
2. Wai, T., Teoli, D. & Shoubridge, E.A. The mitochondrial DNA genetic bottleneck results from replication of a subpopulation of genomes. *Nature genetics* **40**, 1484-1488 (2008).
3. Gorman, G.S. *et al.* Prevalence of nuclear and mitochondrial DNA mutations related to adult mitochondrial disease. *Annals of Neurology* **77**, 753-759 (2015).
4. Srivastava, S. & Moraes, C.T. Manipulating mitochondrial DNA heteroplasmy by a mitochondrially targeted restriction endonuclease. *Human molecular genetics* **10**, 3093-3099 (2001).
5. Tanaka, M. *et al.* Gene therapy for mitochondrial disease by delivering restriction endonuclease SmaI into mitochondria. *Journal of biomedical science* **9**, 534-541 (2002).
6. Chiang, A.C.Y., McCartney, E., O'Farrell, P.H. & Ma, H. A Genome-wide Screen Reveals that Reducing Mitochondrial DNA Polymerase Can Promote Elimination of Deleterious Mitochondrial Mutations. *Current Biology* **29**, 4330-4336.e4333 (2019).
7. Stewart, J.B. & Chinnery, P.F. Extreme heterogeneity of human mitochondrial DNA from organelles to populations. *Nature Reviews Genetics* (2020).
8. Ahier, A. *et al.* Affinity purification of cell-specific mitochondria from whole animals resolves patterns of genetic mosaicism. *Nature cell biology* **20**, 352-360 (2018).
9. Lin, Y.F. *et al.* Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response. *Nature* **533**, 416-419 (2016).
10. Hahn, A. & Zuryn, S. The Cellular Mitochondrial Genome Landscape in Disease. *Trends Cell Biol* **29**, 227-240 (2019).
11. Pereira, C.V. & Moraes, C.T. Current strategies towards therapeutic manipulation of mtDNA heteroplasmy. *Front Biosci (Landmark Ed)* **22**, 991-1010 (2017).
12. Larsson, N.G. & Clayton, D.A. Molecular genetic aspects of human mitochondrial disorders. *Annu Rev Genet* **29**, 151-178 (1995).
13. Schon, E.A., DiMauro, S., Hirano, M. & Gilekerson, R.W. Therapeutic prospects for mitochondrial disease. *Trends Mol Med* **16**, 268-276 (2010).
14. Greaves, L.C. *et al.* Clonal expansion of early to mid-life mitochondrial DNA point mutations drives mitochondrial dysfunction during human ageing. *PLoS genetics* **10**, e1004620 (2014).
15. Bender, A. *et al.* High levels of mitochondrial DNA deletions in substantia nigra neurons in aging and Parkinson disease. *Nature genetics* **38**, 515-517 (2006).
16. Gitschlag, B.L. *et al.* Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in *C. elegans*. *Cell metabolism* **24**, 91-103 (2016).
17. Nargund, A.M., Pellegrino, M.W., Fiorese, C.J., Baker, B.M. & Haynes, C.M. Mitochondrial Import Efficiency of ATFS-1 Regulates Mitochondrial UPR Activation. *Science* **337**, 587-590 (2012).
18. Shpilka, T. *et al.* UPR(mt) scales mitochondrial network expansion with protein synthesis via mitochondrial import in *Caenorhabditis elegans*. *Nature communications* **12**, 479 (2021).
19. Nargund, A.M., Fiorese, C.J., Pellegrino, M.W., Deng, P. & Haynes, C.M. Mitochondrial and nuclear accumulation of the transcription factor ATFS-1 promotes OXPHOS recovery during the UPR(mt). *Molecular cell* **58**, 123-133 (2015).

20. Bernstein, S.H. *et al.* The mitochondrial ATP-dependent Lon protease: a novel target in lymphoma death mediated by the synthetic triterpenoid CDDO and its derivatives. *Blood* **119**, 3321-3329 (2012).
21. Tsang, W.Y. & Lemire, B.D. Stable heteroplasmy but differential inheritance of a large mitochondrial DNA deletion in nematodes. *Biochem Cell Biol* **80**, 645-654 (2002).
22. Lin, Y.-F. *et al.* Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response. *Nature* **533**, 416-419 (2016).
23. Gitschlag, B.L. *et al.* Homeostatic responses regulate selfish mitochondrial genome dynamics in *C. elegans*. *Cell metabolism* **24**, 91-103 (2016).
24. Cristina, D., Cary, M., Lunceford, A., Clarke, C. & Kenyon, C. A Regulated Response to Impaired Respiration Slows Behavioral Rates and Increases Lifespan in *Caenorhabditis elegans*. *PLoS genetics* **5**, e1000450 (2009).
25. Xu, C. *et al.* Genetic inhibition of an ATP synthase subunit extends lifespan in *C. elegans*. *Sci Rep* **8**, 14836 (2018).
26. Deng, P. *et al.* Mitochondrial UPR repression during *Pseudomonas aeruginosa* infection requires the bZIP protein ZIP-3. *Proceedings of the National Academy of Sciences* **116**, 6146-6151 (2019).
27. Nargund, A.M., Fiorese, C.J., Pellegrino, M.W., Deng, P. & Haynes, C.M. Mitochondrial and Nuclear Accumulation of the Transcription Factor ATFS-1 Promotes OXPHOS Recovery during the UPR. *Molecular cell* (2015).
28. Amrita, Christopher, Mark, Deng, P. & Cole Mitochondrial and Nuclear Accumulation of the Transcription Factor ATFS-1 Promotes OXPHOS Recovery during the UPRmt. *Molecular cell* **58**, 123-133 (2015).
29. Gustafsson, C.M., Falkenberg, M. & Larsson, N.-G. Maintenance and Expression of Mammalian Mitochondrial DNA. *Annual review of biochemistry* **85**, 133-160 (2016).
30. Rauthan, M., Ranji, P., Aguilera Pradenas, N., Pitot, C. & Pilon, M. The mitochondrial unfolded protein response activator ATFS-1 protects cells from inhibition of the mevalonate pathway. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 5981-5986 (2013).
31. Bota, D.A. & Davies, K.J.A. Lon protease preferentially degrades oxidized mitochondrial aconitase by an ATP-stimulated mechanism. *Nature cell biology* **4**, 674-680 (2002).
32. Liu, T. *et al.* DNA and RNA binding by the mitochondrial lon protease is regulated by nucleotide and protein substrate. *The Journal of biological chemistry* **279**, 13902-13910 (2004).
33. Chen, S.H., Suzuki, C.K. & Wu, S.H. Thermodynamic characterization of specific interactions between the human Lon protease and G-quartet DNA. *Nucleic acids research* **36**, 1273-1287 (2008).
34. Matsushima, Y., Goto, Y.I. & Kaguni, L.S. Mitochondrial Lon protease regulates mitochondrial DNA copy number and transcription by selective degradation of mitochondrial transcription factor A (TFAM). *Proceedings of the National Academy of Sciences* **107**, 18410-18415 (2010).
35. Göke, A. *et al.* Mrx6 regulates mitochondrial DNA copy number in *S. cerevisiae* by engaging the evolutionarily conserved Lon protease Pim1. *Molecular Biology of the Cell*, mbc.E19-08-0470 (2019).
36. Quirós, P.M., Langer, T. & López-Otín, C. New roles for mitochondrial proteases in health, ageing and disease. *Nature Reviews Molecular Cell Biology* **16**, 345-359 (2015).

37. Valenci, I., Yonai, L., Bar-Yaacov, D., Mishmar, D. & Ben-Zvi, A. Parkin modulates heteroplasmy of truncated mtDNA in *Caenorhabditis elegans*. *Mitochondrion* **20**, 64-70 (2015).
38. Chiang, A.C.-Y., McCartney, E., O'Farrell, P.H. & Ma, H. A genome-wide screen reveals that reducing mitochondrial DNA polymerase can promote elimination of deleterious mitochondrial mutations. *Current Biology* **29**, 4330-4336. e4333 (2019).
39. Kukat, C. *et al.* Cross-strand binding of TFAM to a single mtDNA molecule forms the mitochondrial nucleoid. *Proc Natl Acad Sci U S A* **112**, 11288-11293 (2015).
40. King, M.P. & Attardi, G. Human cells lacking mtDNA: repopulation with exogenous mitochondria by complementation. *Science* **246**, 500-503 (1989).
41. Bruno, C. *et al.* A stop-codon mutation in the human mtDNA cytochrome c oxidase I gene disrupts the functional structure of complex IV. *Am J Hum Genet* **65**, 611-620 (1999).
42. Moraes, C.T. *et al.* Mitochondrial DNA deletions in progressive external ophthalmoplegia and Kearns-Sayre syndrome. *N Engl J Med* **320**, 1293-1299 (1989).
43. Moraes, C.T., Schon, E.A., DiMauro, S. & Miranda, A.F. Heteroplasmy of mitochondrial genomes in clonal cultures from patients with Kearns-Sayre syndrome. *Biochem Biophys Res Commun* **160**, 765-771 (1989).
44. Yusoff, A.A.M., Abdullah, W.S.W., Khair, S. & Radzak, S.M.A. A comprehensive overview of mitochondrial DNA 4977-bp deletion in cancer studies. *Oncol Rev* **13**, 409 (2019).
45. Lee, H.C., Pang, C.Y., Hsu, H.S. & Wei, Y.H. Differential accumulations of 4,977 bp deletion in mitochondrial DNA of various tissues in human ageing. *Biochim Biophys Acta* **1226**, 37-43 (1994).
46. Munch, C. & Harper, J.W. Mitochondrial unfolded protein response controls matrix pre-RNA processing and translation. *Nature* **534**, 710-713 (2016).
47. Lin, Y.F. *et al.* Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response. *Nature* (2016).
48. Lu, B. *et al.* Phosphorylation of human TFAM in mitochondria impairs DNA binding and promotes degradation by the AAA+ Lon protease. *Molecular cell* **49**, 121-132 (2013).
49. Matsushima, Y., Goto, Y. & Kaguni, L.S. Mitochondrial Lon protease regulates mitochondrial DNA copy number and transcription by selective degradation of mitochondrial transcription factor A (TFAM). *Proceedings of the National Academy of Sciences of the United States of America* **107**, 18410-18415 (2010).
50. Gilkerson, R.W. *et al.* Mitochondrial autophagy in cells with mtDNA mutations results from synergistic loss of transmembrane potential and mTORC1 inhibition. *Human molecular genetics* **21**, 978-990 (2012).

ACKNOWLEDGMENTS

We thank the Caenorhabditis Genetics Center for providing *C. elegans* strains (funded by NIH Office of Research 362 Infrastructure Programs (P40 OD010440). We thank Carlos Moraes for the KSS and Giovanni Manfredi for the CoxI G6930A cybrid cell lines.

Funding: This work was supported by HHMI, the Mallinckrodt Foundation and National Institutes of Health grants (R01AG040061 and R01AG047182) to C.M.H., (R01GM115911 and R01AI117839) to S.A.W., (R01GM111706 and R35GM130320) to P.C. and (F31HL147482) to K.L. **Data and materials availability:** All data in the paper are present in the paper or the Supplementary Materials. The ChIP-sequencing data have been deposited to the Gene Expression Omnibus database under the BioProject accession code PRJNA590136. The next-generation sequencing data have been deposited in the NCBI Sequence Read Archive database under the BioProject accession code PRJNA517630.

AUTHOR CONTRIBUTIONS

Q.Y. and C.M.H. planned the experiments. Q.Y., Y.D., T.S., N.N., R.Z and P.C. generated the worm strains. Q.Y. performed the *C. elegans* and cybrid mtDNA analysis including ChIP and respiratory function. P.L., K.L. and S.W. performed and analyzed mtDNA sequencing. Q.Y., N.S.A. and C.M.H. wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing financial interests.

Supplemental Materials:

Materials and Methods

Figures S1-S7

Tables S1-S2 (.xlsx)

References

Fig. 1

Fig. 1 | OXPHOS dysfunction increases mtDNAs through ATFS-1.

a, ATFS-1/UPR^{mt} signaling schematic in healthy cells. **b**, Schematic comparing *C.elegans* wildtype homoplasmic and Δ mtDNA (*uaDf5* deletion) strains. **c**, Photomicrographs of *hsp-6_{pr}::gfp* and *hsp-6_{pr}::gfp;uaDf5* worms (Scale bar 0.1 mm). **d**, Quantification of total mtDNA in homoplasmic wildtype, *uaDf5* worms and *uaDf5* worms raised on *atfs-1*(RNAi) (*n* = 3). **e**, Photomicrographs of *hsp-6_{pr}::gfp* worms raised on control (ctrl), *cco-1* or *cyc-1*(RNAi). **f**, photomicrographs of wildtype, *isp-1(qm150)* and *clk-1(qm30);hsp-6_{pr}::gfp* worms. (Scale bar 0.1 mm). **g**, Quantification of total mtDNA in homoplasmic wildtype and *atfs-1(null)* worms raised on control(RNAi) or nuclear-encoded OXPHOS genes (*n* = 3). **h**, Quantification of mtDNA in wildtype, *isp-1(qm150)*, *clk-1(qm30)* and *nduf-2.1(fc21)* mutant worms raised on control(RNAi) or *atfs-1*(RNAi) (*n* = 3-4). (“*n*” means samples from independent biological replicates and each sample contains 40-60 animals; every dot stands for averaged value from 3 technical replicates); One-way ANOVA was used in **d,g** and **h**; error bars mean $\pm$ S.E.M. **p*<0.05, ***p*<0.01, ****p*<0.001.

Fig. 2 | Defective OXPHOS impedes the degradation ATFS-1 and facilitates mitochondrial localized ATFS-1 and POLG binding to mtDNAs.

a, ATFS-1 ChIP-seq profile of the entire mitochondrial genome (mtDNA) of wildtype worms raised on *spg-7*(RNAi) (adapted from¹⁹). **b**, ATFS-1 immunoblots of wildtype worms raised on control or *cco-1*(RNAi) after fractionation into total lysate (T), post-mitochondrial supernatant (S), and mitochondrial pellet (M). Tubulin (Tub) and the OXPHOS component (NDUFS3) are used as loading controls. Arrow is mitochondrial-localized ATFS-1. **c**, Quantification of mtDNA following ATFS-1 ChIP-mtDNA in homoplasmic wildtype worms and homoplasmic *atfs-1*(null) worms raised on control or *cco-1*(RNAi) ($n = 4$). **d**, POLG immunoblots of wildtype worms raised on control or *cco-1*(RNAi) after mitochondrial fractionation. **e**, Quantification of total mtDNA following POLG ChIP-mtDNA in wildtype or *atfs-1*(null) homoplasmic worms ($n = 3$). **f**, Quantification of mtDNA following ATFS-1 ChIP-mtDNA in homoplasmic *atfs-1^{nuc(-)}* worms and homoplasmic *atfs-1*(null) worms raised on control or *cco-1*(RNAi) ($n = 3$). **g**, Quantification of total mtDNA following POLG ChIP-mtDNA in homoplasmic *atfs-1^{nuc(-)}* and *atfs-1*(null) worms raised on control or *cco-1*(RNAi) ($n = 3$). **h**, Quantification of total mtDNA in *atfs-1^{nuc(-)}* homoplasmic worms raised on control(RNAi) or *cco-1*(RNAi)) ($n = 3$). **i**, Quantification of total mtDNA in wildtype, *clk-1(qm30)* and *clk-1(qm30);atfs-1^{nuc(-)}* homoplasmic wildtype worms ($n = 4$, One-way ANOVA). (“ n ” means independent biological replicates and each sample pooled from large population in **c,e-g**; contains 40-60 animals in **h** and **i**; every dot stands for averaged value from 3 technical replicates); Two-tailed Student’s t test was used except to **i**; error bars mean $\pm$ S.E.M. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Fig. 3 | LONP-1 limits ATFS-1 binding to wildtype mtDNAs and impairs replication.

a, FLAG immunoblots of LONP-1^{FLAG} worms following fractionation into total lysate (T), post-mitochondrial supernatant (S), and mitochondrial pellet (M). Tubulin (Tub) and the OXPHOS component NDUFS3 are loading controls. **b**, Quantification of mtDNA from homoplasmic LONP-1^{FLAG} worms following ChIP-mtDNA using FLAG or control (Mock) antibody ($n = 3$). **c**, LONP-1 consensus binding motif within mtDNA. **d**, ChIP-seq profile of mtDNA from homoplasmic LONP-1^{FLAG} worms raised on control(RNAi) using FLAG antibody (red). ATFS-1 ChIP-seq profile from homoplasmic worms raised on *spg-7*(RNAi) (blue) (as in Fig.2a). **e**, Quantification of total mtDNA following ATFS-1 ChIP-mtDNA in wildtype or *atfs-1(null)* homoplasmic worms raised on control(RNAi) or *lonp-1*(RNAi) ($n = 3$). **f**, Quantification of mtDNA in homoplasmic wildtype and *atfs-1(null)* worms raised on control(RNAi) or *lonp-1*(RNAi) ($n = 4$). **g**, Quantification of total mtDNA following POLG ChIP-mtDNA in wildtype homoplasmic worms raised on control(RNAi) or *lonp-1*(RNAi) ($n = 3$). **h**, Schematic of the relationship between LONP-1 activity, mitochondrial ATFS-1 accumulation and mtDNA replication. (“ n ” means independent biological replicates and each sample pooled from large populations in **b,e,g**; each sample contains 40-60 animals in **f**; every dot stands for averaged value from 3 technical replicates); Two-tailed Student’s t test was used; error bars mean $\pm$ S.E.M. * $p < 0.05$, ** $p < 0.01$.

Fig. 4

Fig. 4 | ATFS-1 and POLG primarily interact with Δ mtDNAs in heteroplasmic worms.

a, Δ mtDNA quantification as determined by qPCR in *uaDf5* worms, *atfs-1(null);uaDf5*, *pdr-1(tm598);uaDf5* and *atfs-1(null);pdr-1(tm598);uaDf5* worms ($n = 3-4$, One-way ANOVA). **b**, Images of TMRE-stained micrographs of wildtype worms raised on control(RNAi) or *cco-1*(RNAi). Scale bar, 10 μ M. **c**, Images of TMRE-stained micrographs of heteroplasmic (Δ mtDNA) worms raised on control(RNAi), or wildtype worms raised on control or *spg-7*(RNAi). Scale bar, 10 μ M. **d**, Immunoblots of wildtype worms raised on control or *spg-7*(RNAi) and heteroplasmic (Δ mtDNA) worms raised on control(RNAi) after fractionation into total lysate (T), post-mitochondrial supernatant (S), and mitochondrial pellet (M). Tubulin (Tub) and the OXPHOS component (NDUFS3) are used as loading controls. Arrow is mitochondrial-localized ATFS-1. **e**, Quantification of total mtDNA following ATFS-1 ChIP-mtDNA in homoplasmic wildtype or Δ mtDNA worms ($n = 3$ or 5, two-tailed Student's t test). **f**, Workflow of ATFS-1 or POLG ChIP-mtDNA and quantification of wildtype mtDNA and Δ mtDNA in heteroplasmic worms. **g**, Quantification of wildtype mtDNA and Δ mtDNA by qPCR following ATFS-1 ChIP-mtDNA in heteroplasmic worms. Post-lysis/Input Δ mtDNA ratio was 60% ($n = 4$). **h**, Quantification of wildtype mtDNA and Δ mtDNA by qPCR following POLG ChIP-mtDNA in heteroplasmic worms. Post-lysis/Input Δ mtDNA ratio was 54% ($n = 3$). **i**, Quantification of wildtype mtDNA and Δ mtDNA following TFAM IP-mtDNA in heteroplasmic worms. Post-lysis/Input Δ mtDNA ratio was 53% ($n = 4$). ("n" means independent biological repeats and each sample contains 40-60 animals in **a**; each sample pooled from large populations in **e,g-i**; every dot stands for averaged value from 3 technical replicates in **a,e** or strands for every technical replicates in **g-i**); error bars mean $\pm$ S.E.M. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$.

Fig.5

Fig. 5 | LONP-1 is required to maintain heteroplasmy.

a, Δ mtDNA and wildtype mtDNA quantification by qPCR following LONP-1 ChIP-mtDNA in heteroplasmic worms. Post-lysis/Input Δ mtDNA ratio was 59% ($n = 3$). **b**, Quantification of wild type mtDNA by qPCR in heteroplasmic worms raised on control(RNAi) or *lonp-1*(RNAi) ($n = 3$). **c**, Quantification of Δ mtDNA by qPCR in heteroplasmic *uaDf5* worms raised on control(RNAi) or *lonp-1*(RNAi) ($n = 3$). **d**, Δ mtDNA and wild-type mtDNA quantification by qPCR following ATFS-1 ChIP-mtDNA in heteroplasmic worms raised on control(RNAi) or *lonp-1*(RNAi) ($n = 4$). **e**, Δ mtDNA and wild-type mtDNA quantification by qPCR following POLG ChIP-mtDNA in heteroplasmic worms raised on *lonp-1*(RNAi). Post-lysis/Input Δ mtDNA ratio was 26% ($n = 4$). (“ n ” means independent biological repeats and each sample within each biological replicate corresponds to a sample pooled from large populations in **a,d,e**; each biological replicate corresponds to a sample pooled from from 40-60 animals in **b,c**; every dot stands for averaged value from 3 technical replicates in **b,c** or strands for every technical replicates in **a,d** and **e**); Two-tailed Student’s t test was used in **b,c**; error bars mean $\pm$ S.E.M. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Fig. 6

Fig. 6 | LONP1 inhibition improves heteroplasmy and OXPHOS function in cybrid cells.

a, Schematic comparing human wildtype, KSS deletion (Δ mtDNA) and CoxI G6930A mtDNAs. **b**, LONP1 immunoblots from KSS heteroplasmic cells treated with *hLONP1* or control (NC) siRNA. Tubulin (Tub) serves as a loading control. **c**, WT mtDNA quantification in KSS cells treated with control or *hLONP1* siRNA ($n = 3-4$). **d**, Quantification of KSS Δ mtDNA in cells treated with control or *hLONP1* siRNA ($n = 3$). **e**, Chemical structure of CDDO. **f**, Quantification of G6930A mtDNA percentage following treatment with DMSO, 0.1 μ M CDDO, or 0.25 μ M CDDO at the indicated time points up to 130 days. * $p < 0.05$ ($n = 3$). **g**, Δ mtDNA quantification in KSS heteroplasmic cells treated with DMSO, 0.1 μ M CDDO, or 0.25 μ M for 4 or 13 weeks. **h**, Cell viability of 143b(WT) and CoxI G6930A cells exposed to various concentrations of CDDO for 72 hours ($n = 3$ in 143b(WT) cell, $n = 4$ in CoxI G6930A cell). (“ n ” means independent biological replicates; every dot stands for averaged value from 3 technical replicates in **c,d,f** or stand for averaged value from biological replicates in **h**); Two-tailed Student’s t test was used in **c,d** and **f**; error bars mean $\pm$ S.E.M. * $p < 0.05$, ** $p < 0.01$.

Fig. 7

Fig. 7 | Pharmacological inhibition of LONP1 improves heteroplasmy and OXPHOS function in cybrid cells.

a, Percentage of CoxI G6930A mtDNA in cells treated with DMSO, 0.1 μ M or 0.25 μ M CDDO for 3, 5, 10 or 18.5 weeks, $n = 3$; error bars mean $\pm$ S.E.M.; * $p < 0.05$ (Student's t-test). **b**, Oxygen consumption rates (OCR) of CoxI G6930A cells treated with DMSO, 0.1 μ M or 0.25 μ M CDDO for 3, 5, 10 or 18.5 weeks ($n \geq 10$). **c**, Quantification of basal respiration ($n \geq 10$). **d**, Quantification of respiratory capacity. (" n " means independent biological repeats; every dot stands for averaged value from 4 technical replicates in **c,d**); error bars mean $\pm$ S.E.M. * $p < 0.05$.