**SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS
ALTER THE PATTERNING, MORPHOLOGY AND ABUNDANCE OF ROOT HAIRS
IN ROOT TIPS OF *A. THALIANA* SEEDLINGS**

Iva McCarthy-Suárez^{1,*}

¹*Hormone Signalling and Plant Plasticity Group. Instituto de Biología Celular y Molecular de Plantas (CSIC-Universidad Politécnica de Valencia). C/ Ingeniero Fausto Elio s/n (46022), Valencia, Spain.*

¹ In leave of absence. *Corresponding author. Email: ivmcsua@upvnet.upv.es.

ABSTRACT

In spite of the known role of gibberellins (GAs), and of their antagonistic proteins, the DELLAs, in leaf hair production, no investigations, however, have assessed their hypothetical function in the production of root hairs. To this aim, the effects of supra-physiological levels of GAs/DELLAs on the spatial patterning of gene expression of the root hair (CPC) and root non-hair (GL2, EGL3 and WER) epidermal cell fate markers, as well as on the distribution, morphology and abundance of root hairs, were studied in root tips of 5-day-old *A. thaliana* seedlings. Results showed that excessive levels of GAs/DELLAs impaired the spatial patterning of gene expression of the root hair/non-hair epidermal cell fate markers, as well as the arrangement, shape and frequency of root hairs, giving rise to ectopic hairs and ectopic non-hairs, two-haired cells, two-tipped hairs, branched hairs, longer and denser hairs near the root tip under excessive DELLAs, and shorter and scarcer hairs near the root tip under excessive GAs. However, when the *gai-1* (GA-insensitive-1) DELLA mutant protein was specifically over-expressed at the root epidermis, no changes in the patterning or abundance of root hairs occurred. Thus, these results suggest that, in seedlings of *A. thaliana*, the GAs/DELLAs might have a role in regulating the patterning, morphology and abundance of root hairs by acting from the sub-epidermal tissues of the root.

KEYWORDS

Gibberellins, DELLAs, root hair patterning, root hair morphology, root hair number.

1. INTRODUCTION

The epidermal cell organization in roots of *A. thaliana* seedlings, consisting of single rows of hair-bearing cells (trichoblasts, which lay over the cleft between two cortical cells) alternating with double rows of hairless cells (atrachoblasts, which lay over just one cortical cell) has been shown to be genetically determined by a complex network of transcription factors and positional signals, such as CAPRICE (CPC), GLABRA2 (GL2), WEREWOLF (WER) and ENHANCER OF GLABRA3 (EGL3), and regulated by auxin, ethylene (ET), abscisic acid (ABA), nitric oxide (NO), brassinosteroids (BRs), cytokinins (CKs) and strigolactones (SLs) (Silverman *et al.*, 1998; Cao *et al.*, 1999; Van Hengel *et al.*, 2004; Lombardo *et al.*, 2006; Kappusamy *et al.*, 2009; Schiefelbein *et al.*, 2009; Niu *et al.*, 2011; Salazar-Henao *et al.*, 2016). These hormones, in turn, seem to act downstream of the *GL2* gene network, permitting root cells to have fate plasticity, i.e., ability to change to the alternative differentiation route at a relatively late state, as it is not cell lineage, but position, and sometimes even a position-independent mechanism, what seems to continuously determine cell fate (Grierson and Schiefelbein, 2002; Schiefelbein *et al.*, 2009; Yu *et al.*, 2017). Moreover, these hormones mediate the changes in the root hair patterning associated to the plant responses to soil stress without altering the expression of the epidermal cell fate markers WER and GL2 (Schmidt *et al.*, 2000; Yang *et al.*, 2007; Martín-Rejano *et al.*, 2011).

Given that the GAs/DELLAs have a role in trichome (leaf hair) production in *A. thaliana* (Chien and Sussex, 1996; Traw and Bergelson, 2003) and participate in microtubule (MT) cytoskeleton organization (Locascio *et al.*, 2013), which is essential for the growth of trichomes and root hairs and for establishing the identity and shape of root cells (Bao *et al.*, 2001), and because there are no reports concerning the hypothetical implication of GAs/DELLAs in the root hair patterning, this study aimed to investigate the effect of excessive levels of these hormones on the distribution and abundance of root hairs in seedlings of *A. thaliana*. In addition, because changes in the levels of auxin, ET, ABA, NO, BRs and SLs have been correlated to alterations in root hair morphology in response to nutritional stresses such as low availability of P, B or Fe in the soil (longer and branched root hairs) (Schmidt *et al.*, 2000; Yang *et al.*, 2007; Martín-Rejano *et al.*, 2011), this work also aimed to determine whether the GAs / DELLAs might have a role in regulating the morphology of root hairs in seedlings of *A. thaliana*. To this aim, the spatial expression of the GUS or GFP-fused transcripts of the root hair (CPC) and non-hair (GL2, EGL3, WER) epidermal cell fate markers, as well as the arrangement, shape and density of root hairs, were studied in *A. thaliana* seedlings grown for 5 days under (or harbouring) excessive levels of GAs/DELLAs. Finally, to locate the tissue from which these hormones might hypothetically affect the patterning of root hairs, the root hair distribution was studied in 5-day-old mutant seedlings resulting from expressing the *gai-1* DELLA dominant allele in different tissues of the root (UAS (GAL4-UPSTREAM ACTIVATION SEQUENCE) expression directed

system lines; Dr. Jim Haselhoff's laboratory). Results of this study suggested that the GAs/DELLAs might be involved in regulating the patterning, morphology and abundance of root hairs in *A. thaliana* seedlings.

2. MATERIALS and METHODS

2.1. Plant Material and Growth Conditions

Arabidopsis thaliana Col (0) seeds were sterilized (70 % Ethanol (v/v) and 0.01 % Triton X-100 (v/v)), sown on half-strength MS medium plates (0.8 % (w/v) agar and 1 % (w/v) sucrose), stratified for 3-4 days (4°C, darkness), germinated, and grown vertically (22 °C; 5-7 days) under continuous white light (Percival growth chamber E-30B) (<http://www.percival-scientific.com>) as described by Lee and Schiefelbein (1999).

2.2. Hormone and Chemical treatments

Stock solutions of paclobutrazol (PAC, 10 mM in acetone 100 % (v/v)), GA₄ (1 mM in 100 % ethanol (v/v)) or GA₃ (50 mM in 100% ethanol (v/v)) were conveniently diluted and added to MS agar medium or water (in the case of liquid incubation experiments) to obtain a final concentration of 0.5 μM PAC, 1 μM GA₄, and 30 μM GA₃.

2.3. Mutant Lines

The spatial patterning of gene expression of the hair (CPC) and non-hair (GL2, EGL3, WER) epidermal cell fate markers in roots of *A. thaliana* seedlings was studied by using their GUS or GFP-fused promoter lines (*CPCpro::GUS*, *GL2pro::GUS*, *EGL3pro::GUS* and *WERpro::GFP*) as well as those derived from crossing lines harbouring constitutively excessive levels of GAs / DELLAs with the *GL2pro::GUS* line (*GID1b ox x GL2pro::GUS*, *gai-1 x GL2pro::GUS*, *HSp::gai-1 x GL2pro::GUS*, *pGAI::gai-1:GR x GL2pro::GUS* and *SCR::gai-1:GR x GL2pro::GUS* (*Ler x GL2pro::GUS* background)). The effect of transient increases in the levels of the *gai-1* dominant DELLA on the root hair distribution in *A. thaliana* seedlings was examined by using the heat-shock inducible *HSp::gai-1* (which over-expresses the DELLA *gai-1* upon heat shock) and dexametasone (DEXA)-inducible *pGAI::gai-1:GR* and *SCR::gai-1:GR* (with glucocorticoid-binding domain) mutant lines. The *HSp::gai-1* mutant seedlings were grown at 37°C for 4 h (heat shock) and then at 22°C for 2 h (recovery period), whereas the *pGAI::gai-1:GR* and *SCR::gai-1:GR* mutant seedlings were incubated in 0.1, 0.2, 0.5, 1.2 or 10 μM DEXA for a minimum of 6h. The root hair distribution was also studied in mutants with excessive levels of GAs/DELLAs (*gai-1*, *GAI-ox* (*GAI*-over-expressing), *QD* (*quadruple DELLA mutant*), *5X* (*quintuple DELLA mutant*), *GID1b-ox* (which over-expresses the GA receptor *GID1b* (*GIBBERELLIN INSENSITIVE DWARF1*), in mutants over-expressing *gai-1* in different tissues of the root [*ML1::gai-1* (epidermis) and *UAS* expression directed system (*GAL4-UPSTREAM ACTIVATION SEQUENCE*) mutants: *UAS::gai-1 x C24* (control, background); *UAS::gai-1 x J0951* (epidermis of the meristematic zone (MZ)); *UAS::gai-1 x J2812* (MZ epidermis and cortex); *UAS::gai-1 x N9142* (cortex of the elongation zone (EZ)); *UAS::gai-1 x M0018* (MZ cortex and endodermis); *UAS::gai-1 x J0571* (MZ cortex and endodermis); *UAS::gai-1 x Q2393* (all tissues but the endodermis); *UAS::gai-1 x Q2500* (MZ endodermis/pericycle); *UAS::gai-1 x J0121* (EZ pericycle); *UAS::gai-1 x J0631* (all tissues of the EZ); *UAS::gai-1 x J3281* (vessels)], and in the *wer*, *cpc* and *35S::CPC* (cauliflower mosaic virus 35S promoter) mutants.

2.4. GUS activity assay

GUS (β-glucuronidase) staining of the *GL2pro::GUS*, *CPCpro::GUS* and *EGL3pro::GUS* reporter lines was performed as described by Frigerio *et al.*, (2006), but using 8mM instead of 2 mM potassium ferro/ferricyanide and incubating the seedlings (15 min to 2h) in the reaction mixture at 4 °C instead of 37 °C.

2.5. Microscopy

The patterning of the hair/non-hair epidermal cell types in roots of *A. thaliana* seedlings was studied by staining the roots with 0.67 mg/ml propidium iodide, by observing the root tips under a Nikon Eclipse E6000 microscope, and by calculating the percentage of hairs/non-hairs at the trichoblast/atrichoblast positions (Dr. Benedicte Desvoyes' method). The *GL2pro::GUS* expression patterning in cross sections of root tips was studied on ultra-thin cross sections of plastic resin-embedded roots as previously described at Dr. Schiefelbein Lab Protocols (<http://www.mcdm.lsa.umich.edu/labs/schiefel/protocols.html>). Seedlings were included in 1% agarose in 0.1M sodium phosphate buffer, pH 6.8, and stained for GUS activity. Root-containing blocks were then cut, fixed with 4% para-formaldehyde in PBS, dehydrated in ethanol series (15%, 30%, 50%, 75%, 95% and 100%, 1h each), kept in 100% ethanol overnight, incubated in Technovit® 7100 infiltration solution for 2 days, inserted in gelatine capsules, and embedded for 9 days in Technovit® 7100 plastic resin (Heraeus Kultzer, Germany). Ultramicrotome (Ultracut E, Reichert Jung, Germany) cross sections of resin-embedded roots were then stained with 0.06% (w/v) toluidine blue and observed under a Nikon Eclipse E600 microscope. The *WERpro::GFP* expression was visualized by using a Leica Confocal Microscope (excitation: 488nm; detection: 500-530nm band-path filter for GFP).

3. RESULTS

3.1. Excessive levels of GAs/DELLAs altered the root hair patterning in seedlings of *A. thaliana*:

To assess whether the GAs/DELLAs might have a role in the root hair patterning of *A. thaliana* seedlings, the spatial gene expression of the root hair (CPC) and non-hair (GL2, EGL3, WER) epidermal cell fate markers was studied in seedlings of the *GL2pro::GUS*, *CPCpro::GUS*, *EGL3pro::GUS* and *WERpro::GFP* transgenic lines grown for 5 days under supra-physiological levels of GAs/DELLAs (Fig. 1A). Results showed that growth under excessive levels of GAs/DELLAs altered the normal patterning of gene expression the root hair/non-hair epidermal cell fate markers (Fig. 1A). This was confirmed by analysing the spatial expression of *GL2* in the *GID1b-ox* (which over-expresses the GA receptor *GID1b*), *gai-1*, *HSp::gai-1* (which over-expresses the DELLA *gai-1* upon exposure to heat (37°C, 4h)), and DEXA-inducible *pGAI::gai-1:GR* and *SCR::gai-1:GR* mutants (Fig. 1A). Moreover, the alteration of the *GL2pro::GUS* expression pattern under excessive levels of GAs/DELLAs was corroborated in ultra-thin sections of resin-embedded roots (Fig. 1B).

An analysis of the distribution of root hairs and non-hairs relative to their position over the cortex cells showed that excessive levels of GAs/DELLAs impaired the correct positioning of the root hairs/non-hairs, giving rise to ectopic root hairs (at the atrichoblast position) and ectopic root non-hairs (at the trichoblast position) (Figs. 2A and 2B; Tables 1 and 2). Interestingly, treatment with GA₄ (1 µM) reduced the percentage of ectopic root hairs in the hairy *35S::CPC* mutant, whereas treatment with PAC (0.5 µM) slightly decreased the percentage of ectopic root non-hairs in the bald *cpc* mutant (Table 2). In accordance with these changes, growing *A. thaliana* seedlings under supra-physiological levels of GAs/DELLAs for 5 days altered the arrangement of root hairs in root tips, giving rise to ectopic hairs (in a non-hair row), ectopic non-hairs (in a hair row) and adjacent hair rows (Figs. 3A and 3B). This was confirmed in the *gai-1*, *QD*, *5X*, *GID1b-ox*, *pGAI::gai-1:GR* and *SCR::gai-1:GR* mutants (Figs. 3A and 3B).

To ascertain from which particular tissue of the root the GAs/DELLAs might be affecting the root hair patterning, the distribution of root hairs and the positioning of the root hair/non-hair cells over the root cortex cells were studied in *A. thaliana* transgenic seedlings expressing the *gai-1* DELLA allele in different tissues of the root (Figs. 2B, 3A and 3B; Table 2). Results showed that the root hair distribution changed when *gai-1* was over-expressed at the cortex, endodermis or pericycle of the meristematic (MZ) or elongation (EZ) zones of the root (J2812 >> *gai-1*, M0018 >> *gai-1*, Q2500 >> *gai-1*, J0121 >> *gai-1*, Q2393 >> *gai-1* and J0631 >> *gai-1* lines), but not when *gai-1* was over-expressed at the root epidermis (J0951 >> *gai-1* and *ML1::gai-1* lines) (Fig. 3A). Moreover, ectopic hairs, ectopic non-hairs and adjacent hair rows appeared when *gai-1* was over-expressed at the cortex (J2812 >> *gai-1* and N9142 >> *gai-1* lines), endodermis (M0018 >> *gai-1* and J0571 >> *gai-1* lines) or pericycle (J0121 >> *gai-1* line) of the root, or in all root tissues but the endodermis (Q2393 >> *gai-1* line) (Figs. 2B, 3A and 3B). However, when *gai-1* was over-expressed at the root vessels (J3281 >> *gai-1* line), the production of root hairs and the growth of the root stopped (Fig. 3A).

3.2. Excessive levels of GAs/DELLAs altered the morphology, length and abundance of hairs in root tips of *A. thaliana* seedlings:

Excessive levels of GAs/DELLAs also modified the morphology of trichoblasts and root hairs in root tips of *A. thaliana* seedlings, frequently giving rise to two-haired cells, two-tipped hairs and branched hairs (Fig 4). In addition, excessive levels of GAs/DELLAs altered the length and density of root hairs. Whereas high levels of DELLAs increased the length and number of hairs near the root tip, high levels of GAs had the opposite effect (Figs. 5A and 5B; Table 3). Moreover, hair abundance in root tips of *A. thaliana* seedlings increased when *gai-1* was over-expressed at the cortex (J2812 >> *gai-1*), endodermis (M0018 >> *gai-1*) or pericycle (J0121 >> *gai-1*) of the root, but not when *gai-1* was over-expressed at the epidermis of the root MZ (J0951 >> *gai-1*) or the cortex of the root EZ (N9142 >> *gai-1*) (Table 3). Also, treatment of the bald *cpc* mutant with PAC slightly increased the hair frequency (and length) near the root tip, whereas treatment of the hairy *wer* and *35S::CPC* mutants with GA₄ reduced it (Fig. 5B; Table 3).

High levels of GAs/DELLAs also altered the abundance of hairs in the radial dimension of the root tips (Tables 4 and 5). The number of root hairs per root cross section, calculated as the summary of root hairs at the trichoblast and atrichoblast positions (or the summary of root hairs and ectopic root hairs per root cross section) increased under excessive DELLAs (PAC, *gai-1*) but decreased in the *5X* mutant (Table 5). On the other hand, the number of root non-hairs per root cross section, calculated as the summary of root non-hairs at the atrichoblast and trichoblast positions (or the summary of root non-hairs and ectopic root non-hairs per root cross section), decreased under excessive DELLAs, but experienced an enhancement in the *5X* mutant (Table 5). Thus, the estimated abundance of root hairs in the radial dimension of root tips seemed to increase under excessive DELLAs, but to decrease under excessive GAs.

4. DISCUSSION

4.1. The GAs/DELLAs might regulate the root hair patterning in seedlings of *A. thaliana*:

Whereas the role of GAs/DELLAs in the production and distribution of leaf hairs has been well studied (Telfer *et al.*, 1997; Perazza *et al.*, 1998), their hypothetical function in the determination and arrangement of root hairs has not been examined up to date. To this aim, the effects of high levels of GAs/DELLAs on the spatial gene expression of the hair (CPC) and non-hair (GL2, WER and EGL3) markers of root epidermal cell fate, as well as on the distribution of root hairs, were analysed in seedlings of *A. thaliana*. Results showed that excessive levels of GAs/DELLAs impaired the spatial gene expression of the root hair/non-hair epidermal cell fate markers and disarranged the normal distribution of root hairs, what suggested that the GAs/DELLAs might be involved in regulating the root hair patterning in seedlings of *A. thaliana*. In fact, stable or inducible mutants with low (*gai-1*, *HSp::gai-1*, *pGAI::gai-1:GR*, *SCR::gai-1:GR*) or high (*QD*, *5X*, *GID1b-ox*) levels of GAs showed not only a random expression of *GL2* at the MZ and EZ of the root, known as the cell fate-decision zones (Pernas *et al.*, 2010), but also a disarrangement of the root hairs. Because neither the *GL2* spatial expression nor the distribution of root hairs suffered changes when the *gai-1* DELLA was over-expressed at the root epidermis (*ML1::gai-1* x *GL2pro::GUS*, *ML1::gai-1* and *UAS::gai-1* x J0951 transgenic lines), it was concluded that the GAs/DELLAs do not seem to affect the root hair patterning in *A. thaliana* seedlings by acting on this cell layer of the root, but on tissues placed underneath. In fact, expression of *gai-1* at the cortex, endodermis or pericycle of the root MZ altered the root hair patterning.

Interestingly, expressing *CPC* at the stele rescues the phenotype of the hairless *cpc* mutant, what suggests that epidermal cell differentiation might be controlled from the internal tissues of the root (Rishmawi *et al.*, 2014). Therefore, the results of this study suggest that, as it was previously reported for auxins, ET, ABA, NO, BRs and SLs (Schiefelbein, 2003), the GAs/DELLAs might regulate the root hair patterning in seedlings of *A. thaliana* independently from the gene network for the root epidermal cell fate specification, although confirmatory studies might be required.

The reason why excessive levels of GAs/DELLAs disarranged the root hair patterning in *A. thaliana* seedlings might have been, in part, related to their effects on the cytoskeleton of MT. The MT cytoskeleton, consisting in polymers of α and β tubulin, is essential for the appropriate distribution of positional signals during development (Schiefelbein, 2003). Also, the orientation of MT participates in the determination of epidermal cell fate (Pietra, 2014). Thus, MT lay randomly in trichoblasts but transversally in atrichoblasts (Dugardeyn and Van der Straeten, 2008). Hormone-induced reorganization of MT is also necessary for root

hair initiation (Bao *et al.*, 2001; Schiefelbein, 2003). Interestingly, the GAs/DELLAs regulate MT organization by interacting with prefoldin, a protein required for the folding of tubulin (Locascio *et al.*, 2013). As a result of this interaction, MT are organized in the presence of GAs, like in root or mesocotyl epidermal cells, and disorganized in the presence of DELLAs (Perazza *et al.*, 1998; Bouquin *et al.*, 2002; Locascio *et al.*, 2013). On the other hand, mutants impaired in MT assembly have an altered root hair patterning (Bouquin *et al.*, 2002). The *lue1* mutant, which lacks a MT-severing and cell wall (CW) biosynthesis-related katanin protein, and whose MT are disorganized, is allelic to ectopic root hair 1 (*erh1*) and has an altered root hair patterning (Bouquin *et al.*, 2002; Webb *et al.*, 2002). In addition, *lue1* presents an inappropriate regulation of the GA biosynthesis-related AtGA20ox activity and responds to GAs (Schneider *et al.*, 1997; Bouquin *et al.*, 2002).

Ectopic root hairs have also been described in TUA6/AS transgenic lines under-expressing α -tubulin genes, in plants treated with MT polymerization-inhibiting drugs or with trichostatin A (TSA, a histone deacetylase (HDA) inhibitor), during the inducible expression of MT-interacting phospholipase-D (PLD) activity, as well as in mutants of MT severing/reorganization-related proteins, such as HDA, COBRA, SABRE and katanin p60 (Schiefelbein *et al.*, 1997; Bao *et al.*, 2001; Bouquin *et al.*, 2002; Sedbrook, 2004; Wang, 2005; Xu *et al.*, 2005; Li *et al.*, 2006, 2015; Chen *et al.*, 2015; Pietra *et al.*, 2015). In fact, the katanin complex is required for the specification of root epidermal cells (Webb *et al.*, 2002). In addition, the katanin P60-related alteration of MT organisation affects the composition and deposition of the CW (Sedbrook, 2004). Histone deacetylation also participates in cellular patterning, because TSA-induced histone acetylation modifies GL2, WER and CPC expression and localization and induces ectopic root hairs (Xu *et al.*, 2005; Cui and Benfey, 2009). Lack of SABRE function equally destabilizes the expression of cell fate markers, including WER and GL2 (Pietra *et al.*, 2015). In addition, a delocalized expression of GL2 has been documented for the *jdk* (jackdaw) and *scm* (scrambled) mutants (Hassan *et al.*, 2010; Pietra, 2014).

Therefore, the MT participate in cell identity specification (Webb *et al.*, 2002). Cell identity, in turn, mediates the root responses to abiotic stress (Dinneny *et al.*, 2008). Thus, ectopic root hairs and non-hairs have been described in *A. thaliana* seedlings exposed to gamma irradiation, Cd or As, and during P deficiency, although without quantitative changes in the *WER* and *GL2* expression (Ma *et al.*, 2001; Nagata *et al.*, 2004; Yang *et al.*, 2007; Bahmani *et al.*, 2016). Moreover, stress down-regulates the actin and tubulin gene expression (Sánchez-Calderon *et al.*, 2013). In turn, a reduced expression of the α -tubulin gene results in MT disassembly, with MT laying in an aberrant way, and in their reorganization (Bao *et al.*, 2001).

Consequently, the root hair patterning responds to environmental signals (Salazar-Henao *et al.*, 2016). For instance, the photoperiod and thermoperiod control the root hair patterning in tomato (Tsai *et al.*, 2004). Interestingly, the GAs participate in thermotolerance (Alonso-Ramirez *et al.*, 2009). Thus, the results of this study suggest that the GAs/DELLAs might regulate, in part, the root hair patterning in *A. thaliana* seedlings by altering MT organization. In root cells, excessive levels of DELLAs might disorganize the cytoskeleton of MT, thereby impairing the link between positional signals and cell fate, whereas excessive levels of GAs might stabilize it.

Results of this study also point at a possible role for the DELLAs in regulating the root hair patterning in response to nutritional deficiencies. The random disposition of root hairs under excessive levels of DELLAs might favour the foraging of scarce or non-mobile minerals in deficient soils. Thus, altering the root hair patterning by modulating the levels of GAs/DELLAs might constitute a mechanism used by plants for increasing the possibilities of acquiring non-available minerals, such as P or Fe, in deficient soils. In fact, plant deficiencies in P, B or Fe disarrange the root hair patterning and induce ectopic root hairs (Schmidt *et al.*, 2000; Péret *et al.*, 2011; Janes *et al.*, 2018). Moreover, low availability of P increases the levels of DELLAs and reduces the levels of GAs in roots (Jiang *et al.*, 2007).

Results of this study also suggest that the GAs/DELLAs might affect the root hair patterning in *A. thaliana* seedlings by acting not at the epidermis, where the gene network for the root hair/non-hair epidermal cell fate operates, but at tissues placed underneath (cortex, endodermis and pericycle). However, confirmatory studies are still needed to uncover why the epidermal expression of *gai-1* did not modify the root hair patterning in *A. thaliana* seedlings, in spite that the DELLAs promote the disorganization of MT in root epidermal cells. Moreover, the fact that only one DELLA (*gai-1*) was over-expressed in this study, and that expression of *gai-1* at the epidermis (*MLI::gai-1*, J0951 >> *gai-1*) induced longer and branched root hairs, suggests that the effects of the GAs/DELLAs on the root epidermal cells and/or the root hair patterning in seedlings of *A. thaliana* might be different depending on the particular concentration at which these hormones might be present.

4.2. The GAs/DELLAs might regulate the shape, length and abundance of hairs in root tips of *A. thaliana* seedlings:

Supra-physiological levels of GAs/DELLAs in roots of *A. thaliana* seedlings also induced two-haired cells, two-tipped hairs and branched hairs. Multiple hairs per root epidermal cell, two-tipped hairs and branched hairs have also been reported in the SUPERCENTIPEDE (*scn1*) mutant, with supernumerary root hair initiation sites, in TUA6/AS *A. thaliana* transgenic lines under-expressing α -tubulin genes, in root hair defective 3, 4 and 6 (*rhd3*, *rhd4*, *rhd6*) and *PLD* mutants, in plants treated with MT-depolymerizing oryzalin, MT-disorganizing 1-butanol (a *PLD*-inhibitor) or MT-stabilizing taxol, in *ROP2* (proteins controlling MT organization) over-expressing plants, and in plants subjected to Fe or NO₃⁻ deficiency (Schiefelbein and Somerville, 1990; Schiefelbein *et al.*, 1993; Masucci and Schiefelbein, 1994; Gilroy and Jones, 2000; Schmidt *et al.*, 2000; Bao *et al.*, 2001; Foreman and Dolan, 2001; Grierson and Schiefelbein, 2002; Jones *et al.*, 2002; Gardiner *et al.*, 2003; Müller and Schmidt, 2004; Carol and Dolan, 2006; Ishida *et al.*, 2008; Shin *et al.*, 2011; Pietra, 2014). Interestingly, hormone-induced reorganization of MT is required for the morphogenesis of root hairs (Bao *et al.*, 2001; Schiefelbein, 2003). In turn, the phenotype of root hair branching, due to changes in actin distribution and dynamics, has been related to the induction of genes for GA biosynthesis and CW modification, and reported during legume-rhizobium symbiosis (i.e., soybean infected with *Bradyrhizobium japonicum*), in plants treated with MT-inhibiting drugs, and in mutants of genes necessary for a correct growth of root hairs, such as *TIP1* (involved in the biosynthesis of CW components and probably in the arrangement of actin filaments) and *RHD3* (Schiefelbein and Somerville, 1990; Schiefelbein *et al.*, 1993; Bao *et al.*, 2001; Salazar-Henao *et al.*, 2016).

The disruption of MT also affects trichome branching (Gilroy and Jones, 2000), as actin regulates the shape and growth of trichomes (Rodríguez-Serrano *et al.*, 2014). In addition, the GAs promote trichome branching and influence CW growth (Telfer *et al.*, 1997; Perazza *et al.*, 1998). Thus, the *spy5* mutant (with high levels of GAs and which also displays ectopic root hairs) has over-branched trichomes (Perazza *et al.*, 1998; Mutanwad *et al.*, 2020). On the other hand, during trichome development, the number of branches and the level of endo-reduplication, which is induced by GAs, are closely related (Perazza *et al.*, 1998; Kondorosi *et al.*, 2001).

In this study, excessive levels of DEllAs in *A. thaliana* seedlings also induced longer root hairs near the root tip. Interestingly, nutrient availability prevents root hair elongation (Tsai *et al.*, 2004), whereas deficiencies in P, B or Mg induce root hair elongation, being the higher levels of DEllAs the mediators of the extra-elongation of root hairs (Péret *et al.*, 2011; Liu *et al.*, 2018). Elongated root hairs have also been described in plants exposed to gamma irradiation, Cd or As, as well as in polyploids (Nagata *et al.*, 2004; Setter *et al.*, 2015; Bahmani *et al.*, 2016; Salazar-Henao *et al.*, 2016). Conversely, shorter root hairs have been reported in mutants of the *TIP1*, *PLD ζ 1-PLD ζ 2*, and *RSL4* (a component of GAs signalling) genes (Schiefelbein *et al.*, 1993; Li *et al.*, 2006; Péret *et al.*, 2011). Moreover, the GAs are necessary for root hair elongation, as the *ga 1-3* mutant (deficient in GAs) produces shorter root hairs (Péret *et al.*, 2011). However, the GAs might act at a later stage of root hair development, as apparently, in this study, high levels of GAs did not stimulate root hair elongation near the root tip as much as the high levels of DEllAs did. Therefore, the changes induced, in this study, by excessive levels of GAs/DEllAs on the shape and length of root hairs in seedlings of *A. thaliana* might have been related to the effect of these hormones on the MT cytoskeleton and/or the CW biosynthesis of the root epidermal cells.

Regarding root hair abundance, it is known that nutrient availability inhibits root hair production (Tsai *et al.*, 2004). Excess of Na⁺ reduces root hair abundance (Dinneney *et al.*, 2008), whereas deficiencies in P, Fe or B increase the frequency of root hairs, mainly by inducing ectopic root hair cells (Schiefelbein, 2003; Martín-Rejano *et al.*, 2011; Péret *et al.*, 2011; Shin *et al.*, 2011; Salazar-Henao *et al.*, 2016; Janes *et al.*, 2018). An increased density of root hairs has also been reported in *ROP2* over-expressing plants, in *arm* (*ctl1*; cellulose biosynthesis-related) and *sabre* mutants, in plants exposed to Cd, Vd or As, and in polyploids (Jones *et al.*, 2002; Pietra, 2014; Lin *et al.*, 2015; Bahmani *et al.*, 2016; Salazar-Henao *et al.*, 2016). Interestingly, the levels of GAs determine trichome number (Perazza *et al.*, 1998). In turn, HDA19 controls the response of the root hair density to low P (Chen *et al.*, 2015).

Because of the GAs/DEllAs are involved in plant stress responses (Alonso-Ramírez *et al.*, 2009), the results of this study suggest that these hormones might have a role in regulating the response of the root hair abundance to nutrient availability. In fact, in this study, root hairs near the root tip were denser and longer under excessive DEllAs, but scarcer and shorter under excessive GAs. With this respect, it is known that root hairs grow closer to the root MZ under mechanic stress or B deficiency (Okamoto *et al.*, 2008; Martín-Rejano *et al.*, 2011). Also, the abundance and length of root hairs respond to environmental signals

(Salazar-Henao *et al.*, 2016). Light signalling, for instance, influences root hair length (Grierson and Schiefelbein, 2002). In turn, the photo-period conditions affect the biosynthesis and/or sensibility of GAs (Telfer *et al.*, 1997).

As PLD inhibitors break the organization of MT, which is essential for the correct directionality, elongation and morphology of root hairs (Gardiner *et al.*, 2003), then, the morphological alterations of root hairs observed in this study point to a possible impairment, by excessive levels of GAs/DELLAs, of the actin microfilaments, the cytoskeleton of MT, and the ROP GTPase proteins. In fact, hair cell morphogenesis requires α -tubulin and Rho-like GTPase activity, which, in turn, interacts with katanin P60 to promote MT ordering (Foreman and Dolan, 2001; Lin *et al.*, 2013). Moreover, the SABRE protein (involved in MT organisation and the stabilization of epidermal patterning factors) acts upstream of ROPs (Pietra, 2014).

ACKNOWLEDGEMENTS

Special thanks to Dr. Miguel Ángel Blázquez and Dr. David Alabadí for supporting the writing of this paper. This study was performed at the Blázquez-Alabadí laboratory (Hormone Signalling and Plant Plasticity Group) of the Instituto de Biología Celular y Molecular de Plantas (IBMCP)-UPV-CSIC, Valencia, Spain. Thanks also to Dr. Benedicte Desvoves for the training in the hair/non-hair cell counting (Dr. Crisanto Gutiérrez Lab). I. McCarthy-Suárez also acknowledges a JAE-Doc postdoctoral fellowship ((2008-2011), C.S.I.C., Valencia, Spain). This paper is dedicated to the loving memory of Dr. Francisco Culiáñez-Macià, Principal Investigator at the IBMCP-UPV-CSIC, Valencia, Spain.

REFERENCES

- Alonso-Ramírez A, Rodríguez D, Reyes D, Jiménez JA, Nicolás G, López-Climent M, GómezCárdenas A and Nicolás C** (2009) Evidence for a role of gibberellins in salicylic acid-modulated early plant responses to abiotic stress in *Arabidopsis* seeds. *Plant Physiology* **150**: 1335-1344.
- Bahmani R, Kim DG, Kim JA and Hwang S** (2016) The density and length of root hairs are enhanced in response to cadmium and arsenic by modulating gene expression involved in fate determination and morphogenesis of root hairs in *Arabidopsis*. *Frontiers in Plant Science* **7**: 1-16.
- Bao Y, Kost B and Chua NH** (2001) Reduced expression of α -tubulin genes in *Arabidopsis thaliana* specifically affects root growth and morphology, root hair development and root gravitropism. *The Plant Journal* **28**: 145-157.
- Bouquin T, Mattson O, Naested H, Foster R and Mundy J** (2002) The *Arabidopsis luel* mutant defines a katanin p60 ortholog involved in hormonal control of microtubule orientation during cell growth. *Journal of Cell Science* **116**: 791-801.
- Cao XF, Linstead P, Berger F, Kieber J and Dolan L** (1999) Differential ethylene sensitivity of epidermal cells is involved in the establishment of cell pattern in the *Arabidopsis* root. *Physiologiae Plantarum* **106**: 311-317.
- Carol RJ and Dolan L** (2006) The roles of reactive oxygen species in cell growth: Lessons from root hairs. *Journal of Experimental Botany* **57**: 1829-1834.
- Chen CY, Wu K and Schmidt W** (2015) The histone deacetylase HDA19 controls root cell elongation and modulates a subset of phosphate starvation responses in *Arabidopsis*. *Scientific reports* **5**: 15708. DOI: 10.1038/srep15708.
- Chien JC and Sussex IM** (1996) Differential regulation of trichome formation on the adaxial and abaxial surfaces by gibberellins and photoperiod in *Arabidopsis thaliana* (L.) heyneh. *Plant Physiology* **111**: 1321-1328.

- Cui H and Benfey P** (2009) Interplay between scarecrow, GA and like heterochromatin protein 1 in ground tissue patterning in the *Arabidopsis* root. *The Plant Journal* **58**: 1016-1027.
- Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J and Benfey PN** (2008) Cell identity mediates the responses of *Arabidopsis* roots to abiotic stress. *Science* **320**: 942-945.
- Dugardeyn J and Van der Straeten D** (2008) Ethylene: Fine tuning plant growth and development by stimulation and inhibition of elongation. *Plant Science* **175**: 59-70.
- Foreman J and Dolan L** (2001) Root hairs as a model system for studying plant cell growth. *Annals of Botany* **88**: 1-7.
- Frigerio M, Alabadí D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P and Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signalling in *Arabidopsis*. *Plant Physiology* **142**: 553-563.
- Gardiner J, Collings DA, Harper JDI and Marc J** (2003) The effects of the phospholipase D-antagonist 1butanol on seedlings development and microtubule organisation in *Arabidopsis*. *Plant Cell Physiology* **44**: 687-696.
- Gilroy S and Jones DL** (2000) Through form to function: Root hair development and nutrient uptake. *Trends in Plant Science* **5**: 56-60.
- Grierson C and Schiefelbein J** (2002) Root hairs. The *Arabidopsis* book 2-22. Editors: C.R. Somerville and E.M. Meyerowitz. Publisher: American Society of Plant Biologists. Rockville, MD.
- Hassan H, Scheres B and Blilou I** (2010) Jackdaw controls epidermal patterning in the *Arabidopsis* root meristem through a non-cell autonomous mechanism. *Development* **137**: 1523-1529.
- Ishida T, Kurata T, Okada K and Wada T** (2008) A genetic regulatory network in the development of trichomes and root hairs. *Annual Review of Plant Biology* **59**: 365-386.
- Janes G, von Wangenheim D, Cowling S, Kerr I, Band L, French AP and Bishop A** (2018) Cellular patterning of *Arabidopsis* roots under low phosphate conditions. *Frontiers in Plant Science* **9**: 735.
- Jiang C, Gao X, Liao L, Harberd NP and Fu X** (2007) Phosphate starvation root architecture and anthocyanin accumulation responses are modulated by the gibberellin-DELLA signalling pathway in *Arabidopsis*. *Plant Physiology* **145**: 1460-1470.
- Jones MA, Shen JJ, Fu Y, Li H, Yang Z and Grierson CS** (2002) The *Arabidopsis* ROP2 GTPases is a positive regulator of root hair initiation and tip growth. *The Plant Cell* **14**: 763-776.
- Kappusamy KT, Chen AY and Nemhauser JL** (2009) Steroids are required for epidermal cell fate establishment in *Arabidopsis* roots. *Proceedings of the National Academy of Sciences (PNAS)* **106**: 80738076.
- Kondorosi E, Roudiera F and Gendreau E** (2001) Plant cell size control: growing by ploidy? *Current Opinion in Plant Biology* **3**: 488-492.
- Lee MM and Schiefelbein J** (1999) WEREWOLF, a MYB-related protein in *Arabidopsis*, is a positiondependent regulator of epidermal cell patterning. *Cell* **99**: 473-483.
- Li M, Qin C, Welti R and Wang X** (2006) Double knockouts of phospholipase D ζ 1 and D ζ 2 in *Arabidopsis* affect root elongation during phosphate-limited growth but do not affect root hair patterning. *Plant Physiology* **140**: 761-770.

- Li DX, Chen WQ, Xu ZH and Bai SN** (2015) Histone deacetylase 6-defective mutants show increased expression and acetylation of enhancer of tryptychon and caprice1 and glabra2 with small but significant effects on root epidermis cellular pattern. *Plant Physiology* **168**: 1448-1458.
- Lin D, Cao L, Zhou Z, Zhu L, Ehrhardt D, Yang Z and Fu Y** (2013) Rho GTPase signalling activates microtubule severing to promote microtubule ordering in *Arabidopsis*. *Current Biology* **23**: 290-297.
- Lin CY, Huang LY, Chi WC, Huang TL, Kakimoto T, Tsai CR and Huang HJ** (2015) Pathways involved in vanadate-induced root hair formation in *Arabidopsis*. *Physiologiae Plantarum* **153**: 137-148.
- Liu M, Bi J, Jin C** (2018) Developmental responses of root hairs to Mg deficiency. *Plant Signal and Behaviour* Vol 13, Number 9, e1500068, doi: 10.1080/15592324.2018.1500068
- Locascio A, Blázquez MA and Alabadí D** (2013) Dynamic regulation of cortical microtubule organization through prefoldin-DELLA interaction. *Current Biology* **23**: 804-809.
- Lombardo MC, Graziano M, Polacco JC and Lamattina L** (2006) Nitric oxide functions as a positive regulator of root hair development. *Plant signalling and Behaviour* **1**: 28-33.
- Ma Z, Bielenberg GD, Brown KM and Lynch JP** (2001) Regulation of root hair density of phosphorus availability in *Arabidopsis thaliana*. *Plant Cell and Environment* **24**: 459-467.
- Martín-Rejano EM, Camacho-Cristóbal JJ, Herrera-Rodríguez MB, Rexach J and Navarro-Gochicoa MT** (2011) Auxin and ethylene are involved in the responses of root system architecture to low boron supply in *Arabidopsis* seedlings. *Physiologiae Plantarum* **142**: 170-178.
- Masucci JD and Schiefelbein JW** (1994) The RHD6 mutation of *Arabidopsis thaliana* alters root hair initiation through an auxin and ethylene associated process. *Plant Physiology* **106**: 1335-1346.
- Müller M and Schmidt W** (2004) Environmentally induced plasticity of root hair development in *Arabidopsis*. *Plant Physiology* **134**: 409-419.
- Mutanwad KV, Zangl I, Lucyshyn D** (2020) The Arabidopsis O-fucosyltransferase SPINDLY regulates root hair patterning independently of gibberellin signalling. *Development* **147**(19): dev192039. Doi: 10.1242/dev.192039.
- Nagata T, Todoriki S and Kikuchi** (2004) Radial extension of root cells and elongation of root hairs of *Arabidopsis thaliana* induced by massive doses of gamma irradiation. *Plant Cell Physiology* **45**: 1557-1565.
- Niu Y, Jin C, Jin G, Zhou Q, Lin X, Tang C and Zhang Y** (2011) Auxin modulates the enhanced development of root hairs in *Arabidopsis thaliana* (L.) Heyhn. under elevated CO₂. *Plant, Cell and Environment* **34**: 1304-1317.
- Okamoto T, Tsurumi S, Shibasaki K, Obana Y, Takaji H, Oono Y and Rahman A** (2008) Genetic dissection of hormonal responses in the roots of *Arabidopsis* grown under continuous mechanical impedance. *Plant Physiology* **146**: 1651-1662.
- Perazza D, Vachon G and Herzog M** (1998) Gibberellins promote trichome formation by up-regulating *GLABROUS1* in *Arabidopsis*. *Plant Physiology* **117**: 375-383.
- Péret B, Clément M, Nussaume L and Desnos T** (2011) Root developmental adaptation to phosphate starvation: Better safe than sorry. *Trends in Plant Science* **16**: 442-450.
- Pernas M, Ryan E and Dolan L** (2010) Schizorrrhyza controls tissue system complexity in plants. *Current Biology* **20**: 812-823.
- Pietra S** (2014) Characterization of new players in planar polarity establishment in *Arabidopsis*. Doctoral thesis Umea Plant Science Centre Fysiologisk Botanik.

- Pietra S, Lang P and Grebe M** (2015) *SABRE* is required for stabilization of root hair patterning in *Arabidopsis thaliana*. *Physiologiae Plantarum* **153**: 440-453.
- Rishmawi L, Pesch M, Juengst C, Schauss AC, Schrader A and Hülskamp M** (2014) Non-cell autonomous regulation of root hair patterning genes by WRKY75 in *Arabidopsis*. *Plant Physiology* **165**: 186-195.
- Rodríguez-Serrano M, Pazmiño DM, Sparkes I, Rochetti A, Hawes C, Romero-Puertas and Sandalio LM** (2014) 2,4-dichlorophenoxyacetic acid promotes S-nitrosylation and oxidation of actin affecting cytoskeleton and peroxisomal dynamics. *Journal of Experimental Botany* **65**: 4783-4793.
- Salazar-Henao JE, Vélez-Bermúdez IC and Schmidt W** (2016) The regulation and plasticity of root hair patterning and morphogenesis. *Development* **143**: 1848-1858.
- Sánchez-Calderón L, Ibarra-Cortés ME and Zepeda-Jazo I** (2013) Root development and abiotic stress adaptation. Abiotic stress-plant responses and applications in agriculture. Chapter 5, 135-168. Intech Open Science.
- Schiefelbein J** (2003) Cell fate specification in the epidermis: A common patterning mechanism in the root and the shoot. *Current opinion in Plant Biology* **6**: 74-78.
- Schiefelbein JW and Somerville C** (1990) Genetic control of root hair development in *Arabidopsis thaliana*. *The Plant Cell* **2**: 235-243.
- Schiefelbein J, Galway M, Masucci J and Ford G** (1993) Pollen tube and root hair tip growth is disrupted in a mutant of *Arabidopsis thaliana*. *Plant Physiology* **103**: 979-985.
- Schiefelbein J, Masucci JD and Wang H** (1997) Building a root: The control of patterning and morphogenesis during root development. *The Plant Cell* **9**: 1089-1098.
- Schiefelbein J, Kwak SH, Wieckowski Y, Barron C and Bruex A** (2009) The gene regulatory network for root epidermal cell-type pattern formation in *Arabidopsis*. *Journal of Experimental Botany* **60**: 1515-1521.
- Schmidt W, Tittel J and Schikora A** (2000) Role of hormones in the induction of iron deficiency responses in *Arabidopsis* roots. *Plant Physiology* **122**: 1109-1118.
- Schneider K, Wells B, Dolan L and Roberts K** (1997) Structural and genetic analysis of epidermal cell differentiation in *Arabidopsis* primary roots. *Development* **124**: 1789-1798.
- Sedbrook JD** (2004) MAPS in plant cells: delineating microtubule growth dynamics and organisation. *Current Opinion in Plant Biology* **7**: 632-640.
- Setter MG, Schmid and Ludewig U** (2015) Uncovering genes and ploidy involved in the high diversity in root hair density, length and response to local scarce phosphate in *Arabidopsis thaliana*. *PLoS ONE* **10**: e0120604.
- Shin LJ, Huang HE, Chang H, Lin YN, Feng TY and Ger MJ** (2011) Ectopic ferredoxin I protein promotes root hair growth through induction of reactive oxygen species in *Arabidopsis thaliana*. *Journal of Plant Physiology* **168**: 434-440.
- Silverman FP, Assiahmah AS and Bush DS** (1998) Membrane transport and cytokinin action in root hairs of *Medicago sativa*. *Planta* **205**: 23-31.
- Telfer A, Bollman KM and Poethig RS** (1997) Phase change and the regulation of trichome distribution in *Arabidopsis thaliana*. *Development* **124**: 645-654.
- Traw MB and Bergelson J** (2003) Interactive effects of jasmonic acid, salicylic acid, and gibberellin on induction of trichomes in *Arabidopsis*. *Plant Physiology* **133**: 1367-1375.

Tsai SL, Harris PJ and Lovell PH (2004) Bands of root hairs are produced in tomato (*Lycopersicon esculentum*) in response to specific combinations of thermoperiods and photoperiods. *New Zealand Journal of Crop Horticultural Sciences* **32**: 121-129.

Van Hengel AJ, Barber C and Roberts K (2004) The expression patterns of arabinogalactan-protein AtAGP30 and GLABRA2 reveal a role for abscisic acid in the early stages of root epidermal patterning. *The Plant Journal* **39**: 70-83.

Wang X (2005) Regulatory functions of phospholipase D and phosphatidic acid in plant growth, development and stress responses. *Plant Physiology* **139**: 566-573.

Webb M, Jouannic S, Foreman J, Linstead P and Dolan L (2002) Cell specification in the *Arabidopsis* root epidermis requires the activity of ECTOPIC ROOT HAIR 3-a katanin P60 protein. *Development* **129**: 123-131.

Xu CR, Liu C, Wang YL, Li LC, Chen WQ, Xu ZN and Bai SN (2005) Histone deacetylation affects expression of cellular patterning genes in the *Arabidopsis* root epidermis. *Proceedings of the National Academy of Sciences (PNAS)* **20**: 14469-14474.

Yang T, Savage N and Schmidt W (2007) Plasticity of root epidermal cell fate in response to nutrient starvation. 18th International Conference on *Arabidopsis* Research. P-116. Abstract available at: https://www.arabidopsis.org/news/2007_18thICAR_CompleteProgram.pdf.

Yu Q, Li P, Liang N, Wang H, Xu M and Wu S (2017) Cell fate specification in *Arabidopsis* roots requires coordinative action of lineage instruction and positional reprogramming. *Plant Physiology* **175**: 816-827.

FIGURE LEGENDS

Fig. 1A. Spatial gene expression of the root hair (*CPC) and non-hair (GL2, *EGL3, WER) epidermal cell fate markers in 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** *GL2pro::GUS* (MS), 20X; **B)** *GL2pro::GUS* (0.5 μ M PAC), 20X; **C)** *GL2pro::GUS* (1 μ M GA₄), 20X; **D)** *GL2pro::GUS* (30 μ M GA₃), 20X; **E)** *Ler x GL2pro::GUS* (MS), 20X; **F)** *gai-1 x GL2pro::GUS* (MS), 20X; **G)** *GID1b-ox x GL2pro::GUS* (24h in H₂O; liquid incubation experiment; leaky line), 20X; **H)** *GID1b-ox x GL2pro::GUS* (24h in 1 μ M GA₄; liquid incubation experiment), 20X; **I)** *HSp::gai-1 x GL2pro::GUS* (22°C, 4h), 20X; **J)** *HSp::gai-1 x GL2pro::GUS* (37°C, 4h), 20X; **K)** *pGAI::gai-1:GR x GL2pro::GUS* (24h in MS; leaky line), 20X; **L)** *pGAI::gai-1:GR x GL2pro::GUS* (24h in 10 μ M DEXA), 20X; **M)** *SCR::gai-1:GR x GL2pro::GUS* (24h in MS; leaky line), 20X; **N)** *SCR::gai-1:GR x GL2pro::GUS* (24h in 10 μ M DEXA), 20X; **O)** *CPCpro::GUS* (MS), 20X; **P)** *CPCpro::GUS* (0.5 μ M PAC), 20X; **Q)** *CPCpro::GUS* (1 μ M GA₄), 20X; **R)** *EGL3pro::GUS* (MS), 20X; **S)** *EGL3pro::GUS* (0.5 μ M PAC), 20X; **T)** *EGL3pro::GUS* (1 μ M GA₄), 20X; **U)** *WERpro::GFP* (MS), 40X; **V)** *WERpro::GFP* (0.5 μ M PAC), 40X; **W)** *WERpro::GFP* (1 μ M GA₄), 40X. In control seedlings, *GL2* is expressed in root non-hair (atrachoblast) cells. *CPC protein is expressed in root non-hair cells, but migrates to root hair cells. *EGL3 protein is expressed in root hair cells, but migrates to root non-hair cells.

Fig. 1B. Spatial gene expression of the root non-hair epidermal cell fate marker *GL2* in cross sections of resin-embedded roots of *A. thaliana* seedlings grown for 5 days under excessive levels of GAs/DELLAs. **A)** *GL2pro::GUS* (MS); **B)** *GL2pro::GUS* (0.5 μ M PAC): Lack of *GL2* expression in an atrichoblast cell; **C)** *GL2pro::GUS* (0.5 μ M PAC): Ectopic expression of *GL2* in a trichoblast cell; **D)** *GL2pro::GUS* (1 μ M GA₄): Ectopic expression of *GL2* in a trichoblast cell and lack of *GL2* expression in an atrichoblast cell; **E)** *GL2pro::GUS* (1 μ M GA₄): Lack of *GL2* expression in an atrichoblast cell; **F)** Ectopic root hair cell in *gai-1*. Magnification: 40X.

Fig. 2A. Ectopic root hairs and ectopic root non-hairs in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** Col(0) (MS) correct hair (epidermis); **B)** Col(0) (MS), correct hair (cortex); **C)** Col(0) (0.5 μ M PAC) ectopic hair (epidermis); **D)** Col(0) (0.5 μ M PAC) ectopic hair (cortex); **E)** Col(0) (1 μ M GA₄) ectopic hair (epidermis); **F)** Col(0) (1 μ M GA₄) ectopic hair (cortex); **G)** Col(0) (MS) correct non-hair (epidermis); **H)** Col(0) (MS) correct non-hair (cortex); **I)** Col(0) (0.5 μ M PAC) ectopic non-hair (epidermis); **J)** Col(0) (0.5 μ M PAC) ectopic non-hair (cortex); **K)** Col(0) (1 μ M GA₄) ectopic non-hair (epidermis); **L)** Col(0) (1 μ M GA₄) ectopic non-hair (cortex); **M)** *Ler*, correct hair and non-hair (epidermis); **N)** *Ler*, correct hair and non-hair (cortex); **O)** *gai-1*, ectopic hair (epidermis); **P)** *gai-1*, ectopic hair (cortex); **Q)** *QD*, ectopic hair (epidermis); **R)** *QD*, ectopic hair (cortex); **S)** *5X*, ectopic non-hair (epidermis); **T)** *5X*, ectopic non-hair (cortex); **U)** *pGAI::gai-1:GR* (10 μ M DEXA), ectopic non-hair and ectopic hair (epidermis); **V)** *pGAI::gai-1:GR* (10 μ M DEXA), ectopic non-hair and ectopic hair (cortex); **W)** *HSp::gai-1*, 2d after heat shock (37°C, 4h), ectopic hair (epidermis); **X)** *HSp::gai-1*, 2d after heat-shock (37°C, 4h), ectopic hair (cortex). Magnification: 20X.

Fig. 2B. Ectopic hairs and non-hairs in root tips of 5-day-old *A. thaliana* seedlings over-expressing the DELLA *gai-1* in different tissues of the root. **A)** *UAS::gai-1* x J2812, ectopic hairs (epidermis); **B)** *UAS::gai-1* x J2812, ectopic hairs (cortex); **C)** *UAS::gai-1* x J2812 (ectopic non-hair, epidermis); **D)** *UAS::gai-1* x J2812 (ectopic non-hair, cortex); **E)** *UAS::gai-1* x Q2393 (ectopic hair, epidermis); **F)** *UAS::gai-1* x Q2393 (ectopic hair, cortex); **G)** *UAS::gai-1* x Q2393 (ectopic non-hair, epidermis); **H)** *UAS::gai-1* x Q2393 (ectopic non-hair, cortex); **I)** *UAS::gai-1* x Q2500 (ectopic hair, epidermis); **J)** *UAS::gai-1* x Q2500 (ectopic hair, cortex); **K)** *UAS::gai-1* x J0121 (ectopic non-hair, epidermis); **L)** *UAS::gai-1* x J0121 (ectopic non-hair, cortex). Magnification: 20X.

Fig. 3A. Arrangement of hairs in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** Col(0) (MS), 10X; **B)** Col(0) (0.5 μ M PAC), 10X; **C)** Col(0) (30 μ M GA₃), 10X; **D)** *Ler*, 10X; **E)** *gai-1*, 10X; **F)** *QD*, 10X; **G)** *5X*, 10X; **H)** *GID1b-ox* (MS, leaky line), lateral root, 10X; **I)** *pGAI::gai-1:GR* (30h in MS; leaky line), 10X; **J)** *pGAI::gai-1:GR* (30h in 10 μ M DEXA), 10X; **K)** *SCR::gai-1:GR* (72h in MS; leaky line), 10X; **L)** *SCR::gai-1:GR* (48h in 10 μ M DEXA), 10X; **M)** *UAS::gai-1* x C24, 10X; **N)** *UAS::gai-1* x J0951, 10X; **O)** *UAS::gai-1* x J2812, 10X; **P)** *UAS::gai-1* x N9142, 10X; **Q)** *UAS::gai-1* x M0018, 10X; **R)** *UAS::gai-1* x Q2500, 10X; **S)** *UAS::gai-1* x Q2393, 10X; **T)** *UAS::gai-1* x J0121, 10X; **U)** *UAS::gai-1* x J0631, 10X; **V)** *UAS::gai-1* x J0571, 10X; **W)** *UAS::gai-1* x J3281, 4X; **X)** *ML1::gai-1*, 10X.

Fig. 3B. Adjacent hair rows in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** Col(0) (0.5 μ M PAC), 20X; **B)** *gai-1* (lateral root), 10X; **C)** *QD*, 10X; **D)** *HSp::gai-1*, 2 days after heat shock (37°C, 4h), 20X; **E)** *pGAI::gai-1:GR* (MS, leaky line), 10X; **F)** *SCR::gai-1:GR* (MS, leaky line), 20X; **G)** *UAS::gai-1* x J2812, 10X; **H)** *UAS::gai-1* x M0018, 10X; **I)** *UAS::gai-1* x Q2393, 10X.

Fig. 4. Morphology of trichoblasts and root hairs in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** Two-haired cell in PAC (0.5 μ M), 20X; **B)** Cell with two hair bulges in PAC (0.5 μ M), 20X; **C)** Two-haired cells in *gai-1*, 10X; **D)** Two-haired cells in *QD*, 10X; **E)** Two-haired cells and branched hairs in *UAS::gai-1* x Q2393, 10X; **F)** Two-tipped hairs in PAC (0.5 μ M), 20X; **G)** Two-tipped hairs in GA₄ (1 μ M), 20X; **H)** Two-tipped hairs in *gai-1*, 20X; **I)** Two-tipped hairs in *QD*, 20X; **J)** Two-tipped hairs in *pGAI::gai-1:GR* (10 μ M DEXA), 20X; **K)** Two-tipped hairs in *UAS::gai-1* x J0121, 10X; **L)** Two-tipped and branched hairs in PAC (0.5 μ M), 20X; **M)** Branched hairs in PAC (0.5 μ M), 20X; **N)** Branched hairs in *gai-1*, 20X; **O)** Branched hairs in *QD*, 20X; **P)** Branched hairs in *pGAI::gai-1:GR* (10 μ M DEXA), 20X; **Q)** Branched hairs in *SCR::gai-1:GR* (MS; leaky line), 20X; **R)** Branched hairs in *UAS::gai-1* x J0951, 20X; **S)** Branched hairs in *UAS::gai-1* x J2812, 20X; **T)** Branched hairs in *UAS::gai-1* x N9142, 10X; **U)** Branched hairs in *UAS::gai-1* x Q2393, 10X; **V)** Branched hairs in *UAS::gai-1* x J0121, 10X; **W)** Branched hairs in *UAS::gai-1* x J0631, 10X; **X)** Branched hairs in *ML1::gai-1*, 20X.

Fig. 5A. Length and abundance of hairs in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** Col(0) (MS); **B)** Col(0) (0.5 μ M PAC); **C)** Col(0) (1 μ M GA₄); **D)** *Ler*; **E)** *gai-1*; **F)** *QD*; **G)** *5X*; **H)** *GID1b-ox* (MS; leaky line); **I)** *HSp::gai-1* (24h at 22°C); **J)** *HSp::gai-1* (24h after heat-shock (37°C for 4h)); **K)** *pGAI::gai-1:GR* (30h in MS; leaky line); **L)** *pGAI::gai-1:GR* (30h in 10 μ M DEXA); **M)** *SCR::gai-1:GR* (24h in MS; leaky line); **N)** *SCR::gai-1:GR* (24h in 10 μ M DEXA).

DEXA); **O**) *UAS::gai-1* x C24; **P**) *UAS::gai-1* x J0951; **Q**) *UAS::gai-1* x J2812; **R**) *UAS::gai-1* x N9142; **S**) *UAS::gai-1* x M0018; **T**) *UAS::gai-1* x J0571; **U**) *UAS::gai-1* x Q2500; **V**) *UAS::gai-1* x Q2393; **W**) *UAS::gai-1* x J0631; **X**) *UAS::gai-1* x J0121. Magnification: 4X.

Fig. 5B. Length and abundance of hairs in root tips of *wer*, *cpc* and *35S::CPC* mutant seedlings grown for 5 days under excessive levels of GAs/DELLAs. **A**) *wer* mutant (MS), lens; **B**) *wer* mutant (0.5 μ M PAC), lens; **C**) *wer* mutant (1 μ M GA₄), lens; **D**) *wer* mutant (MS), microscope; **E**) *wer* mutant (0.5 μ M PAC), microscope; **F**) *wer* mutant (1 μ M GA₄), microscope; **G**) *cpc* mutant (MS), lens; **H**) *cpc* mutant (0.5 μ M PAC), lens; **I**) *cpc* mutant (1 μ M GA₄), lens; **J**) *cpc* mutant (MS), microscope; **K**) *cpc* mutant (0.5 μ M PAC), microscope; **L**) *cpc* mutant (1 μ M GA₄), microscope; **M**) *35S::CPC* mutant (MS), lens; **N**) *35S::CPC* mutant (1 μ M GA₄), lens. Magnification: 2.5 X (lens), 10X (microscope).

FIGURE 1A

FIGURE 1B

FIGURE 2A

Figure 2B

FIGURE 3B

FIGURE 4

FIGURE 5A

FIGURE 5B

TABLES

Table 1. Distribution of hair and non-hair cells at the trichoblast/atrichoblast positions in roots of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Numbers in parenthesis refer to the number of cells analysed. At least 15-20 roots were used per treatment.

	Trichoblast position		Atrichoblast position	
	Hair cell (%)	Non-hair cell (%)	Hair cell (%)	Non-hair cell (%)
Col (0) (MS)	97.5 ± 0.7 (73)	2.5 ± 0.7 (2)	0 ± 0 (0)	100 ± 0 (75)
PAC (0.5 µM)	77.4 ± 5.7 (109)	22.6 ± 5.7 (32)	36.7 ± 8.2 (80)	63.3 ± 8.2 (138)
GA ₄ (1 µM)	81 ± 2.7 (83)	19 ± 2.7 (20)	12.5 ± 3.5 (5)	87.5 ± 3.5 (35)
PAC (0.5 µM) + GA ₄ (1 µM)	94 ± 4.2 (71)	6 ± 4.2 (5)	5 ± 2.8 (4)	95 ± 2.8 (71)
Ler	95.8 ± 2.2 (167)	4.2 ± 2.2 (7)	4.5 ± 3.5 (6)	95.5 ± 3.5 (120)
<i>gai-1</i>	82.7 ± 4.5 (75)	17.3 ± 4.5 (16)	40.4 ± 5 (55)	59.6 ± 5 (81)
<i>QD</i>	78.8 ± 4.5 (126)	21.2 ± 4.5 (34)	24 ± 4.9 (38)	76 ± 4.9 (122)
<i>pGAI::gai-1:GR</i> (MS)	93.5 ± 2.1 (41)	6.5 ± 2.1 (3)	25 ± 7.1 (10)	75 ± 7.1 (30)
<i>pGAI::gai-1:GR</i> (10 µM DEXA)	78 ± 2.8 (38)	22 ± 2.8 (11)	50.5 ± 6.4 (22)	49.5 ± 6.4 (22)
<i>SCR::gai-1:GR</i> (MS)	83.8 ± 3.3 (36)	16.2 ± 3.3 (7)	35 ± 6.2 (13)	65 ± 6.2 (25)
<i>SCR::gai-1:GR</i> (0.1 µM DEXA)	67 ± 12.7 (30)	33 ± 12.7 (15)	15 ± 7.1 (6)	85 ± 7.1 (34)

Table 2. Percentage of ectopic root hair/non-hair cells at the trichoblast/atrichoblast positions in roots of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. *Seedlings analysed at 48h after a heat-shock experiment (37°C, 4h). *GL2pro::GUS* (22°C): control seedlings grown at 22°C for 4h. *GL2pro::GUS* (37°C): control seedlings grown at 37°C for 4h (ectopic root hair cells might have appeared due to heat stress). *Hsp::gai-1* x *GL2pro::GUS* (37°C): inducible *gai-1* mutant seedlings grown at 37°C for 4h. The number of ectopic root hair and non-hair cells from a single experiment is shown in parenthesis.

	Trichoblast position		Atrichoblast position	
	Number of epidermal cells examined	% Ectopic root non-hair cells	Number of epidermal cells examined	% Ectopic root hair cells
<i>Ler</i>	41	2 (1)	30	7 (2)
<i>5X</i>	20	35 (7)	20	0 (0)
<i>GAI-ox</i>	15	7 (1)	15	53 (8)
<i>GL2pro::GUS</i> (22°C) *	30	0 (0)	29	0 (0)
<i>GL2pro::GUS</i> (37°C)*	30	7 (2)	30	30 (9)
<i>Hsp::gai-1</i> x <i>GL2pro::GUS</i> (37°C)*	29	28 (8)	27	44 (12)
<i>wer</i>	28	32 (9)	36	50 (18)
<i>wer</i> (1 µM GA ₄)	30	20 (6)	29	72 (21)
<i>cpc</i>	28	64 (18)	28	21 (6)
<i>cpc</i> (0.5 µM PAC)	28	54 (15)	28	4 (1)
<i>35S::CPC</i>	30	23 (7)	30	60 (18)
<i>35S::CPC</i> (1 µM GA ₄)	30	13 (4)	29	17 (5)
<i>SCR::gai-1:GR</i> (MS)	21	19 (4)	20	30 (6)
<i>SCR::gai-1:GR</i> (0.2 µM DEXA)	19	21 (4)	19	26 (5)
<i>SCR::gai-1:GR</i> (0.5 µM DEXA)	20	30 (6)	18	22 (4)
<i>SCR::gai-1:GR</i> (1.2 µM DEXA)	10	20 (2)	10	40 (4)
<i>UAS::gai 1</i> x <i>C24</i> (control)	50	0 (0)	44	0 (0)
<i>ML1::gai-1</i>	41	2 (1)	30	3 (1)
<i>UAS::gai-1</i> x <i>J0951</i>	60	0 (0)	60	0 (0)
<i>UAS::gai-1</i> x <i>J2812</i>	30	10 (3)	30	50 (15)
<i>UAS::gai-1</i> x <i>Q2393</i>	9	22 (2)	16	63 (10)

Table 3. Length and abundance of hairs in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Analyses of hair length and abundance were performed on electron micrographs of root tips of *A. thaliana* seedlings (4X). (*) Seedlings analysed at 24h and 48h after a heat-shock experiment (37°C, 4h). Analyses of hair abundance performed at 31.5X (lens; Control, PAC and GA₄), 3.2X (lens; *Ler*, *gai-1* and *QD*) or 4X (microscope; other mutants and *UAS::gai-1* lines).

	Hairs analysed	Root Hair Length (µm)	Roots examined	N° of Root Hairs per field
Col (0) (MS)	69	209 ± 121 (100%)	17	38 ± 8 (100 %)
Col (0) (0.5 µM PAC)	94	270 ± 128 (129 %)	19	54 ± 12 (142 %)
Col (0) (1 µM GA ₄)	37	178 ± 93 (85 %)	18	31 ± 8 (82 %)
<i>Ler</i>	45	201 ± 99 (100 %)	5	43 ± 7 (100 %)
<i>gai-1</i>	120	397 ± 186 (198 %)	3	56 ± 1 (130 %)
<i>QD</i>	25	139 ± 85 (69 %)	3	24 ± 5 (56 %)
<i>GID1b-ox</i>	14	80 ± 25 (40 %)	6	28 ± 8 (88 %)
<i>GID1b-ox</i> (30 µM GA ₃)	10	64 ± 35 (32 %)	3	18 ± 6 (57 %)
<i>Hsp::gai-1</i> (22 °C) at 24h (*)	6	55 ± 14 (100 %)	1	18 ± 0 (100 %)
<i>Hsp::gai-1</i> (37 °C) at 24h (*)	11	405 ± 208 (201 %)	2	32 ± 4 (178 %)
<i>Hsp::gai-1</i> (37 °C) at 48h (*)	23	411 ± 165 (204 %)	3	83 ± 31 (459 %)
<i>pGAI::gai-1:GR</i> (MS, 30h)	40	270 ± 118 (100 %)	4	56 ± 7 (100 %)
<i>pGAI::gai-1:GR</i> (0.5 µM DEXA, 30h)	57	314 ± 177 (116 %)	8	79 ± 17 (142 %)
<i>SCR::gai-1:GR</i> (MS, 3d)	30	245 ± 87 (100 %)	3	49 ± 20 (100 %)
<i>SCR::gai-1:GR</i> (0.5 µM DEXA, 3d)	35	507 ± 173 (207 %)	5	76 ± 31 (154 %)
<i>wer</i> (MS)	24	192 ± 88 (100 %)	3	91 ± 12 (100 %)
<i>wer</i> (0.5 µM PAC)	8	243 ± 134 (127 %)	3	125 ± 29 (137 %)
<i>wer</i> (1 µM GA ₄)	6	133 ± 23 (70 %)	3	70 ± 5 (77 %)
<i>cpc</i> (MS)	7	104 ± 29 (100%)	3	17 ± 1 (100 %)
<i>cpc</i> (0.5 µM PAC)	9	213 ± 92 (204 %)	3	18 ± 2 (106 %)
<i>cpc</i> (1 µM GA ₄)	8	88 ± 51 (85 %)	7	11 ± 3 (65 %)
<i>UAS::gai-1</i> x C24 (control)	20	161 ± 105 (100 %)	2	48 ± 23 (100 %)
<i>UAS::gai-1</i> x J0951	34	240 ± 118 (149 %)	3	49 ± 13 (101 %)
<i>UAS::gai-1</i> x J2812	59	243 ± 118 (151 %)	9	77 ± 26 (161 %)
<i>UAS::gai-1</i> x J0571	25	586 ± 273 (364 %)	2	60 ± 11 (125 %)
<i>UAS::gai-1</i> x M0018	90	685 ± 195 (425 %)	10	92 ± 26 (192 %)
<i>UAS::gai-1</i> x Q2500	37	680 ± 189 (422 %)	2	96 ± 12 (200 %)
<i>UAS::gai-1</i> x Q2393	48	272 ± 146 (169 %)	4	67 ± 26 (140 %)
<i>UAS::gai-1</i> x N9142	21	195 ± 97 (121 %)	2	30 ± 1 (63 %)
<i>UAS::gai-1</i> x J0121	47	233 ± 120 (145 %)	5	67 ± 13 (140 %)
<i>UAS::gai-1</i> x J0631	8	386 ± 129 (240 %)	2	96 ± 15 (200 %)

Table 4. Percentage and estimated number of epidermal cells at the trichoblast/atrichoblast positions per root cross section in 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Analyses performed on electron micrographs of cross sections of resin-embedded roots (40X).

	Number of root cross sections examined	% Epidermal Cells at Trichoblast position	% Epidermal Cells at Atrichoblast position	Average number of epidermal cells per root cross section	Predicted N° of epidermal cells at the Trichoblast position per root radial section	Predicted N° of epidermal cells at the Atrichoblast position per root radial section
Control	19	35.5 ± 0.8	64.5 ± 0.8	23 ± 1	8 (100 %)	15 (100 %)
PAC (0.5 µM)	25	29.8 ± 2	70.2 ± 2	27 ± 2	8 (100 %)	19 (127 %)
GA ₄ (1 µM)	20	36 ± 3	64 ± 3	23 ± 2	8 (100 %)	15 (100 %)
Ler	20	39.1 ± 3.8	60.9 ± 3.8	21 ± 2	8 (100 %)	13 (100 %)
<i>gai-1</i>	19	34.9 ± 1	65.1 ± 0.9	23 ± 1	8 (100 %)	15 (115 %)
QD	31	40.8 ± 6.2	59.2 ± 6.2	23 ± 1	9 (113 %)	14 (108 %)
5X	22	41.5 ± 2.4	58.5 ± 2.4	20 ± 3	8 (100 %)	12 (92 %)

Table 5. *Estimated number of root hairs and non-hairs per root cross section in root tips of 5-day-old A. thaliana seedlings grown under (or harbouring) excessive levels of GAs/DELLAs.* Calculations were made by considering the data of Table 1 (distribution of hair and non-hair cells at the trichoblast/atrichoblast positions), Table 2 (percentage of ectopic root hair/non-hair cells at the atrichoblast/trichoblast positions) and Table 4 (average number of epidermal cells per root cross section, estimated number of epidermal cells at the trichoblast position per root cross section, and estimated number of epidermal cells at the atrichoblast position per root cross section). Estimated number of root hairs per root cross section = [hairs at the trichoblast position + hairs at the atrichoblast position]. Estimated number of root non-hairs per root cross section = [non-hairs at the atrichoblast position + non-hairs at the trichoblast position].

	Trichoblast position		Atrichoblast position		Estimated N° of Root Hairs per root cross section	Estimated N° of Non-root hairs per root cross section
	Hairs per root cross section	Non-hairs per root cross section	Hairs per root cross section	Non-hairs per root cross section		
Control	8	0	0	15	8 (100 %)	15 (100 %)
PAC (0.5 µM)	6	2	7	12	13 (163 %)	14 (93 %)
GA ₄ (1 µM)	6	2	2	13	8 (100 %)	15 (100 %)
Ler	8	0	1	12	9 (100 %)	12 (100 %)
<i>gai-1</i>	7	1	6	9	13 (144 %)	10 (83 %)
QD	7	2	3	11	10 (111 %)	13 (108 %)
5X	5	3	0	12	5 (56 %)	15 (125 %)