Characterization of BLUF-photoreceptors present in *Acinetobacter nosocomialis*.

Inés Abatedaga^{1¶}, Bárbara Pérez Mora^{2¶}, Marisel Tuttobene², Gabriela Müller², Daiana Biancotti², Claudio D. Borsarelli¹, Lorena Valle^{1*}, Maria A. Mussi^{2*}

* co-corresponding authors

E-mail: mussi@cefobi-conicet.gov.ar (MAM); lvalle@unse.edu.ar (LV)

[¶] These authors contributed equally to this work

¹ Instituto de Bionanotecnología del NOA (INBIONATEC-CONICET), Universidad Nacional de Santiago del Estero (UNSE), Santiago del Estero, Argentina

² Centro de Estudios Fotosintéticos y Bioquímicos (CEFOBI-CONICET), Universidad Nacional de Rosario (UNR), Rosario, Argentina

Abstract

Acinetobacter nosocomialis is a Gram-negative opportunistic pathogen, whose ability to cause disease in humans is well recognized. Blue light has been shown to modulate important physiological traits related to persistence and virulence in this microorganism. In this work, we characterized the three Blue Light sensing Using FAD (BLUF) domain-containing proteins encoded in the *A. nosocomialis* genome, which account for the only "traditional" light sensors present in this microorganism. By focusing on a light-modulated bacterial process such as motility, the temperature dependence of light regulation was studied, as well as the expression pattern and spectroscopic characteristics of the different *A. nosocomialis* BLUFs. Our results show that the three BLUF-containing proteins encode active photoreceptors, despite only two of them are stable in the light-regulatory temperature range when expressed recombinantly. *In vivo*, only the *A. baumannii*'s ortholog AnBLUF65 is expressed, which is active in a temperature range from 15 °C to 37 °C. In turn, AnBLUF46 is an active photoreceptor between 15 °C to 32 °C *in vitro*, but is not expressed in *A. nosocomialis* in the conditions tested. Intra-protein interactions were analyzed using 3D models built based on *A. baumannii*'s photoreceptor, to support spectroscopic data and profile intra-protein residue interactions. A general scheme is presented on how hydrophobic/aromatic interactions may contribute to the stability of dark/light- adapted states, indicating the importance of these interactions in BLUF photoreceptors.

Introduction

Acinetobacter nosocomialis is a Gram-negative coccobacillus, member of the *Acinetobacter calcoaceticus*-*Acinetobacter baumannii* (ACB) complex [1]. While *A. baumannii* predominates over all other members of the ACB complex in terms of incidence, poorer clinical outcomes, and antibiotic resistance rates, *A. nosocomialis* has gained recognition also as a clinically relevant human pathogen [2].

Although *Acinetobacter* spp. are primarily associated with pneumonia, they are also frequent causes of wound and burn infections, meningitis, urinary tract infections, and sepsis [3]. There is a growing trend for these isolates to display high levels of antibiotic resistance, with some being resistant to all clinically available antibiotics [4].

The ability of *A. nosocomialis* to cause disease in humans is well-recognized [5-7]. Many potential virulence factors have been identified in *A. nosocomialis* and include a CTFR inhibitory factor (Cif), a protein O-glycosylation system, a type-I secretion system, a type-II secretion system, secretion of outer membrane vesicles, the OmpA protein, the CpaA protease, and quorum sensing [2].

We have previously shown that *A. nosocomialis* is able to sense and respond to light modulating biofilm formation and motility at 24 °C [8]. Also, we have shown that light modulates persistence, metabolism, the ability to grow under iron limiting conditions and virulence in this microorganism [9]. The genome of *A. nosocomialis* RUH2624 encodes three Blue Light sensing Using FAD (BLUF)-domain containing proteins, as the only "traditional" light sensors [8]. Extensive work performed on *A. baumannii* showed that this microorganism encodes only one BLUF-type photoreceptor, designated BlsA, which functions at low-environmental temperatures up to 24 °C and is regulated both at the transcriptional level as well as the activity of the photocycle [10, 11]. Also other BLUFs from *Acinetobacter* have been characterized based on light induced phenotypes, gene knockouts and transcriptomic analyses [12, 13]. In this work, we present evidence indicating that regulation

of motility by light in *A. nosocomialis* is maintained in a wide range of temperatures from 24 to 37 °C. Recombinant expression, purification and characterization of the different BLUF-domain containing-proteins showed that the three of them encode active photoreceptors; however only AnBLUF46 and AnBLUF65 are stable. Interestingly, only *anbluf65* is expressed *in vivo* and exhibits a stable photocycle in the temperature-range at which light regulates motility in *A. nosocomialis*. Spectroscopic characterization and analyses of 3D models built for these proteins provide insights into the intra-protein signaling process connecting the widely characterized BLUF photophysics with the subtle re-arrangements located in the C-terminal part of these proteins. Finally, proteomic analyses revealed that light mainly regulates proteins related to signalling and cellular metabolism.

Material and Methods

Bacterial Strains, Plasmids, and Media.

Luria-Bertani (LB) broth (Difco) and agar (Difco) were used to grow and maintain bacterial strains. Broth cultures were incubated at the indicated temperatures either statically or with shaking at 200 rpm.

Blue light treatments.

Blue light treatments were performed as described in our previous studies [8-11, 14-18]. Briefly, cells were incubated at different temperatures in the dark or under blue light emitted by 9-light-emitting diode (LED) arrays, with an intensity of 6 to 10 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Each array was built using 3-LED module strips emitting blue light, with emission peaks centered at 462 nm, determined using the LI-COR LI-1800 spectroradiometer [15].

Cell Motility assays.

Cell motility was tested on 1% tryptone, 0.5% NaCl, 0.3% agarose plates inoculated on the surface by depositing 3 μl of Tryptic Soy Broth (TSB) cultures grown to an optical density at 660 nm (OD_{660}) of 0.3. The plates were incubated in the presence or absence of blue light at the indicated temperatures for 24 hours or else specified. Three independent experiments were performed.

Analyses of gene expression by qRT-PCR.

Retrotranscription and qRT-PCR analysis were done as described in Tuttobene et al., 2019, using primers listed in Table 1. Data are presented as NRQ (Normalized Relative Quantities) calculated by the qBASE method [19], using *recA* and *rpoB* genes as normalizers. The *anbluf65* transcript levels of each sample were normalized to the *rpoB* transcript level for each cDNA sample. Relative gene expression to *rpoB* was calculated using the comparative $2^{-\Delta\text{CT}}$ method [20]. Each cDNA

sample was run in technical triplicate and repeated in at least three independent sets of samples.

t-test was used to determine whether two values were significantly different comparing data within each temperature assayed. *p*-values: *, *p* < 0.01; **, *p* < 0.001.

Table 1. Primers used in this study.

Name	SEQUENCE (5'-3')	REFERENCE
rpoBF_rt	ACTTGCGTGCTGGTGTTCTTT	This study
rpoBR_rt	ACGCGCATGCATCTTGTCATCA	This study
1065F_rt	GACTGTGTTATGCCAGCCAACGAA	This study
1065R_rt	CTCCCATCTTGAGAAAGAAAGCCTCG	This study
98085F_rt	CAATGCGTAGAAGGTCAGAAAG	This study
98085R_rt	GCTGGAATAGTAGAGCTATTTCAGTAA	This study
46F_rt	ACCGTCCGAACAGCGGTTATTT	This study
46R_rt	TGATCCGCTTTCAAATTGGGTTGA	This study
46F	GGACATATGAGTTTAATAGGCTTTATG	This study
46R	GGATCCTTAAACTTGATATTGATCCG	This study

Cloning, synthesis, overexpression and purification of AnBLUFs.

In the case of gene *anbluf46*, a PCR product amplified from *A. nosocomialis* RUH2624 genomic DNA using primers 46F and 46R (Table 1), which contain *Nde*I and *Bam*HI restriction sites, respectively, was cloned into pGEM®T- Easy vector and then subcloned into the *Nde*I and *Bam*HI sites of pET28-TEV. In turn, *anbluf65* and *anbluf85* coding sequences were directly synthesized and subcloned into pET 28 a(+) (Genscript, USA). Thus, the different proteins were overexpressed as N-terminal His-tag fusions. Plasmids were transformed into *E. coli* BL21 (pLys) cells, which were cultured in LB broth supplemented with chloramphenicol and kanamycin at 37 °C until they reached an OD₆₀₀ of 0.6 to 0.7. Overexpression of the His-tagged proteins was induced with 0.5 mM IPTG at 15 °C. After incubation for 5 h at 15 °C to avoid the formation of inclusion bodies, the cells were collected by centrifugation, suspended in lysis buffer (20 mM Tris [pH 8.0], 500 mM NaCl, 1 mM β- mercaptoethanol), and disrupted using liquid nitrogen and a mortar. Cell debris were removed by centrifugation at 20,000 x g for 30 min at 4 °C, and the supernatant was loaded onto a nickel nitrilotriacetic acid (Ni-NTA)-agarose column (Invitrogen). The column was washed

sequentially with lysis buffer containing 20 mM and 40 mM imidazole, respectively, and the His-tagged protein was eluted with the same buffer containing 250 mM imidazole (elution buffer). PD-Minitrap G-25 (GE Healthcare) columns were used to desalt and exchange the buffer to 20 mM Tris (pH 8.0) and 200 mM NaCl (working buffer). Vivaspin 500 (Sartorius) centrifugal filters (cutoff of 10 kDa) were used to concentrate the purified proteins. The purity of the overexpressed His-tagged protein was confirmed by sodium dodecyl sulfate (SDS)-PAGE 16% gels.

Sequence analyses.

Protein sequence alignments were performed using CLUSTALW (<https://www.genome.jp/tools-bin/clustalw>), and the alignments were visualized with Jalview 2 [21]. AnBLUF65 and AnBLUF46 3D modeled structures were performed using the Swiss-Model workspace/GMQE [22] in a search for template mode. In all cases, the best fit corresponded to the BlsA PDB structures (6W6Z for the dark-adapted form and 6W72 for the light-adapted form). Models were visualized and handled with PyMOL (by Schrödinger). Ring 2.0 webserver [23] was used to explore the specific $\pi\pi$ stacking interactions in all protein models as well as other non-covalent interactions.

Spectroscopic measurements.

Absorption spectra were registered using an Ocean Optics modular UV-Vis spectrophotometer USB2000+. The assays were done using a 5x5 mm quartz cuvette (Hellma, Germany), placed in a Quantum Northwest FLASH 300 cuvette holder connected to a Peltier-based temperature controller, containing 250 μ l of air-saturated protein solution in 20 mM TRIS, 200 mM NaCl, pH 8.0 buffer. Scattering effects on the absorption spectra were corrected using a|e – UV-Vis-IR Spectral Software 1.2, FluorTools (www.fluortools.com).

Light-adapted state for AnBLUFs (l-AnBLUF) was obtained by blue light irradiation of the dark-adapted form (d-AnBLUF) with a 1 W Royal Blue LED (Luxeon Star LEDs) emitting at 443 ± 20 nm. Absorbance changes followed at 510 nm (ΔA) vs time (t) were recorded during dark-light cycles at

different temperatures. The light-adapted-state formation time, τ_{BLUF} , and the back-recovery time through thermal recovery to the dark-adapted state, τ_{rec} , were determined using the exponential equations 1 and 2, respectively, where A is the pre-exponential factor and ΔA_0 is the initial absorbance:

$$\Delta A = \Delta A_0 + A e^{\frac{t}{\tau_{BLUF}}} \quad (1)$$

$$\Delta A = \Delta A_0 + A e^{\frac{-t}{\tau_{rec}}} \quad (2)$$

Fluorescence emission spectra were recorded with a Hitachi F-2500 spectrofluorometer equipped with a Hamamatsu R-928 photomultiplier. Neutral density filters (10 %T) were placed onto the excitation output beam to minimize photochemical processes during acquisition. Emission spectra were obtained by selective excitation of FAD cofactor at 460 nm. Temperature control was performed using a circulating fluid bath (Haake F3) connected to the cuvette holder.

Protein extraction.

A. nosocomialis was grown stagnantly in LB at 37 °C under blue light or in the dark for 24 hs, and the procedure was repeated to generate three independent biological replicates. The cells were recovered by centrifugation at 7,000 g for 10 min at 4 °C. The cells were then resuspended in 500 μ l of extraction buffer (25 mM Tris pH 7.0), 0.5% Tween 20, 2 mM EDTA, 5 mM, β -mercaptoethanol) and lysed by sonication while keeping samples in ice. Cell debris and non lysed cells were collected by centrifugation at 12,000 rpm for 15' at 4 °C and the supernatants were carefully decanted into clean tubes and stored at -80 °C. Total protein content present in the supernatant was quantified using bicinchoninic acid (BCA) (Thermo Scientific, Germany). Then, 30 μ g total proteins were loaded in SDS-PAGE (10% stacking gel and 5% running gel), and allowed to separate electrophoretically only 1 cm long within the separation gel. The gel was incubated for at

least 3 hs in fixing solution: 30% v/v ethanol, 2% H₃PO₄, and after being washed with MiliQ water, it was incubated for 1 h in staining solution: 18% v/v methanol, 17% p/v (NH₄)₂SO₄ y 2% H₃PO₄ under vigorous shaking. Then, Coomassie G250 powder (0.5 g/L) was added and further incubated for 1 or 2 days until stained proteins were visible, which were then cut from the gel and sent to the Proteomics Core Facility of CEQUIBIEM at the University of Buenos Aires where protein digestion and Mass Spectrometry analysis were performed. Samples were resuspended in 50 mM (NH₄)HCO₃ pH 8.0, digested overnight with sequencing-grade modified trypsin (Promega) and desalted with Zip-Tip C18 (Merck Millipore). Proteins were analyzed by nanoHPLC (EASY-nLC 1000, Thermo Scientific, Germany) coupled to a mass spectrometer with Orbitrap technology (Q-Exactive with High Collision Dissociation cell and Orbitrap analyzer, Thermo Scientific, Germany). Peptide Ionization was performed by electrospray. Data were analyzed with Proteome Discoverer 2.1 software (Thermo Scientific, Germany) for identification and area quantitation of each protein. Protein identification was performed using *Acinetobacter nosocomialis* strain Ab1 protein collection as reference (UP000244598; <https://www.uniprot.org/uniprot/?query=proteome:UP000244598>) since a fully annotated *A. nosocomialis* RUH2624 proteome is not currently available. Missing value imputation method [24] was applied to the analyzed results, and Perseus software v1.6.1.3 (Max Planck Institute of Biochemistry) was used to perform the statistical tests. Proteins showing a fold change (FC) ≥ 1.5 between light and dark conditions and *p*-value < 0.05 were considered differentially produced.

Proteomics Bioinformatics.

Biological functional analyses of differentially over-represented proteins detected by proteomic approaches were categorized according to their molecular function, biological process, and cellular component, by using the Blast2GO tool.

Results and Discussion

Light modulates motility in a wide range of temperatures in *A. nosocomialis* RUH2624.

We have recently shown that in *A. baumannii* ATCC 17978, light modulates motility from low to moderate temperatures up to 24 °C [11]. In this work, we assayed the effect of light on motility at different temperatures in *A. nosocomialis* RUH2624. Our results show that this microorganism grows at the inoculation point under blue light at 25 °C, and despite of moving increasingly as temperature raises, it only reaches 10-20 % of plate coverage at the highest temperatures such as 32 and 37 °C (Figs 1A and 1B). On the contrary, the bacteria moved covering the whole plates at all temperatures assayed in the dark (Figs 1A and 1B). Thus, in contrast to *A. baumannii*, in *A. nosocomialis* RUH2624 light modulates motility not only at environmental temperatures, but within a wide range of temperatures including those found in warm-blooded animals such as 37 °C.

Fig 1. Effects of blue light and temperature on *A. nosocomialis* motility and *anbluf65* expression. (A) Cells of RUH2624 strain were inoculated on the surface of motility plates and incubated at the indicated temperatures. Plates were inspected and photographed after incubation in darkness (D) or in the presence of blue light (L) at the indicated temperatures. (B) Quantification of cell motility estimated as the percentage of plate coverage, i.e., the percentage of the Petri plate area covered with bacteria, in motility plates inoculated with RUH2624 wild type and incubated at the indicated temperatures. Three independent experiments were performed in each case. The area of plates covered with bacteria was measured with ImageJ (NIH), and then the percentage of plate coverage was calculated. The mean and standard deviation is informed. Different letters indicate significant differences as determined by ANOVA followed by Tukey's multiple comparison test ($p < 0.05$). For those conditions at which the bacteria just reached the edge of the plate a value of 100% is informed. (C). cDNA from *A. nosocomialis* RUH2624 cells grown in motility plates in

the presence of blue light (L) or in darkness (D) at different temperatures was used as template for quantitative real-time PCR using *anBLUF65* specific primers. Transcription of *rpoB* was used as a constitutively expressed internal control. The results are representative of three independent experiments. The mean and standard deviation are shown. *t*-test was used to determine whether two values were significantly different comparing data within each temperature assayed. *p*-values: *, *p* < 0.01; **, *p* < 0.001.

***A. nosocomialis* RUH2624 encodes three BLUF-type putative photoreceptors.**

Comparative sequence analyses have shown the presence of three BLUF domain-containing proteins encoded in the *A. nosocomialis* RUH2624 genome [8]. These putative photoreceptors received accession numbers EEW98085, EEX00046, and EEW01065 in the Genbank database and will be further referred to here as AnBLUF46, AnBLUF65 and AnBLUF85, respectively .

Sequence alignments of these three BLUF-domains containing proteins with other well characterized BLUF domains such as Slr1694 from *Synechocystis* sp., Tll0078 from *Thermosynechococcus elongatus*, AppA and BlrB from *Rhodobacter sphaeroides*, and BlsA from *A. baumannii*, confirmed the presence of 22 highly conserved residues in the N-terminus of canonical BLUF domains according to pfam04940 (Fig 2A). Several BLUF domains belonging to different species from the genus *Acinetobacter* have been described so far as functional blue light photoreceptors [10-12, 15]. Alignment of these functional *Acinetobacter* BLUF-photoreceptor sequences show a higher level of conserved residues in their N-terminus domain (33 residues, Fig 2B), while the analysis of all γ -*Proteobacteria*, including 30 *Acinetobacter* sequences deposited in pfam04940 displays lower conservation (14 residues, Fig 2C) in their primary structure and in some cases, insertions between the conserved Tyr and Gln (Fig 2A), though with a low occupancy level (Fig 2B).

Fig 2. Comparative analysis of BLUF domains (A). Multiple alignment of structurally characterized BLUF domains members of pfam04940 generated with ClustalW and adjusted with Jalview. Residues are colored according to the percentage of identity and the conservation level (darker, highly conserved). Asterisks show residues with high or very high conservation. (B) Alignment logo of *Acinetobacter* BLUFs directly or indirectly characterized as photoactives, i.e. BlsA (*A. baumannii*); AnBLUF65 (156 amino acids), AnBLUF 46 (147 amino acids) and AnBLUF85 (150 amino acids); Q7BC36, Q6FAI1 and Q6FAH9 (*A. baylyi* ADP1).(C) Alignment logo performed with 341 BLUF sequences from 204 γ - *Proteobacteria* species, obtained from PF04940 (Pfam). Asterisks show residues with high or very high conservation interacting with flavin (red) or not (black).

Phylogenetic analyses indicate that AnBLUF65 and *A. baumannii*'s BlsA BLUF domains are grouped in a monophyletic cluster [8], thus indicating that they are orthologs. *anbluf65*'s genomic context shares many features with *blsA*'s (Fig 3). *anbluf65*'s 5' upstream region is highly conserved with respect to the 5' upstream region of BlsA in *A. baumannii* ATCC 17978 (Fig 3), as well as with different strains of *A. nosocomialis* (S1 Fig). Interestingly, the presence of insertion sequence ISAb27 was detected in the intergenic region upstream of the Bof coding sequence in *A. nosocomialis* RUH2624 (Fig 3A and S1 Fig). Whether ISAb27 is providing a hybrid promoter or disrupting the transcription of a putative operon that includes *anbluf65* is a possibility that deserves further experimentation. *A. nosocomialis* RUH2624 *anbluf65* 3' downstream region is more variable compared both with BlsA or with other strains of *A. nosocomialis* (Fig 3 and S1 Fig). Gene *anbluf46* is located in a different genome location, surrounded by genes coding for a putative succinyl-CoA:3-ketoacid-coenzyme A transferase, a LysR transcriptional regulator, a hypothetical protein, a putative heat shock, a conjugal protein, and another hypothetical protein on the 5' upstream region (Fig 3A). On the 3' downstream region there are encoded: a putative protein with a metallo- β -lactamase fold, an H⁺/gluconate symporter, a hydroxybutyrate dehydrogenase, an AraC transcriptional regulator, and an aromatic acid/H⁺ symporter (Fig 3A).

261 *anbluf46* does not seem to be part of an operon, as both flanking genes are encoded in the
 262 opposite direction (Fig 3A). Comparing *anbluf46* genomic context in strain RUH2624 with other
 263 strains of the same species, it is observed that the 3' downstream region is highly conserved while
 264 the 5' upstream region is variable, showing the presence of transposases in different locations (S2
 265 Fig). *anbluf85* was only found encoded in *A. nosocomialis* RUH2624 and UBA873 strains (100%
 266 identity). In RUH2624, *anbluf85* is flanked by a 5' upstream region coding for an acetyltransferase
 267 and a putative conjugal transfer pilus assembly protein TraB. On the 3' downstream region there
 268 are encoded a hypothetical protein, a FAM199X domain-containing protein, and a relaxase
 269 MobA/MobL (Fig 3A). RUH2624 *anbluf85* genomic context is different from that in UBA873 (S3
 270 Fig). In the latter, only the 3' downstream region is shown as the contigs are not assembled yet (S3
 271 Fig). Sequence comparisons indicate the presence of an *anbluf85* homolog in *Acinetobacter lwoffii*
 272 strain M2a plasmid pAVAc116 as well as in *Acinetobacter* sp. ACNIH1 plasmid pACI-148e with
 273 100% identity at nucleotide level. Interestingly, also an *anbluf85* homolog with 98.13% identity at
 274 the nucleotide level was reported to be present in *Acinetobacter baumannii* isolate KAR [25].
 275 Moreover, 100% identity homologs at the aminoacidic level are also present in *A. baumannii*
 276 strains A30, A10, and MSHR_A189, two in *A. radioresistens* strains 50v1 and TG29429; and in *A.*
 277 *baumannii* strains MRSN7353 and ARLG1306 showing 1 aminoacidic difference with AnBLUF85.
 278 Thus *anbluf85* is not widely distributed but its presence is observed discretely in some
 279 *Acinetobacter* species, and the possibility of horizontal gene transfer is suggested.

280 **Fig 3. Genomic context of the three BLUF photoreceptors present in *A. nosocomialis* and comparison with**
 281 **BlsA of *A. baumannii*.** (A). AnBLUF46, AnBLUF85 and AnBLUF65 genetic surroundings in *A. nosocomialis*
 282 RUH2624 strain. (B). BlsA genetic surroundings in *A. baumannii* strain ATCC 17978. Coding-sequences (CDSs)
 283 are located in their corresponding frame. The different photoreceptors are indicated as pink arrows.
 284 Different colors in CDSs indicate different functions. Gene annotations are indicated as numbers above the
 285 schemes, with the following codes: 1-DUF2171; 2-Acyl-CoA dehydrogenase; 3-GlcNAc-PI de-N-acetylase; 4-

286 Methyltransferase domain; 5-Glycosyltransferase; 6-BOF- Class 2b aminoacyl-tRNA synthetases--NirD/
 287 YgiW/YdeI family stress tolerance protein; 7-HP; 8-RhtB - Homoserine/Threonine/Homoserine Lactona
 288 translocator; 9-AraC transcriptional regulator; 10- Poly(R)-hydroxyalkanoic acid synthase; 11-GltS-
 289 Sodium/glutamate symporter; 12- DUF815; 13-FdaR- Fatty acid transcriptional regulator; 14-DDE
 290 endonuclease domain, putative transposase; 15-Helix-turn-helix of DDE superfamily endonuclease; 16-
 291 Proton antiporter-2 (CPA2) family; 17-NDAB-Rossmann Superfamily- Oxidoreductase; 18-DoxX-like family;
 292 19-Succinyl-CoA:3-ketoacid-coenzyme A transferase subunit B; 20-Succinyl-CoA:3-ketoacid-coenzyme A
 293 transferase subunit A; 21-LysR family transcriptional regulator; 22- HP; 23-DUF3298-Putative heat-shock
 294 protein; 24-DUF2846-Putative conjugal domain; 25-HP; 26-MBL fold metallo-hydrolase; 27-H+/gluconate
 295 symporter; 28-3-Hydroxybutyrate dehydrogenase; 29-AraC family transcriptional regulator-RmlC-cupin
 296 protein; 30-Aromatic acid:H⁺ symporter (AAHS); 31-Conjugal transfer pilus assembly protein TraB; 32-N-
 297 Acetyltransferase; 33-HP; 34-FAM199X; 35-Relaxase-MobA/MobL family protein. HP: hypothetical protein.
 298

299 **Only AnBLUF65 and AnBLUF46 are stable photoreceptors.**

300 We evaluated next whether the three BLUF-domain containing genes encode active
 301 photoreceptors. For this purpose, genes *anbluf65*, *anbluf46* and *anbluf85* were recombinantly
 302 expressed and purified to assess their functionality.

303 Fig 4 shows the UV-vis spectra of AnBLUF65 (92% identity with BlsA) and AnBLUF46 (45% identity
 304 with BlsA) at 15 °C in working buffer. Before blue light irradiation, both dark-adapted states of
 305 AnBLUF65 (d-AnBLUF65) and AnBLUF46 (d-AnBLUF46) showed the typical absorption band of the
 306 fully oxidized flavin corresponding to the transition $S_0 \rightarrow S_1$ with $\lambda_{\max} \approx 460$ nm and shoulder
 307 approximately at 483 nm. Upon blue-LED illumination, the band is red-shifted, characteristic of the
 308 formation of the light-adapted state of BLUF proteins (l-BLUF) [26], with absorption maximum
 309 wavelength shift of 8 and 12 nm for l-AnBLUF65 and l-AnBLUF46, respectively (Fig 4A and C). The
 310 differential absorption spectrum (ΔA) between the light- and dark-adapted states of AnBLUF65

showed positive changes at 508 and 472 nm, and negative changes at 455 and 427 nm. AnBLUF46 also showed positive changes at 511 and 476 nm, whereas negative bands were less defined (insets Fig 4A and C). The overall results indicate that AnBLUF65 and AnBLUF46 are active photoreceptors. Gene *anbluf85* was weakly expressed and the partially purified product was barely stable precipitating at T higher than 12 °C, with a photocycle presenting a ≈ 7 nm red shift (S4 Fig).

Fig 4. Normalized steady-state visible absorption-emission spectra changes of photoreceptors by blue light illumination, in working buffer solution at 15 °C. (A and B). dAnBLUF65 (black line) and lAnBLUF 65 (blue line). (C and D). dAnBLUF46 (black line) and lAnBLUF46 (red line). Inset: differential absorption spectra (lAnBLUF-dAnBLUF).

Fluorescence emission spectra obtained by excitation at 460 nm at 15 °C for d-AnBLUF65 and d-AnBLUF46 also depicted a BLUF-like behavior, with an emission maximum at 512 and 519 nm, respectively (Fig 4B and D). The fluorescence bands of both proteins were blue-shifted compared with free FAD in the same working buffer [10, 11] confirming the existence of specific interactions between the isoalloxazine group of the flavin with the surrounding residues, and reducing solvent accessibility within the flavin binding pocket [27]. When the fluorescence spectrum of the light-adapted and the dark-adapted forms of both photoreceptors were compared, they exhibited the typical red shift of approximately 5 nm of the fluorescence maximum in addition to a loss of vibratory structure and a lower emission intensity, suggesting a more relaxed environment around the cofactor [28-30] [10].

Only *anbluf1065* is expressed in *A. nosocomialis* RUH2624.

The expression levels of the three BLUF-type photoreceptors encoded in *A. nosocomialis* RUH2624 were analyzed on cells recovered from motility plates incubated at a wide range of temperatures (25–35 °C) under blue light or in the dark by qRT-PCR as described previously [14]. We were able

to detect expression of *anbluf1065* only, the *blsA* ortholog in *A. nosocomialis* [8]. *anbluf65* was expressed at 25, 30, and 35 °C, with approximately 9, 8, and 7.5-fold higher expression in the dark than under blue light (Fig 1C); a response previously observed for BlsA [11, 15]. Moreover, expression levels were higher at lower temperatures (Fig 1C). Thus, our results suggest that expression of the putative *anbluf65* photoreceptor correlates with regulation of motility by light.

AnBLUF65's photoreceptor activity is operative in a temperature range that correlates with regulation of motility by light.

A. nosocomialis modulates motility in response to blue light in a similar fashion as observed for *A. baumannii*, but in a wider range of temperatures. Table 2 summarizes the temperature dependency of visible absorption and emission fluorescent properties of flavin in purified AnBLUF65 and AnBLUF46 photoreceptors. Absorption maxima of the dark-adapted forms of both proteins were blue-shifted when temperature increased, reaching in the case of d-AnBLUF46 the value of free FAD in solution (≈ 450 nm). The ratio between the shoulder absorbance at ≈ 480 nm and the absorption maximum (A_{sh}/A_{max}) diminished with temperature increments, as did the redshift upon blue light illumination, suggesting a distancing between the conserved Tyr to the isoalloxazine ring [31]. Fluorescence emission properties of the cofactor were significantly affected upon formation of l-AnBLUF65 at 24 and 37 °C, with a redshift in response to blue light, as described before at 15 °C or when temperature increased, implying a relaxation in the hydrogen bond network surrounding the isoalloxazine, and/or changes in the polarity in the chromophore cavity as a consequence of increased solvent accessibility. Widening of the emission spectrum full-width half-maximum (FWHM) in response to the formation of the light-adapted state for AnBLUF65 was observed at 24 and 37 °C, and this effect was also evident when temperature increased. Overall, these results indicate that the flavin environment in AnBLUF65 is affected by temperature, as it has been shown for BlsA [10, 11]. In contrast, emission maximum wavelength

and FWHM for the dark- or the light-adapted states of AnBLUF46 did not show significant changes neither in response to blue light at 24 and 32 °C, nor to temperature increments. In fact, d- and l-AnBLUF46 isoalloxazine pockets appeared to be more polar than the same states of AnBLUF65. Chromophore environments in AnBLUF65 and AnBLUF46 were somewhat dissimilar and appeared to react in a different way to the formation of the light-adapted state. This fact goes in good agreement with structural predictions from models of these two proteins generated using the crystallographic structure of dark- and light-adapted states of BlsA (PDBs 6W6Z and 6W72, respectively) as templates (Fig 5). Fig 5A and 5B show the location of water molecules (blue dots) for the dark-adapted form of BlsA (d-BlsA) and the light-adapted (l-BlsA) respectively, determined by Chitrakar *et al.* [32]. The flavin pocket access site has more affinity and/or accessibility to water molecules in the light-adapted form, supporting our previous characterization of chromophore behavior inside BlsA cavity by the flavin fluorescence emission (Fig 5A and B)[10]. Modeled structures of AnBLUF65 (Fig 5C and 5D), which is BlsA ortholog as described before, show that the environment of the cofactor is conserved compared with BlsA. This means that analogously to BlsA, the flavin pocket in the light-adapted form of AnBLUF65 is more solvent-accessible. In contrast, the solvent accessibility to the chromophore cavity between d- and l-AnBLUF46 is indistinguishable (Fig 5E and F). In addition, the flavin binding site in AnBLUF46 seems more solvent-accessible than in BlsA/AnBLUF65. Hence, these data stress out how temperature and blue light differentially affect them.

Table 2. Absorption-emission properties of AnBLUF46 and AnBLUF65 at different temperatures.

	T (°C)	Absorption			Emission			
		$\lambda_{\max}$	$A_{\text{sh}}/A_{\max}$	ΔA	$\lambda_{\max}$	d-AnBLUF	$\lambda_{\max}$	l-AnBLUF
		d-AnBLUF (nm)	d-AnBLUF	(a.u)	d-AnBLUF (nm)	FWHM (nm)	lBLUF (nm)	FWHM (nm)
AnBLUF65	15	460	0.86	8	512	64	517	74

	24	459	0.87	6	514	65	519	74
	37	457	0.79	3	520	74	523	76
AnBLUF46	15	459	0.81	13	519	68	522	68
	24	457	0.81	13	523	75	523	75
	32	450	0.72	8	523	75	523	76

380 Maximum absorption wavelength ($\lambda_{\max}$), shoulder to maximum absorbance ratio ($A_{\text{sh}}/A_{\max}$), redshift
381 maximum upon BL illumination (ΔA), maximum emission wavelength ($\lambda_{\max}$) and full-width at half maximum
382 (FWHM) for dark and light-adapted proteins.

383 **Fig 5. Surface representation of *Acinetobacter* BLUFs and FMN (yellow sticks) into the site access pocket.**

384 Internal surface associated with the cavity is also depicted. (A) BlsA dark-adapted, PDB 6W6Z; (B) BlsA light-
385 adapted, PDB 6W72, (C) d-AnBLUF65; (D) I-AnBLUF65; (E) d-AnBLUF46 and (F) I-AnBLUF46.- Models were
386 generated with the Swiss-Model server, using PDB 6W6Z as template for figure C and E and PDB 6W72 for
387 figure D and F. Water molecules are depicted as blue dots.

388

389 Insets in Fig 4 show difference spectra both for AnBLUF65 and AnBLUF46 (IAnBLUF-dAnBLUF). In
390 previous work, we have used the ΔA at 510 nm at different temperatures to follow the kinetics of
391 formation of the light-adapted state of each protein, and in darkness, the recovery time back to
392 the dark-adapted state, through cycles of blue light illumination. This kinetic profile was used to
393 determine the quantum yield of the protein light-adapted formation, Φ_{IAnBLUF} . Fig 6 presents the
394 kinetic profiles of AnBLUF65 (Figures 6A and 6B) and AnBLUF46 (Fig 6C and 6D) tested at two
395 temperatures. For AnBLUF65 the kinetic curves were monitored at 24 °C and 37 °C, which
396 correspond to environmental and warm-blooded host temperatures, respectively. However,
397 AnBLUF46 was tested at 24 °C and 32 °C, the maximum temperature at which this photoreceptor
398 presented activity. Table 3 summarizes the light-adapted-state formation time τ_{IAnBLUF} and the
399 recovery time back to the dark-adapted state τ_{rec} for both BLUFs as a function of temperature
400 along with the photoactivation quantum yield of light-adapted state, Φ_{IAnBLUF} .

Table 3. Kinetic parameters observed for AnBLUF65, AnBLUF46, and BlsA at 24 °C and 37 °C.

	Temperature (°C)	τ_{I-BLUF} (s)	Φ_{I-BLUF}	τ_{rec} (s)
AnBLUF65	24	106±8	0.36±0.04	233±8
	37	80±15	0.12±0.03	240±60
AnBLUF46	24	70±4	0.11±0.01	950±90
	32	40±5	0.08±0.02	115±20
	37	ND	ND	ND
BlsA*	24	70±2	0.17±0.01	111±10
	26	28±9	0.11±0.02	40±9
	37	ND	ND	ND

ND: not detectable. * [11]

Fig 6. Kinetic profile of absorbance changes at 510 nm. (A and B) AnBLUF65 at 24 and 37 °C. (C and D)

AnBLUF46 at 24 and 32 °C. Dark adapted protein, dAnBLUF was illuminated using blue light LED at 443 ± 20 nm (blue arrows). After maximal conversion to IAnBLUF, blue light was turned off (black arrow) and protein back to dAnBLUF.

Temperature increments had a significant effect on the $\tau_{IAnBLUF}$ for both proteins, with reductions of 25 and 40% for $\tau_{IAnBLUF65}$ and $\tau_{IAnBLUF46}$, respectively. AnBLUF46 had similar values to BlsA at 24 °C (Table 3), however, $\tau_{IAnBLUF65}$ was larger than that of BlsA. Analyses of τ_{rec} show that AnBLUF46 takes 15 mins to recover, nine times slower than BlsA; while AnBLUF65 was only two-fold slower than the *A. baumannii* protein. As expected for a thermal process, $\tau_{recAnBLUF46}$ was nine times faster when temperature increased to 32 °C, becoming similar to the value of $\tau_{recBlsA}$ at 24 °C. However, this was not the case of AnBLUF65, whose $\tau_{recAnBLUF65}$ remained constant at 24 and 37 °C. This behavior prompts us to speculate whether this recovery time might have some physiological meaning since AnBLUF65 presents conformational changes at least in its binding cavity upon illumination, but the time to return to the dark form is conserved.

Despite the high sequence similarity between AnBLUF65 and BlsA (92%), the behavior of the former was significantly different compared with that of the latter, regarding the efficiency to respond to blue light generating the light-adapted form. In fact, Φ_{AnBLUF65} was more than two-fold higher than that of BlsA at 24 °C, which can be interpreted as AnBLUF65 being more efficient to form the light-adapted state. The τ_{AnBLUF65} was also significantly higher than that for BlsA, although to a lesser extent, and the $\tau_{\text{recAnBLUF65}}$ was two-fold higher than τ_{recBlsA} , probably because the adapted form is more stabilized in AnBLUF65 than in BlsA. Another contrasting effect between such similar proteins is that AnBLUF65 is less prone to aggregate upon temperature increments than BlsA, which shows this effect macroscopically above 30 °C, while AnBLUF65 does not aggregate at 37 °C showing only incipient turbidity in the buffer solution. While Φ_{BlsA} reaches non-detectable values at T higher than 28 °C, Φ_{AnBLUF65} at 37 °C is 0.12 ± 0.03 . In contrast, Φ_{AnBLUF46} was not significantly impacted between 24 °C and 32 °C. Nevertheless, AnBLUF46 lost all photoactivity at T>32 °C. These results suggest that AnBLUF65 is more efficient to form the light-adapted state than AnBLUF46 at 24 °C.

Intramolecular interactions and their role in intra-protein signaling.

So far, we have analyzed these photoreceptors describing intrinsic phenomena occurring mostly in the N-terminal side of the protein, where the flavin resides in its cavity. These data have profiled AnBLUF65 and AnBLUF46 in the first part of the photo-reception process. Next, we aimed to expand our analyses to cover the subtle structural rearrangements between the dark and the light-adapted states of the protein. Intra-protein changes in hydrophobic and aromatic residue interactions driven upon illumination have been suggested by Chitrakar et al, 2020 to explain how the photo-signal might be translated from the flavin surroundings to the variable domain. Based on the similarity between BlsA and AnBLUF65, we modeled structures for both *A. nosocomialis*

proteins, AnBLUF65 and AnBLUF46 (Fig 5), using BlsA as template, which is appropriate since BLUF's in *Acinetobacter* genus share high sequence similarity. We used RING 2.0 [23] to profile all intramolecular interactions, such as π - π stacking, H-bonding, Van der Waals (VDW) interactions, and salt bridges, in AnBLUF65 and AnBLUF46, as presented in S1 Table. The analysis of π - π stacking interactions amongst aromatic residues helped us to profile aromatic networks for both proteins, in both dark- and light-adapted forms.

The first aromatic cluster found surrounds the flavin cofactor and involves the hydrophobic side of the isoalloxazine ring (ring I), the conserved Tyr, a Phe and a His for AnBLUF65 as shown in Fig 7. This His belongs to the already described motif Asp-X-Arg-His and it has been suggested that the protonation/deprotonation of the conserved Gln via His is energetically favourable [33]. In AnBLUF65, His73 forms π - π stacks with Tyr7 and with Phe49 in addition to other interactions i.e. H-bonds that have been described before. After blue light illumination this cluster does not show a significant impact on its C α main chain or in the angle of the aromatics involved. This might be necessary to conserve the hydrophobicity in the flavin pocket [34].

Fig 7. Residues involved in π - π stacking interactions in AnBLUF65 dark-(orange) and light-adapted (blue) state.

The next aromatic network found is in the variable domain (C-terminal) and contains two aromatic tetrads, that are formed by six Phe in AnBLUF65 (Fig 7). Upon blue light illumination, distances between aromatic centroids in these tetrads are shortened in all cases in AnBLUF65. Phe106 loses its π - π stack interaction with Phe115, being left only with VDW interactions. Thus, the formation of the light adapted state of AnBLUF65 requires a local re-arrangement of α -helices in the C-terminal and the aromatic network in this area is strengthened.

A Tyr-Tyr motif, highly conserved in *Acinetobacter* BLUFs (Fig 2B) also participates in the aromatic networks described before. AnBLUF65 has Tyr44 interacting with Phe49 on the chromophore side

of the protein, while Tyr43 interacts through π - π stacking with Phe128 on the other side of the β -sheet, towards the variable domain (Fig 7). The notable re-arrangement is found towards the variable domain may be possible because an Asp residue located in the loop between the Tyr-Tyr motif and the first Phe-Phe motif, rotates upon blue light illumination, losing its contact with Phe115. Thus, one way to explain these findings is that the aromatic clusters found in the variable domain might internally stabilize the structure of this protein upon the formation of the light-adapted state. In contrast, the aromatic cluster in the flavin/N-terminal side of these proteins does not suffer any alteration when changing states. Taken together, it suggests that intramolecular hydrophobic/aromatic interactions might have a role in the elasticity of the protein monomer to go back and forth between the two states.

Quantitative differential proteomic profiling in response to light in

***A. nosocomialis* RUH2624 at 37 °C.**

In previous studies, we demonstrated that light can modulate important physiological traits related to bacterial physiology and virulence in *A. baumannii*, *S. aureus*, *P. aeruginosa* and *A. nosocomialis* at temperatures found in warm-blooded hosts [8, 9, 14, 15, 17, 18, 35]. To broaden our knowledge of the response to light in *A. nosocomialis* at 37 °C, a quantitative comparative proteomic analysis was conducted to obtain an overall representation of the total cellular changes that occur in RUH2624 cells at the protein level upon blue light illumination at the mentioned temperature. We aimed to use proteomics as a tool to identify those proteins that could be present in different amounts under light and dark conditions and how these proteins could affect *A. nosocomialis*' physiology.

In total, 38 proteins were over-represented (fold change (FC) > 1.5 and *p-value* < 0.05, see Tables 4 and 5) when cells were grown under blue light or dark state at 37 °C (Fig 8A). Twenty proteins were accumulated in higher amounts in cells cultured under blue light with respect to darkness

(Fig 8A and B), while eighteen proteins were found to be in a greater amount under dark state (Fig 8A and C). When categorized using BLAST2GO suite, these proteins were dispersed across a wide variety of functions, according to gene ontology terms (Fig 9). In the “Biological Process” and “Molecular Function” sections the majority of the over-represented proteins fell into the metabolic/cellular process and catalytic activity categories, respectively, indicating changes in the metabolism of this bacterium in response to light. Considering “Cellular Component”, over-represented proteins under light conditions appear mainly distributed in the plasma and outer membranes, while in darkness most of them are cytoplasmatic (Tables 4 and 5).

Table 4. Proteins with increased abundance in the light compared to dark at 37 °C.

Accession Number	Description	p-value	FC
FA0A2T7FSK9	Probable allantoicase	2.82E-05	62.0
K9B4X7	Thiol disulfide reductase thioredoxin	4.79E-02	32.1
A0A2T7FKW8	TonB-dependent receptor	1.29E-02	27.4
A0A2T7FJ60	Glutathione S-transferase	1.48E-02	24.0
A0A0A7XMP5	DUF4442 domain-containing protein	3.98E-04	18.1
A0A2T7FKT5	Aldehyde dehydrogenase	2.69E-05	10.5
A0A2T7FHA5	Peptidylprolyl isomerase (Fragment)	2.95E-02	10.3
A0A0A7XME6	Biopolymer transporter ExbD OS=Acinetobacter nosocomialis	1.15E-05	9.3
A0A2T7FQW1	NAD(P)/FAD-dependent oxidoreductase	4.68E-02	4.8
K9C8I9	Methyltransferase	2.63E-03	4.1
A0A2T7FR20	Queuine tRNA-ribosyltransferase	9.55E-03	2.8
A0A2T7FM31	Acetyl-CoA C-acetyltransferase	9.33E-03	1.9
K9BUJ6	1-acyl-sn-glycerol-3-phosphate acyltransferase	4.07E-02	1.7
A0A0Q1LRF3	DnaA regulatory inactivator Hda	1.48E-02	1.6
K9BNN6	3-isopropylmalate dehydratase small subunit	4.37E-03	1.6
A0A2T7FS76	Inorganic triphosphatase	1.74E-02	1.6
A0A0R0WRI3	Heme-binding protein	3.02E-02	1.6
A0A2T7FQL7	Phosphoribosylamine--glycine ligase	4.90E-02	1.5
A0A2T7FKA3	Gamma-glutamyl phosphate reductase	3.09E-02	1.5
A0A2T7FKU3	Ribosomal RNA large subunit methyltransferase K/L	4.37E-02	1.5

500

Table 5. Proteins with increased abundance in dark compared to light at 37 °C

Accession Number	Description	p-value	FC
A0A2T7FRV8	UDP-N-acetylglucosamine 2-epimerase (Non-	2.75E-02	1.7

	hydrolyzing)		
A0A2T7FJB8	Phenylacetate-CoA oxygenase/reductase subunit PaaK	3.80E-02	1.7
A0A2T7FI74	S-(hydroxymethyl)glutathione dehydrogenase	2.34E-02	1.8
A0A0R1BYG0	50S ribosomal protein L30	5.37E-03	1.9
A0A2T7FPU2	Imidazolonepropionase	1.12E-02	2.5
A0A2T7FP64	Uncharacterized protein	1.02E-02	3.0
A0A2T7FQH8	3-hydroxy-2-methylbutyryl-CoA dehydrogenase	3.24E-02	3.1
K9AWZ8	Endopeptidase La	6.61E-03	6.1
A0A2T7FKJ2	Ribonuclease H	1.15E-03	8.7
A0A2T7FJC4	Ligand-gated channel protein	3.02E-02	10.0
A0A2T7FPJ6	ABC transporter ATP-binding protein	7.59E-05	15.4
A0A2L1VLW9	Probable potassium transport system protein kup	3.24E-02	21.3
A0A2T7FHT8	DNA-directed DNA polymerase	7.24E-03	21.4
A0A2T7FRB2	CoA transferase	3.55E-03	23.4
K9BAG9	Chromosome partitioning protein ParA	3.47E-03	31.0
A0A0A7XIC2	DUF541 domain-containing protein	1.70E-05	47.0
K9ATF3	Peptidase S41	1.12E-02	49.3
A0A2T7FS51	Aminopeptidase	9.12E-14	77.9

Fig 8. Comparison of over-represented proteins detected by nanoHPLC under light and dark conditions for *A. nosocomialis* RUH2624. (A) Scatter plot. (B) Heatmap of proteins produced in higher amounts under blue light. (C) Heatmap of proteins produced in higher amounts under dark condition.

Fig 9. Gene ontology categories after Blast2Go analysis of all differential produced proteins. L and D correspond to over-represented proteins under light and dark conditions, respectively, for *A. nosocomialis* RUH2624.

Proteins with increased abundance in light compared to darkness.

Light regulates the synthesis of proteins involved in a wide range of cellular functions. As mentioned before, these differentially over-represented proteins include some playing roles in metabolism, stress responses, and virulence in different bacterial pathogens.

Two proteins, PurD and Alc, related to purine metabolism were over-represented ($FC_{PurD} = 1.50$, $FC_{Alc} = 63.0$) in the presence of blue light. PurD is a phosphoribosylamine-glycine ligase and participates in the synthesis of inosine 5'-monophosphate (IMP) [36]. Purine metabolism plays an important role in microorganisms, recycling carbon and nitrogen compounds and regulating

metabolism [37]. Several previous reports have described the importance of nucleotide biosynthesis by bacteria triggering infections. For instance, certain auxotroph mutants of *Salmonella*, *S. aureus*, or *S. pneumoniae* were avirulent in murine infection models [38, 39]. Besides, other studies demonstrate that limiting amounts of nucleotide bases in human serum, force pathogens to rely on *de novo* nucleotide biosynthesis [40].

Alc, which encodes an allantoicase, is involved in (S)-allantoin degradation and catalyzes the conversion of allantoate to (S)-ureidoglycolate and urea. Previous studies have found that nitrogen controls several pathways involved in secondary metabolism [41]. In general, secondary metabolites are produced by the cells as a response to environmental cues. For example, a notable decrease in antibiotic production was observed in *S. coelicolor* M145 as a response to the excess of intracellular ammonium generated during degradation of allantoin when the microorganism was grown in allantoin as the sole nitrogen source [42].

Regarding proteins involved in amino acids biosynthesis, LeuD and ProA are present in higher amounts in the presence of blue light ($FC_{\text{LeuD}} = 1.62$, $FC_{\text{ProA}} = 1.50$). LeuD catalyzes the isomerization between 2-isopropylmalate and 3-isopropylmalate, and is part of L-leucine biosynthesis [43].

ProA (gamma-glutamyl phosphate reductase protein) participates in step 2 of the subpathway that synthesizes L-glutamate-5-semialdehyde from L-glutamate and is involved in L-proline biosynthesis [44]. There are shreds of evidence that proline metabolism has complex roles in a variety of biological processes, including cell signaling, stress protection, and energy production [45]. Proline can also contribute to the pathogenesis of various disease-causing organisms. Besides, some organisms use proline directly for the biosynthesis of secondary metabolites with antibacterial or antifungal properties [46].

An interesting protein that is over-represented in the light state is SurA, a chaperone involved in the correct folding and assembly of outer membrane proteins (OMPs), and may act both in early periplasmic as well as in late outer membrane-associated steps of protein maturation [47]. A great

variety of outer membrane proteins are porins and autotransporters that facilitate transport and other essential functions, and act as virulence factors [48]. In *P. aeruginosa*, a significantly lower amount of many porins were detected in the outer membrane (OM) of the conditional *surA* mutant, including members of the OprD family (OprO, OprN, OprP, and OprD) [49]. Also, it was found that siderophore receptors were absent or less abundant in the OM upon depletion of SurA [49]. Related to this point, in the present work a TonB-dependent receptor and a biopolymer transporter ExbD, which are part of a complex that energizes specific high-affinity receptors involved in regulating iron uptake, were found in a greater amount under blue light ($FC_{\text{TonB}} = 27.35$, $FC_{\text{ExbD}} = 9.27$). Consistently, a more robust growth under iron-deprived conditions, i.e., in the presence of the iron chelator 2,2'-dipyridyl (DIP), was observed in *A. nosocomialis* under blue light at 37 °C while it was severely affected in the dark [9]. Therefore, the higher abundance of SurA, TonB-dependent receptor, and ExbD under blue light could contribute to the enhanced iron acquisition observed in this condition, and this is an important feature since iron is a pathogenicity determinant essential for the success of bacterial infections [50].

Other proteins such as queuine tRNA-ribosyltransferase (Tgt) and ribosomal RNA large subunit methyltransferase K/L (RlmL) showed an increase in abundance under blue light. Tgt catalyzes the exchange of a guanine 34 with the queuine precursor 7-aminomethyl-7-deazaguanine (PreQ1) in specific tRNAs containing anticodones G(guanine)-U(uracil)-N (tRNA-Asp, -Asn, -His and -Tyr), where N is one of the four canonical nucleotides [51]. It was shown that a functional Tgt is required for efficient pathogenicity of *Shigella* bacteria [52]. A null-mutation in the *tgt* gene strongly reduces translation of *virF*-mRNA, a transcriptional activator required for the expression of a large number of *Shigella* pathogenicity genes [53]. On the other hand, RlmL K/L specifically methylates the guanine in position 2445 and the guanine in position 2069 (m7G2069) of 23S rRNA. The rRNA methyltransferases (rRNA MTases), especially those acting on 23S rRNA, are associated with development of antibiotic resistance in a wide variety of bacteria, and some of them are

recurrent human pathogens [54]. Inactivation of 23S rRNA MTases function has been shown to affect negatively translation and cell physiology [55].

Finally, a glutathione S-transferase was over-represented in the light condition. The glutathione S-transferases (GSTs) are a family of proteins that conjugate glutathione to the sulfur atom of cysteine in various compounds [56]. GST can bind to a variety of hydrophobic compounds endogenous and xenobiotic alkylating agents with high affinity, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, protecting cells from oxidative damage [57].

Taken together, these observations suggest that light could play a main role in the control of *A. nosocomialis* physiology at 37 °C, particularly modulating pathogenesis and allowing cells to respond and adapt to environmental signals.

Proteins with increased abundance in the dark compared to light.

Among the proteins with higher abundance in the dark is present the phenylacetate-CoA oxygenase, PaaK subunit. Phenylacetate-CoA oxygenase is comprised of a complex composed of 5 proteins responsible for the hydroxylation of phenylacetate-CoA (PA-CoA), which is the second catabolic step in phenylacetic acid (PAA) degradation [58]. Interestingly, we have shown that *A. nosocomialis* RUH2624 growth is stimulated in the dark when PAA is present as the sole carbon source at 37 °C [9], supporting the notion that the PAA catabolic pathway is modulated by light. Modulation of the PAA catabolic pathway has been shown to influence *A. baumannii*'s pathogenesis. In fact, inhibition of this pathway resulted in increased neutrophil migration to the site of infection and bacterial clearance [58] [59].

Another protein that appears over-represented in the dark is the nonhydrolyzing UDP-N-acetylglucosamine 2-epimerase (FC= 1.7). This enzyme catalyzes the reversible interconversion of UDP-N-acetylglucosamine (UDP-GlcNAc) to UDP-N-acetylmannosamine (UDP-ManNAc) [60], being

this compound an intermediate in the biosynthesis of a variety of bacterial capsular polysaccharides (CPSs) [61]. In several pathogenic strains, such as *B. anthracis*, *N. meningitides* and *S. aureus*, CPSs are important virulence factors that protects bacteria from the immune system of a host and harsh environmental conditions [62-64]. In this sense, the UDP-N-acetylglucosamine 2-epimerase could contribute to cell-surface polysaccharide synthesis in *A. nosocomialis*, protecting the cells in the absence of light. The probable potassium transport Kup protein, found in a greater quantity in dark, could also favor the adaptation to rapidly changing external conditions.

Conclusion

In this work, we show that light regulates motility in *A. nosocomialis* in a wide range of temperatures, which go from environmental to temperatures found in warm-blooded hosts (23 to 37 °C). This temperature dependence is different from that observed for regulation of motility by light in *A. baumannii* ATCC 17978, which was shown to occur only in the low to environmental temperature range (18 to 24 °C). We hypothesized that this could be due to an unequal endowment of *blsA* homologs, whose intrinsic characteristics may allow them to function in other temperature ranges. In this work, we show that the three BLUF domain-containing genes present in the *A. nosocomialis* RUH2624 genome [8], encode active photoreceptors. But only two of them, AnBLUF65 and AnBLUF46 are stable proteins when produced *in vitro* in the temperature range analyzed: (15-37 °C) and (15-32 °C), respectively. The fact that AnBLUF65 is the only of the three BLUF domain-containing proteins that is expressed *in vivo* along with the photo-regulatory temperature range, strongly suggests that it participates in the modulation of motility by light. Spectroscopic characterizations of AnBLUF65 and AnBLUF46 *in vitro*, indicate that AnBLUF65 is more efficient to form the light-adapted state than AnBLUF46 at 24 °C. And although AnBLUF65 efficiency is negatively affected by temperature increments, the protein remains active at 37 °C, accordingly with being the only blue light photoreceptor expressed in *A. nosocomialis*. 3D models of these proteins were presented and discussed. The relative solvent accessibility to the cofactor pocket derived from flavin emission fluorescence correlates well with the tertiary structure modeled for both proteins and both states, dark and light-adapted. We have additionally characterized intramolecular interactions to profile the underlying phenomena upon formation of the light-adapted state. The presence of aromatic clusters networks on either side of the β -sheet has been described. We propose that the rupture of an Asp-X interaction via H-bonding observed upon illumination in the loop between the Tyr-Tyr and the first set of Phe-Phe motifs, allows the

aromatic tetrads in the variable domain to displace and to adopt the light-adapted state of this part in AnBLUF65 and AnBLUF46 (Fig 10). Thus, the calculated strengthening of the π - π tetrads in AnBLUF65 and the displacements of these residues between the dark and the light-adapted forms in BlsA as observed by Chitrakar et al 2020, reinforce the idea that intramolecular signaling events involve, at least in part, hydrophobic/aromatic interactions. The extent to which these networks are present in other members of the genus *Acinetobacter*, and other microorganisms remain to be further explored.

Fig 10: Intra-protein aromatic networks in BLUFs from *A. nosocomialis*. Black ribbons represent β sheet dividing protein in two areas: Flavin/N-terminal area and variable domain. Solid lines correspond to main chain covalent bonds. Dashed lines represent relative distances between aromatic centroids for dark (black) and blue light (blue).

Overall, in this work, we have characterized different BLUF photoreceptors and show that they are operative at different temperature ranges. Despite diverse BLUF-coding genes are encoded in *A. nosocomialis*, not all are physiologically functional. It is worth mentioning that AnBLUF65 is the only BLUF-protein found to be expressed in *A. nosocomialis*, and the possibility raises that the other BLUF-photoreceptors are expressed under different conditions, such as growth media, temperatures, presence of a host, etc. The presence of AnBLUF85 only in two different *A. nosocomialis* strains and discretionally in other species of the *Acinetobacter* genus is striking, in addition to the fact that this photoreceptor is not stable and not expressed in its host. The overall information suggests that this gene might be cryptic; despite it does not seem to be subjected to genetic drift as its sequence is conserved in different species, and questions arise regarding its evolutionary origin and functionality.

Acknowledgment

This work was supported by grants from the Agencia Nacional de Promoción Científica y Tecnológica (PICT 2018- 00793) to MAM, the Consejo Nacional de Investigaciones Científicas y Tecnológicas (PIP 2017-0493) to IA and ASaCTel (Ministerio de Ciencia, Tecnología e Innovación Productiva de la Provincia de Santa Fe) IO-021-18 to MAM. We would like to thank Dr. Javier M. González for his critical reading of the manuscript and suggestions. Also, Lic. Alba Loto for her assistance with protein overexpression and purification. IA, CDB and MAM are career investigators of CONICET. BPM and MRT are fellows from the same institution. LV is a Universidad Nacional de Santiago del Estero investigator.

References

1. Nemec A, Krizova L, Maixnerova M, van der Reijden TJ, Deschaght P, Passet V, et al. Genotypic and phenotypic characterization of the *Acinetobacter calcoaceticus*-*Acinetobacter baumannii* complex with the proposal of *Acinetobacter pittii* sp. nov. (formerly *Acinetobacter* genomic species 3) and *Acinetobacter nosocomialis* sp. nov. (formerly *Acinetobacter* genomic species 13TU). *Research in microbiology*. 2011;162(4):393-404. Epub 2011/02/16. doi: 10.1016/j.resmic.2011.02.006. PubMed PMID: 21320596.
2. Knight DB, Rudin SD, Bonomo RA, Rather PN. *Acinetobacter nosocomialis*: Defining the Role of Efflux Pumps in Resistance to Antimicrobial Therapy, Surface Motility, and Biofilm Formation. *Frontiers in microbiology*. 2018;9:1902. Epub 2018/09/07. doi: 10.3389/fmicb.2018.01902. PubMed PMID: 30186249; PubMed Central PMCID: PMC6111201.
3. Dijkshoorn L, Nemec A, Seifert H. An increasing threat in hospitals: multidrug-resistant *Acinetobacter baumannii*. *Nature reviews Microbiology*. 2007;5(12):939-51. Epub 2007/11/17. doi: 10.1038/nrmicro1789. PubMed PMID: 18007677.
4. Xie R, Zhang XD, Zhao Q, Peng B, Zheng J. Analysis of global prevalence of antibiotic resistance in *Acinetobacter baumannii* infections disclosed a faster increase in OECD countries. *Emerging microbes & infections*. 2018;7(1):31. doi: 10.1038/s41426-018-0038-9. PubMed PMID: 29535298; PubMed Central PMCID: PMC5849731.
5. Wisplinghoff H, Paulus T, Lugenheim M, Stefanik D, Higgins PG, Edmond MB, et al. Nosocomial bloodstream infections due to *Acinetobacter baumannii*, *Acinetobacter pittii* and *Acinetobacter nosocomialis* in the United States. *J Infect*. 2012;64(3):282-90. Epub 2012/01/03. doi: 10.1016/j.jinf.2011.12.008. PubMed PMID: 22209744.
6. Chusri S, Chongsuvivatwong V, Rivera JI, Silpapojakul K, Singkhamanan K, McNeil E, et al. Clinical outcomes of hospital-acquired infection with *Acinetobacter nosocomialis* and *Acinetobacter pittii*. *Antimicrobial agents and chemotherapy*. 2014;58(7):4172-9. Epub 2014/05/14. doi: 10.1128/AAC.02992-14. PubMed PMID: 24820079; PubMed Central PMCID: PMC4068534.
7. Huang L, Chen TL, Lee YT, Lee MH, Kuo SC, Yu KW, et al. Risk factors for imipenem-nonsusceptible *Acinetobacter nosocomialis* bloodstream infection. *J Microbiol Immunol Infect*. 2014;47(4):311-7. Epub 2013/05/07. doi: 10.1016/j.jmii.2013.02.002. PubMed PMID: 23642729.

689 8. Golic A, Vaneechoutte M, Nemec A, Viale AM, Actis LA, Mussi MA. Staring at the cold sun:
690 blue light regulation is distributed within the genus *Acinetobacter*. *PloS one*. 2013;8(1):e55059.
691 doi: 10.1371/journal.pone.0055059. PubMed PMID: 23358859; PubMed Central PMCID:
692 PMC3554667.

693 9. Tuttobene M, Perez J, Pavesi E, Biancotti D, Cribb P, Altilio M, et al. Light modulates
694 important pathogenic determinants and virulence in *Acinetobacter baumannii*, *Pseudomonas*
695 *aeruginosa* and *Staphylococcus aureus* ESKAPE pathogens. . *Journal of bacteriology*. 2020;En
696 revisión.

697 10. Abatedaga I, Valle L, Golic AE, Muller GL, Cabruja M, Moran Vieyra FE, et al. Integration of
698 Temperature and Blue-Light Sensing in *Acinetobacter baumannii* Through the BlsA Sensor.
699 *Photochemistry and photobiology*. 2017;93(3):805-14. doi: 10.1111/php.12760. PubMed PMID:
700 28500705.

701 11. Golic AE, Valle L, Jaime PC, Alvarez CE, Parodi C, Borsarelli CD, et al. BlsA Is a Low to
702 Moderate Temperature Blue Light Photoreceptor in the Human Pathogen *Acinetobacter*
703 *baumannii*. *Frontiers in microbiology*. 2019;10:1925. Epub 2019/09/10. doi:
704 10.3389/fmicb.2019.01925. PubMed PMID: 31497002; PubMed Central PMCID: PMC6712483.

705 12. Bitrian M, Gonzalez RH, Paris G, Hellingwerf KJ, Nudel CB. Blue-light-dependent inhibition
706 of twitching motility in *Acinetobacter baylyi* ADP1: additive involvement of three BLUF-domain-
707 containing proteins. *Microbiology (Reading)*. 2013;159(Pt 9):1828-41. Epub 2013/07/03. doi:
708 10.1099/mic.0.069153-0. PubMed PMID: 23813679.

709 13. Schramm STJ, Place K, Montana S, Almuzara M, Fung S, Fernandez JS, et al. Genetic and
710 Phenotypic Features of a Novel *Acinetobacter* Species, Strain A47, Isolated From the Clinical
711 Setting. *Frontiers in microbiology*. 2019;10:1375. Epub 2019/07/06. doi:
712 10.3389/fmicb.2019.01375. PubMed PMID: 31275288; PubMed Central PMCID: PMC6591377.

713 14. Muller GL, Tuttobene M, Altilio M, Martinez Amezcaga M, Nguyen M, Cribb P, et al. Light
714 Modulates Metabolic Pathways and Other Novel Physiological Traits in the Human Pathogen
715 *Acinetobacter baumannii*. *Journal of bacteriology*. 2017;199(10). doi: 10.1128/JB.00011-17.
716 PubMed PMID: 28289081; PubMed Central PMCID: PMC5405214.

717 15. Mussi MA, Gaddy JA, Cabruja M, Arivett BA, Viale AM, Rasia R, et al. The opportunistic
718 human pathogen *Acinetobacter baumannii* senses and responds to light. *Journal of bacteriology*.
719 2010;192(24):6336-45. doi: 10.1128/JB.00917-10. PubMed PMID: 20889755; PubMed Central
720 PMCID: PMC3008525.

721 16. Ramirez MS, Traglia GM, Perez JF, Muller GL, Martinez MF, Golic AE, et al. White and blue
722 light induce reduction in susceptibility to minocycline and tigecycline in *Acinetobacter* spp. and
723 other bacteria of clinical importance. *Journal of medical microbiology*. 2015;64(Pt 5):525-37. doi:
724 10.1099/jmm.0.000048. PubMed PMID: 25737436.

725 17. Tuttobene MR, Cribb P, Mussi MA. BlsA integrates light and temperature signals into iron
726 metabolism through Fur in the human pathogen *Acinetobacter baumannii*. *Scientific reports*.
727 2018;8(1):7728. doi: 10.1038/s41598-018-26127-8. PubMed PMID: 29769610; PubMed Central
728 PMCID: PMC5955987.

729 18. Tuttobene MR, Fernandez-Garcia L, Blasco L, Cribb P, Ambroa A, Muller GL, et al. Quorum
730 and Light Signals Modulate Acetoin/Butanediol Catabolism in *Acinetobacter* spp. *Frontiers in*
731 *microbiology*. 2019;10:1376. doi: 10.3389/fmicb.2019.01376. PubMed PMID: 31281296; PubMed
732 Central PMCID: PMC6595428.

733 19. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. qBase relative
734 quantification framework and software for management and automated analysis of real-time
735 quantitative PCR data. *Genome biology*. 2007;8(2):R19. doi: 10.1186/gb-2007-8-2-r19. PubMed
736 PMID: 17291332; PubMed Central PMCID: PMC1852402.

737 20. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
738 quantitative PCR and the 2⁻($\Delta\Delta C_T$) Method. *Methods*. 2001;25(4):402-8. Epub
739 2002/02/16. doi: 10.1006/meth.2001.1262. PubMed PMID: 11846609.

740 21. Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2--a multiple
741 sequence alignment editor and analysis workbench. *Bioinformatics*. 2009;25(9):1189-91. Epub
742 2009/01/20. doi: 10.1093/bioinformatics/btp033. PubMed PMID: 19151095; PubMed Central
743 PMCID: PMCPMC2672624.

744 22. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al. SWISS-
745 MODEL: homology modelling of protein structures and complexes. *Nucleic acids research*.
746 2018;46(W1):W296-W303. Epub 2018/05/23. doi: 10.1093/nar/gky427. PubMed PMID: 29788355;
747 PubMed Central PMCID: PMCPMC6030848.

748 23. Piovesan D, Minervini G, Tosatto SC. The RING 2.0 web server for high quality residue
749 interaction networks. *Nucleic acids research*. 2016;44(W1):W367-74. Epub 2016/05/21. doi:
750 10.1093/nar/gkw315. PubMed PMID: 27198219; PubMed Central PMCID: PMCPMC4987896.

751 24. Webb-Robertson BJ, Wiberg HK, Matzke MM, Brown JN, Wang J, McDermott JE, et al.
752 Review, evaluation, and discussion of the challenges of missing value imputation for mass
753 spectrometry-based label-free global proteomics. *Journal of proteome research*. 2015;14(5):1993-

2001. Epub 2015/04/10. doi: 10.1021/pr501138h. PubMed PMID: 25855118; PubMed Central
PMCID: PMCPMC4776766.

25. Potron A, Poirel L, Croize J, Chanteperdrix V, Nordmann P. Genetic and biochemical
characterization of the first extended-spectrum CARB-type beta-lactamase, RTG-4, from
Acinetobacter baumannii. *Antimicrobial agents and chemotherapy*. 2009;53(7):3010-6. Epub
2009/04/22. doi: 10.1128/AAC.01164-08. PubMed PMID: 19380596; PubMed Central PMCID:
PMCPMC2704689.

26. Fujisawa T, Masuda S. Light-induced chromophore and protein responses and mechanical
signal transduction of BLUF proteins. *Biophysical reviews*. 2018;10(2):327-37. Epub 2017/12/14.
doi: 10.1007/s12551-017-0355-6. PubMed PMID: 29235080; PubMed Central PMCID:
PMCPMC5899715.

27. Valle L, Vieyra FE, Borsarelli CD. Hydrogen-bonding modulation of excited-state properties
of flavins in a model of aqueous confined environment. *Photochemical & photobiological sciences*
: Official journal of the European Photochemistry Association and the European Society for
Photobiology. 2012;11(6):1051-61. Epub 2012/03/22. doi: 10.1039/c2pp05385c. PubMed PMID:
22434390.

28. Kraft BJ, Masuda S, Kikuchi J, Dragnea V, Tollin G, Zaleski JM, et al. Spectroscopic and
mutational analysis of the blue-light photoreceptor AppA: a novel photocycle involving flavin
stacking with an aromatic amino acid. *Biochemistry*. 2003;42(22):6726-34. Epub 2003/06/05. doi:
10.1021/bi030055o. PubMed PMID: 12779327.

29. Villegas JM, Valle L, Moran Vieyra FE, Rintoul MR, Borsarelli CD, Rapisarda VA. FAD binding
properties of a cytosolic version of *Escherichia coli* NADH dehydrogenase-2. *Biochim Biophys Acta*.
2014;1844(3):576-84. Epub 2014/01/15. doi: 10.1016/j.bbapap.2013.12.021. PubMed PMID:
24418395.

30. Tanwar M, Nahar S, Gulati S, Veetil SK, Kateriya S. Molecular determinant modulates
thermal recovery kinetics and structural integrity of the bacterial BLUF photoreceptor. *FEBS
letters*. 2016;590(14):2146-57. Epub 2016/05/28. doi: 10.1002/1873-3468.12227. PubMed PMID:
27230699.

31. Valle L, Abatedaga I, Vieyra FE, Bortolotti A, Cortez N, Borsarelli CD. Enhancement of
photophysical and photosensitizing properties of flavin adenine dinucleotide by mutagenesis of
the C-terminal extension of a bacterial flavodoxin reductase. *Chemphyschem*. 2015;16(4):872-83.
Epub 2015/02/03. doi: 10.1002/cphc.201402774. PubMed PMID: 25641205.

786 32. Chitrakar I, Iuliano JN, He Y, Woroniecka HA, Tolentino Collado J, Wint JM, et al. Structural
787 Basis for the Regulation of Biofilm Formation and Iron Uptake in *A. baumannii* by the Blue-Light-
788 Using Photoreceptor, BlsA. *ACS Infect Dis.* 2020;6(10):2592-603. Epub 2020/09/15. doi:
789 10.1021/acsinfecdis.0c00156. PubMed PMID: 32926768.

790 33. Khrenova K. DT, Nemukhinac A. Molecular mechanism of the dark-state recovery in BLUF
791 photoreceptors. *Chemical Physics Letters.* 2017;676. doi: doi.org/10.1016/j.cplett.2017.03.035.

792 34. Lanzarotti E, Biekofsky RR, Estrin DA, Marti MA, Turjanski AG. Aromatic-aromatic
793 interactions in proteins: beyond the dimer. *Journal of chemical information and modeling.*
794 2011;51(7):1623-33. Epub 2011/06/15. doi: 10.1021/ci200062e. PubMed PMID: 21662246.

795 35. Pezza A, Tuttobene M, Abatedaga I, Valle L, Borsarelli CD, Mussi MA. Through the eyes of a
796 pathogen: light perception and signal transduction in *Acinetobacter baumannii*. *Photochemical &*
797 *photobiological sciences : Official journal of the European Photochemistry Association and the*
798 *European Society for Photobiology.* 2019;18(10):2363-73. Epub 2019/07/11. doi:
799 10.1039/c9pp00261h. PubMed PMID: 31290528.

800 36. Park YJ, Song ES, Kim YT, Noh TH, Kang HW, Lee BM. Analysis of virulence and growth of a
801 purine auxotrophic mutant of *Xanthomonas oryzae* pathovar *oryzae*. *FEMS microbiology letters.*
802 2007;276(1):55-9. Epub 2007/09/25. doi: 10.1111/j.1574-6968.2007.00909.x. PubMed PMID:
803 17888004.

804 37. Kilstrup M, Hammer K, Ruhdal Jensen P, Martinussen J. Nucleotide metabolism and its
805 control in lactic acid bacteria. *FEMS microbiology reviews.* 2005;29(3):555-90. Epub 2005/06/07.
806 doi: 10.1016/j.femsre.2005.04.006. PubMed PMID: 15935511.

807 38. McFarland WC, Stocker BA. Effect of different purine auxotrophic mutations on mouse-
808 virulence of a Vi-positive strain of *Salmonella dublin* and of two strains of *Salmonella typhimurium*.
809 *Microb Pathog.* 1987;3(2):129-41. Epub 1987/08/01. doi: 10.1016/0882-4010(87)90071-4.
810 PubMed PMID: 2849016.

811 39. Mei JM, Nourbakhsh F, Ford CW, Holden DW. Identification of *Staphylococcus aureus*
812 virulence genes in a murine model of bacteraemia using signature-tagged mutagenesis. *Molecular*
813 *microbiology.* 1997;26(2):399-407. Epub 1998/02/12. doi: 10.1046/j.1365-2958.1997.5911966.x.
814 PubMed PMID: 9383163.

815 40. Samant S, Lee H, Ghassemi M, Chen J, Cook JL, Mankin AS, et al. Nucleotide biosynthesis is
816 critical for growth of bacteria in human blood. *PLoS pathogens.* 2008;4(2):e37. Epub 2008/02/20.
817 doi: 10.1371/journal.ppat.0040037. PubMed PMID: 18282099; PubMed Central PMCID:
818 PMCPMC2242838.

819 41. Hobbs G. FC, Gardner D., Flett F., Oliver S. Pigmented antibiotic production by
820 *Streptomyces coelicolor* A3(2): kinetics and the influence of nutrients. *Microbiology*. 1990;136(11).
821 doi: doi.org/10.1099/00221287-136-11-2291.

822 42. Navone L, Casati P, Licona-Cassani C, Marcellin E, Nielsen LK, Rodriguez E, et al. Allantoin
823 catabolism influences the production of antibiotics in *Streptomyces coelicolor*. *Applied*
824 *microbiology and biotechnology*. 2014;98(1):351-60. Epub 2013/12/03. doi: 10.1007/s00253-013-
825 5372-1. PubMed PMID: 24292080.

826 43. Burns RO, Umbarger HE, Gross SR. The Biosynthesis of Leucine. Iii. The Conversion of
827 Alpha-Hydroxy-Beta-Carboxyisocaproate to Alpha-Ketoisocaproate. *Biochemistry*. 1963;2:1053-8.
828 Epub 1963/09/01. doi: 10.1021/bi00905a024. PubMed PMID: 14087358.

829 44. Csonka LN, Leisinger T. Biosynthesis of Proline. *EcoSal Plus*. 2007;2(2). Epub 2007/04/01.
830 doi: 10.1128/ecosalplus.3.6.1.4. PubMed PMID: 26443591.

831 45. Trovato M. FG, Signorelli S., Funck D. . Proline metabolism and its functions in
832 development and stress tolerance. . In: Springer, editor. In *Osmoprotectant-Mediated Abiotic*
833 *Stress Tolerance in Plants* 2019. p. 41-72.

834 46. Christgen SL, Becker DF. Role of Proline in Pathogen and Host Interactions. *Antioxid Redox*
835 *Signal*. 2019;30(4):683-709. Epub 2017/12/16. doi: 10.1089/ars.2017.7335. PubMed PMID:
836 29241353; PubMed Central PMCID: PMC6338583.

837 47. Sklar JG, Wu T, Kahne D, Silhavy TJ. Defining the roles of the periplasmic chaperones SurA,
838 Skp, and DegP in *Escherichia coli*. *Genes Dev*. 2007;21(19):2473-84. Epub 2007/10/03. doi:
839 10.1101/gad.1581007. PubMed PMID: 17908933; PubMed Central PMCID: PMC6338583.

840 48. Nikaïdo H. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol*
841 *Mol Biol Rev*. 2003;67(4):593-656. Epub 2003/12/11. doi: 10.1128/mmbr.67.4.593-656.2003.
842 PubMed PMID: 14665678; PubMed Central PMCID: PMC6338583.

843 49. Klein K, Sonnabend MS, Frank L, Leibiger K, Franz-Wachtel M, Macek B, et al. Deprivation
844 of the Periplasmic Chaperone SurA Reduces Virulence and Restores Antibiotic Susceptibility of
845 Multidrug-Resistant *Pseudomonas aeruginosa*. *Frontiers in microbiology*. 2019;10:100. Epub
846 2019/03/09. doi: 10.3389/fmicb.2019.00100. PubMed PMID: 30846971; PubMed Central PMCID:
847 PMC6338583.

848 50. Lee K, Lee KM, Go J, Ryu JC, Ryu JH, Yoon SS. The ferrichrome receptor A as a new target
849 for *Pseudomonas aeruginosa* virulence attenuation. *FEMS microbiology letters*. 2016;363(11).
850 Epub 2016/05/18. doi: 10.1093/femsle/fnw104. PubMed PMID: 27190289.

51. Morris RC, Elliott MS. Queuosine modification of tRNA: a case for convergent evolution. Mol Genet Metab. 2001;74(1-2):147-59. Epub 2001/10/11. doi: 10.1006/mgme.2001.3216. PubMed PMID: 11592812.

52. Durand JM, Bjork GR, Kuwae A, Yoshikawa M, Sasakawa C. The modified nucleoside 2-methylthio-N6-isopentenyladenosine in tRNA of *Shigella flexneri* is required for expression of virulence genes. Journal of bacteriology. 1997;179(18):5777-82. Epub 1997/09/19. doi: 10.1128/jb.179.18.5777-5782.1997. PubMed PMID: 9294434; PubMed Central PMCID: PMC179466.

53. Durand JM, Dagberg B, Uhlin BE, Bjork GR. Transfer RNA modification, temperature and DNA superhelicity have a common target in the regulatory network of the virulence of *Shigella flexneri*: the expression of the *virF* gene. Molecular microbiology. 2000;35(4):924-35. Epub 2000/02/26. doi: 10.1046/j.1365-2958.2000.01767.x. PubMed PMID: 10692168.

54. LaMarre JM, Howden BP, Mankin AS. Inactivation of the indigenous methyltransferase RlmN in *Staphylococcus aureus* increases linezolid resistance. Antimicrobial agents and chemotherapy. 2011;55(6):2989-91. Epub 2011/03/30. doi: 10.1128/AAC.00183-11. PubMed PMID: 21444696; PubMed Central PMCID: PMC3101465.

55. Liiv A, Karitkina D, Maivali U, Remme J. Analysis of the function of *E. coli* 23S rRNA helix-loop 69 by mutagenesis. BMC molecular biology. 2005;6:18. Epub 2005/08/02. doi: 10.1186/1471-2199-6-18. PubMed PMID: 16053518; PubMed Central PMCID: PMC1190176.

56. Arias M, Fleischner G., Kirsch R., Mishkin S., Gatmaitan Z. . In: I. M. Arias WB, Jakoby, editor. In Glutathione: Metabolism and Function 1976. p. 175-88.

57. Sheehan D, Meade G, Foley VM, Dowd CA. Structure, function and evolution of glutathione transferases: implications for classification of non-mammalian members of an ancient enzyme superfamily. Biochem J. 2001;360(Pt 1):1-16. Epub 2001/11/07. doi: 10.1042/0264-6021:3600001. PubMed PMID: 11695986; PubMed Central PMCID: PMC1222196.

58. Cerqueira GM, Kostoulas X, Khoo C, Aibinu I, Qu Y, Traven A, et al. A global virulence regulator in *Acinetobacter baumannii* and its control of the phenylacetic acid catabolic pathway. The Journal of infectious diseases. 2014;210(1):46-55. doi: 10.1093/infdis/jiu024. PubMed PMID: 24431277.

59. Bhuiyan MS, Ellett F, Murray GL, Kostoulas X, Cerqueira GM, Schulze KE, et al. *Acinetobacter baumannii* phenylacetic acid metabolism influences infection outcome through a direct effect on neutrophil chemotaxis. Proceedings of the National Academy of Sciences of the

883 United States of America. 2016;113(34):9599-604. doi: 10.1073/pnas.1523116113. PubMed PMID:
884 27506797; PubMed Central PMCID: PMC5003227.

885 60. Samuel J, Tanner ME. Mechanistic aspects of enzymatic carbohydrate epimerization. Nat
886 Prod Rep. 2002;19(3):261-77. Epub 2002/07/26. doi: 10.1039/b100492l. PubMed PMID:
887 12137277.

888 61. Cress BF, Englaender JA, He W, Kasper D, Linhardt RJ, Koffas MA. Masquerading microbial
889 pathogens: capsular polysaccharides mimic host-tissue molecules. FEMS microbiology reviews.
890 2014;38(4):660-97. Epub 2014/01/01. doi: 10.1111/1574-6976.12056. PubMed PMID: 24372337;
891 PubMed Central PMCID: PMCPMC4120193.

892 62. Velloso LM, Bhaskaran SS, Schuch R, Fischetti VA, Stebbins CE. A structural basis for the
893 allosteric regulation of non-hydrolysing UDP-GlcNAc 2-epimerases. EMBO Rep. 2008;9(2):199-205.
894 Epub 2008/01/12. doi: 10.1038/sj.embor.7401154. PubMed PMID: 18188181; PubMed Central
895 PMCID: PMCPMC2246411.

896 63. Hurlburt NK, Guan J, Ong H, Yu H, Chen X, Fisher AJ. Structural characterization of a
897 nonhydrolyzing UDP-GlcNAc 2-epimerase from *Neisseria meningitidis* serogroup A. Acta
898 Crystallogr F Struct Biol Commun. 2020;76(Pt 11):557-67. Epub 2020/11/03. doi:
899 10.1107/S2053230X20013680. PubMed PMID: 33135674; PubMed Central PMCID:
900 PMCPMC7605110.

901 64. Mann PA, Muller A, Wolff KA, Fischmann T, Wang H, Reed P, et al. Chemical Genetic
902 Analysis and Functional Characterization of Staphylococcal Wall Teichoic Acid 2-Epimerases
903 Reveals Unconventional Antibiotic Drug Targets. PLoS pathogens. 2016;12(5):e1005585. Epub
904 2016/05/06. doi: 10.1371/journal.ppat.1005585. PubMed PMID: 27144276; PubMed Central
905 PMCID: PMCPMC4856313.

906

907

Supporting information

S1 Fig. Genomic context of AnBLUF65 homologs present in different strains of *A. nosocomialis*.

(A). Coding-sequences (CDSs) are located in their corresponding frame. AnBLUF65 is indicated as pink arrow. Different colors indicate different functions. Gene annotations are indicated as numbers above the schemes, with the following codes: 1- AnBLUF65, 2- DUF2171, 3- Acyl-CoA-dehydrogenase, 4- GlcNAc-PI-de-N-acetylase, 5- methyltransferase domain, 6- glycosyltransferase, 7- DDE endonuclease domain, putative transposase, 8- helix-turn-helix for DDE superfamily endonuclease, 9- BOF- class 2b aminoacyl-tRNA synthetases- NirD/YgiW/Y family stress tolerance protein, 10- HP, 11- poly (R)-hydroxyalkanoic acid synthase, 12- sodium/glutamate symporter, 13- proton antiporter-2 (CPA2) family, 14- NDAB- Rossmann Superfamily- Oxidoreductase, 15- DoxX-like family, 16- LysR family transcriptional regulator, 17- AKR15A family of aldo-keto reductase, 18- pyrabactin resistance 1 (PYR1) receptor, 19- type I secretion target GGXGXDXXX repeat protein, 20- Paax domain, 21- glutathione-regulated potassium-efflux system protein KefC, 22- IS3 family transposase, 23- L- asparagine transporter.

S2 Fig. Genomic context of AnBLUF46 homologs present in different strains of *A. nosocomialis*.

(A). Coding-sequences (CDSs) are located in their corresponding frame. AnBLUF46 is indicated as pink arrow. Different colors indicate different functions. Gene annotations are indicated as numbers above the schemes, with the following codes: 1- AnBLUF46, 2- aromatic acid:H⁺ symporter (AAHS), 3- AraC family transcriptional regulator RmlC-cupin protein, 4- 3-hydroxybutyrate dehydrogenase, 5- H⁺/gluconate symporter, 6- MBL fold metallo-hydrolase, 7- HP, 8- DUF2846- putative conjugal domain, 9- DUF3298- putative heat shock protein, 10- HP, 11- LysR family transcriptional regulator, 12- succinyl-CoA:3- ketoacid-coenzyme A transferase subunit A, 13- succinyl-CoA:3-ketoacid-coenzyme A transferase subunit B, 14- multidrug efflux MFS transporter, 15- pimeloyl-ACP methyl ester carboxylesterase, 16- IS5 transposase, 17- TetR/AcrR

family transcriptional regulator, 18- DoxX family protein, 19- IS transposase, 20- anion permease
 ArsB/NhaD, 21- 3-oxoacyl-ACP reductase FabG, 22- nuclear transport factor 2 family protein, 23-
 fermentarion-respiration switch protein FrsA, 24- DDE transposase domain, 25- Bcr/CflA family
 drug resistance efflux transporter.

S3 Fig. Genomic context of AnBLUF85 homologs present in different strains of *A. nosocomialis*.

(A). Coding-sequences (CDSs) are located in their corresponding frame. AnBLUF85 is indicated as
 pink arrow. Different colors indicate different functions. Gene annotations are indicated as
 numbers above the schemes, with the following codes: 1- AnBLUF85, 2- relaxase-MobA/MobL
 family protein, 3- FAM199X, 4-HP, 5- N-acetyltransferasa, 6- conjugal transfer pilus assembly
 protein TraB, 7-RepB initiator replication protein, 8- potassium transporter, 9-mRNA-degrading
 endonuclease RelE, 10- chromate transport protein ChrA.

**S4 Fig: Visible absorption spectra changes of AnBLUF85 photoreceptor by blue light
 illumination, in working buffer solution at 12 °C. dAnBLUF85 (black line) and lAnBLUF 85 (pale
 green)**

A

25°C

26°C

27°C

32°C

37°C

L

D

B

C

anbluf65

Figure

A

<i>Slr1694</i>	1	- - - - -	- - - - -	MSLYRL	IYSSQG	IPN - -	LQPQDL	KDILESSQRN	31
<i>TII0078</i>	1	- - - - -	- - - - -	MGLHRL	IYLS	CATDG - -	LSYPDL	RDIMAKSEVN	31
<i>AnBLUF65</i>	1	- - - - -	- - - - -	MNVRL	CYASQRN	NENNEDL	LQDL	RDILTEARDF	32
<i>BlsA</i>	1	- - - - -	- - - - -	MNVRL	CYASQRNE	KNEDL	LQDL	RDILTEARDF	32
<i>AnBLUF85</i>	1	- - - - -	- - - - -	MNDFRL	LYVSK	IKSNINPVN -	DLYD	ILTEAVAF	32
<i>AnBLUF46</i>	1	- - - - -	- - - - -	MSLIGF	MYASKTN	SEHSQIKQDL	IDIL	TEAVKF	33
<i>Appa</i>	1	MQHDL	EADV	TMTGSD	LVSCCYRSLA	APDL	TLRDL	LDIVETSQA	44
<i>BlrB</i>	1	- - - - -	- - - - -	MDEL	VS	LTYSRVRLADP	VADIVQ	IMRASRVR	32

<i>Slr1694</i>	32	NPANG	ITGL	LCYSKPAF	LQV	LEGECEQV	NETYHR -	IVQDERHHS	74
<i>TII0078</i>	32	NLRDG	ITGM	LCYGNGMF	LQT	LEGDRQKV	SETYAR -	ILKDPRHHS	74
<i>AnBLUF65</i>	33	NDLNE	ICGV	LYYADNAFF	QC	LEGEKEV	VERLF	EK - IQKDQRHHS	75
<i>BlsA</i>	33	NDLNG	ICGV	LYYADNAFF	QC	LEGEQEV	VERLF	EK - IQKDQRHYN	75
<i>AnBLUF85</i>	33	NSLNE	IYGV	LYYGNDYF	IQCV	EGQKDK	VEYLF	YEK - ILKDPRHKN	76
<i>AnBLUF46</i>	34	NSQND	ITGV	LYYGNGYF	LQY	LEGEKEQV	ETLFYKS	ILKDSRHQN	77
<i>Appa</i>	45	NARAQL	TGAL	FYSQGV	FFQW	LEGRPA	AAVAEVMTH -	IQRDRRHSN	87
<i>BlrB</i>	33	NLRLG	ITG	ILLYNGVHF	VQT	IEGPR	SACDEL	FRL - ISADPRHQE	75

bioRxiv preprint doi: <https://doi.org/10.1101/2021.07.23.453528>; this version posted July 23, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

<i>Slr1694</i>	75	PQ	IE	CMPI	IRRRNF	EVWSM	QA	ITVNDL	STEQVK	TLVLKYS	SGFTT	118
<i>TII0078</i>	75	AE	IVE	FKA	IEERTF	INWSM	RLVQLG	EMDS	DTIRRL	RLKYS	SPAAT	118
<i>AnBLUF65</i>	76	VKW	LCTYS	IEENS	FQRWS	MKYVQR	NTNIEA	FFLKM	GEST	FNPI	L	119
<i>BlsA</i>	76	IKW	LCTYS	IDEHS	FQRWS	MKYVQR	NTNIEA	FFLNM	GENT	FNPL	L	119
<i>AnBLUF85</i>	77	CEV	LSFEN	IEKYL	FSAWH	MKYAMI	HKEV	IDFF	KHYH	NDEF	NPYL	120
<i>AnBLUF46</i>	78	CE	IIF	LEP	SEQRL	FKHWS	MKFAP	INTK	IKDFF	FHHV	DEFNPYL	121
<i>Appa</i>	88	VE	ILAE	EEP	IAKRR	FAGWH	MLSC	SEADM	RS	LGLAES	RQ - - - - -	125
<i>BlrB</i>	76	I	LA	FDLEP	ITARR	FPDWS	MRIV	SRKEL	RALAP	DLER	LDLSG	PED 119

<i>Slr1694</i>	119	L	RPSAM	DPEQ	CLN	FL	LDIA	KIYEL	SDNFF	LDL - - - - -	150
<i>TII0078</i>	119	FQ	PRSM	TAEQ	CFR	FL	KELY	DMSQGS - - - - -	143		
<i>AnBLUF65</i>	120	L	NQQNL	KFF	LNELL	IAEQ	TKVD	TVKKV	G	MVN	NRGVNPF 156
<i>BlsA</i>	120	L	NQQNL	KFF	LNELL	IAEQ	TKMNT	VKKV	G	MVN	NRGVNPF 156
<i>AnBLUF85</i>	121	L	NSST	IPAF	I - ELL	STHG	DSFS	ILK	SKALKK - - - - -	150	
<i>AnBLUF46</i>	122	L	NTNS	IPSF	I - ELL	VDQ - - - - -	PNL	KADQYQV - - - - -	147		
<i>Appa</i>	- - - - -	- - - - -	- - - - -	- - - - -	- - - - -	- - - - -	- - - - -	- - - - -	- - - - -	- - - - -	
<i>BlrB</i>	120	V	AELH	RT	IAAS	L	SKG	DAD	ASG - - - - -	140

Figure

A

B

Figure

Figure

A **B** **C** **D** **E** **F**

Figure

Figure

Figure

BIOLOGICAL PROCESS

L

D

MOLECULAR FUNCTION

L

D

CELLULAR COMPONENT

L

D

Figure

AnBLUF65

AnBLUF46

Figure

RUH2624
ABUH381
PR362
UBA8118
KAN02
XH551
UBA9443
XH653
ABBL032
Ab22222
28F

M2

AC1530

J1A

FDAARGOS_129

Ab1
KNA01
NIPH2119

6411

SSA3

NIPH386

AMA38
ARLG1930
AMA22

Figure

bioRxiv preprint doi: <https://doi.org/10.1101/2021.07.23.453528>; this version posted July 23, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure