Evaluation of the sensitivity and specificity of three diagnostic tests for *Coxiella burnetii* infection in cattle and buffaloes in Punjab (India) using Bayesian latent class analysis

E. Meletis^a, R. Keshavamurthy^{b,c}, B.B. Singh^{b,d}, R.S. Aulakh^b, N.K. Dhand^d, P. Kostoulas^{a,*}

^aFaculty of Public and One (Integrated) Health, University of Thessaly, Mavromichali st., Karditsa, 43100, Greece (Email: emeletis@outlook.com [Meletis]; pkost@uth.gr [Kostoulas])

^bSchool of Public Health & Zoonoses, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, Punjab, 141004, India (Email: r.keshavamurthy@wsu.edu [Keshavamurthy]; bbsdhalwal@gmail.com [Singh]; rsaulakh@rediffmail.com [Aulakh])

^cWashington State University, Paul G. Allen School for Global Animal Health

^dSydney School of Veterinary Science, The University of Sydney, Camden, NSW, Australia (Email: navneet.dhand@sydney.edu.au [Dhand])

*Corresponding author: Eleftherios Meletis (Email address: emeletis@outlook.com; Postal address: Mavromichali st., Karditsa, 43100, Greece)

Abstract

Q Fever is a zoonotic disease of significant animal and public health concern, caused by *Coxiella burnetii* (*C. burnetii*), an obligate intracellular bacterium. This study was done to evaluate the sensitivity (*Se*) and specificity (*Sp*) of three diagnostic methods to diagnose *C. burnetii* infection in cattle and buffaloes in Punjab, India: an indirect ELISA method applied in serum samples and a trans-Polymerase Chain Reaction (trans-PCR) technique applied in milk samples and genital swabs. Bayesian Latent Class Models were developed following the STARD-BLCM reporting guidelines. Conditional independence was assumed between the tests, given (i) the different biological principle of ELISA and trans-PCR and (ii) the fact that the trans-PCR was performed on different tissues. The ELISA method in the serum samples showed the highest *Se* of 0.97 (95% Probability Intervals (PIs): 0.93; 0.99) compared to the trans-PCR method applied in milk samples 0.76 (0.62; 0.87) and genital swabs 0.7 (0.55; 0.82). The *Sps* of all tests were high, with trans-PCR in genital swabs recording the highest *Sp* of 0.99 (0.98; 1), while the *Sp* of trans-PCR in milk samples and ELISA in serum samples were 0.98 (0.96; 0.99) and 0.95 (0.93; 0.97) respectively. The study results show that none of the applied tests are perfect, therefore, a testing regimen based on the diagnostic characteristic of the tests may be considered for diagnosis of *C. burnetii*.

1. Introduction

Q Fever is a zoonotic disease that was first described by Edward Holbrook Derrick (Derrick, 1937) in Queensland. Q fever cases have been reported worldwide, except in New Zealand and Antarctica (Hilbink et al., 1993; Kaplan and Bertagna, 1955) and according to the OIE Terrestrial Animal Health Code, OIE countries and territories are obligated to report occurrences of the disease (OIE, 2018). Q stands for “query” and this designation was applied when the causative agent of the disease was unknown (OIE, 2018).

Coxiella burnetii (*C. burnetii*), an obligate intracellular Gram-negative coccobacillary bacterium was identified as the causative agent of Q fever in 1938 (Raoult and Marrie, 1995; McDade, 1990). *C. burnetii*, as a Gram-negative bacterium can display two different phenotypes. Phase I bacteria are highly virulent, while Phase II bacteria are avirulent (Roest et al., 2013). *C. burnetii*, has been isolated from many domestic and wild animals, birds and arthropods; however, cattle, buffaloes, sheep, goats and humans are commonly affected (Babudieri, 1959). The species under investigation in this study are cattle and buffaloes. Further, many Ixodidae and Argasidae ticks are considered reservoirs of the bacterium (Mauren and Raoult, 1999). Even though prevalence studies of *C. burnetii* in ticks yield negligible estimates, bacterium transmission to tick species is considered feasible (Sprong et al., 2010).

Circulation of the bacterium has been described in wild animals’ populations and arthropods and dispersion of the bacterium in domestic populations can occur through air, direct contact and animal secretions/excretions (e.g., vaginal discharge, placenta, milk, feces, urine, saliva, amniotic fluid) (Angelakis and Raoult, 2010).

Limited information on the pathogenesis of *C. burnetii* in domestic animals is available, while under laboratory conditions several studies based on different animal models (e.g., guinea pigs, mice) have been conducted (Roest et al., 2013). In most cases, inhalation of aerosols or dust contaminated with birth fluids is described as the main route of infection (Waag, 2007). The pathogenesis and associated histopathological findings depend on the route of infection (Roest et al., 2013). Large *C. burnetii* concentrations are present in the infected placenta and amniotic fluid, while infected cows can shed the microorganism in milk for up to 32 months (Marrie, 2003). Guatteo et al. (2006) studied three different shedding routes - milk, vaginal mucus, feces - of the bacterium. Study results indicated no predominant *C. burnetii* shedding route and for the majority of shedder cows one shedding route was identified (Guatteo et al., 2006).

In livestock, the disease is usually subclinical, but the clinical form is associated with reproductive complications such as abortions, stillbirth, weak calves, repeat breeding and general clinical signs (e.g., anorexia) (Lang and Marie, 1990; EFSA, 2010; Keshavamurthy et al., 2019). Thus, *C. burnetii* excretion via feces, vaginal mucus and milk has been reported, sometimes independent of an abortion history (Roest et al., 2013).

Again, the host's immune response to limit the infection has been studied in animal models. Specifically, during infection macrophages are the major target cells, while T-cells -associated with cellular immunity- and B-cells -associated with humoral immunity-, are critical for *C. burnetii* clearance after infection and tissue damage prevention, respectively. Antibody detection differs between the two Phases and the species (Roest et al., 2013).

Further, for the species under investigation in this study – cattle and buffaloes- the course of infection is considered similar (Lucchese et al., 2016).

In humans, the disease is observed in the (i) acute form, where flu-like symptoms, atypical pneumonia, hepatitis and cardiac involvement may be present and (ii) chronic form, that is more severe, than the acute, and fatal without appropriate therapy (Arricau-Bouvery and Rodolakis, 2005). Therefore, *C. burnetii* is characterized as a microorganism of great animal and public health concern.

Since there are no pathognomonic characteristics associated with *C. burnetii* infection, diagnosis poses a challenge. Many diagnostic tests, that are based on either the detection of the immune response of the host e.g., ELISA that detects antibodies directed against *C. burnetii* or the microorganism like Polymerase Chain Reaction (PCR) that detect bacterial DNA, have been used in epidemiological studies for *C. burnetii* to obtain estimates of the incidence and prevalence (Klemmer et al., 2018). However, prevalence estimation depends on the test's sensitivity (*Se*) and specificity (*Sp*), therefore, accurate diagnostic accuracy measures are important.

C. burnetii infection was first described in India in 1952 in a cattle herd and the first human case was reported in 1953 (Kalra and Taneja, 1954). Since then, the disease has been reported in several studies in India and an increasing trend in prevalence is reported, outlining *C. burnetii* as a potential threat to public health (Randhawa et al., 1973; Sodhi et. al., 1980; Vaidya et al., 2010; Keshavamurthy et. al., 2019; Shome et al., 2019; Yadav et al., 2019; Dhaka et al., 2020). Reported prevalence estimates from studies in India, conducted in a frequentist framework, assume that the applied diagnostic tests have perfect *Se* and *Sp* or imperfect, but known, measures of test accuracy. Further, a wide range of hosts, that *C. burnetii* is proven to affect, have been studied to better understand the microorganism's epidemiology (Vaidya et al., 2010).

119 Objective of this study is the diagnostic evaluation of the applied tests to detect *C. burnetii*
 120 infection in cattle and buffalo animals in Punjab, a state in northern India. Since no gold
 121 standard is reported for *C. burnetii*, the study was conducted in a Bayesian framework,
 122 following the STARD-BLCM guidelines (Kostoulas et al., 2017). The STARD-BLCM
 123 checklist is available as a supplementary material (see S1 Appendix).
 124
 125

2. Materials and methods

2.1. Ethics approval

This study was approved by the Institutional Animal Ethical Committee, Guru Angad Dev Veterinary and Animal Sciences University (GADVASU/2017/IAEC/42/02).

2.2. Study design

The study design is presented in detail elsewhere (Keshavamurthy et al., 2019). Briefly, a multi-stage sampling design was performed. Twenty-two villages, one per district, of the state were selected. Further, the number of households sampled in each village was selected, proportional to the number of households in a village. Overall, 179 households (cattle or buffalo herds) participated in the study. We worked towards collecting samples from all the animals for each household. However, many farmers were reluctant to provide samples from some of their animals. We recorded a response rate of 72.5% at the household level and 53.4% at the animal level, respectively (Keshavamurthy et al., 2019).

The sampling unit in this study were cattle and buffaloes. In the analysis cattle and buffaloes were considered as one population, referred as domestic bovine population, because (i) both species are members of Bovidae family (Michelizzi et al., 2010) and (ii) the course of the *C. burnetii* infection is similar in both species (Lucchese et al., 2016). Under this setting, the target population of the study was the domestic bovine population in the Punjab state.

2.3. Sample collection

Blood and genital swab samples were collected from the selected cattle and buffaloes. In addition, milk samples, from lactating female animals, were collected. Blood samples were collected aseptically from the jugular vein. Puncture area was cleaned with 70% alcohol and venipuncture was done using a fresh needle. Approximately 5 ml of blood was withdrawn from each animal in a sterile vacutainer. For the molecular study, genital swabs were collected from both male and female animals using sterile cotton swabs. Vaginal swab samples from female animals were collected by carefully inserting the swab into the vaginal cavity about 10 cm through followed by gently rotating the swab. In males, preputial swabs were collected by swabbing the penile and preputial surface. For the collection of milk samples from the lactating animals, udder and teats were cleaned using germicidal teat dip and three to four streams of milk was discarded before sampling to minimize risk of sample contamination. About 15 ml of milk sample was collected using sterile screw-capped vials.

All samples were transported to the laboratory on the sampling day and stored at -20°C. The sera were separated within 24h in the sterile cryovials before storing at -20°C until screened (Keshavamurthy et al., 2019).

Overall, 610 blood samples, 610 genital swabs and 361 milk samples were collected from the study population. Therefore, the samples were split into two subpopulations; subpopulation-1 (subp_1) includes lactating female animals and subpopulation-2 (subp_2) includes male and non-lactating female animals. Overall, 400 of the sampled animals (65.5.%) were lactating and 221 (36.2%) were pregnant.

2.4. Laboratory tests

Milk samples and genital swabs were screened with PCR to detect bacterial DNA. Specifically, a trans-PCR assay was used to detect *C. burnetii* particles based on two transposon-like repetitive regions of the microorganism: Trans1: 5'-TAT GTA TCC ACC GTA GCC AGT C-3' and Trans2: 5'-CCC AAC AAC ACC TCC TTA TTC-3' (Willems et al., 1994).

The serum samples were screened for *C. burnetii* antibodies using the commercial Q Fever indirect ELISA kit with Phase I and Phase II (ELISA Kit for serodiagnosis of Q Fever in cattle and small ruminants, Monowell, Bio-X Diagnostics, Rochefort, Belgique). The two Phases capture both the acute and chronic infection form. In particular, IgG antibody titers against Phase I antigens are elevated during the acute phase, whereas IgG antibody titers against Phase I and Phase II are elevated during the chronic phase (ELISA Kit for serodiagnosis of Q Fever in cattle and small ruminants, Monowell, Bio-X Diagnostics, Rochefort, Belgique). ELISA quantifies the immune response of the host against *C. burnetii* and does not provide information about the presence or absence of the bacterium. Broadly, serological techniques are useful for screening purposes e.g., monitor the vaccination effectiveness. This is not applicable in India, because a vaccination program for *C. burnetii* does not exist (Dhaka et al., 2020).

2.5. Bayesian latent class analysis

The diagnostic accuracy of the applied tests was estimated using a Bayesian latent class model (BLCM). Traditionally, in the absence of a “gold standard”, latent class models (Hui and Walter, 1980) can be used to obtain unbiased estimates. Over the last decades, Bayesian framework has been applied in latent class analyses, due to their flexibility, incorporation of prior knowledge and software availability (Enøe et al., 2000; Branscum et al., 2005). To ensure

transparency and extrapolation of the study results, the STARD-BLCM reporting guidelines for diagnostic accuracy studies that use BLCMs were followed (Kostoulas et al., 2017).

2.5.1. Definition of infection status

Explicit description of the biological principle of each applied test is crucial towards the structure of any BLCM model. Latent variables are hidden, or unknown and probabilistic estimates can be made for them in conjunction with what the tests actually detect (Walter and Irwig, 1988). In this study, the applied PCR technique in the milk samples (PCR-Milk) and the genital swabs (PCR-Genital) detects bacterial DNA i.e., presence or absence of the *C. burnetii* microorganism and ELISA measures antibodies titers i.e., immune host response (IgG ELISA). Therefore, two distinct biological principles are defined implying different latent states of the sampling unit i.e., it is possible that presence of the bacteria in the host does not always trigger host immune response.

Estimations were based on the cross-classified results (Table 1) of the applied tests in the two subpopulations described above.

Table 1. Cross-classified results of IgG ELISA, PCR-Genital and PCR-Milk

Lactating animals		PCR-Genital ¹				Total
IgG ELISA ³		Positive		Negative		
		PCR-Milk ²		PCR-Milk		
		Positive	Negative	Positive	Negative	
	Positive	1	0	3	23	27
	Negative	0	3	7	324	334
Total		1	3	10	347	361
Non-lactating animals		PCR-Genital				Total
IgG ELISA		Positive		Negative		
	Positive	0		6		6
	Negative	0		243		243
Total		0		249		249

Polymerase Chain Reaction (PCR) in genital swabs

²PCR-Milk: PCR in milk samples

³IgG ELISA: ELISA in serum samples

2.5.2. Model assumptions

BLCM models, in the absence of a gold standard, for Se and Sp estimation can be constructed under different assumptions. An applied set of assumptions adopted by Hui and Walter model (two tests – two populations) state that (i) the population is divided into two or more subpopulations in which two or more tests are evaluated (ii) Se and Sp of each test remain constant across both species and both subpopulations and (iii) all tests are conditionally independent given infection status (Toft et al., 2005). According to the literature, previous Bayesian latent-class analyses for *C. burnetii* infection indicate that the Se and Sp of ELISA do not vary between species (Lucchese et al., 2016). Conditional independence can be assumed, on the basis that ELISA and PCR do not measure similar biological processes (Gardner et al.,

2000). Also, PCR-Milk and PCR-Genital were applied to different organs, therefore, presence (absence) of the infectious agent to one organ does not imply presence (absence) to other organs. Even though, the existence of distinct difference of the true prevalence between the subpopulations is proven to influence the precision of the estimates; this is not applicable in this case, i.e., the two subpopulations have the same true prevalence.

2.5.3. Model description

Bayesian modelling extracts the posterior probability given prior information and the likelihood function. The likelihood is computed through a statistical model for the observed data.

The models for the two subpopulations were structured assuming that the various test combinations follow the multinomial distribution. Specifically,

$$y_subp_1[1:Q, 1:Q, 1:Q] \sim \text{dmulti}(p1[1:Q, 1:Q, 1:Q], n1)$$

$$y_subp_2[1:Q, 1:Q] \sim \text{dmulti}(p2[1:Q, 1:Q], n2)$$

where y_subp_1 and y_subp_2 are the counts of various test combinations, $Q = \{1,2\}$ the dichotomized test result i.e., 1 for positive and 2 for negative, $n1$ & $n2$ the two population sizes and $p1$ & $p2$ the probabilities of observing each test combination.

Based on this notation the frequencies of possible test outcomes can be calculated as followed:

$$p1[1,1,1] = pi * Se_{ELISA} * Se_{Genital}^{PCR} * Se_{Milk}^{PCR} + (1 - pi) * (1 - Sp_{ELISA}) * (1 - Sp_{Genital}^{PCR}) * (1 - Sp_{Milk}^{PCR})$$

$$p1[1,1,2] = pi * Se_{ELISA} * Se_{Genital}^{PCR} * (1 - Se_{Milk}^{PCR}) + (1 - pi) * (1 - Sp_{ELISA}) * (1 - Sp_{Genital}^{PCR}) * Sp_{Milk}^{PCR}$$

$$p2[1,1] = pi * Se_{ELISA} * Se_{Genital}^{PCR} + (1 - pi) * (1 - Sp_{ELISA}) * (1 - Sp_{Genital}^{PCR})$$

$$p2[2,2] = pi * (1 - Se_{ELISA}) * (1 - Se_{Genital}^{PCR}) + (1 - pi) * Sp_{ELISA} * Sp_{Genital}^{PCR}$$

Under this setting, the parameters to be estimated are seven (i.e., Se_{ELISA} , $Se_{PCR-Genital}$, $Se_{PCR-Milk}$, Sp_{ELISA} , $Sp_{PCR-Genital}$, $Sp_{PCR-Milk}$ and pi (true prevalence)), while the degrees of freedom offered by the data are seven. Therefore, identifiability criteria are met and an uniform, noninformative, Beta prior distribution $Be(I, I)$ can be adopted for all parameters of interest. However, degrees of freedom being higher than or equal to the number of parameters to be estimated is a necessary but not always sufficient condition to ensure identifiability. In this analysis, due to the sparsity of the observed data (i.e., zero cell observations for some of the cross-classified results – see Table 1) the ability of the model to estimate the associated parameters diminishes. Hence, informative priors were also introduced.

Prior information was supplied by one of the authors (B.B.S.), an epidemiologist, expert on zoonoses and co-leader of a national project on “Epidemiology, burden and control of zoonotic diseases in India”. Generally, not much is known about the differences in the Se and Sp of similar diagnostic tests in cattle and buffalo populations. A latent class analysis conducted using a Bayesian approach was performed to understand the differences in Se and Sp of ELISA in cattle and goats (Lucchese et al., 2016). The authors reported ELISA Se and Sp values to be 0.97 and 0.98 in cattle and 0.98, 0.83 in goats, respectively (Lucchese et al., 2016). In detail,

estimates of the mean and the 95th percentile for all parameters were provided. Informative prior Beta distributions were calculated using the PriorGen R Package (Kostoulas, 2018) (Table 2).

Table 2. Mean and 95th percentiles for the sensitivity (*Se*) and specificity (*Sp*) priors of PCR and ELISA and the corresponding Beta distributions, *Be(a, b)*. The prior information was provided by one of the co-authors (B.B.S.).

Test	Parameter	Mean (%)	95 th percentile (%)	Be (a, b)
PCR	<i>Se</i> *	75	85	<i>Be(32.58, 10.86)</i>
	<i>Sp</i> **	95	99	<i>Be(40.78, 2.15)</i>
ELISA	<i>Se</i>	97	99	<i>Be(122.51, 3.79)</i>
	<i>Sp</i>	90	95	<i>Be(11.74, 1.3)</i>

2.5.4. Markov Chain Monte Carlo convergence and software

Models were run in the freeware program OpenBUGS (Spiegelhalter et al., 2007). Parameter estimates were based on analytical summaries of 100,000 iterations of two chains after a burn-in phase of 5000 iterations. The tools described in Toft et al., 2007, were monitored to ensure occurrence of convergence. The OpenBUGS code for the final model is available as a supplementary material (see S2 Appendix).

2.5.5. Sensitivity analysis

The influence of the data and the priors to the posterior estimates was examined by running the same model without informative prior values also. Different models under different set of assumptions were constructed to inspect the validity of the applied set of assumptions.

308 Specifically, to ensure constant Se and Sp across species models to examine this assumption
 309 were structured.
 310
 311 Model selection was based on the DIC (Deviance Information Criterion) dialog box in
 312 OpenBUGS environment. The model with the smallest DIC is the model that best fits the data
 313 i.e., the model that would best predict a replicate dataset of the same structure as the currently
 314 observed (Spiegelhalter et al., 2002).

3. Results

The posterior medians and 95% probability intervals (PIs) for the Se and Sp of each diagnostic test are summarized in Table 3.

Table 3. Posterior medians and 95% probability intervals (PrIs) for the Se and Sp of each diagnostic test using informative Beta prior distributions for the Se and Sp of each diagnostic test described in Table 2.

Test	Parameter	Posterior medians and 95% PrIs
IgG ELISA	Se	0.97 (0.93; 0.99)
	Sp	0.95 (0.93; 0.97)
PCR-Genital	Se	0.7 (0.55; 0.82)
	Sp	0.99 (0.98; 1)
PCR-Milk	Se	0.76 (0.62; 0.87)
	Sp	0.98 (0.96; 0.99)

IgG ELISA showed the highest Se with median 0.97 (95% PIs: 0.93; 0.99) compared to PCR-Milk 0.76 (0.62; 0.87) and PCR-genital 0.7 (0.55; 0.82). The Sp s of all tests were high, with PCR-Genital recording the highest Sp median of 0.99 (0.98; 1), while the Sp of PCR-Milk and IgG ELISA were 0.98 (0.96; 0.99) and 0.95 (0.93; 0.97) respectively.

The acquired estimates without informative prior distributions are shown in S1 Table. Under this setting, the Sp estimates are similar in both scenarios. However, the Se estimates were lower; the Se of IgG ELISA was 0.63 (0.17; 0.98), PCR-Genital 0.18 (0.02; 0.77) and PCR-Milk was 0.6 (0.13; 0.98).

338 Sensitivity analysis was performed to validate the assumption of constant accuracy across both
 339 species. Parameter estimates produced by applying a model only to cattle were not substantially
 340 different, validating the assumption of constant accuracy across both species (S2 Table).

341
 342 The model described in Section 2.5.3. was the one with the smallest DIC compared to the
 343 ones that were introduced in the Sensitivity Analysis section.

344

4. Discussion

In this study, BLCMs were used to estimate the Se and Sp of a trans-PCR applied in genital swabs and milk samples to detect *C. burnetii* DNA and an ELISA applied in serum samples that detects antibodies against *C. burnetii*, in cattle and buffaloes in Punjab, India. BLCMs account for the absence of a gold standard and allow parameter estimation merging two components (i) model structure, based on the observed data and (ii) incorporation of prior information (Kostoulas et al. 2017).

The study results show that all three tests are highly specific, with PCR-Genital yielding the higher Sp [0.99 (0.98; 1)], followed by PCR-Milk [0.98 (0.96; 0.99)] and IgG ELISA [0.95 (0.93; 0.99)]. On the other hand, IgG ELISA has the highest Se [0.97 (0.93; 0.99)], followed by PCR-Milk [0.76 (0.62; 0.87)] and PCR-Genital [0.7 (0.55; 0.82)]. This seems reasonable, as serological techniques are in general more sensitive than molecular diagnostic techniques, due to several reasons e.g., cross-reaction (Joseph et al., 1995). Overall, the posterior medians and 95% PIs for the PCR-Milk and PCR-Genital are comparable, indicating that both tests have the same diagnostic accuracy. Further, the Sps for these two tests, are not “prior-driven/dependent”, since both under informative and uniform, noninformative priors the posterior estimates are approximately the same. However, the reported estimates for the Ses of PCR-Milk, and especially PCR-Genital, seem to differ under noninformative and informative prior distributions [informative prior distributions; PCR-Milk 0.76 (0.62; 0.87) and PCR-Genital 0.7 (0.55; 0.82); noninformative prior distributions; PCR-Milk 0.6 (0.13; 0.98) and PCR-Genital 0.18 (0.02; 0.77)]. This is mainly due to the scarcity of the data i.e., small number of animals both positive to PCR-Milk and PCR-Genital. Further, the reported medians for the Se of PCR-Milk and PCR-Genital, using informative prior distributions, are included in the 95% PIs for the Ses under uniform, noninformative prior distributions. Implementation of informative prior

distributions allows shrinkage of the 95% PI for the Ses. Therefore, the final Se estimates for the two PCRs can be considered reliable. As far as IgG-ELISA, even though, a high Sp is recorded, the Se seems to be “prior-driven/dependent”. However, the information provided by the data may not be enough, since only thirty-three animals were tested positive in IgG-ELISA [1 PCR-Milk+, PCR-Genital+, 3 PCR Milk+, PCR-Genital-, 23 PCR-Milk-, PCR-Genital-, 6 PCR-Genital-]. Therefore, IgG-ELISA Se posterior estimates using uniform, noninformative prior distribution cannot be considered reliable. Again, the median for the Se of IgG-ELISA, using informative priors is included in the 95% PI for the Se under a uniform, noninformative prior setting.

Studies on validation of diagnostic tests for *C. burnetii* infection using BLCMs have been conducted in cattle, goats, sheep etc. (Luchesse et al., 2016; Paul et al., 2013). The reported Se and Sp of the tests in our study are comparable between studies and similar between different species e.g., sheep and goats. (Abiri et al., 2019). The posterior medians and 95% PIs for the diagnostic characteristics of IgG-ELISA are similar to those reported in the literature and comparable with the estimates provided by the commercial ELISA kits manufacturers used for detection of antibodies against *C. burnetii* in serum samples from cattle (Luchese et al., 2016; Serrano-Pérez et al., 2015). The PCR method applied in milk samples has been evaluated in cattle (Nusinovici et al., 2015), in a Bayesian framework. The results from our study are similar with the ones reported in Nusinovici et al. (2015). The PCR method applied in genital swabs in cattle has not been evaluated; instead, PCR has been used for bacterial DNA detection in the farm environment (Nusinovici et al., 2015). On the other hand, PCR-Genital and PCR-Milk has been described in the sheep and goats (Abiri et al., 2019). The Sp for both PCRs and for the Se of PCR-Genital are similar, while the reported median and 95% PI for the Se of PCR-Milk in Abiri et al. (2019) is lower [0.42 (0.32; 0.59)].

Conditional independence between PCR-Milk and PCR-ELISA was assumed, since, the method was applied to different organs, even though it is based on the same biological principle. Further, primary shedding route has not been identified for *C. burnetii* and isolation of the bacterium from more than two organs is rare (Table 1) (Guatteo et al., 2006), i.e., presence (absence) of the infectious agent in the genital tract does not imply presence (absence) to milk. The conditional independence assumption for a PCR method applied in milk and vaginal secretions in sheep and goat samples to detect *C. burnetii* has been adopted (Abiri et al., 2019). Therefore, this assumption can be considered valid. Moreover, the specified model has seven degrees of freedom and seven parameters of interest. If conditional independence was not assumed, then two extra parameters for the covariance terms will be added, and the identifiability criteria will not be met (degrees of freedom higher than or equal to the number of parameters of interest), hence, our model would not converge. Therefore, introducing two covariance terms, accounting for conditional dependence between PCR-Milk & PCR-Genital would result in an unidentifiable model. Even though, in our case we adopt the results using informative prior distributions, we do so, due to the scarcity of the data (zero cell observations for animals positive to IgG-ELISA, PCR-Milk, PCR-Genital). Thus, informative prior distributions are added, instead of noninformative, to overcome the sparsity of the data. As shown in S2 Table the specified model using uniform, noninformative priors converges, but results to 95% PIs with high width.

Furthermore, the course of infection in the two species under investigation was considered similar, hence the diagnostic tests properties were assumed constant across species. Applying the model only to the cattle population yields posterior estimates similar to the ones after applying the final selected model (S2 Table). In India, risk factor investigation studies present

contrasting results in the risk of occurrence of *C. burnetii* infection in bovine populations. A recent study conducted in Bihar and Assam states of India reported higher risk of infection for buffaloes than in cattle (28.0% compared to 13.6%, $p=0.042$) at the species level (Shome et al., 2019). However, only 25 buffaloes as compared to 719 cattle were included in this study (Shome et al, 2019). On the other hand, similar investigations in Punjab reported that cattle (adjusted Odds Ratio 3.37, 95% Confidence interval 1.23-9.20, $p=0.02$) were associated with larger odds of *C. burnetii* positive animal status than buffaloes (Keshavamurthy et al., 2020). Based on these results, our assumption that the course of infection and disease occurrence does not vary much in cattle and buffaloes at the species level is valid and there are other factors that need further investigation.

In this analysis, usage of informative priors improved the fit of the model, as indicated by the DIC. This is essential to account for the sparsity of the observed data. Assuming noninformative priors generates estimates with very wide PIs that do not allow safe conclusions about the tests' accuracy.

5. Conclusion

This study was conducted to estimate the *Se* and *Sp* of three tests used for *C. burnetii* detection in Punjab, India. IgG ELISA achieved the highest *Se*, while PCR-Genital had the highest *Sp*. Using BLCMs, none of the applied tests showed perfect *Se* and *Sp*, and therefore, could not be used alone to diagnose *C. burnetii* infection in domestic bovine populations. Further, it is proven that the diagnostic accuracy of the tests does not vary between the two bovine species. In conclusion, better information on *C. burnetii* infection can be provided using a combination of diagnostic tests based on different biological principles i.e., detection of bacterial DNA and immune host response.

Acknowledgments

The India Council for Agricultural Research, Government of India, New Delhi through the “Outreach Programme for Zoonotic Investigations” are thanked for their financial support of the present study.

Declarations of interest

None

References

- Abiri Z, Khalili M, Kostoulas P, Sharifi H, Rad M, Babaei H. Bayesian estimation of sensitivity and specificity of a PCR method to detect *Coxiella burnetii* in milk and vaginal secretions in sheep and goat samples. J. Dairy Sci. 2019;102: 4954-4959.
- Angelakis E, Raoult D. Q fever. Vet. Mic. 2010;140: 297-309.
- Arricau-Bouvery N, Rodolakis A. Is Q fever an emerging or re-emerging zoonosis?. Vet. Res. 2005;36: 327-349.
- Babudieri B. Q fever: a zoonosis. Adv. Vet. Sci. 1959;5: 81
- Branscum AJ, Gardner IA, Johnson WO. Estimation of diagnostic-test sensitivity and specificity through Bayesian modeling. Pre. Vet. Med. 2005;68: 145-163.
- Burnet FM, Freeman M. Experimental Studies on the Virus of "Q" Fever. Med. J. Aus. 1937;2: 299-305.
- Derrick EH. "Q" Fever, a New Fever Entity: Clinical Features, Diagnosis and Laboratory Investigation. Med. J. Aus. 1937;2: 281-299.
- Dhaka P, Malik SVS, Yadav JP, Kumar M, Barbuddhe SB, Rawool DB. Apparent prevalence and risk factors of coxiellosis (Q fever) among dairy herds in India. PloS One, 2020;15: e0239260.

483

484 EFSA Panel on Animal Health and Welfare (AHAW). Scientific Opinion on Q fever. EFSA

485 Journal. 2010;8: 1595.

486

487 Enøe C, Georgiadis MP, Johnson WO. Estimation of sensitivity and specificity of diagnostic

488 tests and disease prevalence when the true disease state is unknown. Pre. Vet. Med. 2000;45:

489 61-81.

490

491 Gardner IA, Stryhn H, Lind P, Collins MT. Conditional dependence between tests affects the

492 diagnosis and surveillance of animal diseases. Prev. Vet. Med. 2000;45: 107-122.

493

494 Guatteo R, Beaudeau F, Berri M, Rodolakis A, Joly A, Seegers H. Shedding routes of *Coxiella*

495 *burnetii* in dairy cows: implications for detection and control. Vet. Res. 2006;37: 827-833.

496

497 Hilbink F, Penrose M, Kovacova E, Kazar J.. Q fever is absent from New Zealand. Int. J.

498 Epidemiol. 1993;22: 945-949.

499

500 Hui, S.L., Walter, S.D., 1980. Estimating the error rates of diagnostic tests. Biometrics, 167-

501 171.

502

503 Joseph L, Gyorkos TW, Coupal L. Bayesian estimation of disease prevalence and the

504 parameters of diagnostic tests in the absence of a gold standard. Am. J. Epidemiol. 1995;141:

505 263-272.

506

507 Kalra SL, Taneja BL. Q fever in India: a serological survey. Indian J. Med. Res. 1954;42: 315-
508 318.
509

510 Kaplan MM, Bertagna P. The geographical distribution of Q fever. Bull. World Health Organ.
511 1955;13: 829.
512

513 Keshavamurthy R, Singh BB, Kalambhe DG, Aulakh RS, Dhand NK. Prevalence of *Coxiella*
514 *burnetii* in cattle and buffalo populations in Punjab, India. Prev. Vet. Med. 2019;166: 16-20.
515

516 Keshavamurthy R, Singh BB, Kalambhe, DG, Aulakh RS, Dhand NK. Identification of risk
517 factors associated with *Coxiella burnetii* infection in cattle and buffaloes in India. Prev. Vet.
518 Med. 2020;181: 105081.
519

520 Klemmer J, Njeru J, Emam A, El-Sayed A, Moawad AA, Henning K, et al. Q fever in Egypt:
521 Epidemiological survey of *Coxiella burnetii* specific antibodies in cattle, buffaloes, sheep, goats
522 and camels. PloS One. 2018;13: e0192188.
523

524 Kostoulas P. PriorGen: Generates Prior Distributions for Proportions. R 364 Packag. version,
525 2018;1: 365.
526

527 Kostoulas P, Nielsen SS, Branscum AJ, Johnson WO, Dendukuri N, Dhand NK, et al. STARD-
528 BLCM: Standards for the Reporting of Diagnostic accuracy studies that use Bayesian Latent
529 Class Models. Prev. Vet. Med. 2017;138: 37-47.
530

Lang GH, Marrie TJ. Coxiellosis (Q fever) in animals. In Q Fever: the disease. CRC Press, New York, USA; 1990. pp. 23-48.

Lucchese L, Capello K, Barberio A, Ceglie L, Guerrini E, Zuliani F, et al. Evaluation of serological tests for Q fever in ruminants using the latent class analysis. Clin. Res. Infect. Dis. 2016;3: 1-5.

Marrie TJ. Coxiella burnetii pneumonia. Eur. Respir. J. 2003;21: 713-719.

Maurin M, Raoult DF. Q fever. Am. Soc. Microbiol. 1999;12: 518-553.

McDade JE, Marrie TJ. Historical Aspects of Q Fever. In Q Fever: the disease. CRC Press, New York, USA; 1990. pp. 5-21.

Michelizzi VN, Dodson MV, Pan Z, Amaral MEJ, Michal JJ, McLean DJ, et al. Water buffalo genome science comes of age. Int. J. Biol. Sci. 2010;6: 333.

Nusinovici S, Madouasse A, Hoch T, Guatteo R, Beaudeau F. Evaluation of two PCR tests for *Coxiella burnetii* detection in dairy cattle farms using latent class analysis. PloS One. 2015;10: e0144608.

Paul S, Toft N, Agerholm JS, Christoffersen AB, Agger JF. Bayesian estimation of sensitivity and specificity of Coxiella burnetii antibody ELISA tests in bovine blood and milk. Prev. Vet. Med. 2013;109: 258-263.

556 Randhawa AS, Dhillon SS, Jolley WB. Serologic prevalence of Q fever in the state of Punjab,
557 India. Am. J. Epidemiol. 1973;97: 131-134.

558

559 Raoult D, Marrie T. Q fever. Clin. Infect. Dis. 1995;489-495.

560

561 Roest HI, Bossers A, van Zijderveld FG, Rebel JM. Clinical microbiology of *Coxiella burnetii*
562 and relevant aspects for the diagnosis and control of the zoonotic disease Q fever. Vet. Q.
563 2013;33: 148-160

564

565 Serrano-Pérez B, Almería S, Tutusaus J, Jado I, Anda P, Monleón E, Badiola J, et al. *Coxiella*
566 *burnetii* total immunoglobulin G, phase I and phase II immunoglobulin G antibodies, and
567 bacterial shedding in young dams in persistently infected dairy herds. J. Vet. Diagn. Invest.
568 2015;27: 167-176.

569

570 Shome R, Deka RP, Miles L, Sahay S, Grace S, Lindahl JF. *Coxiella* seroprevalence and risk
571 factors in large ruminants in Bihar and Assam, India. Acta tropica, 2019;194: 41-46.

572

573 Sodhi SS, Joshi DV, Sharma DR, Baxi KK. Seroprevalence of brucellosis and Q fever in dairy
574 animals. Zbl. Vet. Med. B. 1980;27: 683-685.

575

576 Spiegelhalter DJ, Best NG, Carlin BP, Van Der Linde A. Bayesian measures of model
577 complexity and fit. J. R. Stat. Soc. Series b (Stat Methodol). 2002;64: 583-639.

578

579 Spiegelhalter D, Thomas A, Best N, Lunn D. OpenBUGS user manual, version 3.0. 2. MRC
580 Biostatistics Unit, Cambridge. 2007.

581

582 Sprong H, Tjisse-Klasen E, Langelaar M, De Bruin A, Fonville M, Gassner F, et al. Prevalence
583 of *Coxiella burnetii* in ticks after a large outbreak of Q fever. *Zoonoses Public Health*. 2012;59:
584 69-75.

585

586 Toft N, Jørgensen E, Højsgaard S, Diagnosing diagnostic tests: evaluating the assumptions
587 underlying the estimation of sensitivity and specificity in the absence of a gold standard. *Prev.*
588 *Vet. Med.* 2005;68: 19-33.

589

590 Toft N., Innocent GT, Gettinby G, Reid SW. Assessing the convergence of Markov Chain
591 Monte Carlo methods: an example from evaluation of diagnostic tests in absence of a gold
592 standard. *Prev. Vet. Med.* 2007;79: 244-256.

593

594 Yadav JP, Malik SVS, Dhaka P, Kumar M, Bhoomika S, Gourkhede D, et al. Seasonal variation
595 in occurrence of *Coxiella burnetii* infection in buffaloes slaughtered in India. *Biol. Rhythm Res.*
596 2019;1-7.

597

598 Vaidya VM, Malik SVS, Bhilegaonkar KN, Rathore RS, Kaur S, Barbuddhe SB. Prevalence of
599 Q fever in domestic animals with reproductive disorders. *Comp. Immunol. Microbiol. Infect.*
600 *Dis.* 2010;33: 307-321.

601

602 Waag DM. *Coxiella burnetii*: host and bacterial responses to infection. *Vaccine*, 2007;25:
603 7288-7295.

604

Walter SD, Irwig LM. Estimation of test error rates, disease prevalence and relative risk from misclassified data: a review. J. Clin. Epidemiol. 1988;41: 923-937.

Willems H, Thiele D, Frölich-Ritter R, Krauss H. Detection of *Coxiella burnetii* in cow's milk using the polymerase chain reaction (PCR). J. Vet. Med. Series B. 1994;41: 580-587.

World Organisation for Animal Health (OIE) Terrestrial Manual. Chapter 3.1.16 Q Fever. 2018; Available from: https://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/3.01.16_Q_FEVER.pdf

Supporting information

S1 Table. S1_Table_file. Posterior medians and 95% PrIs for the sensitivity (Se) and specificity (Sp) of each diagnostic test using noninformative Beta prior distributions for all parameters of interest.

S2 Table. S2_Table_file. Posterior medians and 95% PrIs for the Se and Sp of each diagnostic test applying the model only to cattle population, assuming (i) informative Beta prior distributions for the Se and Sp of each diagnostic test and (ii) noninformative Beta prior distributions for all parameters of interest.

S1 Appendix. STARD-BLCM Checklist.

S2 Appendix. OpenBUGS Code