Evolutionary rescue in one dimensional stepping stone models

Matteo Tomasini^{*, †, ‡, 1} and Stephan Peischl^{*, ‡}

^{*}Interfaculty Bioinformatics Unit, University of Bern, 3012 Bern, Switzerland

[†]Computational and Molecular Population Genetics Laboratory, Institute of Ecology and Evolution, University of Bern, 3012 Bern, Switzerland

[‡]Swiss Institute for Bioinformatics, 1015 Lausanne, Switzerland

¹Current address: Department of Marine Sciences and Linnaeus Centre for Marine Evolutionary Biology, University of Gothenburg, 405 30 Gothenburg, Sweden

July 27, 2021

Abstract

Genetic variation and population sizes are critical factors for successful adaptation to novel environmental conditions. Gene flow between sub-populations is a potent mechanism to provide such variation and can hence facilitate adaptation, for instance by increasing genetic variation or via the introduction of beneficial variants. On the other hand, if gene flow between different habitats is too strong, locally beneficial alleles may not be able to establish permanently. In the context of evolutionary rescue, intermediate levels of gene flow are therefore often optimal for maximizing a species chance for survival in metapopulations without spatial structure. To which extent and under which conditions gene flow facilitates or hinders evolutionary rescue in spatially structured populations remains unresolved. We address this question by studying the differences between evolutionary rescue in the island model and in the stepping stone model in a habitat gradually deteriorating. We show that spatial structure affects the interactions between positive and negative effects of gene flow on evolutionary rescue, but that ultimately the most important factor is the gene flow between deteriorated and intact environment, as opposed to gene flow between subpopulations. As such,

many spatially complex scenarios can be understood in terms of two big demes under divergent selection exchanging individuals. As an implication, the insights gained from two-deme models can apply to spatially explicit metapopulations and improve our understanding of evolutionary rescue in fragmented habitats.

Introduction

Recent environmental or climatic changes have severe consequences on the survival of many species [Thomas et al., 2004]. Rapid and severe environmental change might cause extinction of resident populations due to genetic maladaptation to the new environmental conditions [Cahill et al., 2013]. A population may survive such environmental change if it adapts quickly enough via standing genetic variation or *de novo* mutations, a process that has been coined evolutionary rescue [Gomulkiewicz and Holt, 1995, Alexander et al., 2014, Carlson et al., 2014]. Theoretical studies of the mechanisms that can facilitate or hinder survival of populations experiencing environmental change can provide critical insights in the context of adaption and survival in the face of global climatic change, but also in medically relevant scenarios such as the evolution of antibiotic or other treatment resistance (*e.g.* Alexander et al. [2014], Carlson et al. [2014]).

It has been demonstrated theoretically as well as experimentally that evolutionary rescue in single, isolated populations depends strongly on two factors: initial population size and standing genetic variation [Lynch, 1993, Bell and Gonzalez, 2009, Samani and Bell, 2010, Bell and Gonzalez, 2011, Bell, 2013, Ramsayer et al., 2013, Carlson et al., 2014]. In large populations or in populations with higher mutation rates mutations that ultimately rescue a population have higher chances to arise and establish before the population goes extinct [Lynch, 1993, Gomulkiewicz and Holt, 1995, Bell, 2013]. Similarly, initially high standing genetic variation increases the chance that a mutation that can save the population may already be present in the population [Gomulkiewicz and Holt, 1995, Barrett and Schluter, 2008, Vander Wal et al., 2013]. Other factors, such as the competition between individuals, the rate of environmental decay or gene flow from other populations can have more subtle effects and affect the chance of rescue positively or negatively [Uecker et al., 2014, Marrec and Bitbol, 2020].

Evolutionary rescue has been thoroughly studied in idealized laboratory conditions and key theoretical

predictions have been confirmed [Carlson et al., 2014]. However, these idealized laboratory conditions are unlikely to reflect the spatio-temporal demographics of natural populations or the intricate spatial and population structure within and across human bodies during the evolution of antibiotic or treatment resistance [Bell, 2017]. For instance, in the face of structured populations it has been demonstrated both experimentally [Bell and Gonzalez, 2009, 2011, Gonzalez and Bell, 2013] and theoretically [Uecker et al., 2014, Czappon et al., 2021] that gene flow between sub-populations can facilitate evolutionary rescue. The benefits of spatial structures and dispersal to adaptation and evolutionary rescue have also been demonstrated in other contexts. For example, dispersal can help to spread beneficial mutations causing rapid population differentiation [Feder et al., 2019], and heterogeneity in environmental conditions caused by imperfect drug penetration allowed the evolution of multi-drug resistance [Moreno-Gamez et al., 2015]. In the context of antibiotic resistance evolution, interesting examples come from *Pseudomonas aeruginosa* where the formation of biofilms are hallmarks of disease progression and chronic infections [Botelho et al., 2019]. Experimental studies have shown that the spatial structure and environmental heterogeneity in biofilms may also contribute to an increased likelihood of resistance evolution as compared to well-mixed populations [Ahmed et al., 2018].

First steps towards understanding how dispersal can help beneficial mutations to cause evolutionary rescue were done by Uecker et al. [2014] (in an island model) and by Tomasini and Peischl [2020] (in a two-deme model). A critical assumption of these modelling attempts is that gene flow occurs at the same rate between any pair of demes. Experimental and theoretical work suggests, however, that the spatial organization of sub-populations, as well as biased dispersal, modulate the effect of gene flow [Bell and Gonzalez, 2011, Gonzalez and Bell, 2013, Czappon et al., 2021]. Whether and how spatial structure might change our understanding of the role of gene flow during adaptation to severe environmental change remains an unresolved question.

In this work we extend the work of Uecker et al. [2014] (in an island model) and Tomasini and Peischl [2020] to study the role of gene flow on evolutionary rescue in spatially extended populations using stochastic simulations. In particular, we contrast two extreme cases of gene flow: long-range migration (*i.e.* the island model [Wright, 1943], where migrants are exchanged among all demes at equal rates) and short-range migration (*i.e.* the stepping stone model [Kimura and Weiss, 1964], where migrants move

Figure 1: In figure A, a schematic of the island model (local migration), in figure B a schematic of the stepping stone model (global migration). Schematic for $D = 4$.

only between neighboring demes).

We find that short- and long-range dispersal can both have a beneficial effect on evolutionary rescue, and that gene flow is beneficial for evolutionary rescue under quite similar conditions. More precisely, under certain conditions, we find that the island model and the stepping stone model appear equivalent if we consider an effective migration rate that quantifies migration between deteriorated and non-deteriorated areas of the environment. If rescue mutations are strongly deleterious in one environment but not (close to) lethal, we find however qualitative differences between the two models. This happens because positive and negative effects of gene flow dominate for different migration rates in different scenarios. Our findings suggest that the number of individuals flowing between deteriorated and non-deteriorated parts of the environment is a key quantity in spatially or otherwise structured populations, rather than the rate of gene flow between demes.

Materials and Methods

Model

We consider a structured population consisting of D demes organized along a one-dimensional lattice (see figure 1). Demes are labeled $i = 1, \dots, D$ and are filled with haploid wild-type individuals with discrete and non-overlapping generations. Each deme can be in one of two environmental states: the

original unperturbed state (or “old” state) and the deteriorated state (or “new” state). In our model, initially the whole habitat is at demographic equilibrium in the original environment. For the sake of explanation, we assume deterioration from left to right (see figure 1) although the model is independent of the direction of deterioration. Every θ generations the leftmost deme in the unperturbed state switches to the deteriorated state such that the whole habitat gradually deteriorates at a constant rate of one deme each θ generations. In the deteriorated part of the environment, growth rates of wild-type individuals are negative but beneficial mutations can arise and save the population from extinction by restoring positive growth rates.

Population size in deme i at time t is denoted $N_i(t)$ ($i \in \{1, \dots, D\}$, $t = 0, 1, 2, \dots$). In the original environment, density regulation is implemented such that all demes are at carrying capacity, κ , at the end of each generation. This model of instantaneous growth or culling is a good approximation to logistic density regulation if growth rates are sufficiently large relative to migration rates [Beverton and Holt, 1957, Tomasini and Peischl, 2020]. Initially all demes are at carrying capacity such that $N_i(0) = \kappa$, for all i . We define the total carrying capacity of the habitat as $K_{\text{tot}} = D\kappa$. We set $K_{\text{tot}} = 2 \times 10^4$ through the rest of the paper (but see [Carlson et al., 2014] for more comprehensive studies of evolutionary rescue for different population sizes). This means that varying D corresponds to varying κ (as the two are inversely proportional). We call m the forward migration rate, that is the proportion of individuals in each deme that migrate [Karlin, 1982, Vuilleumier et al., 2010]. We consider two modes of migration: (i) global migration, allowing an individual to migrate to any deme in the habitat (the island model, Wright [1943]; figure 1A), and (ii) local migration, that allows an individual to migrate only between neighboring demes (the stepping stone model, Kimura and Weiss [1964]; figure 1B). The boundaries of the habitat (to the left of the first deme and to the right of the last deme) are reflecting, that is, individuals cannot leave the habitat.

Fitness is determined by a single locus with two alleles: a wild-type allele and a mutant allele. We assume that each individual has Poisson-distributed offspring with the mean proportional to its fitness. In the original environment, fitness of wild-type individuals is set to $w_{\text{wt}}^{(o)} = 1$ and the fitness of a mutant is $w_{\text{mu}}^{(o)} = 1 - s$, where $0 < s < 1$ is the selection coefficient against the mutant. Mutations are therefore counter-selected in the original habitat. Absolute fitness in the original habitat is then

given by $w_l^{(o)}\kappa/N'_i(t)$ ($l \in \{\text{wt}, \text{mu}\}$), where $N'_i(t)$ is the population size after selection and migration but before density regulation. Mutations occur with probability u per generation and we ignore back mutations. Previous work has shown that the probability of rescue depends monotonically on the product of population size and mutation rate [Orr and Unckless, 2008], hence for the rest of this work, we set $u = 1/K_{\text{tot}}$ such that on average a single mutation occurs per generation in the whole habitat. Because our focus is to study the interaction of gene flow, spatial structure and demographic dynamics on evolutionary rescue, we choose a simple genetic architecture that allows us to investigate our model in detail without having to account for clonal interference [Gerrish and Lenski, 1998], the distribution of fitness effects [Anciaux et al., 2018], or recombination and multi-step rescue [Uecker and Hermisson, 2016, Osmond et al., 2020].

At a time $t = 0$, the first (*i.e.* leftmost in figure 1) deme ($i = 1$) switches its environmental state to the deteriorated environment. In a deteriorated deme, wild-type individuals have an absolute fitness of $w_{\text{wt}}^{(n)} = 1 - r$, with $0 < r < 1$, and the resident population starts to decline. In the new environment, the mutant has an absolute fitness $w_{\text{mu}}^{(n)} = 1 + z$, with $z > 0$. We generally chose a value of z large enough to just restore positive growth after deterioration. The relative fitness of mutants is then $(1 + z)/(1 - r) = 1 + (r + z)/(1 - r)$. In the range of parameters that we chose, $(r + z)/(1 - r)$ is often larger than the cost of having the mutation in the non-deteriorated area, s – as we should expect for example with mutations involved with antibiotic resistance (see *e.g.* Melnyk et al. [2015]). For example, when $z = 0.02$ and $r = 0.5$, which we use for the discussion of most of our results, the relative fitness of a mutant in a deteriorated deme is larger than 1, and thus always larger than s . At time $t = \theta$, deme $i = 2$ (at that point the leftmost deme in the old state) deteriorates, at $t = 2\theta$ the third deme, and so on until the whole habitat is deteriorated. We call the period between two subsequent deterioration events an epoch and θ is the length of an epoch. The deterioration of the whole habitat takes $\Theta = (D - 1)\theta$ generations (see table 1 for a full list of all parameters of the model).

Simulations

We performed stochastic simulations. At the beginning of the simulation, D demes are filled each with κ individuals ($K_{\text{tot}} = \kappa D = 2 \times 10^4$). Before any deterioration occurs, we leave a burn-in time of 500

Table 1: **List and description of all parameters**

Parameter	Description
D	Number of demes
$N_i(t)$	Number of individuals in deme i
κ	Carrying capacity per deme
$K_{\text{tot}} = \kappa \cdot D$	Total carrying capacity of the habitat, fixed at 2×10^4 through the paper
m	Rate of migration per individual
$w_{\text{wt}}^{(o)} = 1$	Fitness of a wild-type individual in the old environment
$w_{\text{wt}}^{(n)} = 1 - r$	Fitness of a wild-type individual in the new environment
$w_{\text{mu}}^{(o)} = 1 - s$	Fitness of a mutant individual in the old environment
$w_{\text{mu}}^{(n)} = 1 + z$	Fitness of a mutant individual in the new environment
$0 < s \leq 1$	Disadvantage against a mutant copy in the old environment
$0 < z \ll 1$	Advantage of a mutant copy in the new environment
$0 < r < 1$	Stress against the wild-type population in the new environment
θ	Speed of deterioration of the environment, length of an epoch
$\Theta = \theta(D - 1)$	Deterioration time of the whole habitat
$u = 1/K_{\text{tot}}$	Mutation probability (wild type to mutant)

generations to generate standing genetic variation.

Reproduction We implement reproduction and selection by sampling from a Poisson distribution with the mean of the distribution given by the absolute fitness of wild-type and mutant individuals. Thus the number of wild-type and mutant individuals (here simplified with the subscript $l \in \{\text{wt}, \text{mu}\}$) changes as

$$N_i^l \rightarrow N_i^{(l,1)} \sim \text{Pois}(N_i^l \cdot w_l^i),$$

where the superscript (1) represents the first step of the generation (reproduction). For simplicity, in this section we used w_l^i as opposed to $w_l^{(o)}$ or $w_l^{(n)}$, as the fitness depends on the state of deme i – *e.g.* at time $0 < t < \theta$, $w_{\text{mu}}^1 = 1 + z$ and $w_{\text{mu}}^2 = 1 - s$.

Mutation After reproduction, we implement mutation as a binomial process with sampling probability

$\pi = u$:

$$N_i^{(\text{mu},1)} \rightarrow N_i^{(\text{mu},2)} \sim \text{Bin}(N_i^{\text{wt},1}, u) + N_i^{(\text{mu},1)},$$

$$N_i^{(\text{wt},1)} \rightarrow N_i^{(\text{wt},2)} = N_i^{(\text{wt},1)} - (N_i^{(\text{mu},2)} - N_i^{(\text{mu},1)}),$$

where the superscript ⁽²⁾ represents the second step of the generation (mutation). For wildtypes, the term $N_i^{(\text{mu},2)} - N_i^{(\text{mu},1)}$ is simply the number of new mutants generated via binomial sampling (and thus removed from the wildtype pool).

Migration in the island model In the island model, the total number of migrants is determined in each deme with a binomial process with sampling probability $\pi = m$, then the whole pool of migrants is re-distributed over all demes with a multinomial distribution. The process is done in the same way separately for mutants and wild types, but for simplicity we will skip the type subscript l . For each deme i , the number of migrants $N_i^{(\text{emigr})}$ is given by

$$N_i^{(\text{emigr})} \sim \text{Bin}(N_i^{(2)}, m) ,$$

and the total number of individuals in the pool is

$$N^{(\text{pool})} = \sum_i N_i^{(\text{emigr})} .$$

The individuals in the migrant pool, $N^{(\text{pool})}$, are then redistributed multinomially, with each individual having the probability $1/D$ of migrating to each deme.

$$N_i^{(\text{immigr})} \sim \text{Mul}(N^{(\text{pool})}, 1/D) .$$

Finally, for each type l (subscript omitted), the number of individuals in each deme after migration is

$$N_i^{(2)} \rightarrow N_i^{(3)} = N_i^{(2)} - N_i^{(\text{emigr})} + N_i^{(\text{immigr})} ,$$

where the superscript ⁽³⁾ represents the third step in the generation (migration).

Migration in the stepping stone model In the stepping stone model, individuals migrating to the right and individuals migrating to the left are picked as a binomial process with probability $\pi = m/2$

(subscript l omitted):

$$N_i^{(\text{migr, left})} \sim \text{Bin}(N_i^{(2)}, m/2),$$

$$N_i^{(\text{migr, right})} \sim \text{Bin}(N_i^{(2)}, m/2).$$

Finally, the number of individuals in deme i after migration for each type is updated as

$$N_i^{(2)} \rightarrow N_i^{(3)} = N_i^{(2)} - N_i^{(\text{migr, right})} - N_i^{(\text{migr, left})} + N_{i-1}^{(\text{migr, right})} + N_{i+1}^{(\text{migr, left})}.$$

Note that in our formulation the walls of the habitat are reflecting, thus in the first deme $N_i^{(\text{migr, left})} = 0$ and in the last deme $N_i^{(\text{migr, right})} = 0$.

Density regulation The last phase of the life cycle is density regulation, and it only occurs in non-deteriorated demes. We define a new population of K_i individuals, where $K_i \sim \text{Pois}(\kappa)$, and we sample the number of mutants and wildtypes according to the frequencies after migration, $N^{(l,3)} / (N^{(\text{mu},3)} + N^{(\text{wt},3)})$.

Burn-out phase After the whole habitat is deteriorated, we let the system evolve for 500 more generations. The expected extinction time in the absence of evolutionary rescue is much shorter than this burn-off. We consider the population rescued if the mutant population reaches $K_{\text{tot}}/2$.

We performed 1000 replicates for each parameter combination. Simulation were written in Python and the code can be found on <https://github.com/mtomasini/EvoRescueFragmentation>. To test the accuracy of our simulations, we compared them against known cases (see Appendix B). Our simulation recovers expected results for the case with $m = 0$ (figure S2), $\theta = 0$ (figure S3) and $D = 2$ (figure S4).

Results

Optimal migration rate depends on cost of rescue mutations

From previous work on island models we know that there are three different effects of migration on evolutionary rescue. First, migration provides a demographic rescue effect where gene flow from the original habitat delays extinction in the deteriorated habitat, thus increasing the probability for rescue

mutations to occur and spread in the deteriorated habitat. For larger migration rates, rescue mutations are unlikely to establish because they are removed from demes where they are beneficial [Tomasini and Peischl, 2018] (i.e., gene swamping occurs [Bulmer, 1972, Lenormand, 2002]). Finally, when migration rates are very large, migration will deplete the original habitat such that density regulation leads to higher absolute fitness in the original habitats, which increases the establishment probability of rescue mutations [Tomasini and Peischl, 2018]. This effect has been called relaxed competition [Uecker et al., 2014] and is especially potent if the fitness cost of rescue mutations is low and if growth rates are high. Figure 2 shows the probability of rescue as a function of the migration rate for different combinations of selection intensities and varying numbers of demes. We identify three qualitatively different scenarios, depending on the strength of selection against rescue mutations in unperturbed environments. If the cost of the mutation in the original environment is relatively low ($s = 0.1$, Figures 2A and B) we see that increasing the rate of migration generally increases the probability of rescue in both the stepping stone as well as the island model. The reason is that for low costs of the rescue mutation we mainly see the positive effects of migration - demographic rescue and relaxed competition - whereas gene swamping can be neglected. The stepping stone and the island model show qualitatively similar patterns. The only difference is that for a given migration rate the positive effect of gene flow is larger in the island model as compared to the stepping stone model, and this difference increases with increasing number of demes. This is sensible because in the stepping stone model gene flow is restricted between neighboring demes, such that larger rates of gene flow are necessary to obtain the same positive effect of gene flow as compared to the island model.

Figures 2C and D show what happens when the cost of a mutation in the non-deteriorated environment is relatively high, but not lethal ($s = 0.5$). For $D = 4$ demes, we observe the effects of demographic rescue, gene swamping and relaxed competition. The interplay of demographic rescue and gene swamping causes a local maximum of the rescue probability for intermediate migration rates (around $m = 0.01 - 0.1$ in Figure 2C). Relaxed competition creates another increase in the probability of rescue when migration rates become for very large (around $m > 0.5$ in Figure 2C). As before, the island and the stepping stone model appear qualitatively similar for four demes. However, when the number of demes increases ($D = 16$ in Figure 2D) the two models show qualitatively different behavior. In the stepping stone

Figure 2: Comparison between stepping stone and island model. We show P_{rescue} as a function of migration rate for (A) $D = 4$, $s = 0.1$, (B) $D = 16$, $s = 0.1$ (C) $D = 4$, $s = 0.5$, (D) $D = 16$, $s = 0.5$, (E) $D = 4$, $s = 1.0$, (F) $D = 16$, $s = 1.0$. For all figures, $r = 0.5$, $z = 0.02$, and $\Theta = 4800$.

model the ranges where relaxed competition and gene swamping dominate start to overlap, such that the chance of evolutionary rescue monotonically increases with the migration rate (Figure 2D), whereas the two distinct peaks in the probability of rescue remain visible in the island model.

Finally, when rescue mutations are lethal in the old environment ($s = 1$), relaxed competition does not occur because lethal mutations do not survive until density regulation in the original habitat. We thus find that intermediate migration rates maximize the chances of survival (Figures 2E and F), which is due to two contrasting effects of demographic rescue and gene swamping [Uecker et al., 2014, Tomasini and Peischl, 2020]. Again, the main difference between the island and the stepping stone model is that the optimal migration rate for evolutionary rescue is larger in the stepping stone model than in the island model, and this difference increases with increasing number of demes.

Effective migration rate between habitats can explain differences between models

Gene swamping and relaxed competition are modulated by the amount of migration between different types of habitats, rather than the rate of migration between different demes. Roughly speaking, migration between demes that share the same environmental conditions does not affect the probability of establishment of a rescue mutation and has very little impact on local population sizes. We thus define an “effective migration rate” $\tilde{m}$ for the stepping stone model that quantifies the rate at which individuals migrate between the non-deteriorated and the deteriorated habitats. While this effective migration rate is difficult to calculate precisely, we present a simple heuristic derivation in Appendix A that shows that the effective migration rate between habitats in the stepping stone model is $\tilde{m} = m \cdot 2/D$.

We find that our heuristic effective migration rate can indeed explain the differences between the stepping stone model and the island model in most cases (Figure 3). Only when the cost of the rescue mutation is intermediate ($s = 0.5$) and the number of demes is large ($D = 16$) do we see notable differences between the two models. Apparently the reason for this is that while the effective migration rate captures the effects of demographic rescue and gene swamping quite well, relaxed competition depends on the migration rate in a more complex way in the stepping stone model. Indeed, when we perform simulations where we reduce the effect of relaxed competitions via Beverton-Holt density regulation [Beverton and Holt,

1957] with low growth rates, the discrepancy between the stepping stone model and the island model disappears (see figure S5 in the supplemental material).

Results are not strongly affected by the choice of parameters

Figure 4 shows contour plots of the probability of evolutionary rescue P_{res} as a function of the migration rate as well as other parameters of the model; we picked a set of parameters for which we expect qualitative differences between the models (highly fragmented habitat, $D = 16$, and when not otherwise specified, $s = 0.5$). Figures 4A, C and E show the probability of rescue in the island model; we recover findings previously shown for evolutionary rescue in a two-deme model and in an island model [Uecker et al., 2014, Tomasini and Peischl, 2020] and show that intermediate migration rates are optimal for a large range of parameters. Rescue is more likely for slower environmental change θ (figure 4A), for low stress r (figure 4C), and for a mutation weakly selected against in the old environment (low s , figure 4E). We can also see the strong effect of relaxed competition for low s and high migration rates m (figure 4E). Uecker et al. [2014] have studied the effects of different parameters for the island model in further detail, thus here we will focus on differences with the stepping stone model.

Figures 4B, D and F show the contour plots for the probability of rescue in the stepping stone model. In contrast with the island model, these figures show that large migration rates tend to maximize the probability for rescue in the stepping stone model. In general, we observe patterns similar to those found in the island model, although for higher migration rates, as we would expect when considering the difference in effective migration rates between the models. Rescue is more likely for larger values of θ , and smaller values of r and s . We can see that the qualitative differences that we identified in the previous sections are robust for a wide range of parameters, in particular over the whole range of r , for $100 < \theta < 500$ and for $0.2 < s < 0.9$. This suggests that the three cases that we studied in detail above (low s , intermediate s and high s) are representative of the majority of the range of parameters, with little change due to other parameters.

Figure 3: Evolutionary rescue as a function of the “effective migration rate”. We adjusted the migration rate in the stepping stone model, m_S , with respect to the migration rate in the island model, m_I , as $m_I = m_S \cdot 2/D$. Chosen examples are the same as in figure 2: (A) $D = 4$, $s = 0.1$, (B) $D = 16$, $s = 0.1$ (C) $D = 4$, $s = 0.5$, (D) $D = 16$, $s = 0.5$, (E) $D = 4$, $s = 1.0$, (F) $D = 16$, $s = 1.0$. For all figures, $r = 0.5$, $z = 0.02$, and $\Theta = 4800$.

Figure 4: We show the contour plots of (A, C, E) the probability of rescue for the island model P_{res}^I , (B, D, F) the probability of rescue for the stepping stone model P_{res}^S , as a function of the migration rate m . We vary parameters one at a time: (A-B) the length of an epoch θ ; (C-D) the harshness of environmental change r ; (E-F) the cost s of the mutation in the unperturbed environment. Other parameters are $D = 16$, $z = 0.02$, and when not varying, $\theta = 200$, $r = 0.5$ and $s = 0.5$.

Fragmentation can increase or decrease probability of rescue

Next we investigate the effect of increased habitat fragmentation in more detail. Since the effect of fragmentation is negligible in the island model [Uecker et al., 2014], we focus on the stepping stone model

instead. By increasing fragmentation we mean increasing the number of demes over which a population is distributed in each run, while keeping the total population size (and the time it takes such that the whole habitat is deteriorated) constant. In other words, we split a large, well mixed population into an increasing number of smaller sub-populations. In figure 5 we look at the effect of fragmentation given different values of m and s . Figure 5 shows that in contrast to the island model, the number of demes

Figure 5: Probability of rescue as a function of the number of demes, for different values of migration rate and different values of s . For all figures, gold is $m = 3.2 \times 10^{-4}$, orange is $m = 1 \times 10^{-2}$, magenta is $m = 2.154 \times 10^{-2}$, purple is $m = 1 \times 10^{-1}$ and blue is $m = 6.8129 \times 10^{-1}$. (A) $s = 0.1$, (B) $s = 0.5$, (C) $s = 1.0$. In all figures, $r = 0.5$, $z = 0.02$, $\Theta = 4800$.

can have quite dramatic effects on the probability of rescue. Only when gene flow is very low (yellow line in figures 5A, B and C, corresponding to $m = 3.2 \times 10^{-4}$) fragmentation has no effect on the probability for rescue. This is unsurprising, because with small migration populations evolve almost independently from one another. In such a scenario, evolutionary rescue depends on the total number of individuals in the habitat K_{tot} and not on how they are distributed over the environment. For larger values of gene flow ($m = 1 \times 10^{-2}$ (orange line), $m = 2.154 \times 10^{-2}$ (magenta line), and $m = 1 \times 10^{-1}$ (purple line) in Figure 5), we generally see that increasing fragmentation has negative effects on the rescue probability. The reason is simply that the positive demographic rescue effect of gene flow on evolutionary rescue

becomes smaller with increasing number of demes (see also figures 2 and 3, and the derivation of the effective migration rate). When gene flow is very high (blue line, corresponding to $m = 6.8129 \times 10^{-1}$; for comparison, this is the second to last simulated point in figure 2), we generally see a rather strong increase in the rescue probability with increasing number of demes. This is due to a transition from gene swamping being dominant for high migration and low number of demes (where the effective migration rate between habitats is large) to relaxed competition being dominant if the number of demes is large (see also figures 2D and 3D). The only exception is when the cost of rescue mutation is low, where we find certain rescue in virtually all simulation runs if the migration rate is high (5A). The reason is simply that when a mutation is weakly selected against in the old environment, relaxed competition outweighs gene swamping across the whole range of values of the number of demes D .

Random deterioration pattern

So far, we have considered evolutionary rescue in the island and the stepping stone models for scenarios where deterioration follows a spatial gradient (*i.e.* from deme 1 to deme D). While the order of deterioration of the demes is not relevant in the island model (because migration connects all the demes in the same way), it is important in the stepping stone model. This is once again because what determines the positive and negative effects of migration are the effective migration rates between types of environment, rather than between demes. In the stepping stone model with gradual deterioration, there is never more than $\kappa m/2$ individuals moving between regions; to this effect, we can see the lastly deteriorated deme as a sort of valve that controls the flow of individuals between regions. If deterioration occurs randomly though, there will be more than just two demes exchanging migrants from one region to the other. In figure 6 we observe that when deterioration is random, island model and stepping stone model show very few differences. In particular, for mild mutation cost ($s = 0.1$, figures 6A and B) no difference at all can be noted. For cases with larger s we observe slight quantitative differences that are probably to attribute to slight differences in the flow of mutants between deteriorated and non-deteriorated area. In the island model, the local population in each deme depends on the environmental state of the deme (deteriorated or non-deteriorated, see *e.g.* figure S6E and F); however in the stepping stone model with random deterioration local density in each deme depends on whether the neighboring demes are in the

same state, both in a different state, or split between non-deteriorated and deteriorated. The average effect on evolutionary rescue remains qualitatively the same.

Figure 6: Comparison between stepping stone and island model with random deterioration. We show P_{rescue} as a function of migration rate for (A) $D = 4$, $s = 0.1$, (B) $D = 16$, $s = 0.1$ (C) $D = 4$, $s = 0.5$, (D) $D = 16$, $s = 0.5$, (E) $D = 4$, $s = 1.0$, (F) $D = 16$, $s = 1.0$. For all figures, $r = 0.5$, $z = 0.02$, and $\Theta = 4800$.

Discussion

In this paper we analyzed how the interplay between spatial structure, gene flow and fragmentation affects evolutionary rescue. We compared the island model and the stepping stone model with environmental

conditions deteriorating over space and time.

In the island model the number of demes in the population does not influence rescue (as shown previously by Uecker et al. [2014]), while in the stepping stone model fragmentation strongly affects evolutionary rescue (figures 2, 4). In both models, rescue is the result of the interaction of three previously identified effects of gene flow on adaptation and thus rescue [Uecker et al., 2014]: gene swamping [Bulmer, 1972, Lenormand, 2002], demographic rescue [Brown and Kodric-Brown, 1977], and relaxed competition [Uecker et al., 2014]. Gene swamping and demographic rescue depend mostly on the flow of individuals between the non-deteriorated and the deteriorated areas of the environment, while relaxed competition is also strongly affected by the cost s of the rescuing mutation in the old environment.

We showed that for very high or very low cost of the rescuing mutation s , the stepping stone model can be thought of as a two-deme model with individuals moving between deteriorated and non-deteriorated regions, and we defined an effective migration rate that allows to compare island model to stepping stone model (figure 3A, B, E, F). This approximation breaks down for intermediate cost s and high fragmentation D (figure 3D).

Our main result was to show that, ultimately, the spatial structure of the habitat (*i.e.* how demes are connected) matters only in function of how it regulates the flow of individuals moving between the deteriorated and the non-deteriorated region; in the island model, fragmentation of a habitat does not affect its connectivity, while in the stepping stone model, fragmentation decreases the flow between areas due to decrease of local population. As such, all of these models can be understood as two-deme models – models where what is important is the exchange between different environments within the habitat. As a direct consequence of this, theoretical insights of two-deme models (*e.g.* Tomasini and Peischl [2020]) – generally presenting less analytical complexity – can also provide insights for models with more complex spatial structure.

We also analyzed how fragmentation affects evolutionary rescue, and showed that the effect of fragmentation is highly dependent on gene flow m and cost of mutation s (figure 5). As expected, for low migration rates fragmentation has no effect, as each deme evolves almost independently from the other demes. For intermediate migration, we find that rescue tends to decrease with increasing fragmentation, because demographic rescue becomes less important as deme carrying capacity decreases. For high migration,

rescue increases with increasing fragmentation, as relaxed competition becomes more and more dominant and gene swamping loses importance. When the mutation is weakly selected in the old environment, relaxed competition dominates for high migration rates, for any rate of fragmentation.

Finally, we showed that the two models are almost indistinguishable if demes are deteriorated randomly over space: this is due to the fact that with random deterioration, migration between deteriorated and non-deteriorated areas is not limited to the lastly deteriorated deme, but occurs on the interface of all the regions of the environment where two different environments are neighbors. Then, this is yet another case where the stepping stone model can be approximated by a two-deme model.

Because of the importance of gene flow between types of environment (as opposed to gene flow between demes), we argue that experimental set-ups should not ignore the interaction between migration rates and spatial structure; in particular, it is important to understand what happens when gene flow is modified directly (by the experimenter) or indirectly (by the effects of fragmentation). For example, studies such as [Bell and Gonzalez \[2011\]](#), analyzing the different rates of rescue events in an experimental island model or a stepping stone model, only consider spatial structure as an explanatory variable, and not the amount of gene flow – thus conflating the two. This may also help explain why some experimental set-ups have observed contrasting results about the net effect of gene flow on evolutionary rescue [[Carlson et al., 2014](#)].

The fact that spatially explicit models can often be interpreted as two-demes model, if confirmed experimentally, could lead to improvement of conservation strategies of fragmented habitats. Our results suggest that knowledge about the exact spatial arrangement of sub-populations is not always required to understand rescue processes, instead one only needs to consider the rate at which gene flow occurs between different types of habitats.

When deterioration does not occur gradually in space (random deterioration), we show that spatial structure plays no role and evolutionary rescue occurs as in the island model (figure 6). This is because when demes deteriorate randomly, the exchange of individuals between the deteriorated and the non-deteriorated area occurs at many demes, as opposed to only one in the stepping stone model deteriorating gradually in space. This suggests also that evolutionary rescue in two or three spatial dimensions should look more like evolutionary rescue in an island model, even when migration is limited in space:

this follows from the understanding that the deteriorated and non-deteriorated area are in contact over a larger “contact surface” than the one provided by the stepping stone model on a one-dimensional lattice. The exploration of different geometries in space, such as two- and three-dimensional lattices, could also yield a better understanding of the role of effects such as isolation by distance [Wright, 1943], as these geometries are arguably more general than one-dimensional lattices. However, the general conclusions of our study are not expected to change significantly in two- or three-dimensional habitats, as the effects we see (gene swamping, demographic rescue, relaxed competition) are not unique properties of a one-dimensional lattice.

We have assumed that migration is either between all demes, or only between neighboring demes. These are two extreme scenarios and an interesting extension would be to consider models where individuals disperse according to a distribution of migration distances. A better suited framework to explore such a theoretical model could be continuous space models (*e.g.* like in Barton [1987], Kirkpatrick and Peischl [2013]). For a narrow migration kernel, we expect to recover the behavior of the stepping stone model even in continuous space. As we increase the width of the kernel (*i.e.* the distance traveled by migrants) we would expect this model to be more similar to the island model. But since we do not expect gene flow to be isotropic in natural systems, because *e.g.* of physical barriers over the habitat and habitat range limits, it is difficult to predict whether chances of evolutionary rescue would increase or decrease when widening the migration kernel, due to the effects on evolutionary rescue when gene flow is not isotropic (see *e.g.* Kawecki and Holt [2002], Tomasini and Peischl [2020] for a discussion of asymmetric gene flow). In addition, Czuppon et al. [2021] studied the effects of biased migration on local adaptation and evolutionary rescue, using the same setting as in our current work. They show that dispersal schemes in which migration has a larger chance to occur towards deteriorated habitats increase the probability of rescue. This is in accordance to what we have found, as these migration schemes correspond to cases where the demographic positive effect of migration are stronger, and the negative effects are weaker.

In our model, density regulation determines how the probability of evolutionary rescue changes with high gene flow. When we choose a different mode of regulation (*e.g.* Beverton-Holt dynamics [Beverton and Holt, 1957]), the discrepancy between the island model and the stepping stone model disappears when accounting for the migration rate between environments (see figure S5 in the appendix). While

we assumed that density is regulated only in the non-deteriorated environment, density regulation in the deteriorated environment is also expected to influence the chances of evolutionary rescue [Holt and Gomulkiewicz, 1997, Gomulkiewicz et al., 1999].

In our study, we considered single rescue mutations, thus ignoring multi-step rescue [Uecker and Hermisson, 2016] or the effect of the distribution of fitness effects and the fitness-landscape [Anciaux et al., 2018, Osmond et al., 2020]. It would be interesting to study the interactions of genetic architecture and spatial structure on evolutionary rescue, which would require including recombination rates and other effects (*e.g.* clonal interference, Gerrish and Lenski [1998]). Our results should however be a good approximation for populations that reproduce sexually with strong recombination, such that rescue mutations evolve independently. We do not expect our model to be predictive for organisms reproducing with low recombination rates if multiple rescue mutations are required for restoring positive growth rates, as we neglected competition between beneficial mutations [Gerrish and Lenski, 1998] or competition in other ecological contexts (*e.g.* Osmond and de Mazancourt [2013], Klausmeier et al. [2020]). It has also been shown that the shape of the distribution of fitness effects as well as the recombination rate between loci harbouring rescue mutations can play an important role [Uecker and Hermisson, 2016, Anciaux et al., 2018, Osmond et al., 2020].

Ultimately, our understanding of how dispersal interacts with evolution to save populations from extinction is still incomplete. While we analyzed the positive and negative effects due to how dispersal modifies the probability of rescue through demographic changes, we ignored several other eco-evolutionary feedbacks [Thompson and Fronhofer, 2019]. We also ignored the fact that dispersal rates can themselves be subjected to evolution, especially when a species' range is shifting (*e.g.* Phillips and Shine [2006], Peischl and Gilbert [2020]). While such complexities are beyond the scope of the present study, they are paramount to reach a complete understanding of the feedback between evolution and dispersal, and how it affects the survival of species at risk of extinction.

Evolutionary rescue is still difficult to observe in natural populations [Carlson et al., 2014] with only a few recent putative examples (*e.g.* Epstein et al. [2016], Clayton and Spicer [2020], Muleta et al. [2021]). Direct comparison of our results to observed cases of evolutionary rescue in natural populations is therefore challenging. Laboratory model systems with automated dispersal show that gene flow can have different

effects on evolutionary rescue, depending on whether dispersal is implemented locally or globally [Bell, 2013]. Such systems provide exciting opportunities to validate our theoretical results. Indirect evidence for our observation that antibiotic resistance should evolve more easily in populations with strong spatial structure comes from biofilm producing bacteria, in which antibiotic resistance is particularly widespread and difficult to control (see *e.g.* Vasudevan et al. [2019]). Furthermore, loss of flagellar function can confer a fitness advantage in the presence of an antibiotic in *Pseudomonas aeruginosa* [Rundell et al., 2020], suggesting further interactions between the evolution of antibiotic resistance and dispersal related traits. Our model could hence shed light on the difficulties to avoid the evolution of treatment resistance in spatially structured populations such as biofilms.

In conclusion, a better understanding of the role of spatial structure and the mode of dispersal would be helpful not only for conservation strategies of fragmented natural populations, but also in the context of public health and the evolution of antibiotic resistance.

Authors contributions

MT and SP designed the project. MT and SP did analytical work. MT did all the simulations, figures, and wrote the first draft of the manuscript. MT and SP contributed to writing the manuscript.

Acknowledgments

We thank Laurent Excoffier, Joachim Hermisson and Katie Peichel for stimulating discussions on this subject, and for helpful comments on the first manuscript. We also thank Lindi Wahl and Nick Barton for editorial work on an earlier version of this manuscript, as well as anonymous reviewers.

Data archiving

The code used to perform simulations and generate figures, as well as the supporting information for this article can be found on github: <https://github.com/mtomasini/EvoRescueFragmentation>.

References

- M. N. Ahmed, A. Porse, M. O. A. Sommer, N. Høiby, and O. Ciofu. Evolution of antibiotic resistance in biofilm and planktonic pseudomonas aeruginosa populations exposed to subinhibitory levels of ciprofloxacin. *Antimicrobial agents and chemotherapy*, 62(8):e00320–18, 2018.
- H. K. Alexander, G. Martin, O. Y. Martin, and S. Bonhoeffer. Evolutionary rescue: linking theory for conservation and medicine. *Evolutionary applications*, 7(10):1161–1179, 2014.
- Y. Anciaux, L.-M. Chevin, O. Ronce, and G. Martin. Evolutionary rescue over a fitness landscape. *Genetics*, 209(1):265–279, 2018.
- R. D. Barrett and D. Schluter. Adaptation from standing genetic variation. *Trends in ecology & evolution*, 23(1):38–44, 2008.
- N. Barton. The probability of establishment of an advantageous mutant in a subdivided population. *Genetics Research*, 50(1):35–40, 1987.
- G. Bell. Evolutionary rescue and the limits of adaptation. *Phil. Trans. R. Soc. B*, 368(1610):20120080, 2013.
- G. Bell. Evolutionary rescue. *Annual Review of Ecology, Evolution, and Systematics*, 48(1): 605–627, 2017. doi: 10.1146/annurev-ecolsys-110316-023011. URL <https://doi.org/10.1146/annurev-ecolsys-110316-023011>.
- G. Bell and A. Gonzalez. Evolutionary rescue can prevent extinction following environmental change. *Ecology letters*, 12(9):942–948, 2009.
- G. Bell and A. Gonzalez. Adaptation and evolutionary rescue in metapopulations experiencing environmental deterioration. *Science*, 332(6035):1327–1330, 2011.
- R. J. H. Beverton and S. J. Holt. *On the dynamics of exploited fish populations*, volume 19. Ministry of Agriculture, Fisheries and Food, 1957.
- J. Botelho, F. Grosso, and L. Peixe. Antibiotic resistance in pseudomonas aeruginosa—mechanisms, epidemiology and evolution. *Drug Resistance Updates*, 44:100640, 2019.

- J. H. Brown and A. Kodric-Brown. Turnover rates in insular biogeography: effect of immigration on extinction. *Ecology*, 58(2):445–449, 1977.
- M. G. Bulmer. Multiple niche polymorphism. *The American Naturalist*, 106(948):254–257, 1972.
- A. E. Cahill, M. E. Aiello-Lammens, M. C. Fisher-Reid, X. Hua, C. J. Karanewsky, H. Y. Ryu, G. C. Sbeglia, F. Spagnolo, J. B. Waldron, O. Warsi, and J. J. Wiens. How does climate change cause extinction? *Proceedings of the Royal Society B: Biological Sciences*, 280(1750):20121890, 2013. doi: 10.1098/rspb.2012.1890. URL <https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2012.1890>.
- S. M. Carlson, C. J. Cunningham, and P. A. Westley. Evolutionary rescue in a changing world. *Trends in Ecology & Evolution*, 29(9):521–530, 2014.
- K. A. Clayton and J. I. Spicer. Evidence for physiological niche expansion of an intertidal flatworm: evolutionary rescue in the wild. *Marine Ecology Progress Series*, 651:85–95, 2020.
- P. Czappon, F. Blanquart, H. Uecker, and F. Débarre. The effect of habitat choice on evolutionary rescue in subdivided populations. *The American Naturalist*, 197(6):000–000, 2021.
- B. Epstein, M. Jones, R. Hamede, S. Hendricks, H. McCallum, E. P. Murchison, B. Schönfeld, C. Wiench, P. Hohenlohe, and A. Storfer. Rapid evolutionary response to a transmissible cancer in tasmanian devils. *Nature communications*, 7:12684, 2016.
- A. F. Feder, P. S. Pennings, J. Hermisson, and D. A. Petrov. Evolutionary dynamics in structured populations under strong population genetic forces. *G3: Genes, Genomes, Genetics*, 9(10):3395–3407, 2019.
- P. J. Gerrish and R. E. Lenski. The fate of competing beneficial mutations in an asexual population. *Genetica*, 102:127, 1998.
- R. Gomulkiewicz and R. D. Holt. When does evolution by natural selection prevent extinction? *Evolution*, 49(1):201–207, 1995.

- R. Gomulkiewicz, R. D. Holt, and M. Barfield. The effects of density dependence and immigration on local adaptation and niche evolution in a black-hole sink environment. *Theoretical population biology*, 55(3):283–296, 1999.
- A. Gonzalez and G. Bell. Evolutionary rescue and adaptation to abrupt environmental change depends upon the history of stress. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 368(1610), 2013. ISSN 0962-8436. doi: 10.1098/rstb.2012.0079. URL <http://rstb.royalsocietypublishing.org/content/368/1610/20120079>.
- R. D. Holt and R. Gomulkiewicz. How does immigration influence local adaptation? a reexamination of a familiar paradigm. *The American Naturalist*, 149(3):563–572, 1997.
- S. Karlin. Classifications of selection-migration structures and conditions for a protected polymorphism. *Evol. Biol*, 14(61):204, 1982.
- T. J. Kawecki and R. D. Holt. Evolutionary consequences of asymmetric dispersal rates. *The American Naturalist*, 160(3):333–347, 2002.
- M. Kimura and G. H. Weiss. The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics*, 49(4):561–576, 1964.
- M. Kirkpatrick and S. Peischl. Evolutionary rescue by beneficial mutations in environments that change in space and time. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1610):20120082, 2013.
- C. A. Klausmeier, M. M. Osmond, C. T. Kremer, and E. Litchman. Ecological limits to evolutionary rescue. *Philosophical Transactions of the Royal Society B*, 375(1814):20190453, 2020.
- T. Lenormand. Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, 17(4):183–189, 2002.
- M. Lynch. Evolution and extinction in response to environmental change. *Biotic interactions and global change*, pages 234–250, 1993.
- L. Marrec and A.-F. Bitbol. Adapt or perish: Evolutionary rescue in a gradually deteriorating environment. *Genetics*, 216(2):573–583, 2020.

- A. H. Melnyk, A. Wong, and R. Kassen. The fitness costs of antibiotic resistance mutations. *Evolutionary applications*, 8(3):273–283, 2015.
- S. Moreno-Gamez, A. L. Hill, D. I. Rosenbloom, D. A. Petrov, M. A. Nowak, and P. S. Pennings. Imperfect drug penetration leads to spatial monotherapy and rapid evolution of multidrug resistance. *Proceedings of the National Academy of Sciences*, 112(22):E2874–E2883, 2015.
- K. T. Muleta, T. Felderhoff, N. Winans, R. Walstead, J. R. Charles, J. S. Armstrong, S. Mamidi, C. Plott, J. P. Vogel, P. G. Lemaux, et al. The recent evolutionary rescue of a staple crop depended on over half a century of global germplasm exchange. *bioRxiv*, 2021.
- H. A. Orr and R. L. Unckless. Population extinction and the genetics of adaptation. *The American Naturalist*, 172(2):160–169, 2008.
- M. M. Osmond and C. de Mazancourt. How competition affects evolutionary rescue. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1610):20120085, 2013.
- M. M. Osmond, S. P. Otto, and G. Martin. Genetic paths to evolutionary rescue and the distribution of fitness effects along them. *Genetics*, 214(2):493–510, 2020.
- S. Peischl and K. J. Gilbert. Evolution of dispersal can rescue populations from expansion load. *The American Naturalist*, 195(2):349–360, 2020.
- B. L. Phillips and R. Shine. An invasive species induces rapid adaptive change in a native predator: cane toads and black snakes in australia. *Proceedings of the Royal Society B: Biological Sciences*, 273(1593):1545–1550, 2006.
- J. Ramsayer, O. Kaltz, and M. E. Hochberg. Evolutionary rescue in populations of pseudomonas fluorescens across an antibiotic gradient. *Evolutionary applications*, 6(4):608–616, 2013.
- E. Rundell, N. Commodore, A. Goodman, and B. Kazmierczak. A screen for antibiotic resistance determinants reveals a fitness cost of the flagellum in pseudomonas aeruginosa. *Journal of Bacteriology*, 202(6), 2020.
- P. Samani and G. Bell. Adaptation of experimental yeast populations to stressful conditions in relation to population size. *Journal of evolutionary biology*, 23(4):791–796, 2010.

- C. D. Thomas, A. Cameron, R. E. Green, M. Bakkenes, L. J. Beaumont, Y. C. Collingham, B. F. Erasmus, M. F. De Siqueira, A. Grainger, L. Hannah, et al. Extinction risk from climate change. *Nature*, 427 (6970):145, 2004.
- P. L. Thompson and E. A. Fronhofer. The conflict between adaptation and dispersal for maintaining biodiversity in changing environments. *Proceedings of the National Academy of Sciences*, 116(42): 21061–21067, 2019.
- M. Tomasini and S. Peischl. Establishment of locally adapted mutations under divergent selection. *Genetics*, 209(3):885–895, 2018. doi: 10.1534/genetics.118.301104.
- M. Tomasini and S. Peischl. When does gene flow facilitate evolutionary rescue? *Evolution*, 74(8): 1640–1653, 2020. doi: 10.1111/evo.14038.
- H. Uecker and J. Hermisson. The role of recombination in evolutionary rescue. *Genetics*, 202(2):721–732, 2016.
- H. Uecker, S. P. Otto, and J. Hermisson. Evolutionary rescue in structured populations. *The American Naturalist*, 183(1):E17–E35, 2014.
- E. Vander Wal, D. Garant, M. Festa-Bianchet, and F. Pelletier. Evolutionary rescue in vertebrates: evidence, applications and uncertainty. *Phil. Trans. R. Soc. B*, 368(1610):20120090, 2013.
- S. Vasudevan, H. A. Joseph, S. S. Swamy, and A. P. Solomon. Antibiotic resistance in biofilms. In *Introduction to Biofilm Engineering*, pages 205–224. ACS Publications, 2019.
- S. Vuilleumier, J. Goudet, and N. Perrin. Evolution in heterogeneous populations: from migration models to fixation probabilities. *Theoretical population biology*, 78(4):250–258, 2010.
- S. Wright. Isolation by distance. *Genetics*, 28(2):114–138, 1943.