Competitive advantage of oral streptococci for colonization of the middle ear mucosa

Kristin M. Jacob and Gemma Reguera[#]

Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing, Michigan, USA

#Corresponding Author:

Gemma Reguera

567 Wilson Rd., Rm. 6190

Biomedical & Physical Sciences building

East Lansing, MI 48824

Tel.: (517)884-5401

e-mail: reguera@msu.edu.

Running title: otic streptococci

Keywords: middle ear, Eustachian tube, biofilms, otic microbiome, oral cavity, otopathogen, mucin

Abstract

The intermittent aeration of the middle ear seeds its mucosa with saliva aerosols and selects for a distinct community of commensals adapted to the otic microenvironment. We gained insights into the selective forces that enrich for specific groups of oral migrants in the middle ear mucosa by investigating the phylogeny and physiology of 19 strains enriched (*Streptococcus*) or transiently present (*Staphylococcus*, *Neisseria* and actinobacterial *Micrococcus* and *Corynebacterium*) in otic secretions. Phylogenetic analyses of full length 16S rRNA sequences resolved close relationships between the streptococcal strains and oral commensals as well as between the transient migrants and known nasal and oral species. Physiological functions that facilitate mucosal colonization (swarming motility, surfactant production) and nutrition (mucin and protein degradation) were widespread in all the otic cultivars, as was the ability of most of the isolates to grow both aerobically and anaerobically. However, streptococci stood out for their enhanced biofilm-forming abilities under oxic and anoxic conditions and for their efficient fermentation of mucosal substrates into lactate, a key metabolic intermediate in the otic trophic webs. Additionally, the otic streptococci inhibited the growth of common otopathogens, an antagonistic interaction that could exclude competitors and protect the middle ear mucosa from infections by transient pathobionts. These adaptive traits allow streptococcal migrants to colonize the otic mucosa and grow microcolonies with syntrophic anaerobic partners, establishing trophic webs with other commensals similar to those formed by the oral ancestors in buccal biofilms.

Importance

The identification of a diverse microbiome in otic secretions from healthy young adults challenged the entrenched dogma of middle ear sterility and underscored previously unknown roles for oral commensals in the seeding of otic biofilms. By comparing the physiology of novel

lineages of streptococci and transient (peri)oral species isolated from otic secretions, we identified adaptive behaviors that allow specific oral streptococcal species to successfully colonize the mucosa of the middle ear. We also describe antagonistic properties of the otic streptococci that help them outcompete transient nasal and oral migrants, including known otopathogens. This knowledge is important to predictively understand the functionality of the otic communities, their interactions with the host mucosa and the outcome of infections.

Introduction

The oral cavity provides a heterogenous landscape of surfaces and microenvironments (teeth, gingiva, tongue, cheek, hard and soft palate, etc.) for the growth of microbial communities (1). The availability of dietary substrates supports the growth and diversification of the oral inhabitants, making these communities some of the richest and most diverse in the human body (2). Many of these microbes readily disperse with saliva into perioral regions (1) and, from there, to other parts of the digestive tract (3). Dispersal is also enhanced by the spread of saliva aerosols from the oropharynx (back of the throat) into the respiratory tract (3). Indeed, the air inhaled through the nasal passages or the mouth disperses the saliva aerosols from the oropharynx into the esophagus and trachea and, from there, to the lower parts of the aerodigestive tract. Similarly, air exhaled from the lungs disperses saliva aerosols from the oropharynx into the nasal cavity. The opening of the extended tube of the middle ear (the Eustachian tube, Fig. 1) into the distal part of the nasopharynx also promotes the entry of saliva aerosols with exhaled air (4). The Eustachian tube is passively collapsed at rest to sound proof the tympanic cavity and minimize microbial entry, yet it opens when we swallow to draw in air from the lower airways and ventilate the tympanic cavity (Fig. 1) (4). The periodic aperture and collapse of the Eustachian tube promotes the intermittent aeration of the tympanic cavity, relieves negative pressure across the eardrum and secretes into the nasopharynx excess mucus and fluids (4). It also introduces saliva aerosols into the middle ear and establishes fluctuating redox conditions optimal for the growth of strict and facultative anaerobes (5). Consistent with this, otic secretions collected at the orifice of the Eustachian tube are enriched in obligate or facultatively anaerobic genera within the phyla Bacteroidetes (*Prevotella* and *Alloprevotella*), Fusobacteria (*Fusobacterium* and *Leptotrichia*) and Firmicutes (*Veillonella* and *Streptococcus*) (5). The co-enrichment in otic secretions of Bacteroidetes, streptococci and *Veillonella* spp. suggests that they grow syntrophically as in oral biofilms (5). In this model (Fig.

1), *Bacteroidetes* could degrade mucin glycoproteins and other mucosal proteins into substrates (sugars and peptides) that oral streptococci can ferment under both aerobic and anaerobic conditions (6). This metabolic capacity could produce lactate for *Veillonella* fermentation into propionate, acetate, CO₂ and H₂, as described in oral biofilms (7, 8). The lactate dependency of *Veillonella* spp. could also support metabolic interactions with *Bacteroidetes* partners that ferment simple sugars into lactate (9). The co-aggregative properties of oral streptococci during the formation of the dental plaque (2, 10) could facilitate the formation of microcolonies in the middle ear as well and promote metabolic exchange with the strictly anaerobic *Bacteroidetes* and *Veillonella* partners (5). Through their collective metabolism (Fig. 1), otic microcolonies of *Bacteroidetes*, streptococci and *Veillonella* could degrade and ferment host-derived mucins and proteins into short chain fatty acids such as propionate and acetate, potentially contributing to mucosal health as in other body sites (11).

Fig 1: Illustration of the human ear anatomy (left) and trophic webs within bacterial microcolonies in the middle ear mucosa (right). The human ear is divided in three compartments (outer, middle and inner). The eardrum separates the outer ear canal from the tympanic cavity of the middle ear, which extends as a tube (Eustachian tube) into the nasopharynx to draw in air and drain otic secretions. The microbiome sequenced from otic secretions of healthy young adults (5) supports the establishment of an otic trophic web (inset) for the degradation of host mucins and proteins by *Bacteroidetes* into substrates (sugars and peptides) that *Streptococcus* and *Veillonella* cooperatively ferment into short chain fatty acids (SCFs) via lactate.

The presence of bacterial microcolonies in biopsy specimens of the mucosal lining of the tympanic cavity (12, 13) suggests that some oral migrants penetrated the mucus layer, attached

to the underlying epithelium and formed biofilms. Swarming motility could have allowed these successful colonizers to move through the viscous mucus medium and evade immune attack (14). Crossing the mucus barrier is also important to avoid clearance, which in the middle ear is mediated by the movement of cilia (mucociliary clearance) and the pumping force exerted by muscles contracting and relaxing around the Eustachian tube when swallowing (muscular clearance) (15, 16). Once they reach the mucosal epithelium, successful colonizers attach to it and grow as microcolonies on the mucosal surface to avoid clearance (17). This points at biofilm formation as a critical selective factor for growth and reproduction in the otic mucosa. Biofilms are also important to create anoxic microenvironments for anaerobic partners to grow despite the periodic redox fluctuations experienced in the middle ear. Because the aperture of the Eustachian tube is triggered by swallowing, air only enters the middle ear every minute during the wake hours or every five minutes during sleep (4). Facultative anaerobes such as *Streptococcus* have a growth advantage under these conditions, and could co-aggregate as in oral biofilms (18) to establish anaerobic niches for their anaerobic syntrophic partners (e.g., *Bacteroidetes* and *Veillonella* species). Nutrient foraging is also important for the establishment of biofilms in the middle ear mucosa. In the absence of dietary nutrients, the otic communities are more likely to sustain their trophic webs with host-derived nutrients such as proteins and mucin glycoproteins secreted to the mucosa (19). Motility in the viscous environment of the mucosa could facilitate nutrient foraging (20) while the ability of the bacteria to secrete hydrolytic enzymes (mucinases and proteases) could allow them to break down the mucin glycoproteins and mucosal proteins into sugars and amino acids (21).

Phylogenetic analyses of cultivars recovered from otic secretions also support an oral ancestry of the middle ear communities (5). These cultivars include streptococcal strains closely related yet phylogenetically distinct from oral ancestors, consistent with the diversification of oral taxa into lineages better suited for growth and reproduction in the middle ear mucosa (5). Cultivars recovered from otic secretions also include nasal and oral species from genera

(*Staphylococcus*, *Neisseria* and actinobacterial *Micrococcus* and *Corynebacterium*) that transiently disperse through the oral and perioral regions (5). Streptococcal and staphylococcal species are, for example, among the most prominent members in the oral and nasal microbiomes, respectively (3, 22). Both groups disperse in the aerodigestive tract and are predicted to enter the middle ear during the intermittent cycles of aperture of the Eustachian tube. Yet, while streptococci are one of the most abundant groups in otic secretions, staphylococcal-like sequences are seldom detected (5). This suggests that streptococcal migrants have a competitive advantage over the transient staphylococcal species during the colonization of the middle ear mucosa. To test this, we sequenced and partially assembled the genomes of 19 cultivars representing otic commensals (*Streptococcus*) and transient genera (*Staphylococcus*, *Neisseria*, *Micrococcus* and *Corynebacterium*) (5) and used the full length 16S rRNA sequences to identify their closest relatives. We then screened the cultivars for adaptive traits predicted to be important for mucosal colonization (swarming, surfactant production, biofilm formation) and for growth under conditions (redox, nutritional, etc.) relevant to the middle ear microenvironment. Our study revealed similar adaptive traits for mucosal growth in most of the isolates but aggregative and metabolic properties of streptococci that could facilitate their attachment to the mucosal epithelium and syntrophic growth within microcolonies. These same properties are also present in their closest oral ancestors, with whom they share the ability to establish trophic webs with anaerobes and antagonize transient members from the oral and perioral regions. These findings provide novel insights into the adaptive responses that sustain the growth and functionality of otic communities as well as interactions with the host mucosa and transient migrants that influence the outcome of infections.

Results

Phylogenetic analysis supports the oral ancestry of otic streptococcal commensals

Cultivars recovered from otic secretions collected from healthy young adults include genera enriched in the otic secretions (*Streptococcus*) as well as transient or lowly abundant groups (*Staphylococcus*, *Neisseria*, and the actinobacterial genera *Micrococcus* and *Corynebacterium*) (5). Phylogenetic analysis of partial 16S rDNA sequences amplified from these isolates revealed close relationships with oral (oropharyngeal and buccal) strains isolated from the same individuals but lacked the resolution needed for species-level demarcation (5). Thus, we sequenced and partially assembled the genomes of 19 otic cultivars to retrieve full-length 16S rDNA sequences for each of the isolates. A species sequence identity cutoff of >98.7% (23) matched each otic isolate to more than one species within each genus (Table 1 shows the top identity hit for each strain). Phylogenetic inference methods resolved, however, close evolutionary ties with specific species that reside or are transiently isolated from the oral cavity (Fig. 2).

Fig. 2: 16S rRNA gene phylogeny of otic cultivars. Maximum-likelihood tree constructed with full-length 16S rRNA sequences from the otic isolates and the closest reference strains (accession number in parentheses). The scale bar indicates 5% sequence divergence filtered to a conservation threshold above 79% using the Living Tree Database (24, 25). Bootstrap probabilities by 1000 replicates at or above 50% are denoted by numbers at each node. The circles identify catalase-positive isolates.

The nearest neighbor to most of the *Streptococcus* sequences (seven of them) was *Streptococcus salivarius* (subspecies *salivarius* and *thermophilus*) (Fig. 2). Genomic divergence (size and gene content) is high for species and subspecies within the *Salivarius* group (26). As a result, strains of *S. salivarius* can have very different metabolic and physiological characteristics or even habitat/host preferences despite high 16S rRNA sequence identity (26, 27). Thus, the separate clustering of the 7 otic strains of *S. salivarius* could reflect substantial divergence of the otic subclade from an oral ancestor. The rest of the otic streptococcal sequences clustered separately from close oral relatives within the *Mitis* (L0020-02 and *Streptococcus*

parasanguinis), Viridans (L0023-02 and *Streptococcus pseudoneumoniae*) and the Lancefield's group B streptococcus or GBS (L0023-01 and L0023-03 and *Streptococcus agalactiae*) groups (27, 28) (Fig. 2). Hence, 16S rRNA phylogeny supports the oral ancestry of the streptococcal cultivars but also reveals a level of divergence that is consistent with the ecological diversification of niche-adapted otic lineages proposed in earlier studies (5).

The 16S rRNA sequence identity of the non-streptococcal strains also produced more than one match to species of *Staphylococcus*, *Neisseria*, *Micrococcus* and *Corynebacterium* (Table 1). Consistent with the classification of three of the isolates as *Staphylococcus* spp., all were catalase negative, and all branched within subclades of staphylococcal 16S rRNA sequences (Fig. 2). The closest neighbors to the three otic staphylococci were species (*Staphylococcus hominis*, *Staphylococcus aureus*, and *Staphylococcus epidermidis*) that are highly represented in the nasal passages (22). Their nasal abundance facilitates their dispersal in the contiguous oral cavity (29) and their transient detection in oral and perioral regions (29, 30), including otic secretions (5). Transience also explains the recovery of *Neisseria* strains closely related to *Neisseria perflava*, *Neisseria subflava* and *Neisseria flavescens* (Fig. 2), which are species that colonize the mucosa of the oropharynx (31, 32) and disperse via saliva aerosols (33). Neutral community models predict, however, a similar distribution of *Neisseria* species in otic secretions compared to oral and perioral source communities, supporting the idea that they are transient migrants (5). The otic isolates also included two actinobacterial *Micrococcus* and *Corynebacterium* strains (Table 1). The *Micrococcus* isolate was catalase-positive, a general phenotypic trait of the genus (34), and branched closely to *Micrococcus yunnanensis* (Fig. 2). This is a soil *Micrococcus* species (35) that, like other environmental micrococci, transiently disperses with air in the human aerodigestive tract (36). The second actinobacterial isolate was closely related to *Corynebacterium pseudodiphthericum* (Fig. 2). *Corynebacterium* commensals are prominent members of the nasal microbiomes and antagonists of nasal pathobionts, including some of the most important otopathogens (37). Their

abundance in the nasal microflora explains their detection in oral and perioral regions (38). However, actinobacteria only account for ~1% of the operational taxonomic units (OTUs) in otic secretions, suggesting they are negatively selected for growth and reproduction in the middle ear mucosa (5).

Surfactant-mediated swarming motility is widespread among the otic cultivars

Successful colonization of any respiratory mucosa requires bacterial migrants to move rapidly across the mucus layer in order to avoid immune attack and clearance (17). Some flagellated bacteria can reach the underlying epithelial lining by rapidly swarming in groups through the viscous mucoid layer, a process that is stimulated by the lubricating effect of mucin glycoproteins (20). Swarming behaviors can be identified in laboratory plate assays that test the expansion of microcolonies on a soft agar (0.4-0.5%) surface (20). Thus, we tested the ability of the 19 otic isolates to swarm on the surface of 0.5% tryptone soy agar (TSA) plates in reference to the robust swarmer *Pseudomonas aeruginosa* PA01 (39). Figure 3A shows the progression of the swarming expansion for each strain over time, which we calculated as the average expansion zone in triplicate plate assays (Table 2). Although *P. aeruginosa* showed large zones of swarming expansion already at 18 h, we only detected swarming activity in the otic isolates after 42 or 62 h of colony growth (Fig. 3A). Lag phases are not unusual prior to swarming on agar plates for cells that must reprogram their physiology to grow on the surface of the agar-solidified medium (20). Consistent with this, the strains that grew faster on the semisolid TSA plates (three staphylococcal and the two actinobacterial isolates) produced visible zones of swarming expansion at 42 h, while the slowest growers (*N. perflava* L0023-05 and L0023-06) required 62 h of incubation (Table 2). Notably, most of the streptococci grew well in tryptone soy broth (TSB), yet they aggregated strongly when growing on the surface of soft-agar plates (Fig. 3B). These aggregative strains also had delayed swarming phenotypes on 0.5% TSA plates (Table 2) but it was possible to rescue the swarming delay on plates with lower (0.4%) agar

concentration (Fig. 3B). For example, the streptococcal strain L0022-03 did not swarm on 0.5% TSA plates until after 62 h (Table 2) but expanded 0.28 cm away from the edge of the macrocolony after 42 h of growth on 0.4% TSA plates (Fig. 3B). Temperate swimmers often require a softer agar surface to overcome frictional forces between the cell and the surface (20). This is because lowering the agar concentration facilitates water movement to the surface and immerses the cells in a layer of liquid that stimulates swarming (20).

Fig. 3. Swarming motility and surfactant production by otic cultivars in reference to *Pseudomonas aeruginosa* PA01. (A) Average surfactant production (halo of mineral oil dispersal around 24h colonies grown on 1.5% TSA) and size of swarming expansion (0.5% TSA plates at 18, 42 and 62 h) measured in triplicate replicates of the otic isolates (*Streptococcus* in gray; *Staphylococcus* in orange; *Neisseria* in purple; and actinobacterial strains of *Corynebacterium* and *Micrococcus* in green) and the positive control (*P. aeruginosa* PA01 in white). (B) Representative images of swarming (0.4% TSA, 42 h) and surfactant (1.5% TSA, 24 h) plate assays for *P. aeruginosa* PA01 (positive control, boxed) and otic strains of *Streptococcus*, *Corynebacterium* and *Micrococcus* (scale bars, 0.5 cm). The edge of the surfactant halo is highlighted with a dashed white line.

The need for some bacteria to express cellular components (flagella, exopolysaccharide, surfactants, etc.) mediating swarming on semisolid agar can also delay the appearance of expansion zones (20). Surfactants are particularly important to reduce frictional resistance between the surface of swarming cells and the underlying substratum (20). Furthermore, their concentration and diffusion in soft-agar medium controls the extent of swarming expansion (40). Hence, we also screened the cultivars for surfactant production. To do this, we spot-plated each

isolate on hard agar (1.5%) TSA plates and airbrushed a fine mist of mineral oil droplets onto colonies grown at 37°C for 24 h. This atomized oil assay instantaneously reveals halos of oil droplet dispersal around surfactant-producing strains and allows for semiquantitative estimation of the levels of surfactant production, even when present at concentrations too low to be detected by traditional methods such as the water drop collapse assay (40). We detected haloes of oil dispersal around 9 of the isolates at 24 h (Table 2). Furthermore, we observed positive correlations between surfactant production and swarming ability on 0.5% TSA for most strains (Fig. 3A). For example, the actinobacterial isolates, which were the most robust swimmers, produced the highest levels of surfactant (Table 2). By contrast, temperate swimmers such as the streptococcal isolates produced low or undetectable levels of surfactants under the experimental conditions. As an exception, the staphylococcal isolates swarmed robustly on the soft agar plates (Fig. 3) although they did not produce detectable halos of mineral oil dispersion (Table 2). Staphylococcal cells lack flagellar locomotion and thus do not have canonical (i.e., flagella-driven) swarming behaviors. These bacteria can however passively ‘spread’ on soft agar surfaces (41) through the coordinated synthesis of lubricating peptides known as phenol-soluble modulins (PSMs) (42). PSM surfactants accumulate very close to the colony edge (43). Hence, they are unlikely to produce a halo of oil dispersal in the atomized assay used for testing.

Redox and nutritional advantage of otic streptococci in the middle ear mucosa

Successful colonizers of the middle ear mucosa face sharp redox fluctuations due to the brief (400 milliseconds) and infrequent (approximately every minute when we swallow) aperture of the Eustachian tube (4). The enrichment in otic secretions of anaerobic metabolisms further supports the idea that conditions of oxygen limitation prevail in the middle ear (5). For this reason, we tested the ability of the otic cultivars to grow under aerobic or anaerobic conditions (Fig. 4A). All the isolates grew well in oxic and anoxic TSB medium, except for two *Neisseria*

strains (L0023-05 and L0023-06) that grew slowly in the oxic broth. These two strains also flocculated extensively in oxic medium, an aggregative behavior exhibited by microaerophiles in response to elevated (and toxic) oxygen concentrations (44). By contrast, the streptococcal and staphylococcal strains had similar growth rates aerobically and anaerobically (0.56 ± 0.23 and 0.49 ± 0.12 doubling times, respectively), suggestive of a competitive advantage for growth and reproduction under sharp redox fluctuations. The actinobacterial strains also grew aerobically and anaerobically but differed in their redox preference. Both isolates doubled every ~ 0.5 h under anoxic conditions. However, generation times increased in aerobic cultures of *Micrococcus* L0020-05 (~ 0.7 h) whereas *Corynebacterium* L0020-06 grew more rapidly than any other strain under these conditions (0.15 h average generation time). This aerobic preference matches well the enrichment of *Corynebacterium* species in the aerated nasal passages (22) and the reduced abundance of this group in otic secretions (5).

Fig. 4: Growth of otic isolates as a function of oxygen availability and host nutrients (protein and mucin). (A) Average doubling times of otic isolates growing in triplicate TSB cultures aerobically or anaerobically at 37°C. Data points are color-coded for *Streptococcus* (gray), *Staphylococcus* (orange), *Neisseria* (purple) and actinobacterial genera *Micrococcus* and *Corynebacterium* (green). The flocculating strains of *Neisseria* are labeled. (B) Protease and mucinase activity (haloes of milk casein or porcine gastric mucin degradation, respectively) of representative otic isolates and *P. aeruginosa* PA01 (positive control, boxed). The milk casein plates were photographed without staining after 24 h of incubation at 37°C. The mucin plates were incubated for 48 h and stained with 0.1% amido black prior to photography. Scale bars, 0.5 cm.

In addition to redox fluctuations, bacteria colonizing the otic mucosa must cope with a scarcity of nutrients. The limited carriage of dietary substrates in saliva aerosols reduces nutrient availability in the middle ear, exerting selective pressure on colonizers to use host-derived nutrients such as mucosal proteins and mucin glycoproteins (5). To test this, we screened the otic isolates for their ability to secrete enzymes (proteases and mucinases) needed to break down the host nutrients into readily assimilated substrates (amino acids and sugars). For these experiments, we spot-plated the cultivars onto TSA plates supplemented with 5% lactose-free skim milk (protease assay) or 0.5% porcine gastric mucin (mucinase assay) for 24 h to identify zones of substrate degradation around the colonies. Figure 4B shows typical results for representative otic strains and the positive control *P. aeruginosa* PA01, while Table 3 shows the results (presence or absence of a halo of substrate degradation) in triplicate plate assays for each strain. Notably, all the isolates were able to degrade mucin to various degrees after 24 h of incubation at 37°C. Three aggregative strains of *S. salivarius* (L0021-01, L0022-03 and L0022-04) produced only faint mucin clearings after 24 h (+/- in Table 3) but the zone of degradation became more prominent after extending the incubation period to 48 h. By contrast, protease activity was only detected in the streptococcal and staphylococcal groups (Table 3). Extracellular proteases typically have low substrate selectivity and, thus, can cleave a wide range of substrates (45). Non-selectivity is particularly advantageous in the middle ear mucosa, as it allows residents to scavenge proteins and the mucin protein backbone as nitrogen sources (19). In addition to providing a metabolic advantage, proteases facilitate mucosal penetration, control mucus viscosity, modulate host immune responses, and can prevent the establishment of competitors (46). Therefore, protease secretion may confer on streptococcal and staphylococcal migrants a competitive advantage for growth and reproduction in the otic mucosa.

Metabolic advantage of streptococci for syntrophic growth in biofilms

The presence of bacterial microcolonies on the epithelial surface of biopsy specimens collected from the tympanic cavity of healthy individuals (13) points at biofilm formation as an important adaptive response for successful colonization of the middle ear mucosa. Based on this, we investigated the ability of the otic isolates to form biofilms under aerobic and anaerobic conditions. These assays used crystal violet to stain 24-h biofilms formed at the bottom of the microtiter plates that we previously used for planktonic growth studies (Fig. 4A). We then solubilized the biofilm-associated dye to estimate the biofilm biomass from the absorbance of the solution at 550 nm (Fig. 5A). All but two streptococcal strains (*S. pseudopneumoniae* L0023-02 and *S. agalactiae* L0023-03) formed robust biofilms under aerobic conditions. Among these streptococcal biofilm formers, four strains (*S. salivarius* L0021-04 and L0021-05, *S. parasanguinis* L0020-02, and *S. agalactiae* L0023-01) clustered separately with a staphylococcal isolate (*S. aureus* L0021-02) for their ability to also form robust biofilms in anoxic media (Fig. 5A). This contrasts with strains of *S. salivarius* (L0021-01, L0022-03, L0022-04, L0022-05 and L0022-06) that had a biofilm growth advantage in oxic medium only. The enhanced biofilm abilities of these isolates correlated well with pH drops below 5 in the culture broth (Fig. 5B) due to the accumulation of lactate as a fermentation byproduct ($p=0.03$) (Fig. 5C). Culture acidification likely triggered biofilm formation in the streptococcal cultures, because cells entered stationary phase (0.62 ± 0.05 OD₆₀₀) once the pH dropped below 5. These results point at lactate accumulation and broth acidification as the trigger of the planktonic-to-biofilm transition in the otic streptococci. This response is similar to that described for oral streptococcal commensals, which also produce lactic acid as the main fermentation byproduct (47) and stop growing once the pH drops to inhibitory levels, usually at or below 5 (48). As a result, commensal oral streptococci co-aggregate with lactate-utilizing bacteria such as *Veillonella* (49). The abundance of not only streptococci but also *Veillonella* sequences in otic secretions is suggestive of similar trophic webs in the middle ear mucosa within microcolonies (5). Such syntrophic partnerships with *Veillonella*, a strict anaerobe, are especially favored within biofilms.

Thus, aggregative streptococci are well suited for biofilm growth in the otic mucosa and for syntrophic cooperation with anaerobic partners.

Fig. 5: Adaptive responses promoting the establishment of otic trophic webs. (A) Biofilm biomass (crystal violet staining, measured as absorbance at 550 nm, A₅₅₀) of otic isolates in oxic (blue) and anoxic (white) cultures. The dashed circles identify two separate clusters of isolates with highest biofilm-forming abilities. (B) Correlation between biofilm formation and pH in oxic cultures showing . The circle highlights a cluster of strains with highest biofilm-forming activities and lowest pH. (C) Lactate and acetate production (mM) in stationary-phase cultures grown in oxic (black) and anoxic (white) media. The asterisks show significant differences between oxic and anoxic values ($p \leq 0.05$, *; $p \leq 0.01$, **; $p \leq 0.001$, ***). All data points in A–C are average values of three independent biological experiments and are color-coded for *Streptococcus* (gray), *Staphylococcus* (orange), *Neisseria* (purple) and actinobacterial genera *Micrococcus* and *Corynebacterium* (green).

Antagonistic interactions of otic streptococci with common otopathogens

Commensal oral streptococci mediate intra- and interspecies antagonistic interactions in oral biofilms that are critical to dental and mucosal health (18). Given their oral ancestry, we screened the otic streptococci for their ability to inhibit the growth of known otopathogens (*Streptococcus pneumoniae*, *Moraxella catharralis*, and non-typeable *Haemophilus influenzae*). For these assays, we followed the same protocol as in other plate assays and spot-plated overnight cultures on TSA plates before incubating them at 37°C for 24 h to allow the colonies to grow. We then covered the plates with a soft (0.7%) agar overlay containing a diluted cell suspension of each otopathogen in a growth medium suitable for their growth. After incubation of the overlaid plates for an additional 24 h, we examined the overlays for zones of growth inhibition on top and around the underlying streptococcal colonies. Growth inhibition in this assay may be the result of nutrient competition, secretion of growth inhibitors or both. Fig. 6

shows representative plate assays for all the otic strains against each otopathogen. Notably, all the otic streptococci inhibited the growth of *S. pneumoniae* and *M. catarrhalis*, producing large zones of clearing beyond the colony edge. The extended region of growth inhibition in the soft-agar overlay is consistent with the secretion of a diffusible inhibitory compound. We also observed antagonistic effects against *H. influenzae*, but they were less pronounced and strain-specific (Fig. 6).

Fig. 6: Growth inhibitory effect of otic streptococci against common otopathogens. TSA plates containing 24-h colonies of the otic streptococci were incubated for 24h with soft-agar overlays of the otopathogens *Streptococcus pneumoniae*, *Moraxella catarrhalis*, *Haemophilus influenzae* and *Staphylococcus aureus*. The plates show clear areas of growth inhibition of the otopathogen on top and/or around antagonistic streptococcal colonies underneath (scale bar, 0.5 cm). The symbols indicate average size of the growth inhibition halo around the underlying streptococcal colony in triplicate plate assays (+, <0.4; ++, >0.4; +/-, ~0.1 but not always reproducible).

We also used the plate assay to screen for potential antagonism of the otic streptococci towards the nasopharyngeal staphylococci that disperse in the aerodigestive tract. As a test strain, we used *S. aureus* subsp. *aureus* JE2 (50), a plasmid-cured derivative of the epidemic community-associated methicillin-resistant *S. aureus* (CA-MRSA) isolate USA300 (51). We observed antagonism by all the non-salivarius isolates (Fig. 6), suggesting species-specific mechanism for growth inhibition by these streptococcal groups (*S. pseudopneumoniae*, *S.*

parasanguinis and *S. agalactiae*). The ability of non-salivarius streptococci to inhibit the growth of *S. aureus* is not uncommon. Despite being catalase positive, *S. aureus* is sensitive to hydrogen peroxide produced by *S. pneumoniae* in the nasal mucosa (52). This is because hydrogen peroxide is converted into a highly toxic hydroxyl radical ($\cdot\text{OH}$) that rapidly kills *S. aureus* (53). The fact that the closest relatives to the non-salivarius otic streptococci all have catalase-independent mechanisms for anti-oxidative stress resistance (54) and release hydrogen peroxide as a byproduct of their metabolism (55-57) suggests similar mechanisms for interspecies interference with *S. aureus*.

Discussion

The recovery from otic secretions of close relatives of nasal and oral bacteria (Fig. 2) highlights the role that saliva aerosols play in the dispersal of bacteria throughout the aerodigestive tract. Human saliva carries bacteria shed from oral surfaces such as teeth and gums and spreads them to distant mucosae (58, 59). The constant flux of saliva to the oropharynx (back of the throat) facilitates the formation of aerosols and their carriage to the middle ear every time the Eustachian tube opens and exhaled air is drawn in (5). Not surprisingly, phylogenetic analysis of full-length 16S rRNA sequences recovered from 19 otic cultivars resolved close evolutionary relationships with species that reside or transiently disperse in the oral cavity (Fig. 2). Particularly important were the ancestral ties between the otic streptococci, the most prominent residents of the middle ear communities (5), and pioneer species of oral biofilms (Fig. 2). Most of the otic streptococci were closely related to *S. salivarius*, one of the first colonizers of the human oral cavity after birth and an abundant commensal throughout the life of the host (60). This bacterium disperses in the aerodigestive tract via saliva and forms aggregates that survive stomach passage (61). This allows Salivarius species to enter the small intestine and colonize its mucosa (62). The abundance of *S. salivarius* in saliva also increases its dispersal potential in saliva aerosols, the primary mechanism for seeding of the otic mucosa (5). Aggregation

facilitates immunoescape and successful colonization of the otic mucosa. It also promotes coaggregation with anaerobic syntrophic partners and the formation of otic trophic webs (Fig. 1) that mirror those described in oral biofilms. Additionally, oral *S. salivarius* strains mediate antagonistic interactions with virulent streptococci that are critical to prevent tooth decay, periodontal disease, and the spread of respiratory pathogens such as the otopathogen *S. pneumoniae* (63, 64). We see similar interspecies interference of otic *S. salivarius* strains towards common otopathogens (Fig. 6), suggesting similar roles for these middle ear residents in disease prevention.

The non-salivarius otic streptococci were also close relatives of oral species (Fig. 2). For example, one of the isolates (L0020-02) was closely related to *S. parasanguinis*, a bacterium that groups with species of the Mitis group based on 16S rRNA gene sequence analysis and shares with them many of their phenotypic characteristics (27). Like *S. salivarius*, *S. parasanguinis* is one of the early colonizers of the oral cavity (18) and disperses in saliva (65). It produces fimbriae to firmly attach to the biofilms (66), which facilitates its co-dispersal in syntrophic oral aggregates. Also recovered from otic secretions were strains closely related to streptococcal commensals such as *S. pseudopneumoniae* (L0023-02; Viridans group) and *S. agalactiae* (L0023-01 and L0023-03; GBS group). Although the oral ancestors of these otic streptococci have been linked to infective processes in the aerodigestive tract and other body sites (67, 68), the otic relatives readily inhibited the growth of the three most common otopathogens (*S. pneumoniae*, *M. catarrhalis*, and non-typeable *H. influenzae*) (Fig. 6). Furthermore, they were the only otic streptococci that interfered with the growth of *S. aureus* (Fig. 6). Antagonism towards *S. aureus* may involve the production of hydrogen peroxide as a metabolic byproduct, as noted for related oral streptococcal species (55-57). Hydrogen peroxide also functions as a signaling molecule for the co-aggregation of non-salivarius streptococci in syntrophic biofilms (56). Future studies will need to evaluate the role of these streptococcal

lineages in producing hydrogen peroxide as a signal for intra and interspecies co-aggregation and as an antagonist of bacterial competitors.

The results presented in this study identified physiological traits of the otic streptococci that could facilitate the colonization of the middle ear mucosa and the formation of syntrophic biofilms. The otic streptococci were temperate swimmers on soft agar plates (Fig. 3), a motile behavior that could allow them to penetrate the otic mucus layer and reach the underlying mucosal epithelium. However, not all the isolates secreted surfactants (Table 2). While endogenous surfactants are often needed for bacteria swarming on semisolid agar surfaces, they are not always needed for efficient swarming through the native mucus layers (20). This is particularly true for bacteria colonizing the middle ear mucosa, where host surfactants abound (69). Furthermore, surfactant production by bacterial colonizers may be undesirable in the otic mucosa as it would change the rheological properties of the mucus and interfere with critical mucosal functions such as antimicrobial activity, immunomodulation and Eustachian tube mechanics (69). The latter is carefully controlled by the secretion of host surfactants that regulate the viscosity and surface tension of the mucus layer (70). This chemical mechanism ensures that the surface tension of the mucus is kept sufficiently low (58 mN/m) to facilitate the rhythmic aperture of the collapsed Eustachian tube and adequate ventilation and decompression of the tympanic cavity (69). Disruption of surfactant homeostasis increases the pressure needed to open the Eustachian tube, risking barotrauma and making the middle ear mucosa more vulnerable to infections (69).

An important finding of our study was the identification of similar phenotypic traits among streptococcal and staphylococcal cultivars recovered from otic secretions that could give them both a competitive advantage during the colonization of the middle ear mucosa. For example, the streptococcal and staphylococcal isolates grew optimally with and without oxygen (Fig. 4A) and secreted mucins and proteases (Table 3), giving them a competitive advantage for growth and reproduction in the middle ear mucosa. Moreover, both groups had fermentative

metabolisms and produced lactate (Fig. 5C), a critical intermediate in the syntrophic otic communities (5). This contrasts with otic *Neisseria* (L0020-05 and L0020-06) closely related to *N. flavescens* and *N. subflava*, whose microaerophilic metabolism led to poor growth and extensive flocculation in oxic broth (Fig. 4A). These two *Neisseria* species form a separate clade with members of the family Neisseriaceae that populate the tongue dorsum (71) and although they readily disperse via saliva into the oropharynx (32), they are not positively selected in the middle ear (5). Furthermore, the *Neisseria* cultivars were along with the two actinobacterial isolates (*Micrococcus* spp. L0020-05 and *Corynebacterium* spp. L0020-06) the only cultivars that did not secrete proteases on casein plates (Table 3). Furthermore, they did not form robust biofilms (Fig. 5A) nor did they produce fermentative byproducts such as lactate (Fig. 5C), a key metabolic intermediate in the otic trophic webs (Fig. 1). Not surprisingly, despite being the most abundant nasal phylum (38), actinobacterial sequences are poorly represented in otic secretions (5).

The most notable difference between the staphylococcal and streptococcal isolates was, however, the co-aggregative behavior of most *Streptococcus*, which is critical for the formation of microcolonies with anaerobic syntrophic partners (Fig. 1). Aggregative behavior may in fact be the most successful strategy of oral streptococci during the colonization of the middle ear mucosa. Most of the streptococcal isolates autoaggregated when growing on agar-solidified media (Fig. 3B), a phenotype associated with increased resistance to antimicrobials and immunoescape (72). Aggregation allows oral streptococci to recognize and partner with other bacteria during the formation of biofilms (49). Thus, streptococci coaggregate with actinomyces to colonize the tooth surface and recruit other bacteria during the formation of the dental plaque (18, 73). Metabolic interactions between lactic acid-producing strains of *Streptococcus* and *Veillonella* spp., which ferment lactate, are critical for coaggregation during the early stages of biofilm formation on oral surfaces (49). Fusobacteria mediate early coaggregation as well, forming physical bridges across the biofilms and promoting the attachment of non-

coaggregating aerobes and anaerobes (18, 73). These syntrophic interactions sustain the growth of the dental plaque throughout all dentition stages and the formation of subgingival biofilms in the predentate and postdentate states (59). The widespread presence and abundance of syntrophic co-aggregates in the oral cavity promotes their co-dispersal in saliva (74) and provides a mechanism for their co-immigration in saliva aerosols. These coaggregates readily establish otic trophic webs (Fig. 1) similar to those described in oral biofilms (5). Consistent with this, all the streptococcal isolates characterized in this study were highly aggregative (Fig. 4), formed robust biofilms (Fig. 5A) and produced lactate as primary byproduct of their fermentative metabolism (Fig. 4C). These adaptive traits allow streptococci to grow and reproduce in the middle ear mucosa with obligate anaerobic, syntrophic partners such as *Prevotella*, *Fusobacterium* and *Veillonella* (5). The syntrophic microcolonies metabolize and ferment host mucins and proteins in the otic mucosa (Fig. 1), indirectly controlling the viscoelastic properties of the mucus layer and Eustachian tube functionality (69). The detection of a differential gradient of mucin gene expression along the tympanic cavity and Eustachian tube (69) suggests a high degree of spatial heterogeneity in bacterial colonization as well. Shaped like an inverted flask (4), the posterior region of the Eustachian tube is more readily seeded with saliva aerosols during the cycles of tubal aperture. Concentration of streptococcal aggregates in this region closer to the nasopharyngeal opening of the Eustachian tube could provide increased protection against otopathogens, which typically reside in nasal reservoirs. Future research should therefore consider the mechanisms that allow otic streptococci to co-aggregate with syntrophic partners, their spatial distribution in the otic mucosa and antagonistic interactions with transient migrants. This knowledge is important to understand the functionality of the otic communities and how they influence host functions and the outcome of infections.

Methods

Bacterial strains and culture conditions.

The bacterial strains used in this study include 19 cultivars isolated from otic secretions (5). Briefly, the samples were collected with a single swab from the left and right nasopharyngeal openings of the Eustachian tube in 4 young (19-32 years old), healthy adults recruited as part of a larger study approved by the Michigan State University Biomedical and Physical Health Review Board (IRB # 17-502). The cultivars were isolated as single colonies on Tryptic Soy Agar (TSA) plates (30g/L of Tryptic Soy Broth from Sigma Aldrich and 15g/L of Bacto Agar from BD) grown at 37°C. The isolates were routinely grown overnight in 5 ml of Tryptic Soy Broth (TSB) at 37°C with gentle agitation. For growth studies, we transferred mid-log phase ($OD_{600} \sim 0.5$) TSB cultures twice (initial OD_{600} of 0.1) to prepare a stationary phase (~ 0.9 - 1.0 OD_{600}) inoculum for growth assays in Corning® 96-well clear round bottom TC-treated microplate (Corning 3799). Growth was initiated with the addition of 18 μ l of the inoculum to 162 μ l of TSB per well and monitored spectrophotometrically every 30 min (OD_{630} readings after 0.1 sec of gentle agitation) while incubating the plates at 37°C inside a PowerWave HT (BioTek) plate reader. Each plate contained a well with TSB medium without cells to use as a blank. Growth in anoxic medium was monitored in a similar way but in a plate reader housed inside an 855-ABC Portable Anaerobic Chamber (Plas Labs, Inc.) containing a headspace of $N_2:CO_2$ (80:20).

DNA sequencing and phylogenetic analyses.

For taxonomic and phylogenetic analyses, we grew 19 otic isolates (Table 1) in 2 ml of TSB at 37°C for 24 h and harvested the cells by centrifugation (25,000 x g for 5 min) in an Eppendorf 5417R refrigerated centrifuge prior to extracting the genomic DNA with a FastDNA™ Spin kit (MP Biomedicals). Library preparation with an Illumina Nextera kit and whole genome sequencing in an Illumina NextSeq 550 platform were at the Microbial Genome Sequencing Center (MiGS; Pittsburgh, PA). We used the FastQC tool from the Babraham Institute (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) for sequence quality control and Trimmomatics (75) for cleaning/trimming of the Illumina short

reads. After assembling the genomes *de novo* with the Spades assembler (76), we identified the 16S rRNA gene sequences in the contigs with the BAseic Rapid Ribosomal RNA Predictor (Barnap) (<https://github.com/tseemann/barnap>). The 16S rRNA gene sequences were deposited in the GenBank database under individual accession numbers (Table 1). We used the full-length 16S rRNA sequences to identify the closest species (% identity) in the GenBank database using the nucleotide Basic Local Alignment Search Tool (BLAST) at the U.S. National Center of Biological Information (NCBI) using an identity species cutoff value of 98.7% (23). We retrieved the 16S rRNA gene sequences from the closest type strains listed in the SILVA rRNA database (<https://www.arb-silva.de>) and aligned them to the otic sequences with the MUSCLE program in the MEGA X software (77). We used the alignment to build a maximum-likelihood phylogenetic tree and calculate bootstrap confidence values for each node using 1,000 replications. The tree shows bootstrap values above 50% (78).

Catalase assay. Frozen stocks of the otic isolates were directly streaked on 1.5% (w/v) TSA plates to grow individual colonies at 37°C overnight. We spread each colony onto a microscope slide and added a drop of freshly prepared 3% hydrogen peroxide. Catalase-positive strains breakdown the hydrogen peroxide into water and oxygen gas, which generates bubbles. Lack or weak production of bubbles is used to designate a strain as catalase-negative.

Swarming motility and surfactant detection assays. We screened each otic isolate for their ability to move on soft (0.5% and, when indicated, 0.4% w/v agar) TSA plates, as a modification of a previously described assay (79). For these assays, we first grew each isolate and the positive control (*P. aeruginosa* PA01) in TSB at 37°C overnight (OD₆₀₀ ~1) and prepared a diluted TSB inoculum (OD₆₀₀ 0.1). We pipetted a 5-μl drop of the diluted culture onto the surface of the soft agar plates and allowed it to absorb until completely dry (~30 min). We then incubated the plates at 37°C and photographed the areas of growth at 18, 42 and 62 h against a ruler using a dissecting scope (Leica MZ6) at a magnification of 0.8X and 1X. The photographs

were then analyzed with the ImageJ software (80) to measure the colony diameter over time and calculate the area expansion (swarming distance) from the initial inoculation spot.

We also screened the ability of the otic isolates to produce surfactants with a previously described atomized oil assay (40). For this, we plated a 5- μ l drop of the diluted TSB culture (OD₆₀₀ of 0.1) on agar-solidified (1.5% w/v) TSA medium, allowed the inoculum to absorb for ~30 min, and incubated the plates at 37°C for 24 h. Using an airbrush (type H; Paasche Airbrush Co., Chicago, IL), we applied a fine mist of mineral oil onto the plate surface. Surfactant-producing colonies readily display a halo of mineral oil dispersal whose size provides a semiquantitative measure of surfactant secretion (40). Photography and halo diameter visualization was as described above for swarming assays, except that we measured the size of the oil dispersal zone from the colony edge. All strains were tested in three independent swarming and surfactant assays plates to calculate the average and standard deviation values.

Protease and mucinase plate assays. We used TSA plates containing 5% lactose-free, skim milk (Fairlife, LLC) or 0.5% Type II porcine gastric mucin (Sigma Aldrich) to screen the otic isolates for mucinase and protease secretion, respectively, using *P. aeruginosa* PA01 as a positive control. For these assays, we spot-plated 5 μ l of overnight TSB cultures and incubated at 37°C for 24 h, as described earlier for the surfactant assays. Strains that secrete proteases to the medium degrade the milk's casein and produce a clear halo around the area of growth after 24 h of incubation. Mucinase producers have zones of mucin lysis around or under the colony that show as zones of discoloration after staining with 7ml of 0.1% amido black for 30 min and destaining with 14 ml of 2.5 M acetic acid for 30 min. When indicated, plates were incubated for 48 h to confirm emerging phenotypes. Each strain was tested in triplicate plates and each was photographed on a lightboard (A4 LED Light Box 9x12 Inch Light Pad, ME456) with an iPhone 11 at 2.4x magnification.

Organic acid detection in culture supernatant fluids.

We grew triplicate stationary phase cultures of the otic isolates in oxic and anoxic TSB medium at 37°C and harvested the culture supernatant fluids by centrifugation (14,000 rpm, 10 min). We measured the pH of the supernatant fluids (5 ml) with a pH probe (Thermo Scientific™ Orion™ 720A+ benchtop pH meter) and stored 1 ml of the samples at –20°C for chemical analyses by high performance liquid chromatography (HPLC). Once thawed, we filter-sterilized 250 µl of the supernatant fluid into 1-ml HPLC vials and measured their organic acid content in a Shimadzu 20A HPLC equipped with an Aminex HPX-87H column and a Micro-Guard cation H⁺ guard column (Bio-Rad, Hercules, CA) at 55°C, as previously described (81). As controls, we included samples with TSB medium and standard solutions of acetate, lactate and pyruvate (provided at 1, 2, 5, 10 and 20 mM concentrations).

Biofilm assays.

We used a previously described assay (82) to test the ability of the otic cultivars to form biofilms in Corning® 96-well clear round bottom TC-treated microplates (Corning 3799). We first grew overnight cultures in TSB with gentle agitation (~200 rpm) and used them to prepare a diluted cell suspension (OD₆₀₀ ~0.1) for inoculation (18 µl) into TSB medium (162 µl per well). Each isolate was tested in 8 replicate wells. After incubating the plates at 37°C for 24 h, we removed the planktonic culture, washed the wells with ddH₂O and stained the surface-attached cells with 0.1% (w/v) crystal violet. We then rinsed the wells with water and let the stained biofilms to dry overnight at room temperature before solubilizing the biofilm-associated crystal violet with 180 µl of 30% glacial acetic acid and measured the crystal violet in the solution spectrophotometrically at 550 nm (82).

Growth inhibition plate assays.

We screened the otic streptococcal isolates for their ability to inhibit the growth of bacterial species (*S. pneumoniae*, *M. catarrhalis*, and non-typeable *H influenzae*) commonly associated with infections of the middle ear (83). As test strains, we used *S. pneumoniae* ATCC 6303 and *M. catarrhalis* ATCC 25238 (from the laboratory strain

collection of Dr. Martha Mulks, Department of Microbiology and Molecular Genetics, Michigan State University) and a non-typeable *H. influenzae* (NTHi) strain isolated by Dr. Poorna Viswanathan in the teaching lab of the Department of Microbiology and Molecular Genetics (Michigan State University). The NTHi strain was confirmed prior to experimental use by multiplex PCR confirmation, as described previously (84). We also included for testing the laboratory strain *S. aureus* JE2, which was kindly provided by Dr. Neal Hammer (Department of Microbiology and Molecular Genetics, Michigan State University). The otic streptococci and *S. aureus* JE2 were routinely grown in 5 ml TSB at 37°C with gentle agitation to prepare overnight cultures for the plate assays. *S. pneumoniae* and *M. catarrhalis* were grown at 37°C overnight in 5 mL of brain heart infusion (BHI) broth (Sigma-Aldrich) without agitation. The NTHi reference strain of *H. influenzae* was also grown statically at 37°C but in supplemented BHI (sBHI) (85), which contains (per L): 30 g BHI, 0.01 mg hemin (Bovine, Sigma Aldrich), and 0.002 mg β -Nicotinamide adenine dinucleotide sodium salt (Sigma Aldrich). All incubations were in a 37°C incubator with a 5% CO₂ atmosphere except for *S. aureus*, which were in air.

We used the spot-on-lawn method (86) to investigate antagonistic interactions between the otic streptococci and test strains. We first spotted 5 μ l of a diluted (OD₆₀₀ 0.1) overnight culture of each streptococcal strain onto a 1.5% (w/v agar) TSA plate and allowed it to dry for 30 min at room temperature before incubating at 37°C for 24 h to grow the colonies. We then overlaid the plates with a warm (55°C) 8-ml layer of soft-agar (0.7%, w/v, final concentration) medium (TSA, BHI or sBHI) containing the test strain (OD₆₀₀ 0.1). The general procedure to make 0.7% agar overlays was to autoclave 6 ml of 1% agar-solidified growth medium, cool down the melted agar in a 55°C water bath, add 2 ml of the test strain culture to a final OD₆₀₀ of 0.1, and mix by inversion before pouring over the TSA plate surface with the otic colonies. To make sBHI overlays, we added the chemical supplements to 6 ml of warm (55°C), melted 1% (w/v) agar BHI before mixing with 2 ml of an overnight NTHi culture to a final OD₆₀₀ of 0.1. The overlays were allowed to solidify at room temperature before incubating for an additional 24 h at

37°C in an incubator with or without (*S. aureus* overlay) 5% CO₂. These culture conditions promoted the growth of the test strains as a turbid lawn in the overlays after 24 h, except for areas of growth inhibition (halos or clear zones) on top and around colonies of antagonistic streptococci growing underneath. At the end of the incubation period, we photographed the overlaid plates with a dissecting scope (0.63x objective) against a ruler and used the ImageJ program (4) to measure the size of the growth inhibition zone from the streptococcal colony edge underneath in triplicate biological replicates.

Acknowledgements

The authors would like to thank Dr. Michaela TerAverst and Nicholas Tefft at Michigan State University for assistance with the HPLC analyses and Drs. Batsirai Mabvakure and Heather Blankenship at the Michigan Department of Health and Human Services Bureau of Laboratory for guidance on Illumina short read assembly and contig gene analysis. We are also thankful to Drs. Martha Mulks, Poorna Viswanathan and Neal Hammer for providing the test strains used in the growth inhibition plate assays.

Funding

This research was funded by grants N00014-17-2678 and N00014-20-1-2471 from the Microbiome program at the Office of Naval Research (ONR) to GR. KJ acknowledges support from a summer 2020 G. D. Edith Hsiung and Margaret Everett Kimball Endowed fellowship from the department of Microbiology and Molecular Genetics at Michigan State University. The funders had no role in study design, data collection and analysis, decision to publish or preparation of manuscript.

References

1. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, Lakshmanan A, Wade WG. 2010. The human oral microbiome. *J Bacteriol* 192:5002-17.

- 674 2. Welch JLM, Dewhirst FE, Borisy GG. 2019. Biogeography of the oral microbiome: The
675 site-specialist hypothesis. *Annu Rev Microbiol* doi:10.1146/annurev-micro-090817-
676 062503.
- 677 3. Gao L, Xu T, Huang G, Jiang S, Gu Y, Chen F. 2018. Oral microbiomes: more and more
678 importance in oral cavity and whole body. *Protein Cell* 9:488-500.
- 679 4. Bluestone CD. 2005. The Eustachian Tube: Structure, function, and role in the middle
680 ear. BC Decker Inc, Hamilton, Ontario.
- 681 5. Lee J-Y, Jacob KM, Kashefi K, Reguera G. 2021. Oral seeding and niche-adaptation of
682 middle ear biofilms in health. *Biofilm* 3:100041.
- 683 6. Kandler O. 1983. Carbohydrate metabolism in lactic acid bacteria. *Antonie Van*
684 *Leeuwenhoek* 49:209-24.
- 685 7. Cotter PD, Hill C. 2003. Surviving the acid test: responses of gram-positive bacteria to
686 low pH. *Microbiol Mol Biol Rev* 67:429-53.
- 687 8. Delwiche EA, Pestka JJ, Tortorello ML. 1985. The *Veillonellae*: Gram-negative cocci with
688 a unique physiology. *Annu Rev Microbiol* 39:175-93.
- 689 9. Rios-Covian D, Salazar N, Gueimonde M, de los Reyes-Gavilan CG. 2017. Shaping the
690 metabolism of intestinal *Bacteroides* population through diet to improve human health.
691 *Front Microbiol* 8.
- 692 10. Palmer RJ, Diaz PI, Kolenbrander PE. 2006. Rapid succession within the *Veillonella*
693 population of developing human oral biofilm in situ. *J Bacteriol* 188:4117.
- 694 11. Blaak EE, Canfora EE, Theis S, Frost G, Groen AK, Mithieux G, Nauta A, Scott K, Stahl
695 B, van Harsselaar J, van Tol R, Vaughan EE, Verbeke K. 2020. Short chain fatty acids in
696 human gut and metabolic health. *Benef Microbes* 11:411-455.
- 697 12. Hall-Stoodley L, Hu FZ, Gieseke A, Nistico L, Nguyen D, Hayes J, Forbes M, Greenberg
698 DP, Dice B, Burrows A, Wackym PA, Stoodley P, Post JC, Ehrlich GD, Kerschner JE.

2006. Direct detection of bacterial biofilms on the middle-ear mucosa of children with chronic otitis media. JAMA 296:202-11.
13. Tonnaer EL, Mylanus EA, Mulder JJ, Curfs JH. 2009. Detection of bacteria in healthy middle ears during cochlear implantation. Arch Otolaryngol Head Neck Surg 135:232-7.
14. Janssen WJ, Stefanski AL, Bochner BS, Evans CM. 2016. Control of lung defence by mucins and macrophages: ancient defence mechanisms with modern functions. Eur Respir J 48:1201-1214.
15. Honjo I, Hayashi M, Ito S, Takahashi H. 1985. Pumping and clearance function of the eustachian tube. Am J Otolaryngol 6:241-244.
16. Sade J. 1971. Mucociliary flow in the middle ear. Ann Otol Rhinol Laryngol 80:336-41.
17. Siegel SJ, Weiser JN. 2015. Mechanisms of bacterial colonization of the respiratory tract. Annu Rev Microbiol 69:425-444.
18. Kolenbrander PE. 2000. Oral microbial communities: biofilms, interactions, and genetic systems. Annu Rev Microbiol 54:413-37.
19. Reddy MS, Murphy TF, Faden HS, Bernstein JM. 1997. Middle ear mucin glycoprotein: Purification and interaction with nontypable *Haemophilus influenzae* and *Moraxella catarrhalis*. Otolaryngol Head Neck Surg 116:175-80.
20. Partridge JD, Harshey RM. 2013. Swarming: Flexible roaming plans. J Bacteriol 195:909-918.
21. Tailford LE, Crost EH, Kavanaugh D, Juge N. 2015. Mucin glycan foraging in the human gut microbiome. Front Genet 6:81.
22. Bomar L, Brugger SD, Lemon KP. 2018. Bacterial microbiota of the nasal passages across the span of human life. Curr Opin Microbiol 41:8-14.
23. Stackebrandt E, Ebers J. 2006. Taxonomic parameters revisited: tarnished gold standards. Microbiol Today 33:152–155.

24. Munoz R, Yarza P, Ludwig W, Euzéby J, Amann R, Schleifer KH, Glockner FO, Rossello-Mora R. 2011. Release LTPs104 of the All-Species Living Tree. Syst Appl Microbiol 34:169-70.
25. Yarza P, Richter M, Peplies J, Euzéby J, Amann R, Schleifer KH, Ludwig W, Glockner FO, Rossello-Mora R. 2008. The All-Species Living Tree project: A 16S rRNA-based phylogenetic tree of all sequenced type strains. Syst Appl Microbiol 31:241-250.
26. Delorme C, Abraham AL, Renault P, Guédon E. 2015. Genomics of *Streptococcus salivarius*, a major human commensal. Infect Genet Evol 33:381-92.
27. Facklam R. 2002. What happened to the streptococci: Overview of taxonomic and nomenclature changes. Clin Microbiol Rev 15:613-+.
28. Huch M, De Bruyne K, Cleenwerck I, Bub A, Cho GS, Watzl B, Snauwaert I, Franz CM, Vandamme P. 2013. *Streptococcus rubneri* sp. nov., isolated from the human throat. Int J Syst Evol Microbiol 63:4026-32.
29. Ohara-Nemoto Y, Haraga H, Kimura S, Nemoto TK. 2008. Occurrence of staphylococci in the oral cavities of healthy adults and nasal oral trafficking of the bacteria. J Med Microbiol 57:95-9.
30. McCormack MG, Smith AJ, Akram AN, Jackson M, Robertson D, Edwards G. 2015. *Staphylococcus aureus* and the oral cavity: an overlooked source of carriage and infection? Am J Infect Control 43:35-7.
31. Knapp JS, Hook EW, 3rd. 1988. Prevalence and persistence of *Neisseria cinerea* and other *Neisseria* spp. in adults. J Clin Microbiol 26:896-900.
32. Liu G, Tang CM, Exley RM. 2015. Non-pathogenic *Neisseria*: Members of an abundant, multi-habitat, diverse genus. Microbiology 161:1297-1312.
33. Bassis CM, Erb-Downward JR, Dickson RP, Freeman CM, Schmidt TM, Young VB, Beck JM, Curtis JL, Huffnagle GB. 2015. Analysis of the upper respiratory tract

- microbiotas as the source of the lung and gastric microbiotas in healthy individuals. mBio 6:e00037-15.
34. Kocur M, Kloos WE, Schleifer K-H. 2006. The genus *Micrococcus*, p 961-971. In Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (ed), The Prokaryotes: Volume 3: Archaea Bacteria: Firmicutes, Actinomycetes doi:10.1007/0-387-30743-5_37. Springer New York, New York, NY.
35. Zhao G-Z, Li J, Qin S, Zhang Y-Q, Zhu W-Y, Jiang C-L, Xu L-H, Li W-J. 2009. *Micrococcus yunnanensis* sp. nov., a novel actinobacterium isolated from surface-sterilized *Polyspora axillaris* roots. Int J Syst Evol Microbiol 59:2383-2387.
36. Kooker JM, Fox KF, Fox A. 2012. Characterization of *Micrococcus* strains isolated from indoor air. Mol Cell Probes 26:1-5.
37. Brugger SD, Bomar L, Lemon KP. 2016. Commensal–pathogen interactions along the human nasal passages. PLOS Pathogens 12:e1005633.
38. Koskinen K, Reichert JL, Hoier S, Schachenreiter J, Duller S, Moissl-Eichinger C, Schopf V. 2018. The nasal microbiome mirrors and potentially shapes olfactory function. Sci Rep 8:1296.
39. Köhler T, Curty LK, Barja F, van Delden C, Pechère J-C. 2000. Swarming of *Pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. J Bacteriol 182:5990-5996.
40. Burch AY, Shimada BK, Browne PJ, Lindow SE. 2010. Novel high-throughput detection method to assess bacterial surfactant production. Appl Environ Microbiol 76:5363-72.
41. Kaito C, Sekimizu K. 2007. Colony spreading in *Staphylococcus aureus*. J Bacteriol 189:2553-2557.
42. Tsompanidou E, Denham EL, Becher D, de Jong A, Buist G, van Oosten M, Manson WL, Back JW, van Dijk JM, Dreisbach A. 2013. Distinct roles of phenol-soluble modulins

- in spreading of *Staphylococcus aureus* on wet surfaces. Appl Environ Microbiol 79:886-95.
43. Pollitt EJG, Diggle SP. 2017. Defining motility in the *Staphylococci*. Cell Mol Life Sci 74:2943-2958.
44. Bible AN, Khalsa-Moyers GK, Mukherjee T, Green CS, Mishra P, Purcell A, Aksenova A, Hurst GB, Alexandre G. 2015. Metabolic adaptations of *Azospirillum brasilense* to oxygen stress by cell-to-cell clumping and flocculation. Appl Environ Microbiol 81:8346-57.
45. Wandersman C. 1989. Secretion, processing and activation of bacterial extracellular proteases. Mol Microbiol 3:1825-31.
46. Khan R, Petersen FC, Shekhar S. 2019. Commensal bacteria: An emerging player in defense against respiratory pathogens. Front Immunol 10.
47. Smith PA, Sherman JM. 1942. The lactic acid fermentation of streptococci. J Bacteriol 43:725-731.
48. Sheng J, Marquis RE. 2006. Enhanced acid resistance of oral streptococci at lethal pH values associated with acid-tolerant catabolism and with ATP synthase activity. FEMS Microbiology Lett 262:93-98.
49. Mashima I, Nakazawa F. 2015. Interaction between *Streptococcus* spp. and *Veillonella tobetsuensis* in the early stages of oral biofilm formation. J Bacteriol 197:2104-2111.
50. Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW. 2013. A genetic resource for rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus* genes. mBio 4:e00537-12.
51. Diep BA, Gill SR, Chang RF, Phan TH, Chen JH, Davidson MG, Lin F, Lin J, Carleton HA, Mongodin EF, Sensabaugh GF, Perdreau-Remington F. 2006. Complete genome sequence of USA300, an epidemic clone of community-acquired methicillin-resistant *Staphylococcus aureus*. Lancet 367:731-9.

52. Regev-Yochay G, Trzcinski K, Thompson CM, Malley R, Lipsitch M. 2006. Interference between *Streptococcus pneumoniae* and *Staphylococcus aureus*: In vitro hydrogen peroxide-mediated killing by *Streptococcus pneumoniae*. J Bacteriol 188:4996-5001.
53. Wu X, Gordon O, Jiang W, Antezana BS, Angulo-Zamudio UA, Del Rio C, Moller A, Brissac T, Tierney ARP, Warncke K, Orihuela CJ, Read TD, Vidal JE. 2019. Interaction between *Streptococcus pneumoniae* and *Staphylococcus aureus* generates $\cdot\text{OH}$ radicals that rapidly kill *Staphylococcus aureus* strains. J Bacteriol 201.
54. Chen Z, Wang X, Yang F, Hu Q, Tong H, Dong X. 2017. Molecular insights into hydrogen peroxide-sensing mechanism of the metalloregulator MntR in controlling bacterial resistance to oxidative stresses. J Biol Chem 292:5519-5531.
55. Wilson CB, Weaver WM. 1985. Comparative susceptibility of group B streptococci and *Staphylococcus aureus* to killing by oxygen metabolites. J Infect Dis 152:323-9.
56. Duan D, Scofield JA, Zhou X, Wu H. 2016. Fine-tuned production of hydrogen peroxide promotes biofilm formation of *Streptococcus parasanguinis* by a pathogenic cohabitant *Aggregatibacter actinomycetemcomitans*. Environ Microbiol 18:4023-4036.
57. Redanz S, Cheng X, Giacaman RA, Pfeifer CS, Merritt J, Kreth J. 2018. Live and let die: Hydrogen peroxide production by the commensal flora and its role in maintaining a symbiotic microbiome. Mol Oral Microbiol 33:337-352.
58. Crielaard W, Zaura E, Schuller AA, Huse SM, Montijn RC, Keijser BJ. 2011. Exploring the oral microbiota of children at various developmental stages of their dentition in the relation to their oral health. BMC Med Genomics 4:22.
59. Mason MR, Chambers S, Dabdoub SM, Thikkurissy S, Kumar PS. 2018. Characterizing oral microbial communities across dentition states and colonization niches. Microbiome 6:67.
60. Kaci G, Goudercourt D, Dennin V, Pot B, Doré J, Ehrlich SD, Renault P, Blottière HM, Daniel C, Delorme C. 2014. Anti-Inflammatory properties of *Streptococcus salivarius*, a

- commensal bacterium of the oral cavity and digestive tract. Appl Environ Microbiol 80:928-934.
61. Hakalehto E, Vilpponen-Salmela T, Kinnunen K, von Wright A. 2011. Lactic acid bacteria enriched from human gastric biopsies. ISRN Gastroenterol 2011:109183.
62. Wang M, Ahrné S, Jeppsson B, Molin G. 2005. Comparison of bacterial diversity along the human intestinal tract by direct cloning and sequencing of 16S rRNA genes. FEMS Microbiol Ecol 54:219-31.
63. Van Hoogmoed CG, Geertsema-doornbusch GI, Teughels W, Quirynen M, Busscher HJ, Van der Mei HC. 2008. Reduction of periodontal pathogens adhesion by antagonistic strains. Oral Microbiol Immunol 23:43-48.
64. Van Hoogmoed CG, Geertsema-Doornbusch GI, Teughels W, Quirynen M, Busscher HJ, Van der Mei HC. 2008. Reduction of periodontal pathogens adhesion by antagonistic strains. Oral Microbiol Immunol 23:43-8.
65. Wang K, Lu W, Tu Q, Ge Y, He J, Zhou Y, Gou Y, Van Nostrand JD, Qin Y, Li J, Zhou J, Li Y, Xiao L, Zhou X. 2016. Preliminary analysis of salivary microbiome and their potential roles in oral lichen planus. Sci Rep 6:22943.
66. Garnett JA, Simpson PJ, Taylor J, Benjamin SV, Tagliaferri C, Cota E, Chen Y-YM, Wu H, Matthews S. 2012. Structural insight into the role of Streptococcus parasanguinis Fap1 within oral biofilm formation. Biochem Biophys Res Commun 417:421-426.
67. Keith ER, Podmore RG, Anderson TP, Murdoch DR. 2006. Characteristics of *Streptococcus pseudopneumoniae* isolated from purulent sputum samples. J Clin Microbiol 44:923-7.
68. Raabe VN, Shane AL. 2019. Group B Streptococcus (*Streptococcus agalactiae*). Microbiol Spectr 7:10.1128/microbiolspec.GPP3-0007-2018.
69. McGuire JF. 2002. Surfactant in the middle ear and eustachian tube: a review. Int J Pediatr Otorhinolaryngol 66:1-15.

70. Ghadiali SN, Banks J, Swarts JD. 2002. Effect of surface tension and surfactant administration on Eustachian tube mechanics. *J Appl Phys* 93:1007-1014.
71. Donati C, Zolfo M, Albanese D, Tin Truong D, Asnicar F, Iebba V, Cavalieri D, Jousson O, De Filippo C, Huttenhower C, Segata N. 2016. Uncovering oral *Neisseria* tropism and persistence using metagenomic sequencing. *Nature Microbiol* 1:16070.
72. Trunk T, Khalil HS, Leo JC. 2018. Bacterial autoaggregation. *AIMS Microbiol* 4:140-164.
73. Kriebel K, Hieke C, Müller-Hilke B, Nakata M, Kreikemeyer B. 2018. Oral biofilms from symbiotic to pathogenic interactions and associated disease –Connection of periodontitis and rheumatic arthritis by peptidylarginine deiminase. *Front Microbiol* 9.
74. Belstrom D. 2020. The salivary microbiota in health and disease. *J Oral Microbiol* 12:1723975.
75. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114-20.
76. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455-477.
77. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Mol Biol Evol* 35:1547-1549.
78. Hall BG. 2018. Phylogenetic trees made easy: a how-to manual, Fifth edition. ed. Sinauer Associates is imprint of Oxford University Press, New York.
79. Quinones B, Dulla G, Lindow SE. 2005. Quorum sensing regulates exopolysaccharide production, motility, and virulence in *Pseudomonas syringae*. *Mol Plant Microbe Interact* 18:682-93.
80. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9:671-5.

81. Duhl KL, Tefft NM, TerAvest MA. 2018. *Shewanella oneidensis* MR-1 utilizes both sodium- and proton-pumping NADH dehydrogenases during aerobic growth. Appl Environ Microbiol 84:e00415-18.
82. Merritt JH, Kadouri DE, O'Toole GA. 2005. Growing and analyzing static biofilms. Curr Protoc Microbiol Chapter 1:Unit 1B 1.
83. Schilder AG, Chonmaitree T, Cripps AW, Rosenfeld RM, Casselbrant ML, Haggard MP, Venekamp RP. 2016. Otitis media. Nat Rev Dis Primers 2:16063.
84. World Health O, Centers for Disease C, Prevention. 2011. Laboratory methods for the diagnosis of meningitis caused by *Neisseria meningitidis*, *Streptococcus pneumoniae*, and *Haemophilus influenzae*: WHO manual, 2nd ed ed. World Health Organization, Geneva.
85. Poje G, Redfield RJ. 2003. General methods for culturing *Haemophilus influenzae*. Methods Mol Med 71:51-6.
86. Schillinger U, Lücke FK. 1989. Antibacterial activity of *Lactobacillus sake* isolated from meat. Appl Environ Microbiol 55:1901-1906.

Table 1: Taxonomic classification (reference strain) of otic strains based on the % identity (ID) of their full-length 16S rRNA sequence.

Strain	GenBank no.	Reference Strain (Accession; % ID)
<i>Streptococcus</i>		
L0020-02	MW866489	<i>Streptococcus parasanguinis</i> (NR_024842.1; 99.47)
L0021-01	MW866494	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0021-04	MW866496	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0021-05	MW866497	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0022-03	MW866499	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0022-04	MW866500	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0022-05	MW866501	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0022-06	MW866502	<i>Streptococcus salivarius</i> (NR_042776.1; 99.81)
L0023-01	MW866503	<i>Streptococcus agalactiae</i> (NR_040821.1; 100)
L0023-02	MW866504	<i>Streptococcus oralis</i> (NR_117719.1; 99.47)
L0023-03	MW866505	<i>Streptococcus agalactiae</i> (NR_040821.1; 100)
<i>Staphylococcus</i>		
L0020-04	MW866491	<i>Staphylococcus hominis</i> (NR_036956.1; 99.61)
L0021-02	MW866495	<i>Staphylococcus aureus</i> (NR_037007.2; 99.87)
L0021-06	MW866498	<i>Staphylococcus saccharolyticus</i> (NR_113405.1; 99.4)
<i>Micrococcus</i>		
L0020-05	MW866492	<i>Micrococcus luteus</i> (NR_075062.2, 99.61)
<i>Corynebacterium</i>		
L0020-06	MW866493	<i>Corynebacterium pseudodiphthericum</i> (NR_042137.1; 99.47)
<i>Neisseria</i>		
L0020-03	MW866490	<i>Neisseria perflava</i> (NR_114694; 99.93)
L0023-05	MW866507	<i>Neisseria perflava</i> NR_117694.1; 99.74)
L0023-06	MW866506	<i>Neisseria perflava</i> NR_117694.1; 99.74)

Table 2. Coaggregation, swarming motility and surfactant production of otic isolates in reference to positive control (*P. aeruginosa* PA01).

Species/closest relative	Strain	Aggregation ^a	Surfactant ^b	Swarming ^c		
				18 h	42 h	62 h
<i>Streptococcus</i>						
<i>S. parasanguinis</i>	L0020-02	+	–	–	–	0.07 (0.10)
<i>S. salivarius</i>	L0021-01	+	0.23 (0.19)	–	–	0.16 (0.05)
<i>S. salivarius</i>	L0021-04	+	0.46 (0.09)	–	–	0.05 (0.12)
<i>S. salivarius</i>	L0021-05	+	–	–	–	0.05 (0.07)
<i>S. salivarius</i>	L0022-03	+	0.28 (0.19)	–	–	0.19 (0.01)
<i>S. salivarius</i>	L0022-04	+	–	–	–	0.20 (0.08)
<i>S. salivarius</i>	L0022-05	+	0.18 (0.12)	–	–	0.20 (0.03)
<i>S. salivarius</i>	L0022-06	+	0.09 (0.06)	–	–	0.18 (0.02)
<i>S. agalactiae</i>	L0023-01	+	–	–	0.21 (0.22)	0.23 (0.02)
<i>S. pseudopneumoniae</i>	L0023-02	+	–	–	–	0.07 (0.10)
<i>S. agalactiae</i>	L0023-03	+	0.31 (0.11)	–	0.19 (0.01)	0.20 (0.06)
<i>Staphylococcus</i>						
<i>S. hominis</i>	L0020-04	–	–	–	0.37 (0.01)	0.63 (0.08)
<i>S. aureus</i>	L0021-02	–	–	–	0.53 (0.01)	0.79 (0.01)
<i>S. epidermidis</i>	L0021-06	–	–	–	0.46 (0.03)	0.64 (0.02)
<i>Actinobacteria</i>						
<i>M. yunnanensis</i>	L0020-05	–	0.86 (0.13)	–	0.54 (0.05)	0.75 (0.04)
<i>C. pseudodiphthericum</i>	L0020-06	–	2.64 (0.05)	–	0.41 (0.06)	0.72 (0.03)
<i>Neisseria</i>						
<i>N. perflava</i>	L0020-03	–	–	–	0.44 (0.03)	0.66 (0.01)
<i>N. flavescens</i>	L0023-05	+	–	–	–	0.23 (0.32)
<i>N. flavescens</i>	L0023-06	+	–	–	–	0.25 (0.35)
<i>P. aeruginosa</i>	PA01	–	1.37 (0.06)	1.24 (0.35)	1.58 (0.53)	2.18 (0.65)

^a Aggregative (+) or uniform (–) growth of cultures spotted on 0.5% TSA plates.

^b Average (and standard deviation) of triplicate surfactant haloes (cm) measured as the zone of mineral oil dispersion around colonies grown at 37°C on 1.5% TSA plates. (–, not detected).

^c Average (and standard deviation) of triplicate swarming expansion zones (cm) around colonies grown at 37°C on soft agar (0.5%) TSA plates for 18, 42 and 62 h. (–, not detected).

Table 3. Protease and mucinase enzymatic activity of otic isolates in reference to positive control (*P. aeruginosa* PA01). Presence (+) or absence (–) of a halo of degradation in TSA plates supplemented with skim milk (protease assay) or mucin (mucinase assay) after 24 h of growth. The presence of a faint halo is indicated with “+/-”.

Species/closest relative	Strain	Protease	Mucinase
<i>Streptococcus</i>			
<i>S. parasanguinis</i>	L0020-02	–	+
<i>S. salivarius</i>	L0021-01	–	+/-
<i>S. salivarius</i>	L0021-04	+	+
<i>S. salivarius</i>	L0021-05	+	+
<i>S. salivarius</i>	L0022-03	+	+/-
<i>S. salivarius</i>	L0022-04	+	+/-
<i>S. salivarius</i>	L0022-05	+	+
<i>S. salivarius</i>	L0022-06	+	+
<i>S. agalactiae</i>	L0023-01	+	+
<i>S. pseudopneumoniae</i>	L0023-02	–	+
<i>S. agalactiae</i>	L0023-03	+	+
<i>Staphylococcus</i>			
<i>S. hominis</i>	L0020-04	+	+
<i>S. aureus</i>	L0021-02	+	+
<i>S. epidermidis</i>	L0021-06	+	+
<i>Actinobacteria</i>			
<i>Micrococcus yunnanensis</i>	L0020-05	–	+
<i>Corynebacterium pseudodiphthericum</i>	L0020-06	–	+
<i>Neisseria</i>			
<i>N. perflava</i>	L0020-03	–	+
<i>N. flavescens</i>	L0023-05	–	+
<i>N. flavescens</i>	L0023-06	–	+
<i>P. aeruginosa</i>	PA01	+	+