The temperature-regulated DEAD-box RNA helicase CrhR interactome: Autoregulation and photosynthesis-related transcripts

Anzhela Migur^{1,§}, Florian Heyl², Janina Fuss^{3,§}, Afshan Srikumar⁴, Bruno Huettel³,
Claudia Steglich¹, Jogadhenu S. S. Prakash⁴, Richard Reinhardt³, Rolf Backofen²,
George W. Owtrim⁵, Wolfgang R. Hess^{1*}

¹Faculty of Biology, University of Freiburg, Schänzlestr. 1, D-79104 Freiburg, Germany

²Department of Computer Science, University of Freiburg, Georges-Koehler-Allee 106,
D-79110 Freiburg, Germany

³Max Planck-Genome-Centre Cologne, Carl-von-Linné-Weg 10, D-50829 Köln,
Germany

⁴Department of Biotechnology & Bioinformatics, School of Life Sciences, University of
Hyderabad, Hyderabad, India

⁵Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada
T6G 2E9

Running title: CrhR DEAD-box RNA helicase interactome

**Corresponding author:* Wolfgang R. Hess; Phone: 0049-761-2032796; Fax: 0049-761-
2032745; Email: wolfgang.hess@biologie.uni-freiburg.de

[§]Current address: RNA Synthetic Biology Group, Helmholtz Institute for RNA-based
Infection Research (HIRI), Würzburg, Germany

[§]Current address: Institute of Clinical Molecular Biology, University of Kiel, Kiel,
Germany

27 Email addresses:

28 anzhela.migur@helmholtz-hiri.de

29 j.fuss@ikmb.uni-kiel.de

30 reinhardt@mpipz.mpg.de

31 huettel@mpipz.mpg.de

32 claudia.steglich@biologie.uni-freiburg.de

33 backofen@informatik.uni-freiburg.de

34 heylyf@informatik.uni-freiburg.de

35 syamsunderp@yahoo.com

36 afshan.sri17@gmail.com

37 gowttrim@ualberta.ca

38 wolfgang.hess@biologie.uni-freiburg.de

39

40 Date of submission: July 26, 2021

41 Number of tables and figures: 7

42 Total Word count: ~12,800

43 Word count from the start of the introduction to the end of the Discussion - excluding

44 Materials and methods, References, Tables and Figure Legends: 5648

45 Supplementary data: 3 figures, 5 tables

46

47 *Conflicts of Interest:* The authors declare the absence of conflicts of interest.

48

49

Highlight

The cyanobacterial DEAD-box RNA helicase CrhR binds mainly photosynthesis-associated and redox-controlled transcripts connecting its regulation, localization and phenotypes of mutants for the first time with a set of potential RNA targets.

Abstract

RNA helicases play crucial functions in RNA biology. In plants, RNA helicases are encoded by large gene families, performing roles in abiotic stress responses, development, the post-transcriptional regulation of gene expression as well as house-keeping functions. Several of these RNA helicases are targeted to the organelles, mitochondria and chloroplasts. Cyanobacteria are the direct evolutionary ancestors of plant chloroplasts. The cyanobacterium *Synechocystis* 6803 encodes a single DEAD-box RNA helicase, CrhR, that is induced by a range of abiotic stresses, including low temperature. Though the $\Delta crhR$ mutant exhibits a severe cold-sensitive phenotype, the physiological function(s) performed by CrhR have not been described. To identify transcripts interacting with CrhR, we performed RNA co-immunoprecipitation with extracts from a *Synechocystis* *crhR* deletion mutant expressing the FLAG-tagged native CrhR or a K57A mutated version with an anticipated enhanced RNA binding. The composition of the interactome was strikingly biased towards photosynthesis-associated and redox-controlled transcripts. A transcript highly enriched in all experiments was the *crhR* mRNA, suggesting an auto-regulatory molecular mechanism. The identified interactome explains the described physiological role of CrhR in response to the redox poise of the photosynthetic electron transport chain and characterizes CrhR as an enzyme with a diverse range of transcripts as molecular targets.

Keywords: chloroplasts, CrhR RNA helicase, cyanobacteria, gene expression regulation, photosynthesis, small regulatory RNA, redox regulation, RNA:RNA interaction

79 **Abbreviations:** co-IP, co-immunoprecipitation; CrhR, cyanobacterial RNA helicase
80 redox; ETC, electron transport chain, HLR1, high light regulatory 1; OCP, orange
81 carotenoid protein; sRNA, small regulatory RNA; TSS, transcriptional start site

82

83

84

Introduction

DEAD-box RNA helicases

The synthesis, maturation, modification and decay of RNA molecules and their interaction with each other and with other cellular components is central to the molecular basis of life. The largest family of enzymes involved in the metabolism of RNA molecules are enzymes belonging to the Superfamily I (SF1) and II (SF2) RNA helicases (Bourgeois *et al.*, 2016). DEAD-box RNA helicases, named after the conserved DEAD (Asp-Glu-Ala-Asp) amino acid motif in their core motif, form the largest and most complex group of SF2 RNA helicases (Jarmoskaite and Russell, 2011). DEAD-box RNA helicases are RNA-dependent ATPases (Rocak and Linder, 2004) characterized by the presence of twelve highly conserved motifs (Redder *et al.*, 2015). These motifs form the motor core of the helicase, which binds specifically adenine nucleotides and single-stranded RNA in a sequence-independent manner. The main function of these proteins is the conformational rearrangement of RNA by unwinding short double stranded regions. The unwinding reaction can be performed within one RNA molecule or between two duplex-forming RNAs. Besides the unwinding reaction, some RNA helicases are able to anneal single stranded RNAs (Chamot *et al.*, 2005; Yang and Jankowsky, 2005). In addition, some RNA helicases act as RNA clamps or facilitate RNA-protein complex dissociation without duplex unwinding (Jankowsky *et al.*, 2001; Fairman *et al.*, 2004).

Abiotic stresses and DEAD-box RNA helicases in plants and cyanobacteria

Cold stress, one of the most common stress conditions in nature, frequently induces the expression or activity of DEAD-box RNA helicases. At low temperature, RNA secondary structures are thermodynamically stabilized, which may interfere with their function. RNA helicases can rescue the RNA's functions by rearranging its secondary structures (Jones *et al.*, 1996). This role makes certain RNA helicases more relevant or even conditionally essential at low temperature. For instance, deletion of *csdA* or *srmB* in *Escherichia coli* leads to a cold-sensitive phenotype (Redder *et al.*, 2015). If all four

DEAD-box RNA helicase genes are deleted, *Bacillus subtilis* is not viable at low temperature (16°C), although the strain grows well at 37°C (Lehnik-Habrink *et al.*, 2013).

Most organisms encode several DEAD-box RNA helicases possessing non-complementary roles associated with a variety of physiological functions. Common functions in all organisms involve ribosome biogenesis, RNA turnover and translation, and responses to multiple stress conditions (Py *et al.*, 1996; de la Cruz *et al.*, 1999; Schneider and Schwer, 2001; Rogers *et al.*, 2002; Macovei *et al.*, 2012). Plant genomes typically possess larger and more diverse RNA helicase gene families than observed in other systems (Linder and Owttrim, 2009). Hence, their relevance in RNA secondary structure rearrangement under different environmental conditions or in plant development presents interesting lines for investigation. For example, *Arabidopsis thaliana* encodes 58 DEAD-box RNA helicases (Boudet *et al.*, 2001), many of which are essential as they are not functionally complementary (Mingam *et al.*, 2004). RNA helicases in plants fulfil roles in the defense against viruses (Wu and Nagy, 2020), abiotic stress responses to high salinity (Capel *et al.*, 2020), low temperature (Lu *et al.*, 2020; Wang *et al.*, 2020), and in development (Gong *et al.*, 2005). Interestingly, although not complementary, some plant helicases are involved in the same processes, suggesting multiple, independent RNA structure rearrangements are associated with a single physiological response (Huang *et al.*, 2016). In addition, several RNA helicases are targeted to the organelles, mitochondria and chloroplasts (Matthes *et al.*, 2007; Nawaz and Kang, 2017; Nawaz *et al.*, 2018). However, the expanded DEAD-box RNA helicase families make the precise functional characterization of RNA helicases challenging in plants.

Cyanobacteria are the direct evolutionary ancestors of plant chloroplasts (Mereschkowsky, 1905; Margulis, 1981; Martin and Kowallik, 1999; Ponce-Toledo *et al.*, 2017). The endosymbiosis of a cyanobacterium not only led to the chloroplast but also had pivotal impact on the composition of the plant nuclear genome (Martin *et al.*, 2002). Therefore, analysis of gene function in cyanobacteria is also informative at the higher plant level.

The cyanobacterium *Synechocystis* encodes a single DEAD-box RNA helicase, CrhR

The unicellular cyanobacterium *Synechocystis* sp. PCC 6803 (from here: *Synechocystis*) encodes the DEAD-box RNA helicase CrhR, for cyanobacterial RNA helicase redox (Rosana *et al.*, 2012a). As defined by a 50 amino acids sequence motif, CrhR is the archetype protein of a new clade within the DEAD-box RNA helicase family (Whitford *et al.*, 2021). In contrast to the situation in plants and also in many other bacteria, CrhR (*crhR/slr0083*) is the only DEAD-box RNA helicase encoded in *Synechocystis* (Redder *et al.*, 2015). Functionally, CrhR can therefore be expected to be of particular importance as it likely performs multiple functions that from an evolutionary perspective could have been distributed to different members of the complex family of RNA helicases that we observe in plants today (Kiefer *et al.*, 2020). *crhR* (*slr0083*) was originally characterized as a salt and cold-shock inducible protein (Vinnemeier and Hagemann, 1999; Kujat and Owtrim, 2000). Cold stress-inducible RNA helicases were also investigated in the cyanobacterium *Anabaena* sp. PCC 7120 (Chamot *et al.*, 1999) and *Synechococcus* sp. WH 7803 (Gierga *et al.*, 2012). Detailed analysis indicated that a low temperature stabilization of both transcript and protein contribute to the observed low temperature induction of CrhR (Rosana *et al.*, 2012a) and that a *crhR* inactivation mutant was severely impaired morphologically and physiologically at lower but not at higher temperatures (Rosana *et al.*, 2012b). In addition to low temperature, *crhR* is induced by a range of abiotic stresses that reduce the electron transport chain, independent of temperature shift (Vinnemeier and Hagemann, 1999; Kujat and Owtrim, 2000; Ritter *et al.*, 2020). The gene *crhR* (*slr0083*) is located in a dicistronic operon together with *rimO* (*slr0082*), whose putative protein product has 38% identity with RimO, a ribosomal protein S12 methylthiotransferase (UniProtKB P0AEI4), and 29% identity with the paralogous tRNA methylthiolase MiaB (Uni-ProtKB P0AEI1) from *E. coli* (Rosana *et al.*, 2020). The auto-regulated enhanced operon discoordination and processing of the *crhR* mRNA from the dicistronic operon RNA adds further complexity to its regulation (Rosana *et al.*, 2012; Rosana *et al.*, 2020). In addition, upon temperature upshift CrhR undergoes rapid repression via conditional proteolysis at the

post-translational level through an unknown mechanism (Tarassova *et al.*, 2014). In localization studies, CrhR localized to the thylakoid membranes and co-sedimented with degradosome and polysome complexes (Rosana *et al.*, 2016), consistent with the recent co-fractionation analysis using Grad-seq (Riediger *et al.*, 2021).

Although the molecular effects of *crhR* deletion or inactivation were studied at both the transcriptome (Prakash *et al.*, 2010; Georg *et al.*, 2019) and proteome level (Rowland *et al.*, 2011), the direct RNA targets of CrhR and interacting protein partners have not been identified. To monitor transcriptome-wide binding of CrhR, we immuno-precipitated the RNA species interacting with a FLAG-tagged version of the native RNA helicase expressed in a $\Delta crhR$ background (subsequently called CrhR_{WT}). We performed UV cross-linking *in vivo* with CrhR_{WT} cultures grown at the standard growth temperature of 30°C or exposed to 20°C for 2 h (low temperature stress). A possible obstacle in the analysis of RNA helicases can be their transient interaction with RNA molecules followed by rapid ATP-dependent dissociation (Linder and Jankowsky, 2011). Therefore, we introduced a K57A mutation located in the predicted Walker A ATP-binding motif I (GTGKT) that is conserved in CrhR (Tanner and Linder, 2001). Mutation of the conserved lysine or the last threonine in this motif is known to interfere with the ATPase activity of DEAD-box RNA helicases (Cordin *et al.*, 2006). Plasmid encoded K57A was conjugated into the $\Delta crhR$ background yielding strain CrhR_{K57A}. This strain was used for comparison applying UV cross-linking *in vivo* at 30°C followed by co-immunoprecipitation (co-IP) and RNA sequence analysis as for the CrhR_{WT} strain.

Altogether, 119 RNA sequences were significantly enriched in at least one experiment. Functional analysis of the specifically enriched RNAs indicated a striking preference for CrhR interaction with transcripts associated with photosynthesis but also transcripts associated with RNA metabolism, among them *crhR* itself. The enrichment of *crhR*-transcripts with the tagged CrhR protein is consistent with an auto-regulatory mechanism in which CrhR controls its own expression at the post-transcriptional level. Broader implications were revealed by the potential RNA targets having previously been characterized as controlled by the transcription factor RpaB (Riediger *et al.*, 2019), which is of interest in conjunction with the known ETC redox poise regulation of *crhR*

expression (Kujat and Owtrim, 2000; Ritter *et al.*, 2020). Overall, the results connect the known association of CrhR with both the thylakoid membrane and ribosomes and its cold-induced and redox poise-controlled expression with a specific set of potential RNA targets comprising a subset of the *Synechocystis* transcriptome.

Materials and methods

Bacterial strains

The *Synechocystis* $\Delta crhR$ mutant (Prakash *et al.*, 2010) was used for the construction of three different strains. To establish ectopic expression of tagged CrhR, the triple FLAG-tag was introduced at the 5'-end of the wild type *crhR*/*slr0083* gene by PCR in three steps. First, *crhR* was amplified with the primers 3xFL-CrhR-F1 and 3xFL-CrhR-R (see **Table S1** for the sequences of oligonucleotide primers used in this work), re-amplified with primers 3xFL-CrhR-F2 and 3xFL-CrhR-R, followed by another reamplification using the primers 3xFL-CrhR-F3 and 3xFL-CrhR-R. The resulting amplicon 3xFLAG-*crhR* was digested with *NdeI* and *XbaI* and ligated into pJET1.2. The copper inducible promoter, P_{petE} (Zhang *et al.*, 1992) was amplified with the primers petE-FP and petE-RP and cloned into pJET1.2 via Gibson assembly. The *rrnB* terminator was amplified from pBAD using primers *rrnB*-TT-F and *rrnB*-TT-R and ligated into pJET1.2 via *SacI* and *XbaI* restriction sites yielding pJET1.2:: P_{petE} ::3xFLAG-*crhR*::*rrnBT*.

To introduce the K57A mutation into *crhR*, primers *crhR*(K57A)_inverse_fw and *crhR*(K57A)_inverse_rv were used in an inverse PCR to generate CrhR_{K57A}. Both the pJET1.2:: P_{petE} ::3xFLAG-*crhR*::*rrnBT* and CrhR_{K57A} constructs contain the P_{petE} promoter followed by the triple FLAG tag translationally fused to the *crhR* ORF followed by the *rrnB* terminator. To eliminate the *crhR* gene and create the negative control $\Delta crhR_{control}$, the vector pJET1.2:: P_{petE} ::3xFLAG-*crhR*::*rrnBT* was inverse PCR amplified with primers petE-3xF-*rrnB*_neg_ctrl_FP and petE-3xF-*rrnB*_neg_ctrl_RP carrying *SacI* restriction sites on the 5' overhangs, cleaved with *SacI* and self-ligated.

For cloning of the constructs into pVZ321 (Zinchenko *et al.*, 1999), the constructs were amplified from pJET1.2 using the primers petE_foR-pVZ_FP and rrnB_foR-pVZ_RP carrying *EcoRI* restriction sites. The amplified constructs were inserted into the *EcoRI* restriction site of pVZ321. This insertion resulted in disruption of the chloramphenicol resistance cassette of pVZ321, so that the final constructs pVZ321::P_{petE}::3xFLAG-*crhR*::*rrnBT* (strain CrhR_{WT}), pVZ321::P_{petE}::3xFLAG-*crhR*(K57A)::*rrnBT* (strain CrhR_{K57A}) and pVZ321::P_{petE}::3xFLAG::*rrnBT* (strain Δ *crhR*_{control}) possessed only kanamycin resistance. The resulting pVZ321 plasmids were transferred into the Δ *crhR* *Synechocystis* strain (Prakash *et al.*, 2010) via triparental mating with *E. coli* J53/RP4 and TOP10F' (Scholz *et al.*, 2013). Exconjugants were selected on agar plates containing 20 μ g/ml spectinomycin (Sp) and 50 μ g/ml kanamycin (Km) in BG-11 (Rippka *et al.*, 1979).

Culture conditions

E. coli strains were grown in liquid LB medium (10 g l⁻¹ bacto-tryptone, 5 g l⁻¹ bacto-yeast extract, 10 g l⁻¹ NaCl), with continuous agitation or on agar-solidified (1.5% [w/v] bacto agar) LB, supplemented with appropriate antibiotics, at 37°C.

Synechocystis strains were cultivated in the presence of appropriate antibiotics either in Erlenmeyer flasks in BG-11 medium (Rippka *et al.*, 1979) or in 100 ml two-tier vessel CellDEG cultivators (CellDEG GmbH (Bähr *et al.*, 2016)) at the indicated temperatures in fresh water organisms (FWO) medium with shaking. FWO medium consists of 50 mM NaNO₃, 15 mM KNO₃, 2 mM MgSO₄, 0.5 mM CaCl₂, 0.025 mM H₃BO₃, 0.15 mM FeCl₃/Na₂EDTA, 1.6 mM KH₂PO₄, 2.4 mM K₂HPO₄, 10 mM NaHCO₃, 10 nM MnCl₂, 1 nM ZnSO₄, 2 nM Na₂MoO₄, 2 nM CuSO₄, 30 pM CoCl₂. High CO₂ concentrations were delivered to the flasks from a carbonate buffer (1.447 M K₂CO₃ and 12.486 M KHCO₃) through a highly gas-permeable polypropylene membrane. The CellDEG cultivators were inoculated at an OD₇₅₀ ~0.8 and a light intensity of 150 μ mol photons m⁻² s⁻¹. Light intensity was adjusted to culture density; therefore, 24 h after the start of inoculation light intensity was increased to 300 μ mol photons m⁻² s⁻¹ and to 500 μ mol photons m⁻² s⁻¹.

after another 24 h. To induce gene expression from the Cu²⁺-responsive promoter P_{petE} in the CellDEG system, 6 μM CuSO₄ was added to the medium. The cells were harvested at OD₇₅₀ ~20, which was reached upon 72 hours of cultivation.

FLAG-tag affinity purification, UV crosslinking and RNA co-IP

Synechocystis cultures were grown in CellDEG cultivators to an OD₇₅₀ ~20 at 30°C. For cold stress, cultures were exposed to 20°C for 2 h. The cells were placed in a 20 x 20 cm Petri dish and irradiated three times on ice with 500 mJ/cm² UV-C light (254 nm) in a Stratalinker® 2400 for a total irradiation time of 3 min. The treated cells were harvested by centrifugation (4,000 x g, 10 min, 4°C). Cell pellets were resuspended in precooled TBS buffer (50 mM Tris-HCl (pH 7.5)), containing cOmplete™ Protease Inhibitor Cocktail (Roche), washed once with TBS and centrifuged again. The washed and cross-linked cells were suspended in TBS buffer and 0.5 volume of glass beads were added. Cells were disrupted mechanically applying 5 cycles at 6,500 rpm with 10 s breaks on ice between the cycles in a Precellys® 24 homogenizer. The cell lysate was separated by centrifugation (13,000 x g, 15 min, 4°C) into two fractions, one containing membrane proteins and the other containing soluble proteins. The latter was incubated with 50 μl of Anti-FLAG® M2 magnetic beads (Sigma Aldrich) to allow binding of FLAG-tagged proteins at 4°C for 4 hours. Subsequently, beads were washed twice in TBS-containing cOmplete™ Protease Inhibitor Cocktail followed by a wash step that removed the non-crosslinked RNA by incubation in 500 μl RNA elution buffer (25 mM Tris-HCl (pH7.5), 2 M NaCl) for 15 min at room temperature. The crosslinked RNA was released by digestion of the FLAG-tagged protein with 20 μg of proteinase K for 10 min at room temperature, purified using the RNA Clean & Concentrator-5 Kit (Zymo Research) and utilized for the generation of cDNA libraries for Illumina sequencing as described below. All co-immunoprecipitation experiments were performed in biological duplicates.

RNA preparation

Synechocystis cell pellets were disrupted by incubation for 15 min at 65 °C in PGTX as described by Pinto *et al.* (2009). Chloroform/isoamyl alcohol (IAA) (24:1) extraction was performed for 10 min at room temperature followed by centrifugation at 3,270 x g for 15 min at room temperature for phase separation. The chloroform/IAA extraction was repeated on the aqueous phase, followed by isopropanol precipitation. Precipitated RNA was pelleted by centrifugation (13,000 x g, 30 min, 4°C), washed with 70% ethanol, air-dried and resuspended in nuclease-free water. RNA concentration was determined on a NanoDrop ND-1000 spectrophotometer. RNA purity and quality was evaluated on a Fragment Analyzer parallel capillary electrophoresis system (Agilent).

Preparation of cDNA libraries

To initiate cDNA library preparation, 0.3-5 µg of total RNA was treated with 4 U of TURBO DNase (Life Technologies) in 2 consecutive incubation steps, each at 37°C for 15 min as indicated by the manufacturer. Next, RNA clean-up and separation into small (17-200 nt) and large (>200 nt) RNA fractions was performed using RNA Clean & Concentrator columns (Zymo Research). The large RNA fraction was fragmented according to the protocol by Pfeifer-Sancar *et al.* (2013). After RNA Clean & Concentrator column purification, both fractions were combined and all subsequent steps performed as previously described (Pfeifer-Sancar *et al.*, 2013) except that the clean-up of RNA 5'-phosphatase (Epicentre)-treated samples was performed by Clean & Concentrator column purification. For RNA adapter ligation, the UGA linker (Supplementary **Table S1**) was used. After RNA adapter ligation and cDNA synthesis, the samples were excised from 2% agarose gels and purified using the NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel) using NTC buffer for solubilization of the gel slices. After PCR amplification, residual primers were removed by adding 10 µl of ExoSAP-IT (USB) to 50 µl of PCR, and with samples incubated for 15 min at 37°C followed by heat inactivation of the enzyme for 15 min at 85°C. The samples were cleaned-up using a NucleoSpin Gel and PCR Clean-up kit. The quality of RNA and DNA

was analyzed on a Fragment Analyzer parallel capillary electrophoresis system (Agilent). Sequencing of libraries was performed on a HiSeq 3000 sequencer with a 2 x 150 bp read mode.

Bioinformatics analyses

Analysis of the raw reads from the RNA sequencing was performed using the Galaxy web platform (<https://usegalaxy.eu/> (Afgan *et al.*, 2018)). Illumina paired-end reads were trimmed, adapter sequences and reads shorter than 14 nt filtered out using cutadapt 1.16 (Martin, 2011). The remaining reads were mapped to the chromosome and plasmids of *Synechocystis* using bowtie2 version 2.3.4.3 with the following selected parameters for paired-end reads: -l 0 -X 500 --fr --no-mixed --no-discordant --very-sensitive (Langmead and Salzberg, 2012). Peak calling of the mapped reads was performed by PEAKachu 0.1.0.2 with the parameters: --pairwise_replicates --norm_method deseq --mad_multiplier 2.0 --fc_cutoff 1 --padj_threshold 0.05. The total number of reads as well as those that mapped to the *Synechocystis* genome are listed in **Supplementary Material, Table S1**.

RNA secondary structures were predicted using the RNAfold WebServer as part of the ViennaRNA Websuite (Gruber *et al.*, 2015) with default parameters. Structures were visualized for publication in the VARNA Applet version 3.93 (Darty *et al.*, 2009).

Generation of recombinant CrhR

To obtain recombinant, C-terminal His-tagged CrhR, the *crhR* gene was amplified from the constructs pVZ321::P_{petE}::3xFLAG-*crhR*::*rrnBT* with primers pQE70_*crhR*-fw and pQE70_*crhR*-rv (**Table S1**). The pQE70 vector was amplified with primers pQE_GIBSOOn-fw and pQE_GIBSOOn-rv. The fragments were mixed and transformed into *E. coli* Top10 F' using AQUA cloning (Beyer *et al.*, 2015) yielding strain pQE70:*crhR*-6xHis. For recombinant protein expression, the construct pQE70::*crhR*-6xHis was transformed into *E. coli* M15. Overnight cultures were diluted 1:100 in fresh LB medium

supplemented with 50 µg/ml ampicillin, and grown to an OD₆₀₀ of 0.7 at 37°C. Protein expression was induced by adding IPTG to 1 mM final concentration. Three hours after induction, cells were harvested by centrifugation at 6,000 x g for 10 min at room temperature. Cell pellets were resuspended in lysis buffer (50 mM NaH₂PO₄ (pH 8), 1 M NaCl, 10% glycerol, 15 mM imidazole, cOmplete™ Protease Inhibitor Cocktail (Roche)) and lysed using the One Shot constant cell disruption system (Constant Systems Limited, U.K.) at 2.4 kbar. Cell lysates were cleared by centrifugation at 13,000 g for 30 min at 4°C and the lysate was filtered through 0.45 µm Supor-450 filters (Pall). Recombinant proteins were immobilized on a HiTrap Talon crude 1 ml column (GE Healthcare), equilibrated with buffer A (50 mM NaH₂PO₄ (pH 8), 500 mM NaCl), and eluted with elution buffer B (50 mM NaH₂PO₄ (pH 8), 500 mM imidazole, 500 mM NaCl). The protein concentration was calculated with a Bradford assay using bovine serum albumin as the standard.

Cy3 labelling of RNA

To initiate Cy3 RNA labelling for EMSA analysis, two volumes of 6 mM KIO₄ were added to 4 µg RNA and incubated at room temperature in the dark for 1 h. The RNA was precipitated at -20°C for 1 h by addition of one volume of ethylene glycol/H₂O (1:1), 2.9 volumes of 100% ethanol, and 0.1 volume of 3.3 M NaCl and pelleted at 13,000 x g for 30 min at 4°C. The RNA pellet was washed with 70 % ethanol, air dried, and resuspended in 10 µl of 50 mM Cy3 dye dissolved in DMSO (Thermo Fisher Scientific) and incubated at 37°C in the dark for 2 h. Two volumes of 0.1 M Tris-HCl (pH 7.5) were added followed by the addition of 20 mM freshly prepared NaBH₄ and the reaction mix incubated at 4°C in the dark for 30 min. The reaction was quenched and RNA precipitated by ethanol precipitation.

Electrophoretic mobility shift assays

Binding of different amounts of recombinant CrhR (0.1 to 5 pmol) to 0.2 pmol of Cy3 labelled RNA was performed in 20 mM HEPES-KOH (pH 8.3) buffer containing 3 mM

MgCl₂, 1 mM DTT, 500 µg/ml BSA. As a substrate competitor, 1 µg of LightShift Poly (dIdC) (Thermo Fischer Scientific) was included in each assay. The reaction was incubated at room temperature for 15 min prior to separation on 2% agarose-TAE gels. The signals were visualized with a Laser Scanner Typhoon FLA 9500 (GE Healthcare) using a green light laser at a wavelength of 532 nm and a Cy3 filter (LPG, DGR1, BPG1.

Results

Physiological consequences of manipulation of CrhR sequence and abundance

To gain insight into the CrhR interactome, we performed immunoprecipitation and isolation of tagged CrhR-RNA complexes from *in vivo* UV cross-linked *Synechocystis* cells. A previously characterized helicase deletion mutant, $\Delta crhR$, generated by replacement of the *crhR* gene (*slr0083*) by a spectinomycin resistance gene (Prakash *et al.*, 2010), served as the genetic background in which three strains were constructed. One, ectopically expressing the triple FLAG-tagged CrhR (CrhR_{WT}), and the second expressing the triple FLAG-tagged CrhR(K57A) mutant (CrhR_{K57A}) from the conjugative vector pVZ321 under control of the copper inducible promoter, P_{petE} (Zhang *et al.*, 1992) (**Fig. 1A**). The introduced K57A mutation should interfere with ATP binding and ATPase activity of CrhR (Tanner and Linder, 2001) and therefore delay or reduce its capability to dissociate from bound RNA. This mutation is predicted to stabilize CrhR-target transcript interaction, thereby aiding our ability to identify especially low-abundant transcripts. As a control, a third strain was constructed with pVZ321 carrying a short reading frame encoding a triple FLAG tag and identical promoter and terminator sequences as the other two strains ($\Delta crhR$ _{control}). Three experiments were performed with these strains (**Fig. 1B**). The proper regulation and expression of FLAG-tagged CrhR proteins was confirmed by western blotting (**Fig. S1**). We noticed that under identical conditions the expression level of CrhR_{K57A} was higher than that of CrhR_{WT}, although (with the exception of the single substitution) both constructs including the promoter, 5' and 3' sequences, were identical. Therefore, the protein carrying the K57A substitution appears to be stabilized in the absence of functional CrhR.

We followed the growth of the three generated strains and wild type over 25 days at 20°C. Growth of $\Delta crhR_{\text{control}}$ was dramatically reduced compared to wild type cells, while growth of $CrhR_{\text{WT}}$ was similar to the unmodified wild type (**Fig. 2A**). Thus, ectopically transcribed, FLAG-tagged CrhR in the $CrhR_{\text{WT}}$ strain complemented the *crhR* deletion at this temperature. Interestingly, complementation in strain $CrhR_{\text{K57A}}$ led to a partial restoration of the cold-sensitive phenotype, i.e. growth of $CrhR_{\text{K57A}}$ was reduced compared to the wild type but not to the same extent as of $\Delta crhR_{\text{control}}$ (**Fig. 2A**). In addition, a strong reduction in pigment content in $CrhR_{\text{K57A}}$ and in $\Delta crhR_{\text{control}}$ was apparent compared to the other two strains (**Fig. 2B, C**). These variations were congruent with previously identified changes in light-harvesting pigment composition upon *crhR* inactivation (Rosana *et al.*, 2012b). We concluded that the FLAG tag did not interfere with the enzymatic and regulatory functions of CrhR and therefore these lines were suitable for more detailed analyses.

RNA co-immunoprecipitations

The RNA helicase CrhR performs unwinding and annealing of RNA (Chamot *et al.*, 2005), reactions for which mainly transient interactions with RNA can be assumed including binding to RNA followed by rapid dissociation. This makes the experimental mapping of an RNA helicase interactome more challenging than for an RNA-binding protein. Therefore, we performed covalent crosslinking *in vivo* to enrich RNA-CrhR complexes sufficient for RNA and library preparation. $CrhR_{\text{WT}}$ and $CrhR_{\text{K57A}}$ cultures grown at the standard growth temperature of 30°C were irradiated with 254 nm UV-C light at 500 mJ/cm² (Holmqvist *et al.*, 2016). To identify potentially different interacting partners, the experiment was also performed on $CrhR_{\text{WT}}$ cultures exposed to cold stress at 20°C for 2 hours. In parallel, total transcriptome data were generated for all strains and conditions (see overview in **Fig. 1B**). The total and mapped read numbers are listed in **Supplementary Material, Table S2**. CrhR interacting RNAs were predicted with the peak calling algorithm PEAKachu (Holmqvist *et al.*, 2016). In total, we obtained 98 different peaks with statistically significant enrichment (*p*_{adj} < 0.05, log₂FC > 0) in the three experiments corresponding to 64 different genes (**Supplementary Material,**

Tables S3 to S5). CrhR interacting RNAs were most abundant in experiment 3 using CrhR_{K57A} as bait (75 peaks), while 28 and 16 peaks were identified in the experiments 1 (20°C) and 2 (30°C) using CrhR_{WT}. The higher number of peaks using CrhR_{K57A} as bait matches the assumption that the K57A substitution would stabilize CrhR-RNA interactions. In addition, the somewhat higher expression level of CrhR_{K57A} compared to CrhR_{WT} (**Fig. S1**) may have contributed as well.

Interestingly, the co-IPs frequently identified only specific regions of RNA transcripts which could provide insight into both the molecular mechanism and also the functional roles CrhR is performing. Typical patterns of enriched transcripts are visualized in **Fig. 3**. The predominant detected interactions were associated with segments containing the 5'-untranslated region (5' UTR), start codon and a few codons into the reading frame (**Fig. 3A**). A related pattern also included the 5' UTR, start codon and initial codons but actually matched the entire intergenic space between two genes in an operon (**Fig. 3B**). A strikingly different pattern was observed in which multiple sharp peaks indicated a multitude of interactions along the length of the dicistronic operon, as illustrated for the *psaA-psaB* operon (**Fig. 3C**). Nevertheless, as observed for other target transcripts, the most pronounced interactions corresponded to the 5' UTR, start and first codons of *psaA*, the first gene of this operon (**Fig. 3C**). We noticed that the read distribution in the enriched RNA segments resembled somewhat the total RNA-seq read distribution. The reason likely is that the CrhR-mRNA interaction lowers the accessibility for endoribonucleases and thereby protects the respective transcript segments from degradation *in vivo*.

Transcripts encoding photosynthesis-associated proteins or with predicted roles in redox regulation are enhanced in the CrhR interactome

Functional analysis over the complete dataset using GO terms (Ashburner *et al.*, 2000; Gene Ontology Consortium, 2021) identified “photosynthetic membrane” ($q_{\text{adj}} = 1.18\text{E-}04$), “thylakoid” ($q_{\text{adj}} = 2.74\text{E-}04$) and “photosystem” ($q_{\text{adj}} = 2.74\text{E-}04$) as the dominating cellular components encoded by the enriched transcripts.

Accordingly, detailed analysis indicated that most of the enriched transcripts were associated with photosynthesis (**Table 1**; for the exact coordinates of recovered transcripts, enrichment factors and adjusted p values in each of the experiments, see **Supplementary Material, Tables S3 to S5**). Several of these transcripts encode protein components of the two photosystems, PSI and PSII. PSI proteins included *psaL/slr1655* for PSI subunit XI, *slr1834/psaA* and *slr1835/psaB* encoding the P700 apoprotein subunits Ia and A2 (**Fig. 3C**), *ssr2831/psaE* encoding the PSI reaction center subunit IV. Transcripts encoding PSII subunits included *slr0906/psbB* for the CP47 reaction center protein, *slr0927/psbD*, *slr1311/psbA2* and *smr0009* for the reaction center proteins D2, D1 and PsbN.

Several enriched transcripts encoding proteins that are not directly components of PSI or PSII but that are associated with photosynthesis were also detected. Transcripts in this category included *slr0228/ftsH2* encoding the FtsH2 protease crucial for the replacement of photodamaged D1 protein during the PSII repair cycle (Silva *et al.*, 2003; Komenda *et al.*, 2010), *ssl0020/petF* encoding the major plant-like ferredoxin Fed1 (Mazouni *et al.*, 2003), *slr0199/petE* encoding plastocyanin and *ssr2016* encoding the *Synechocystis* Pgr5 (proton gradient regulation 5) homolog. In plants and cyanobacteria, Pgr5 contributes to cyclic electron flow from PSI to the plastoquinone pool (Yeremenko *et al.*, 2005; Dann and Leister, 2019), electron transfer that is associated with *crhR* expression (Kujat and Owttrim, 2000; Ritter *et al.*, 2020). Another gene in this category was *trxA* (*slr0623*), encoding thioredoxin A, the most abundant of the four thioredoxins in *Synechocystis* (Hishiya *et al.*, 2008), and suggests CrhR is involved in maintaining intracellular redox status by regulating protein expression associated with scavenging reactive oxygen species (ROS).

Light harvesting is an integral aspect of photosynthesis, the primary contributor to which are phycobilisomes in cyanobacteria. Among several mRNAs encoding phycobilisome (PBS) proteins in the dataset (**Table 1**) was *apcC/ssr3383* encoding the PBS 7.8 kDa linker polypeptide. The interacting RNA sequence encompassed 146 nt of the 5' UTR and 54 of the 67 *apcC* codons (**Fig. 4**), importantly ending at the start of the region from which the regulatory sRNA ApcZ originates. ApcZ targets the *ocp/slr1963* transcript

encoding the water-soluble orange carotenoid protein OCP (Zhan *et al.*, 2021). Again, interaction of this transcript with CrhR is associated with redox homeostasis since OCP functions to quench excess excitation energy absorbed by phycobilisomes, and also directly scavenges singlet oxygen ROS ($^1\text{O}_2$) (Kerfeld *et al.*, 2003). Downstream of the photosynthetic electron transport chain (ETC), CrhR interaction with the *slr0148* transcript was detected (**Table 1**). The gene *slr0148* encodes the ferredoxin *fed5* (Angeleri *et al.*, 2018), homologs of which are widely distributed in cyanobacteria and plants (Cassier-Chauvat and Chauvat, 2014). The Fed2 to Fed9 gene family functions to distribute electrons to a range of metabolic pathways each of which are associated with cyanobacterial tolerance to a different but overlapping range of environmental stresses (Cassier-Chauvat and Chauvat, 2014). In addition, *slr0148* is the terminal gene of 5 consecutive genes *slr0144* to *slr0148*, all of whose mRNA fragments were enriched by co-IP with CrhR_{K57A} (**Table S5**). These five genes are also jointly down-regulated after cold shift, together with the two following genes *slr0149* to *slr0151* (Georg *et al.*, 2019).

In addition, we identified CrhR-interacting transcripts of genes more distantly related to photosynthesis, among them *slr0374* (**Fig. 3C**), encoding the stress-responsive AAA+ protease protein Ycf46 (Singh and Sherman, 2002). Homologs of Ycf46 are highly conserved in all cyanobacterial lineages and most algal chloroplast genomes. Ycf46 proteins are involved in the regulation of CO₂ utilization in photosynthesis but the precise function is unknown (Jiang *et al.*, 2015). Transcript peaks were also recovered for *slr1674*, encoding a DUF760-containing protein described to be associated with the thermal acclimation of PSII (Rowland *et al.*, 2010).

Identification of photosynthesis-associated transcripts is consistent with the published photosynthetic acclimation and loss-of-function phenotypes observed in *crhR* deletion and inactivation mutants upon temperature downshift (Rosana *et al.*, 2012b; Sireesha *et al.*, 2012). One should note that most of these transcripts are abundant and might be easier detected in our co-IP approach than less abundant transcripts. However, not all identified transcripts are abundant, e.g. the transcripts for genes such as *slr0442* encoding a 'target gene of Sycrp1' or *slr0587* belong to the set of less abundant transcriptional units according to the RNA-Seq analyses by Kopf *et al.* (2014).

Nevertheless, while we detected some low abundance transcripts, the nature of the experiments and rigorous exclusion criteria employed would naturally have resulted in the exclusion of some lower expressed transcripts.

CrhR interactome comparison with omic data

Here, we recovered in the three experiments 119 peaks belonging to 90 different genes. Previously, differential gene expression due to downshifts in temperature and the molecular effects of *crhR* deletion or inactivation have been studied at transcriptome (Prakash *et al.*, 2010; Georg *et al.*, 2019) and proteome level (Rowland *et al.*, 2011), which generated differing regulons. Indeed 68/119 recovered peaks belong to genes for which significant differential regulation due to temperature downshifts were described at RNA level, including several transcripts encoding PSI proteins. The here identified mRNAs *psaAB*, *psaD*, *psaE* and *psaL* belong to some of the most strongly down-regulated genes at the lower temperature (see **Tables S3 to S5** for details).

Previous 2D gel electrophoresis of soluble proteins identified 16 proteins differentially expressed in the $\Delta crhR$ mutant at 34°C and 25 proteins at 24°C (Rowland *et al.*, 2011). We found that 15 of the 90 different genes for which mRNAs were enriched in the CrhR co-IP had been identified in the analysis by Rowland *et al.* (2011) as differently expressed at protein level. This includes several proteins of the light-harvesting system including our detection of *apcA*, *B* and *E* transcripts coding for allophycocyanin subunits, *cpcA* and *B* encoding phycocyanin subunits and *apcC*, *cpcC1* and *cpcG1* encoding the linker polypeptides ApcC, CpcC1 and CpcG1 as well as the cold-inducible OCP.

Another category of proteins identified in the analysis by Rowland *et al.* (2011) are those involved in protein production and gene expression. Corresponding to the proteome described by Rowland *et al.* (2011), we identified the CrhR-interacting transcripts encoding GroEL-1 and GroEL-2 chaperones, ribosomal proteins Rps1a, Rps4, Rps14, Rpl21, Rpl28 and Slr0923 and the hypothetical protein Slr0552 (transcribed in a dicistron with *slr0551* encoding RNase J). Finally, the additional strong upregulation of the cold-inducible *slr0082* encoding methylthiotransferase RimO in $\Delta crhR$ was described

(Rowland *et al.*, 2011). While we did not detect the *slr0082* mRNA in our CrhR co-IPs, its upregulation can be explained via an operon discoordination mechanism that depends on CrhR (see below).

Transcripts encoding proteins involved in housekeeping functions and RNA metabolism as potential CrhR targets

Another category of transcripts enriched in the CrhR co-IPs were related to housekeeping functions of the cell. Examples included *groEL1/slr2076* and *groEL2/sll0416* encoding the 60 kDa chaperonins 1 and 2 that match previous reports on the CrhR-dependent upregulation of *groEL1* and *groEL2* in *Synechocystis* (Prakash *et al.*, 2010). Additionally, transcripts encoding the ribosomal proteins, *rplU/slr1678* and *rpl26/ssr1604* for the 50S ribosomal proteins L21 and L28, and *rps14/slr0628* and *rps1a/slr1356* for the 30S ribosomal proteins S14 and S1a, belonging to four separate operons, were detected. Of specific interest in relation to *crhR* expression, a transcript enriched specifically from cold stressed cultures was *desC/sll0541* encoding the $\Delta 9$ fatty acid desaturase DesC was identified as a potential CrhR target.

A regulatory gene of interest is *sigG/slr1545* encoding the Type 3 alternative sigma factor SigG (Huckauf *et al.*, 2000), especially because it was recovered in all three experiments suggesting a stable interaction with CrhR (**Fig. 3A**). SigG is associated with the oxidative stress response caused by high light (Huckauf *et al.*, 2000) and is ectopically upregulated in a $\Delta rpaA$ mutant in the dark (Köbler *et al.*, 2018). Furthermore, interacting transcripts encoding enzymes associated with RNA turnover and metabolism were identified, including *rnj/slr0551* encoding ribonuclease J, an enzyme possessing both endonuclease and a robust 5'-exoribonuclease activity (Cavauiolo *et al.*, 2020), *slr0053* encoding the rRNA maturation RNase YbeY and, as an enzyme of RNA metabolism, *crhR* itself.

The *crhR* mRNA is a target for auto-regulation through CrhR

The *crhR* transcript was consistently enriched in all three *crhR* lines irrespective of temperature (**Fig. 5**). The enriched peak regions within the *crhR* transcript identify three potential regions of preferred interaction: the major sequence overlaps the translational start codon and extends 216 to 270 nt into the coding region, a second region from approximately codon 125 to 219 and a third extending from approximately codon 350 to 450 of the 492 codon ORF.

To confirm CrhR-*crhR* mRNA interaction *in vitro*, 150 nt of the *crhR* transcript, corresponding to the primary interacting region, was tested in an electrophoretic mobility shift assay (EMSA) using His-tagged recombinant CrhR produced in *E. coli* (**Fig. 5B**). Enhanced formation of a prominent retarded protein-RNA complex was directly proportional to CrhR concentration. We conclude that the *crhR* mRNA is a preferred target of CrhR and thus an auto-regulatory effect appears likely.

Discussion

Involvement of CrhR in a CrhR-dependent auto-regulatory negative feedback loop

The DEAD-box RNA helicase CrhR is one of the major cold shock proteins of *Synechocystis* (Kujat and Owtrim, 2000; Prakash *et al.*, 2010; Rowland *et al.*, 2011; Rosana *et al.*, 2012b,a; Sireesha *et al.*, 2012). Consistent with observations in many other organisms that shift to lower temperature increases expression of DEAD-box RNA helicases (Owtrim, 2013), we observed previously an increased *crhR* transcript abundance after temperature downshift to 20°C, with a log₂FC of 1.4 in the wild type and 2.6 for the 5' portion of the gene not deleted in a partial *crhR* inactivation mutant (Georg *et al.*, 2019). The higher induction in the partial *crhR* inactivation mutant pointed at a possible feedback effect of CrhR that was active in the wild type but absent in the mutant.

Regulation of *crhR* expression is complex, involving transcriptional and post-transcriptional effects (Rosana *et al.*, 2012a), and an auto-regulated operon

discoordination and processing mechanism was proposed (Rosana *et al.*, 2020). Complicating the regulation, *crhR* is encoded in the *rimO/slr0082-crhR/slr0083* (*rimO-crhR*) dicistronic operon (Rosana *et al.*, 2012a). However, in wild type *Synechocystis* cells, accumulation of the full-length dicistronic transcript is essentially undetectable; *rimO* and *crhR* mRNAs co-occur predominantly as monocistronic transcripts (Rosana *et al.*, 2012a, 2020). Interestingly, in the absence of functional CrhR, accumulation of the entire *rimO-crhR* operon transcript is elevated compared to wild type cells (Rosana *et al.*, 2020). This observation led to the assumption that CrhR may facilitate processing of the *rimO-crhR* operon transcript, although CrhR is not necessary and sufficient for the processing activity (Rosana *et al.*, 2020). The discovery of an RNase E processing site ~138 nt upstream of the *crhR* coding sequence then suggested that RNase E might play a role in processing of the operon. We previously showed that the *rimO-crhR* dicistronic transcript was cleaved by RNase E ~138 nt upstream of the *crhR* start codon *in vitro* (Rosana *et al.*, 2020). Interestingly, both specific RNA sequence and structure were required for RNase E substrate recognition and cleavage. Thus, we propose that CrhR may assist RNase E by interacting with the *rimO-crhR* transcript and altering its structure to the one that favors cleavage by RNase E (**Fig. 6**). In this scenario, CrhR would auto-regulate its own expression by enhancing processing of the *rimO-crhR* operon by RNase E. Moreover, CrhR could also be involved in the further processing of monocistronic *crhR* (Rosana *et al.*, 2012a, 2020). In support of this hypothesis, northern and microarray analysis previously indicated that the absence of functional CrhR RNA helicase activity enhances stabilization of *crhR* transcripts at 30 and 20°C (Georg *et al.*, 2019; Rosana *et al.*, 2020). From the degradation constants, we concluded that at 20°C, alteration of RNA levels in the *crhR* helicase deficient mutant could be largely explained by regulation occurring at post-transcriptional level (Georg *et al.*, 2019).

In combination, these observations infer a CrhR-dependent negative feedback loop, in which binding of CrhR to *crhR* transcripts leads to auto-regulation of expression.

The CrhR interactome *in vivo*

DEAD-box RNA helicases have been characterized as drivers and regulators of gene expression. For many DEAD-box RNA helicases, both the preferred targets as well as the mechanism of target recognition have remained unknown. This is relevant as the diversity of functions of DEAD-box RNA helicases and sequences of the auxiliary domains suggests that there is no single mechanism of specific target recognition or pathway association. Determination of the RNA helicase interactome is more challenging than for other RNA-interacting proteins since after binding, the helicase will rapidly modify the structure and dissociate. This potentially impedes the ability to co-IP interacting transcripts compared to RNA-binding proteins such as CsrA, Hfq or ProQ (Hör *et al.*, 2018). To mitigate this challenge, we employed the CrhR variant, CrhR_{K57A}, in which RNA residence time was predicted to be extended, since the Walker A ATP binding domain is inactivated.

This consideration was supported by the enhanced recovery of interacting transcripts from the CrhR_{K57A} strain, i.e. experiment 3 (**Fig. 1**) suggesting that RNA binding was most efficient in the CrhR_{K57A} strain. Moreover, the partial complementation of the $\Delta crhR$ phenotype by CrhR_{K57A} (**Fig. 2**) supported the view that this variant had reduced RNA helicase enzymatic activity. The overlap in the enriched targets identified in the three experiments consists of transcripts belonging to only four genes (*crhR*, *psaA*, *sigG* and *ycf46*) as shown in the Venn diagram in **Fig. S2**.

Functionally, the prime targets of CrhR interaction were consistently observed to include transcripts associated with photosynthesis, light harvesting or directly encoding core photosystem proteins (**Table 1**). These findings are consistent with observations that the rapid cessation of photosynthesis upon temperature downshift from 30 to 20°C is the major physiological consequence created by the absence of CrhR RNA helicase activity (Rosana *et al.*, 2012b). A vital aspect relevant to photosynthesis is the reported inability of the $\Delta crhR$ mutant to properly perform light-induced state transitions, which occur as part of the normal acclimation response to downshifts in temperature in the wild type (Sireesha *et al.*, 2012). Our data provide deeper insight into the molecular basis of these observations. In contrast to the monomeric PSI structures in plants and algae, PSI in

many cyanobacteria is mainly trimeric, in some strains tetrameric (Li *et al.*, 2019). The trimeric structure enlarges the antenna system under low-light conditions, while the dynamic regulation of the PSI trimer-to-monomer ratio impacts the speed of state transitions (Aspinwall *et al.*, 2004). The PSI reaction center protein subunit XI encoded by *psaL* is crucial for the multimerization status of PSI (Aspinwall *et al.*, 2004; Li *et al.*, 2019). Here, we observed a pronounced interaction between CrhR and a segment of the *psaL* mRNA that contains the 5' UTR and the first 30 codons (**Fig. S3**). This region furthermore contains the interaction site for the photosynthesis regulatory RNA1, PsrR1 (Georg *et al.*, 2014). Previous analyses demonstrated that the proportion of PSI trimers is reduced in $\Delta crhR$ relative to wild type upon temperature downshift (Sireesha *et al.*, 2012). Therefore, the interaction between the cold-induced CrhR and this region is predicted to be necessary for the required expression of *psaL*, likely by facilitating translation, as diagramed in **Fig. S3**.

A role of CrhR in the biology of photosynthesis-relevant mRNAs is further consistent with the demonstrated localization of ribosome-associated CrhR at the thylakoid membrane (Rosana *et al.*, 2016) and the observation that *psbA* and *psaA* mRNAs are transported to the thylakoid membrane and possibly translated by thylakoid-associated ribosomes (Tyystjärvi *et al.*, 2001; Mahbub *et al.*, 2020). These findings point to a possible involvement of CrhR in physiological secondary structure rearrangements of the identified mRNAs that could influence translation on thylakoid membrane associated ribosomes.

Since *Synechocystis* encodes only a single DEAD-box RNA helicase, it is likely that CrhR operates in multiple pathways. These roles could be associated with different stresses, independent of temperature, as observed for the *Staphylococcus aureus* helicase, CsdA (Oun *et al.*, 2013; Khemici *et al.*, 2020). Transcripts recovered for ribosomal proteins and ribonucleases support this idea. Indeed, a role in translation regulation is supported by the observation that CrhR interacts with the *rps1a/slr1356* transcript encoding Rps1a. In previous analyses, Rps1a sedimented away from the majority of ribosomal proteins (Riediger *et al.*, 2021) and was reported to perform a role in the Shine-Dalgarno-independent initiation of translation (Mutsuda and Sugiura, 2006;

Nakagawa *et al.*, 2010), another process which may hence be related to CrhR helicase activity.

Another intriguing example for the possible involvement of CrhR in functionally relevant secondary structure re-arrangements was observed for *apcC/ssr3383* where the recovered reads terminated in the region from which the regulatory sRNA ApcZ originates. ApcZ is an sRNA, which targets the *ocp/slr1963* mRNA encoding the water-soluble OCP by inhibiting its translation (Zhan *et al.*, 2021). If expressed, OCP directly senses light intensity and induces thermal energy dissipation under stress conditions (Muzzopappa and Kirilovsky, 2020). Therefore, its expression is tightly controlled and, in this context, ApcZ can inhibit or delay the production of OCP. This regulation is also relevant under nitrogen starvation conditions, when ApcZ is induced from a transcription start site (TSS) that is activated by the transcription factor NtcA (Zhan *et al.*, 2021). However, ApcZ is also detectable as a transcript under other conditions where it derives from processing of the *apcABC* mRNA. Interestingly, the co-IP coverage included the first 30 nt of ApcZ, matching precisely the upstream portion of the helical structure located at the 5' end of ApcZ (**Fig. 4**). Therefore, it is tempting to speculate that CrhR interaction and helicase activity is either required for i) ApcZ processing, ii) to open the ApcZ structure for interaction, or possibly iii) to facilitate ApcZ-*ocp* duplex formation. These functions are supported by the observation that the *ocp* mRNA was identified as a potential CrhR target in our co-IP experiment (**Table 1**), that it accumulates at a higher level upon temperature downshift (Georg *et al.*, 2019) and by the ability of CrhR to catalyze both RNA duplex unwinding and annealing (Chamot *et al.*, 2005). In conjunction, the OCP protein level was significantly increased at the lower temperature (Rowland *et al.*, 2011).

Overlaps in the regulon controlled by RpaB and the set of transcripts interacting with CrhR

Several of the genes whose transcripts interacted with CrhR were previously predicted or demonstrated to play a role in redox signaling or are regulated in a redox-dependent

manner. This applies first and foremost to *crhR* itself (Kujat and Owttrim, 2000; Ritter *et al.*, 2020). Another is thioredoxin A, the homolog of plant m-type thioredoxin and therefore sometimes also called TrxM, which plays a major role in redox regulation in *Synechocystis* (Hishiya *et al.*, 2008). A protein interacting with TrxM *in vitro* is the redox-responsive transcription factor RpaB (Kadowaki *et al.*, 2015). RpaB recognizes the “high light regulatory 1” (HLR1) promoter element, a pair of imperfect 8-nt long direct repeats (G/T)TTACA(T/A) (T/A) separated by two random nucleotides (Eriksson *et al.*, 2000; Kappell and van Waasbergen, 2007). Binding of RpaB to HLR1 promoter motifs yields one of two different outcomes, repression or activation under low light depending on the distance between the location of these elements and the TSS (reviewed by Riediger *et al.*, 2018). RpaB is the response regulator (RR) of the histidine kinase Hik33 in a two-component signal transduction system that controls >150 target promoters in *Synechocystis*, regulating expression of diverse genes associated with photosynthesis (Riediger *et al.*, 2019). Thus, there is a striking correspondence that approximately two thirds of the transcripts recovered in our CrhR-co-IP analysis belong to the RpaB regulon (**Table 1**). Identification of RpaB regulon members as CrhR-interacting transcripts therefore suggests a scenario in which CrhR contributes to the post-transcriptional regulation of RpaB regulon expression. Together, these observations support the function of CrhR as an important post-transcriptional regulatory element within the network of redox-dependent signaling and regulation.

Acknowledgements

We thank the Freiburg Galaxy Team for maintaining this great bioinformatics resource.

Author contributions

WRH designed the study, CS established the cDNA library preparation and supported the bioinformatic analysis; JF, BH and RR sequenced cDNA libraries; AM, FH, RB and WRH analyzed data; AS and JSSP constructed the *Synechocystis* strains. All other

experiments were carried out by AM. AM, GWO and WRH wrote the paper with contributions from all authors.

Funding

This work was supported by grants from the Baden-Wuerttemberg Foundation BWST_NCRNA_008, by the German Research Foundation (DFG) - 322977937/GRK2344 and by the Federal Ministry of Education and Research (BMBF) program RNAProNet, grant 031L0164B to WRH and RB, by the EU ITN “Photo.COMM” to support the stay of AS in the lab of WRH, a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant, grant 171319 to GWO and a Department of Biotechnology (DBT) of India grant BT/PR13616/BRB/10/774/2010 to JSSP.

Data availability

The RNA-seq data and raw sequence data from co-IP analyses have been deposited in the SRA database <https://www.ncbi.nlm.nih.gov/sra/> and are openly available under the accession numbers SAMN14615974 to SAMN14615989 and SAMN14651279 to SAMN14651286.

References

- Afgan E, Baker D, Batut B, et al.** 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Research* **46**, W537–W544.
- Angeleri M, Zorina A, Aro E-M, Battchikova N.** 2018. Interplay of SpkG kinase and the Slr0151 protein in the phosphorylation of ferredoxin 5 in *Synechocystis* sp. strain PCC 6803. *FEBS Letters* **592**, 411–421.
- Ashburner M, Ball CA, Blake JA, et al.** 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nature Genetics* **25**, 25–29.
- Aspinwall CL, Sarcina M, Mullineaux CW.** 2004. Phycobilisome mobility in the cyanobacterium *Synechococcus* sp. PCC7942 is influenced by the trimerisation of photosystem I. *Photosynthesis Research* **79**, 179.
- Bähr L, Wüstenberg A, Ehwald R.** 2016. Two-tier vessel for photoautotrophic high-density cultures. *Journal of Applied Phycology* **28**, 783–793.
- Beyer HM, Gonschorek P, Samodelov SL, Meier M, Weber W, Zurbriggen MD.** 2015. AQUA cloning: A versatile and simple enzyme-free cloning approach. *PLOS ONE* **10**, e0137652.
- Boudet N, Aubourg S, Toffano-Nioche C, Kreis M, Lecharny A.** 2001. Evolution of intron/exon structure of DEAD helicase family genes in *Arabidopsis*, *Caenorhabditis*, and *Drosophila*. *Genome Research* **11**, 2101–2114.
- Bourgeois CF, Mortreux F, Auboeuf D.** 2016. The multiple functions of RNA helicases as drivers and regulators of gene expression. *Nature Reviews Molecular Cell Biology* **17**, 426–438.
- Capel C, Albaladejo I, Egea I, et al.** 2020. The res (restored cell structure by salinity) tomato mutant reveals the role of the DEAD-box RNA helicase SIDEAD39 in plant development and salt response. *Plant, Cell & Environment* **43**, 1722–1739.
- Cassier-Chauvat C, Chauvat F.** 2014. Function and regulation of ferredoxins in the cyanobacterium, *Synechocystis* PCC6803: recent advances. *Life* **4**, 666–680.
- Cavaiuolo M, Chagneau C, Laalami S, Putzer H.** 2020. Impact of RNase E and RNase J on global mRNA metabolism in the cyanobacterium *Synechocystis* PCC6803. *Frontiers in Microbiology* **11**, 1055.
- Chamot D, Colvin KR, Kujat-Choy SL, Owtrim GW.** 2005. RNA structural rearrangement via unwinding and annealing by the cyanobacterial RNA helicase, CrhR. *The Journal of Biological Chemistry* **280**, 2036–2044.

803 **Chamot D, Magee WC, Yu E, Owttrim GW.** 1999. A cold shock-induced cyanobacterial
804 RNA helicase. *Journal of Bacteriology* **181**, 1728–1732.

805 **Cordin O, Banroques J, Tanner NK, Linder P.** 2006. The DEAD-box protein family of
806 RNA helicases. *Gene* **367**, 17–37.

807 **de la Cruz J, Kressler D, Linder P.** 1999. Unwinding RNA in *Saccharomyces*
808 *cerevisiae*: DEAD-box proteins and related families. *Trends in Biochemical Sciences* **24**,
809 192–198.

810 **Dann M, Leister D.** 2019. Evidence that cyanobacterial Sll1217 functions analogously
811 to PGRL1 in enhancing PGR5-dependent cyclic electron flow. *Nature Communications*
812 **10**, 5299.

813 **Darty K, Denise A, Ponty Y.** 2009. VARNA: Interactive drawing and editing of the RNA
814 secondary structure. *Bioinformatics* **25**, 1974–1975.

815 **Eriksson J, Salih GF, Ghebremedhin H, Jansson C.** 2000. Deletion mutagenesis of
816 the 5' *psbA2* region in *Synechocystis* 6803: Identification of a putative cis element
817 involved in photoregulation. *Molecular Cell Biology Research Communications* **3**, 292–
818 298.

819 **Fairman ME, Maroney PA, Wang W, Bowers HA, Gollnick P, Nilsen TW, Jankowsky**
820 **E.** 2004. Protein displacement by DExH/D 'RNA helicases' without duplex unwinding.
821 *Science* **304**, 730–734.

822 **Gene Ontology Consortium.** 2021. The Gene Ontology resource: enriching a GOld
823 mine. *Nucleic Acids Research* **49**, D325–D334.

824 **Georg J, Dienst D, Schürgers N, et al.** 2014. The small regulatory RNA SyR1/PsrR1
825 controls photosynthetic functions in cyanobacteria. *The Plant Cell* **26**, 3661–3679.

826 **Georg J, Rosana ARR, Chamot D, Migur A, Hess WR, Owttrim GW.** 2019.
827 Inactivation of the RNA helicase CrhR impacts a specific subset of the transcriptome in
828 the cyanobacterium *Synechocystis* sp. PCC 6803. *RNA Biology* **16**, 1205–1214.

829 **Gierga G, Voss B, Hess WR.** 2012. Non-coding RNAs in marine *Synechococcus* and
830 their regulation under environmentally relevant stress conditions. *The ISME Journal* **6**,
831 1544–1557.

832 **Gong Z, Dong C-H, Lee H, Zhu J, Xiong L, Gong D, Stevenson B, Zhu J-K.** 2005. A
833 DEAD box RNA helicase is essential for mRNA export and important for development
834 and stress responses in *Arabidopsis*. *The Plant Cell* **17**, 256–267.

835 **Gruber AR, Bernhart SH, Lorenz R.** 2015. The ViennaRNA Web Services. In: Picardi
836 E, ed. *Methods in Molecular Biology. RNA Bioinformatics*. New York, NY: Springer, 307–
837 326.

838 **Hihara Y, Sonoike K, Kanehisa M, Ikeuchi M.** 2003. DNA microarray analysis of
839 redox-responsive genes in the genome of the cyanobacterium *Synechocystis* sp. strain
840 PCC 6803. *Journal of Bacteriology* **185**, 1719–1725.

841 **Hishiya S, Hatakeyama W, Mizota Y, Hosoya-Matsuda N, Motohashi K, Ikeuchi M,**
842 **Hisabori T.** 2008. Binary reducing equivalent pathways using NADPH-thioredoxin
843 reductase and ferredoxin-thioredoxin reductase in the cyanobacterium *Synechocystis*
844 sp. strain PCC 6803. *Plant & Cell Physiology* **49**, 11–18.

845 **Holmqvist E, Wright PR, Li L, Bischler T, Barquist L, Reinhardt R, Backofen R,**
846 **Vogel J.** 2016. Global RNA recognition patterns of post-transcriptional regulators Hfq
847 and CsrA revealed by UV crosslinking in vivo. *The EMBO Journal* **35**, 991–1011.

848 **Hör J, Gorski SA, Vogel J.** 2018. Bacterial RNA biology on a genome scale. *Molecular*
849 *Cell* **70**, 785–799.

850 **Huang C-K, Shen Y-L, Huang L-F, Wu S-J, Yeh C-H, Lu C-A.** 2016. The DEAD-box
851 RNA helicase AtRH7/PRH75 participates in pre-rRNA processing, plant development
852 and cold tolerance in *Arabidopsis*. *Plant and Cell Physiology* **57**, 174–191.

853 **Huckauf J, Nomura C, Forchhammer K, Hagemann M.** 2000. Stress responses of
854 *Synechocystis* sp. strain PCC 6803 mutants impaired in genes encoding putative
855 alternative sigma factors. *Microbiology* **146**, 2877–2889.

856 **Jankowsky E, Gross C, Shuman S, Pyle A.** 2001. Active disruption of an RNA-protein
857 interaction by a DExH/D RNA helicase. *Science* **291**, 121–125.

858 **Jarmoskaite I, Russell R.** 2011. DEAD-box proteins as RNA helicases and
859 chaperones: DEAD-box proteins. *Wiley Interdisciplinary Reviews: RNA* **2**, 135–152.

860 **Jiang H-B, Song W-Y, Cheng H-M, Qiu B-S.** 2015. The hypothetical protein Ycf46 is
861 involved in regulation of CO₂ utilization in the cyanobacterium *Synechocystis* sp. PCC
862 6803. *Planta* **241**, 145–155.

863 **Jones PG, Mitta M, Kim Y, Jiang W, Inouye M.** 1996. Cold shock induces a major
864 ribosomal-associated protein that unwinds double-stranded RNA in *Escherichia coli*.
865 *Proceedings of the National Academy of Sciences USA* **93**, 76–80.

866 **Kadowaki T, Nishiyama Y, Hisabori T, Hihara Y.** 2015. Identification of OmpR-family
867 response regulators interacting with thioredoxin in the cyanobacterium *Synechocystis*
868 sp. PCC 6803. *PloS ONE* **10**, e0119107.

869 **Kappell AD, van Waasbergen LG.** 2007. The response regulator RpaB binds the high
870 light regulatory 1 sequence upstream of the high-light-inducible *hliB* gene from the
871 cyanobacterium *Synechocystis* PCC 6803. *Archives of Microbiology* **187**, 337–342.

872 **Khemici V, Prados J, Petrignani B, Di Nolfi B, Bergé E, Manzano C, Giraud C,**
873 **Linder P.** 2020. The DEAD-box RNA helicase CshA is required for fatty acid
874 homeostasis in *Staphylococcus aureus*. *PLoS Genetics* **16**, e1008779.

875 **Kiefer M, Nauerth BH, Volkert C, Ibberson D, Loreth A, Schmidt A.** 2020. Gene
876 function rather than reproductive mode drives the evolution of RNA helicases in sexual
877 and apomictic *Boechera*. *Genome Biology and Evolution* **12**, 656–673.

878 **Köbler C, Schultz S-J, Kopp D, Voigt K, Wilde A.** 2018. The role of the *Synechocystis*
879 sp. PCC 6803 homolog of the circadian clock output regulator RpaA in day-night
880 transitions. *Molecular Microbiology* **110**, 847–861.

881 **Komenda J, Knoppová J, Krynická V, Nixon PJ, Tichý M.** 2010. Role of FtsH2 in the
882 repair of Photosystem II in mutants of the cyanobacterium *Synechocystis* PCC 6803
883 with impaired assembly or stability of the CaMn(4) cluster. *Biochimica Biophysica Acta*
884 **1797**, 566–575.

885 **Kopf M, Klähn S, Scholz I, Matthiessen JKF, Hess WR, Voß B.** 2014. Comparative
886 analysis of the primary transcriptome of *Synechocystis* sp. PCC 6803. *DNA Research*
887 **21**, 527–539.

888 **Kujat SL, Owtrim GW.** 2000. Redox-regulated RNA helicase expression. *Plant*
889 *Physiology* **124**, 703–714.

890 **Langmead B, Salzberg SL.** 2012. Fast gapped-read alignment with Bowtie 2. *Nature*
891 *Methods* **9**, 357–359.

892 **Lehnik-Habrink M, Rempeters L, Kovacs AT, Wrede C, Baierlein C, Krebber H,**
893 **Kuipers OP, Stülke J.** 2013. DEAD-box RNA helicases in *Bacillus subtilis* have multiple
894 functions and act independently from each other. *Journal of Bacteriology* **195**, 534–544.

895 **Li M, Calteau A, Semchonok DA, Witt TA, Nguyen JT, Sassoon N, Boekema EJ,**
896 **Whitelegge J, Gugger M, Bruce BD.** 2019. Physiological and evolutionary implications
897 of tetrameric photosystem I in cyanobacteria. *Nature Plants* **5**, 1309–1319.

898 **Linder P, Jankowsky E.** 2011. From unwinding to clamping — the DEAD box RNA
899 helicase family. *Nature Reviews Molecular Cell Biology* **12**, 505–516.

900 **Linder P, Owtrim GW.** 2009. Plant RNA helicases: linking aberrant and silencing RNA.
901 *Trends in Plant Science* **14**, 344–352.

902 **Lu C-A, Huang C-K, Huang W-S, Huang T-S, Liu H-Y, Chen Y-F.** 2020. DEAD-box
903 RNA helicase 42 plays a critical role in pre-mRNA splicing under cold stress. *Plant*
904 *Physiology* **182**, 255–271.

905 **Macovei A, Vaid N, Tula S, Tuteja N.** 2012. A new DEAD-box helicase ATP-binding
906 protein (OsABP) from rice is responsive to abiotic stress. *Plant Signaling & Behavior* **7**,
907 1138–1143.

908 **Mahbub M, Hemm L, Yang Y, et al.** 2020. mRNA localization, reaction centre
909 biogenesis and thylakoid membrane targeting in cyanobacteria. *Nature Plants* **6**, 1179–
910 1191.

911 **Margulis L.** 1981. *Symbiosis in cell evolution: Life and its environment on the early*
912 *earth*. W.H.Freeman & Co Ltd.

913 **Martin M.** 2011. Cutadapt removes adapter sequences from high-throughput
914 sequencing reads. *EMBnet.journal* **17**, 10.

915 **Martin W, Kowallik K.** 1999. Annotated English translation of Mereschkowsky's 1905
916 paper 'Über Natur und Ursprung der Chromatophoren im Pflanzenreiche'. *European*
917 *Journal of Phycology* **34**, 287–295.

918 **Martin W, Rujan T, Richly E, Hansen A, Cornelsen S, Lins T, Leister D, Stoebe B,**
919 **Hasegawa M, Penny D.** 2002. Evolutionary analysis of *Arabidopsis*, cyanobacterial, and
920 chloroplast genomes reveals plastid phylogeny and thousands of cyanobacterial genes
921 in the nucleus. *Proceedings of the National Academy of Sciences USA* **99**, 12246–
922 12251.

923 **Matthes A, Schmidt-Gattung S, Köhler D, Forner J, Wildum S, Raabe M, Urlaub H,**
924 **Binder S.** 2007. Two DEAD-box proteins may be part of RNA-dependent high-
925 molecular-mass protein complexes in *Arabidopsis* mitochondria. *Plant Physiology* **145**,
926 1637–1646.

927 **Mazouni K, Domain F, Chauvat F, Cassier-Chauvat C.** 2003. Expression and
928 regulation of the crucial plant-like ferredoxin of cyanobacteria. *Molecular Microbiology*
929 **49**, 1019–1029.

930 **Mereschkowsky C.** 1905. Über Natur und Ursprung der Chromatophoren im
931 Pflanzenreiche. *Biol. Centralbl.* **25**, 593–604.

932 **Mingam A, Toffano-Nioche C, Brunaud V, Boudet N, Kreis M, Lecharny A.** 2004.
933 DEAD-box RNA helicases in *Arabidopsis thaliana*: establishing a link between
934 quantitative expression, gene structure and evolution of a family of genes: AtRH
935 expression by real-time Q-PCR. *Plant Biotechnology Journal* **2**, 401–415.

936 **Mutsuda M, Sugiura M.** 2006. Translation initiation of cyanobacterial *rbcS* mRNAs
937 requires the 38-kDa ribosomal protein S1 but not the Shine-Dalgarno sequence:
938 development of a cyanobacterial in vitro translation system. *The Journal of Biological*
939 *Chemistry* **281**, 38314–38321.

- 940 **Muzzopappa F, Kirilovsky D.** 2020. Changing color for photoprotection: The Orange
941 Carotenoid Protein. *Trends in Plant Science* **25**, 92–104.
- 942 **Nakagawa S, Niimura Y, Miura K, Gojobori T.** 2010. Dynamic evolution of translation
943 initiation mechanisms in prokaryotes. *Proceedings of the National Academy of Sciences*
944 USA **107**, 6382–6387.
- 945 **Nawaz G, Kang H.** 2017. Chloroplast-or mitochondria-targeted DEAD-box RNA
946 helicases play essential roles in organellar RNA metabolism and abiotic stress
947 responses. *Frontiers in plant science* **8**, 871.
- 948 **Nawaz G, Lee K, Park SJ, Kim Y-O, Kang H.** 2018. A chloroplast-targeted cabbage
949 DEAD-box RNA helicase BrRH22 confers abiotic stress tolerance to transgenic
950 Arabidopsis plants by affecting translation of chloroplast transcripts. *Plant Physiology*
951 and Biochemistry: PPB **127**, 336–342.
- 952 **Oun S, Redder P, Didier J-P, François P, Corvaglia A-R, Buttazzoni E, Giraud C,**
953 **Girard M, Schrenzel J, Linder P.** 2013. The CshA DEAD-box RNA helicase is
954 important for quorum sensing control in *Staphylococcus aureus*. *RNA Biology* **10**, 157–
955 165.
- 956 **Owtrim GW.** 2013. RNA helicases. *RNA Biology* **10**, 96–110.
- 957 **Pfeifer-Sancar K, Mentz A, Rückert C, Kalinowski J.** 2013. Comprehensive analysis
958 of the *Corynebacterium glutamicum* transcriptome using an improved RNAseq
959 technique. *BMC Genomics* **14**, 888.
- 960 **Pinto F, Thapper A, Sontheim W, Lindblad P.** 2009. Analysis of current and
961 alternative phenol based RNA extraction methodologies for cyanobacteria. *BMC*
962 *Molecular Biology* **10**, 79.
- 963 **Ponce-Toledo RI, Deschamps P, López-García P, Zivanovic Y, Benzerara K,**
964 **Moreira D.** 2017. An early-branching freshwater cyanobacterium at the origin of
965 plastids. *Current Biology* **27**, 386–391.
- 966 **Prakash JSS, Krishna PS, Sirisha K, Kanesaki Y, Suzuki I, Shivaji S, Murata N.**
967 2010. An RNA helicase, CrhR, regulates the low-temperature-inducible expression of
968 heat-shock genes *groES*, *groEL1* and *groEL2* in *Synechocystis* sp. PCC 6803.
969 *Microbiology* **156**, 442–451.
- 970 **Py B, Higgins CF, Krisch HM, Carpousis AJ.** 1996. A DEAD-box RNA helicase in the
971 *Escherichia coli* RNA degradosome. *Nature* **381**, 169–172.
- 972 **Redder P, Hausmann S, Khemici V, Yasrebi H, Linder P.** 2015. Bacterial versatility
973 requires DEAD-box RNA helicases. *FEMS Microbiology Reviews* **39**, 392–412.

974 **Riediger M, Hihara Y, Hess WR.** 2018. From cyanobacteria and algae to land plants:
975 The RpaB/Ycf27 regulatory network in transition. *Perspectives in Phycology* **5**, 13–25.

976 **Riediger M, Kadowaki T, Nagayama R, Georg J, Hihara Y, Hess WR.** 2019.
977 Biocomputational analyses and experimental validation identify the regulon controlled by
978 the redox-responsive transcription factor RpaB. *iScience* **15**, 316–331.

979 **Riediger M, Spät P, Bilger R, Voigt K, Maček B, Hess WR.** 2021. Analysis of a
980 photosynthetic cyanobacterium rich in internal membrane systems via gradient profiling
981 by sequencing (Grad-seq). *The Plant Cell* **33**, 248–269.

982 **Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY.** 1979. Generic
983 assignments, strain histories and properties of pure cultures of cyanobacteria. *Journal of*
984 *General Microbiology* **111**, 1–61.

985 **Ritter SPA, Lewis AC, Vincent SL, Lo LL, Cunha APA, Chamot D, Ensminger I,**
986 **Espie GS, Owtrim GW.** 2020. Evidence for convergent sensing of multiple abiotic
987 stresses in cyanobacteria. *Biochimica et Biophysica Acta* **1864**, 129462.

988 **Rocak S, Linder P.** 2004. DEAD-box proteins: the driving forces behind RNA
989 metabolism. *Nature Reviews Molecular Cell Biology* **5**, 232–241.

990 **Rogers GW, Komar AA, Merrick WC.** 2002. eIF4A: the godfather of the DEAD box
991 helicases. *Progress in Nucleic Acid Research and Molecular Biology* **72**, 307–331.

992 **Rosana ARR, Chamot D, Owtrim GW.** 2012a. Autoregulation of RNA helicase
993 expression in response to temperature stress in *Synechocystis* sp. PCC 6803. *PLoS*
994 *ONE* **7**, e48683.

995 **Rosana ARR, Ventakesh M, Chamot D, Patterson-Fortin LM, Tarassova O, Espie**
996 **GS, Owtrim GW.** 2012b. Inactivation of a low temperature-induced RNA helicase in
997 *Synechocystis* sp. PCC 6803: Physiological and morphological consequences. *Plant*
998 *and Cell Physiology* **53**, 646–658.

999 **Rosana ARR, Whitford DS, Fahlman RP, Owtrim GW.** 2016. Cyanobacterial RNA
1000 helicase CrhR localizes to the thylakoid membrane region and cosediments with
1001 degradosome and polysome complexes in *Synechocystis* sp. strain PCC 6803. *Journal*
1002 *of Bacteriology* **198**, 2089–2099.

1003 **Rosana ARR, Whitford DS, Migur A, Steglich C, Kujat-Choy SL, Hess WR, Owtrim**
1004 **GW.** 2020. RNA helicase-regulated processing of the *Synechocystis rimO-crhR* operon
1005 results in differential cistron expression and accumulation of two sRNAs. *Journal of*
1006 *Biological Chemistry* **19**, 6372–6386.

1007 **Rowland JG, Pang X, Suzuki I, Murata N, Simon WJ, Slabas AR.** 2010. Identification
1008 of components associated with thermal acclimation of photosystem II in *Synechocystis*
1009 sp. PCC6803. *PLoS One* **5**, e10511.

1010 **Rowland JG, Simon WJ, Prakash JSS, Slabas AR.** 2011. Proteomics reveals a role
1011 for the RNA helicase *crhR* in the modulation of multiple metabolic pathways during cold
1012 acclimation of *Synechocystis* sp. PCC6803. *Journal of Proteome Research* **10**, 3674–
1013 3689.

1014 **Schneider S, Schwer B.** 2001. Functional domains of the yeast splicing factor Prp22p.
1015 *The Journal of Biological Chemistry* **276**, 21184–21191.

1016 **Scholz I, Lange SJ, Hein S, Hess WR, Backofen R.** 2013. CRISPR-Cas systems in
1017 the cyanobacterium *Synechocystis* sp. PCC6803 exhibit distinct processing pathways
1018 involving at least two Cas6 and a Cmr2 protein. *PloS One* **8**, e56470.

1019 **Silva P, Thompson E, Bailey S, Kruse O, Mullineaux CW, Robinson C, Mann NH,**
1020 **Nixon PJ.** 2003. FtsH is involved in the early stages of repair of photosystem II in
1021 *Synechocystis* sp PCC 6803. *The Plant Cell* **15**, 2152–2164.

1022 **Singh AK, Sherman LA.** 2002. Characterization of a stress-responsive operon in the
1023 cyanobacterium *Synechocystis* sp. strain PCC 6803. *Gene* **297**, 11–19.

1024 **Sireesha K, Radharani B, Krishna PS, Sreedhar N, Subramanyam R, Mohanty P,**
1025 **Prakash JSS.** 2012. RNA helicase, CrhR is indispensable for the energy redistribution
1026 and the regulation of photosystem stoichiometry at low temperature in *Synechocystis* sp.
1027 PCC6803. *Biochimica et Biophysica Acta* **1817**, 1525–1536.

1028 **Tanner NK, Linder P.** 2001. DExD/H box RNA helicases: from generic motors to
1029 specific dissociation functions. *Molecular Cell* **8**, 251–262.

1030 **Tarassova OS, Chamot D, Owtrim GW.** 2014. Conditional, temperature-induced
1031 proteolytic regulation of cyanobacterial RNA helicase expression. *Journal of*
1032 *Bacteriology* **196**, 1560–1568.

1033 **Tyystjärvi T, Herranen M, Aro EM.** 2001. Regulation of translation elongation in
1034 cyanobacteria: membrane targeting of the ribosome nascent-chain complexes controls
1035 the synthesis of D1 protein. *Molecular Microbiology* **40**, 476–484.

1036 **Vinnemeier J, Hagemann M.** 1999. Identification of salt-regulated genes in the genome
1037 of the cyanobacterium *Synechocystis* sp. strain PCC 6803 by subtractive RNA
1038 hybridization. *Archives of Microbiology* **172**, 377–386.

1039 **Wang B, Chai H, Zhong Y, Shen Y, Yang W, Chen J, Xin Z, Shi H.** 2020. The DEAD-
1040 box RNA helicase SHI2 functions in repression of salt-inducible genes and regulation of
1041 cold-inducible gene splicing. *Journal of Experimental Botany* **71**, 1598–1613.

1042 **Whitford DS, Whitman BT, Owtrim GW.** 2021. Genera specific distribution of DEAD-
1043 box RNA helicases in cyanobacteria. *Microbial Genomics*.

1044 **Wu C-Y, Nagy PD.** 2020. Role reversal of functional identity in host factors: Dissecting
1045 features affecting pro-viral versus antiviral functions of cellular DEAD-box helicases in
1046 tombusvirus replication. *PLoS Pathogens* **16**, e1008990.

1047 **Yang Q, Jankowsky E.** 2005. ATP- and ADP-dependent modulation of RNA unwinding
1048 and strand annealing activities by the DEAD-box protein DED1. *Biochemistry* **44**,
1049 13591–13601.

1050 **Yeremenko N, Jeanjean R, Prommeenate P, Krasikov V, Nixon PJ, Vermaas WFJ,**
1051 **Havaux M, Matthijs HCP.** 2005. Open reading frame *ssr2016* is required for antimycin
1052 A-sensitive photosystem I-driven cyclic electron flow in the cyanobacterium
1053 *Synechocystis* sp. PCC 6803. *Plant & Cell Physiology* **46**, 1433–1436.

1054 **Zhan J, Steglich C, Scholz I, Hess WR, Kirilovsky D.** 2021. Inverse regulation of light
1055 harvesting and photoprotection is mediated by a 3'end-derived sRNA in cyanobacteria.
1056 *The Plant Cell* **33**, 358–380.

1057 **Zhang L, McSpadden B, Pakrasi HB, Whitmarsh J.** 1992. Copper-mediated regulation
1058 of cytochrome c553 and plastocyanin in the cyanobacterium *Synechocystis* 6803. *The*
1059 *Journal of Biological Chemistry* **267**, 19054–19059.

1060 **Zinchenko VV, Piven IV, Melnik VA, Shestakov SV.** 1999. Vectors for the
1061 complementation analysis of cyanobacterial mutants. *Russ J Genet.* **35**, 228–32.

1062

Tables

Table 1. Abundant transcripts recovered in CrhR co-IPs. Assignment to one of the three experiments (**Fig. 1**) is given by the numbers in the first column, followed by gene IDs according to Genbank file NC_000911 and localization of the peak within the coding region (CDS) or UTR. The final two rows provide the annotation of the interacting RNAs and information on their transcriptional regulation, either activation (A) or repression (R) by the transcription factor RpaB under low light (Riediger *et al.*, 2019) or with “redox”, if they were classified as redox-responsive genes (Hihara *et al.*, 2003; Ritter *et al.*, 2020). Detailed information on peak locations, enrichment factors and adjusted p values generated by the PEAKachu algorithm (Holmqvist *et al.*, 2016) are given in **Tables S3** to **S5**.

E	ID	Annotation	RpaB/ Redox
1, 2, 3	slr1834 5'UTR, CDS	PsaA, PSI P700 apoprotein subunit Ia, Fig. 3C	A,R
1, 3	slr1655 5'UTR, CDS	PSI subunit XI, PsaL, Fig. S3	A
3	slr0927 CDS, 3'UTR	PSII protein PsbD/D2	R
1	smr0009 CDS, 3'UTR	photosystem II reaction center protein PsbN	R
3	slr0906 CDS	PSII reaction center protein PsbB/CP47	R
3	slr1311 5'UTR, CDS, 3'UTR	PSII reaction center protein PsbA/D1	R
3	slr1867 5'UTR, CDS	PSII reaction center protein PsbA/D1	R
1	slr1674 5'UTR, CDS	protection of PSII from thermal damage	R, redox
3	slr0144 to slr0148 CDS	operon of PSII assembly proteins, Fed5	A
3	slr0342 5'UTR, CDS	PetB, cytochrome <i>b₆</i>	R
1	ssl0020 5'UTR, CDS	ferredoxin I, PetF	R
2, 3	slr0199 5'UTR, CDS	plastocyanin, PetE	
3	slr2076 CDS	groEL1 chaperonin	
3	slr2067 5'UTR, CDS	allophycocyanin alpha subunit ApcA	
3	slr1986 CDS	allophycocyanin beta subunit ApcB	
3	ssr3383 5'UTR, CDS	allophycocyanin-associated linker ApcC, Fig. 4	
3	slr0335 5'UTR, CDS	PBS LCM core-membrane linker ApcE	A
3	slr2051 5'UTR, CDS 3'UTR	PBS rod-core linker CpcG1	
3	slr0009 to slr0012 CDS	RuBisCO operon <i>rbcLXS</i>	
1, 2, 3	slr0373-slr0374 CDS	Ycf46, CO ₂ utilization, Fig. 3B	redox

1	ssr2016 CDS	Pgr5, soluble electron acceptor	R
1, 2	slr0228 CDS	FtsH2 protease	R
1	slr1963 CDS	Water-soluble carotenoid protein OCP	redox
1, 3	slr0623 5'UTR, CDS	thioredoxin A	
1, 2, 3	slr1545 5'UTR, CDS	Extracytoplasmic function (ECF) sigma factor SigG, Fig. 3A	A
1, 2, 3	slr0083 5'UTR, CDS	DEAD-box RNA helicase CrhR, Fig. 5	redox
1	slr0923 5'UTR, CDS	YCF65; 30S ribosomal protein 3 PSRP-3	R
1	slr1638 5'UTR, CDS	Hypothetical protein	R
1, 2	slr0551 CDS	ribonuclease J	redox
1	slr0541 5'UTR	<i>desC/des9</i> , Acyl-CoA desaturase	A
1	slr0006 5'UTR, CDS	putative aminotransferase AspC	A
1, 3	slr1841 CDS	Porin, major outer membrane protein	R
1	slr0915 CDS, 3'UTR; <i>trnL</i> intron; 6803t34 exon 2; slr0917 5'UTR	group I intron, tRNA(Leu) ^{UAA} exon 2, endonuclease; <i>bioF</i> , 8-amino-7-oxononanoate synthase	R
3	6803t34 exon 1, <i>trnL</i> intron, slr0915 5'UTR, CDS	tRNA(Leu) ^{UAA} exon 1, group I intron, intron endonuclease	R

1075

1076

Figure legends

Fig. 1. Overview on the experimental flow. (A) Schematic representation of the cyanobacterial strains created for the chosen strategy. A deletion mutant, $\Delta crhR$, previously generated by replacement of the *crhR* gene (*slr0083*) by a spectinomycin (Sp^R) resistance gene by homologous recombination (Prakash *et al.*, 2010) served as the initial platform. In this background, strains were engineered that express the native form of CrhR ($CrhR_{WT}$) or a K57A substitution ($CrhR_{K57A}$), both translationally fused to an N-terminal triple FLAG tag in the conjugative vector pVZ321 under control of the copper-inducible P_{petE} promoter. A strain expressing only the 3xFLAG-tag was constructed as a control ($\Delta crhR_{control}$). **(B)** Experimental design. Experiment 1: co-IP of the UV-crosslinked RNA- $CrhR_{WT}$ complexes was performed from *Synechocystis* $CrhR_{WT}$ cultures exposed to cold stress at 20°C for 2 h or, Experiment 2, grown at 30°C. Experiment 3: co-IP of UV-crosslinked RNA- $CrhR_{K57A}$ complexes was performed from cultures grown at 30°C. In parallel, to control for non-specific enrichment, an RNA mock co-IP was performed with the $\Delta crhR_{control}$ strain. Cultures in Experiments 1 to 3 were grown in FWO medium using the CellDEG system. Each experiment was performed in biological duplicates and cDNA libraries were prepared from the enriched RNA and from the total RNA of all respective cultures omitting the co-IP step.

Fig. 2. Phenotypic characterization of wild type, $CrhR_{WT}$, $CrhR_{K57A}$ and the $\Delta crhR_{control}$ strains. (A) Growth of the wild type (WT), $CrhR_{WT}$, $CrhR_{K57A}$ and $\Delta crhR_{control}$ strains was determined spectrophotometrically by measuring the OD_{750} of cultures grown at 20°C in FWO medium. The experiments were performed in biological triplicates and standard deviations are indicated. **(B)** Pigmentation in the wild type (WT), $CrhR_{WT}$, $CrhR_{K57A}$ and $\Delta crhR_{control}$ cultures at an OD_{750} of 0.52 grown at 20°C in BG-11 medium. Spectra were normalized to 750 nm. **(C)** Visual appearance of the four strains at the indicated OD_{750} . Strains were grown in the presence of antibiotics and 2 μM Cu^{2+} .

Fig. 3. Peak pattern of transcripts enriched in CrhR co-IPs. (A) The region containing the 5' UTR, start codon and following codons were enriched for gene *slr1545* encoding the RNA polymerase sigma factor SigG. This peak was significant in all three experiments, using CrhR_{K57A} or CrhR_{WT} as bait proteins at the standard growth temperature of 30°C (read coverage in red and brown) as well as with CrhR_{WT} at 20°C (cold stress, coverage in blue). **(B)** Reads yielding a peak in the middle of the first gene and extending into the entire intergenic space between two genes and including the 5' UTR, start codon and following codons of the second gene were recovered for the dicistronic operon *slr0373* and *slr0374*. **(C)** Multiple sharp peaks indicating a multitude of interactions between CrhR and the dicistronic *psaAB* mRNA. *psaAB* encode the PSI P700 apoprotein subunits Ia and A2. In all three panels, the enrichment from reads recovered from CrhR co-IP experiments is shown (red, brown and blue) together with the coverage from RNA-Seq analyses of the respective total transcriptomes in shades of grey as indicated. The numbers along the x-axis correspond to chromosomal nucleotide positions in *Synechocystis* (Genbank accession number NC_000911.1), while the normalized numbers of reads are given on the y-axis.

Fig. 4. CrhR and *apcC* transcript interaction. Reads recovered from CrhR co-IP experiments cover the 5' UTR and almost the entire coding sequence of *apcC/ssr3383* that encodes the 7.8 kDa allophycocyanin-associated linker polypeptide ApcC. Only the reads recovered in experiment 3 using CrhR_{K57A} as bait protein were significantly enriched (red) but an enrichment was also observed in CrhR_{WT} cells grown at 30°C (brown). The sRNA ApcZ originates from within the *apcC* coding sequence (boxed). The 5' end of ApcZ was previously mapped to position 1431853, 46 nt upstream of the *apcC* stop codon (Zhan *et al.*, 2021). The secondary structure of ApcZ is shown in the inset. The interaction decreases in the region from which ApcZ originates, colored light green in both the transcript and in the corresponding segment of ApcZ secondary structure where it corresponds to half of the first helix of the predicted secondary structure (nt 1 to

30). The RNAfold algorithm was used for the secondary structure prediction (Gruber *et al.*, 2015).

Fig. 5. CrhR interaction with its own transcript *in vivo* and *in vitro*. (A) Enrichment of *crhR* mRNA in the UV-crosslinking RNA co-IP from *Synechocystis* CrhR_{WT} grown at 20°C or 30°C (experiments 1 and 2). The colored plots show the read coverage of the enriched RNAs, as indicated. (B) EMSA showing binding of recombinant His-tagged CrhR to the *crhR* mRNA. For *crhR* transcript synthesis, a 150 nt DNA fragment starting from the start codon of the *crhR*/*slr0083* ORF was amplified with primers EMSA_CrhR-T7_Fw (carrying the T7 promoter sequence followed by two Gs) and EMSA_CrhR-T7_Rv (Table S1), matching to the genomic positions 2887644 to 2887793. The resulting 150 nt transcript was Cy3 labelled. Binding of 0.2 pmol of the Cy3 labelled transcripts with the indicated amounts of purified recombinant His-tagged CrhR protein was performed in the presence of poly dI-dC (1 µg).

Fig. 6. Hypothetical model for CrhR self-regulation. The translational start region of the *crhR* transcript contains a predicted secondary structure and coincides with a sharp peak of mRNA fragments recovered in all three co-IP experiments (Fig. 5A) and was verified to interact with CrhR in EMSA assays (Fig. 5B). Within this structure the AUG signal for ribosome interaction is buried (black oval) and is followed by a 9-nt sequence that is complementary to the 3' end of the upstream *slr0082* (*rimO*) gene (red boxes). Initially, *crhR* is transcribed as a dicistronic message together with *rimO*. Operon discoordination occurs by RNase E cleavage three or four nt upstream the *rimO* stop codon within the RNase E cleavage motif (black arrows), as shown previously (Rosana *et al.*, 2020). We propose that an abundance of CrhR enhances its interaction with its own mRNA by clamping the shown structure, effectively suppressing translation. Upon stress-induced activation of CrhR RNA helicase activity, this RNA secondary structure is rearranged, making the 9-nt element available for interaction with the complementary region in *rimO* (red double arrow). Further CrhR action generates the conserved single-

1163 stranded RNA motif required for processing by RNase E cleavage. The resulting
 1164 monocistronic *crhR* transcript is stabilized and the detected dominating mRNA form
 1165 while the remaining *rimO* transcript segment showed a drastically reduced half-life
 1166 (Rosana *et al.*, 2012a, 2020). Hence, CrhR is proposed to feedback regulate its own
 1167 expression at two post-transcriptional levels, via operon discoordination-stabilization and
 1168 at the level of translation.

1169

A **B**

Fig. 1. Overview on the experimental flow. (A) Schematic representation of the cyanobacterial strains created for the chosen strategy. A deletion mutant, *ΔcrhR*, previously generated by replacement of the *crhR* gene (*slr0083*) by a spectinomycin (Sp^R) resistance gene by homologous recombination (Prakash *et al.*, 2010) served as the initial platform. In this background, strains were engineered that express the native form of CrhR (CrhR_{WT}) or a K57A substitution (CrhR_{K57A}), both translationally fused to an N-terminal triple FLAG tag in the conjugative vector pVZ321 under control of the copper-inducible *P_{petE}* promoter. A strain expressing only the 3xFLAG-tag was constructed as a control (ΔcrhR_{control}). **(B)** Experimental design. Experiment 1: co-IP of the UV-crosslinked RNA-CrhR_{WT} complexes was performed from *Synechocystis* CrhR_{WT} cultures exposed to cold stress at 20°C for 2 h or, Experiment 2, grown at 30°C. Experiment 3: co-IP of UV-crosslinked RNA-CrhR_{K57A} complexes was performed from cultures grown at 30°C. In parallel, to control for non-specific enrichment, an RNA mock co-IP was performed with the ΔcrhR_{control} strain. Cultures in Experiments 1 to 3 were grown in FWO medium using the CellDEG system. Each experiment was performed in biological duplicates and cDNA libraries were prepared from the enriched RNA and from the total RNA of all respective cultures omitting the co-IP step.

Fig. 2. Phenotypic characterization of wild type, $CrhR_{WT}$, $CrhR_{K57A}$ and the $\Delta crhR_{control}$ strains. (A) Growth of the wild type (WT), $CrhR_{WT}$, $CrhR_{K57A}$ and $\Delta crhR_{control}$ strains was determined spectrophotometrically by measuring the OD₇₅₀ of cultures grown at 20°C in FWO medium. The experiments were performed in biological triplicates and standard deviations are indicated. (B) Pigmentation in the wild type (WT), $CrhR_{WT}$, $CrhR_{K57A}$ and $\Delta crhR_{control}$ cultures at an OD₇₅₀ of 0.52 grown at 20°C in BG-11 medium. Spectra were normalized to 750 nm. (C) Visual appearance of the four strains at the indicated OD₇₅₀. Strains were grown in the presence of antibiotics and 2 μM Cu²⁺.

Fig. 3. Peak pattern of transcripts enriched in CrhR co-IPs. (A) The region containing the 5' UTR, start codon and following codons were enriched for gene *slr1545* encoding the RNA polymerase sigma factor SigG. This peak was significant in all three experiments, using CrhR_{K57A} or CrhR_{WT} as bait proteins at the standard growth temperature of 30°C (read coverage in red and brown) as well as with CrhR_{WT} at 20°C (cold stress, coverage in blue). (B) Reads yielding a peak in the middle of the first gene and extending into the entire intergenic space between two genes and including the 5' UTR, start codon and following codons of the second gene were recovered for the dicistronic operon *slr0373* and *slr0374*. (C) Multiple sharp peaks indicating a multitude of interactions between CrhR and the dicistronic *psaAB* mRNA. *psaAB* encode the PSI P700 apoprotein subunits Ia and A2. In all three panels, the enrichment from reads recovered from CrhR co-IP experiments is shown (red, brown and blue) together with the coverage from RNA-Seq analyses of the respective total transcriptomes in shades of grey as indicated. The numbers along the x-axis correspond to chromosomal nucleotide positions in *Synechocystis* (Genbank accession number NC_000911.1), while the normalized numbers of reads are given on the y-axis.

Fig. 4. CrhR and *apcC* transcript interaction. Reads recovered from CrhR co-IP experiments cover the 5' UTR and almost the entire coding sequence of *apcC/ssr3383* that encodes the 7.8 kDa allophycocyanin-associated linker polypeptide ApcC. Only the reads recovered in experiment 3 using CrhR_{K57A} as bait protein were significantly enriched (red) but an enrichment was also observed in CrhR_{WT} cells grown at 30°C (brown). The sRNA ApcZ originates from within the *apcC* coding sequence (boxed). The 5' end of ApcZ was previously mapped to position 1431853, 46 nt upstream of the *apcC* stop codon (Zhan *et al.*, 2021). The secondary structure of ApcZ is shown in the inset. The interaction decreases in the region from which ApcZ originates, colored light green in both the transcript and in the corresponding segment of ApcZ secondary structure where it corresponds to half of the first helix of the predicted secondary structure (nt 1 to 30). The RNAfold algorithm was used for the secondary structure prediction (Gruber *et al.*, 2015).

Fig. 5. CrhR interaction with its own transcript *in vivo* and *in vitro*. **(A)** Enrichment of *crhR* mRNA in the UV-crosslinking RNA co-IP from *Synechocystis* CrhR_{WT} grown at 20°C or 30°C (experiments 1 and 2). The colored plots show the read coverage of the enriched RNAs, as indicated. **(B)** EMSA showing binding of recombinant His-tagged CrhR to the *crhR* mRNA. For *crhR* transcript synthesis, a 150 nt DNA fragment starting from the start codon of the *crhR/sl0083* ORF was amplified with primers EMSA_CrhR-T7_Fw (carrying the T7 promoter sequence followed by two Gs) and EMSA_CrhR-T7_Rv (Table S1), matching to the genomic positions 2887644 to 2887793. The resulting 150 nt transcript was Cy3 labelled. Binding of 0.2 pmol of the Cy3 labelled transcripts with the indicated amounts of purified recombinant His-tagged CrhR protein was performed in the presence of poly dI-dC (1 µg).

Fig. 6. Hypothetical model for CrhR self-regulation. The translational start region of the *crhR* transcript contains a predicted secondary structure and coincides with a sharp peak of mRNA fragments recovered in all three co-IP experiments (Fig. 5A) and was verified to interact with CrhR in EMSA assays (Fig. 5B). Within this structure the AUG signal for ribosome interaction is buried (black oval) and is followed by a 9-nt sequence that is complementary to the 3' end of the upstream *slr0082* (*rimO*) gene (red boxes). Initially, *crhR* is transcribed as a dicistronic message together with *rimO*. Operon discooordination occurs by RNase E cleavage three or four nt upstream the *rimO* stop codon within the RNase E cleavage motif (black arrows), as shown previously (Rosana *et al.*, 2020). We propose that an abundance of CrhR enhances its interaction with its own mRNA by clamping the shown structure, effectively suppressing translation. Upon stress-induced activation of CrhR RNA helicase activity, this RNA secondary structure is rearranged, making the 9-nt element available for interaction with the complementary region in *rimO* (red double arrow). Further CrhR action generates the conserved single-stranded RNA motif required for processing by RNase E cleavage. The resulting monocistronic *crhR* transcript is stabilized and the detected dominating mRNA form while the remaining *rimO* transcript segment showed a drastically reduced half-life (Rosana *et al.*, 2012a, 2020). Hence, CrhR is proposed to feedback regulate its own expression at two post-transcriptional levels, via operon discooordination-stabilization and at the level of translation.