Modeling minimum viable population size with multiple genetic problems of small populations

Peter NABUTANYI and Meike J. WITTMANN

Bielefeld University

Abstract

An important goal for conservation is to define minimum viable population (MVP) sizes for long-term persistence. Although many MVP size estimates focus on ecological processes, with increasing evidence for the role of genetic problems in population extinction, conservation practitioners have also increasingly started to incorporate inbreeding depression (ID). However, small populations also face other genetic problems such as mutation accumulation (MA) and loss of genetic diversity through genetic drift that are usually factored into population viability assessments only via verbal arguments. Comprehensive quantitative theory on interacting genetic problems is missing. Here we develop eco-evolutionary quantitative models that track both population size and levels of genetic diversity. Our models assume a biallelic multilocus genome whose loci can be under either a single or interacting genetic forces. In addition to mutation-selection-drift balance (for loci facing ID and MA), we include three forms of balancing selection (for loci where variation is lost through genetic drift). We define MVP size as the lowest population size that avoids an eco-evolutionary extinction vortex after a time sufficient for an equilibrium allele frequency distribution to establish. Our results show that MVP size decreases rapidly with increasing mutation rates for populations whose genomes are only under balancing selection, while for genomes under mutation-selection-drift balance, the MVP size increases rapidly. MVP sizes also increase rapidly with increasing number of loci under

19 the same or different selection mechanisms until a point is reached at which even arbitrarily
20 large populations cannot survive anymore. In the case of fixed number of loci under selection,
21 interaction of genetic problems did not necessarily increase MVP sizes. To further enhance our
22 understanding about interaction of genetic problems, there is need for more empirical studies
23 to reveal how different genetic processes interact in the genome.

24 Introduction

25 In conservation management, the desire to protect threatened populations or species given limited
 26 available resources gave birth to the concept of a minimum viable population (MVP) size (Shaffer,
 27 1981). Minimum viable population size is commonly defined as the lowest size that leads to popu-
 28 lation persistence for a specified duration with a set persistence probability, e.g., persistence for at
 29 least 1000 years with 99% probability or persistence for at least 50 generations with a probability
 30 of 90% (Shaffer, 1981; Traill et al., 2007). Alternatively, MVP size can be defined as the lowest size
 31 above which a substantial increase in extinction time occurs (Reed & Bryant, 2000) or in terms of
 32 fitness thresholds instead of persistence probability (Reed, 2005). So far, the choice of time and
 33 probability thresholds is arbitrary, varies among researchers, and across populations (Traill et al.,
 34 2007). This lack of consensus can make it hard to quantify relative extinction risk among different
 35 populations.

36 The viability of a small population and thus its MVP size depends on various ecological and genetic
 37 processes. Relevant ecological processes include demographic and environmental stochasticity, as
 38 well as Allee effects, e.g. through mate or pollen limitation at low density. In this study, however,
 39 we focus on the genetic processes shaping population viability and MVP sizes. In small populations,
 40 genetic problems such as inbreeding depression (ID), mutation accumulation (MA), and loss of
 41 genetic variation are inevitable (Lynch et al., 1995; Higgins & Lynch, 2001; Estes et al., 2004;
 42 Frankham, 2005; Coron et al., 2013). For example, increased homozygosity of deleterious alleles
 43 leads to a decrease in various fitness components (Spielman et al., 2004a; Bensch et al., 2006;
 44 Blomqvist et al., 2010; Palomares et al., 2012). Generally, genetic diversity is known to positively
 45 correlate with population size and influences population fitness (Reed & Frankham, 2003; Reed,
 46 2005; Leimu et al., 2006; Hensen & Wesche, 2006). For instance, over 70% of threatened species have
 47 lower heterozygosity than their taxonomically related non-threatened taxa (Spielman et al., 2004b).
 48 However, diversity at neutral marker loci and high population size are not always a good predictor
 49 of population fitness (Yates et al., 2019; Teixeira & Huber, 2021). Other factors such as the genetic
 50 architecture of the population may have to be invoked for a better prediction of population fitness
 51 (Grossen et al., 2020; Kyriazis et al., 2021; Kardos & Luikart, 2021). But all in all, even if it is not

52 always clear what aspects of diversity are most important and how they should be measured, there
53 is increasing evidence for the importance of genetic diversity in population extinction (Frankham,
54 2005).

55 To date, inbreeding depression is the only genetic problem of small population size that is routinely
56 included in MVP studies, e.g. those that employ the software package *VORTEX* (Lacy, 1993). Thus,
57 other genetic problems such as mutation accumulation and loss of genetic diversity through drift
58 are ignored although they are known to be important contributors to extinction risk (Frankham,
59 2005). In nature, these problems may occur simultaneously in a small population. However, to the
60 best of our knowledge, no studies exist that quantify the expected interaction of the different genetic
61 problems. At best, the interaction is verbally projected to increase extinction risk.

62 Another key factor for long-term population persistence is how genetic diversity is maintained.
63 It is now clear that both mutation-selection-drift equilibrium and balancing selection mechanisms
64 maintain genetic variation in populations (e.g., Hughes & Nei, 1988; Sharp & Agrawal, 2018), but
65 balancing selection is still ignored in MVP studies. Balancing selection comes in various forms
66 such as heterozygote advantage (HA), negative frequency-dependent selection (FD), and fluctuating
67 selection with dominance reversals (FS), among others. The different forms of balancing selection
68 can act in isolation or in some combination (e.g., Glémin et al., 2005; Hedrick, 2012; Bergland et al.,
69 2014; Chen et al., 2015; Chapman et al., 2019; Mérot et al., 2020).

70 Our study has thus three goals: 1. to develop methods to account for genetic problems other than
71 inbreeding depression in MVP studies, including loss of genetic variation at loci under balancing
72 selection, 2. to quantify the effect of the interaction of two or more genetic problems on extinction
73 risk, and 3. to find ways to reduce the arbitrariness of time and persistence probability thresholds
74 in MVP analyses. To this end, we developed eco-evolutionary models using three approaches, i.e.,
75 individual-based model (IBM), diffusion approximation, and equilibrium-based Markov chain ap-
76 proaches. We defined MVP size as the minimum size below which an eco-evolutionary extinction
77 vortex occurs (Nabutanyi & Wittmann, 2021). An eco-evolutionary vortex occurs when the per-
78 capita population growth rate turns negative. For the stochastic model, we used 95% persistence
79 probability, but lower or higher probabilities did not substantially affect our MVP size estimates. To

80 reduce the arbitrariness of time thresholds used in most definitions of MVP size, we either estimated
81 MVP size based on time to attain genetic equilibria or even used a long-term stable equilibrium
82 perspective.

83 General Model

84 Our models considered a diploid hermaphroditic population initially at carrying capacity K with
85 discrete-time population growth and non-overlapping generations. We assumed a genome with
86 multiple unlinked bi-allelic loci. The genome was acted on by either one selection mechanism at
87 all loci or multiple mechanisms simultaneously, each at a set of loci. We developed three different
88 approaches to quantifying minimum viable population size (Fig. 1). The time-iterating approaches
89 (one stochastic, one deterministic) tracked absolute fitness W and population size N until either the
90 population went extinct or a set maximum number of generations (in this study, 4000 generations)
91 was reached. The MVP size was the lowest population size needed for at least 1 individual to survive
92 at the end of the simulation. For the stochastic IBM, this condition had to be satisfied in at least
93 95% of the replicates. We also used a time-independent approach that assumed a population at
94 equilibrium with a stationary distribution of allele frequencies and constant population size. Here
95 we defined MVP size as the minimum size required for absolute fitness W at equilibrium to be at
96 least 1.

Figure 1: Overview of the three approaches to quantifying minimum viable population (MVP) sizes. For each population, we start with the required population parameters such as carrying capacity K , which also acts as the initial population size N_0 , intrinsic growth rate r , the number of loci on the genome n , and initial allele frequency x_0 . We then chose which selection mechanism(s) acts on the genome and simulated the population via either time-iterating approaches (i.e., diffusion approximation approach (DAA) or individual-based model (IBM)) or the equilibrium-based Markov chain approach (MCA). In the IBM, we tracked genotypes (and fitness) of each individual at each locus, and in DAA, we tracked allele frequency distribution $f(x, t)$ from which offspring viability V_t , absolute fitness W_t , and population size N_t were then determined. The iterations stopped when either the population went extinct or the maximum pre-set number of generations T was reached (whichever came first). In the MCA, we calculated the transition matrix for the number of allele copies at a constant population size from which the equilibrium allele-frequency distribution $f(x)$, viability V , and absolute fitness W , were sequentially determined. Starting with $K = 10$, we incremented K by 10 in the MVP loop until either the MVP criteria was satisfied or $K > 150000$ (whichever came first).

97 Selection Mechanisms

98 We considered five selection mechanisms, i.e., fluctuating selection with dominance reversals (FS),
 99 negative frequency-dependent selection (FD), heterozygote advantage (HA), inbreeding depression
 100 (ID), and mutation accumulation (MA). For HA, ID, and MA, the relative fitnesses of the three
 101 genotypes AA , Aa , and aa at a given locus had the general form $w_{AA} = 1 - s_A$, $w_{Aa} = 1 - h \cdot s_a$, and
 102 $w_{aa} = 1 - s_a$, with the three mechanisms differing in the magnitude of selection (s_i) and dominance
 103 (h) coefficients (Fig. 2a). In HA, $0 < s_A, s_a < 1$ and $h = 0$ such that the heterozygotes had higher
 104 relative fitness than either homozygote. For both ID and MA, $s_A = 0$, $0 < s_a \leq 1$ and $0 \leq h \leq 1$.
 105 Thus, allele A was considered the wild-type allele whose homozygote had a maximum relative fitness
 106 while a was the deleterious allele. The difference between our example scenarios for ID and MA
 107 arose in the magnitude of s_a and h , i.e., small h and potentially large s_a for ID while larger h and
 108 smaller s_a for MA (Fig. 2).

Figure 2: Relative fitness of the genotypes under different selection mechanisms. (a) For heterozygote advantage (HA), inbreeding depression (ID), and mutation accumulation (MA), the relative fitness is constant through generations. (b) For fluctuating selection with dominance reversals (FS), the relative fitness temporally fluctuates cyclically. We implemented this mechanism such that the heterozygote genotype is not the fittest in every generation but in any complete cycle of κ generations (we arbitrary used $\kappa = 50$), it emerged with a higher relative geometric mean fitness than either homozygote.

For FS and FD, the relative fitnesses of genotypes fluctuated over time. In FS, the fluctuations depended on the temporal changes in selection coefficients (Fig. 2b and Appendix S1). With FD, the fluctuations depended on allele frequency whereby a genotype with a rarer allele was more fit. Thus, we set $w_{AA,t} = 1 - s_A x_t$, $w_{aa,t} = 1 - s_a(1 - x_t)$ and $w_{Aa,t} = 1 - \frac{x_t(s_A - s_a) + s_a}{2}$, where s_i was the maximum selection coefficient against the respective homozygote and x_t is the frequency of allele A . Three of the genetic mechanisms (HA, FS, and FD) give rise to balancing selection such that in an infinite population, both alleles can be maintained even without mutation. For these mechanisms, the mutation rates were assumed to be symmetric, i.e., $\mu_A = \mu_a$, where μ_A is the mutation rate for allele A to allele a and vice versa for μ_a . On the other hand, ID and MA do not yield balancing selection and genetic diversity at loci under these mechanisms is due to the mutation-selection-drift

equilibrium. Here we assumed that back mutations are very rare and set $\mu_a = \frac{\mu_A}{100}$ for both ID and MA.

Individual-Based Model (IBM) Simulations

The initial population size was drawn from a Poisson distribution with mean K . Each initial individual was randomly assigned two gene copies at each of the n loci. The initial A allele frequency was 0.5 for all balancing selection mechanisms and 0.95 for ID and MA.

In each generation, each parent produced a random number of offspring drawn from a Poisson distribution with mean $E = e^r$, where r is the intrinsic per-capita growth rate. The focal parent randomly chose a mating partner from all the available individuals including itself as a second parent for each offspring independently. For each locus, each partner independently and randomly passed on one of the available alleles to the respective offspring locus. Finally, the offspring alleles mutated with probability μ_A or μ_a .

Assuming multiplicative fitness across loci, each offspring was viable, i.e. survived, with probability, $V = \prod_{l=1}^n w_{g,l,t}$, where $w_{g,l,t}$ denotes the fitness of genotype $g \in \{AA, Aa, aa\}$ at locus l and generation t . After all individuals had reproduced, the surviving offspring replaced the parent generation. If there were more than K surviving offspring, only K offspring were randomly selected to replace the parents.

Each simulation run was iterated until either the population went extinct or generation 4000 was reached. This time was long enough for stationary allele frequency distributions to be attained under most selection scenarios except at very low mutation rates in ID and MA (see Results and Discussion sections).

Deterministic Approximations to the Individual-Based Model

There were two main steps in the deterministic approximations, namely, approximating the distribution f of allele frequency x and determining the absolute fitness W using f . For the first step, f was determined by either the diffusion approximation approach (DAA) or the Markov chain approach (MCA).

145 The diffusion approximation involved numerically solving the diffusion equation

$$\frac{\partial f(x, t)}{\partial t} = -\frac{\partial [a(x, t)f(x, t)]}{\partial x} + \frac{1}{2} \frac{\partial^2 [b(x, t)f(x, t)]}{\partial x^2} \quad (1)$$

146 for $f(x, t)$ in each generation subject to some initial and boundary conditions (Zhao et al., 2013;
147 Xu et al., 2019; Nabutanyi & Wittmann, 2021), where $a(x, t)$ and $b(x, t)$ are the mean and variance
148 of the change in allele frequency during an infinitesimal time interval, derived for each selection
149 mechanism (Appendix S2.1).

150 For the MCA, we assumed constant population size N and computed the transition probabilities of
151 allele A from i copies at time t to j copies at time $t + 1$ ($i, j \in \{0, 1, 2, \dots, 2N\}$). Assuming random
152 sampling, the number of copies of the A allele in the next generation follows a binomial distribution
153 with parameters $2N_t$ and x'' (the allele frequency after the action of selection and mutation). Due
154 to numerical problems at large N ($N > 500$) with the binomial distribution, we approximated
155 the transition probabilities using the Poisson distribution (when one of the alleles is rare) or the
156 normal distribution (Appendix S2.2). Given the transition probability matrix, we then obtained the
157 allele frequency distribution $f(x)$ as the eigenvector corresponding to the largest eigenvalue of the
158 transition matrix. Appendix S2.2.1 also shows how MCA can be made time-dependent.

159 In the second step, we began by estimating the mean offspring viability V from $f(x)$. We let $n_{\mathcal{M}}$ be
160 the number of loci under genetic mechanism $\mathcal{M}$. Then, assuming multiplicative fitness across loci
161 and independence between loci, the fitness from $\mathcal{M}$, is given by

$$V_{\mathcal{M}} = \prod_{i=1}^{n_{\mathcal{M}}} w(x) = w(x)^{n_{\mathcal{M}}}, \quad (2)$$

162 where $w(x)$ is the fitness at allele frequency x at a specific locus given by

$$w(x) = x^2 w_{AA} + 2x(1-x)w_{Aa} + (1-x)w_{aa}. \quad (3)$$

163 Because the exact allele frequency x at a given locus is not explicitly known, we calculated $V_{\mathcal{M}}$ using

the expected value of $w(x)$ under the allele-frequency distribution:

$$V_{\mathcal{M}} \approx \left(\sum f(x)w(x) \right)^{n_{\mathcal{M}}}, \quad (4)$$

where the sum is over all allele-frequency bins. Also, assuming independence between selection mechanisms, then, the offspring survival fitness $V = \prod_{\mathcal{M}} V_{\mathcal{M}}$.

The second step was completed by computing the absolute fitness, $W = VE$. Note that for DAA, the distribution f is calculated for every generation and so is $V_{\mathcal{M}}$ and W . Also, the population size N_{t+1} was determined from the previous generation t as

$$N_{t+1} = N_t \times \min\{W_t, \frac{K}{N_t}\}. \quad (5)$$

Results

In this section, we focus on results from HA, FS, ID, and MA. The results from FD are to a large extent similar to those from HA (see supplement Appendices S3 to S7).

For both isolated and interacting selection mechanisms, the absolute fitness initially decreased while the population remained at the carrying capacity K (Fig. 3, Appendix S3.1). For relatively high K , absolute fitness decreased to an equilibrium value of at least 1, while populations remained at K . However, at lower K , absolute fitness decreased below a minimum threshold of 1 (gray line on Fig. 3 row 2) while the populations remained at K , before rapidly decreasing to extinction at the time when absolute fitness just fell below 1. For scenarios with FS, there were expected periodic fluctuations in both population size and absolute fitness, and the absolute fitness fell below 1 in some generations without causing population extinction (Fig. 3 column 4). The lowest carrying capacity that avoided population extinction was taken as the MVP size (indicated in red, Fig. 3 row 1). The MVP size estimate from scenarios that involved ID (and MA) increased with increasing simulation time (Appendix S3.1.1)

In cases involving more than one selection mechanism, the individual mechanisms reduced maximum

185 offspring viability (which is 1) by different amounts that varied through time (Fig. 3, row 3). The
186 rate at which locus fitness decreased (and attained equilibrium) also differed between constituent
187 mechanisms.

Figure 3: Example trajectories under various carrying capacities for; Column 1: HA, Column 2: ID, Column 3: HA-ID and Column 4: HA-FS while for Row 1: population size, Row 2: absolute fitness and Row 3: offspring viability fitness for each constituent mechanism. The trajectories were obtained using the diffusion approximation approach (see Appendix S3.2 for some IBM trajectories). The gray horizontal line is the threshold absolute fitness of 1. For balancing selection, $s_a = s_A = 0.005$, $\mu_A = \mu_a = 0.0005$ and for ID $s_A = 0$, $s_a = 0.05$, $h = 0.01$, $\mu_A = 0.0005$, $\mu_a = \frac{\mu_A}{100}$. Other shared parameters are $n = 100$, $K = N_0$, $r = \ln(1.53)$. For scenarios with two selection mechanisms (columns 3 and 4), 50 loci were under each mechanism. For population trajectories from other scenarios, see Appendix S3.

Figure 4: Effect of increasing number of loci under selection on MVP size. The genome from populations with at most 100 loci (below the vertical gray line) was acted on by the same selection mechanism (indicated at the start of the line) (a) HA, (b) FS, (c) ID and (d) MA. Those genomes with above 100 loci, the mechanism indicated at the end of the line acted on the remaining loci, e.g., the line with HA—ID label implies the first 100 loci were acted on by HA while ID acted on the remaining loci. See Appendix S4.3 for results from more scenarios. Parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a = 0.0005$, for ID and MA, $s_A = 0$, $\mu_a = \frac{\mu_A}{100}$ while $h = 0.01$, $s_a = 0.05$ for ID and $h = 0.5$, $s_a = 0.005$ for MA. Shared parameters were $K = N_0$, $r = \ln(1.53)$.

188 In Fig. 4, we simulated populations with different number of loci under selection. We first varied
189 the number of loci under selection between 1 and 100 for each mechanism separately. Thus, genomes
190 with at most 100 loci under selection were acted on by one selection mechanism. The additional
191 loci were either under the same mechanism or one of the other three selection mechanisms. The
192 MVP size initially remained constant at 10 (our lowest tested population size) as the number of loci
193 increased until a certain number was reached where a rapid increase was registered. Even under
194 the same selection mechanism, the rate at which log MVP size increased decreased after a certain
195 number of loci. As more loci were added, log MVP size again increased rapidly, producing an
196 inflection point. Note though that on a linear scale, MVP size increases more and more rapidly with
197 increasing number of loci and there is no inflection point, except for MA which according to the MCA
198 mechanism still exhibits an inflection point (Appendix S4.3). At a certain critical number of loci,
199 MVP sizes exhibited an asymptotic increase (to infinity), indicating that no finite population size
200 was sufficient to keep the population at equilibrium. Note that we tested K values from 10 to 150000
201 (using DAA) and numbers of loci from 1 to 1000 for all the scenarios. Unlike for ID and MA, the
202 critical number of loci required for an asymptotic increase in MVP size was independent of mutation
203 rates for balancing selection mechanisms (Appendices S4 and S4.3). Also, MCA produces higher
204 MVP estimates than DAA in some cases of ID and MA scenarios, but the divergence disappears as
205 more loci are added.

206 The effect of mutation rate on MVPs varied strongly among genetic mechanisms (Fig. 5). For
207 genomes that were acted on by only balancing selection mechanisms (HA, FS) either separately or
208 simultaneously, the MVP size estimates generally decreased rapidly with mutation rates (Fig. 5).
209 By contrast, MVP estimates from genomes acted on by either ID, MA, or both increased rapidly.
210 With a fixed total number of loci under selection, in genomes with more than one mechanism, the
211 joint MVP size was in most cases closer to the harmonic (or geometric) mean of the MVP estimates
212 obtained when the constituent selection mechanisms acted in isolation than to the arithmetic mean
213 (Appendix S5.3). However, there were also exceptions where the joint MVP size was close to or
214 even larger than the arithmetic mean of single MVP sizes.

215 In most scenarios that contained only balancing selection, the estimates from DAA and MCA roughly

216 coincided. However, as mutation rates decrease in ID and MA, there were substantial quantitative
 217 differences at lower mutation rates between MVP estimates from DAA and MCA (Fig. 5 row 1).
 218 The IBM estimates (Fig. 5 row 2) were (slightly) higher than estimates from DAA, and converged
 219 better to the MCA and DAA estimates at higher MVP sizes. Also, the IBM estimates were fairly
 220 insensitive to levels of survival probability thresholds (Appendix S5.1).

Figure 5: Comparison of MVP size estimates across approaches with varying forward mutation rate, μ_A . Row 1: Results obtained from the Markov chain approach (MCA; dashed lines) and the diffusion approximation (DAA; solid lines). Genomes under (a) isolated mechanisms, (b) paired mechanisms and (c) three and four mechanisms. Row 2: Comparison of MVP size estimates from the IBM, DAA and MCA. (d) HA only, (e) ID only and (f) HA-FS-FD-ID-MA. The estimates from IBM assumed 95% survival probability of 60 replicates. In both rows, parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a$, and for ID and MA $s_A = 0$, $\mu_a = \frac{\mu_A}{100}$. ID and MA differed in h and s_a with row 1: $h = 0.5$, $s_a = 0.005$ for MA and $h = 0.01$, $s_a = 0.05$ for ID while row 2: $s_a = 0.005$ for both ID and MA but $h = 0.5$ for MA and $h = 0$ for ID. Other shared parameters were $K = N_0$, $r = \ln(1.53)$. The total number of loci n on the genome is 100 except for combinations with 3 mechanisms where it is 99 for purposes of equal distribution of loci between mechanisms.

Discussion

Small populations can be faced with multiple genetic problems simultaneously. In this study, we quantified the effect of such interactions on extinction risk by estimating MVP sizes. MVP size rapidly increased as more loci under selection were added. However, when the total number of loci in the genome was fixed, MVP size estimates were in most but not all of the scenarios we considered dominated by the mechanism that yielded the lowest estimate when acting alone on the genome.

Number of loci under selection

As the number of loci under selection increased, cumulative selection pressure on the genome increased and as a result, offspring viability decreased, thus increasing MVP size. Under balancing selection mechanisms, the critical number of loci above which no arbitrary population size survived was independent of mutation rates and decreased with increasing selection coefficients (Appendix S4). This is because, in an infinite population under symmetric balancing selection and symmetric mutation rates, equilibrium fitness is independent of mutation rates. Moreover, heterozygosity increases with increasing mutation rate, which increases population fitness. Thus MVP estimates are lower at higher mutation rates. By contrast, the critical number of loci under mutation-selection-balance was independent of the selection coefficient but decreased with an increase in mutation rates. This is in agreement with classical studies showing that equilibrium fitness only depends on mutation rate (μ_A) and not on selection coefficient (s_a) if $\mu_A \ll s_a$ (Haldane, 1937; Agrawal & Whitlock, 2012; Felsenstein, 2015).

Most of the predominance of the mechanism that yielded the lowest MVP estimate in the fixed number of loci scenario can be explained by the relationship between viability per locus and population size. In Appendix S6, we assume that viability increases linearly with population size, and show that the joint MVP is approximately equal to the harmonic mean of the MVPs obtained from the constituent mechanisms when acting in isolation. We also illustrate scenarios where the joint MVP can be even higher than the harmonic, geometric, and arithmetic means as also observed in some cases in our results.

247 DAA, MCA, and IBM

248 In many regions of parameter space, the three approaches largely agreed in their MVP size estimates
 249 both qualitatively and quantitatively, but there were also discrepancies and each approach has its
 250 advantages and disadvantages. The relatively low MVP estimates from the iterating approaches in
 251 ID and MA at low mutation rates can be attributed to lack of attainment of a stationary allele
 252 frequency distribution within the simulated period of time. For example, for mutation rate of
 253 $\mu_A = 0.00005$ and $\mu_a = \frac{\mu_A}{100}$, equilibrium is attained after at least $\frac{1}{\mu_A + \mu_a} \approx 20,000$ generations from
 254 mutation alone (see Felsenstein, 2015). The equilibrium-based MCA estimates did not depend on
 255 the length of the simulation period and the initial conditions. Therefore, MCA estimates can be
 256 considered as the "true" MVP estimates if one is interested in the possibility of a long-term stable
 257 population. However, MCA may give overly optimistic predictions for population survival if the
 258 population considered starts far from genetic equilibrium (Appendix S7; see also Kardos & Luikart,
 259 2021). For example, during founder events, population fragmentation, and bottlenecks, deleterious
 260 allele frequencies may be above equilibrium values such that populations may be unable to replace
 261 themselves in the early generations and go extinct before reaching a genetic equilibrium. The time-
 262 dependent approaches could account for such transient dynamics, but for this the initial genetic
 263 composition of the population needs to be known.

264 Unlike in the DAA and MCA approaches where allele frequencies evolve completely independently
 265 across loci, in the IBM evolution at one locus can potentially interfere with evolution at another locus,
 266 making the fitnesses of the genotypes at the focal locus more variable. IBMs also have the advantage
 267 that they take into account demographic stochasticity (which in some cases led to higher MVP size
 268 estimates than for the MCA and DAA approach). Therefore IBM gives a better representation of
 269 natural populations. However, IBMs are extremely time-consuming especially at higher K since
 270 many replicates are required to infer MVP sizes. Similarly, obtaining the transition matrix in the
 271 MCA is computationally impossible for higher K . On the other hand, the accuracy in DAA depends
 272 the discretization of allele frequency, x and can perform poorly for very small population sizes. Thus
 273 each of the approaches will be most useful in a different context: MCA to get long-term predictions
 274 for small populations, IBM for small populations when demographic stochasticity and inter-locus

interactions should be accounted for, and DAA for large populations.

Choice of the MVP Definition Thresholds

In the case of more than one threatened population, managers may have to prioritise allocation of available limited resources. Therefore, there is need to establish relative viability status of the populations e.g., by comparing required MVP sizes to current population sizes. Equilibrium-based MVP sizes could help make MVP sizes more comparable across populations, while still taking into account specific population parameters. In the IBM, our estimates were largely insensitive to survival thresholds (Appendix S5.1), possibly due to low sources of stochasticity in the model. We expect survival thresholds to be important in estimates that consider more ecological stochastic processes such as environmental stochasticity.

Furthermore, we chose an extinction threshold of 1 individual for the time-iterating approaches. However, larger thresholds could also be chosen for dioecious populations and populations that suffer from Allee effects due to mate-finding difficulties or other ecological problems at low population size. For the MCA with equilibrium assumptions, a quasi-extinction threshold of absolute fitness 1 was used. The absolute fitness of 1 acts as a boundary between population persistence ($W > 1$) and rapid population decline ($W < 1$) in an eco-evolutionary extinction vortex (Nabutanyi & Wittmann, 2021).

Concluding Remarks and Recommendations

Since our models assumed an idealized randomly mating population, the estimates in this paper should be compared to the genetically derived effective population sizes in the literature such as the long-term 500 estimate in the 50/500 rule of thumb (Franklin, 1980; Frankham et al., 2014). Our estimates range from 10 to about 5000 before the critical number of loci is reached. Some studies have proposed that census MVP values would need to be up to 10 times higher than the effective MVP sizes (Frankham, 1995). This would translate our estimates to the range 100 to 50000, which generally lies in the range reported by most studies, (Harcourt, 2002; Reed et al., 2003; Traill et al., 2007; Wang et al., 2019).

301 In nature, both mutation-selection balance and balancing selection contribute to standing genetic
302 variation (Sharp & Agrawal, 2018), potentially with the different genetic mechanisms playing a
303 larger role at different life stages. For instance, variation in offspring viability traits may often be
304 under mutation-selection balance while variation in traits linked to sexual reproduction may more
305 often be under some form(s) of balancing selection (Sharp & Agrawal, 2018). Our observations were
306 based on independence between loci. Epistasis and linkage disequilibrium may give different results.
307 Therefore, to improve extinction risk assessment, a better understanding of the underlying genetic
308 processes in a given population is needed. Also, balancing selection may be more common in natural
309 populations than earlier proclaimed (Chapman et al., 2019; Mérot et al., 2020), and thus we suggest
310 that it would make sense to incorporate balancing selection mechanisms into software tools designed
311 to evaluate population persistence.

312 To better understand how interacting genetic problems affect population persistence, empirical stud-
313 ies are needed. There are many technical challenges in obtaining the necessary detailed genetic infor-
314 mation, e.g., identifying which and how many loci are responsible for specific fitness traits, and which
315 genetic mechanism(s) maintain variation at which loci. However, current studies are already heading
316 in the right direction. For example, laboratory experiments with *Drosophila melanogaster* can now
317 distinguish the contributions of different genetic mechanisms in maintaining variation at a given
318 locus (Charlesworth, 2015; Sharp & Agrawal, 2018). In addition, modern sequencing techniques
319 can allow quantitative trait loci to be mapped to different fitness components and genomic regions
320 with traits that influence fitness components (such as survival and fecundity) can be identified from
321 neutral quantitative trait loci (Ågren et al., 2013).

322 To conclude, our results revealed that if the number of loci under selection is fixed, a mixture of
323 genetic processes is not necessarily worse than having the same genetic mechanism acting at all
324 loci. However, if the total number of loci under selection increases, for example because genetic
325 mechanisms that were previously neglected are now included, estimated extinction risk and MVP
326 sizes can dramatically increase. These observations call for inclusion of all genetic processes in
327 assessing the viability of populations. Populations normally also face other ecological problems
328 such as environmental stochasticity and Allee effects, and their consideration is expected to further

increase extinction risk and MVP size estimates (Tanaka, 2000; Wittmann et al., 2018).

Supporting Information

In the supplementary material, we briefly show the mathematical derivation for FS (Appendix S1). In Appendix S2, we provide more details about DAA and MCA. Appendices S3 through S7 provide some additional results that also involve FD. Also, we show how estimates of minimum viable population size depend on fitness per locus (Appendix S6) and how equilibrium fitness and critical number of loci can be estimated (Appendix S4).

References

- Agrawal, A.F. & Whitlock, M.C. (2012) Mutation load: the fitness of individuals in populations where deleterious alleles are abundant. *Annual Review of Ecology, Evolution, and Systematics* 43, 115–135.
- Ågren, J., Oakley, C.G., McKay, J.K., Lovell, J.T. & Schemske, D.W. (2013) Genetic mapping of adaptation reveals fitness tradeoffs in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences* 110, 21077–21082.
- Bensch, S., Andrén, H., Hansson, B., Pedersen, H.C., Sand, H., Sejberg, D., Wabakken, P., Åkesson, M. & Liberg, O. (2006) Selection for heterozygosity gives hope to a wild population of inbred wolves. *PloS One* 1, e72.
- Bergland, A.O., Behrman, E.L., O'Brien, K.R., Schmidt, P.S. & Petrov, D.A. (2014) Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLoS Genetics* 10, e1004775.
- Blomqvist, D., Pauliny, A., Larsson, M. & Flodin, L.Å. (2010) Trapped in the extinction vortex? Strong genetic effects in a declining vertebrate population. *BMC Evolutionary Biology* 10, 33.
- Chapman, J.R., Hill, T. & Unckless, R.L. (2019) Balancing selection drives the maintenance of

- 352 genetic variation in *Drosophila* antimicrobial peptides. *Genome biology and evolution* 11, 2691–
353 2701.
- 354 Charlesworth, B. (2015) Causes of natural variation in fitness: evidence from studies of *Drosophila*
355 populations. *Proceedings of the National Academy of Sciences* 112, 1662–1669.
- 356 Chen, J., Nolte, V. & Schlötterer, C. (2015) Temperature stress mediates decanalization and domi-
357 nance of gene expression in *Drosophila melanogaster*. *PLoS Genetics* 11, e1004883.
- 358 Coron, C., Méléard, S., Porcher, E. & Robert, A. (2013) Quantifying the mutational meltdown in
359 diploid populations. *The American Naturalist* 181, 623–636.
- 360 Durrett, R. (2008) *Probability models for DNA sequence evolution*. Springer-Verlag, Ithaca, New
361 York.
- 362 Estes, S., Phillips, P.C., Denver, D.R., Thomas, W.K. & Lynch, M. (2004) Mutation accumula-
363 tion in populations of varying size: the distribution of mutational effects for fitness correlates in
364 *Caenorhabditis elegans*. *Genetics* 166, 1269–1279.
- 365 Felsenstein, J. (2015) Theoretical evolutionary genetics, genome 562. *University of Washington,*
366 *Seattle, Washington* pp. 135–155.
- 367 Frankham, R. (1995) Effective population size/adult population size ratios in wildlife: a review.
368 *Genetics Research* 66, 95–107.
- 369 Frankham, R. (2005) Genetics and extinction. *Biological Conservation* 126, 131–140.
- 370 Frankham, R., Bradshaw, C.J. & Brook, B.W. (2014) Genetics in conservation management: revised
371 recommendations for the 50/500 rules, red list criteria and population viability analyses. *Biological*
372 *Conservation* 170, 56–63.
- 373 Franklin, I.R. (1980) Evolutionary changes in small populations. In: *Michael E. Soulé and B. M.*
374 *Wilcox, Eds., Conservation Biology: An Evolutionary-Ecological Perspective* pp. 135–149.
- 375 Glémin, S., Gaude, T., Guillemin, M.L., Lourmas, M., Olivieri, I. & Mignot, A. (2005) Balancing
376 selection in the wild: testing population genetics theory of self-incompatibility in the rare species
377 *Brassica insularis*. *Genetics* 171, 279–289.

- 378 Grossen, C., Guillaume, F., Keller, L.F. & Croll, D. (2020) Purging of highly deleterious mutations
379 through severe bottlenecks in Alpine ibex. *Nature communications* 11, 1–12.
- 380 Haldane, J. (1937) The effect of variation of fitness. *The American Naturalist* 71, 337–349.
- 381 Harcourt, A.H. (2002) Empirical estimates of minimum viable population sizes for primates: tens
382 to tens of thousands? *Animal Conservation* 5, 237–244.
- 383 Hedrick, P.W. (2012) What is the evidence for heterozygote advantage selection? *Trends in Ecology*
384 *& Evolution* 27, 698–704.
- 385 Hensen, I. & Wesche, K. (2006) Relationships between population size, genetic diversity and fitness
386 components in the rare plant *Dictamnus albus* in Central Germany. *Biodiversity and Conservation*
387 15, 2249–2261.
- 388 Higgins, K. & Lynch, M. (2001) Metapopulation extinction caused by mutation accumulation. *Pro-*
389 *ceedings of the National Academy of Sciences* 98, 2928–2933.
- 390 Hughes, A.L. & Nei, M. (1988) Pattern of nucleotide substitution at major histocompatibility com-
391 plex class I loci reveals overdominant selection. *Nature* 335, 167–170.
- 392 Kardos, M. & Luikart, G. (2021) The genetic architecture of fitness drives population viability during
393 rapid environmental change. *The American Naturalist* 197, 511–525.
- 394 Kimura, M. et al. (1955) Stochastic processes and distribution of gene frequencies under natural
395 selection. *Cold Spring Harbor Symposia on Quantitative Biology* 20, 33–53.
- 396 Kyriazis, C.C., Wayne, R.K. & Lohmueller, K.E. (2021) Strongly deleterious mutations are a primary
397 determinant of extinction risk due to inbreeding depression. *Evolution Letters* 5, 33–47.
- 398 Lacy, R.C. (1993) VORTEX - a computer simulation model for population viability analysis. *Wildlife*
399 *Research* 20, 45–65.
- 400 Leimu, R., Mutikainen, P., Koricheva, J. & Fischer, M. (2006) How general are positive relationships
401 between plant population size, fitness and genetic variation? *Journal of Ecology* 94, 942–952.

- 402 Lynch, M., Conery, J. & Burger, R. (1995) Mutation accumulation and the extinction of small
403 populations. *The American Naturalist* 146, 489–518.
- 404 Mérot, C., Llaurens, V., Normandeau, E., Bernatchez, L. & Wellenreuther, M. (2020) Balancing
405 selection via life-history trade-offs maintains an inversion polymorphism in a seaweed fly. *Nature*
406 *communications* 11, 1–11.
- 407 Nabutanyi, P. & Wittmann, M.J. (2021) Models for eco-evolutionary extinction vortices under bal-
408 ancing selection. *The American Naturalist* 197, 336–350.
- 409 Palomares, F., Godoy, J.A., López-Bao, J.V., Rodríguez, A., Roques, S., Casas-Marce, M., Revilla,
410 E. & Delibes, M. (2012) Possible extinction vortex for a population of Iberian lynx on the verge
411 of extirpation. *Conservation Biology* 26, 689–697.
- 412 Reed, D.H. (2005) Relationship between population size and fitness. *Conservation biology* 19, 563–
413 568.
- 414 Reed, D.H. & Bryant, E.H. (2000) Experimental tests of minimum viable population size. *Animal*
415 *Conservation* 3, 7–14.
- 416 Reed, D.H. & Frankham, R. (2003) Correlation between fitness and genetic diversity. *Conservation*
417 *Biology* 17, 230–237.
- 418 Reed, D.H., O’Grady, J.J., Brook, B.W., Ballou, J.D. & Frankham, R. (2003) Estimates of min-
419 imum viable population sizes for vertebrates and factors influencing those estimates. *Biological*
420 *Conservation* 113, 23–34.
- 421 Shaffer, M.L. (1981) Minimum population sizes for species conservation. *BioScience* 31, 131–134.
- 422 Sharp, N.P. & Agrawal, A.F. (2018) An experimental test of the mutation-selection balance model
423 for the maintenance of genetic variance in fitness components. *Proceedings of the Royal Society B*
424 285, 20181864.
- 425 Spielman, D., Brook, B.W., Briscoe, D.A. & Frankham, R. (2004a) Does inbreeding and loss of
426 genetic diversity decrease disease resistance? *Conservation Genetics* 5, 439–448.

- 427 Spielman, D., Brook, B.W. & Frankham, R. (2004b) Most species are not driven to extinction before
428 genetic factors impact them. *Proceedings of the National Academy of Sciences* 101, 15261–15264.
- 429 Tanaka, Y. (2000) Extinction of populations by inbreeding depression under stochastic environments.
430 *Population Ecology* 42, 55–62.
- 431 Teixeira, J.C. & Huber, C.D. (2021) The inflated significance of neutral genetic diversity in conser-
432 vation genetics. *Proceedings of the National Academy of Sciences* 118, e2015096118.
- 433 Traill, L.W., Bradshaw, C.J. & Brook, B.W. (2007) Minimum viable population size: a meta-analysis
434 of 30 years of published estimates. *Biological conservation* 139, 159–166.
- 435 Wang, T., Fujiwara, M., Gao, X. & Liu, H. (2019) Minimum viable population size and population
436 growth rate of freshwater fishes and their relationships with life history traits. *Scientific Reports*
437 9, 3612.
- 438 Wittmann, M.J., Stuis, H. & Metzler, D. (2018) Genetic Allee effects and their interaction with
439 ecological Allee effects. *Journal of Animal Ecology* 87, 11–23.
- 440 Xu, S., Chen, M., Liu, C., Zhang, R. & Yue, X. (2019) Behavior of different numerical schemes for
441 random genetic drift. *BIT Numerical Mathematics* pp. 1–25.
- 442 Yates, M., Bowles, E. & Fraser, D. (2019) Small population size and low genomic diversity have no
443 effect on fitness in experimental translocations of a wild fish. *Proceedings of the Royal Society B*
444 286, 20191989.
- 445 Zhao, L., Yue, X. & Waxman, D. (2013) Complete numerical solution of the diffusion equation of
446 random genetic drift. *Genetics* 194, 973–985.

1 Appendices

1 S1 Brief Description of Fluctuating Selection with Reversal 2 of Dominance (FS)

3 Briefly, FS was derived as follows (see Nabutanyi & Wittmann (2021) for a detailed description).
4 The relative fitnesses of the genotypes were $w_{AA,t} = (1 - s_{A,t})/\mathcal{C}$, $w_{aa,t} = (1 - s_{a,t})/\mathcal{C}$ and $w_{Aa,t} =$
5 $(1 - h_t \cdot s_{A,t} - (1 - h_t) \cdot s_{a,t})/\mathcal{C}$, where $s_{A,t} = s_A \cdot \sin(2\pi \cdot t/\kappa)$ and $s_{a,t} = s_a \cdot \sin(\pi + 2\pi \cdot t/\kappa)$
6 are temporally fluctuating selection coefficients and $h_t = 0.5 - c \cdot \sin(2\pi \cdot t/\kappa)$ is the temporally
7 fluctuating dominance coefficient. Here, $\mathcal{C} = 1 + \max(s_a, s_A)$ is the normalization constant such
8 that the genotype fitness lies between 0 and 1, s_A and s_a determine the amplitude of selection
9 and $0 \leq c \leq 0.5$ determines the strength of dominance. We arbitrarily used $c = 0.4$ and $\kappa = 50$
10 throughout the paper, where κ is the number of generations in a complete cycle.

11 S2 Diffusion Approximation and Markov Chain Approaches

12 S2.1 Diffusion Approximation Approach (DAA)

13 Here, $f(x, t)$ at a given locus was approximated using the diffusion equation (e.g in Kimura et al.
14 (1955))

$$\frac{\partial f(x, t)}{\partial t} = -\frac{\partial [a(x, t)f(x, t)]}{\partial x} + \frac{1}{2} \frac{\partial^2 [b(x, t)f(x, t)]}{\partial x^2}, \quad (\text{S2.1})$$

15 where $a(x, t)$ and $b(x, t)$ refer to the mean and variance of the change in allele frequency during an
16 infinitesimal time interval. For all the mechanisms, we took

$$b(x, t) = \frac{1}{2N_t} x(1 - x) \quad (\text{S2.2})$$

(e.g see Durrett (2008)) while $a(x, t)$ depended on the mechanism. For example, assuming s_a and s_A are very small that higher order terms involving s_i^3 and above are negligible, for HA, ID and MA,

$$a(x, t) = x(1 - x)(s_a(1 - x + 2hx) - hs_a - s_Ax)(1 + s_Ax^2 - s_a(1 - x)(x - 2hx - 1)) + \mu_a(1 - x) - \mu_Ax. \quad (\text{S2.3})$$

The three mechanisms differed in h , s_a and s_A . For FS (see Nabutanyi & Wittmann (2021) supplementary Section A.3),

$$a(x, t) = x(1 - x)(x(1 - 2h_t) + h_t)(1 + s_{a,t})(s_a - s_{A,t}) + \mu_a(1 - x) - \mu_Ax \quad (\text{S2.4})$$

and for FD,

$$a(x, t) = x(1 - x)(s_a(1 - x) - s_Ax)(1 + s_Ax^2 + s_a(1 - x)^2)/2 + \mu_a(1 - x) - \mu_Ax, \quad (\text{S2.5})$$

where, h_t , h , μ_a , μ_A , s_A , s_a , $s_{a,t}$, and $s_{A,t}$ are as defined under "Selection Mechanisms" subsection.

We adopted the numerical scheme in Zhao et al. (2013); Xu et al. (2019) to solve Equation (1) subject to the initial condition $f(x, 0) = x_0$ and the boundary condition that conserves probability at all times. The scheme basically discretizes both time t and allele frequency x over the interval 0 and 1 inclusive. The step-length for x is $\frac{1}{m}$, where m is an integer above 0 (used $m = 20$ for all the results) and step-length for time being $\tau > 0$ (in the whole paper, we used $\tau = 1$ so as to correspond to a full generation). The numerical scheme eventually gives a vector of probabilities $f(x_i, t)$ with $i \in \{0, 1, \dots, m\}$ for each time point as the distribution $f(x, t)$.

S2.2 Markov Chain Approach (MCA)

This approach involved finding the transition probabilities from i copies of a given allele at time t to j copies of the same allele at $t + 1$. Let X be a random variable representing the number of copies of allele A in a population of size N_t at time t . In a diploid population with two alleles at a given locus, there are $2N_t$ alleles in total, and X can take on $\{0, 1, 2, \dots, 2N_t\}$. The allele copies in the next generation $t + 1$ were randomly sampled (with replacement) from those present

at the current generation t . Therefore, X follows a binomial distribution with parameters $2N_t$ and x'' (the allele frequency after selection and mutation acting). Due to computational failure of the binomial distribution for large $N_t > 500$, we approximated the binomial distribution with one of two distributions, the Poisson distribution with parameter $2x''N_{t+1}$ (or $2(1-x'')N_{t+1}$ when one of the alleles is rare (expected copy number < 35) (Fig. S2.1), and otherwise the normal distribution with parameters $2x''N_{t+1}$ and $2x''(1-x'')N_{t+1}$ as mean and variance respectively. In summary, the probability of having j allele copies at time $t+1$ given that i copies are present at time t yields a transition matrix $\mathbf{P}_{ij}$,

$$\mathbf{P}_{ij} = \begin{cases} \text{Bin}(j; 2N_{t+1}, x''), & \text{if } N_{t+1} \leq 500, \\ \text{Poi}(j; 2x''N_{t+1}), & \text{if } N_{t+1} > 500 \ \& \ 2N_{t+1}x'' < 35, \\ \text{Poi}(2N - j; 2(1-x'')N_{t+1}), & \text{if } N_{t+1} > 500 \ \& \ 2N_{t+1}(1-x'') < 35, \\ \mathcal{N}(j; 2N_{t+1}, 2x''(1-x'')N_{t+1}), & \text{elsewhere.} \end{cases} \quad (\text{S2.6})$$

where $\text{Bin}(\dots)$, $\text{Poi}(\dots)$ and $\mathcal{N}(\dots)$ are the probability mass functions of the binomial, Poisson and normal distributions respectively. Here, $i = \{0, 1, \dots, 2N_t\}$, $j = \{0, 1, \dots, 2N_{t+1}\}$ and x'' is the frequency of allele A at the end of a given generation t after possible selection and mutation, i.e., $x = \frac{i}{2N_t}$, $x' = \frac{x^2 w_{AA} + x(1-x)w_{Aa}}{x^2 w_{AA} + 2x(1-x)w_{Aa} + (1-x)^2 w_{aa}}$ and $x'' = (1 - \mu_A)x' + \mu_a(1 - x')$. We then normalise each row such that the transition probabilities always sum to 1.

Assuming a population at equilibrium with a constant size N , $\mathbf{P}_{ij}$ is a $(2N+1) \times (2N+1)$ square matrix. The equilibrium distribution of allele copies was obtained by getting the eigenvector corresponding to the largest real part of the eigenvalues (which is always equal to 1) of matrix $\mathbf{P}_{ij}$, which was then normalised to get $f(x, t)$.

S2.2.1 Distribution of Allele Frequencies via Iterating through Generations for the Markov Chain Approach

Similar to the diffusion approximation, the MCA can also be made time-dependent by starting with some initial allele frequency distribution and evaluating the distribution in every generation (Section S2.2.1). Here, the transition matrix is not necessarily a square matrix and allows varying population

58 sizes between generations.

59 Here, matrix $\mathbf{P}_t$ (subscript t due to change with time) is calculated every generation and is not
60 necessarily a square matrix. $\mathbf{P}$ is generally a $2N_t + 1 \times 2N_{t+1} + 1$ matrix filled according to Equation
61 (S2.6). At each generation $t \geq 0$, the (row vector) distribution of allele copies X is determined as

$$X_{t+1} = X_t \mathbf{P}_t, \quad (\text{S2.7})$$

62 where we start with some initial allele frequency distribution X_0 . The distribution is used in a
63 similar way as in the diffusion approximation to determine population size (Equation (5)). However,
64 this approach is computationally more expensive than the DAA and equilibrium-based MCA.

Figure S2.1: Densities for the distribution of allele frequencies with different values of probability, x . The upper panel is obtained from the binomial ($n = 2N, p = x$), Poisson ($\lambda = 2Nx$), and normal ($\mu = 2Nx, \sigma^2 = 2Nx(1 - x)$) distributions, while for lower panel, we subtract the random draws (using same parameters) from $2N$; where $N = 500$ and number of draws = 500000. The green vertical dotted line is at either 0 or $2N$ to indicate cases where the normal distribution yield non-feasible observations (allele copies below 0 and above $2N$). From these plots, we consider less than 35 allele copies to be rare enough to use the Poisson approximation (as this threshold yields a good fit to the binomial and avoids negative samples or samples above $2N$ at extreme allele frequencies.)

S3 Population trajectories

S3.1 Population Trajectories from DAA

Figure S3.1: Isolated selection mechanisms: Trajectories for various carrying capacities from the diffusion approximation approach. Row 1: population size and Row 2: absolute fitness. The horizontal gray line in Row 2 corresponds to the threshold absolute fitness of 1. Parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a = 0.0005$, $\kappa = 50$ and for MA $s_A = 0$, $s_a = 0.005$, $h = 0.5$, $\mu_A = 0.0005$, $\mu_a = \frac{\mu_A}{100}$. Other shared parameters were $n = 100$, $K = N_0$, $r = \ln(1.53)$.

Figure S3.2: Interacting mechanisms: Trajectories for different carrying capacities from the diffusion approximation for Row 1: HA-MA and Row 2: FS-ID. Column 1: population size, Column 2: absolute fitness, Column 3: contribution of constituent selection mechanisms to offspring viability. The gray horizontal line in Column 2 is the threshold absolute fitness of 1. Under balancing selection: $s_a = s_A = 0.005, \mu_A = \mu_a = 0.0005, \kappa = 50$ and for ID and MA $s_A = 0, \mu_A = 0.0005, \mu_a = \frac{\mu_A}{100}$ with $h = 0.5, s_a = 0.005$ for MA and $h = 0.01, s_a = 0.05$ for ID. Shared parameters: $n = 100, K = N_0, r = \ln(1.53)$. The number of loci were equally shared between constituent mechanisms.

67 S3.1.1 Longer-time trajectories

Figure S3.3: Longer-time population and fitness trajectories (run for 40000 generations) for various carrying capacities using the diffusion approximation approach. The gray horizontal line is the threshold absolute fitness of 1. The red line shows the MVP size estimate. In cases of more than one selection mechanism, the number of loci under each mechanism was equal. Parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a = 0.0005$, $\kappa = 50$ and for ID and MA $s_A = 0$, $\mu_A = 0.0005$, $\mu_a = \frac{\mu_A}{100}$ with $h = 0.5$, $s_a = 0.005$ for MA and $h = 0.01$, $s_a = 0.05$ for ID. Other shared parameters were $n = 100$, $K = N_0$, $r = \ln(1.53)$.

S3.2 Population Trajectories from the IBM

Figure S3.4: Individual-based simulations of 60 trajectories (in gray) for HA at $K = 170$ (column 1) and HA-FS-FD-ID-MA at $K = 60$ (column 2). The thick lines are for various carrying capacities using the diffusion approximation approach. Parameters for balancing selection are $s_a = s_A = 0.005$, $\mu_A = \mu_a = 0.00022$, $\kappa = 50$ and for ID and MA $s_A = 0$, $s_a = 0.005$, $\mu_A = 0.00022$, $\mu_a = \frac{\mu_A}{100}$ with $h = 0.5$ for MA and $h = 0.0$ for ID. Other shared parameters were $n = 100$, $K = N_0$, $r = \ln(1.53)$. In column 2, the number of loci under each mechanism was equal (i.e., 20 loci each).

69 S4 Critical Number of Loci and Equilibrium Fitness

70 S4.1 Equilibrium Fitness

71 We start with single-locus dynamics where the mean fitness is obtained from the deterministic
72 equilibrium allele frequency x (which would be reached in a population of infinite size) as

$$\bar{V}(x) = x^2 w_{AA} + 2x(1-x)w_{Aa} + (1-x)w_{aa}, \quad (\text{S4.1})$$

73 where w_{ij} , $i, j \in \{A, a\}$ is the equilibrium genotype fitness (geometric mean fitness in a complete
74 cycle for FS) and $\bar{V}$ is the mean viability fitness for a given mechanism. Assuming equilibrium
75 conditions and for symmetric selection, the equilibrium mean fitness for HA is given by

$$\bar{V} = 1 - \frac{s}{2}, \quad (\text{S4.2})$$

76 where s is the selection coefficient against homozygotes.

77 For a purely recessive deleterious allele, the fitness is given by (assuming Hardy-Weinberg proportions
78 in an infinite population; Felsenstein (2015))

$$\bar{V} = 1 - (1 - \bar{x})s_a, \quad (\text{S4.3})$$

79 where s_a the selection coefficients against the homozygote of the recessive allele a and $\bar{x}$ is the
80 frequency of wild-type allele A approximated by

$$\bar{x} = 1 - \sqrt{\frac{\mu_A}{s_a}}. \quad (\text{S4.4})$$

81 However, when the deleterious allele is not completely recessive,

$$\bar{V} = 1 - 2\bar{x}(1 - \bar{x})hs_a - (1 - \bar{x})^2s_a, \quad (\text{S4.5})$$

82 where $h > 0$ is the dominance coefficient and this time $\bar{x}$ is approximated by

$$\bar{x} = \frac{\mu_A}{hs_a}. \quad (\text{S4.6})$$

83 Note that when $s_a < \mu_A$ in Equation S4.4 and $hs_a < \mu_A$ in Equation S4.6 gives $\bar{V} > 1$ which does
 84 not make sense. So, the approximations fail for these cases because the equilibrium allele frequency
 85 of the wild-type allele will be below 0. Ideally, in such cases, selection against deleterious allele
 86 is so weak and yet their inflow via mutation is so high and thus it will result into fixation of the
 87 deleterious allele.

88 In Fig. S4.1, we show effect of increasing selection coefficient and mutation rates on MVP estimates
 89 for loci under mutation-selection-drift equilibrium. At a given mutation rate, as s_a (or hs_a) increases,
 90 the MVP decreases rapidly. Also, for a given selection coefficient, increase in mutation increases
 91 MVP size estimates. These observations are expected as shown from Equations (S4.4) and (S4.6).
 92 The increase in s_a reduces the frequency of the deleterious allele. Thus strong purging of deleterious
 93 alleles eventually lead to a higher equilibrium fitness. On the other hand, increase in μ_A increases
 94 frequency of the deleterious mutation at equilibrium leading to low equilibrium fitness.

Figure S4.1: MVP estimates for populations under different selection pressure from inbreeding depression with fixed number of loci $n = 100$. On the x-axis, we have the forward mutation rate μ_A . Other parameters are $\mu_a = \frac{m_A}{100}$, $K = N_0$, $r = \ln(1.53)$.

95 S4.2 Critical number of loci

96 It is possible to have the critical number of loci n_c , above which no arbitrary size of a population
 97 can survive. Here, n_c is obtained based on the assumption that absolute fitness of 1 is the threshold

below which populations go extinct i.e., $\bar{V}^n E < 1$, n is the number of loci under selection,

$$n_c = \frac{-\ln(E)}{\ln(\bar{V})}, \quad (\text{S4.7})$$

while for two mechanisms, $\bar{V}_1^{n_1} \bar{V}_0^{n_0} E < 1$, n_i being number of loci under respective selection mechanism.

$$n_{1c} = \frac{\ln(\frac{1}{(\bar{V}_0^{n_0} \times E)})}{\ln \bar{V}_1} = -\frac{n_0 \ln(\bar{V}_0) + \ln(E)}{\ln \bar{V}_1}, \quad (\text{S4.8})$$

where n_0 is the number of loci originally occupied by a genetic mechanism whose equilibrium fitness per locus is $\bar{V}_0$ and n_{1c} is the critical number of loci added from the new genetic mechanism whose mean fitness per locus is $\bar{V}_1$, both evaluated at the respective equilibrium frequencies $\bar{x}_0$ and $\bar{x}_1$ while E is the expected number of offspring per individual. So, n_{1c} gives the maximum additional loci that can be acted on to have a finite population size surviving. We illustrate how n_c varies with selection coefficient, fitness and mutation rate in Figure S4.2

Figure S4.2: (a) How absolute fitness changes with increasing number of loci under selection (assuming infinite population size). Parameters used $s_a = s_A = 0.005$ for HA and FS, $s_a = 0.05$, $s_A = 0$, $\mu = 0.001$, $h = 0$ for ID (b) Shows how critical number of loci (that results into an asymptotic increase in MVPs) varies with selection coefficients in HA (from Equation (S4.7)). (c) Shows how the critical number of loci under selection varies with mutation rate μ_A for different selection coefficients for MA. Parameters used $h = 0$, $\bar{x} = \sqrt{\mu/s_a}$ for the deleterious allele.

107 S4.3 Effect of Number of Loci on MVP estimates from the MCA

Figure S4.3: Effect of increasing number of loci under selection on MVP size. Same data as in 4 but the y-axis not log-scaled but limited to a maximum of 10000. The genome from populations with at most 100 loci (below the vertical gray line) was acted on by the same selection mechanism (indicated at the start of the line) (a) HA, (b) FS, (c) ID and (d) MA. Those genomes with above 100 loci, the mechanism indicated at the end of the line acted on the remaining loci, e.g., the line with HA—ID label implies the first 100 loci were acted on by HA while ID acted on the remaining loci. Parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a = 0.0005$, for ID and MA, $s_A = 0$, $\mu_a = \frac{\mu_A}{100}$ while $h = 0.01$, $s_a = 0.05$ for ID and $h = 0.5$, $s_a = 0.005$ for MA. Shared parameters were $K = N_0$, $r = \ln(1.53)$.

Figure S4.4: Effect of mutation rates on MVP sizes and critical number of polymorphic loci above which no arbitrary finite population survives. The results used the diffusion approximation approach (DAA). Mutation rates only affects non-balancing selection mechanisms. Genomes with less than 100 loci (separated by the vertical gray line) are acted on by the same selection mechanism (a) HA, (b) FS, (c) ID and (d) MA. Those with above 100 loci, the mechanism indicated at the end of the line acts on the remaining loci, e.g., the line with HA—ID label implies the first 100 loci were acted on by HA while ID acts on the possible available loci. Here, $\mu = 0.0001$ for unbroken lines and $\mu = 0.0005$ for broken lines. Other parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a$, $\kappa = 50$, for ID and MA, $s_A = 0$, $\mu_a = \frac{\mu_A}{100}$ while $h = 0.01$, $s_a = 0.05$ for ID and $h = 0.5$, $s_a = 0.005$ for MA. Shared parameters were $K = N_0$, $r = \ln(1.53)$.

Figure S4.5: Effect of increasing number of loci under selection on MVP size. Everything is as in Fig. S4.4 but this is shown for the Markov chain mechanism.

S5 MVP Estimates from Fixed Number of Loci

S5.1 MVP Estimates from IBM with Fixed Number of Loci

Figure S5.1: MVP estimates from the IBM. Row 1: HA only, Row 2: ID only, and Row 3: HA-FS-FD-ID-MA. 1st column: MVP estimates for different mutation rates measured at different levels of survival probability. 2nd column: Mean survival time of populations at different carrying capacities estimated under different mutation rates. 3rd column: Coefficient of variation in survival times of populations at different carrying capacities estimated under different mutation rates. Parameters for balancing selection; $s_a = s_A = 0.005$, $\mu_A = \mu_a$, $\kappa = 50$, for ID and MA $s_A = 0$, $s_a = 0.005$, $\mu_a = \frac{\mu_A}{100}$ while $h = 0.5$ for MA and $h = 0$ for ID. Other shared parameters were $K = N_0$, $r = \ln(1.53)$ and a total of 100 loci were equally shared by the constituent mechanisms.

Figure S5.1 shows MVP size estimates from the IBM. The MVP size estimates from the IBM were qualitatively similar to those estimated by DAA and MCA (Fig. 5 in the main text). Quantitatively, estimates from the IBM were (slightly) higher than corresponding estimates (Fig. 5 row 2). The MVP estimates were fairly insensitive to levels of survival probability threshold (Fig. S5.1 column 1). As expected, the mean time to extinction increased rapidly with increasing carrying capacity K (Fig. S5.1 column 2) Also, there was more uncertainty in the extinction times in populations with lower carrying capacities than their counterparts with large capacities (Fig. S5.1 column 3).

S5.2 MVP Estimates with Fixed Number of Loci Including those from Negative Frequency-Dependent Selection (FD)

Figure S5.2: MVP estimates from the Markov chain approach (with broken thick lines) and the diffusion approximation (with unbroken thin lines). Parameters $s_a = s_A = 0.005$, $K = N_0$, $r = \ln(1.53)$, $\mu_A = \mu_a$, $\kappa = 50$ except for ID and MA where $s_A = 0$, $\mu_a = \frac{\mu_A}{100}$ and also for ID, $h = 0.5$. The total number of loci on the genome is $n = 100$ except for genomes with 3 selection mechanisms where $n = 99$ for purposes of equal distribution of loci.

S5.3 Arithmetic, Geometric and Harmonic Mean of the MVPs from Constituent Selection Mechanisms

Figure S5.3: Comparison of the estimated MVPs (on x axis) to the geometric, harmonic and arithmetic means (on y axis) of the constituent isolated selection mechanisms for a given scenario using (a) the diffusion approximation and (b) the Markov chain approaches. The different colours show different calculated means (black, red, and green points for geometric, harmonic, and arithmetic means respectively) while the solid lines are the regression lines for each calculated mean. Parameters were as in Fig. S5.2. The points are from HAFS, HAID, FSMA, MAID, HAFSMA, HAFSID, FSMAID, and HAFSMAID.

In Fig. S5.3, we compare the joint MVP to the different means of the MVPs estimated from the isolated selection mechanisms. Generally, the observed MVPs under a mixture of mechanisms are closer to the harmonic and geometric means than the arithmetic mean of the MVPs calculated for the mechanisms in isolation with the same total number of loci. To a large extent, the joint MVPs are (slightly) above the calculated harmonic means and (slightly) below the geometric means. We briefly illustrate why joint MVP is closer to the harmonic mean in Section S6.

127 **S6 Relationship between MVP estimates and fitness across** 128 **loci**

Figure S6.1: Viability fitness per locus at different population sizes for the different mechanisms from the Markov chain approach at equilibrium (Equation (S2.6)). Parameters: $s_a = s_A = 0.005$, $\mu_A = \mu_a$, $\kappa = 50$ for balancing selection mechanisms and for MA: $s_A = 0$, $s_a = 0.005$, $h = 0.5$, $\mu_a = \frac{\mu_A}{100}$, and ID: $s_A = 0$, $s_a = 0.05$, $h = 0.01$, $\mu_a = \frac{\mu_A}{100}$. Panel (b) zooms in on the upper viability range in (a).

Figure S6.2: Logistic curve fitting $V = V_{min} + \frac{V_{max} - V_{min}}{1 + e^{-m(N - N_m)}}$, where V is the viability per locus, V_{min} and V_{max} are the minimum and maximum viability per locus, and N_m is the population size N of the viability sigmoid midpoint, and m is steepness of the curve. Illustration of the relationship between minimum viable population sizes in a logistic curve setting (as observed from Figure S6.1). To get viability for a single mechanism with $n = 100$ loci, we simply use V^n (blue and red solid lines), and for two mechanisms, we use $V_1^{n/2} \times V_2^{n/2}$ (magenta solid line). The corresponding minimum viable population sizes with a critical viability $v_{crit} = 1/E = 1/1.53$ are shown as dashed vertical lines. The circle, diamond, and triangle indicate the harmonic, geometric, and arithmetic mean of the two separate minimum viable population sizes, respectively.

Figure S6.3: Illustration of the relationship between minimum viable population sizes in a simple linear setting. Assuming approximately additive effects of small fitness reductions, the viability with 50% of loci under mechanism 1 and 50% under mechanism 2 for a given population size (magenta solid line) is the arithmetic mean of viabilities for the two separate mechanisms acting at 100% of loci (blue and red solid lines). The corresponding minimum viable population sizes with a critical viability $v_{crit} = 1/E = 0.92$ are shown as dashed vertical lines. The circle, diamond, and triangle indicate the harmonic, geometric, and arithmetic mean of the two separate minimum viable population sizes, respectively.

Here we use a simple setting to illustrate possible relationships between MVP sizes under single versus multiple genetic mechanisms. For this section, we assume that the relationship between population size and viability at low population sizes can be well approximated by a linear relationship (Figures S6.1 and S6.2). Assume that we have two genetic mechanisms ($i = 1, 2$) that when acting at n loci have viability

$$V = v_{min,i} + m_i \cdot N. \quad (\text{S6.1})$$

The respective minimum viable population size then is the population size at which viability takes the critical value $v_{crit} = 1/E$ at which absolute fitness is 1. Thus,

$$v_{crit} = v_{min,i} + m_i \cdot MVP_i \quad (\text{S6.2})$$

136

$$\Leftrightarrow MVP = \frac{v_{crit} - v_{min,i}}{m_i}. \quad (\text{S6.3})$$

137 As long as deviations of viability from 1 are small, loci make roughly additive contributions to
138 reducing viability and thus the viability relationship with half of the loci under mechanisms 1 and
139 half of the loci under mechanism 2 is

$$V = \frac{v_{min,1} + v_{min,2}}{2} + \frac{m_1 + m_2}{2} \cdot N \quad (\text{S6.4})$$

140 and the corresponding minimum viable population size is

$$MVP_{joint} = \frac{v_{crit} - \frac{v_{min,1} + v_{min,2}}{2}}{\frac{m_1 + m_2}{2}} = \frac{C_1 + C_2}{\frac{C_1}{MVP_1} + \frac{C_2}{MVP_2}}, \quad (\text{S6.5})$$

141 where $C_i = v_{crit} - v_{min,i}$, $i = 1, 2$.

142 We now consider three different exemplary cases (Fig. S6.3). In the first case (Fig. S6.3 A), the two
143 mechanisms have the same minimum viability $v_{min,1} = v_{min,2} = v_{min}$ (i.e., $C_1 = C_2$) as was also
144 approximately the case in Fig. S6.1. Then

$$MVP_{joint} = \frac{v_{crit} - v_{min}}{\frac{m_1 + m_2}{2}} = \frac{1}{\frac{1}{2} \cdot \frac{m_1}{v_{crit} - v_{min}} + \frac{m_2}{v_{crit} - v_{min}}} = \frac{2}{\frac{1}{MVP_1} + \frac{1}{MVP_2}}. \quad (\text{S6.6})$$

145 That is, the joint minimum viable population size (magenta vertical line in Fig. S6.3 A) is equal
146 to the harmonic mean (indicated by a black dot in Fig. S6.3 A) of MVP sizes under mechanism 1
147 and 2. Intuitively, this is the case because minimum viable population size is inversely proportional
148 to the rate at which viability increases with increasing population size. This is approximately true
149 even when the viability functions are nonlinear and thus explains why we often obtain joint MVP
150 sizes close to the harmonic mean of individual MVP sizes.

151 However, this equality does not hold if the two mechanisms have different v_{min} (i.e., $C_1 \neq C_2$).
152 Depending on the relation of the two slopes and intercepts, we can then get scenarios where the
153 joint MVP is larger than the harmonic mean of MVPs but still smaller than the arithmetic mean
154 (Fig. S6.3 B) or scenarios where the joint MVP is even larger than the arithmetic mean of MVPs
155 (Fig. S6.3 C). Also for general nonlinear viability functions, all cases are possible and the exact
156 relationship between MVPs will depend on the precise shape of the curves.

157 **S7 Initial allele frequency**

158 Figure S7.1 shows how initial allele frequency depends affects MVP estimates at different mutation
 159 rates for different isolated mechanisms. Initial allele frequency has large effects in FS. For loci under
 160 mutation-selection-drift balance (ID and MA), MVPs are insensitive to initial allele frequency unless
 161 the frequency of the deleterious allele is extreme like above 90%. Similarly, for HA, only very extreme
 162 allele frequencies have an effect on MVPs.

Figure S7.1: Effect of initial allele frequency on MVP size estimates. The blue lines show MVP trajectories at given initial allele frequency. The parameters for balancing selection were $s_a = s_A = 0.005$, $\mu_A = \mu_a$, $\kappa = 50$, for ID and MA $s_A = 0$, $s_a = 0.005$, $\mu_a = \frac{\mu_A}{100}$ while $h = 0.5$ and $h = 0$ were respectively for MA and ID. Other shared parameters were $K = N_0$, $r = \ln(1.53)$. The gray shaded parts have their MVP estimates above 10000.