The gut bacteria of an invasive amphibian respond to the dual challenges of range-expansion and parasite attack

Jia Zhou^{1,6*}, Tiffanie Maree Nelson², Carlos Rodriguez Lopez³, Shao Jia Zhou^{1,4}, Georgia Ward-Fear⁵, Katarina C. Stuart⁶, Richard Shine⁵, Lee Ann Rollins⁶

¹School of Agriculture, Food and Wine, University of Adelaide, Waite Campus, PMB 1 Glen Osmond, South Australia, Australia, 5064.

²Queensland Facility for Advanced Bioinformatics, School of Medicine, Menzies Health Institute Queensland, Building G40, Level 9, Gold Coast Campus, Griffith University, Southport, QLD, Australia, 4215.

³Environmental Epigenetics and Genetics Group, Department of Horticulture, College of Agriculture, Food and Environment, University of Kentucky, Lexington, KY, USA, 40546.

⁴Robinson Research Institute, University of Adelaide, South Australia, Australia, 5005

⁵Department of Biological Sciences, Macquarie University, North Ryde, NSW, Australia, 2109.

⁶Ecology & Evolution Research Centre, School of Biological, Earth and Environmental Sciences, University of New South Wales, Biological Sciences South (E26) UNSW, Kensington, Australia, 2052

*** Correspondence:**

Jia Zhou

E-mail: jjazhou0116@gmail.com

Keywords: microbiome, Cane toad, 16S rRNA, host-parasite

23 Word count: 6334.

24

25 Abstract

26 Gut bacterial communities influence, and are influenced by, the behaviour and ecology of their hosts.
 27 Those interactions have been studied primarily in humans and model organisms, but we need field
 28 research to understand the relationship between an organism's gut bacteria and its ecological
 29 challenges, such as those imposed by rapid range expansion (as in invasive species) and the presence
 30 of host-manipulating parasites. Cane toads (*Rhinella marina*) provide an excellent model system in
 31 this respect, because the species' ongoing colonization of Australia has enforced major changes in
 32 phenotypic traits (including behaviour), and lungworm parasites (*Rhabdias pseudosphaerocephala*)
 33 modify host gut function in ways that enhance the viability of lungworm larvae. We collected female
 34 toads from across the species' invasive range and studied their morphology, behaviour, parasite
 35 infection status and gut bacterial community. Range-core *versus* range-edge toads differed in
 36 morphology, behaviour, gut bacterial composition and predicted gut bacterial function but did not
 37 differ in the occurrence of parasite infection nor in the intensity of infection. Toads infected with
 38 lungworms differed from uninfected conspecifics in gut bacterial composition and diversity. Our
 39 study demonstrates strong associations between gut bacterial community and host ecology and
 40 behaviour.

41

42 Introduction

43 The bacterial community within an organism's intestines can be strongly influenced by host
 44 behaviour and ecology, such as habitat selection and diet [1–5]. But that interaction runs both ways
 45 because gut bacteria can influence behaviour and ecology of the host. For example, transferring gut
 46 contents can modify the recipient host's behaviour (exploratory behaviour, *Mus musculus* [6];
 47 emotional reactivity, *Coturnix japonica* [7]). Similarly, altering gut microbial communities by
 48 administering antibiotics or altering dietary composition triggered aggressive behaviour in leaf-
 49 cutting ants (*Acromyrmex echinator* [8]). Remarkably, changes in only a single bacterial species
 50 within the gut can affect behaviour of the host (e.g., *Drosophila melanogaster* [9]; *Danio rerio* [10];
 51 *M. musculus* [11]). Gut microbiota can also affect mating choices [12] and foraging [13–15].

To date, most evidence for effects of intestinal bacteria on host behaviour comes from studies on humans and “model organisms”. To elucidate the functional significance of this phenomenon, we need to extend such studies to free-ranging animals, incorporating a wider range of taxa [16]. Invasive species offer good models for such research, because novel challenges in the invaded range create an opportunity to compare closely-related organisms exposed to profoundly different environments [17,18]. Host-parasite relationships also may be disrupted during biological invasions, due to processes such as “enemy release” (loss of co-evolved native pathogens from the native range [19]). Parasites can manipulate host behaviour and physiology in ways that enhance parasite fitness but reduce host fitness [20]. Thus, if the gut bacterial community provides a mechanism for such effects, parasite-infected individuals should exhibit different gut bacteria than uninfected conspecifics.

These ideas suggest two predictions: (i) that the gut bacterial community should differ between populations of an invasive species (e.g., between the range-core and the invasion-front); and (ii) that the gut bacterial community should differ between parasitized hosts and non-parasitized conspecifics. The colonization of Australia by cane toads (*Rhinella marina*) provides a robust opportunity to test these predictions. Since their release in north-eastern Australia in 1935, toads have dispersed into areas that are much hotter and more seasonally arid than in the native range or the initial release sites [21]; and toads have brought with them a native-range nematode lungworm (*Rhabdias pseudosphaerocephala*) that can have devastating impacts on host viability, and induces behavioural and physiological changes in the host [22]. Notably, infected toads produce copious watery faeces [22]; hence, we expect lungworm-infected toads to exhibit different gut bacterial communities than uninfected individuals.

Methodology

Study species, sample collection and behavioural assays

Cane toads are native to South America, and were introduced into Australia in 1935 as a biocontrol for pests of sugar cane crops [23]. As the toads spread through tropical Australia, they fatally poisoned many native predators [23]. Toads from range-core populations (eastern Australia) differ from invasion-front conspecifics (in north-western Australia) in phenotypic traits that confer increased dispersal ability, such as endurance [24], limb morphology [25], boldness and exploratory behaviour [17,26]. Toads from the invasion-front also have lower rates of infection of the co-

introduced parasitic lungworm [27]. Drivers of variation in invasion-related behaviours in this species include genetics, morphology, habitat, diet, prior experience and parasites [17,28–31]. However, the possible role of gut bacteria as a potential driver of behavioural shifts across the invasive range has not been studied.

We hand-captured 60 adult females from three sites on the invasion-front and three sites in the range-core (Table S1). We conducted brief behavioural assays upon collection including: (i) struggle score (number of kicks after being captured until toad remains still for 5 seconds) and struggle likelihood; and (ii) righting effort (time to right itself, number of kicks within two minutes after toad is placed on its dorsal side, and righting effort likelihood [17]. These measures are predictive of traits including speed and stamina (K. Stuart, pers. comm.), suggesting that these simple assays may reveal a toad's dispersal potential. We then placed animals into individual, moist, calico bags and weighed, measured (snout urostyle length; SUL) and euthanised them by injecting tricaine methanesulfonate (MS222) buffered with bicarbonate of soda.

We dissected the toads and scored the presence of two types of toad parasite: the gut-encysted physalopterine larvae [32] and adult lungworms [22,33]. Lungworm larvae pass through the gut, but are less easily detected and reliably counted than are adult lungworms. From each toad, we removed 0.3cm of colon near the cloaca (including gut contents) and preserved it in 95% ethanol (see Supplementary Material for justification of sampling protocols).

Analyses

We compared host morphology and behaviour between regions (range-core *versus* invasion-front), and as a function of infection status (lungworm infected *versus* non-infected). Because body length (SUL) and mass were correlated, we only included SUL as our measure of host morphology in further analyses. We used a t-test to compare mean SUL between regions and infected/non-infected toads. For associations between region or infection status with host characteristics or behavioural traits, we used SUL as a covariate in generalized linear models (GLM). See Supplementary Material for details of statistical analyses.

Laboratory methods and data pre-processing for characterizing gut bacterial community composition and predicted functions are described in Supplementary Material. Briefly, we calculated within-individual (alpha) bacterial diversity and between-site (beta) bacterial diversity. For the latter variable, we subset our data to include the Core50 gut community (Amplified Sequence Variants (ASVs) present in a minimum of 50% of toads from each site [2]). We predicted bacterial functions and generated pathway abundance based on Core50 ASVs. We compared bacterial composition and predicted function between regions, and between lungworm-infected and non-infected toads. We identified differences in individual ASVs and predicted bacterial functions between range-core and invasion-front toads and identified associations between host characteristics (including infection with parasites) with bacterial communities and predicted bacterial functions. Analyses were conducted in QIIME2 [34], PICRUST2 [35], and R packages in R version 4.0.2 [36].

Results

Ecological traits

Wild-caught invasion-front toads were larger than range-core toads (Tables S2, S3; mean SUL $t = 2.54$, $df = 53.90$, $p = 0.014$). Neither counts nor presence of parasites (lungworm and gut) differed significantly across the range (Table S2). Range-core toads were more likely to struggle (Tables S2, S3; $p = 0.008$, 95% CI: core [-0.174, 0.005], edge [-0.023, 0.199]) and, in those that did struggle, the number of struggle movements was higher for range-core toads (Tables S2, S3; $p = 0.002$, 95% CI: core [-0.057, 0.026], edge [0.046, 0.2]). Range-core toads also were more likely to attempt to right themselves ($p = 0.036$, 95% CI: core [-0.092, 0.04], edge [-0.006, 0.19]), but righting effort and righting time did not differ significantly between geographic regions (Tables S2, S3).

Because there were no significant differences in prevalence or intensity of lungworm between the range-core and invasion-front toads (Table S2), we combined samples to analyse correlates of lungworm infection. Infected toads were similar in SUL to non-infected toads (Tables S3, 4; $t = 0.86$, $df = 57.19$, $p = 0.393$), with no significant behavioural differences between the two groups (Table S3, 4).

Gut bacterial community composition and predicted bacterial function

Alpha diversity did not differ significantly between regions (Supplementary Material), but beta diversity of bacterial taxonomic communities differed between regions (Figures 1A, S1; $R^2 = 0.050$,

F = 3.050, $p < 0.001$) and sampling sites (Table S5; all p -values < 0.001). Among 230 ASVs that were assigned to family level, the abundance of 124 ASVs differed between the colons of range-core *versus* invasion-front toads (Table S6). The number of significantly different ASVs in each phylum were: Bacteroidetes (60 ASVs), Firmicutes (55 ASVs), Proteobacteria (7 ASVs), Actinobacteria (1 ASVs), Verrucomicrobia (1 ASV) (Table S6, Figure 2A).

Among the identified 474 predicted bacterial functions, we found significant differences between invasion-front and range-core toads (Figure 1B; $R^2 = 0.064$, $F = 4.110$, p -value = 0.002). Pairwise tests between sampling sites indicated that Kununurra toads had different bacterial functions to Rossville (p -value = 0.009) and Lucinda toads (p -value = 0.046), but no other sites differed functionally from each other (Table S7; all other p -values > 0.05). In total, 84 predicted bacterial functions differed between invasion-front and range-core toads (Table S8, Figure 2B). Range-core toads had more abundant bacterial function in the superpathway of pyrimidine ribonucleosides degradation ($\log_2\text{FoldChange} = 5.98$) and less abundant bacterial function in phosphopantothenate biosynthesis III ($\log_2\text{FoldChange} = -4.98$), superpathway of sialic acids and CMP-sialic acids biosynthesis ($\log_2\text{FoldChange} = -4.89$) and factor 420 biosynthesis ($\log_2\text{FoldChange} = -4.72$) than did invasion-front toads (Table S8, Figure 2B). Among the 30 most abundant functions, range-core toads had lower bacterial function in urate biosynthesis/inosine 5'-phosphate degradation ($\log_2\text{FoldChange} = -0.10$) than did invasion-front toads (Figure 3, Table S8).

Associations between ecological traits and intestinal bacteria

To assess correlates of gut bacterial composition and function, we compared characteristics of individual hosts to bacterial variation. Only the occurrence of lungworms was significantly associated with the bacterial composition ($R^2 = 0.128$, $p = 0.02$) (Table 1; Figure S2A, B).

In a redundancy analysis combining ecological traits measured here, the model that explained the most variation in the bacterial community assemblage included only the occurrence of lungworms (AIC = 178.58). The best model to explain variation in predicted bacterial functions included the likelihood of righting (AIC = 53.613), the occurrence of lungworms (AIC = 54.297) and righting time (AIC = 56.912). The combination of these three factors explained 17.8% of total variation in predicted bacterial functions (Figure 4).

In explicit tests of whether bacterial community assemblages differed in infected *versus* non-infected toads, we found a significant association with lungworm occurrence (Table 2, $p=0.005$). Intensity of lungworm infection was not significantly associated with gut bacterial community but did have a significant interaction with region in this analysis (Table 2, $p=0.04$).

Discussion

Bacteria influence animal behaviour in diverse ways [16,37], but the ecological drivers of variation in gut bacterial composition remain largely unstudied. Our analyses of cane toads from two regions within their invasive range documents substantial variation in community assemblage and function of gut bacteria. Importantly, that variation was associated with two traits that we predicted to influence gut bacterial assemblages: invasion history and parasite infection. Interestingly, toad behaviour differed across the invasive range, and toad righting behaviour was associated with bacterial function but not with parasite infection.

Geographic divergence in gut bacteria

First, we consider the differences in gut bacteria between toads from the invasion-front and the range-core. Although these populations have been separated by less than a century, the toads have diverged remarkably in morphology, physiology and behaviour and much of that divergence is heritable [38,39]. Some of those shifts likely reflect evolutionary pressures for increased rates of dispersal, due to adaptive (natural selection) and non-adaptive (spatial sorting) mechanisms [40,41]. Other geographically variable aspects of toad phenotypes likely are responses to different climatic conditions in the newly-invaded regions (hot, seasonally arid) compared to the range-core (cooler, more equable climate) [42]. Similar geographic divergence has been reported for the microbiome on the toad's skin [43]. Our data illustrate that the invasion of Australia by cane toads has been accompanied by substantial divergence in gut bacterial communities. Alpha diversity in gut bacteria was similar in invasion-front and range-core individuals, but there were differences in both the gut bacteria composition and predicted bacterial function between toad populations across the species' Australian invasive range. Predicted bacterial functions better explained cane toad righting behaviour than did gut bacterial community composition. Intriguingly, similarity between gut bacterial communities between individuals within regions in Australia is related to the similarity of their host's epigenome, and this relationship is strengthened in populations where genetic diversity is lowest, such as on the invasion front [44]. Relationships between gut bacterial communities and their hosts

are complex, and that a clear understanding of these relationships requires careful consideration of numerous environmental, host and gut bacterial factors.

The diversity and composition of bacterial communities differed between range-core and invasion-front toads, despite an overall similarity in their dominant phyla and alpha diversity. ASVs in the family Veillonellaceae were higher at the invasion-front (Figure 2A). The abundance of this bacterial family may influence host metabolic regulation. For example, in Brandt's voles (*Lasiopodomys brandtii*) exposed to colder temperatures, voles which huddled had more Veillonellaceae and more short-chain fatty acids (SCFAs) in their intestines than did non-huddling voles [45]. This family produces SCFAs such as propionic acid [46,47], which can increase locomotor activity [48]. The link to host metabolic regulation suggests that invasion-front toads might fuel their invasion in this way [24]. ASVs from another family of SCFA-producing bacteria, Clostridiaceae [49], were also higher in invasion-front toads than those from the range-core. Furthermore, the family Veillonellaceae may be associated with host sociality. A reduction of Veillonellaceae has been observed in children with Autism Spectrum Disorder, often known for desiring social isolation [50]. Higher abundance of Veillonellaceae in invasion-front toads could foster their "bolder" personality, retaining a higher propensity for exploration and risk-taking [26,51].

Several other ASVs that differed across the toad's range also may affect behaviour. ASVs from the family Peptococcaceae, more common in invasion-front toads (Figure 2A), are related to host neurotransmitter levels (noradrenaline linking visual awareness to external world events [52]). For example, Peptococcaceae levels in the caecum are positively correlated with noradrenaline levels in mice [53]. ASVs from family Bacillaceae, lower in invasion-front toads (Figure 2A), might be related to host anxiety (e.g., abundant in methamphetamine-treated rats [54], and in exercised *versus* sedentary mice [55]). Abundant Bacillaceae might induce anxiety-like behaviours, thus intensifying the stress response [54] and decreasing exploratory behaviour in new environments [56]. In summary, invasion-front toads possessed gut bacterial communities that in other studies have been associated with SCFAs production and neurotransmitters. That pattern supports the idea that gut microbes in invasion-front toads may increase locomotor ability, alertness and propensity for exploration and risk-taking. In comparison, range-core toads possessed bacterial taxa that have been associated with anxiety, and a decreased propensity to explore.

Geographic variation was less obvious in the predicted bacterial functional groups than in community composition (Figure 1), consistent with the hypothesis that bacterial function is more conservative than taxonomic composition (e.g. in fire salamanders [2]). Different gut microbiota can have similar bacterial functions, increasing resilience and functional stability [2,3,57]. Despite this broad similarity, bacterial functions differed between range-core and invasion-front toads. These differences included those involved in functional pathways related to food sources and metabolism. Invasion-front toads had less bacterial function in the superpathway of pyrimidine ribonucleosides degradation, which provides a nitrogen source for microbes [58] and plays an important role in perturbations in the uridine monophosphate (UMP) biosynthetic pathways. This pathway allows the bacterial cell to sense signals such as starvation, nucleic acid degradation, and availability of exogenous pyrimidines, and to adapt the production of the extracellular matrix to changing environmental conditions [59]. This function might help to explain the disappearance of Verrucomicrobia as a dominant taxon. As for microbe metabolism, invasion-front toads have higher abundance of bacterial functions in factor 420 biosynthesis, critical to bacterial metabolism and mediating important redox transformations involved in bacterial persistence, antibiotic biosynthesis, pro-drug activation, and methanogenesis [60].

We also detected geographic variation in bacterial functional pathways that contribute to host health. Invasion-front toads exhibited bacterial functions beneficial to host health and immunity: (i) phosphopantothenate biosynthesis (involved in bacterial production of coenzyme A [61]); and (ii) superpathway of sialic acids biosynthesis (involved in immunity including acting as host receptors and pathogen decoys for viruses and bacteria [62] and especially critical for preventing neural tissue damage [63]). Despite this abundance of health-promoting bacterial functions, these toads may also face health challenges. Invasion-front toad bacteria had a higher abundance of urate biosynthesis function (urate biosynthesis/inosine 5'-phosphate degradation, the only significantly different one out of the top 30 abundant functions), which affects serum urate levels [64]. High levels of urate can result in the formation of needle-like crystals of urate in the joints (gout), perhaps related to severe spinal arthritis in invasion-front cane toads [65].

Associations between lungworms and host gut bacteria

Pathogens and parasites impact the composition of the host microbiota and can modify host behaviour in a manner that improves parasite transmission and survival [66–68]. Lungworms can affect cane toad locomotor performance and reduce host endurance, curtailing oxygen supply from

infected lungs [69]. Lungworms also can alter a cane toad's thermal preference and manipulate the timing and location of defecation, thereby enhancing lungworm egg production and larval survival [22]. Lungworms are reported to lag behind their host on the invasion-front by 2-3 years [27] and to affect righting behaviour (prolongs righting time [70]). In the current study, although we collected invasion-front toads in recently invaded areas, we found no difference in lungworm presence or intensity between the invasion-front *versus* range-core toads, nor did we find behavioural differences in lungworm-infected *versus* uninfected toads.

Infection by parasitic lungworms was associated with differences in gut bacteria. Here, the direction of causation is less ambiguous than is the case for geographic variation in the gut bacteria. It seems unlikely that a toad's bacteria affect its probability of carrying adult lungworms, although bacterial-driven differences in habitat selection might create such a link. Instead, we suggest that the presence of lungworms induces a shift in gut bacteria. Consistent with that hypothesis, experimental trials have shown that lungworms modify gastric function in their hosts, changing the volume and consistency of faeces produced in ways that enhance survival of larval lungworms [22]. Shifts in the microbiome inside the gut might be either causes or consequences of that shift in gastric function. Moreover, *C. elegans* are known to prefer specific bacterial foods [71], suggesting that lungworm larvae may also feed selectively on bacteria in the gut, generating differences in bacterial communities between lungworm-infected toads *versus* non-infected conspecifics. Additionally, gut bacteria may affect lungworms via microbiome-induced shifts in host immunity [72].

Associations between host behaviours and gut bacteria

Interestingly, behaviours including righting effort likelihood and righting time were associated more closely with predicted gut bacterial functions than with bacterial taxonomic composition. Multiple identified taxa may share the same bacterial function, or one taxon may contribute to multiple bacterial functions, obscuring the relationship between host behaviour and bacterial taxonomic composition. Nonetheless, these relationships we found between gut bacterial function and righting behaviours may be related to toad health and/or rearing conditions. A dampened stress response (lower corticosterone levels) in invasion-front toads [73] could result from higher abundance of bacterial functions beneficial to host health and immunity, especially the superpathway of sialic acids biosynthesis [63]. Further, invasion-front toads are more reluctant to flee in simulated predation trials [74]. Dampened stress responses can be related to more exploratory behaviour [56], and to greater dispersal ability [26]. Rearing conditions also affect righting behaviour [17]. Although manipulative

studies are needed to clarify causal relationships between stress responses, proactive behaviours, and gut bacterial functions, our results indicate that host behaviour and gut bacterial functions are related, suggesting that gut bacteria may be an important driver of invasion.

Our study has identified patterns rather than testing alternative hypotheses about underlying causal processes. To clarify causal mechanisms underlying the geographic divergence in gut bacteria across the toads' Australian range, future studies could use reciprocal transplantation to examine if (and how) their gut bacteria respond to novel environmental conditions. Breeding these animals, and raising their offspring under common-garden conditions, could reveal the degree to which a toad's gut bacteria is driven by host genetics *versus* their rearing conditions [75,76]. To clarify the hypothesis that changes in gut bacteria mediate the ability of lungworm parasites to modify host gut function, we could implant colon contents from infected into uninfected toads. In short, our discovery of strong associations between gut bacteria and important facets of toad ecology provides the opportunity to move to hypothesis-testing experimental studies.

Our research illustrates that during invasion, as a species expands across a novel and variable landscape, a complex relationship between host behaviour, its parasite community, and its microbiome may unfold. A clearer understanding of these relationships and how they influence the rate of expansion are key to understanding the role of the holobiont during invasion [77]. Such studies also will advance our understanding of co-evolution and may facilitate innovative approaches to invasive species management.

Ethics

Approved by University of Adelaide Animal Ethics Committee (S-2018-056).

Author Contributions

Designed research: JZ, TMN, CRL, SJZ, LAR; performed research: JZ, CRL, GWF, KS, LAR; analyzed data: JZ, TMN, CRL, SJZ, KS, LAR; drafted manuscript: JZ, RS, TMN, CRL, SJZ, GWF, KS, LAR.

Funding

330 This project was supported by the Holsworth Wildlife Research Endowment and University of
331 Adelaide Graduate Research Scholarship to JZ, and a UNSW Scientia Fellowship to LAR. CMRL is
332 partially supported by the National Institute of Food and Agriculture (AFRI Competitive Grant
333 1018617 and Hatch Program 1020852).

334 **Acknowledgments**

335 We thank Dr. Alice Russo and Ms. Rita Kurpiewski for field assistance, and Dr. Eve Slavich (UNSW
336 Stats Central) for statistical advice.

337 **Data Availability Statement**

338 Supplementary methods and results available online. Code available at:
339 <https://github.com/jiazhou0116/gut-microbiome-analyses-2>. Sequence data are available in NCBI
340 Sequence Read Archive (PRJNA670039). Raw ecological data available on Dryad
341 (doi:10.5061/dryad.v15dv41tw).

342

Bibliography

- [1] Hopkins WA. Amphibians as models for studying environmental change. *ILAR J* 2007;48:270–7. doi:10.1093/ilar.48.3.270.
- [2] Bletz MC, Goedbloed DJ, Sanchez E, Reinhardt T, Tebbe CC, Bhuju S, et al. Amphibian gut microbiota shifts differentially in community structure but converges on habitat-specific predicted functions. *Nat Commun* 2016;7:13699. doi:10.1038/ncomms13699.
- [3] Huang B-H, Chang C-W, Huang C-W, Gao J, Liao P-C. Composition and functional specialists of the gut microbiota of frogs reflect habitat differences and agricultural activity. *Front Microbiol* 2017;8:2670. doi:10.3389/fmicb.2017.02670.
- [4] Scott KP, Gratz SW, Sheridan PO, Flint HJ, Duncan SH. The influence of diet on the gut microbiota. *Pharmacol Res* 2013;69:52–60. doi:10.1016/j.phrs.2012.10.020.
- [5] Zmora N, Suez J, Elinav E. You are what you eat: diet, health and the gut microbiota. *Nat Rev Gastroenterol Hepatol* 2019;16:35–56. doi:10.1038/s41575-018-0061-2.
- [6] Bercik P, Denou E, Collins J, Jackson W, Lu J, Jury J, et al. The intestinal microbiota affect central levels of brain-derived neurotropic factor and behavior in mice. *Gastroenterology* 2011;141:599–609, 609.e1. doi:10.1053/j.gastro.2011.04.052.
- [7] Kraimi N, Calandreau L, Zemb O, Germain K, Dupont C, Velge P, et al. Effects of gut microbiota transfer on emotional reactivity in Japanese quails (*Coturnix japonica*). *J Exp Biol* 2019;222. doi:10.1242/jeb.202879.
- [8] Teseo S, van Zweden JS, Pontieri L, Kooij PW, Sørensen SJ, Wenseleers T, et al. The scent of symbiosis: gut bacteria may affect social interactions in leaf-cutting ants. *Anim Behav* 2019;150:239–54. doi:10.1016/j.anbehav.2018.12.017.
- [9] Schretter CE, Vielmetter J, Bartos I, Marka Z, Marka S, Argade S, et al. A gut microbial factor modulates locomotor behaviour in *Drosophila*. *Nature* 2018;563:402–6. doi:10.1038/s41586-018-0634-9.
- [10] Borrelli L, Aceto S, Agnisola C, De Paolo S, Dipineto L, Stilling RM, et al. Probiotic modulation of the microbiota-gut-brain axis and behaviour in zebrafish. *Sci Rep* 2016;6:30046. doi:10.1038/srep30046.
- [11] Savignac HM, Kiely B, Dinan TG, Cryan JF. Bifidobacteria exert strain-specific effects on stress-related behavior and physiology in BALB/c mice. *Neurogastroenterol Motil* 2014;26:1615–27. doi:10.1111/nmo.12427.
- [12] Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I, Rosenberg E. Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 2010;107:20051–6. doi:10.1073/pnas.1009906107.
- [13] Vogel H, Shukla SP, Engl T, Weiss B, Fischer R, Steiger S, et al. The digestive and defensive

- 379 basis of carcass utilization by the burying beetle and its microbiota. Nat Commun
380 2017;8:15186. doi:10.1038/ncomms15186.
- 381 [14] Wong AC-N, Wang Q-P, Morimoto J, Senior AM, Lihoreau M, Neely GG, et al. Gut
382 Microbiota Modifies Olfactory-Guided Microbial Preferences and Foraging Decisions in
383 *Drosophila*. Curr Biol 2017;27:2397–2404.e4. doi:10.1016/j.cub.2017.07.022.
- 384 [15] Jones JC, Fruciano C, Marchant J, Hildebrand F, Forslund S, Bork P, et al. The gut
385 microbiome is associated with behavioural task in honey bees. Insectes Soc 2018;65:419–29.
386 doi:10.1007/s00040-018-0624-9.
- 387 [16] Davidson GL, Raulo A, Knowles SCL. Identifying Microbiome-Mediated Behaviour in Wild
388 Vertebrates. Trends Ecol Evol (Amst) 2020. doi:10.1016/j.tree.2020.06.014.
- 389 [17] Stuart KC, Shine R, Brown GP. Proximate mechanisms underlying the rapid modification of
390 phenotypic traits in cane toads (*Rhinella marina*) across their invasive range within Australia.
391 Biological Journal of the Linnean Society 2019;126:68–79. doi:10.1093/biolinnean/bly150.
- 392 [18] Wagener C, Mohanty NP, Measey J. The gut microbiome facilitates ecological adaptation in
393 an invasive vertebrate. BioRxiv 2020. doi:10.1101/2020.12.10.418954.
- 394 [19] Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. Is invasion success explained by the
395 enemy release hypothesis? Ecol Lett 2004;7:721–33. doi:10.1111/j.1461-0248.2004.00616.x.
- 396 [20] Poulin R. Parasite manipulation of host behavior: an update and frequently asked questions.
397 Advances In The Study Of Behavior, vol. 41, Elsevier; 2010, p. 151–86. doi:10.1016/S0065-
398 3454(10)41005-0.
- 399 [21] Tingley R, Ward-Fear G, Schwarzkopf L, Greenlees MJ, Phillips BL, Brown G, et al. New
400 weapons in the toad toolkit: A review of methods to control and mitigate the biodiversity
401 impacts of invasive cane toads (*rhinella marina*). Q Rev Biol 2017;92:123–49.
402 doi:10.1086/692167.
- 403 [22] Finnerty PB, Shine R, Brown GP. Survival of the feces: Does a nematode lungworm
404 adaptively manipulate the behavior of its cane toad host? Ecol Evol 2018;8:4606–18.
405 doi:10.1002/ece3.3870.
- 406 [23] Shine R. The ecological impact of invasive cane toads (*Bufo marinus*) in Australia. Q Rev Biol
407 2010;85:253–91. doi:10.1086/655116.
- 408 [24] Llewelyn J, Phillips BL, Alford RA, Schwarzkopf L, Shine R. Locomotor performance in an
409 invasive species: cane toads from the invasion front have greater endurance, but not speed,
410 compared to conspecifics from a long-colonised area. Oecologia 2010;162:343–8.
411 doi:10.1007/s00442-009-1471-1.
- 412 [25] Hudson CM, Vidal-García M, Murray TG, Shine R. The accelerating anuran: evolution of
413 locomotor performance in cane toads (*Rhinella marina*, *Bufo* spp.) at an invasion front. Proc R
414 Soc Lond, B, Biol Sci 2020;287:20201964. doi:10.1098/rspb.2020.1964.
- 415 [26] Gruber J, Brown G, Whiting MJ, Shine R. Geographic divergence in dispersal-related

- behaviour in cane toads from range-front versus range-core populations in Australia.
Behavioral Ecology And 2017.
- [27] Phillips BL, Kelehear C, Pizzatto L, Brown GP, Barton D, Shine R. Parasites and pathogens lag behind their host during periods of host range advance. *Ecology* 2010;91:872–81. doi:10.1890/09-0530.1.
- [28] Child T, Phillips BL, Shine R. Abiotic and biotic influences on the dispersal behavior of metamorph cane toads (*Bufo marinus*) in tropical Australia. *J Exp Zool A Ecol Genet Physiol* 2008;309:215–24. doi:10.1002/jez.450.
- [29] Selechnik D, Richardson MF, Shine R, DeVore JL, Ducatez S, Rollins LA. Increased adaptive variation despite reduced overall genetic diversity in a rapidly adapting invader. *Front Genet* 2019;10:1221. doi:10.3389/fgene.2019.01221.
- [30] Brown GP, Kelehear C, Pizzatto L, Shine R. The impact of lungworm parasites on rates of dispersal of their anuran host, the invasive cane toad. *Biol Invasions* 2016;18:103–14. doi:10.1007/s10530-015-0993-1.
- [31] Hudson CM, Brown GP, Shine R. Evolutionary shifts in anti-predator responses of invasive cane toads (*Rhinella marina*). *Behav Ecol Sociobiol (Print)* 2017;71:134. doi:10.1007/s00265-017-2367-4.
- [32] Kelehear C, Jones HI. Nematode larvae (order Spirurida) in gastric tissues of Australian anurans: a comparison between the introduced cane toad and sympatric native frogs. *J Wildl Dis* 2010;46:1126–40. doi:10.7589/0090-3558-46.4.1126.
- [33] Baker MR. The free-living and parasitic development of *Rhabdias* spp. (Nematoda: Rhabdiasidae) in amphibians. *Can J Zool* 1979;57:161–78. doi:10.1139/z79-014.
- [34] Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet C, Al-Ghalith GA, et al. QIIME 2: Reproducible, interactive, scalable, and extensible microbiome data science 2018. doi:10.7287/peerj.preprints.27295v1.
- [35] Douglas GM, Maffei VJ, Zaneveld J, Yurgel SN, Brown JR, Taylor CM, et al. PICRUSt2: An improved and extensible approach for metagenome inference. *BioRxiv* 2019. doi:10.1101/672295.
- [36] R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2020.
- [37] Davidson GL, Cooke AC, Johnson CN, Quinn JL. The gut microbiome as a driver of individual variation in cognition and functional behaviour. *Philos Trans R Soc Lond B, Biol Sci* 2018;373. doi:10.1098/rstb.2017.0286.
- [38] Rollins LA, Richardson MF, Shine R. A genetic perspective on rapid evolution in cane toads (*Rhinella marina*). *Mol Ecol* 2015;24:2264–76. doi:10.1111/mec.13184.
- [39] Shine R. The impact of an invasive amphibian: the cane toad *Rhinella marina*. In: Heatwole H, Rowley J, editors. *Status of Conservation and Decline of Amphibians: Australia, New Zealand,*

and Pacific Islands, Victoria, Australia: CSIRO Publishing; 2018, p. 107–24.

- [40] Shine R, Brown GP, Phillips BL. An evolutionary process that assembles phenotypes through space rather than through time. *Proc Natl Acad Sci USA* 2011;108:5708–11. doi:10.1073/pnas.1018989108.
- [41] Brown GP, Kelehear C, Shine R. The early toad gets the worm: cane toads at an invasion front benefit from higher prey availability. *J Anim Ecol* 2013;82:854–62. doi:10.1111/1365-2656.12048.
- [42] Kosmala GK, Brown GP, Shine R. Thin-skinned invaders: geographic variation in the structure of the skin among populations of cane toads (*Rhinella marina*). *Biological Journal of the Linnean Society* 2020;131:611–21. doi:10.1093/biolinnean/blaa128.
- [43] Weitzman CL, Kaestli M, Gibb K, Brown GP, Shine R, Christian K. Disease exposure and antifungal bacteria on skin of invasive cane toads, australia. *Emerging Infect Dis* 2019;25:1770–1. doi:10.3201/eid2509.190386.
- [44] Zhou J, Tesfamichael K, Zhou SJ, Rollins LA, Rodriguez Lopez CM. Genetic similarity enhances the strength of the relationship between gut bacteria and host DNA methylation. *BioRxiv* 2021. doi:10.1101/2021.07.10.451923.
- [45] Zhang X-Y, Sukhchuluun G, Bo T-B, Chi Q-S, Yang J-J, Chen B, et al. Huddling remodels gut microbiota to reduce energy requirements in a small mammal species during cold exposure. *Microbiome* 2018;6:103. doi:10.1186/s40168-018-0473-9.
- [46] Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol* 2017;19:29–41. doi:10.1111/1462-2920.13589.
- [47] Gamage HKAH, Tetu SG, Chong RWW, Ashton J, Packer NH, Paulsen IT. Cereal products derived from wheat, sorghum, rice and oats alter the infant gut microbiota in vitro. *Sci Rep* 2017;7:14312. doi:10.1038/s41598-017-14707-z.
- [48] Thomas RH, Meeking MM, Mephram JR, Tichenoff L, Possmayer F, Liu S, et al. The enteric bacterial metabolite propionic acid alters brain and plasma phospholipid molecular species: further development of a rodent model of autism spectrum disorders. *J Neuroinflammation* 2012;9:153. doi:10.1186/1742-2094-9-153.
- [49] Hugenholtz F, Davids M, Schwarz J, Müller M, Tomé D, Schaap P, et al. Metatranscriptome analysis of the microbial fermentation of dietary milk proteins in the murine gut. *PLoS One* 2018;13:e0194066. doi:10.1371/journal.pone.0194066.
- [50] Kang D-W, Park JG, Ilhan ZE, Wallstrom G, Labaer J, Adams JB, et al. Reduced incidence of *Prevotella* and other fermenters in intestinal microflora of autistic children. *PLoS One* 2013;8:e68322. doi:10.1371/journal.pone.0068322.
- [51] González-Bernal E, Brown GP, Shine R. Invasive cane toads: social facilitation depends upon an individual's personality. *PLoS One* 2014;9:e102880. doi:10.1371/journal.pone.0102880.
- [52] Gelbard-Sagiv H, Magidov E, Sharon H, Hendler T, Nir Y. Noradrenaline modulates visual

- 490 perception and late visually evoked activity. *Curr Biol* 2018;28:2239–2249.e6.
491 doi:10.1016/j.cub.2018.05.051.
- 492 [53] Houlden A, Goldrick M, Brough D, Vizi ES, Lénárt N, Martinecz B, et al. Brain injury induces
493 specific changes in the caecal microbiota of mice via altered autonomic activity and
494 mucoprotein production. *Brain Behav Immun* 2016;57:10–20. doi:10.1016/j.bbi.2016.04.003.
- 495 [54] Ning T, Gong X, Xie L, Ma B. Gut Microbiota Analysis in Rats with Methamphetamine-
496 Induced Conditioned Place Preference. *Front Microbiol* 2017;8:1620.
497 doi:10.3389/fmicb.2017.01620.
- 498 [55] Choi JJ, Eum SY, Rampersaud E, Daunert S, Abreu MT, Toborek M. Exercise attenuates
499 PCB-induced changes in the mouse gut microbiome. *Environ Health Perspect* 2013;121:725–
500 30. doi:10.1289/ehp.1306534.
- 501 [56] Golla A, Østby H, Kermen F. Chronic unpredictable stress induces anxiety-like behaviors in
502 young zebrafish. *Sci Rep* 2020;10:10339. doi:10.1038/s41598-020-67182-4.
- 503 [57] Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and
504 resilience of the human gut microbiota. *Nature* 2012;489:220–30. doi:10.1038/nature11550.
- 505 [58] West TP. Degradation of pyrimidine ribonucleosides by *Pseudomonas aeruginosa*. *Antonie*
506 *Van Leeuwenhoek* 1996;69:331–5. doi:10.1007/bf00399622.
- 507 [59] Garavaglia M, Rossi E, Landini P. The pyrimidine nucleotide biosynthetic pathway modulates
508 production of biofilm determinants in *Escherichia coli*. *PLoS One* 2012;7:e31252.
509 doi:10.1371/journal.pone.0031252.
- 510 [60] Bashiri G, Antoney J, Jirgis ENM, Shah MV, Ney B, Copp J, et al. A revised biosynthetic
511 pathway for the cofactor F420 in prokaryotes. *Nat Commun* 2019;10:1558.
512 doi:10.1038/s41467-019-09534-x.
- 513 [61] Sibon OCM, Strauss E. Coenzyme A: to make it or uptake it? *Nat Rev Mol Cell Biol*
514 2016;17:605–6. doi:10.1038/nrm.2016.110.
- 515 [62] Varki A, Gagneux P. Multifarious roles of sialic acids in immunity. *Ann N Y Acad Sci*
516 2012;1253:16–36. doi:10.1111/j.1749-6632.2012.06517.x.
- 517 [63] Liao H, Klaus C, Neumann H. Control of innate immunity by sialic acids in the nervous tissue.
518 *Int J Mol Sci* 2020;21. doi:10.3390/ijms21155494.
- 519 [64] Sinnott-Armstrong N, Naqvi S, Rivas MA, Pritchard JK. GWAS of three molecular traits
520 highlights core genes and pathways alongside a highly polygenic background. *BioRxiv* 2020.
521 doi:10.1101/2020.04.20.051631.
- 522 [65] Brown GP, Shilton C, Phillips BL, Shine R. Invasion, stress, and spinal arthritis in cane toads.
523 *Proc Natl Acad Sci USA* 2007;104:17698–700. doi:10.1073/pnas.0705057104.
- 524 [66] House PK, Vyas A, Sapolsky R. Predator cat odors activate sexual arousal pathways in brains
525 of *Toxoplasma gondii* infected rats. *PLoS One* 2011;6:e23277.

doi:10.1371/journal.pone.0023277.

- [67] Poulin R. Parasite manipulation of host behavior: an update and frequently asked questions. *Advances In The Study Of Behavior*, vol. 41, Elsevier; 2010, p. 151–86. doi:10.1016/S0065-3454(10)41005-0.
- [68] Gegear RJ, Otterstatter MC, Thomson JD. Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. *Proc Biol Sci* 2006;273:1073–8. doi:10.1098/rspb.2005.3423.
- [69] Pizzatto L, Shine R. Lungworm infection modifies cardiac response to exercise in cane toads. *J Zool* 2012;287:150–5. doi:10.1111/j.1469-7998.2012.00898.x.
- [70] Finnerty PB. The physiological and ecological impacts of lungworm infection on invasive cane toads. Undergraduate thesis. University of Sydney, 2017.
- [71] Abada EA, Sung H, Dwivedi M, Park B-J, Lee S-K, Ahnn J. C. elegans behavior of preference choice on bacterial food. *Mol Cells* 2009;28:209–13. doi:10.1007/s10059-009-0124-x.
- [72] Zheng D, Liwinski T, Elinav E. Interaction between microbiota and immunity in health and disease. *Cell Res* 2020;30:492–506. doi:10.1038/s41422-020-0332-7.
- [73] Brown GP, Kelehear C, Shilton CM, Phillips BL, Shine R. Stress and immunity at the invasion front: a comparison across cane toad (*Rhinella marina*) populations. *Biological Journal of the Linnean Society* 2015;116:748–60. doi:10.1111/bij.12623.
- [74] Hudson CM, Brown GP, Shine R. Evolutionary shifts in anti-predator responses of invasive cane toads (*Rhinella marina*). *Behav Ecol Sociobiol* (Print) 2017;71:134. doi:10.1007/s00265-017-2367-4.
- [75] Zhang C, Zhang M, Wang S, Han R, Cao Y, Hua W, et al. Interactions between gut microbiota, host genetics and diet relevant to development of metabolic syndromes in mice. *ISME J* 2010;4:232–41. doi:10.1038/ismej.2009.112.
- [76] Fan P, Bian B, Teng L, Nelson CD, Driver J, Elzo MA, et al. Host genetic effects upon the early gut microbiota in a bovine model with graduated spectrum of genetic variation. *ISME J* 2020;14:302–17. doi:10.1038/s41396-019-0529-2.
- [77] Arnaud-Haond S, Aires T, Candeias R, Teixeira SJL, Duarte CM, Valero M, et al. Entangled fates of holobiont genomes during invasion: nested bacterial and host diversities in *Caulerpa taxifolia*. *Mol Ecol* 2017;26:2379–91. doi:10.1111/mec.14030.
- [Preprint] Zhou J, Nelson TM, Rodriguez Lopez C, Zhou SJ, Ward-Fear G, Stuart KC, et al. The gut bacteria of an invasive amphibian respond to the dual challenges of range-expansion and parasite attack. *BioRxiv* 2020. doi:10.1101/2020.11.16.385690.

562 Tables and Figures

563 **Table 1a-d.** The association between a) single host factors and microbial community, b) single host
564 factors and predicted microbial function, c) single behavioural trait and microbial community and d)
565 single behavioural trait and predicted microbial function. Significant p-values denoted by an asterisk.

566

	MDS1	MDS2	r ²	Pr(>r)
<i>a. Host factor/microbial community</i>				
SUL	0.170	0.985	0.071	0.119
BodyWeight	-0.009	1.000	0.051	0.231
Lungworms	0.254	-0.967	0.020	0.585
Occurrence of lungworms	0.453	-0.892	0.128	0.023*
<i>b. Host factor/microbial function</i>				
SUL	0.680	0.733	0.021	0.556
BodyWeight	0.610	0.793	0.041	0.311
Lungworms	-0.791	0.612	0.007	0.839
Occurrence of lungworms	-0.827	-0.562	0.059	0.187
<i>c. Behavioural trait/microbial community</i>				
Struggle score	-0.814	0.580	0.044	0.268
Struggle likelihood	-0.880	-0.474	0.021	0.555
Righting effort	-0.124	0.992	0.075	0.121
Righting effort likelihood	0.499	0.867	0.0255	0.474
Righting time	0.933	0.360	0.0615	0.174
<i>d. Behavioural trait/microbial function</i>				

Struggle score	0.376	0.927	0.051	0.221
Struggle likelihood	0.183	0.983	0.017	0.606
Righting effort	0.571	0.821	0.070	0.141
Righting effort likelihood	0.399	0.917	0.035	0.362
Righting time	-0.956	-0.294	0.004	0.894

Table 2. Association between gut microbiota variation and lungworm occurrence and intensity, based on Bray Curtis dissimilarity values for microbial community assemblages.

	Df	SumOfSqs	R2	F	Pr(>F)
Location	1	1.314	0.091	6.026	<0.001***
Lungworm_occurrence	1	0.482	0.033	2.213	0.005**
Lungworms_intensity	1	0.150	0.010	0.688	0.861
Location:Lungworm_occurrence	1	0.331	0.023	1.520	0.075
Location:Lungworm_intensity	1	0.357	0.025	1.638	0.043*
Residual	54	11.774	0.817		
Total	59	14.409	1.000		

Significance codes: 0 '***' ≤0.001 '**' ≤0.01 '*' ≤0.05.

Figure 1. Beta diversity by location. Principle coordinate analysis plot of Bray Curtis distance of microbial community (A) and Bray Curtis distance of predicted functional groups (B) from 60 cane toad individuals of the invasion-front (Kununurra, Old Theda, and Mary Pool) and the range-core (Rossville, Croyden, and Lucinda).

Figure 2. Significantly different bacterial taxa and predicted functions between range-core (QLD) and invasion-front (WA) toads' colon. Significant differences were identified between locations via differential abundance testing based on a negative binomial distribution. The dots represent the average log₂ fold change (x axis) abundance and positive log₂ fold changes signify increased abundance in range-core, and negative log₂ fold changes display increased abundance in invasion-front. Bacterial taxa (A) were classified to the taxonomic level of family (y axis) and coloured by taxonomic level of phylum. Family name in bracket is proposed taxonomy by Greengenes. Only ASVs that could be matched to a known bacterial family and with a log₂FoldChange value higher than 20 or lower than -20 are presented. Predicted functions (B) with a log₂FoldChange value higher than 3 or lower than -3 are presented.

Figure 3. Heatmap for top 30 functional group abundance.

Heatmap indicates the top 30 functional groups in the intestinal samples from range-core and invasion-front toads. Abundance indicates the raw count of functional groups inferred from taxonomic 16S sequences using PICRUST where light blue is high abundance and dark blue is lower abundance. Functional pathways that differ significantly between range-core and invasion-front toads are highlighted in bold. Range-core includes Rossville, Croyden, and Lucinda; invasion-front includes Kununurra, Old Theda, and Mary Pool.

Figure 4. Main variables that affect predicted function differentiation among individuals. CAP (capscale) plot displays the combination of variables that explained the greatest variation in the predicted functions through model selection, using 60 cane toad individuals from the invasion-front (Kununurra, Old Theda, and Mary Pool) and the range-core (Rossville, Croyden, and Lucinda). The final model explained 17.8% of variation in the microbial predicted functions, which includes righting effort likelihood (AIC = 53.613), occurrence of lung worms (AIC = 54.297) and righting time (AIC = 56.912) explained the greatest variation.

A Microbial community – Bray Curtis

B Microbial predicted functions – Bray Curtis

A

ASVs: range-core vs invasion-front

B

Predicted Functions: range-core vs invasion-front