Sex-specific body mass ageing trajectories in adult Asian elephants

Lucas D. Lalande^{1,2,3,a}, Virpi Lummaa^{1,a}, Htoo H. Aung⁴, Win Htut⁴, U. Kyaw Nyein⁴,
Vérane Berger^{1,a,*}, Michael Briga^{1,a,*}

¹ Department of Biology, University of Turku, Turku, Finland;

² Université Bourgogne Franche-Comté, Dijon, France;

³ Present address: Université de Lyon, Université Lyon 1, UMR CNRS 5558,
Villeurbanne CEDEX, France;

⁴ Myanma Timber Enterprise, Ministry of Natural Resources and Environmental
Conservation, West Gyogone Forest Compound, Yangon, Myanmar;

* Equal contribution of these authors

^a Corresponding authors: L. D. Lalande: lucas.lalande@univ-lyon1.fr, V. Lummaa:
virpi.lummaa@gmail.com, V. Berger: verane.berger@orange.fr, M. Briga:
michbriga@gmail.com

Abstract

In species with marked sexual dimorphism and where one sex undergoes stronger intrasexual competition, that sex is expected to age earlier or quicker. Here, we utilise a unique, longitudinal dataset of a semi-captive population of Asian elephants (*Elephas maximus*), a species with marked male-biased intrasexual competition, with males being larger and living shorter, and test the hypothesis that males show earlier and/or faster body mass ageing than females. We show sex-specific body mass ageing trajectories: adult males gained weight up to the age of 48 years old, followed by a decrease in body mass until natural death. In contrast, adult females gained body mass with age until a body mass decline in the last year of life. Our study shows that sex-specific life-histories shape ageing patterns consistent with the predictions of the classical theory of ageing.

Keywords: senescence, sex-differences, terminal decline, *Elephas maximus*, long-term data

30 Introduction

31 Ageing – a decline in organismal functioning with age (Monaghan et al., 2008) – has
 32 been observed in many species (Jones et al., 2014). However, the onset and rates of
 33 ageing differ both between (Jones et al., 2014) and within species (Nussey et al.,
 34 2007) and between sexes (Douhard et al., 2017). A main challenge in ageing
 35 research is to quantify and explain such differences in the onset and rates of ageing
 36 (Rando & Wyss-Coray, 2021).

37 In species with sex-specific intrasexual competition, classic theory predicts that the
 38 sex with the highest intrasexual competition has a shorter lifespan and an earlier
 39 onset and/or higher rate of ageing (Maklakov & Lummaa, 2013). The rationale is that
 40 high intrasexual selection often results in one sex showing conspicuous displays or
 41 aggressive intrasexual behaviours, leading to increased mortality and a *live fast, die*
 42 *young* pace of life (Bonduriansky et al., 2008; Clutton-Brock & Isvaran, 2007;
 43 Maklakov & Lummaa, 2013). Accordingly, in polygynous species with male-biased
 44 intrasexual competition, males often die earlier (Lemaître et al., 2020) and age earlier
 45 or faster than females (Beirne et al., 2015; Clutton-Brock & Isvaran, 2007; Douhard et
 46 al., 2017; Nussey et al., 2009; Tidière et al., 2015). However, recent conceptual
 47 developments have shown that this association can be disrupted. This can occur for
 48 example because of condition-dependent extrinsic mortality selecting particularly
 49 high-performing individuals in the population (Chen & Maklakov, 2014) or
 50 canalisation (*i.e.* the more a trait contributes to fitness, the less it should deviate from
 51 optimal trait value, with respect to environmental variation (Flatt, 2005)), thereby
 52 contradicting the theoretically expected earlier or faster ageing in males. The extent
 53 to which such phenomena occur in nature remains unknown.

Here, we used a unique long-term dataset to describe sex-specific body mass ageing trajectories in a nutritionally unsupplemented semi-captive timber population of Asian elephants (*Elephas maximus*) living in their natural environment in Myanmar. Body mass is of interest in the study of ageing because it is positively associated with key life-history traits such as reproduction and lifespan in many non-human species (Hämäläinen et al., 2014). Accordingly, in Asian elephants, seasonal variation in body mass was positively associated with survival the following month (e.g. low body mass was associated with low survival during dry season) (Mumby, Mar, et al., 2015).

However, we know almost nothing about how body mass changes with age in elephants despite the interest in studying ageing in such a long-lived, social and sexually dimorphic non-human species. While females live in kin groups, adult males often roam solitary, undergo a more intense intrasexual competition for dominance and mating (Sukumar, 2003) and hence are bigger, heavier (Mumby, Chapman, et al., 2015), more aggressive and less sociable (Selmann et al., 2019) and shorter-lived than females (respective median lifespans: 30.8 and 44.7 years) (Lahdenperä et al., 2018). Based on this male-biased intrasexual competition and shorter lifespan, and following the classic theory of ageing, we expected males to display earlier and/or faster body mass ageing than females (Bonduriansky et al., 2008; Maklakov & Lummaa, 2013).

Material and methods

Study population

We studied the world's largest semi-captive Asian elephant population consisting of around 3,000 individually-marked elephants owned by the government-

run Myanma Timber Enterprise (MTE) (Leimgruber et al., 2008). Their birth, death, pedigree details, and morphological measurements have been recorded for almost a century by local veterinarians. These elephants are distributed across Myanmar in forest camps and used as riding, transport and drafting animals. Elephants work during the day and, at night, they socialise, mate and forage freely, unsupervised in forests (Oo, 2010; Zaw, 1997). There are no husbandry procedures and timber elephants are never culled. Calves born in captivity are cared for and nursed by their biological mother and allomothers (Lahdenperä et al., 2016; Lynch et al., 2019). Therefore, breeding rates are natural with no reproductive management. Moreover, there is minimal food provisioning, but elephants benefit from veterinary care that consists of treatment of simple injuries and monitoring of working conditions.

Both males and females are used in the workforce, and each working group of six elephants is composed of both sexes. Males and females follow the same government set limitations on taming age, working and retirement age, working-days per year, hours of work per day and tonnage pulled annually apply to both sexes, although it is possible that males might be used for somewhat different working tasks at times (e.g. when use of tusks is required; only males can possess long tusks in Asian elephants). Pregnant females are given a rest period from mid-pregnancy (around 11 months into gestation) until the calf is 1-year-old (Toke Gale, 1974), while they and their calf are being monitored by their mahouts (individual caretakers and riders) throughout this period. Following this break, mothers are used for light work but are kept with calves at heel and able to suckle on demand until the calf is four or five years old (Oo, 2010) at which point calves are assigned a rider, name, logbook and registration number. After the training period, elephants are used for light work

duties until the age of 17, when they enter the full workforce until retirement around age 50. The MTE maintains their care and logbooks until death.

Data collection and selection

Our analyses focused on age- and sex-specific variation in adult body mass from age 18 onwards, omitting the phase during which elephants grow in height (Mumby, Chapman, et al., 2015). From the logbooks, we extracted 1,901 body masses of 347 elephants with known sex, year of birth ('YOB'), alive or dead status, origin (captive-born or wild-caught) and measurement season (hot: Feb-May, monsoon: Jun-Sep, cold: Oct-Jan (Mumby, Mar, et al., 2015)) from 325 working localities ('township'). Study elephants were aged 18 – 72 years (mean = 39.3) and born 1941 – 1999. Age and cohort information were comparable between sexes, with 171 males (n = 1,316) born 1954 – 1999 and aged 18 – 64 years (mean = 37.4), and 322 females (n = 2,570) born 1941 – 1999 and aged 18 – 72 years (mean = 40.2).

When body mass measurements were not available, we estimated body masses using height to the shoulder and chest girth (n = 1,985 measurements on 342 individuals, method in Supplementary Information 1). In Asian elephants, chest girth reflects changes in mass in addition to the fixed effect of structural size during adulthood (details in SI2). In total, we obtained a median of 4.0 measurements/individual [2.5 – 97.5th percentiles: 1.0 – 36.4], followed for a median period of 2.8 years [2.5 – 97.5th: 0.0 – 36.6] on 493 elephants (n = 3,886). Two influential observations measured at age 18 and 23 were removed for one male because of particularly low Δ age (Cook's distance = 0.61 and 0.25, mean of 0.001 on all males). Other observations for this male, all after age 50, were included.

Statistical analyses

We investigated the age- and sex-specific variation in body mass in R version 4.0.5 (R Core Team, 2020), using the log-transformed body mass as a dependent variable with a normal error distribution. We first tested whether there were sex-specific ageing trajectories (table S2) and then compared the sex-specific ageing trajectories using general additive mixed models (GAMs) and general linear mixed models (GLMs), with respectively the functions ‘*gamm*’ of the package ‘*mgcv*’ (Wood, 2017) and the function ‘*lmer*’ of the package ‘*lme4*’ (Bates et al., 2015). GAMs allow more flexible ageing trajectories than GLMs, but the more constrained ageing trajectories in GLMs allow a less descriptive identification of differences in ageing trajectories (figure S1) and both approaches gave consistent conclusions (see results section). We identified the best fitting models using the model selection approach based on the second order Akaike Information Criterion (AICc) as implemented in the package ‘*MuMIn*’ (Bartoń, 2019). In brief, the best fitting model has the lowest AICc value, with other models within 4 Δ AICc being plausible and models become increasingly equivocal up to 14 Δ AICc, after which they become implausible (Burnham et al., 2011). Visual inspection of model residuals confirmed that these fulfilled all assumptions of distribution and homogeneity without any influential data points or outliers (see above).

Within- vs. between-individual change

In all models, we accounted for non-independence of data due to repeated measurements from the same individual by including elephant identity (‘ID’) as a random intercept. The composition of the population can change with age for example due to selective disappearance of certain (e.g. lighter or heavier) individuals, which can affect the age trajectory. In order to alleviate this problem to the best possible in this long-lived species, we decomposed body mass changes with age into

between- and within-individual changes following the approach developed by van de Pol & Verhulst, 2006 and van de Pol & Wright, 2009 using two terms: *i*) the age at last measurement for each individual, which captures the between-individual variations and *ii*) a ' Δ Age' term (age at measurement minus the individual's mean age for all measurements) capturing the within-individual changes with age. We mean-centred and standardised ' Δ Age' so that *i*) individuals measured once all get a Δ Age = 0 and hence contribute to the variance of the Δ Age intercept but not to its slope and *ii*) to avoid collinearity and to have comparable variance for Δ Age and Δ Age² (Bolker, 2008; Zuur et al., 2009). Our dataset included 105 individuals with one measurement after the age of 18 and we included these individuals by giving them Δ Age = 0 (*i.e.* mean-centred) so they do not contribute to the coefficient but do contribute to the variance along the Y axis on Δ Age = 0, diminishing the likelihood of a false positive, and *ii*) do contribute to the coefficient of the age at last measurement term, thereby avoiding a bias in the dataset from selecting only the longer-lived or most monitored individuals.

Testing ageing trajectories

We tested several within-individual ageing trajectories: GAMs can have any shape and for GLMs we tested linear, quadratic, threshold and terminal models (figure S1) and we selected the ageing trajectory with the lowest AICc. For GAMs, we identified the age at which maxima occurred based on the first-order derivative (= 0) using the function 'fderiv' of the package 'gratia' (Simpson & Singmann, 2021). For threshold models (figure S1C), we followed the approaches previously developed in Briga et al., 2019 and Douhard et al., 2017. In brief, we first identified the best-fitting threshold age in a series of models, varying the threshold in the ' Δ Age' term between -35 to 22 years with intervals of one Δ Age (1 mean-centered Δ Age = 4.4 and 4.5 years for

males and females respectively) and estimated the threshold and its confidence intervals using $\pm 4 \Delta AIC$ age range. Then we compared the best-fitting threshold model with the other ageing trajectories. Sometimes, declines in trait value appear shortly before death (terminal decline). We coded a ‘terminal’ change (figure S1D) as a binomial factor for whether an individual died during the year following the measurement. We used a one-year-window to avoid a possible seasonal covariation in weight and because it was the best fitting time-window, but note that models with other time-windows gave consistent conclusions (figure S3).

Accounting for temporal and spatial variation in body mass

As body mass variation can be influenced by seasonal, spatial and within-individual factors, we tested whether body mass values were affected by (i) measured or estimated, (ii) individuals were alive or dead, (iii) captive- or wild-born and (iv) the measurement season. To this end we used a combination of backward elimination and forward selection approaches, *i.e.* first we analysed the models including these covariates using the most complex ageing trajectory, identified that none of these factors improved the model, then we eliminated these co-variables from the model, identified the best fitting ageing trajectory, and then we repeated the final models obtained with these covariates and identified again that none of these improved the model fit or altered the previously obtained conclusions (table S3). In our models, we included as random intercepts individual identity to account for the repeated measurement of the same individual. We also included ‘township’ to account for the spatial clustering of individuals across Myanmar (except in male GLMs which lacked power), although actually adding township worsened the model fit (male GAM: $\Delta AICc = +1.5$; female GAM: $\Delta AICc = +0.6$).

Results

At the measurements' starting age of 18 years, males were on average 253 kg heavier than females, weighing respectively 2,559 kg [95%CI: 2,431 – 2,694] and 2,306 kg [95%CI: 2,258 – 2,355] and this difference was statistically significant ($\Delta AICc = -122.6$ in a GLM with vs without sex as a fixed effect).

We identified the elephants body mass ageing trajectories using general additive models (GAMs) and general linear models (GLMs) and both approaches gave consistent results. Both analyses showed that sexes have different body mass ageing trajectories (GAM: $\Delta AICc = -65.7$, figure 1, table S2; GLM: $\Delta AICc = -47.0$, figure 2, table S2) and hence, we identified the ageing trajectories for both sexes separately.

For males, both GAMs and GLMs indicated a body mass gain from age 18 years until their late-forties or early fifties (GAM maximum: 54 years, 95%CI: 53 – 56, figure S2; GLM maximum: 48.3 years, 4 AICc CI: 46.6 – 52.3, figure 2A), followed by a decline until death (figure 1A, figure 2A, Table 1). In GLMs, this maximum was shown through the best fit of a threshold model ($\Delta AICc = -30.4$ compared to a linear trajectory): males gained mass at a rate of 21 kg/year [95%CI: 19.1 – 23.4] or 0.9% [95%CI: 0.9 – 1.0] of males' mean body mass and then lost mass at a rate of 28 kg/year [95%CI: 14.6 – 41.3] or 1.3% [95%CI: 0.7 – 1.8] of males' mean body mass (figure 2A). Regarding the decline, neither GAMs nor GLMs showed confident statistical support that it was terminal (*i.e.* determined by time before death rather than age): in GAMs, models with and without the terminal term were almost equivalent ($\Delta AICc = +0.01$, Table 1) and in GLMs adding a terminal term worsened the model fit ($\Delta AICc = +5.0$, Table 1, figure S3A).

223 For females, both GAMs and GLMs indicated a body mass gain throughout their lives
 224 until a terminal decline during their last year of life (figure 1B, figure 2B, Table 1).
 225 GLMs indicated a mass gain of 8 kg/year [95%CI: 7.4 – 10.3] or 0.3% [95%CI: 0.3 –
 226 0.4] of females' mean body mass ($\Delta AICc = -6.7$, Table 1). Loss of body mass
 227 occurred in the last year of life (GAM: $\Delta AICc = -12.2$, Table 1, figure 1B; GLM: $\Delta AICc$
 228 $= -4.9$, Table 1, figure 2B, figure S3B) and consisted of 173 kg [95%CI: 80 – 263] or
 229 6.8% [95%CI: 3.2 – 10.4] of their mean body mass. For comparison, the extent of the
 230 terminal decline in males, if any, is just over half that in females at 97 kg [95%CI: -20
 231 – 207] or 4.3% [95%CI: -0.9 – 9.2] of males' mean body mass (quantified in the
 232 aforementioned best-fitting threshold model).

Figure 1. Body mass ageing trajectories of (A) males (n=1,316 measurements on 171 individuals) and (B) females (n=2,570 measurements on 322 individuals) with solid lines showing predictions of the best-fitting GAM models (table 1) and grey areas 95%CI. For females, measurements in the terminal year (red) are significantly lower than measurements at other ages (grey), but note that the association (slope) with Δage is for illustration purposes only and was not statistically tested.

Figure 2. Body mass ageing trajectories of (A) males ($n = 1,316$ measurements on 171 individuals) and (B) females ($n = 2,570$ measurements on 322 individuals) with predictions of the best-fitting GLMs (table 1) with grey areas 95%CI. For males, the thick dashed-line shows the threshold age at onset of the body mass decline (1.9 or 48.3 years) with thin dashed-lines the 4 $\Delta AICc-Cl$ [46.6, 52.3]. For females, measurements in the terminal year (red) are significantly lower (intercept) than measurements at other ages (blue). Note that the terminal slope is for illustration purposes only and was not statistically tested.

249 **Table 1.** Best fitting body mass ageing trajectories (bold) for males and females, using GAMs (top) and GLMs (bottom) with for
 250 each model ageing trajectories ranked from the least to the most complex. AICc: second-order Akaike Information Criterion; Δ AICc:
 251 change in AICc relative to the best fitting model; k: degrees of freedom.

Model type		Model	k	Males AICc	Δ AICc	k	Females AICc	Δ AICc
GAM								
	null	$\log(\text{bm}) \sim 1$	4.0	-2,835.8	386.1	4.0	-5,422.6	217.1
	smooth1	$\log(\text{bm}) \sim s(\Delta\text{age}) + \text{age-last}$	8.1	-3,221.9	0.0	8.7	-5,613.9	25.8
	+terminal	$\log(\text{bm}) \sim \text{terminal} + s(\Delta\text{age}) + \text{age-last}$	9.1	-3,221.9	0.0	10.0	-5,627.3	12.4
	smooth2	$\log(\text{bm}) \sim s(\Delta\text{age}) + \text{age-last} + \text{age-last}^2$	9.1	-3,221.2	0.7	9.7	-5,627.5	12.2
	+terminal	$\log(\text{bm}) \sim \text{terminal} + s(\Delta\text{age}) + \text{age-last} + \text{age-last}^2$	10.1	-3,221.3	0.6	10.9	-5,639.7	0.0
GLM								
	null	$\log(\text{bm}) \sim 1$	3.0	-2,830.3	364.0	4.0	-5,442.9	160.7
	linear	$\log(\text{bm}) \sim \Delta\text{age} + \text{age-last}$	5.0	-3,163.9	30.4	6.0	-5,586.1	17.5
	+terminal	$\log(\text{bm}) \sim \text{terminal} + \Delta\text{age} + \text{age-last}$	6.0	-3,158.7	35.6	7.0	-5,592.3	11.3
	age-last ²	$\log(\text{bm}) \sim \Delta\text{age} + \text{age-last} + \text{age-last}^2$	6.0	-3,156.4	37.9	7.0	-5,598.7	4.9
	+terminal	$\log(\text{bm}) \sim \text{terminal} + \Delta\text{age} + \text{age-last} + \text{age-last}^2$	7.0	-3,151.2	43.1	8.0	-5,603.6	0.0
	Δage^2	$\log(\text{bm}) \sim \Delta\text{age} + \Delta\text{age}^2 + \text{age-last}$	6.0	-3,178.5	15.8	7.0	-5,571.6	32.0
	+terminal	$\log(\text{bm}) \sim \text{terminal} + \Delta\text{age} + \Delta\text{age}^2 + \text{age-last}$	7.0	-3,172.9	21.4	8.0	-5,577.9	25.7
	quadratic	$\log(\text{bm}) \sim \Delta\text{age} + \Delta\text{age}^2 + \text{age-last} + \text{age-last}^2$	7.0	-3,171.0	23.3	8.0	-5,584.3	19.3
	+terminal	$\log(\text{bm}) \sim \text{terminal} + \Delta\text{age} + \Delta\text{age}^2 + \text{age-last} + \text{age-last}^2$	8.0	-3,165.4	28.9	9.0	-5,589.3	14.3
	threshold	$\log(\text{bm}) \sim \Delta\text{age1} + \Delta\text{age2} + \text{age-last}$	7.0	-3,194.3	0.0	8.0	-5,580.5	23.1
	+terminal	$\log(\text{bm}) \sim \text{terminal} + \Delta\text{age1} + \Delta\text{age2} + \text{age-last}$	8.0	-3,189.3	5.0	9.0	-5,586.0	17.6
	threshold (age-last ²)	$\log(\text{bm}) \sim \Delta\text{age1} + \Delta\text{age2} + \text{age-last} + \text{age-last}^2$	8.0	-3,186.7	7.6	9.0	-5,592.6	11.0
	terminal	$\log(\text{bm}) \sim \text{terminal} + \Delta\text{age1} + \Delta\text{age2} + \text{age-last} + \text{age-last}^2$	9.0	-3,181.8	12.5	10.0	-5,596.9	6.7

Discussion

We tested whether in a species with marked male-biased intrasexual competition, males showed an earlier and/or faster rate of body mass ageing than females. Both sexes gained mass during early adulthood. However, males began to lose mass from 48.3 years old onwards. In contrast, females lost body mass generally at an older age, namely in their last year of life. Here we discuss the implications of our results in the light of theories of ageing (Williams, 1957) and of the management of Asian elephants.

Asian elephants show male-biased intra-sexual competition, with males being more aggressive (Seltmann et al., 2019), fighting more for dominance and showing higher rates of mortality at all ages than females, including during early development, as calves and during adulthood (Lahdenperä et al., 2018). In such species, classic theory predicts that males should show an earlier onset or accelerated ageing (Williams, 1957). Indeed, in several polygynous mammals, males display higher rates of body mass declines than females, suggested to be due to their stronger intrasexual competition (Beirne et al., 2015; Clutton-Brock & Isvaran, 2007). Our results are consistent with those studies and with the prediction of the classic theory of ageing. Here, the prediction of the classic theory is not disrupted by high early-life condition-dependent mortality in males (Chen & Maklakov, 2014) or by canalisation (Flatt, 2005).

Our study is subject to a number of limitations when it comes down to identifying why sexes may differ in their ageing trajectories. First, it is possible that male elephants in our timber elephant population are used more for tasks requiring strength or tusks, thereby causing an earlier onset of body mass declines in males than in females.

However, both sexes fall under the same government-set workload, care and retirement regulation, except for females' maternity leave. One substantial difference between sexes is that parental care is concentrated on females, with for example only females being given 'parental leave' following reproduction (Toke Gale, 1974). However, since maternity is energetically expensive and no more favourable than timber working, this is unlikely to lead to the delayed onset of body mass declines in females. An ideal test would be to analyse the effect of timber work and maternity leave on body mass dynamics.

Second, elephants have a specific dentition that consists of molar teeth that eventually wear down at the end of their lives, and pathologic malocclusions or lack of molars can lead to weight loss and death by starvation. Although both sexes have the same molar dental anatomy, it is possible that the earlier onset of body mass declines in males may reflect sex-specific differences in tooth wear rather than sex-specific sexual selection. Indeed, in captive elephants, dental problems have been described well before the age at onset of the terminal decline found in this study (Gaillard et al., 2015) and, if there is sex-specific tooth wear, this could be associated with the earlier onset of body mass decline in males. Third, male elephants have recurring periods of physiological "musth" throughout their adult lives, which can temporarily but profoundly impact the body mass of individual males (Eisenberg et al., 1971) thereby affecting the body mass ageing trajectory. Fourth, in our study, we did not find any evidence for body mass-based selective disappearance, but, as it is often the case in long-lived species, the average longitudinal individual monitoring is short relative to the lifespan of this species (e.g. Global BMI Mortality Collaboration et al., 2016; Prospective Studies Collaboration, 2009), and hence we only have limited power to detect such association. It is possible that there are sex-specific dynamics

of selective disappearance, but whether that is the case in Asian elephants remains to be shown. An analysis with more longitudinal data would be useful to tackle this question. Fifth, for both sexes, it is possible that maximum body mass is set by physiological and ecological constraints as indicated by the weight growth curves found earlier in this population (Mumby, Chapman, et al., 2015). These constraints could be to some extent sex-specific, although at this point, we can only speculate as to why these constraints may drive sex-specific ageing trajectory. Finally, we found a maximum body mass in males but not in females. This sex-specific differences could be driven by the fact that male elephants benefit more than females from being heavy during intrasexual competition (Sukumar, 2003). However, another factor to take into account is that retirement occurs at around 50 years in both sexes, which likely diminishes physical exercise and allows more time for foraging, thereby continuing the weight gain. The reduced intrasexual competition in females relative to males, together with this retirement, could lead to the continued mass gain of females. Given that elephants in the wild do not experience timber labour and retirement, we cannot exclude that the sex-specific body mass ageing trajectories could be different in a wild (non-working) population of Asian elephants compared to those found in our study.

We found that females experienced a terminal body mass decline in the last year of life. Our data contain both males and females among the oldest ages (>50), hence sex-specific terminal decline is unlikely to emerge from differences in lifespan. In European badgers (*Meles meles*), a species in which females outlive males, both sexes displayed terminal body mass declines (Beirne et al., 2015). It is possible that the sex-specific terminal declines in our study resulted from differences in power, with 5 old dead males and 18 old dead females. Indeed, for both males and females, the

coefficient and effect size of the terminal terms were negative, but the effect size in males remained about half of that in females (Cohen's $d_{\text{males}} = -0.045$ [-0.10, 0.01], Cohen's $d_{\text{females}} = -0.071$ [-0.11, -0.03]). Hence, it is possible that the sex-specific terminal effect is driven by power issues and we look forward to testing that with several more years of monitoring.

Terminal declines emphasise that the chronological age is rarely a perfect estimation of the biological age which can better describe the 'true biological state' of an organism (Klemmera & Doubal, 2006). In that sense, terminal decline is a biomarker of health and remaining lifespan. The 'terminal illness' hypothesis refers to the age-independent decrease of a trait value, related to the imminent death of the individual (Coulson & Fairweather, 2001) and such terminal effects were shown for example for body mass and sexual signals in mammals and birds (Beirne et al., 2015; Simons et al., 2016). For which traits or under which conditions to expect terminal declines remains yet poorly understood but our study highlights the importance of studying sex-specific differences in ageing and illustrates the need to improve our understanding of the mechanisms driving the diversity of ageing patterns in the wild.

References

- Bartoń, K. (2019). *MuMIn*: Multi-Model Inference (1.43.15) [Computer software]. <https://CRAN.R-project.org/package=MuMIn>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Beirne, C., Delahay, R., & Young, A. (2015). Sex differences in senescence: The role of intra-sexual competition in early adulthood. *Proceedings of the Royal Society B: Biological Sciences*, 282(1811), 20151086. <https://doi.org/10.1098/rspb.2015.1086>

351 Bolker, B. M. (2008). *Ecological Models and Data in R*. Princeton University Press.

352 <https://press.princeton.edu/books/hardcover/9780691125220/ecological-models-and->

353 [data-in-r](https://press.princeton.edu/books/hardcover/9780691125220/ecological-models-and-data-in-r)

354 Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual selection, sexual

355 conflict and the evolution of ageing and life span. *Functional Ecology*, 22(3),

356 443–453. <https://doi.org/10.1111/j.1365-2435.2008.01417.x>

357 Briga, M., Jimeno, B., & Verhulst, S. (2019). Coupling lifespan and aging? The age at onset

358 of body mass decline associates positively with sex-specific lifespan but negatively

359 with environment-specific lifespan. *Experimental Gerontology*, 119, 111–119.

360 <https://doi.org/10.1016/j.exger.2019.01.030>

361 Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and

362 multimodel inference in behavioral ecology: Some background, observations, and

363 comparisons. *Behavioral Ecology and Sociobiology*, 65(1), 23–35.

364 <https://doi.org/10.1007/s00265-010-1029-6>

365 Chen, H., & Maklakov, A. A. (2014). Condition Dependence of Male Mortality Drives the

366 Evolution of Sex Differences in Longevity. *Current Biology*, 24(20), 2423–2427.

367 <https://doi.org/10.1016/j.cub.2014.08.055>

368 Clutton-Brock, T. H., & Isvaran, K. (2007). Sex differences in ageing in natural populations of

369 vertebrates. *Proceedings of the Royal Society B: Biological Sciences*, 274(1629),

370 3097–3104. <https://doi.org/10.1098/rspb.2007.1138>

371 Coulson, J. C., & Fairweather, J. A. (2001). Reduced reproductive performance prior to death

372 in the Black-legged Kittiwake: Senescence or terminal illness? *Journal of Avian*

373 *Biology*, 32(2), 146–152. <https://doi.org/10.1034/j.1600-048X.2001.320207.x>

374 Douhard, F., Gaillard, J.-M., Pellerin, M., Jacob, L., & Lemaître, J.-F. (2017). The cost of

375 growing large: Costs of post-weaning growth on body mass senescence in a wild

376 mammal. *Oikos*, 126(9), 1329–1338. <https://doi.org/10.1111/oik.04421>

377 Eisenberg, J. F., McKay, G. M., & Jainudeen, M. R. (1971). Reproductive behavior of the
378 Asiatic elephant (*Elephas maximus maximus* L.). *Behaviour*, 38(3), 193–225.
379 <https://doi.org/10.1163/156853971x00087>

380 Flatt, T. (2005). The evolutionary genetics of canalization. *The Quarterly Review of Biology*,
381 80(3), 287–316. <https://doi.org/10.1086/432265>

382 Gaillard, J.-M., Berger, V., Tidière, M., Duncan, P., & Lemaître, J.-F. (2015). Does tooth wear
383 influence ageing? A comparative study across large herbivores. *Experimental*
384 *Gerontology*, 71, 48–55. <https://doi.org/10.1016/j.exger.2015.09.008>

385 Global BMI Mortality Collaboration, Di Angelantonio, E., Bhupathiraju, S., Wormser, D., Gao,
386 P., Kaptoge, S., Berrington de Gonzalez, A., Cairns, B., Huxley, R., Jackson, C.,
387 Joshy, G., Lewington, S., Manson, J., Murphy, N., Patel, A., Samet, J., Woodward,
388 M., Zheng, W., Zhou, M., ... Hu, F. (2016). Body-mass index and all-cause
389 mortality: Individual-participant-data meta-analysis of 239 prospective studies in four
390 continents. *The Lancet*, 388(10046), 776–786. <https://doi.org/10.1016/S0140->
391 [6736\(16\)30175-1](https://doi.org/10.1016/S0140-6736(16)30175-1)

392 Hämäläinen, A., Dammhahn, M., Aujard, F., Eberle, M., Hardy, I., Kappeler, P. M., Perret,
393 M., Schliehe-Diecks, S., & Kraus, C. (2014). Senescence or selective disappearance?
394 Age trajectories of body mass in wild and captive populations of a small-bodied
395 primate. *Proceedings of the Royal Society B: Biological Sciences*, 281(1791),
396 20140830. <https://doi.org/10.1098/rspb.2014.0830>

397 Jones, O. R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C. G., Schaible, R., Casper,
398 B. B., Dahlgren, J. P., Ehrlén, J., García, M. B., Menges, E. S., Quintana-Ascencio, P.
399 F., Caswell, H., Baudisch, A., & Vaupel, J. W. (2014). Diversity of ageing across the
400 tree of life. *Nature*, 505(7482), 169–173. <https://doi.org/10.1038/nature12789>

401 Klemmera, P., & Doubal, S. (2006). A new approach to the concept and computation of
402 biological age. *Mechanisms of Ageing and Development*, 127(3), 240–248.
403 <https://doi.org/10.1016/j.mad.2005.10.004>

404 Lahdenperä, M., Mar, K. U., Courtiol, A., & Lummaa, V. (2018). Differences in age-specific
405 mortality between wild-caught and captive-born Asian elephants. *Nature*
406 *Communications*, 9(1), 3023. <https://doi.org/10.1038/s41467-018-05515-8>

407 Lahdenperä, M., Mar, K. U., & Lummaa, V. (2016). Nearby grandmother enhances calf
408 survival and reproduction in Asian elephants. *Scientific Reports*, 6(1), 27213.
409 <https://doi.org/10.1038/srep27213>

410 Leimgruber, P., Senior, B., Aung, M., Songer, M., Mueller, T., Wemmer, C., & Ballou, J.
411 (2008). Modeling population viability of captive elephants in Myanmar (Burma):
412 Implications for wild populations. *Animal Conservation*, 11, 198–205.
413 <https://doi.org/10.1111/j.1469-1795.2008.00172.x>

414 Lemaître, J.-F., Ronget, V., Tidière, M., Allainé, D., Berger, V., Cohas, A., Colchero, F.,
415 Conde, D. A., Garratt, M., Liker, A., Marais, G. A. B., Scheuerlein, A., Székely, T., &
416 Gaillard, J.-M. (2020). Sex differences in adult lifespan and aging rates of mortality
417 across wild mammals. *Proceedings of the National Academy of Sciences*, 117(15),
418 8546–8553. <https://doi.org/10.1073/pnas.1911999117>

419 Lynch, E. C., Lummaa, V., Htut, W., & Lahdenperä, M. (2019). Evolutionary significance of
420 maternal kinship in a long-lived mammal. *Philosophical Transactions of the Royal*
421 *Society B: Biological Sciences*, 374(1780), 20180067.
422 <https://doi.org/10.1098/rstb.2018.0067>

423 Maklakov, A. A., & Lummaa, V. (2013). Evolution of sex differences in lifespan and aging:
424 Causes and constraints. *BioEssays*, 35, 717–724.
425 <https://doi.org/10.1002/bies.201300021>

426 Monaghan, P., Charmantier, A., Nussey, D. H., & Ricklefs, R. E. (2008). The evolutionary
427 ecology of senescence. *Functional Ecology*, 22(3), 371–378.
428 <https://doi.org/10.1111/j.1365-2435.2008.01418.x>

429 Mumby, H. S., Chapman, S. N., Crawley, J. A. H., Mar, K. U., Htut, W., Thura Soe, A., Aung,
430 H. H., & Lummaa, V. (2015). Distinguishing between determinate and indeterminate

431 growth in a long-lived mammal. *BMC Evolutionary Biology*, 15(1), 214.

432 <https://doi.org/10.1186/s12862-015-0487-x>

433 Mumby, H. S., Mar, K. U., Thitaram, C., Courtiol, A., Towiboon, P., Min-Oo, Z., Htut-Aung, Y.,

434 Brown, J. L., & Lummaa, V. (2015). Stress and body condition are associated with

435 climate and demography in Asian elephants. *Conservation Physiology*, 3(1), cov030.

436 <https://doi.org/10.1093/conphys/cov030>

437 Nussey, D. H., Kruuk, L. E. B., Morris, A., & Clutton-Brock, T. H. (2007). Environmental

438 conditions in early life influence ageing rates in a wild population of red deer. *Current*

439 *Biology*, 17(23), R1000–R1001. <https://doi.org/10.1016/j.cub.2007.10.005>

440 Nussey, D. H., Kruuk, L., Morris, A., Clements, M., Pemberton, J., & Clutton-Brock, T. (2009).

441 Inter- and Intrasexual Variation in Aging Patterns across Reproductive Traits in a Wild

442 Red Deer Population. *The American Naturalist*, 174, 342–357.

443 <https://doi.org/10.1086/603615>

444 Oo, Z. M. (2010). The training methods used in Myanma Timber Enterprise. *Gajah*, 33,

445 58–61.

446 Prospective Studies Collaboration. (2009). Body-mass index and cause-specific mortality in

447 900 000 adults: Collaborative analyses of 57 prospective studies. *The Lancet*,

448 373(9669), 1083–1096. [https://doi.org/10.1016/S0140-6736\(09\)60318-4](https://doi.org/10.1016/S0140-6736(09)60318-4)

449 R Core Team. (2020). *R: A language and environment for statistical computing*. R

450 Foundation for Statistical Computing. <https://www.R-project.org>

451 Rando, T. A., & Wyss-Coray, T. (2021). Asynchronous, contagious and digital aging. *Nature*

452 *Aging*, 1(1), 29–35. <https://doi.org/10.1038/s43587-020-00015-1>

453 Seltmann, M. W., Helle, S., Htut, W., & Lahdenperä, M. (2019). Males have more aggressive

454 and less sociable personalities than females in semi-captive Asian elephants.

455 *Scientific Reports*, 9(1), 2668. <https://doi.org/10.1038/s41598-019-39915-7>

456 Simons, M. J. P., Briga, M., & Verhulst, S. (2016). Stabilizing survival selection on

457 presenescent expression of a sexual ornament followed by a terminal decline.

458 *Journal of Evolutionary Biology*, 29(7), 1368–1378. <https://doi.org/10.1111/jeb.12877>

- 459 Simpson, G. L., & Singmann, H. (2021). *gratia*: Graceful 'ggplot'-based graphics and other
460 functions for GAMs fitted using « mgcv » (0.6.0) [Computer software].
- 461 Sukumar, R. (2003). *The living elephants*: Evolutionary ecology, behavior, and
462 conservation. Oxford University Press.
- 463 Tidière, M., Gaillard, J.-M., Müller, D. W. H., Lackey, L. B., Gimenez, O., Clauss, M., &
464 Lemaître, J.-F. (2015). Does sexual selection shape sex differences in longevity and
465 senescence patterns across vertebrates? A review and new insights from captive
466 ruminants. *Evolution*, 69(12), 3123–3140. <https://doi.org/10.1111/evo.12801>
- 467 Toke Gale, U. (1974). *Burmese timber elephant*. Trade Corporation.
- 468 van de Pol, M., & Verhulst, S. (2006). Age-dependent traits: A new statistical model to
469 separate within- and between-individual effects. *The American Naturalist*, 167(5),
470 766–773. <https://doi.org/10.1086/503331>
- 471 van de Pol, M., & Wright, J. (2009). A simple method for distinguishing within- versus
472 between-subject effects using mixed models. *Animal Behaviour*, 77(3), 753–758.
473 <https://doi.org/10.1016/j.anbehav.2008.11.006>
- 474 Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of senescence.
475 *Evolution*, 11(4), 398–411. <https://doi.org/10.2307/2406060>
- 476 Wood, S. (2017). Generalized Additive Models: An Introduction With R. In *Generalized*
477 *Additive Models: An Introduction with R, Second Edition* (2nd éd., Vol. 66).
478 Chapman and Hall/CRC Press. <https://doi.org/10.1201/9781315370279>
- 479 Zaw, U. K. (1997). Utilization of elephants in timber harvesting in Myanmar. *Gajah*, 17, 9–22.
- 480 Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects*
481 *Models and Extensions in Ecology with R*. Springer-Verlag.
482 <https://doi.org/10.1007/978-0-387-87458-6>

Ethics

The study was performed with the permission from the Myanmar Ministry of Natural Resources and Environmental Conservation and following the University of Turku ethical guidelines. Note that the University of Turku does not provide project numbers.

Data accessibility

The dataset supporting the conclusions of this article will be published on a digital repository and made publicly available upon acceptance.

Authors' contributions

Lucas D. Lalande: Formal analysis, Data curation, Writing – Original draft, Writing – Review & Editing, Visualization, **Virpi Lummaa:** Conceptualization, Supervision, **Htoo H. Aung:** Investigation, Resources, **Win Htut:** Investigation, Resources, **U. Kyaw Nyein:** Investigation, Resources, **Vérane Berger:** Conceptualization, Writing – Review and Editing, Supervision, **Michael Briga:** Conceptualization, Writing – Review and Editing, Supervision.

Competing interests

None. We declare we have no competing interests.

Funding

This work was supported by the Academy of Finland [292368], the European Research Council [ERC-2014-CoG 648766] and the Ella & Georg Ehrnrooth Foundation.

Acknowledgements

We thank the Myanma Timber Enterprise (MTE) and the Myanmar Ministry of Natural Resources and Environmental Conservation for their collaboration and all of the MTE staff enabling this study by recording elephant life events and morphology for so long. Particularly, we thank Mu Mu Thein and Nina Aro for their patience in helping translating countless elephant morphological measurements from burmese to english. We also thank Carly Lynsdale for ensuring our english was fluent. We thank *the Academy of Finland, the European Research Council and the Ella & Georg Ehrnrooth Foundation* for fundings.