PRC1 Sustains the Memory of Neuronal Fate Independent of PRC2 Function

Ayana Sawai¹, Sarah Pfennig¹, Milica Bulajić², Alexander Miller¹, Alireza Khodadadi-Jamayran³, Esteban O. Mazzoni², Jeremy S. Dasen^{1*}

¹Neuroscience Institute, Department of Neuroscience and Physiology, NYU School of Medicine, New York, NY 10016, USA

²Department of Biology, New York University, New York, NY 10003, USA

³Applied Bioinformatics Laboratories, Office of Science and Research, NYU School of Medicine, New York, NY 10016, USA

*Corresponding author: jeremy.dasen@nyumc.org

Abstract

Polycomb repressive complexes (PRCs) 1 and 2 maintain stable cellular memories of early fate decisions by establishing heritable patterns of gene repression. PRCs repress transcription through histone modifications and chromatin compaction, but their roles in neuronal subtype diversification are poorly defined. We unexpectedly found that PRC2 is dispensable to preserve the morphogen-induced positional fates of spinal motor neurons (MNs), while PRC1 is essential for the specification of segmentally-restricted subtypes. Mutation of the core PRC1 component *Ring1* in mice leads to increased chromatin accessibility and ectopic expression of a broad variety of fate determinants, including Hox transcription factors, while neuronal class-specific features are maintained. Loss of MN subtype identities in *Ring1* mutants is due to the suppression of Hox networks by derepressed caudal *Hox* genes. These results indicate that PRC1 can function independently of *de novo* PRC2-dependent histone methylation to maintain chromatin topology and transcriptional memory at the time of neuronal differentiation.

Introduction

Accurate control of gene expression is essential for the specification and maintenance of neural fates during development. Studies of cell type-restricted transcription factors have illuminated the mechanisms by which spatial and temporal regulation of gene expression gives rise to identifiable neuronal subtypes (Doe, 2017; Fishell and Kepecs, 2020; Hobert and Kratsios, 2019; Sagner and Briscoe, 2019; Venkatasubramanian and Mann, 2019). A parallel and critical mechanism of gene regulation is through the post-translational modification of histones, which enables and restricts transcription by modifying chromatin structure (Kishi and Gotoh, 2018; Schuettengruber et al., 2017). The Polycomb group (PcG) is key family of histone-associated proteins that play evolutionarily conserved roles in restricting gene expression during development (Blackledge et al., 2015; Gentile and Kmita, 2020; Simon and Kingston, 2009; Soshnikova and Duboule, 2009). In embryonic stem cells, cell fate determinants are repressed through PRC activities, and PRC-associated histone marks are subsequently removed from loci as cells differentiate (Boyer et al., 2006; Farcas et al., 2012; Tavares et al., 2012). PRC repression is maintained through cell division and after differentiation, and is thought to contribute to stable cellular memories of early patterning events (Ciabrelli et al., 2017; Coleman and Struhl, 2017). In vertebrates, much of our knowledge of how PRCs regulate gene expression has emerged from biochemical studies of PcG proteins or from the activity of these factors in proliferating cells. Despite an in depth understanding of the mechanisms of PRC action, how PcG proteins interact with gene regulatory networks in the CNS remains poorly understood.

The specification of neuronal fates in the vertebrate spinal cord provides a tractable system to elucidate the function of PcG proteins, as the pathways that determine identities are well-defined, and the molecular signatures of many subtypes are known (Butler and Bronner, 2015; Sagner and Briscoe, 2019). One neuronal class where fate specification has been closely examined is the spinal MN. A core set of transcription factors, including Hb9, Isl1/2, and Lhx3, determines class-specific features of MNs, including neurotransmitter identity (Shirasaki and Pfaff, 2002). The subsequent diversification of MNs into hundreds of muscle-specific subtypes is achieved through a conserved network of Hox transcription factors differentially expressed along the rostrocaudal axis (Philippidou and Dasen, 2013). During neural tube patterning, opposing gradients of retinoic acid (RA) and fibroblast growth factors (FGFs) provide spinal progenitors

with a positional identity (Bel-Vialar et al., 2002; Dasen et al., 2003; Liu et al., 2001). These morphogens act, in part, by temporally and spatially depleting Polycomb-associated histone marks from *Hox* clusters (Mazzoni et al., 2013; Narendra et al., 2015). As progenitors exit the cell cycle, MNs continue to express *Hox* genes, where they regulate repertoires of subtype-specific genes (Catela et al., 2016; Dasen et al., 2005; Mendelsohn et al., 2017). Although the role of Hox proteins in the CNS are well-characterized (Parker and Krumlauf, 2020), and are known targets of PRC activities (Gentile and Kmita, 2020), the specific contributions of PRC1 and PRC2 to CNS maturation remain unclear, as few studies have directly compared their functions during embryonic development.

Polycomb repression is initiated by PRC2, which methylates histone H3 at lysine-27, permitting recruitment of PRC1 through subunits that recognize this mark, leading to chromatin compaction at genes targeted for repression (Margueron and Reinberg, 2011; Schuettengruber et al., 2017). PRC1 and PRC2 can also exist in a variety of configurations, which may contribute to neuronal subtype-specific activities. The core subunit of PRC1, Ring1, binds to one of six Polycomb group Ring finger (Pcgrf) proteins (Gao et al., 2012). PRC1 containing Pcgrf4 interacts with Cbx proteins (canonical PRC1) which recognize H3K27me3 (Bernstein et al., 2006; Morey et al., 2012). Variant forms of PRC1 containing Rybp can inhibit incorporation of Cbx proteins into PRC1, and bind target loci independent of H3K27me3 (Tavares et al., 2012; Wang et al., 2010). We previously found that a PRC1 component, Pcgrf4, is required to establish rostral boundaries of *Hox* expression in differentiating MNs (Golden and Dasen, 2012). By contrast, deletion of a core PRC2 component, *Ezh2*, from MN progenitors had no apparent impact on fate specification, possibly due to compensation by *Ezh1* (Shen et al., 2008), or through use of variant PRC1 containing Rybp. These observations raise the question of what are the specific roles of canonical PRC1, variant PRC1, and PRC2 during MN subtype diversification.

To determine the function of PRCs during neural differentiation, we removed core components of these complexes from MN progenitors. We found that depletion of Ring1 proteins, essential constituents of all PRC1 isoforms, causes pronounced changes in transcription factor expression and a loss of Hox-dependent MN subtypes. Surprisingly, neither PRC2 nor variant PRC1 activities are required for rostrocaudal patterning at the time of MN differentiation. Deletion of *Ring1* leads to increased chromatin accessibility and derepression of a broad variety

of cell fate determinants, while class-specific features of MNs are preserved. The derepression of caudal *Hox* genes in *Ring1* mutants leads to the suppression of MN subtype diversification programs. These findings indicate that PRC1 can preserve the memory of early patterning events in the CNS.

Results

PRC1 is Essential for Rostrocaudal Patterning During Neuronal Differentiation

To determine the relative contributions of PRC1 and PRC2 to neuronal specification, we analyzed mice in which core subunit-encoding genes were selectively removed from MN progenitors. *Ezh* genes encode the methyltransferase activity of PRC2, while *Eed* is required to enhance this function. We first generated mice in which both *Ezh* genes are conditionally deleted, by breeding *Ezh1* and *Ezh2* floxed lines to *Olig2::Cre* mice, which targets Cre to MN progenitors (*Ezh^{MNA}* mice) (Hidalgo et al., 2012; Su et al., 2003). We confirmed MN-restricted loss of PRC2 activity by examining the pattern of H3K27me3 at E11.5, which was selectively depleted from progenitors and post-mitotic MNs (Figure 1A,B). Expression of Hb9 and *Vacht*, two general markers of MN identity, were grossly unchanged in *Ezh^{MNA}* mice. (Figure 1C,D,G). We next analyzed expression of Hox proteins in *Ezh^{MNA}* mice and found that the MN columnar subtype determinants Hoxc6, Hoxc9, and Hoxc10, were all expressed in their normal domains (Figure 1E,F, Supplementary Figure 1A-C). Similarly, we observed no changes in Hb9, Hoxc6, Hoxc9, and Hoxc10 expression in MNs after removal of the PRC2 component *Eed*, using *Sox1::Cre* (*Eed^{NEA}* mice), which targets Cre to neuroectoderm (Supplementary Figure 1D-H) (Takashima et al., 2007; Yaghmaeian Salmani et al., 2018). Depletion of PRC2 activity from neural progenitors therefore does not affect MN class specification or rostrocaudal patterning.

We next investigated the function of PRC1, which is thought to repress gene expression in a PRC2-dependent manner. We generated mice in which *Ring1B* is conditionally deleted from MN progenitors using *Olig2::Cre* in a global *Ring1A* mutant background (*Ring1^{MNA}* mice) (Cales et al., 2008; del Mar Lorente et al., 2000). In *Ring1^{MNA}* mice, Ring1B protein was selectively removed from progenitors and post-mitotic MNs (Figure 1H,I). The number of Hb9⁺ MNs and pattern of *Vacht* expression were similar to controls in *Ring1^{MNA}* mice (Figure 1J,K,N). By contrast, Hox proteins normally expressed by forelimb-innervating brachial MNs were not

detected, while thoracic and lumbar Hox proteins were ectopically expressed in more rostral MNs (Figure 1L,M,O-T, Supplementary Figure 1I,J,K). Brachial-level Hox proteins (Hoxc4, Hoxa5, Hoxc6 and Hoxc8), were selectively depleted from MNs, while thoracic and lumbar Hox determinants, (Hoxc9, Hoxc10, and Hoxd10) were derepressed in brachial segments (Figure 1Q-T). At thoracic levels Hoxc9 expression was attenuated and Hoxd10 was ectopically expressed (Supplementary Figure 1J). These results show that loss of *Ring1* leads to a marked rostral shift in Hox pattern, without affecting pan-MN molecular features (Figure 1O).

Canonical PRC1 Regulates Hox Expression in MNs

Ring1 can interact with multiple Pcgf proteins, raising the question of which PRC1 configuration contributes to MN patterning. PRC1 containing Pcgf4 is required to establish Hox boundaries in MNs (Golden and Dasen, 2012), but can exist in two alternative configurations, depending on mutually exclusive incorporation of Cbx (canonical PRC1) or Rybp (variant PRC1) (Gao et al., 2012; Tavares et al., 2012). We examined the function of PRC1 isoforms first by manipulating Rybp and Cbx expression in MNs. We hypothesized that if canonical PRC1 regulates Hox expression, then overexpression of Rybp would inhibit binding of Cbx to Ring1, leading to MN phenotypes similar to *Ring1* mutants (Figure 1U). We used chick *in ovo* neural tube electroporation to express Rybp in postmitotic MNs using the *Hb9* promoter (*Hb9::mRybp*). Expression of *mRybp* under *Hb9* led to ectopic Hoxc9 and Hoxc10 expression at brachial levels (Figure 1V). If Rybp acts by displacing Cbx, then elevating Cbx levels should restore normal Hox expression. To test this, we co-electroporated *Hb9::mRybp* and *Hb9::mCbx2* at equivalent plasmid concentration, and no longer observed ectopic Hoxc9 in brachial segments (Figure 1X).

These observations are consistent with canonical PRC1 containing Pcgf4, Cbx, and Ring1 restricting Hox expression in MNs. Rybp is expressed by MNs at the time of their differentiation (Supplementary Figure 1L), raising the possibility that *Rybp* (or its paralog *Yaf2*) also plays regulatory roles during development. We therefore selectively deleted *Rybp* from MNs using *Olig2::Cre* mice in a *Yaf2*^{-/-} background (*Rybp/Yaf2*^{MNA} mice). Combined deletion of *Rybp* and *Yaf2* did not affect MN generation, Hoxc6, Hoxc9, or Hoxc10 expression (Supplementary Figure 1L). These findings indicate that canonical PRC1 maintains appropriate Hox expression, while variant PRC1 and PRC2 do not contribute to rostrocaudal patterning at the time of MN differentiation.

PRC1 is Required for MN Subtype Diversification

Hox transcription factors play central roles in establishing neuronal subtype identities through regulating expression of subtype-specific genes. To determine the consequences of altered *Hox* expression in *Ring1*^{MNA} mice, we analyzed the molecular profiles and peripheral innervation pattern of MNs. A key Hox target in MNs is the transcription factor *Foxp1*, which is essential for the differentiation of limb-innervating lateral motor column (LMC) and thoracic preganglionic column (PGC) neurons (Dasen et al., 2008; Rousso et al., 2008). In *Ring1*^{MNA} mice, *Foxp1* expression is lost from brachial and thoracic MNs, and markedly reduced in lumbar MNs (Figure 2A,B,G, Supplementary Figure 2C). By contrast, in *Ezh*^{MNA}, *Eed*^{NEA} and *Rybp/Yaf2*^{MNA} mice, *Foxp1* expression was maintained, consistent with the preservation of normal *Hox* profiles in these mutants (Supplementary Figure 1L, 2A,B). In *Ring1*^{MNA} mice, expression of the LMC marker *Raldh2* was lost at brachial levels, and the thoracic PGC marker *nNos* was not detected (Figure 2C, D). Moreover, expression of determinants of respiratory phrenic MNs (*Scip*⁺ *Isl1/2*⁺), was markedly depleted, and likely contributes to the perinatal lethality of *Ring1*^{MNA} mice (Supplementary Figure 2D). These results indicate that *Ring1* deletion leads to a loss of genes acting downstream of Hox function in MNs.

We next assessed the impact of *Ring1* deletion on two MN subtypes that are specified independent of Hox function, the hypaxial and median motor columns (HMC and MMC). In *Ring1*^{MNA} mice, MMC neurons (*Lhx3*⁺ *Hb9*⁺) were generated at normal numbers in thoracic segments, while the number of HMC neurons (*Isl1/2*⁺ *Hb9*⁺ *Lhx3*⁻) was increased in thoracic and brachial segments (Figure 2E,F,H, Supplementary Figure 2E,F). The increase in HMC neurons in *Ring1* mutants is likely due to a reversion of presumptive Hox-dependent subtypes to an HMC fate, similar to mice in which Hox function is disrupted (Dasen et al., 2008; Hanley et al., 2016). Mutation in *Ring1* therefore depletes Hox-dependent subtypes, with the remaining MNs having a more ancestral axial identity (Figure 2P).

As we observed a dramatic loss of segmentally-restricted MN subtypes in *Ring1*^{MNA} mice, we next assessed the impact on peripheral innervation pattern. To trace motor axon projections, we crossed *Ring1*^{MNA} mice to an *Hb9::GFP* reporter, in which all MN axons are labelled with GFP. In control mice there are nine primary trajectories of forelimb-innervating motor axon at

E12.5 (Figure 2I,J) (Catela et al., 2016). In *Ring1*^{MNΔ} mice, only three nerve branches, radial, median, and ulnar were visible but appeared prematurely truncated and unbranched (Figure 2K). In the trunk, innervation of sympathetic chain ganglia was lost, consistent with a loss of PGC fates (Figure 2L,M). Projections to dorsal and ventral axial muscles by MMC and HMC subtypes were maintained in *Ring1*^{MNΔ} mice, although axons were shorter than in controls (Figure 2N, O). Loss of *Ring1* therefore causes severe defects in innervation pattern, while the trajectories of axial MNs are relatively spared.

Loss of *Ring1* Causes Broad Derepression of Developmental Fate Determinants

Polycomb proteins regulate diverse aspects of differentiation by restricting gene expression during development. To investigate changes in gene expression after deletion of *Ring1* genes in an unbiased manner, we performed RNAseq on MNs isolated from *Ring1*^{MNΔ} mice. We purified MNs from control and *Ring1*^{MNΔ};*Hb9::GFP* embryos at E12.5 by flow cytometry (n=4 *Ring1* mutants, and n=4 Cre⁻ controls), and performed RNAseq. Because *Ring1* mutants display segment-specific phenotypes, we collected MNs from brachial, thoracic, and lumbar levels and profiled each population independently.

We identified a total of 1001 upregulated and 641 downregulated genes in *Ring1*^{MNΔ} mice (log₂-FC>2, FDR<0.1), with 391 genes upregulated in all three segmental regions, compared to 46 genes that were commonly downregulated (Figure 3A-C, Supplementary Figure 3A,C, Table 1). Thus, many upregulated genes are shared between each segment, while downregulated genes tended to be segment-specific. Strikingly, 76% of the top 100 derepressed genes (by log₂-FC) at brachial levels encode transcription factors, and include genes normally involved in the specification of spinal interneurons (e.g. *Gata3*, *Shox2*, *Lhx2*), brain regions (*FoxG1*, *Six6*, *Lhx8*), and non-neuronal lineages (Figure 3B).

Although a variety of cell fate determinants were derepressed in *Ring1*^{MNΔ} mice, our RNAseq analyses provide further evidence that core molecular determinants of MN class identity are preserved in *Ring1* mutants. We found no significant changes in transcription factors (*Isl1*, *Isl2*, *Hb9*, *Lhx3*, *Lhx4*), guidance molecules (*Slit2*, *Robo1*, *Robo2*, *DCC*) and neurotransmitter genes (*Chat*, *Aclt*) associated with pan-MN features (Figure 3D). There was a modest decrease in expression *Slc18a3* (*Vacht*) (Figure 3D). Expression of genes that mark other excitatory

(*Slc17a7*, *Slc17a6*) or inhibitory (*Slc32a1*, *Gad2*, *Gad1*,) neuronal classes were not markedly changed in *Ring1* mutants (Supplementary Figure 3B). By contrast, expression of genes associated with MN subtype identities was decreased in *Ring1*^{MNA} mice, including genes that mark limb-innervating (e.g *Foxp1*, *Etv4*, *Raldh2*, *Runx1*), thoracic-specific (*Nos1*, *Cyp26b*), and respiratory (*Alcam*, *Ptn*) subtypes (Figure 3E).

To validate these changes in gene expression, we performed mRNA *in situ* hybridization on a subset of upregulated or downregulated genes. We found that *Gata3*, *Lhx8*, and *Lhx2* were markedly upregulated in MNs of *Ring1*^{MNA} mice (Figure 4A-F), while other genes (*Foxg1*, *Six6*, *Pitx2*, *Shox2*, *Nkx2.1*) showed less prominent, but detectable, ectopic expression (Supplementary Figure 4A-J). We also examined expression of previously uncharacterized genes that were downregulated in all three segmental regions. We found *Rgs4*, *Uts2b*, *Vat1*, and *Gabra2* were expressed by MNs of control embryos, and were downregulated in *Ring1* mutants (Figure 4G-L, Supplementary Figure 4K,L). These results indicate that despite the derepression of multiple fate determinants in *Ring1*^{MNA} mice, core features of MN identity are preserved, but that subtype diversification programs are selectively disrupted.

Selective Derepression of Caudal *Hox* genes in *Ring1*^{MNA} mice

As our preliminary analyses revealed altered expression in a subset of *Hox* genes in *Ring1*^{MNA} mice, we further evaluated *Hox* genes in our RNAseq dataset. Consistent with the analysis of *Hox* protein expression, rostral *Hox* genes were reduced in brachial segments of *Ring1*^{MNA} mice, while *Hox10-Hox13* genes were derepressed (Figure 5A, Supplementary Figure 5A-C). We observed derepression of caudal *Hox* genes from each of the four vertebrate *Hox* clusters, including genes not normally detectable in the ventral spinal cord (e.g *HoxB* genes) (Dasen et al., 2005) (Figure 5A, Supplementary Figure 5A-C). The extent of caudal *Hox* derepression correlated with the relative position of genes within a cluster, with *Hox13* genes displaying the most pronounced derepression in *Ring1*^{MNA} mice (by FC), relative to other caudal *Hox* genes (Figure 5B, Supplementary Figure 3C). The marked derepression of *Hox13* genes was observed in each of the three segmental levels we analyzed, while *Hox10* genes were selectively derepressed in brachial and thoracic segments (Figure 5A, Supplementary Figure 5A-C).

To further validate these findings, we analyzed *Hox13* expression by *in situ* hybridization. In controls *Hoxc13* is restricted to sacral segments, while *Hoxb13* is not detected in MNs (Figure 5C). In *Ring1^{MNA}* mice, both *Hoxb13* and *Hoxc13*, were de-repressed in MNs throughout the rostrocaudal axis (Figure 5C). By contrast, *in situ* hybridization of *Hoxc6* and *Hoxc9* expression revealed reduced expression at brachial and thoracic levels, respectively (Supplementary Figure 5C). Thus, ectopic expression of *Hox13* genes in *Ring1* mutants is associated with a loss of rostral *Hox* gene expression.

***Ring1* is Essential to Maintain MN Chromatin Topology**

Our findings indicate that in the absence of *Ring1* genes, a broad variety of cell fate determinants are ectopically expressed in MNs, while only a subset of caudal *Hox* genes are derepressed. As PRC1 restricts gene expression through chromatin compaction, we investigated whether removal of *Ring1* leads to changes in DNA accessibility at derepressed loci. We used Assay for Transposase-Accessible Chromatin with high-throughput sequencing (ATACseq) to identify genomic regions which have gained or lost accessibility in *Ring1^{MNA}* mice. We purified *Hb9::GFP⁺* MNs at E12.5 from control and *Ring1^{MNA}* embryos at brachial, thoracic, and lumbar levels, and performed ATACseq. In control samples, we observed a progressive opening of caudal *Hox* genes in more rostral segments. For example, the accessibility of *Hox9* genes increases from brachial to thoracic segments, while *Hox10* and *Hox11* genes are more accessible in lumbar segments (Supplementary Figure 6A, B).

To determine which genomic regions gained accessibility in *Ring1* mutants, we compared ATACseq profiles between MNs of control and *Ring1^{MNA}* mice. We identified a total of 2,305 loci that gained accessibility in *Ring1^{MNA}* embryos, 324 (14%) of which were common to all three segmental levels (Figure 6A, Table 2). Common genes included caudal *Hox* genes (*Hoxa13*, *Hoxc13*) and other transcription factors (e.g. *Foxg1*, *Lhx2*, *Pitx2*) that were derepressed in our RNAseq analyses (Figure 6C). We also identified 1,264 loci that lost accessibility in *Ring1^{MNA}* mice, 39 (3%) of which were common to all three segments (Figure 6A). Thus, similar to our RNAseq results, loss of *Ring1* leads to increased accessibility in many genes that are shared among each segment, while genes that lose accessibility tend to be segment-specific.

To assess which genes may be directly repressed by PRC1, we examined the overlap between transcripts that were upregulated and loci that gained accessibility in *Ring1^{MNA}* mice. We found that 18% (148/813) of genes that were ectopically expressed at brachial segments also gained accessibility in *Ring1^{MNA}* mice (Figure 6B). We found 19 genes, including *Hoxa13* and *Hoxc13*, were derepressed and gained accessibility in all three segments (Supplementary Figure 6C). Since lumbar segments still retain features of Hox-dependent subtypes, we also compared the overlap between genes that were upregulated and gained accessibility in brachial and thoracic MNs. We identified 73 genes, 56 (77%) of which encode transcription factors, including each of the caudal *Hox* genes we found by RNAseq (Figure 6B, Supplementary Figure 6C). The gain of accessibility at transcription factor-encoding genes was prominent near transcription start sites (Figure 6C), and regions that gained accessibility in *Ring1^{MNA}* neurons correspond to regions shown to be bound by Ring1B in ES cells (Supplementary Figure 6D) (Bonev et al., 2017).

In agreement with our RNAseq analysis, *Hox13* paralogs showed pronounced increases in accessibility, with each of the four *Hox13* genes gaining accessibility in each of the three segmental levels (Figure 6D,E). By contrast, rostral *Hox4-Hox8* genes, which are transcriptionally downregulated in *Ring1* mutants, were not among that targets that lost accessibility (Figure 6D). These results suggest that reduced expression of rostral *Hox4-Hox9* genes in *Ring1* mutants is not due to a loss in chromatin accessibility.

Hox13 Paralogs Represses Rostral *Hox* genes by Engaging Accessible Chromatin Domains

As PRC1-mediated chromatin compaction is essential for *Hox* repression, it is surprising that *Hox4-Hox9* genes are diminished in *Ring1* mutants, without a noticeable reduction in chromatin accessibility at these loci (Supplementary Figure 6D). Because cross-repressive interactions between *Hox* genes themselves are an important regulatory mechanism determining *Hox* boundaries (Philippidou and Dasen, 2013), it is possible that ectopically expressed *Hox13* genes directly repress multiple *Hox* genes in *Ring1^{MNA}* mice. To test this, we first analyzed MNs derived from ES cells (ESC-MNs) in which *Hoxc13* expression can be induced upon doxycycline (Dox) treatment (Bulajic et al., 2020). ESC-MNs differentiated via RA and Sonic Hedgehog agonist are characterized by expression of *Hox4-Hox5* paralogs. RNAseq analyses revealed that *Hoxa4*, *Hoxa5*, *Hoxc4*, and *Hoxc5* expression were markedly reduced after Dox-induced *Hoxc13* expression in MN progenitors, compared to non-induced ESC-MNs (Figure 7A,

Supplementary Figure 7A-C). This repressive effect also appears to be direct, as ChIPseq analysis indicates that Hoxc13 can bind at multiple *Hox* genes (Figure 7A).

Hox13 proteins can target inaccessible chromatin (Bulajic et al., 2020; Desanlis et al., 2020), while repression by the *Drosophila* Hox protein Ubx is associated with chromatin compaction (Loker et al., 2021). To examine the possible effects of Hox13 proteins on local chromatin structure, we compared accessibility at *Hox* loci in both control and *Ring1* mutants, relative to the location of Hoxc13 binding sites. At *Hox4-Hox5* genes, Hoxc13 sites correspond to regions that are accessible in both control and *Ring1* mutant MNs (Figure 7A, Supplementary Figure 7A-C). Hox13 proteins therefore appear to repress rostral *Hox* genes, in part, through binding at pre-existing accessible regions.

To test whether *Hox13* genes can repress multiple *Hox* paralogs *in vivo*, we used chick neural tube electroporation to express *Hox13* genes within brachial, thoracic, and lumbar segments, and assessed *Hox4-Hox10* gene expression. We electroporated *pCAGGs::mHoxa13-ires-nucGFP* or *pCAGGs::mHoxb13-ires-nucGFP* and found that both cell-autonomously repressed expression of brachial, thoracic, and lumbar Hox proteins (Hoxc4, Hoxa5, Hoxc6, Hoxc8, Hoxc9, and Hoxc10) (Figure 7B-D). By contrast, misexpression of *Hox13* genes did not affect expression of the general MN markers Isl1/2 and Hb9 (Figure 7B-D). These results indicate that Hox13 proteins can directly repress multiple *Hox* genes, and likely contribute to the MN fate specification defects of *Ring1*^{MNΔ} mice.

Discussion

The Polycomb group encompass a large and diverse family of proteins essential for maintaining epigenetic memory of early patterning events. Classically, PcG-mediated repression is thought to depend on recruitment of PRC1 through recognition of histone methylation marks deposited by PRC2 activity. Although alternative, H3K27me3-independent, mechanisms of PRC1 repression have been described (Gao et al., 2012; Tavares et al., 2012), the relative contribution of PRC1 and PRC2 to neuronal fate specification have not been resolved. In this study, we found that genetic depletion of PRC2 components has no discernable impact on neural class or subtype differentiation programs at the time of MN differentiation. By contrast, a core

PRC1 subunit, Ring1, is required to restrict transcription factor expression in the CNS, maintain neuronal subtype-specific chromatin topology, and determine rostrocaudal boundaries of *Hox* expression. Our findings indicate PRC1 plays an key role in regulating expression of cell fate determinants during MN differentiation, safeguarding neurons from acquiring inappropriate gene regulatory programs.

PRC Functions in Neural Development

Polycomb proteins function in diverse aspects of CNS maturation, including the temporal transition from neurogenesis to gliogenesis, maintenance of adult neural stem cell fates, and restriction of gene expression in mature neurons (Di Meglio et al., 2013; Fasano et al., 2009; Hirabayashi et al., 2009; Molofsky et al., 2003; von Schimmelfmann et al., 2016). The role of PRCs in neural subtype diversification has been challenging to study, in part, due the complexity and dynamic temporal regulation of its constituents. By selectively removing core PRC1 and PRC2 subunits from neural progenitors, we determined the relative contributions of these chromatin-associated complexes to neuronal diversification. We found that PRC1 represses a broad variety of cell fate determinants, with the majority of highly derepressed genes in *Ring1* mutants encoding transcription factors. Ectopic expression of Ring1-regulated targets is associated with a disruption in chromatin topology, resulting in a gain in accessibility of derepressed genes.

Despite the derepression of multiple fate determinants in the absence of PRC1 function, expression of transcription factors that define MN class identity (e.g. Hb9, Isl1/2, and Lhx3), are largely unchanged, and MNs do not acquire characteristics of other neuronal classes. This observation suggests that once a class identity has been established, ectopic expression of fate determinants is insufficient to respecify basic neuronal features, such as neurotransmitter identity. In *C. elegans* the ability of transcription factors to reprogram neuronal class identity requires removal of histone deacetylase proteins (Tursun et al., 2011), suggesting that additional regulatory constraints, such as histone acetylation and/or DNA-methylation, restricts the ability of ectopic transcription factors to interfere with class-specific programs in *Ring1* mutants.

While basic elements of MN fate are maintained in the absence of PRC1, their differentiation into molecularly distinct subtypes is severely compromised. The loss of subtype-

specific programs in brachial and thoracic segments can be attributed to the attenuation of *Hox4-*
Hox9 paralog expression, as mice lacking these *Hox* genes show a similar loss in MN subtype
features (Jung et al., 2010; Jung et al., 2014; Philippidou et al., 2012). Selective loss of *Hox*
function in MNs also appears to account for the observation that the majority of downregulated
genes in *Ring1* mutants are segment-specific.

PRC-Dependent and Independent Mechanisms of Rostrocaudal Patterning in the CNS

We found that removal of PRC1 leads to the derepression and increased accessibility of
caudal *Hox* genes. Loss of PRC1 does not result in a sustained derepression of all *Hox* genes, as
Hox genes normally expressed by brachial- and thoracic-level MNs are attenuated in the absence
of *Ring1*. These findings are reminiscent of the function of PRCs in *Drosophila*, where loss of
PRC function leads to ectopic expression of the caudal *Hox* gene *Abd-b* and repression of *Ubx*
expression (Struhl and Akam, 1985; Struhl and White, 1985), likely reflecting cross-repressive
interactions between caudal Hox proteins and rostral *Hox* genes. Notably, reduced rostral *Hox*
gene expression in *Ring1* mutants is not a reflection of reduced chromatin accessibility at these
targets, consistent with a model in which this form of repression is not associated with global
changes in chromatin structure.

In *Ring1* mutants, *Hox13* genes are derepressed in MNs, and appear to contribute to the
loss of MN subtypes. Derepression of *Hox13* genes alone does not appear to account for all of
the specification defects in *Ring1* mutants, as *Hox10* genes were expressed normally in lumbar
segments. *Hoxc9* can also repress *Hox4-Hox8* genes in MNs through direct binding (Jung et al.,
2010), and likely contributes to the *Ring1* mutant phenotype. Derepression of multiple caudal
Hox paralogs therefore appears to have a cumulative effect on repressing brachial and thoracic
Hox genes. In conjunction with studies in *Drosophila*, these findings indicate a deeply-conserved
mechanism of rostrocaudal patterning, in which PRCs establish and maintain rostral *Hox*
boundaries, while Hox cross-repressive interactions define caudal boundaries.

PRC2-Independent Functions of PRC1 in Maintaining Neuronal Subtype Fate

Studies of neuronal and non-neuronal development provide compelling evidence that
PRC2 activity is required to restrict gene expression in early development (Ezhkova et al., 2009;
Gentile et al., 2019; Snitow et al., 2015; Yaghmaeian Salmani et al., 2018). An unexpected

finding from our studies is that MN subtype differentiation and *Hox* boundaries are maintained after removal of PRC2 activity. Mutation of *Ezh* genes or *Eed* depletes H3K27me3 from progenitors and postmitotic neurons, without appreciably affecting MN generation, *Hox* expression, or downstream Hox effectors. These observations indicate that neural progenitors can acquire independence from continuous PRC2 activity to restrict gene expression, whereas there is an absolute requirement for PRC1 function to maintain appropriate transcription factor expression and chromatin organization.

How does PRC1 restrict expression of fate determinants in the absence of PRC2 activity? One possibility is that residual H3K27me3, below the threshold of histological detection, is transmitted through cell division and is sufficient to recruit canonical PRC1 and maintain target gene repression. As cells divide, newly synthesized histones are presumably devoid of H3K27me3 in PRC2 mutants, leading to replication-coupled dilution of H3K27me3. During neural differentiation, the rate of cell division decreases (Kicheva et al., 2014; Wilcock et al., 2007), potentially limiting H3K27me3 reduction, and enabling PRC1 to bind at target loci, even in the absence of *de novo* H3K27 methylation. Although we observed no MN phenotypes in *Ezh* or *Eed* mutants, later-born oligodendrocytes that derive from the same precursors as MNs have been shown to depend on PRC2 function for their differentiation (Wang et al., 2020). This more pronounced effect on glial development likely reflects further dilution of H3K27me3 through cell division in PRC2 mutants.

Stabilization of PRC1 at repressed loci could also maintain target repression independently of *de novo* H3K27 methylation. Repression by canonical PRC1 has been shown to depend on the formation of phase-separated subnuclear structures (Polycomb bodies) assembled through polymerization of Polyhomeotic-like (Phc) and/or Cbx proteins (Isono et al., 2013; Plys et al., 2019; Tsuboi et al., 2018). Mutation of *Phc2* in mice leads to ectopic expression of *Hox* genes, with *Hox13* genes among the most robustly derepressed targets (Isono et al., 2013). One possibility is that PcG-mediated repression may not depend on anchoring of PRC1 through H3K27me3, but is maintained through Phc-mediated chromatin compaction at specific loci.

We suggest that in the early phases of development PRC2 activity defines the sites of maintained H3K27 tri-methylation, initiates PRC1 recruitment, and restricts transcription factor

expression in MNs. These activities are likely deployed prior to neural progenitor specification, as recent studies have demonstrated that the establishment of rostrocaudal positional identities can be specified before neural induction (Metzis et al., 2018). Patterning morphogens, such as RA and FGF, may initially act on stem cells to establish the pattern of H3K27me3 at *Hox* loci prior to neurogenesis. This early rostrocaudal patterning step may ultimately serve to coordinate *Hox* expression in the neural tube and surrounding mesodermally-derived tissues. The subsequent switch to reliance on PRC1 function could reflect a general mechanism of gene regulation in neuronal subtypes that become terminally differentiated during development.

Materials and Methods

Mouse Genetics

Ezh1 flox (Hidalgo et al., 2012), *Ezh2 flox* (Su et al., 2003), *Olig2::Cre* (Dessaud et al., 2007), *Hb9::GFP* (Arber et al., 1999) mice have been previously described. *Ring1A*^{-/-}; *Ring1B flox* (Cales et al., 2008; del Mar Lorente et al., 2000) and *Rybp flox*; *Yaf2*^{-/-} (Hisada et al., 2012; Rose et al., 2016) mouse lines were generated by microinjection of the mouse ESCs expressing these constructs into the blastocyst followed by implantation of these eggs into pseudopregnant foster female mice. Generation of *Ezh*^{MNA} mice was performed by crossing *Ezh1 flox*, *Ezh2 flox* and *Olig2::Cre*. *Ezh*^{MNA} mice are viable at birth but do not survive beyond P20. Generation of *Ring1*^{MNA} mice was performed by crossing *Ring1A*^{-/-}; *Ring1B flox* and *Olig2::Cre*. *Ring1*^{MNA} mice perish at birth. Generation of *Ring1*^{MNA}; *Hb9::GFP* mice was performed by crossing *Ring1A*^{-/-}; *Ring1B flox*, *Olig2::Cre* mice to *Hb9::GFP* mice. Animal work was approved by the Institutional Animal Care and use Committee of the NYU School of Medicine in accordance to NIH guidelines.

Slide Immunohistochemistry

Embryos were fixed in 4% PFA for 1.5-2 hours at 4°C, washed 5-6 times in cold PBS for 15-30 minutes each wash, and incubated overnight in 30% sucrose. Tissue was embedded in OCT, frozen in dry ice, and sectioned at 16 μm on a cryostat. For antibody staining of sections, slides of cryosections were placed in PBS for 5 minutes to remove OCT. Sections were then transferred to humidified trays and blocked for 20-30 minutes in 0.75 ml/slide of PBT (PBS, 0.1% Triton)

containing 1% Bovine serum albumin (BSA). The blocking solution was replaced with primary staining solution containing antibodies diluted in PBT with 0.1% BSA. Primary antibody staining was performed overnight at 4°C. Slides were then washed three times for 5 minutes each in PBT. Fluorophore-conjugated secondary antibodies were diluted 1:500-1:1000 in PBT and filtered through a 0.2 µm syringe filter. Secondary antibody solution was added to slides (0.75 ml/slide) and incubated at room temperature for 1 hour. Slides were washed three times in PBT, followed by a final wash in PBS. Coverslips were placed on slides using 110 µl of Vectashield (Vector Laboratories).

Antibodies against Hox proteins and MN subtypes were generated as described (Dasen et al., 2008; Dasen et al., 2005). Additional antibodies were used as follows: goat anti-Ring1B (Abcam, 1:2000), rabbit anti-Ring1B (Abcam, 1:5000), rabbit anti-Rybp (Abcam 1:2000), rabbit anti-Yaf2 (Abcam, 1:2000), rabbit anti-Cbx2 (Bethyl, 1:5000) , rabbit anti-H3K27me3 (Cell Signaling, 1:2000).

***In situ* mRNA Hybridization**

Probe templates were generated by RT-PCR and incorporated a T7 promoter sequence in the antisense strand. Total RNA was first extracted from eviscerated E12.5 embryos using TRIzol (Invitrogen). Genes of interest were amplified with the One Taq One-Step RT-PCR kit (NEB) using 1 µg of RNA. After amplification by RT-PCR, a second PCR was performed to incorporate a T7 promoter sequence. Antisense riboprobes were generated using the Digoxigenin-dUTP (SP6/T7) labeling kit (Sigma-Aldrich). For *in situ* hybridization, sections were first dried for 10-15 minutes at room temperature, placed in 4% PFA, and fixed for 10 minutes at room temperature. Slides were then washed three times for 3 minutes each in PBS, and then placed in Proteinase K solution (1 µg/ml) for 5 minutes at room temperature. After an additional PFA fixation and washing step, slides were treated in triethanolamine for 10 minutes, to block positive charges in tissue. Slides were then washed three times in PBS and blocked for 2-3 hours in hybridization solution (50% formamide, 5X SSC, 5X Denhardt's solution, 0.2 mg/ml yeast RNA, 0.1 mg/ml salmon sperm DNA). Prehybridization solution was removed, and replaced with 100 µl of hybridization solution containing 100 ng of DIG-labeled antisense probe. Slides were then incubated overnight (12-16 hours) at 72°C. After hybridization, slides were

transferred to a container with 400 ml of 5X SSC and incubated at 72°C for 20 minutes. During this step, coverslips were removed using forceps. Slides were then washed in 400 ml of 0.2X SSC for 1 hour at 72°C. Slides were transferred to buffer B1 (0.1 M Tris pH 7.5, 150 mM NaCl) and incubated for 5 minutes at room temperature. Slides were then transferred to staining trays and blocked in 0.75 ml/slide of B1 containing 10% heat inactivated goat serum. The blocking solution was removed and replaced with antibody solution containing 1% heat inactivated goat serum and a 1:5000 dilution of anti-DIG-AP antibody (Roche). Slides were then incubated overnight at 4°C in a humidified chamber. The following day, slides were washed 3 times, 5 minutes each, with 0.75 ml/slide of buffer B1. Slides were then transferred to buffer B3 (0.1 M Tris pH 9.5, 100 mM NaCl, 50 mM MgCl₂) and incubated for 5 minutes. Slides were then developed in 0.75 ml/slide of B3 solution containing 3.5 µl/ml BCIP and 3.5 µl/ml NBT for 12-48 hours. After color development, slides were washed in ddH₂O and coverslipped in Glycergel (Agilent). A more detailed in situ hybridization protocol is available on our lab website (<http://www.dasenlab.com>).

Wholemout Immunohistochemistry

For wholemount immunohistochemistry embryos were fixed in PFA for 2 hours, then bleached for 24 hours at 4°C in a 10% H₂O₂, 10% DMSO solution prepared in methanol. Embryos were washed three times for 10 minutes each in methanol, followed by five washes for 10 minutes in PBS. Primary antibodies were diluted in staining solution (5% BSA, 20% DMSO in PBS) and specimens were incubated in staining solution on a rotator overnight at room temperature. Samples were then washed three times for 5 minutes each in PBS, followed by four 1 hour washes in PBS. Specimens were then incubated in secondary antibodies diluted in staining solution overnight at room temperature. Samples were then washed three times for 5 minutes each in PBS, followed by four 1 hour washes in PBS, a single 10 min wash in 50% methanol, and three 20 minute washes in 100% methanol. Samples were transferred to glass depression slides and tissue was cleared by incubating samples in BABB solution (1-part benzyl alcohol: 2-parts benzyl benzoate). Confocal images of embryos were obtained from Z-stacks using Zen software (Zeiss). Further details of wholemount staining protocols are available on our lab website: (<http://www.dasenlab.com>).

***In ovo* chick electroporation**

In ovo electroporation were performed on Hamburger Hamilton (HH) stage 13-14 chick embryos and analyzed at HH stage 24-25. Fertilized chicken eggs (Charles River) were incubated in a humidified incubator at 39°C for 40-48 hours until they reached HH13-14. The top of the egg shell was removed and a 1 µg/µl DNA (150-500 ng/µL expression plasmid and pBKS carrier DNA) containing ~0.02% Fast green was injected into the central canal of the neural tube using a sharpened glass capillary tube. Electrodes (Platinum/Iridium (80%/20%), 250 µm diameter, UEPMBGVNXXND, FHC Inc.) were placed on both sides of the neural tube (4 mm separation) and DNA was electroporated using an ECM 830 electroporator (ECM 830, BTX; 25V, 4 pulses, 50 ms duration, 1 second interval). Eggs were sealed with parafilm and incubated for 48 hours prior to fixation. Results shown in figures are representative of at least 3 electroporated embryos from two or more experiments in which electroporation efficiency in MNs was above 60%.

RNA preparation and Library Preparation

RNA was extracted from FACS purified MNs dissected from E12.5 mouse embryo (10-20,000 cells/segment), using the Arcturus Picopure RNA isolation kit. For on-column DNase treatment, Turbo DNase was used (ambion, AM2238). Each samples used separated bar code for libraries. RNA quality and quantity were measured with an Agilent Picochip using a Bioanalyzer, all samples had quality scores between 9-10 RIN. For library preparation 10ng of total RNA was used to generate cDNA, which was amplified with SMARTer Stranded RNA-Seq kit. 100ng of cDNA were used as input to prepare the libraries (Takara, #634837), and amplified by 10 PCR cycles. Samples were run in four 50-nucleotide paired-end read rapid run flow cell lanes with the Illumina HiSeq 4000 sequencer.

RNAseq Data Analyses

Sequencing reads were mapped to the assembled reference genome (mm10) using the STAR aligner (v2.5.0c) (Dobin et al., 2013). Alignments were guided by a Gene Transfer Format (GTF) file. The mean read insert sizes and their standard deviations were calculated using Picard tools (v.1.126) (<http://broadinstitute.github.io/picard>). The read count tables were generated using HTSeq (v0.6.0) (Anders et al., 2015), normalized based on their library size factors using DEseq2 (Love et al., 2014), and differential expression analysis was performed. The Read Per Million (RPM) normalized BigWig files were generated using BEDTools (v2.17.0) (Quinlan and

Hall, 2010) and bedGraphToBigWig tool (v4). To compare the level of similarity among the samples and their replicates, we used two methods: principal-component analysis and Euclidean distance-based sample clustering. All the downstream statistical analyses and generating plots were performed in R environment (v3.1.1) (<https://www.r-project.org/>).

ATACseq

ATACseq was performed following previously described protocols(Buenrostro et al., 2015). DNA was extracted from purified dissected mouse embryonic MNs. Cells were aliquoted and washed twice in ice-cold 1× PBS. Cell pellets were resuspended in 10 mM Tris (pH 7.4), 10 mM NaCl, 3 mM MgCl₂, 0.1% NP-40 (v/v), 0.1% tween20, 0.01% Digitonin and 1% BSA, centrifuged at 500 g for 5 min at 4°C. Pellets were resuspended in 12.5 µl of 2× tagmentation DNA buffer, 1.25 µl Tn5 (Nextera DNA Sample Preparation Kit, FC-121–1030) and 11.25 µl of water, and incubated at 37°C for 30 min. The sample was purified using the MinElute PCR Purification Kit (Qiagen, 28004). PCR enrichment of the library was performed with custom-designed primers and 2× NEB Master Mix. A qPCR reaction with 1× SYBR Green (Invitrogen), custom-designed primers and 2× NEB Master Mix (New England Labs, M0541) was performed to determine the optimal number of PCR cycles (one third of the maximum measured fluorescence)(Buenrostro et al., 2013). The libraries were purified using the AMPure XP beads (Beckman Coulter, A63880). High Sensitivity DNA ScreenTape (Agilent, 5067-5584) was used to verify the fragment length distribution of the library. Library quantification was performed using the KAPA Library Amplification kit on a Roche LightCycler 480. The libraries were sequenced on an Illumina NovaSeq (100 cycles, paired-end).

ATACseq data analysis

All of the reads from the Sequencing experiment were mapped to the reference genome (mm10) using the Bowtie2 (v2.2.4)(Langmead and Salzberg, 2012) and duplicate reads were removed using Picard tools (v.1.126) (<http://broadinstitute.github.io/picard/>). Low quality mapped reads (MQ<20) were removed from the analysis. The read per million (RPM) normalized BigWig files were generated using BEDTools (v.2.17.0) (Quinlan and Hall, 2010) and the bedGraphToBigWig tool (v.4). Peak calling was performed using MACS (v1.4.2)(Zhang et al., 2008) and peak count tables were created using BEDTools. Differential peak analysis was

performed using DESeq2 (Love et al., 2014). ChIPseeker (v1.8.0)(Yu et al., 2015) R package was used for peak annotations and motif discovery was performed using HOMER (v4.10)(Heinz et al., 2010). ngs.plot (v2.47) and ChIPseeker were used for TSS site visualizations and quality controls. KEGG pathway analysis and Gene Ontology (GO) analysis was performed using the clusterProfiler R package (v3.0.0)(Yu et al., 2012). To compare the level of similarity among the samples and their replicates, we used two methods: principal-component analysis and Euclidean distance-based sample clustering. The downstream statistical analyses and generating plots were performed in R environment (v3.1.1) (<https://www.r-project.org/>).

Statistics

Samples sizes were determined based on previous experience and the number of animals and definitions of N are indicated in the main text and figure legends. In figures where a single representative image is shown, results are representative of at least two independent experiments, unless otherwise noted. No power analysis was employed, but sample sizes are comparable to those typically used in the field. Data collection and analysis were not blind. Graphs of quantitative data are plotted as means with standard error of mean (SEM) as error bars, using Prism 8 (Graphpad) software. Unless noted otherwise, significance was determined using unpaired t-test in Prism 8 software, or using adjusted p-values. Exact p-values are indicated, where appropriate, in the main text, figures, and figure legends.

Data Availability

RNAseq and ATACseq data are available through GEO (GSE175503).

Acknowledgements

We thank Kristen D’Elia, Sara Fenstermacher, Jessica Treisman, and Ed Ziff for discussion and comments on the manuscript, and Rachel Kim and Orly Wapinski for assistance. We thank Isabel Hidalgo and Susana Gonzalez for providing *Ezh1**flox* mice, Alexander Tarakhovsky for *Ezh2**flox* mice, Stefan Thor for *Eed* mutant embryos, Haruhiko Koseki for *Ring1A*^{-/-};*Ring1B**flox* ES cells, and Robert Klose for *Rybp**flox*; *Yaf2*^{-/-} ES cells. We also thank Genome Technology Center and Cytometry and Cell Sorting Laboratory at NYU Langone. This work was supported by NIH NINDS grants T32 GM007238, F31 NS087772 to AS, R01NS100897 to EOM, R35 NS116858, R01 NS062822 and R01 NS097550 to JD.

619

620 **Author Contributions**

621 A.S. and J.S.D. designed experiments, interpreted data, and wrote the paper. S.P. performed in
622 situ hybridizations. A.M. and A.S. performed chick neural tube experiments. M.B. and E.O.M
623 designed and performed mouse ESC experiments. A.S. performed all mouse genetic analyses
624 and performed RNAseq and ATACseq. A. K.-J. analyzed ATACseq and RNAseq data.

625

626 **Declaration of Interests**

627 The authors declare no competing interests.

628

References

- Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31, 166-169.
- Arber, S., Han, B., Mendelsohn, M., Smith, M., Jessell, T.M., and Sockanathan, S. (1999). Requirement for the homeobox gene Hb9 in the consolidation of motor neuron identity. *Neuron* 23, 659-674.
- Bel-Vialar, S., Itasaki, N., and Krumlauf, R. (2002). Initiating Hox gene expression: in the early chick neural tube differential sensitivity to FGF and RA signaling subdivides the HoxB genes in two distinct groups. *Development* 129, 5103-5115.
- Bernstein, E., Duncan, E.M., Masui, O., Gil, J., Heard, E., and Allis, C.D. (2006). Mouse polycomb proteins bind differentially to methylated histone H3 and RNA and are enriched in facultative heterochromatin. *Mol Cell Biol* 26, 2560-2569.
- Blackledge, N.P., Rose, N.R., and Klose, R.J. (2015). Targeting Polycomb systems to regulate gene expression: modifications to a complex story. *Nat Rev Mol Cell Bio* 16, 643-649.
- Bonev, B., Mendelson Cohen, N., Szabo, Q., Fritsch, L., Papadopoulos, G.L., Lubling, Y., Xu, X., Lv, X., Hugnot, J.P., Tanay, A., *et al.* (2017). Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* 171, 557-572 e524.
- Boyer, L.A., Plath, K., Zeitlinger, J., Brambrink, T., Medeiros, L.A., Lee, T.I., Levine, S.S., Wernig, M., Tajonar, A., Ray, M.K., *et al.* (2006). Polycomb complexes repress developmental regulators in murine embryonic stem cells. *Nature* 441, 349-353.
- Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013). Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* 10, 1213-1218.
- Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* 109, 21 29 21-21 29 29.
- Bulajic, M., Srivastava, D., Dasen, J.S., Wichterle, H., Mahony, S., and Mazzoni, E.O. (2020). Differential abilities to engage inaccessible chromatin diversify vertebrate Hox binding patterns. *Development* 147.
- Butler, S.J., and Bronner, M.E. (2015). From classical to current: analyzing peripheral nervous system and spinal cord lineage and fate. *Dev Biol* 398, 135-146.
- Cales, C., Roman-Trufero, M., Pavon, L., Serrano, I., Melgar, T., Endoh, M., Perez, C., Koseki, H., and Vidal, M. (2008). Inactivation of the polycomb group protein Ring1B unveils an antiproliferative role in hematopoietic cell expansion and cooperation with tumorigenesis associated with Ink4a deletion. *Mol Cell Biol* 28, 1018-1028.

663 Catela, C., Shin, M.M., Lee, D.H., Liu, J.P., and Dasen, J.S. (2016). Hox Proteins Coordinate
664 Motor Neuron Differentiation and Connectivity Programs through Ret/Gfr alpha Genes. *Cell Rep*
665 *14*, 1901-1915.

666 Ciabrelli, F., Comoglio, F., Fellous, S., Bonev, B., Ninova, M., Szabo, Q., Xuereb, A., Klopp, C.,
667 Aravin, A., Paro, R., *et al.* (2017). Stable Polycomb-dependent transgenerational inheritance of
668 chromatin states in *Drosophila*. *Nat Genet* *49*, 876-886.

669 Coleman, R.T., and Struhl, G. (2017). Causal role for inheritance of H3K27me3 in maintaining
670 the OFF state of a *Drosophila* HOX gene. *Science* *356*.

671 Dasen, J.S., De Camilli, A., Wang, B., Tucker, P.W., and Jessell, T.M. (2008). Hox repertoires
672 for motor neuron diversity and connectivity gated by a single accessory factor, FoxP1. *Cell* *134*,
673 304-316.

674 Dasen, J.S., Liu, J.P., and Jessell, T.M. (2003). Motor neuron columnar fate imposed by
675 sequential phases of Hox-c activity. *Nature* *425*, 926-933.

676 Dasen, J.S., Tice, B.C., Brenner-Morton, S., and Jessell, T.M. (2005). A Hox regulatory network
677 establishes motor neuron pool identity and target-muscle connectivity. *Cell* *123*, 477-491.

678 del Mar Lorente, M., Marcos-Gutierrez, C., Perez, C., Schoorlemmer, J., Ramirez, A., Magin, T.,
679 and Vidal, M. (2000). Loss- and gain-of-function mutations show a polycomb group function for
680 Ring1A in mice. *Development* *127*, 5093-5100.

681 Desanlis, I., Kherdjemil, Y., Mayran, A., Bouklouch, Y., Gentile, C., Sheth, R., Zeller, R.,
682 Drouin, J., and Kmita, M. (2020). HOX13-dependent chromatin accessibility underlies the
683 transition towards the digit development program. *Nat Commun* *11*, 2491.

684 Dessaud, E., Yang, L.L., Hill, K., Cox, B., Ulloa, F., Ribeiro, A., Mynett, A., Novitch, B.G., and
685 Briscoe, J. (2007). Interpretation of the sonic hedgehog morphogen gradient by a temporal
686 adaptation mechanism. *Nature* *450*, 717-720.

687 Di Meglio, T., Kratochwil, C.F., Vilain, N., Loche, A., Vitobello, A., Yonehara, K., Hrycaj,
688 S.M., Roska, B., Peters, A.H., Eichmann, A., *et al.* (2013). Ezh2 orchestrates topographic
689 migration and connectivity of mouse precerebellar neurons. *Science* *339*, 204-207.

690 Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
691 M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* *29*,
692 15-21.

693 Doe, C.Q. (2017). Temporal Patterning in the *Drosophila* CNS. *Annu Rev Cell Dev Biol* *33*,
694 219-240.

695 Ezhkova, E., Pasolli, H.A., Parker, J.S., Stokes, N., Su, I.H., Hannon, G., Tarakhovsky, A., and
696 Fuchs, E. (2009). Ezh2 orchestrates gene expression for the stepwise differentiation of tissue-
697 specific stem cells. *Cell* *136*, 1122-1135.

698 Farcas, A.M., Blackledge, N.P., Sudbery, I., Long, H.K., McGouran, J.F., Rose, N.R., Lee, S.,
699 Sims, D., Cerase, A., Sheahan, T.W., *et al.* (2012). KDM2B links the Polycomb Repressive
700 Complex 1 (PRC1) to recognition of CpG islands. *eLife* 1, e00205.

701 Fasano, C.A., Phoenix, T.N., Kokovay, E., Lowry, N., Elkabetz, Y., Dimos, J.T., Lemischka,
702 I.R., Studer, L., and Temple, S. (2009). Bmi-1 cooperates with Foxg1 to maintain neural stem
703 cell self-renewal in the forebrain. *Genes Dev* 23, 561-574.

704 Fishell, G., and Kepecs, A. (2020). Interneuron Types as Attractors and Controllers. *Annu Rev*
705 *Neurosci* 43, 1-30.

706 Gao, Z., Zhang, J., Bonasio, R., Strino, F., Sawai, A., Parisi, F., Kluger, Y., and Reinberg, D.
707 (2012). PCGF homologs, CBX proteins, and RYBP define functionally distinct PRC1 family
708 complexes. *Mol Cell* 45, 344-356.

709 Gentile, C., Berlivet, S., Mayran, A., Paquette, D., Guerard-Millet, F., Bajon, E., Dostie, J., and
710 Kmita, M. (2019). PRC2-Associated Chromatin Contacts in the Developing Limb Reveal a
711 Possible Mechanism for the Atypical Role of PRC2 in HoxA Gene Expression. *Dev Cell* 50,
712 184-196 e184.

713 Gentile, C., and Kmita, M. (2020). Polycomb Repressive Complexes in Hox Gene Regulation:
714 Silencing and Beyond: The Functional Dynamics of Polycomb Repressive Complexes in Hox
715 Gene Regulation. *Bioessays* 42, e1900249.

716 Golden, M.G., and Dasen, J.S. (2012). Polycomb repressive complex 1 activities determine the
717 columnar organization of motor neurons. *Genes & development* 26, 2236-2250.

718 Hanley, O., Zewdu, R., Cohen, L.J., Jung, H., Lacombe, J., Philippidou, P., Lee, D.H., Selleri,
719 L., and Dasen, J.S. (2016). Parallel Pbx-Dependent Pathways Govern the Coalescence and Fate
720 of Motor Columns. *Neuron* 91, 1005-1020.

721 Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
722 Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription
723 factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* 38,
724 576-589.

725 Hidalgo, I., Herrera-Merchan, A., Ligos, J.M., Carramolino, L., Nunez, J., Martinez, F.,
726 Dominguez, O., Torres, M., and Gonzalez, S. (2012). Ezh1 is required for hematopoietic stem
727 cell maintenance and prevents senescence-like cell cycle arrest. *Cell Stem Cell* 11, 649-662.

728 Hirabayashi, Y., Suzuki, N., Tsuboi, M., Endo, T.A., Toyoda, T., Shinga, J., Koseki, H., Vidal,
729 M., and Gotoh, Y. (2009). Polycomb limits the neurogenic competence of neural precursor cells
730 to promote astrogenic fate transition. *Neuron* 63, 600-613.

731 Hisada, K., Sanchez, C., Endo, T.A., Endoh, M., Roman-Trufero, M., Sharif, J., Koseki, H., and
732 Vidal, M. (2012). RYBP represses endogenous retroviruses and preimplantation- and germ line-
733 specific genes in mouse embryonic stem cells. *Mol Cell Biol* 32, 1139-1149.

734 Hobert, O., and Kratsios, P. (2019). Neuronal identity control by terminal selectors in worms,
735 flies, and chordates. *Curr Opin Neurobiol* 56, 97-105.

736 Isono, K., Endo, T.A., Ku, M., Yamada, D., Suzuki, R., Sharif, J., Ishikura, T., Toyoda, T.,
737 Bernstein, B.E., and Koseki, H. (2013). SAM domain polymerization links subnuclear clustering
738 of PRC1 to gene silencing. *Dev Cell* 26, 565-577.

739 Jung, H., Lacombe, J., Mazzoni, E.O., Liem, K.F., Jr., Grinstein, J., Mahony, S., Mukhopadhyay,
740 D., Gifford, D.K., Young, R.A., Anderson, K.V., *et al.* (2010). Global control of motor neuron
741 topography mediated by the repressive actions of a single hox gene. *Neuron* 67, 781-796.

742 Jung, H., Mazzoni, E.O., Soshnikova, N., Hanley, O., Venkatesh, B., Duboule, D., and Dasen,
743 J.S. (2014). Evolving Hox activity profiles govern diversity in locomotor systems. *Dev Cell* 29,
744 171-187.

745 Kicheva, A., Bollenbach, T., Ribeiro, A., Valle, H.P., Lovell-Badge, R., Episkopou, V., and
746 Briscoe, J. (2014). Coordination of progenitor specification and growth in mouse and chick
747 spinal cord. *Science* 345, 1254927.

748 Kishi, Y., and Gotoh, Y. (2018). Regulation of Chromatin Structure During Neural
749 Development. *Front Neurosci* 12, 874.

750 Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat*
751 *Methods* 9, 357-359.

752 Liu, J.P., Laufer, E., and Jessell, T.M. (2001). Assigning the positional identity of spinal motor
753 neurons: rostrocaudal patterning of Hox-c expression by FGFs, Gdf11, and retinoids. *Neuron* 32,
754 997-1012.

755 Loker, R., Sanner, J.E., and Mann, R.S. (2021). Ubx orchestrates tissue identity through regional
756 and bidirectional changes to chromatin accessibility. *bioRxiv*, 2021.2001.2015.426863.

757 Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
758 dispersion for RNA-seq data with DESeq2. *Genome Biol* 15, 550.

759 Margueron, R., and Reinberg, D. (2011). The Polycomb complex PRC2 and its mark in life.
760 *Nature* 469, 343-349.

761 Mazzoni, E.O., Mahony, S., Peljto, M., Patel, P., Thornton, S.R., McCuine, S., Reeder, C.,
762 Boyer, L.A., Young, R.A., Gifford, D.K., *et al.* (2013). Saltatory remodeling of Hox chromatin
763 in response to rostro-caudal patterning signals. *Nature Neuroscience* *In press*.

764 Mendelsohn, A.I., Dasen, J.S., and Jessell, T.M. (2017). Divergent Hox Coding and Evasion of
765 Retinoid Signaling Specifies Motor Neurons Innervating Digit Muscles. *Neuron* 93, 792-+.

766 Metzis, V., Steinhäuser, S., Pakanavicius, E., Gouti, M., Stamatakis, D., Ivanovitch, K., Watson,
767 T., Rayon, T., Mousavy Gharavy, S.N., Lovell-Badge, R., *et al.* (2018). Nervous System
768 Regionalization Entails Axial Allocation before Neural Differentiation. *Cell* 175, 1105-1118
769 e1117.

770 Molofsky, A.V., Pardal, R., Iwashita, T., Park, I.K., Clarke, M.F., and Morrison, S.J. (2003).
771 Bmi-1 dependence distinguishes neural stem cell self-renewal from progenitor proliferation.
772 *Nature* 425, 962-967.

773 Morey, L., Pascual, G., Cozzuto, L., Roma, G., Wutz, A., Benitah, S.A., and Di Croce, L. (2012).
774 Nonoverlapping functions of the Polycomb group Cbx family of proteins in embryonic stem
775 cells. *Cell Stem Cell* 10, 47-62.

776 Narendra, V., Rocha, P.P., An, D., Raviram, R., Skok, J.A., Mazzoni, E.O., and Reinberg, D.
777 (2015). CTCF establishes discrete functional chromatin domains at the Hox clusters during
778 differentiation. *Science* 347, 1017-1021.

779 Parker, H.J., and Krumlauf, R. (2020). A Hox gene regulatory network for hindbrain
780 segmentation. *Curr Top Dev Biol* 139, 169-203.

781 Philippidou, P., and Dasen, J.S. (2013). Hox genes: choreographers in neural development,
782 architects of circuit organization. *Neuron* 80, 12-34.

783 Philippidou, P., Walsh, C.M., Aubin, J., Jeannotte, L., and Dasen, J.S. (2012). Sustained Hox5
784 gene activity is required for respiratory motor neuron development. *Nat Neurosci* 15, 1636-1644.

785 Plys, A.J., Davis, C.P., Kim, J., Rizki, G., Keenen, M.M., Marr, S.K., and Kingston, R.E. (2019).
786 Phase separation of Polycomb-repressive complex 1 is governed by a charged disordered region
787 of CBX2. *Genes & development* 33, 799-813.

788 Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
789 genomic features. *Bioinformatics* 26, 841-842.

790 Rose, N.R., King, H.W., Blackledge, N.P., Fursova, N.A., Ember, K.J., Fischer, R., Kessler,
791 B.M., and Klose, R.J. (2016). RYBP stimulates PRC1 to shape chromatin-based communication
792 between Polycomb repressive complexes. *eLife* 5.

793 Rouso, D.L., Gaber, Z.B., Wellik, D., Morrissey, E.E., and Novitch, B.G. (2008). Coordinated
794 actions of the forkhead protein Foxp1 and Hox proteins in the columnar organization of spinal
795 motor neurons. *Neuron* 59, 226-240.

796 Sagner, A., and Briscoe, J. (2019). Establishing neuronal diversity in the spinal cord: a time and
797 a place. *Development* 146.

798 Schuettengruber, B., Bourbon, H.M., Di Croce, L., and Cavalli, G. (2017). Genome Regulation
799 by Polycomb and Trithorax: 70 Years and Counting. *Cell* 171, 34-57.

800 Shen, X., Liu, Y., Hsu, Y.J., Fujiwara, Y., Kim, J., Mao, X., Yuan, G.C., and Orkin, S.H. (2008).
801 EZH1 mediates methylation on histone H3 lysine 27 and complements EZH2 in maintaining
802 stem cell identity and executing pluripotency. *Mol Cell* 32, 491-502.

803 Shirasaki, R., and Pfaff, S.L. (2002). Transcriptional codes and the control of neuronal identity.
804 *Annu Rev Neurosci* 25, 251-281.

805 Simon, J.A., and Kingston, R.E. (2009). Mechanisms of polycomb gene silencing: knowns and
806 unknowns. *Nat Rev Mol Cell Biol* 10, 697-708.

807 Snitow, M.E., Li, S., Morley, M.P., Rath, K., Lu, M.M., Kadzik, R.S., Stewart, K.M., and
808 Morrissey, E.E. (2015). Ezh2 represses the basal cell lineage during lung endoderm development.
809 *Development* 142, 108-117.

810 Soshnikova, N., and Duboule, D. (2009). Epigenetic temporal control of mouse Hox genes in
811 vivo. *Science* 324, 1320-1323.

812 Struhl, G., and Akam, M. (1985). Altered distributions of Ultrabithorax transcripts in extra sex
813 combs mutant embryos of *Drosophila*. *EMBO J* 4, 3259-3264.

814 Struhl, G., and White, R.A. (1985). Regulation of the Ultrabithorax gene of *Drosophila* by other
815 bithorax complex genes. *Cell* 43, 507-519.

816 Su, I.H., Basavaraj, A., Krutchinsky, A.N., Hobert, O., Ullrich, A., Chait, B.T., and
817 Tarakhovsky, A. (2003). Ezh2 controls B cell development through histone H3 methylation and
818 Igh rearrangement. *Nat Immunol* 4, 124-131.

819 Takashima, Y., Era, T., Nakao, K., Kondo, S., Kasuga, M., Smith, A.G., and Nishikawa, S.
820 (2007). Neuroepithelial cells supply an initial transient wave of MSC differentiation. *Cell* 129,
821 1377-1388.

822 Tavares, L., Dimitrova, E., Oxley, D., Webster, J., Poot, R., Demmers, J., Bezstarosti, K., Taylor,
823 S., Ura, H., Koide, H., *et al.* (2012). RYBP-PRC1 complexes mediate H2A ubiquitylation at
824 polycomb target sites independently of PRC2 and H3K27me3. *Cell* 148, 664-678.

825 Tsuboi, M., Kishi, Y., Yokozeki, W., Koseki, H., Hirabayashi, Y., and Gotoh, Y. (2018).
826 Ubiquitination-Independent Repression of PRC1 Targets during Neuronal Fate Restriction in the
827 Developing Mouse Neocortex. *Dev Cell* 47, 758-772 e755.

828 Tursun, B., Patel, T., Kratsios, P., and Hobert, O. (2011). Direct conversion of *C. elegans* germ
829 cells into specific neuron types. *Science* 331, 304-308.

830 Venkatasubramanian, L., and Mann, R.S. (2019). The development and assembly of the
831 *Drosophila* adult ventral nerve cord. *Curr Opin Neurobiol* 56, 135-143.

832 von Schimmelmann, M., Feinberg, P.A., Sullivan, J.M., Ku, S.M., Badimon, A., Duff, M.K.,
833 Wang, Z., Lachmann, A., Dewell, S., Ma'ayan, A., *et al.* (2016). Polycomb repressive complex 2
834 (PRC2) silences genes responsible for neurodegeneration. *Nat Neurosci* 19, 1321-1330.

835 Wang, R., Taylor, A.B., Leal, B.Z., Chadwell, L.V., Ilango, U., Robinson, A.K., Schirf, V.,
836 Hart, P.J., Lafer, E.M., Demeler, B., *et al.* (2010). Polycomb group targeting through different
837 binding partners of RING1B C-terminal domain. *Structure* 18, 966-975.

838 Wang, W., Cho, H., Kim, D., Park, Y., Moon, J.H., Lim, S.J., Yoon, S.M., McCane, M., Aicher,
839 S.A., Kim, S., *et al.* (2020). PRC2 Acts as a Critical Timer That Drives Oligodendrocyte Fate
840 over Astrocyte Identity by Repressing the Notch Pathway. *Cell Rep* 32, 108147.

841 Wilcock, A.C., Swedlow, J.R., and Storey, K.G. (2007). Mitotic spindle orientation distinguishes
842 stem cell and terminal modes of neuron production in the early spinal cord. *Development* *134*,
843 1943-1954.

844 Yaghmaeian Salmani, B., Monedero Cobeta, I., Rakar, J., Bauer, S., Curt, J.R., Starkenberg, A.,
845 and Thor, S. (2018). Evolutionarily conserved anterior expansion of the central nervous system
846 promoted by a common PcG-Hox program. *Development* *145*.

847 Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for comparing
848 biological themes among gene clusters. *OMICS* *16*, 284-287.

849 Yu, G., Wang, L.G., and He, Q.Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP
850 peak annotation, comparison and visualization. *Bioinformatics* *31*, 2382-2383.

851 Zhang, Y., Liu, T., Meyer, C.A., Eeckhoutte, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
852 Myers, R.M., Brown, M., Li, W., *et al.* (2008). Model-based analysis of ChIP-Seq (MACS).
853 *Genome Biol* *9*, R137.

854

Figure 1. Roles of PRC1 and PRC2 in determining of Hox expression in spinal MNs.

(A) Core components of PRC2. (B) Brachial spinal sections showing H3K27me3 is depleted from progenitors (boxed region) and post-mitotic MNs (oval) in E11.5 *Ezh^{MNΔ}* (*Ezh1^{flox/flox}; Ezh2^{flox/flox}; Olig2::Cre⁺*) embryos. (C-D) MNs express Hb9 and *Vacht* in *Ezh^{MNΔ}* mice. (E-F) Brachial Hoxc6 expression is normal in *Ezh^{MNΔ}* mice, and no ectopic Hoxc9 is detected. (G) Quantification of MNs: 54±2 Hb9⁺ MNs per section in brachial controls, versus 60±1 in *Ezh^{MNΔ}* mice, n=6 sections, p=0.0696. (H) Core components of PRC1. (I) Ring1B is selectively removed from progenitors (boxed region) and post-mitotic MNs (oval) in E12.5 *Ring1^{MNΔ}* (*Ring1A^{-/-}; Ring1B^{flox/flox}; Olig2::Cre⁺*) mice. (J-K) *Ring1^{MNΔ}* mice express Hb9 and *Vacht*. (L-M) Hoxc6 is lost from brachial MNs and Hoxc9 is ectopically expressed in *Ring1^{MNΔ}* mice. (N) Quantification of MNs: 73±3 Hb9⁺ MNs in controls, versus 68±2 in *Ring1^{MNΔ}* mice, n=6 sections, p=0.0956. (O) Summary of changes in Hox expression in MNs of *Ring1^{MNΔ}* mice. (P-R) Loss of Hoxc8 and ectopic Hoxc10 and Hoxd10 expression in brachial MNs in *Ring1^{MNΔ}* mice. (S-T) Colabeling of Hoxc6/Hoxc9 and Hoxc4/Hoxc10 in *Ring1^{MNΔ}* mice, showing ectopically expressed caudal Hox proteins in place of rostral Hox protein expression in brachial segments. (U) Schematic of Cbx and Rybp interactions in PRC1. (V) Misexpression of Rybp in postmitotic MNs under *Hb9* in chick leads to ectopic Hoxc9 and Hoxc10 expression in brachial MNs. Bolt symbol indicates electroporated side of spinal cord. (X) Co-expression of Rybp and Cbx fails to induce Hoxc9 in brachial MNs. Panels B-F show brachial sections from E11.5 embryos; I-M, P-T brachial sections from E12.5 embryos; V,X brachial chick sections at HH st25. See also Supplementary Figure 1.

Figure 3. *Ring1* is required to restrict transcription factor expression in spinal MNs.

(A) Venn diagrams showing the number of upregulated (left) and downregulated (right) genes in bracial (Br), thoracic (Th), and lumbar (Lu) MNs upon loss of *Ring1* from MNs by RNAseq (log₂-FC>2, FDR<0.1). (B) Heat map of top 100 upregulated genes (by log₂-FC) in *Ring1*^{MNA} mice and control MNs. Genes shown in blue are *Hox* genes and other transcription factor are shown in red. (C) Heat map of 45 genes downregulated in *Ring1*^{MNA} mice in comparison to control MNs. (D) Plots of DESeq2 counts of genes associated with MN class identity. Each data point shows DESeq2 counts for each sample, and segment-specific counts are plotted together. Expression of *Slc18a3* (*Vacht*) is reduced (padj.=0.026477) in *Ring1*^{MNA}. (E) DESeq2 counts of genes associated with specific MN subtype identities were reduced in *Ring1*^{MNA} mice. Counts for LMC and PMC markers are from Br segments, PGC from Th segments. Black bars shown in graphs indicate mean± SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. See also Supplementary Figure 3.

Figure 4. Analyses of misregulated genes in *Ring1* mutants.

(A,C,E) *In situ* mRNA hybridization of selected upregulated genes from *Ring1^{MNA}* RNAseq. Images show sections of brachial, thoracic, and lumbar segments from E12.5 control and *Ring1^{MNA}* mice. (B,D,F) Graphs of DESeq2 counts for each upregulated gene in each segment. Data points show DESeq2 counts from MNs of individual animals from indicated segments. (G, I, K) Analysis of downregulated genes by *in situ* hybridization. *Rgs4* and *Uts2b* displayed elevated expression in specific segmental levels in controls, suggesting that a subset of the commonly-downregulated genes are also Hox-dependent. *Rgs4* expression is normally elevated in LMC neurons (panel G), while *Uts2b* is elevated in thoracic segments of controls (panel I) (H,J,L) Graphs of DESeq2 counts for each downregulated gene in each segment. See also Supplementary Figure 4.

Figure 5. Derepression of caudal *Hox* genes in *Ring1* mutants.

(A) DESeq2 counts of *HoxC*, *HoxA*, and *HoxB* cluster genes in brachial segments in control and *Ring1*^{MNA} mice showing derepression of caudal *Hox* genes. Grey shaded regions highlight *Hox* genes that are derepressed in *Ring1*^{MNA} mice. *Hoxc9* does not show significant derepression, likely because it is normally expressed by caudal brachial MNs. Black bars shown in graphs indicate mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. (B) Comparison of *Hox9-Hox13* paralog gene derepression in brachial segments. Graph shows absolute fold changes of DESeq2 counts, showing a marked increase for caudal *Hox13* paralogs in *Ring1*^{MNA} mice. Each data point shows individual counts for *Ring1* mutants/average of controls. (C) *In situ* of *Hoxb13* and *Hoxc13* mRNA transcripts in E12.5 embryos. *Hoxb13* is normally not detectable in spinal cord, but is derepressed in MNs in *Ring1*^{MNA} mice. *Hoxc13* transcripts are normally restricted to sacral segments but derepressed in rostral segments in *Ring1*^{MNA} mice. See also Supplementary Figure 5.

Figure 6. *Ring1* is essential for maintaining chromatin topology of cell fate determinants.

(A) Proportional Venn diagram showing the number of genes that gain (left) or lose (right) chromatin accessibility in brachial, thoracic and lumbar segments in *Ring1*^{MNA} mice. (B) Venn diagram of upregulated genes from RNAseq and genes that gained accessibility in ATACseq of brachial and thoracic segments. (C) IGV browser views of selected genes that are upregulated and gained chromatin accessibility in controls and *Ring1*^{MNA} mice in brachial segments. (D) IGV browser views of ATACseq tracks in controls and *Ring1*^{MNA} mice in brachial, thoracic, and lumbar segments. *Hox13* genes gain chromatin accessibility in each segment, while reduced brachial *Hoxc6* expression is not associated with a loss of chromatin accessibility. (E) Comparison of IGV browser views of RNAseq and ATACseq tracks of caudal *HoxC*, *HoxA* and *HoxB* genes in brachial segments. See also Supplementary Figure 6.

Figure 7. Hox13 proteins repress multiple *Hox4-Hox10* paralogs.

(A) Effects of *Hoxc13* induction in ESC-MNs. Top panels show IGV browser views (log scale) of RNAseq (red) and ChIPseq tracks (green) at the *HoxC* cluster. Induced *Hoxc13* binds near multiple *HoxC* genes and represses expression of *Hoxc4* and *Hoxc5*. Bottom panels show ATACseq in brachial control and *Ring1* mutant MNs at the *HoxC* cluster (blue). (B) Misexpression of *mHoxa13-ires-nucGFP* represses more rostral *Hox* genes. *Hoxc4*, *Hoxa5*, *Hoxc6* and *Hoxc8* were analyzed in brachial segments, *Hoxc9* in thoracic segments, and *Hoxc10* in lumbar segments. *Hb9* and *Isl1* images are from brachial segments. (C,D) Quantification of percentage of electroporated cells which retained the expression of indicated *Hox* protein and MN markers upon *Hoxa13* or *Hoxb13* misexpression. Quantified electroporated cells were selected from the ventrolateral spinal cord, where MNs reside. *Hb9* and *Isl1/2* quantification shows percentage of GFP⁺ neurons that express either marker, and are shown for Br segments in panel B, and Br and Th segments in C. Data in graphs shown as mean $\pm$ SEM, averaged from n=3 animals, 4 sections each. See also Supplementary Figure 7.

Supplementary Figure 1. Effects of PRC mutations on MN differentiation.

(A-C) Expression of Hoxc9 in thoracic segments and Hoxc10 in lumbar segments is normal in E11.5 *Ezh^{MNA}* mice (*Eed^{flax/flax}, Sox1::Cre⁺*), and no ectopic thoracic Hoxc6 expression is detected. (D,E) In *Eed^{NEA}* mice (H3K27me3 is depleted throughout the spinal cord at E12.5, while Hb9 expression is unaffected. Some H3K27me3 is detected in the floor plate of *Eed^{NEA}* mice (boxed region). (F-H) Brachial Hoxc6, thoracic Hoxc9, and lumbar Hoxc10 are detected normally in *Eed^{NEA}* mice. (I) Brachial Hoxc4 and Hoxa5 are lost from MNs in *Ring1^{MNA}* mice, and Hoxd10 is ectopically expressed in brachial MNs. (J) Hoxc9 expression is reduced in thoracic segments of *Ring1^{MNA}* mice, and Hoxd10 is expressed. (K) Expression of Hoxc10 is retained at lumbar levels of *Ring1^{MNA}* mice. (L) Analysis of E12.5 *Rybp/Yaf2^{MNA}* mice (*Rybp^{flax/flax}, Yaf2^{-/-}, Olig2::Cre⁺*). Rybp expression is selectively depleted from MNs. Hb9 and indicated Hox proteins are expressed normally.

Supplementary Figure 2. Analyses of MN subtypes in PRC mutant mice.

(A,B) Foxp1 expression is unaffected at brachial, thoracic, and lumbar levels in *Ezh^{MNΔ}* and *Eed^{NEΔ}* mice. (C) Reduced Foxp1 expression in lumbar segment of *Ring1^{MNΔ}* mice. (D) Reduced expression of phrenic motor column (PMC) markers (Scip⁺, Isl1/2⁺) in rostral brachial segments of *Ring1^{MNΔ}* mice. (E) Expression of MMC neuron (Hb9, Lhx3) markers in brachial and lumbar segments of *Ring1^{MNΔ}* mice. (F) Expression of HMC markers (Hb9, Isl1/2) in brachial and lumbar segments of *Ring1^{MNΔ}* mice.

Supplementary Figure 3. RNAseq analyses of *Ring1* mutant mice.

(A) PCA plot of RNAseq samples. (B) Plot of neurotransmitter encoding genes in control and *Ring1*^{MNΔ} mice. Commonly used gene names are shown in parenthesis. Expression of *Slc17a7* is increased in *Ring1*^{MNΔ} mice, although the absolute DESeq2 counts were still relatively low (46.3 ± 7.2 in controls versus 111.1 ± 9 in *Ring1*^{MNΔ} mice, $p=0.00012$). Elevated expression of *Gad1* in controls and mutants may be due to presence of spinal interneurons in sorted samples. (C) Volcano plot of upregulated (red) and downregulated (blue) genes. Three caudal *Hox* genes are indicated.

Supplementary Figure 4. Validation of downregulated genes from *Ring1^{MNΔ}* RNAseq.

(A,C,E,G,I) *In situ* hybridization of selected upregulated genes from the *Ring1^{MNΔ}* RNAseq. Images show sections at brachial, thoracic, and lumbar levels from E12.5 control and *Ring1^{MNΔ}* mice. (B,D,F,H,J) Graphs of DESeq2 counts for each upregulated gene in each segment. (K) Analysis of the downregulated gene *Gabra2* by *in situ* hybridization. (L) Graph of DESeq2 counts for *Gabra2* in each segment.

Supplementary Figure 5. Derepression of caudal *Hox* genes in *Ring1*^{MNA} mice.

(A-C) DESeq2 counts of *HoxA*, *HoxB*, *HoxC*, and *HoxD* gene clusters in brachial (A), thoracic (B), and lumbar (C) segments. Gray-shaded regions highlight *Hox* genes that are derepressed in MNs of *Ring1*^{MNA} mice. In lumbar segments, expression of *Hox10* genes is not reduced in *Ring1* mutants, although *Hox13* genes are derepressed. (D) *In situ* hybridization of *Hoxc6* and *Hoxc9* in *Ring1*^{MNA} mice at brachial (Br) and thoracic (Th) levels at E12.5. In brachial segments, expression of *Hoxc6* is lost from MNs, and *Hoxc9* is expressed, similar to the analyses of Hox protein expression. In thoracic segments, *Hoxc9* expression is reduced in MNs.

Supplementary Figure 6. Integrative analysis of RNA and ATACseq results

(A,B) IGV browser view of ATACseq tracks of the *HoxC* cluster (A) and *HoxA* cluster (B) in control mice. The peaks highlighted with green box indicate regions that gain accessibility in more caudal segments. (C) Venn diagram of upregulated genes (RNAseqUP) and genes that gain chromatin accessibility (ATACseqUP) in all segments. Genes that are upregulated and gained accessibility for all segments (19 genes) or Br and Th segments (73 genes) are listed on the right. *Hox* genes are highlighted in blue, other transcription factors in red. (D) IGV browser view comparing ATACseq tracks from brachial segments (control and *Ring1^{MNA}* mice) with Ring1B ChIPseq tracks from ES and neural progenitor cells (NPC). Regions that gain accessibility in *Ring1* mutants correspond to regions bound by Ring1B in ES cells.

Supplementary Figure 7. Comparison of Hoxc13 binding and accessibility in control and *Ring1^{MNA}* MNs

(A) Effects of *Hoxc13* expression in ESC-MNs at the *HoxA* locus. Top panels show IGV browser views (log scale) of RNAseq (red) and ChIPseq tracks (green). Induced *Hoxc13* represses expression of *Hoxa1-Hoxa5* genes and binds near multiple *HoxA* genes. Bottom panels show ATACseq in brachial control and *Ring1* mutant MNs at the *HoxA* cluster (blue). Bottom panels show ATACseq in brachial control and *Ring1* mutant MNs at the *HoxA* cluster. (B,C) Magnified views of *Hoxc13* binding, RNA expression, and ATACseq results at *Hox4-Hox5* genes in the *HoxA* and *HoxC* clusters.