Genome sequencing of the vermicompost strain *Stenotrophomonas maltophilia* UENF-4GII and population structure analysis of the *S. maltophilia* Sm3 genogroup

Francisnei Pedrosa-Silva^a, Filipe P. Matteoli^a, Hemanoel Passarelli-Araujo^{a,b}, Fabio L. Olivares^{c, d}, Thiago M. Venancio^{a, *}

^a Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF), Brazil; ^b Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil; ^c Núcleo de Desenvolvimento de Insumos Biológicos para a Agricultura (NUDIBA), UENF, Brazil; ^d Laboratório de Biologia Celular e Tecidual, Centro de Biociências e Biotecnologia, UENF, Brazil.

* Corresponding author:

Thiago M. Venancio; Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF); Av. Alberto Lamego 2000, P5 / sala 217; Campos dos Goytacazes, Rio de Janeiro, Brazil. E-mail: thiago.venancio@gmail.com.

ABSTRACT

The *Stenotrophomonas maltophilia* complex (Smc) is a cosmopolitan bacterial group that has been proposed an emergent multidrug-resistant pathogen. Taxonomic studies support the genomic heterogeneity of Smc, which comprises genogroups exhibiting a range of phenotypically distinct strains from different sources. Here, we report the genome sequencing and in-depth analysis of *S. maltophilia* UENF-4GII, isolated from vermicompost. This genome harbors a unique region encoding a penicillin-binding protein (*pbpX*) that was carried by a transposon, as well as horizontally-transferred genomic islands involved in anti-phage defense via DNA modification, and pili glycosylation. We also analyzed all available Smc genomes to investigate genes associated with resistance and virulence, niche occupation, and population structure. *S. maltophilia* UENF-4GII belongs to genogroup 3 (Sm3), which comprises three phylogenetic clusters (PC). Pan-GWAS analysis uncovered 471 environment-associated and 791 PC-associated genes, including antimicrobial resistance (e.g. *blaL1* and *blaR1*) and virulence determinants (e.g. *treS* and *katG*) that provide insights on the resistance and virulence potential of Sm3 strains. Together, the results presented here provide the grounds for more detailed clinical and ecological investigations of *S. maltophilia*.

INTRODUCTION

Vermicomposting is a non-thermophilic biodegradation technique used to manage organic waste [1]. The process involves synergistic interactions between earthworms and microorganisms to biodegrade different organic waste into a humus-like material known as vermicompost, a nutrient-rich organic amendment used to enhance soil microbial diversity and plant development [2, 3]. Vermicomposts harbor soil bacteria from various genera, such as *Bacillus*, *Pseudomonas*, *Serratia*, and *Stenotrophomonas*, which may engage in beneficial interactions with plants [4, 5].

Stenotrophomonas maltophilia is a Gram-negative bacillus found in a wide range of natural habitats, including water sources, soils, rhizospheres, animal microbiotas, including humans [6, 7]. *S. maltophilia* has also been used as part of bioremediation and biocontrol strategies [6, 8]. However, *S. maltophilia* has been reported as a global multidrug resistant opportunistic pathogen associated with significant mortality rates of up to 37.5%, mainly because of bacteremia and respiratory tract infections in severely debilitated, immunosuppressed or chronic lung disease patients [9, 10]. *S. maltophilia* is intrinsically resistant to multiple classes of antibiotics, such as aminoglycosides, carbapenems, and macrolides, posing a therapeutic challenge and delay the administration of proper antibiotics [7].

S. maltophilia present high intraspecific variability [11] and, along with the closely related species *S. pavanii*, comprise the *S. maltophilia* complex (Smc) [12-14]. Phylogenetic studies based on multilocus sequencing typing (MLST) and whole-genome sequencing revealed the organization of Smc members in several genogroups [15-17]. A study using whole-genome multilocus sequence typing (wgMLST) and average nucleotide identity (ANI) analyses showed the genetic organization of Smc in 23 monophyletic genogroups with different virulence and resistance characteristics [18]. Among those, genogroup 3 (Sm3) exhibits a myriad of phenotypically distinct strains from different sources. These strains remain poorly explored, hampering investigations on the genetic determinants underlying the physiology and niche occupation of *S. maltophilia* isolates.

Here we present the genome sequencing of *S. maltophilia* UENF-4GII, the first *S. maltophilia* isolated from vermicompost. *S. maltophilia* UENF-4GII belongs to Sm3 and harbors a set of interesting horizontally-transferred regions, including genomic islands (GIs) involved in phage resistance via DNA modification, and pili glycosylation. ANI analysis and SNP-based phylogenetic reconstructions allowed us to reclassify six publicly available *Stenotrophomonas* spp. genomes as *S. maltophilia* from Sm3. We also used the *S. maltophilia* UENF-4GII genome with those of other 66 *S. maltophilia* isolates to thoroughly characterize of the population structure, virulence, and resistance profiles of Sm3. Finally, a pangenome-wide association study (pan-GWAS) of Sm3 allowed us to identify genes associated with niche occupation and phylogenetic clusters.

76

77 METHODS

78 Bacterial isolation and identification

79 The bacterium was isolated from mature manure vermicompost produced from cattle
80 manure at Universidade Estadual do Norte Fluminense Darcy Ribeiro, Brazil. In
81 summary, serial dilutions (10^{-1} to 10^{-7}) were performed on a solution prepared by
82 adding 10 g of vermicompost in 90 mL of saline ($8.5 \text{ g} \cdot \text{L}^{-1}$ NaCl), followed by shaking for
83 60 min. Then, 100 μL of the final dilution (10^{-7}) were taken and spread on plates
84 containing solid Nutrient Broth (NB) with $8 \text{ g} \cdot \text{L}^{-1}$ of NB and $15 \text{ g} \cdot \text{L}^{-1}$ of agar in 1 L of
85 distilled water. After incubation at 30 °C for 7 days, different colony types could be
86 identified and, for purification, individual colonies were transferred to Petri plates with
87 Dygs solid media acquired from Vetec (São Paulo, Brazil). After isolation and purification
88 on Dygs solid medium, a yellowish, circular, convex elevation, punctiform and smooth
89 surface bacterial colony was selected. Light microscopy revealed a Gram-negative strain
90 and the presence of rod-shaped motile was confirmed under phase contrast microscopy.
91 This isolate, named UENF-4GII, was stored in a 16 mL glass flask containing 5 mL of
92 Nutrient Broth solid medium covered with mineral oil and later grown in liquid Dygs
93 medium under rotatory shaker at 150 rpm and 30 °C for 36 h.

94

95 Genome sequencing and annotation

96 Genomic DNA was extracted using QIAamp® DNA Mini Kit (Qiagen) and quantified with
97 an Agilent Bioanalyzer 2100 instrument (Agilent, California, USA). Paired-end libraries
98 were previously prepared with the TruSeq Nano DNA LT Library Prep (Illumina) and
99 sequenced on an Illumina HiSeq 2500 sequencing system at the Life Sciences Core
100 Facility (LaCTAD; UNICAMP, Campinas, Brazil). Sequencing reads (2 x 100 bp) were
101 assembled *de novo* with SPAdes v.10.3.1 [19] and scaffolded with Gfinisher v.1.4 [20],
102 using as references the complete genome of *S. maltophilia* JV3 (GCF_000223885.1) and
103 an alternative assembly generated with Velvet 1.2.10 [21]. The assembly statistics were
104 assessed with QUAST v.3.0 [22]. Genome completeness was assessed with BUSCO v.4.0
105 [23], using the *Gammaproteobacteria* dataset as reference. PlasmidSpades [24] was
106 used to predict plasmids. The assembled genome was annotated with the NCBI
107 Prokaryotic Genome Annotation Pipeline (PGAP) [25]. The UENF-4GII genome was
108 deposited into DDBJ/EMBL/GenBank under the accession number JABUNQ000000000.
109 Genes involved in antimicrobial metabolite biosynthesis were predicted using
110 antiSMASH [26]. Insertion sequences (ISs) and GIs were predicted with ISEscan v.1.5.4
111 [27] and Islandviewer4 [28], respectively. Bacteriophage signatures were analyzed with
112 PHASTER [29].

113

Genome similarity assessment

We downloaded 641 genomes of *Stenotrophomonas* available in the RefSeq database in January, 2021 [30, 31]. Genome completeness was assessed with BUSCO 4.0 [23], using quality score $\geq 90\%$ and the *Gammaproteobacteria* dataset as reference. We excluded assemblies with more than 500 contigs. A preliminary genome distance estimation analysis of isolate UENF-4GII and RefSeq genomes was performed using Mash [32] and a distance tree was generated using MASHtree v.0.50 [33]. All-against-all ANI based on MUMmer alignment (ANIm) was performed with pyani v.0.27 [34]. To assess the concordance between ANI and Mash estimates, we performed a linear correlation analysis using Pearson's correlation coefficient. For this analysis, the Mash distances were converted into Mash scores (1-Mash distance) to allow a direct comparison with ANI values.

Pangenome analysis

The Sm3 genogroup pangenome was performed with Roary v.3.6, using 95% identity threshold to determine orthologous [35]. Core genes were aligned with MAFFT v.7.394 [36]. SNPs were extracted from the core-genome alignment using SNP-sites v.2.3.3 [37] and maximum likelihood phylogenetic reconstructions were performed with IQ-tree [38], with ascertainment bias correction under the model GTR+ASC. The bootstrap support was evaluated using the ultrafast bootstrap method with 1000 replicates [39]. The resulting phylogenetic tree was visualized with iTOL v4 [40].

As a complementary approach, we performed a pangenome-wide association study (pan-GWAS) on the Sm3 dataset using Scoary [92]. The pan-GWAS was computed using the Roary output to find genes associated with isolation source (clinical or non-clinical) and to establish which genes were typical of each phylogenetic cluster (PC), while correcting for population structure using the core-genome phylogenetic tree (command `-n tree`). False-discovery rate was estimated by Benjamini–Hochberg adjusted p-value provided in Scoary. We only reported the results with specificity $> 70\%$ and Benjamini–Hochberg corrected p-value < 0.05 . The binary heatmaps of trait-associated genes were rendered using R package tidyverse [41].

Virulome and resistome analysis

Virulence and antimicrobial resistance genes were predicted using Usearch v.11.0.667 screened against the Virulence Factors of Pathogenic Bacteria Database (VFDB) and the Comprehensive Antibiotic Resistance Database (CARD). Minimum identity and coverage thresholds of 60% and 50% were used in these searches, respectively. The

presence/absence profiles of virulence and resistance-associated genes were rendered using R package tidyverse [41].

RESULTS

Genome analysis of *S. maltophilia* UENF-4GII

During the characterization of culturable bacteria from mature cattle vermicompost, we identified a bacterium that was preliminarily characterized as *Stenotrophomonas* sp. using 16S rRNA sequencing. We submitted the genome of this isolate to whole-genome sequencing (see methods for details). The 30,093,445 paired-end reads were assembled into 3 scaffolds, with an N₅₀ of 1.3 Mbp, encompassing a total of 4.4 Mbp with 66.55% GC content. No plasmids were detected. BUSCO assessment recovered 437 (96.68%) single copy genes of *Gammaproteobacteria* dataset, supporting the completeness and high quality of the assembled genome. The genome harbors 3,932 protein-coding genes, 70 tRNA genes, and 2 rRNA operons.

The *S. maltophilia* UENF-4GII genome carries 18 complete IS elements: nine of the IS481 family, seven of the IS3 family and a single copy of the IS5 and IS21 families (Supplementary figure S1). In addition, we also found an incomplete prophage region that is closely related with a temperate bacteriophage from *Burkholderia pseudomallei* (Burkho_phi1026b) [42]. AntiSMASH prediction revealed that UENF-4GII possesses four gene clusters involved in the production of two bacteriocin-like compounds, an aryl polyene related to Xanthomonadin [43], and a non-ribosomal peptide synthetase (NRPS) gene cluster involved in the biosynthesis of siderophore (Supplementary figure S1). Siderophores are the most important iron uptake systems in *S. maltophilia* [44]. In NRPS gene cluster, we identified six genes (HRE58_11480 to 11500, and HRE58_11505) encoding an specific siderophore of *S. maltophilia* [45]. These enzymes are similar to those involved in enterobactin biosynthesis in enteric bacteria (e.g. *E. coli*) [44].

Genome similarity of UENF-4GII

In order to investigate the genomic relatedness of UENF-4GII strain, we downloaded all available *Stenotrophomonas* genomes in Refseq database (n=641, January 2021), out of which 627 were retained after BUSCO quality filtering (see method for details). Genome identity was computed using Mash and ANI. The mash-distance phylogeny highlights the diversity of the *Stenotrophomonas* genus, corroborating the *S. maltophilia* complex organization in different genogroups (Figure 1A), as previously described by Gröschel et al. [18]. We identified 17 genogroups according to the distance estimation of well-characterized representative genomes, including the affiliation of UENF-4GII to Sm3, along with other 67 clinical and non-clinical strains (Supplementary table S1). In

addition, this analysis allowed us to reclassify six *Stenotrophomonas* spp. as *S. maltophilia* Sm3 isolates, including the environmental strains *Stenotrophomonas* sp. DDT-1 (GCF_001580555.1) and *Stenotrophomonas* sp. Pemsol (GCF_003586545.1), reported as specific biodegraders of organochlorine pesticide (DDT) and polycyclic aromatic hydrocarbons (PAH), respectively. Mash results are supported by ANI analysis, which showed genomic identity above 95% within the same genogroup (Figure 1B). Moreover, the strong positive linear correlation ($r = 0.98$, $p\text{-value} < 2.2e-16$) between ANI and Mash results (Figure 1C) confirm a similar level of resolution for species delineation within the *Stenotrophomonas* genus.

Pangenome analysis

The pangenome analysis of the 67 Sm3 isolates comprises 13,380 genes, with 2,754 core genes (i.e. present in at least 95% of the strains), 1,958 shell genes (present in 15% to 95% of the strains), and 8,668 cloud genes (present in up to 15% of the strains). The heap law estimate supports an open pangenome ($\alpha = 0.65$) (Figure 2A), which typically reflects a high genetic diversity through the acquisition of exogenous DNA [46]. The genomic fluidity (ϕ) Sm3 was estimated in 0.18 (± 0.05), indicating that an average of 18% of unique gene families between in a given pair of genomes.

We also performed a maximum-likelihood phylogenetic reconstruction using SNPs extracted from core genes (i.e. core-genome multilocus sequencing typing, cgMLST). This analysis revealed that Sm3 comprises three distinct and highly supported phylogenetic clusters (PC) (Figure 2B). PC1 ($n = 21$) and PC2 ($n = 15$) encompass clinical and non-clinical strains, while PC3 ($n = 31$) has only clinical strains. Interestingly, 10 of the 21 PC1 strains are non-clinical, including *S. maltophilia* UENF-4GII.

We identified 75 unique genes in *S. maltophilia* UENF-4GII (Supplementary table S2), including *pbpX* (HRE58_12720), which encodes a low molecular weight penicillin-binding protein (LMW-PBP). This gene is closely located to other unique genes: a gene with unknown function (HRE58_12725) and two transposase genes (HRE58_12730, HRE58_12735), flanked by a perfect 26-bp inverted repeat (IR) sequence, supporting their acquisition by horizontal gene transfer (Supplementary figure 2). LMW-PBPs are important enzymes involved in cell-wall recycling and considered the major molecular targets for β -lactam antibiotics [47]. Alterations in the structure of LMW-PBPs are associated with reduced susceptibility to penicillin and other β -lactams and can induce the expression of β -lactamases [47-49]. The presence of an LMW-PBP composite transposon likely confers a competitive advantage for *S. maltophilia* UENF-4GII to thrive in microbial communities containing penicillin.

UENF-4GII genomic islands

To further understand genome plasticity, we identified and analyzed 17 putative GIs (GI1-17) in *S. maltophilia* UENF-4GII, which probably represent recent horizontal gene transfers (Supplementary table S3). In addition, we compared these GIs with other Sm3 genomes (Figure 3). Among these regions, GI3 harbors an operon (HRE58_02625-02660) containing genes encoding GalE, HldD, and glycosyltransferases involved in lipopolysaccharide biosynthesis. This operon is present in 68% of the PC3 genomes. GI16 contains a class C beta-lactamase and GI10 have blue-light sensor genes that are absent in PC3 genomes.

Interestingly, 60% of the *S. maltophilia* UENF-4GII unique genes are located within 7 GIs related with DNA metabolism (GI6 and GI17), virulence (GI7 and GI15) and unknown functions (GI2, GI9 and GI14). GI6 has 9 unique genes, including a restriction-modification (R-M) system with two type I restriction endonucleases (HRE58_05925 and HRE58_05935) and a DNA methyltransferase (HRE58_05920). R-M systems defend the bacterial genomes against bacteriophages and other types of exogenous genetic elements [50] and, hence, their acquisition might be adaptive by boosting resistance against phages [51, 52]. The GI17 has 41,433 bp and comprises 14 unique genes, including a DNA polymerase-like protein (HRE58_18840) and an operon containing an 8-oxoguanine DNA glycosylase (HRE58_18870), a nitroreductase (HRE58_18875), a 7-cyano-7-deazaguanine synthase (HRE58_18880; PreQ(0)), and a nucleoside 2-deoxyribosyltransferase (HRE58_18885). These genes are likely part of a repair system for oxidative stress-mediated DNA damage [53]. PreQ(0) might be involved in the insertion of 7-deazaguanine derivatives in the DNA [54], as part of a defense system against foreign DNA and phages [55].

Finally, 9 of 10 genes from GI15 are unique (Supplementary table S3). This island is involved in the biosynthesis and glycosylation of type IV pilin-like proteins. Hence, this island is likely an important virulence factor that mediate bacterial adherence to biotic and abiotic surfaces. Although pilin glycosylation has been associated with immune evasion of pathogenic bacteria [56], a recent study showed its role in the defense against phage infection [57]. Collectively, these GIs indicate that *S. maltophilia* UENF-4GII is equipped with different horizontally-acquired genes that increased its tolerance to foreign DNA.

Pan-GWAS of Sm3 genogroup

We performed a pan-GWAS to identify genes associated with environment type (clinical and non-clinical; Supplementary table S1) and PCs in Sm3 using Scoary. We found 299 and 172 gene clusters associated with clinical and non-clinical environments, respectively (Supplementary figure S3, Supplementary table S4). Interestingly, 58% of the 471 environment-associated genes encode hypothetical proteins with unknown function.

Among the genes associated with clinical environments, we identified genes encoding a chemotaxis response regulator protein (*cheB*), blue-light sensing protein (*bluF*), and trehalose sintase *treS*, a crucial enzyme involved in trehalose biosynthesis. Several bacteria use trehalose as carbon source and stress protectant [58]. Trehalose metabolism has also been associated with the emergence of virulent human pathogens [59, 60]. Interestingly, *treS* was also reported as a significant gene associated with clinical strains of the *S. maltophilia* complex [61]. Moreover, we identified significant associations with genes involved in bacterial resistance, including multidrug efflux pumps (*mdtA* and *mdtC*), macrolide efflux (*macA*), aminoglycoside-modifying enzyme (*aac*), β -lactamases (*blaL1* and *blaL2*), and a β -lactam sensor gene (*blaR1*). *BlaL1* and *BlaL2* are well known metallo- β -lactamases in *Stenotrophomonas*. They hydrolyze almost all β -lactams, are resistant to all clinically available β -lactamase inhibitors [62, 63] and have been shown to increase pathogenicity in clinical settings [63, 64]. Among the genes associated with non-clinical environments, we also found genes involved in bacterial resistance, particularly in macrolide efflux (*macA* and *macB*), as well as three gene clusters involved in heavy metal resistance including copper (*copB*), cobalt/zinc/cadmium (*czcB*), and mercury (*merR1*).

As most non-clinical strains belong to PC1 and PC3 comprised only clinical strains, most of the environment-related genes are also linked with their respective phylogenetic groups. A total of 252, 97 and 442 gene clusters were associated, respectively, with PC1, PC2 and PC3 (Figure 4, Supplementary table S5). The majority (60%) of the PC-associated genes encoded proteins with unknown functions. PC1 was strongly associated (100% of sensitivity and specificity) with 13 gene clusters, including a type II toxin-antitoxin gene (*pasI*) and c-AMP phosphodiesterase encoding gene (*cpdA*) involved in stress resistance [65, 66]. Further, 11 gene clusters were strongly associated to PC2, including *besA*, involved in iron metabolism [67], and *katG*, related with oxidative stress resistance [68]. The PC3 presented 107 associated gene clusters, with 100% sensitivity and specificity. Among these, we identified genes involved in transcriptional regulation (*cueR*, *comR*, *rstA*, and *cysL*), iron metabolism (*besA*), antibiotic efflux (*mdtA* and *mdtC*), and β -lactam resistance (*blaL1*).

The variants found in different traits (e.g. *besA*, *macA* and *blaL1*) are directly associated with genomic diversity of the PCs. A closer look into the *blaL1* variant of PC3 reveals a molecular heterogeneity with 86% mean similarity with *blaL1* from PC1 and PC2 genomes. Although the functional implications of such diversity is not fully understood, variation in *blaL1* have been associated with β -lactam resistance in *S. maltophilia*, which may contribute to its increased prevalence as a nosocomial pathogen [63, 64, 69].

Virulome and resistome analysis of the Sm3 genogroup

In order to understand the pathogenic potential within the Sm3 genogroup, we systematically investigated the distribution of virulence and antimicrobial resistance genes. The Sm3 virulome contains 54 genes (Supplementary table S6), out of which 35 constitute the core virulome. Most (21 out of 35) of the core virulence genes are involved in motility and adherence, including *pilMNOPQ* and the *pilTU* operon, which encode type IV pili subunits [70], and *tuf* (elongation factor Tu), involved in adhesion to host cells and extracellular matrix components [71]. In addition, we also identified seven type II secretion system (T2SS) genes (e.g. *xpsEFG* and *hxcRS*), which promote the export of enzymes during bacterial colonization. Further, the presence of T2SS genes along with other core virulence genes (e.g. *acpXl*, *hemB*, *hemL*, *csrA*, and *icl*), can be involved in infection and immune evasion capacity of Sm3 strains.

The Sm3 accessory virulome comprises 12 low and 7 high frequency genes, respectively (Figure 5). The high frequency genes are associated with adherence (*pinN*), motility (*flmH*), iron uptake (*bauA*), immune evasion (*rmIB*), and stress tolerance (*clpB*, *kata*, and *katG*). The *clpB* gene encodes a heat shock-inducible chaperone required for bacterial tolerance to a variety of stresses, including heat, osmotic and acidic stress [72]. The *kata* gene encodes a catalase that might confer tolerance against hydrogen peroxide-based disinfectants [18, 73] and is absent in all PC2 strains. Nevertheless, PC2 has an alternative catalase-peroxidase gene (*katG*) that is absent in PC1 and PC3. The *katG* gene is associated with oxidative stress control and has different roles in pathogenic bacteria [68, 74]. Four catalases (KatA1, KatA2, KatMn, and KatE) have been described in *S. maltophilia*, out of which KatA has been considered the main factor in conferring tolerance against hydrogen peroxide [75]. The apparent displacement of *kata* by *katG* in PC2 warrants further investigation and might provide insights the response of *S. maltophilia* to oxidative stress.

The resistome analysis revealed 23 antimicrobial resistance genes (Supplementary table S7), out of which 17 comprise the core resistome, including the metallo- β -lactamase BlaL1 and the inducible Ambler class A β -lactamase BlaL2, as described above [76]. Further, antibiotic efflux pumps (e.g. *SmeABC*, *SmeDEF*, *SmeRS*, *OqxAB*, *GolS*, *EmrE*, and *MexK*) represent the main resistance mechanism encoded by the Sm3 core resistome.

The accessory resistome encompasses six genes (Figure 5). No accessory genes were found in PC2, as well as in most PC1 strains, including *S. maltophilia* UENF-4GII. The accessory resistome contains two classes of aminoglycoside-modifying enzymes: aminoglycoside phosphotransferase (encoded by *aph(6)*-Id and *aph(3'')*Ib); and N-aminoglycoside acetyltransferase (encoded by *aac(6')*-Iz and *aac(6')*-Iak). The *aac(6')*-Iz gene was exclusively found in 25 (81%) PC3 genomes, whereas *aac(6')*-Iak and *aph* genes were found in four PC1 genomes. Only two strains (G4S2 and G4S2-1) have the sulfonamide resistance gene *sul2* and the chloramphenicol resistance gene *floR*. In summary, the Sm3 genomes do not present a large repertoire of acquired resistance

genes. The low frequency of sulfonamide resistance genes suggests that this antibiotic class might be a good therapeutic strategy against *S. maltophilia* infections, as previously proposed [18].

CONCLUSION

In this study, we report the sequencing and in-depth analysis of the *S. maltophilia* UENF-4GII genome. This genome harbors a range of exclusive GIs associated with DNA-modification and anti-phage defense systems. Comparative analysis using pairwise genome identity metrics and cgMLST phylogeny show that *S. maltophilia* UENF-4GII belong to the Sm3 genogroup, which comprises three PCs. Using a pan-GWAS approach, we identified 131 genes as significantly associated with specific Sm3 PCs. Further, 471 genes were specifically associated with clinical and non-clinical environments. These genes could be used as biomarkers in future studies. Sm3 genomes comprise a low number of acquired virulence and resistance genes, although the presence of *katG* and aminoglycoside resistance genes were associated with specific PCs. Collectively, our results provide important information regarding *S. maltophilia* genomic diversity that could provide the grounds for more detailed clinical and ecological investigations.

ACKNOWLEDGEMENTS

This work was supported by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ; grants E-26/203.309/2016 and E-26/203.014/2018), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES; Finance Code 001), and Conselho Nacional de Desenvolvimento Científico e Tecnológico. The funding agencies had no role in the design of the study and collection, analysis, and interpretation of data and in writing.

AUTHOR CONTRIBUTIONS

Conceived the study: FP-S, TMV, FLO; Funding and resources: TMV, FLO; Data analysis: FP-S, HP-A; Interpretation of the results: FP-S, FPM, HP-A, TMV; Wrote the manuscript: FP-S, FPM, HP-A, TMV.

FIGURES

Figure 1. Genomic diversity of *Stenotrophomonas* genus. (A) Mash-distance-based phylogeny of *Stenotrophomonas*, built using 627 publicly genomes and that of UENF-4GII (red color). The *S. maltophilia sensu lato* clade is shaded in light blue. (B) Pairwise average nucleotide identity (ANI) calculated with 627 *Stenotrophomonas* genomes. Colors depict the degree of genome identity. (C) Correlation between ANI and Mash methods. (D) Density plot of pairwise ANI within the Sm3 genogroup.

Figure 2. Pangenome and phylogeny of Sm3 genogroup. (A) Number of gene families in the Sm3 pangenome. The cumulative curve (in dark-red) and alpha value of the Heap law less than one (0.65) supports an open pangenome. (B) cgMLST of Sm3 genogroup showing three phylogenetic groups. SNPs extracted from the core genome were used to build a maximum likelihood phylogenetic tree using IQ-tree (see methods for details). Bootstrap values below and above 70% are represented by orange and gray points, respectively.

401
402 **Figure 3. Circular genome representation of 75 unique genes and 17 genomic islands**
403 **(GIs) identified in *S. maltophilia* UENF-4GII in comparison to other 66 Sm3 strains.** The
404 inner ring represents the UENF-4GII genome GC content. Green, blue and gray rings
405 represent genomes from PC1, PC2 and PC3, respectively. The GIs were predicted and
406 classified as metabolic (red), virulence (orange) or unknown (purple). Unique genes
407 predicted within GIs were highlighted according to the GI category.

408

409 **Figure 4. Distribution of PC-associated genes in *S. maltophilia* Sm3 genogroup.** The
410 cgMLST tree is annotated with two strips representing the phylogenetic cluster (PC) and
411 the environment source. The heatmaps represent the presence (dark-gray) or absence
412 (light-gray) of the genes identified by the pan-GWAS pipeline using phylogenetic cluster
413 (PC) as trait. The neon red square highlights the most strongly associated genes found
414 for each trait (i.e. 100% specificity and sensitivity).

415

Figure 5. Acquired virulome and resistome of *S. maltophilia* Sm3 genogroup. The cgMLST tree is annotated with two strips representing the phylogenetic cluster (PC) identified and the environment source. The binary heatmaps represent the presence (dark-gray) or absence (light-gray/light-blue) of the genes identified.

REFERENCES

1. Lim SL, Lee LH, Wu TY: **Sustainability of using composting and vermicomposting technologies for organic solid waste biotransformation: recent overview, greenhouse gases emissions and economic analysis.** *Journal of Cleaner Production* 2016, **111**:262-278.
2. Domínguez J, Aira M, Kolbe AR, Gómez-Brandón M, Pérez-Losada M: **Changes in the composition and function of bacterial communities during vermicomposting may explain beneficial properties of vermicompost.** *Scientific reports* 2019, **9**(1):1-11.
3. Pathma J, Sakthivel N: **Microbial diversity of vermicompost bacteria that exhibit useful agricultural traits and waste management potential.** *SpringerPlus* 2012, **1**(1):26.
4. Pathma J, Sakthivel N: **Molecular and functional characterization of bacteria isolated from straw and goat manure based vermicompost.** *Applied soil ecology* 2013, **70**:33-47.

- 437 5. Matteoli FP, Passarelli-Araujo H, Reis RJA, da Rocha LO, de Souza EM, Aravind L,
438 Olivares FL, Venancio TM: **Genome sequencing and assessment of plant growth-**
439 **promoting properties of a *Serratia marcescens* strain isolated from vermicompost.**
440 *BMC Genomics* 2018, **19**(n. 1):750.
- 441 6. Ryan RP, Monchy S, Cardinale M, Taghavi S, Crossman L, Avison MB, Berg G, Van Der
442 Lelie D, Dow JM: **The versatility and adaptation of bacteria from the genus**
443 ***Stenotrophomonas*.** *Nature Reviews Microbiology* 2009, **7**(7):514-525.
- 444 7. Brooke JS: ***Stenotrophomonas maltophilia*: an emerging global opportunistic**
445 **pathogen.** *Clinical microbiology reviews* 2012, **25**(1):2-41.
- 446 8. Mukherjee P, Roy P: **Genomic potential of *Stenotrophomonas maltophilia* in**
447 **bioremediation with an assessment of its multifaceted role in our environment.**
448 *Frontiers in Microbiology* 2016, **7**:967.
- 449 9. Falagas ME, Kastoris AC, Vouloumanou EK, Rafailidis PI, Kapaskelis AM, Dimopoulos G:
450 **Attributable mortality of *Stenotrophomonas maltophilia* infections: a systematic**
451 **review of the literature.** *Future microbiology* 2009, **4**(9):1103-1109.
- 452 10. Zemanick ET, Wagner BD, Robertson CE, Ahrens RC, Chmiel JF, Clancy JP, Gibson RL,
453 Harris WT, Kurland G, Laguna TA: **Airway microbiota across age and disease spectrum**
454 **in cystic fibrosis.** *European Respiratory Journal* 2017, **50**(5).
- 455 11. Gherardi G, Creti R, Pompilio A, Di Bonaventura G: **An overview of various typing**
456 **methods for clinical epidemiology of the emerging pathogen *Stenotrophomonas***
457 ***maltophilia*.** *Diagnostic microbiology and infectious disease* 2015, **81**(3):219-226.
- 458 12. Rhee J-Y, Choi JY, Choi M-J, Song J-H, Peck KR, Ko KS: **Distinct groups and antimicrobial**
459 **resistance of clinical *Stenotrophomonas maltophilia* complex isolates from Korea.**
460 *Journal of medical microbiology* 2013, **62**(5):748-753.
- 461 13. Hauben L, Vauterin L, Moore E, Hoste B, Swings J: **Genomic diversity of the genus**
462 ***Stenotrophomonas*.** *International Journal of Systematic and Evolutionary Microbiology*
463 1999, **49**(4):1749-1760.
- 464 14. Svensson-Stadler LA, Mihaylova SA, Moore ER: ***Stenotrophomonas* interspecies**
465 **differentiation and identification by *gyrB* sequence analysis.** *FEMS microbiology*
466 *letters* 2012, **327**(1):15-24.
- 467 15. Ochoa-Sánchez LE, Vinuesa P: **Evolutionary genetic analysis uncovers multiple species**
468 **with distinct habitat preferences and antibiotic resistance phenotypes in the**
469 ***Stenotrophomonas maltophilia* complex.** *Frontiers in microbiology* 2017, **8**:1548.
- 470 16. Patil PP, Kumar S, Midha S, Gautam V, Patil PB: **Taxonogenomics reveal multiple novel**
471 **genomospecies associated with clinical isolates of *Stenotrophomonas maltophilia*.**
472 *Microbial genomics* 2018, **4**(8).
- 473 17. Steinmann J, Mamat U, Abda EM, Kirchhoff L, Streit WR, Schaible UE, Niemann S, Kohl
474 TA: **Analysis of phylogenetic variation of *Stenotrophomonas maltophilia* reveals**
475 **human-specific branches.** *Frontiers in microbiology* 2018, **9**:806.
- 476 18. Gröschel MI, Meehan CJ, Barilar I, Diricks M, Gonzaga A, Steglich M, Conchillo-Solé O,
477 Scherer I-C, Mamat U, Luz CF: **The phylogenetic landscape and nosocomial spread of**
478 **the multidrug-resistant opportunist *Stenotrophomonas maltophilia*.** *Nature*
479 *Communications* 2020, **11**(1):1-12.
- 480 19. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
481 Nikolenko SI, Pham S, Prjibelski AD *et al*: **SPAdes: a new genome assembly algorithm**
482 **and its applications to single-cell sequencing.** *Journal of computational biology : a*
483 *journal of computational molecular cell biology* 2012, **19**(5):455-477.
- 484 20. Guizelini D, Raittz RT, Cruz LM, Souza EM, Steffens MB, Pedrosa FO: **GFinisher: a new**
485 **strategy to refine and finish bacterial genome assemblies.** *Scientific reports* 2016,
486 **6**:34963.
- 487 21. Zerbino R, Birney E: **Velvet: algorithms for de novo short read assembly using de**
488 **Bruijn graphs,** vol. 18; 2008.

- 489 22. Gurevich A, Saveliev V, Vyahhi N, Tesler G: **QUAST: quality assessment tool for**
490 **genome assemblies.** *Bioinformatics* 2013, **29**(8):1072-1075.
- 491 23. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM: **BUSCO:**
492 **assessing genome assembly and annotation completeness with single-copy**
493 **orthologs.** *Bioinformatics* 2015, **31**(19):3210-3212.
- 494 24. Antipov D, Hartwick N, Shen M, Raiko M, Lapidus A, Pevzner PA: **plasmidSPAdes:**
495 **assembling plasmids from whole genome sequencing data.** *Bioinformatics* 2016,
496 **32**(22):3380-3387.
- 497 25. Tatusova T, DiCuccio M, Badretin A, Chetvernin V, Nawrocki EP, Zaslavsky L,
498 Lomsadze A, Pruitt KD, Borodovsky M, Ostell J: **NCBI prokaryotic genome annotation**
499 **pipeline.** *Nucleic acids research* 2016, **44**(14):6614-6624.
- 500 26. Medema MH, Blin K, Cimermancic P, de Jager V, Zakrzewski P, Fischbach MA, Weber T,
501 Takano E, Breitling R: **antiSMASH: rapid identification, annotation and analysis of**
502 **secondary metabolite biosynthesis gene clusters in bacterial and fungal genome**
503 **sequences.** *Nucleic acids research* 2011, **39**(suppl_2):W339-W346.
- 504 27. Xie Z, Tang H: **ISEScan: automated identification of insertion sequence elements in**
505 **prokaryotic genomes.** *Bioinformatics* 2017, **33**(21):3340-3347.
- 506 28. Bertelli C, Laird MR, Williams KP, Group SFURC, Lau BY, Hoad G, Winsor GL, Brinkman
507 FS: **IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets.**
508 *Nucleic acids research* 2017, **45**(W1):W30-W35.
- 509 29. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS: **PHASTER: a better,**
510 **faster version of the PHAST phage search tool.** *Nucleic acids research* 2016,
511 **44**(W1):W16-W21.
- 512 30. O'Leary NA, Wright MW, Brister JR, Ciufu S, Haddad D, McVeigh R, Rajput B, Robbertse
513 B, Smith-White B, Ako-Adjei D *et al*: **Reference sequence (RefSeq) database at NCBI:**
514 **current status, taxonomic expansion, and functional annotation.** *Nucleic Acids Res*
515 2016, **44**(D1):D733-745.
- 516 31. Pruitt KD, Tatusova T, Maglott DR: **NCBI reference sequences (RefSeq): a curated non-**
517 **redundant sequence database of genomes, transcripts and proteins.** *Nucleic Acids*
518 *Res* 2007, **35**(Database issue):D61-65.
- 519 32. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM:
520 **Mash: fast genome and metagenome distance estimation using MinHash.** *Genome*
521 *Biology* 2016, **17**(n. 1):132.
- 522 33. Katz L, Griswold T, Carleton H: **Generating WGS Trees with Mashtree.** In: *Poster*
523 *presented at: American Society for Microbiology Conference on Rapid Applied*
524 *Microbial Next-Generation Sequencing and Bioinformatic Pipelines: 2017*; 2017: 8-11.
- 525 34. Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK: **Genomics and taxonomy**
526 **in diagnostics for food security: soft-rotting enterobacterial plant pathogens.**
527 *Analytical Methods* 2016, **8**(1):12-24.
- 528 35. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D,
529 Keane JA, Parkhill J: **Roary: rapid large-scale prokaryote pan genome analysis.**
530 *Bioinformatics* 2015, **31**(22):3691-3693.
- 531 36. Katoh K, Standley DM: **MAFFT multiple sequence alignment software version 7:**
532 **improvements in performance and usability.** *Molecular biology and evolution* 2013,
533 **30**(4):772-780.
- 534 37. Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, Harris SR: **SNP-sites:**
535 **rapid efficient extraction of SNPs from multi-FASTA alignments.** *Microbial Genomics*
536 2016, **2**(4).
- 537 38. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ: **IQ-TREE: a fast and effective**
538 **stochastic algorithm for estimating maximum-likelihood phylogenies.** *Molecular*
539 *biology and evolution* 2014, **32**(1):268-274.

- 540 39. Minh BQ, Nguyen MAT, von Haeseler A: **Ultrafast approximation for phylogenetic**
541 **bootstrap**. *Molecular biology and evolution* 2013, **30**(5):1188-1195.
- 542 40. Letunic I, Bork P: **Interactive Tree Of Life (iTOL) v4: recent updates and new**
543 **developments**. *Nucleic acids research* 2019.
- 544 41. Wickham H, Averick M, Bryan J, Chang W, McGowan LDA, François R, Grolemond G,
545 Hayes A, Henry L, Hester J: **Welcome to the Tidyverse**. *Journal of Open Source*
546 *Software* 2019, **4**(43):1686.
- 547 42. Yuzenkova J, Nechaev S, Berlin J, Rogulja D, Kuznedelov K, Inman R, Mushegian A,
548 Severinov K: **Genome of Xanthomonas oryzae bacteriophage Xp10: an odd T-odd**
549 **phage**. *Journal of molecular biology* 2003, **330**(4):735-748.
- 550 43. Goel AK, Rajagopal L, Nagesh N, Sonti RV: **Genetic locus encoding functions involved**
551 **in biosynthesis and outer membrane localization of xanthomonadin in Xanthomonas**
552 **oryzae pv. oryzae**. *Journal of bacteriology* 2002, **184**(13):3539-3548.
- 553 44. Kalidasan V, Joseph N, Kumar S, Awang Hamat R, Neela VK: **Iron and virulence in**
554 **Stenotrophomonas maltophilia: all we know so far**. *Frontiers in cellular and infection*
555 *microbiology* 2018, **8**:401.
- 556 45. Nas MY, Cianciotto NP: **Stenotrophomonas maltophilia produces an EntC-dependent**
557 **catecholate siderophore that is distinct from enterobactin**. *Microbiology* 2017,
558 **163**(11):1590.
- 559 46. Tettelin H, Medini D: **The pangenome: Diversity, dynamics and evolution of genomes**.
560 2020.
- 561 47. Ealand CS, Machowski EE, Kana BD: **β -lactam resistance: The role of low molecular**
562 **weight penicillin binding proteins, β -lactamases and Id-transpeptidases in bacteria**
563 **associated with respiratory tract infections**. *IUBMB life* 2018, **70**(9):855-868.
- 564 48. Sarkar SK, Chowdhury C, Ghosh AS: **Deletion of penicillin-binding protein 5 (PBP5)**
565 **sensitises Escherichia coli cells to β -lactam agents**. *International journal of*
566 *antimicrobial agents* 2010, **35**(3):244-249.
- 567 49. Huang Y-W, Wang Y, Lin Y, Lin C, Lin Y-T, Hsu C-C, Yang T-C: **Impacts of penicillin**
568 **binding protein 2 inactivation on β -lactamase expression and muropeptide profile in**
569 **Stenotrophomonas maltophilia**. *Msystems* 2017, **2**(4).
- 570 50. Dimitriu T, Szczelkun MD, Westra ER: **Evolutionary Ecology and Interplay of**
571 **Prokaryotic Innate and Adaptive Immune Systems**. *Current Biology* 2020,
572 **30**(19):R1189-R1202.
- 573 51. Zheng Y, Cohen-Karni D, Xu D, Chin HG, Wilson G, Pradhan S, Roberts RJ: **A unique**
574 **family of Mrr-like modification-dependent restriction endonucleases**. *Nucleic acids*
575 *research* 2010, **38**(16):5527-5534.
- 576 52. Tock MR, Dryden DT: **The biology of restriction and anti-restriction**. *Current opinion in*
577 *microbiology* 2005, **8**(4):466-472.
- 578 53. David SS, O'Shea VL, Kundu S: **Base-excision repair of oxidative DNA damage**. *Nature*
579 2007, **447**(7147):941-950.
- 580 54. Hutinet G, Swarjo MA, de Crécy-Lagard V: **Deazaguanine derivatives, examples of**
581 **crosstalk between RNA and DNA modification pathways**. *RNA biology* 2017,
582 **14**(9):1175-1184.
- 583 55. Thiaville JJ, Kellner SM, Yuan Y, Hutinet G, Thiaville PC, Jumpathong W, Mohapatra S,
584 Brochier-Armanet C, Letarov AV, Hillebrand R: **Novel genomic island modifies DNA**
585 **with 7-deazaguanine derivatives**. *Proceedings of the National Academy of Sciences*
586 2016, **113**(11):E1452-E1459.
- 587 56. Smedley JG, Jewell E, Roguskie J, Horzempa J, Syboldt A, Stolz DB, Castirc P: **Influence**
588 **of pilin glycosylation on Pseudomonas aeruginosa 1244 pilus function**. *Infection and*
589 *immunity* 2005, **73**(12):7922-7931.

- 590 57. Harvey H, Bondy-Denomy J, Marquis H, Sztanko KM, Davidson AR, Burrows LL:
591 **Pseudomonas aeruginosa defends against phages through type IV pilus**
592 **glycosylation. *Nature microbiology* 2018, 3(1):47-52.**
- 593 58. Vanaporn M, Titball RW: **Trehalose and bacterial virulence. *Virulence* 2020,**
594 **11(1):1192-1202.**
- 595 59. Collins J, Robinson C, Danhof H, Knetsch C, Van Leeuwen H, Lawley T, Auchtung J,
596 Britton R: **Dietary trehalose enhances virulence of epidemic *Clostridium difficile*.**
597 ***Nature* 2018, 553(7688):291-294.**
- 598 60. Schwarz S, Van Dijck P: **Trehalose metabolism: a sweet spot for *Burkholderia***
599 ***pseudomallei* virulence. *Virulence* 2017, 8(1):5-7.**
- 600 61. Mercier-Darty M, Royer G, Lamy B, Charron C, Lemenand O, Gomart C, Fourreau F,
601 Madec J-Y, Jumas-Bilak E, Decousser J-W: **Comparative Whole-Genome Phylogeny of**
602 **Animal, Environmental, and Human Strains Confirms the Genogroup Organization**
603 **and Diversity of the *Stenotrophomonas maltophilia* Complex. *Applied and***
604 ***Environmental Microbiology* 2020, 86(10).**
- 605 62. Okazaki A, Avison MB: **Induction of L1 and L2 β -lactamase production in**
606 ***Stenotrophomonas maltophilia* is dependent on an AmpR-type regulator.**
607 ***Antimicrobial agents and chemotherapy* 2008, 52(4):1525.**
- 608 63. Mojica MF, Rutter JD, Taracila M, Abriata LA, Fouts DE, Papp-Wallace KM, Walsh TJ,
609 LiPuma JJ, Vila AJ, Bonomo RA: **Population structure, molecular epidemiology, and β -**
610 **lactamase diversity among *Stenotrophomonas maltophilia* isolates in the United**
611 **States. *MBio* 2019, 10(4).**
- 612 64. Han L, Zhang R-M, Jia L, Bai S-C, Liu X-W, Wei R, Cui C-Y, Sun R-Y, Wang M-G, Cai D-T:
613 **Diversity of L1/L2 genes and molecular epidemiology of high-level carbapenem**
614 **resistance *Stenotrophomonas maltophilia* isolates from animal production**
615 **environment in China. *Infection, Genetics and Evolution* 2020, 86:104531.**
- 616 65. Barth E, Gora KV, Gebendorfer KM, Settele F, Jakob U, Winter J: **Interplay of cellular**
617 **cAMP levels, σ S activity and oxidative stress resistance in *Escherichia coli*.**
618 ***Microbiology* 2009, 155(Pt 5):1680.**
- 619 66. Norton JP, Mulvey MA: **Toxin-antitoxin systems are important for niche-specific**
620 **colonization and stress resistance of uropathogenic *Escherichia coli*. *PLoS Pathog***
621 **2012, 8(10):e1002954.**
- 622 67. Miethke M, Klotz O, Linne U, May JJ, Beckering CL, Marahiel MA: **Ferri-bacillibactin**
623 **uptake and hydrolysis in *Bacillus subtilis*. *Molecular microbiology* 2006, 61(6):1413-**
624 **1427.**
- 625 68. Chouchane S, Giroto S, Kapetanaki S, Schelvis JP, Yu S, Magliozzo RS: **Analysis of heme**
626 **structural heterogeneity in *Mycobacterium tuberculosis* catalase-peroxidase (KatG).**
627 ***Journal of Biological Chemistry* 2003, 278(10):8154-8162.**
- 628 69. Kim Y, Maltseva N, Wilamowski M, Tesar C, Endres M, Joachimiak A: **Structural and**
629 **biochemical analysis of the metallo- β -lactamase L1 from emerging pathogen**
630 ***Stenotrophomonas maltophilia* revealed the subtle but distinct di-metal scaffold for**
631 **catalytic activity. *Protein Science* 2020, 29(3):723-743.**
- 632 70. Comolli JC, Hauser AR, Waite L, Whitchurch CB, Mattick JS, Engel JN: ***Pseudomonas***
633 ***aeruginosa* gene products PilT and PilU are required for cytotoxicity in vitro and**
634 **virulence in a mouse model of acute pneumonia. *Infection and immunity* 1999,**
635 **67(7):3625-3630.**
- 636 71. Harvey KL, Jarocki VM, Charles IG, Djordjevic SP: **The diverse functional roles of**
637 **elongation factor Tu (EF-Tu) in microbial pathogenesis. *Frontiers in microbiology***
638 **2019, 10:2351.**
- 639 72. Pan H, Luan J, He X, Lux R, Shi W: **The clpB gene is involved in the stress response of**
640 ***Myxococcus xanthus* during vegetative growth and development. *Microbiology* 2012,**
641 **158(Pt 9):2336.**

- 642 73. Charoenlap N, Jiramonai L, Chittrakanwong J, Tunsakul N, Mongkolsuk S,
643 Vattanaviboon P: **Inactivation of ahpC renders *Stenotrophomonas maltophilia***
644 **resistant to the disinfectant hydrogen peroxide.** *Antonie Van Leeuwenhoek* 2019,
645 **112(5):809-814.**
- 646 74. Chun H, Choi O, Goo E, Kim N, Kim H, Kang Y, Kim J, Moon JS, Hwang I: **The quorum**
647 **sensing-dependent gene katG of *Burkholderia glumae* is important for protection**
648 **from visible light.** *Journal of bacteriology* 2009, **191(13):4152-4157.**
- 649 75. Li L-H, Shih Y-L, Huang J-Y, Wu C-J, Huang Y-W, Huang H-H, Tsai Y-C, Yang T-C:
650 **Protection from hydrogen peroxide stress relies mainly on AhpCF and KatA2 in**
651 ***Stenotrophomonas maltophilia*.** *Journal of biomedical science* 2020, **27(1):1-9.**
- 652 76. Sánchez MB: **Antibiotic resistance in the opportunistic pathogen *Stenotrophomonas***
653 ***maltophilia*.** *Frontiers in microbiology* 2015, **6:658.**

654