Mutagenic effectiveness and efficiency of gamma rays and sodium azide in M₂ generation of Cowpea [*Vigna unguiculata* (L.) Walp.]

Aamir Raina^{1,2*}, Samiullah Khan¹

¹Mutation Breeding Laboratory, Department of Botany, Aligarh Muslim University, Aligarh, India

²Botany Section, Women's College, Aligarh Muslim University, Aligarh, India

Corresponding author: *aamir854@gmail.com

¹²These authors contributed equally to this work

Abstract

Legumes play a pivotal role in combating the chronic hunger and malnutrition in the developing nations and are also ideal crops to achieve global food and nutrition security. In the era of climate change, erratic rainfalls, depleting arable land and water resource, feeding the rapidly growing population is a challenging task. Like most other pulses, cowpea is a self pollinated, nutritious, versatile and widely adapted grain legume, but harbor a little accessible genetic variability. Lack of sufficient genetic variability and small size of flowers, traditional plant breeding methods are not enough to meet the demands of improvement of cowpea. Hence, induced mutagenesis was employed to induce significant genetic variability across a range of agro-economical traits in two cowpea varieties Gomati VU-89 and Pusa-578 from M₁ to M₄ generations. The success of induced mutagenesis largely depends on the selection of appropriate mutagen, its dose, effectiveness and efficiency. Hence present study was conduct to assess the effectiveness and efficiency of single and combined doses of sodium azide and gamma rays to set an appropriate protocol for induced mutagenesis experimentation in cowpea.

1. Introduction

Pulses are considered as vital constituents in the diets of people and hence are most suitable crops to achieve the global food and nutrition security [1]. The production of pulses in India is much lower as compared to global pulse production (Table 1). Additional there is a progressive decline in the per capita pulses consumption from 1951 (60.7 g/day) to 2017 (52.9 g/day) [2]. The low yielding potential of crops is one of the primary obstacles in achieving the desired goals of production. The common asynchronous and longer maturity period, more flower drop and less seed set, natural and anthropogenic pressures are other obstacles in achieving the higher production of pulses.

Table 1. Average yield of pulses in India and world compared during 2007-2016.

YEAR	PULSES YEILD (Q/ha)	
	INDIA	WORLD
2007	5.970	8.154
2008	6.042	8.607
2009	6.706	9.285
2010	6.496	9.043
2011	6.163	8.769
2012	6.460	9.381
2013	7.162	9.745
2014	7.056	9.436
2015	6.465	9.505
2016	6.644	9.929

Source: FAOSTAT | © FAO Statistics Division, 2017.

Among the ten primary pulse crops as recognised by FAO, Cowpea is an important member on the basis of its use, nutritive value and other desired qualities [3-5]. It is suitable for cultivation in subtropical areas of the world and usually grown alone but can be intercropped with a range of other crops including sorghum, millet, maize, cassava and cotton [6]. Cowpea is considered as a vital constituent of traditional cropping system because of its capability to reinstate soil fertility for succeeding cereal crops grown in rotation with it and enhances soil fertility mostly in smallholder farming systems where very less or no fertilizer is used. Cowpea growing countries in the Asian region are India, Sri Lanka, Bangladesh, Myanmar, China, Korea, Thailand, Indonesia, Nepal, Pakistan, Philippines and Malaysia (Fig 1) [7]. Nigeria is the top most producer of cowpea with an estimated 45% of the world cowpea production (Table 2). The objective of this study was to study effect of mutagens on physiological parameters and their mutagenic effectiveness and efficiency.

Fig 1. Cowpea production.

Table 2: World's top 10 producers of Cowpea (Average 2016).

RANK	COUNTRY	PRODUCTION (mt)
1	Nigeria	3.02
2	Niger	1.98
3	Burkina Faso	0.60
4	Cameroon	0.19
5	United Republic of Tanzania	0.18
6	Sudan	0.16
7	Mali	0.14
8	Kenya	0.14
9	Myanmar	0.11
10	Sri Lanka	0.01

Source: FAOSTAT | © FAO Statistics Division, 2017.

2. Materials and methods

2.1. Experimental materials and Seed Irradiation

In the present study, M_0 seeds were obtained from NBPGR, New Delhi (S Table 1). Initially, LD50 was calculated, the doses of gamma rays and sodium azide were optimised by calculating values based on germination rate and survival percentage [6]. A set of 300 dry and healthy M_0 seeds were pre-soaked in the double distilled water for 9 hours and then treated with 0.01%, 0.02%, 0.03% and 0.04% concentrations of sodium azide (SA) solution for 6 hrs at 25 C at Mutation Breeding Laboratory, Department of Botany, Aligarh Muslim University, Aligarh, India. Another set of 600 M_0 seeds per dose were irradiated with 100, 200, 300 and 400 Gy of gamma radiation using a Cobalt 60 (^{60}Co) gamma source under standard conditions at NBRI, Lucknow, India. Each set of irradiated 300 M_0 seeds were also treated with respective SA concentration to produce M_1 seeds. Each set of seeds were washed in running water.

2.2. Experimental site and crop cultivation

The experimentation were performed at Agriculture field, Aligarh Muslim University. The net field size included 13 (S Fig 1). Cowpea M_1 seeds were sown in 10 replications of 30 seeds to raise the M_1 generation during mid-april 2014-October 2014 along with respective controls. Field was irrigated at regular intervals after seed sowing and; weeding was carried out at seedling and flowering stage. All the M_1 plants were harvested separately at pod stage to collect M_2 seeds. During the mid-april 2015-October 2015, 10 healthy M_2 seeds collected from each M_1 plant were sown treatment wise for raising M_2 generation. In M_3 generation seven and four high yielding stable mutant lines of Gomati VU-89 and Pusa-578, respectively were identified and advanced to M_4 generation during the mid-april 2017-October 2017 [8].

All the recommended cultivation practices were taken care of; such as 15 - 20 N, 50 - 60 P₂O₅ and 50 - 60 K₂O kilogram per hectare were mixed with soil.

S1 Fig. Field plots layout of M₁ generations.

(Source: Raina et al., 2020. <http://creativecommons.org/licenses/by/4.0/>).

2.3. Field Analysis

2.3.1. Germination of Seeds

Germination of seeds was noted in control and treated sets and the number of seeds germinated in each treated and untreated population was counted. After recording germination counts, the percentage of seed germination was calculated on the basis of total number of seeds sown in the field using the following formula.

$$\text{Germination (\%)} = \frac{\text{No. of seeds germinated}}{\text{No. of seeds sown}} \times 100$$

2.3.2. Chlorophyll mutations

These mutations were noted immediately after seed sowing. The identification and categorisation proposed by Gustafsson [9] and Khan [10] was followed using the following formula Sharma [11].

$$\text{Mutation frequency (\%)} = \frac{\text{Number of mutant seedlings}}{\text{Total number of M}_2 \text{ seedlings}} \times 100$$

$$\text{Coefficient of interaction (k)} = \frac{(a + b)}{(a) + (b)}$$

2.4. Pollen fertility

The determination of pollen fertility was necessary to assess the pollen sterility based mutagenic effectiveness and efficiency. At the time of flowering, pollens were collected from 30 random plants per treatment and control in M₁ generation. Pollens were then placed on slides and stained with 1% acetocarmine solution. The uniformly stained pollens were counted as fertile, while as unstained pollens were counted as sterile.

2.5. Mutagenic effectiveness and efficiency

The frequency of mutation induced by unit mutagen dose refers to mutagenic effectiveness whereas percentage of mutations in relation to biological damage refers to mutagenic efficiency. Mutagenic effectiveness and efficiency was calculated by following formulae given by Konzak *et al.* [12].

$$\text{Mutagenic effectiveness (Gamma rays)} = \frac{\text{Rate of mutation (Mp)}}{\text{Dose in Gray (Gy)}}$$

$$\text{Mutagenic effectiveness (SA)} = \frac{\text{Rate of mutation (Mp)}}{\text{Concentration} \times \text{duration of treatment}}$$

$$\text{Mutagenic effectiveness (Gamma rays + SA)} = \frac{\text{Rate of mutation (Mp) Dose}}{\text{of physical mutagen (Gy)} \times \text{concentration of chemical mutagen} \times \text{duration of treatment}}$$

$$\text{Mutagenic efficiency} = \frac{\text{Rate of mutation (Mp)}}{* \text{Biological damage in M}_1 \text{ generation}}$$

*Biological damage: For measuring the biological damage, three different criteria were used; Injury, Sterility and Meiotic abnormalities refers to %age decrease in seedling height (Mp/I), pollen fertility (Mp/S) and meiotic abnormalities (Mp/Me), respectively.

2.6. Morphological mutations

These traits include tall, dwarf, semi dwarf, compact/bushy, prostrate, semi dwarf spreading, one sided branching, axillary branching, broad leaf, narrow leaf, altered leaf architecture, elongated rachis, multiple flower, flower color, open flower, non flowering, late flowering, early maturity, small/narrow pods, bold seeded pods, coat color, coat pattern, shape and surface traits which were examined and modified [13-14]. The characteristics of each mutant is presented in results section. In the field, morphological mutant features for plant height variations, growth habit, leaf, flower, pod and seed were noted.

$$F(\%) = \frac{\text{No. of mutants}}{\text{Total number of plants studied}} \times 100$$

3. Results

3.1. Seed germination

In M_2 generation, compared to the control, the gamma irradiated and SA treated population reflected reduced seed germination (Table 3). In both the varieties, germination decreased with the increase in mutagen dose. In the var. Gomati VU-89, control showed a germination percentage of 93.33%, while it reduced from 88.33 to 80.67% in 100 - 400Gy gamma rays and 87.33 to 81.33% in 0.01-0.04% SA and 87.00 to 77.67% in 100Gy + 0.01% SA - 400Gy+0.04% SA treatments. Variety Pusa-578 showed much reduced seed germination, the germination percentage for control was 92.00%, while it decreased from 85.00 to 80.33% in 100 - 400Gy gamma rays and 88.33 to 76.33% in 0.01 - 0.04% SA and 83.67 to 75.33% in 100Gy + 0.01% SA - 400Gy + 0.04% SA in gamma rays+SA treatments. As compared to M_1 generation, the %age of seed germination was improved in M_2 generation in both the varieties.

Table 3. Effects of gamma rays, SA and their combinations on seed germination and pollen fertility in M₂ generation two varieties of cowpea

Treatment	Gomati VU-89				Pusa-578			
	Seed germination	% inhibition	Pollen fertility (%)	% reduction	Seed germination	% inhibition	Pollen fertility (%)	% reduction
Control	93.33	---	96.53	---	92.00	---	97.77	---
100Gy γ rays	88.33	5.36	90.35	6.41	85.00	7.61	88.12	9.87
200Gy γ rays	86.00	7.86	89.11	7.69	83.33	9.42	86.88	11.14
300Gy γ rays	82.67	11.43	85.40	11.54	81.67	11.23	81.19	16.96
400Gy γ rays	80.67	13.57	81.19	15.90	80.33	12.68	77.23	21.01
Mean	84.42	9.55	86.51	10.38	82.58	10.24	83.35	14.75
0.01% SA	87.33	6.43	90.84	5.90	88.33	3.99	89.11	8.86
0.02% SA	86.00	7.86	87.13	9.74	84.00	8.70	84.65	13.42
0.03% SA	83.33	10.71	81.68	15.38	79.67	13.41	77.72	20.51
0.04% SA	81.33	12.86	77.23	20.00	76.33	16.67	75.00	23.29
Mean	84.50	9.46	84.22	12.76	82.17	10.69	81.62	16.52
100Gy γ rays +0.01% SA	87.00	6.79	91.58	5.13	83.67	9.06	87.13	10.89
200Gy γ rays +0.02% SA	83.67	10.36	86.14	10.77	80.67	12.32	84.41	13.67
300Gy γ rays +0.03% SA	80.33	13.93	83.66	13.33	77.33	15.94	76.49	21.77
400Gy γ rays +0.04% SA	77.67	16.79	74.50	22.82	75.33	18.12	72.03	26.33
Mean	82.17	11.96	83.97	13.01	79.25	13.86	80.01	18.16

3.2. Pollen fertility

Pollen fertility was noted in doses employed individually and in combination (Table 3). Its decrease ranged from 6.41% in 100Gy to 15.90% in 400Gy in gamma rays treatments, while as it ranged from 5.90% in 0.01% SA to 20.00% in 0.04% SA treatments and in case of combination treatments it ranged from 5.13% in 100Gy+0.01% SA to 22.82% in 400Gy+0.04% SA. It ranged from 9.87% in 100Gy to 21.01% in 400Gy in gamma rays treatments, while as it ranged from 8.86% to 23.29 in 0.01% SA-0.04% SA in SA treatments and in case of combination treatments it ranged from 10.89% in 100Gy + 0.01% SA to 26.33% in 400Gy + 0.04% SA. The individual mutagen treatments caused more reduction in pollen fertility than combination treatments in both the varieties.

3.3. Chlorophyll alterations

Chlorophyll mutations are frequently employed for the determination of mutagenic potency in inducing genetic variability as they are vital indicator in the assessment of induced genetic changes of mutagenized population. A concise account of chlorophyll mutants are presented in Table 4 and Fig 2. The spectrum and frequency of the chlorophyll mutants are presented in Table 5. The chlorophyll mutant frequencies were estimated on M₂ seedling basis. In both the varieties, except highest dose in each treatment of gamma rays, SA and gamma rays+SA, mutation frequency increased linearly with the increase in mutagen dose. The combination treatment induced higher mutation frequency in comparison to individual treatments. The chlorophyll mutation frequency ranged from 0.72 to 1.06%, 0.88 to 0.92% and 0.53 to 1.68% in gamma rays, SA and gamma rays+SA treatments, respectively. Var. Pusa-578 responded with the lesser mutation frequency compared to the var. Gomati VU-89. In gamma rays, 0.89 to 1.02% in SA treatment and the range of chlorophyll deficient mutants induced by the combined mutagens was 0.48 to 1.78% in the var. Pusa-578.

Table 4. Characteristic cowpea chlorophyll mutants

Mutant types and their characteristics	Treatment	No. of M ₁ plant progenies	No. of plant progenies segregating in M ₂	% mutated plant
	Gomati VU-89			
1. Albina White leaves; seedlings survive for 10-12 days after germination	Control	270.00	0.00	0.00
	100Gy γ rays	245.00	2.00	0.82
	200Gy γ rays	223.00	11.00	4.93
	300Gy γ rays	208.00	7.00	3.37
	400Gy γ rays	195.00	9.00	4.62
	Total	871.00	29.00	3.33
2. Chlorina Leaves are light green in colour; mostly seedlings survive for 20-25 days but few may survive till maturity	0.01% SA	235.00	2.00	0.85
	0.02% SA	218.00	5.00	2.29
	0.03% SA	203.00	5.00	2.46
	0.04% SA	183.00	6.00	3.28
	Total	839.00	18.00	2.15
3. Xantha Yellow colour leaves due to absence of chlorophyll; seedlings survive for 15-20 days	100Gy γ rays + 0.01% SA	230.00	7.00	3.04
	200Gy γ rays + 0.02% SA	210.00	8.00	3.81
	300Gy γ rays + 0.03% SA	192.00	8.00	4.17
	400Gy γ rays + 0.04% SA	178.00	10.00	5.62
4. Tigrina Leaves with yellow and green patches; seedlings survive for 15-25 days	Total		33.00	4.07
	Pusa-578			
	Control	260.00	0.00	0.00
	100Gy γ rays	235.00	1.00	0.43
5. Viridis Leaves possess viridine green colour; reduced seedling height survive till maturity	200Gy γ rays	213.00	5.00	2.35
	300Gy γ rays	200.00	6.00	3.00
	400Gy γ rays	185.00	6.00	3.24
	Total	833.00	18.00	2.16
	0.01% SA	225.00	1.00	0.44
	0.02% SA	208.00	5.00	2.40
6. Xanthaviridis Leaves possess viridine green colour.	0.03% SA	193.00	4.00	2.07
	0.04% SA	173.00	6.00	3.47
	Total	799.00	16.00	2.00
	100Gy γ rays + 0.01% SA	220.00	6.00	2.73
	200Gy γ rays + 0.02% SA	200.00	6.00	3.00
	300Gy γ rays + 0.03% SA	182.00	7.00	3.85
	400Gy γ rays + 0.04% SA	168.00	6.00	3.57
	Total	770.00	25.00	3.25

195 **Table 5.** Frequency of chlorophyll mutants.

Treatment	N					V	Xv	CMS	F	K	
		A	C	X	T						
Gomati VU-89											
Control	3276.00	-	-	-	-	-	-	-	-	-	
100Gy γ rays	3205.00	12.00	8.00	5.00	1.00	3.00	-	29.00	0.90	-	
200Gy γ rays	3146.00	13.00	7.00	5.00	-	4.00	-	29.00	0.92	-	
300Gy γ rays	3102.00	10.00	8.00	4.00	2.00	4.00	5.00	33.00	1.06	-	
400Gy γ rays	3050.00	9.00	8.00	2.00	-	1.00	2.00	22.00	0.72	-	
0.01%SA	3185.00	13.00	10.00	1.00	3.00	1.00	-	28.00	0.88	-	
0.02%SA	3102.00	14.00	8.00	1.00	-	3.00	2.00	28.00	0.90	-	
0.03%SA	3052.00	10.00	7.00	5.00	4.00	-	2.00	28.00	0.92	-	
0.04%SA	2925.00	9.00	10.00	4.00	-	2.00	1.00	26.00	0.89	-	
100Gy γ rays+0.01% SA	3025.00	11.00	-	-	1.00	-	4.00	16.00	0.53	0.30	
200Gy γ rays+0.02% SA	2956.00	13.00	12.00	-	1.00	-	3.00	29.00	0.98	0.54	
300Gy γ rays+0.03% SA	2854.00	15.00	15.00	6.00	4.00	6.00	2.00	48.00	1.68	0.85	
400Gy γ rays+0.04% SA	2705.00	15.00	13.00	7.00	1.00	7.00	1.00	44.00	1.63	1.02	
Pusa-578											
Control	3147.00	-	-	-	-	-	-	-	-	-	
100Gy γ rays	3097.00	10.00	6.00	0.00	1.00	1.00	-	18.00	0.58	-	
200Gy γ rays	3038.00	11.00	5.00	3.00	-	2.00	-	21.00	0.69	-	
300Gy γ rays	2994.00	9.00	5.00	3.00	2.00	2.00	2.00	23.00	0.77	-	
400Gy γ rays	2942.00	7.00	7.00	0.00	-	2.00	1.00	17.00	0.58	-	
0.01%SA	3077.00	11.00	10.00	1.00	4.00	3.00	-	29.00	0.94	-	
0.02%SA	2994.00	13.00	10.00	2.00	-	2.00	2.00	29.00	0.97	-	
0.03%SA	2944.00	13.00	7.00	3.00	3.00	2.00	2.00	30.00	1.02	-	
0.04%SA	2817.00	11.00	9.00	2.00	-	2.00	1.00	25.00	0.89	-	
100Gy γ rays+0.01% SA	2917.00	10.00	-	-	1.00	-	3.00	14.00	0.48	0.32	
200Gy γ rays+0.02% SA	2848.00	14.00	9.00	-	2.00	-	2.00	27.00	0.95	0.57	
300Gy γ rays+0.03% SA	2746.00	16.00	14.00	6.00	5.00	6.00	2.00	49.00	1.78	1.02	
400Gy γ rays+0.04% SA	2597.00	14.00	14.00	6.00	3.00	5.00	2.00	44.00	1.69	1.15	

196 N= Number of M₂ seedlings, CMS= Chlorophyll mutated seedlings, F= Frequency, K= Coefficient of interaction based on % M₂ chlorophyll mutation
 197 frequency.

Fig 2. Chlorophyll mutants.

a Gomati VU-89 seedling (Control) showing green color of leaf.

b Albina

c Xantha

d Tigrina

e Pusa-578 Control

f Chlorina

g Xanthaviridis

h Viridis

The spectrum of chlorophyll mutations is given in (Table 6). Among the mutants the occurrence of albina type of chlorophyll mutants was the highest in varieties Gomati VU-89 and Pusa-578. Among the doses the frequency of albina mutants was highest in individual and combination doses of gamma rays+SA, whereas SA treatments induced higher frequency of chlorina mutants in varieties Gomati VU-89 and Pusa-578 (Fig 3).

Fig 3. Frequency of chlorophyll mutations.

198 **Table 6.** Chlorophyll mutation frequency.

Mutagen/ Variety							Frequency
	A	C	X	T	V	Xv	
Gomati VU-89							
Gamma radiations	0.35	0.25	0.13	0.02	0.10	0.06	0.90
SA	0.38	0.29	0.09	0.06	0.05	0.04	0.90
Gamma radiations + SA	0.47	0.35	0.11	0.06	0.11	0.09	1.19
Average Frequency	0.40	0.30	0.11	0.04	0.09	0.06	1.00
Pusa-578							
Gamma radiations	0.31	0.19	0.05	0.03	0.06	0.02	0.65
SA	0.41	0.30	0.07	0.06	0.08	0.04	0.96
Gamma radiations + SA	0.49	0.33	0.11	0.10	0.10	0.08	1.21
Average Frequency	0.40	0.27	0.07	0.06	0.09	0.04	0.94

199 *Data based on pooled values of treatments

200254

3.4. Mutagenic effectiveness and efficiency

The mutagen effectiveness and efficiency in a induced mutagenesis determines its usefulness and applicability for crop improvement. Its assessment has been evaluated on the basis of chlorophyll mutation frequency (Tables 7 and 8). The effectiveness and efficiency were highest at the intermediate treatments of mutagens, however lower doses of combined mutagen treatments were most effective and efficient in both the varieties. SA was effective than gamma rays. In both the varieties, effectiveness of single and combined mutagens treatments decreased progressively with the increase in dose in both the varieties. The effectiveness of SA treatments ranged from 25.00 to 41.66 and 16.66 to 41.66 in the varieties Gomati VU-89 and Pusa-578, respectively.

201 **Table 7.** Effectiveness and efficiency of gamma rays, SA and their combinations in cowpea var. Gomati VU-89.

Treatment	(%) Seedling injury (I)	(%) Pollen sterility (S)	%	%	Mutagenic effectiveness	Mutagenic efficiency		
			meiotic abnormalities (Me)	Mutated plant progenies (Mp)		Mp/I	Mp/S	Mp/Me
100Gy γ rays	10.91	6.41	3.19	2.00	0.02	0.18	0.31	0.63
200Gy γ rays	16.37	7.69	5.14	11.00	0.06	0.67	1.43	2.14
300Gy γ rays	18.88	11.54	5.47	7.00	0.02	0.37	0.61	1.28
400Gy γ rays	25.61	15.90	8.51	9.00	0.02	0.35	0.57	1.06
0.01%SA	17.40	5.90	1.63	2.00	33.33	0.11	0.34	1.23
0.02%SA	19.64	9.74	3.50	5.00	41.66	0.25	0.51	1.43
0.03%SA	22.31	15.38	6.02	5.00	27.77	0.22	0.33	0.83
0.04%SA	29.13	20.00	7.63	6.00	25.00	0.21	0.30	0.79
100Gy γ rays+ 0.01% SA	16.89	5.13	5.44	7.00	1.16	0.41	1.37	1.29
200Gy γ rays +0.02% SA	20.35	10.77	6.43	8.00	0.33	0.39	0.74	1.25
300Gy γ rays+0.03% SA	25.05	13.33	9.05	8.00	0.14	0.32	0.60	0.88
400Gy γ rays+0.04% SA	31.79	22.82	10.31	10.00	0.10	0.31	0.44	0.97

The highest efficiency was noted at 200 Gy of gamma rays followed by 100 Gy+0.01% SA and 0.02% SA in both the varieties of cowpea. Mp/Me based efficiency was maximum at 200Gy gamma rays followed by 0.02% SA and 100Gy+0.01% SA in the var. Gomati VU-89. However, the trend was slightly altered in var. Pusa-578 as the highest efficiency was achieved in 0.02% SA than 100Gy+ 0.01% SA and 100 Gy gamma rays. In both the varieties injury and sterility based mutagenic efficiency was found in the decreasing order i.e., gamma rays> gamma rays+ SA> SA. In var. Gomati VU-89 and Pusa-578, meiotic abnormalities based efficiency was higher in gamma rays + SA treatments gamma rays treatments and SA. Meiotic abnormalities based mutagenic efficiency was higher than injury and sterility based efficiency.

Table 8. Effectiveness and efficiency of gamma rays, SA and their combinations in cowpea var. Pusa-578.

Treatments	(%) Seedling injury (I)	(%) Pollen injury (S)	(% abnormalities (Me)	(% plant progenies (Mp)	Mutagenic effectiveness	Mutagenic efficiency		
						Mp/I	Mp/S	Mp/Me
100Gy γ rays	12.28	9.87	3.19	1.00	0.01	0.08	0.10	0.31
200Gy γ rays	17.17	11.14	5.77	6.00	0.03	0.35	0.54	1.04
300Gy γ rays	21.51	16.96	5.83	6.00	0.02	0.28	0.35	1.03
400Gy γ rays	23.47	21.01	8.48	6.00	0.02	0.26	0.29	0.71
0.01%SA	11.99	8.86	1.05	1.00	16.66	0.08	0.11	0.95
0.02%SA	14.84	13.42	3.56	5.00	41.66	0.34	0.37	1.41
0.03%SA	19.18	20.51	5.00	4.00	22.22	0.21	0.20	0.80
0.04%SA	20.12	23.29	6.62	6.00	25.00	0.30	0.26	0.91
100Gy γ rays+ 0.01% SA	13.11	10.89	4.05	5.00	1.00	0.38	0.46	1.23
200Gy γ rays+0.02% SA	17.72	13.67	4.90	6.00	0.25	0.34	0.44	1.22
300Gy γ rays+0.03% SA	21.12	21.77	5.63	7.00	0.13	0.33	0.32	1.24
400Gy γ rays+0.04% SA	26.88	26.33	8.43	8.00	0.06	0.30	0.30	0.95

3.5. Morphological mutations

A diverse range of mutagen induced morphological mutants were obtained in the mutagenized cowpea population in M2 generation, many of which may be useful from yield improvement perspectives. These mutants revealed differences in the traits such as plant height, growth habit, leaf, flower pod and seed. The morphological mutations, detected at different growth stages of M2 plants, were categorically inspected throughout the growing season. The maximum frequency was recorded in gamma rays+SA treatments (5.86 and 6.56%) and the minimum frequency with gamma rays treatments (4.49 and 5.05%), while the SA treatments induced intermediate frequency of morphological mutants (5.38 and 6.05%) in the varieties (Tables 9 and 10). The overall frequencies of morphological mutants were more in Pusa-578 than Gomati VU-89. The spectrum of morphological mutations is given in (Fig 4). The most prominent morphological mutations were found to be associated with the seed and flower and growth habit, plant size and pod length (Tables 11 and 12). The collective treatments produced significant effects on morphological mutations.

Fig 4. Percentage and spectrum of morphological mutations

203 **Table 9.** Frequency and spectrum of morphological mutants.

T	N	Morphological mutant types						TMP	% Mutated	k
		H	G	L	F	P	S			
Control	3155.00	-	-	-	-	-	-	-	-	-
100Gy γ rays	3000.00	8.00	-	6.00	2.00	9.00	4.00	29.00	0.97	-
200Gy γ rays	2941.00	4.00	10.00	13.00	5.00	-	5.00	37.00	1.26	-
300Gy γ rays	2897.00	-	16.00	9.00	7.00	-	6.00	38.00	1.31	-
400Gy γ rays	2845.00	8.00	8.00	-	2.00	6.00	3.00	27.00	0.95	-
Total	11683.00	20.00	34.00	28.00	16.00	15.00	18.00	131.00	4.49	0.00
0.01%SA	2980.00	9.00	-	11.00	12.00	-	3.00	35.00	1.17	-
0.02%SA	2897.00	5.00	-	6.00	5.00	11.00	6.00	33.00	1.14	-
0.03%SA	2847.00	-	12.00	7.00	10.00	-	7.00	36.00	1.26	-
0.04%SA	2720.00	10.00	7.00	13.00	8.00	7.00	4.00	49.00	1.80	-
Total	11444.00	24.00	19.00	37.00	35.00	18.00	20.00	153.00	5.38	0.00
100Gy γ rays +0.01% SA	2820.00	6.00	-	-	15.00	-	9.00	30.00	1.06	0.44
200Gy γ rays+0.02% SA	2751.00	6.00	-	19.00	13.00	12.00	10.00	60.00	2.18	0.80
300Gy γ rays+0.03% SA	2649.00	-	16.00	9.00	-	-	4.00	29.00	1.09	0.37
400Gy γ rays+0.04% SA	2500.00	-	17.00	7.00	-	14.00	0.00	38.00	1.52	0.52
Total	10720.00	12.00	33.00	35.00	28.00	26.00	23.00	157.00	5.86	2.13

204 **Table 10.** Frequency and spectrum of morphological mutants
 205

Treatments	N	P	G	L	F	P	S	TMP	Mutated plants	k
Control	2960.00	-	-	-	-	-	-	-	-	-
100Gy γ rays	2910.00	10.00	-	8.00	2.00	-	9.00	29.00	1.00	-
200Gy γ rays	2851.00	2.00	9.00	14.00	4.00	11.00	10.00	50.00	1.75	-
300Gy γ rays	2807.00	-	14.00	8.00	5.00	-	5.00	32.00	1.14	-
400Gy γ rays	2755.00	7.00	10.00	-	4.00	8.00	3.00	32.00	1.16	-
TOTAL	11323.00	19.00	33.00	30.00	15.00	19.00	27.00	143.00	5.05	-
0.01%SA	2890.00	10.00	-	11.00	12.00	-	4.00	37.00	1.28	-
0.02%SA	2807.00	7.00	13.00	6.00	5.00	9.00	8.00	48.00	1.71	-
0.03%SA	2757.00	-	-	8.00	10.00	9.00	9.00	36.00	1.31	-
0.04%SA	2630.00	11.00	8.00	10.00	10.00	-	7.00	46.00	1.75	-
TOTAL	11084.00	28.00	21.00	35.00	37.00	18.00	28.00	167.00	6.05	-
100Gy γ rays +0.01% SA	2730.00	8.00	-	-	14.00	-	10.00	32.00	1.17	0.46
200Gy γ rays+0.02% SA	2661.00	8.00	-	17.00	15.00	13.00	11.00	64.00	2.41	0.60
300Gy γ rays+0.03% SA	2559.00	3.00	15.00	11.00	-	2.00	5.00	36.00	1.41	0.48
400Gy γ rays+0.04% SA	2410.00	-	14.00	9.00	-	15.00	0.00	38.00	1.58	0.50
TOTAL	10360.00	19.00	29.00	37.00	29.00	30.00	26.00	170.00	6.56	2.04

N = Number of M₂ plants, TMP = Total mutated plants, k = coefficient of interaction based on % M₂ mutated plants.

Table 11. Frequency and spectrum of morphological mutants induced by various mutagens in M₂ Gomati VU-89.

Mutant Type		Gamma rays		SA		Gamma rays + SA		T		GT	
		N	F %	N	F%	N	F%	N	F%	N	F%
Plant height	T	13.00	0.11	15.00	0.13	9.00	0.08	37.00	0.11	56.00	0.17
	D	5.00	0.04	8.00	0.07	2.00	0.02	15.00	0.04		
Growth Habit	SD	2.00	0.02	1.00	0.01	1.00	0.01	4.00	0.01	63.00	0.19
	C	8.00	0.07	7.00	0.06	12.00	0.11	27.00	0.08		
Leaf	P	1.00	0.01	2.00	0.02	5.00	0.05	8.00	0.02		
	SDS	3.00	0.03	4.00	0.03	3.00	0.03	10.00	0.03		
Branch	OSB	7.00	0.06	1.00	0.01	1.00	0.01	9.00	0.03		
	AB	3.00	0.03	2.00	0.02	4.00	0.04	9.00	0.03		
	BL	8.00	0.07	7.00	0.06	6.00	0.06	21.00	0.06	59.00	0.17
	NL	5.00	0.04	5.00	0.04	7.00	0.07	17.00	0.05		
	ALA	2.00	0.02	2.00	0.02	5.00	0.05	9.00	0.03		
	ER	4.00	0.03	6.00	0.05	2.00	0.02	12.00	0.04		
Flower	MF	4.00	0.04	3.00	0.03	3.00	0.03	10.00	0.03	66.00	0.19
	C	2.00	0.02	5.00	0.04	4.00	0.04	11.00	0.03		
	OF	7.00	0.06	7.00	0.06	6.00	0.06	20.00	0.06		
	NF	1.00	0.01	1.00	0.01	2.00	0.02	4.00	0.01		
	LF	6.00	0.05	6.00	0.05	2.00	0.02	14.00	0.04		
	EM	4.00	0.03	2.00	0.02	1.00	0.01	7.00	0.02		
Pod	SNP	7.00	0.06	8.00	0.07	10.00	0.09	25.00	0.07	45.00	0.13
	BP	5.00	0.04	6.00	0.05	9.00	0.08	20.00	0.06		
Seed	CC	7.00	0.06	8.00	0.07	7.00	0.07	22.00	0.06	75.00	0.22
	CP	6.00	0.05	9.00	0.08	8.00	0.07	23.00	0.07		
	SS	10.00	0.09	11.00	0.10	9.00	0.08	30.00	0.09		
Grand Total		120.00	1.03	126.00	1.10	118.00	1.10	364.00	1.08		

206 F % = Mutant frequency of M₂ plants.
207322

208 **Table 12.** Frequency and spectrum of morphological mutants induced by various mutagens in M₂ generation of cowpea var. Pusa-578.

Mutant Type		Gamma rays		SA		Gamma rays + SA		Total		GT	
		N	F %	N	F%	N	F%	N	F%	N	F%
Plant height	T	10.00	0.09	11.00	0.10	8.00	0.08	29.00	0.09	44.00	0.13
	D	4.00	0.04	7.00	0.06	1.00	0.01	12.00	0.04		
	S	1.00	0.01	1.00	0.01	1.00	0.01	3.00	0.01		
Growth habit	C	7.00	0.06	6.00	0.05	11.00	0.11	24.00	0.07	57.00	0.17
	P	4.00	0.04	1.00	0.01	6.00	0.06	11.00	0.03		
	SDS	2.00	0.02	3.00	0.03	2.00	0.02	7.00	0.02		
	OSB	5.00	0.04	2.00	0.02	1.00	0.01	8.00	0.02		
	AB	2.00	0.02	3.00	0.03	2.00	0.02	7.00	0.02		
Leaf	BL	7.00	0.06	6.00	0.05	5.00	0.05	18.00	0.05	50.00	0.15
	NL	4.00	0.04	4.00	0.04	5.00	0.05	13.00	0.04		
	ALA	1.00	0.01	3.00	0.03	4.00	0.04	8.00	0.02		
	ER	5.00	0.04	5.00	0.05	1.00	0.01	11.00	0.03		
Flower	MF	3.00	0.03	2.00	0.02	4.00	0.04	9.00	0.03	64.00	0.20
	C	3.00	0.03	4.00	0.04	5.00	0.05	12.00	0.04		
	OF	6.00	0.05	8.00	0.07	5.00	0.05	19.00	0.06		
	NF	2.00	0.02	2.00	0.02	1.00	0.01	5.00	0.02		
	LF	5.00	0.04	5.00	0.05	1.00	0.01	11.00	0.03		
	EM	3.00	0.03	4.00	0.04	1.00	0.01	8.00	0.02		
Pod	SNP	5.00	0.04	7.00	0.06	9.00	0.09	21.00	0.06	45.00	0.14
	BP	4.00	0.04	5.00	0.05	8.00	0.08	17.00	0.05		
Seed	CC	6.00	0.05	7.00	0.06	6.00	0.06	19.00	0.06	75.00	0.23
	CP	5.00	0.04	8.00	0.07	7.00	0.07	20.00	0.06		
	SS	9.00	0.08	10.00	0.09	8.00	0.08	27.00	0.08		
Grand Total		103.00	0.91	114.00	1.03	102.00	0.98	319.00	0.97		

209 F% = Mutant frequency of M₂ Plants

I. Plant height mutants (Fig 5)

(i) Tall mutants: Such mutants were induced with frequency of 0.11 and 0.09% at lower doses of gamma rays and SA in both the cowpea varieties. The mutants Plants attained a height 180-185 cm compared to 178-183 cm in control plants.

(ii) Dwarf mutants: Such mutants were mostly induced with a 0.04% frequency of the total morphological mutations. These mutants were induced at higher doses of all the treatments except the combination treatments in the var. Gomati VU-89. These mutants showed lesser leaves, reduced pod and seed size and poor yield. The plant height was severely reduced and it ranged from 110-115 cm.

(iii) Semi dwarf mutants: Such mutants were induced at 0.01% frequency. Pod length was increased, however, the seeds were smaller in size.

II. Growth habit mutants (Fig 5)

(i) Compact/Bushy mutants: Such mutants are characterised with 150.55 cm average height, robust growth. The yield per plant was slightly higher and matured five days earlier compared to control plants. Such mutants were mostly induced by the moderate and higher treatments. The bushy mutants appeared at 0.08% of the total morphological mutations in the var. Gomati VU-89 and 0.07% in var. Pusa-578.

(ii) Prostrate mutants: Such mutants were flat on ground, branches spread, possess long internodes, and reflected trailing tendency at the soil surface. They possess, less number of pods containing 6-7 shriveled seeds with rough and hard seed coat. In both the varieties of cowpea prostrate mutants were commonly induced with combination treatments

(iii) Spreading mutants: Such mutants were observed in 200Gy gamma rays treatment (Gomati VU-89) and 0.03% SA treatment (Pusa-578). The leaves were mostly broad and

branches appeared spreading. Seed yield per plant was significantly reduced. Their frequency was 0.03% and 0.02% in varieties Gomati VU-89 and Pusa-578, respectively.

(iv) One sided branching mutants: Mutant plants had thick stems bearing branches on one side and were late in flowering and maturing, low yielded with few pods and shriveled seeds. These mutants were screened in SA doses in both the cowpea varieties and in combination treatment of 200Gy gamma rays + 0.02% SA in the var. Pusa-578.

Fig 5. Morphological mutations.

- a** Control Plant
- b** Tall mutant
- c** Dwarf mutant
- d** Semi dwarf mutant
- e** Spreading mutant.
- f** Bushy mutant.
- g** Axillary branched mutant.
- h** Prostrate mutant

III. Leaf mutants (Fig 6-7)

(i) Broad/Giant leaves mutants: Moderate doses of gamma rays and SA treatments induced higher frequency of broad leaf mutants.

(ii) Narrow leaves mutants: Such mutants possess narrow leaves with pointed leaf tips. Compared to control, the yield was lower, small pods with few seeds per pod. Higher doses of mutagens induce higher frequency of mutants in both the cowpea varieties.

(iii) Altered leaf architecture mutants: Plants showed reduced height (about 95 cm) at maturity. Such mutants were induced by all the mutagen treatments

(iv) Elongated rachis: Mutants possessed increased length of rachis with narrow leaflets. They appeared mostly at the lower mutagen doses with a frequency of 0.04% and 0.03% in the var. Gomati VU-89 and var. Pusa-578, respectively.

Fig 6. Morphological mutations in leaves

a Control leaf

b Broad leaf mutant

c Narrow leaf mutant

d Long rachis mutant

e Trifoliate mutant

f Leaf architecture.

Fig 7. Morphological mutations (plants with altered leaflet number)

g Mutant showing incision of leaf.

h Tetrafoliate with fused terminal leaflets.

i Pentafoiate with variations in leaflet size and shape.

j Tetrafoliate with free leaflets.

k Broad tetrafoliate leaflets.

l Bifoliate leaf.

IV. Flower mutants (Fig 8)

(i) Multiple flower mutants: Such mutants have small pods and shrivelled seeds. Mutants were late in maturity compared to the control. These mutants appeared at frequency of about 0.03% in both the varieties.

(ii) Flower colour mutants: The 300 Gy gamma rays showed higher frequency of white or blue flower colour mutants. Their frequency was recorded as 0.03% in the var. Gomati VU-89 and 0.04% in the var. Pusa-578.

(iii) Open flower mutants: Such mutants appeared at 0.06% frequency of total mutation at moderate and higher doses of SA in var. Gomati VU-89 and Pusa-578.

(iv) Non-flowering/Vegetative mutants: These mutants were noticed in SA and gamma rays+SA treatments.

(v) Late flowering mutants: Higher doses of mutagens induced large number of late flowering mutants.

(vi) Early maturity mutants: Such mutants matured 10-15 days earlier than control and were induced at the lower mutagens doses used singly or in combination. The frequency of such mutants was 0.02% in both the varieties.

Fig 8. Morphological mutations in flowers

a Control flowers

b Petal mutants

c Multiple flowers.

d Open flower mutant.

e Light blue colour flower mutant.

f White flowers mutant.

V. Pod mutants (Fig 9)

(i) Small/Narrow pods: Such mutants possessed small and /or narrow pods compared to long pods in control, observed in combined mutagen treatments. The seed yield per plant was also reduced as compared to control plants.

(ii) Bold seeded pod mutants: The bold size of pods led to significant increase in yield per plant as compared to control. Lower and moderate doses of individual and combined mutagens induced broad spectrum of these mutants.

Fig 9. Morphological mutations in pods

a Normal pod (control).

b Long pods.

c Broad pod with bold seeds.

d Broad pods.

e Pod width mutants.

f Pod length mutants.

VI. Seed mutants (Fig 10)

(i) Seed coat colour mutants: Such mutants are characterized red and black seed coats compared to brown and white seed coats in control plants of the var. Gomati VU-89 and var. Pusa-578, respectively.

(ii) Seed shape and surface mutants: Seeds of control plants of both the varieties were of kidney shaped with rough seed surface and were induced in lower doses. Mutants with globular shape and smooth seed surface were isolated in the mutagen treated population.

Fig 10. Morphological mutations in seed characters)

a Seed coat mutants.

b Bold seed mutants

c Red seed mutants.

d Wrinkled seed mutants.

4. Discussion

A considerable decline in seed germination, seedling height, pollen fertility and plant survival of treated population was observed in mutagen treatments. However, the extent of diminution in the bio-physiological parameters differed among mutagen doses. The diminution in plant survival was recorded in all the mutagen doses, and confirmed the fact that the mutagens interact independently with the genes governing a particular trait and that mutations occur in a random manner. These results are in line with earlier reports on several grain legumes [15], [16], [17], [18], [19], [20], [21]. However, the promoting effects of mutagen doses on bio-physiological parameters were also reported in several other crops [22], [23], [24], [25], [26].

5.1. Seed Germination

The assessment of seed germination gives an idea about the mutagenic potency, the dose at which germination is below 50 per cent is treated as lethal or undesirable. Since, the purpose of this present study was to develop large mutagenized population to facilitate effective screening of the high yielding mutants, therefore, the lethal doses of mutagens were discarded to ensure high frequency of desirable mutation. Decreased germination due to mutagen treatments may be attributed to mutagen induced germination inhibition. The inhibition in seed germination was found to be more or less comparable in gamma rays and SA treatments, while the combined treatments induced higher seed germination inhibition in both the cowpea varieties, which may be attributed to the synergistic effect of the two mutagens. Chauhan and Singh [27] reported that gamma rays interact and cause disruption of tunica zone that leads to the decrease in seed germination. Alterations in hormone synthesizing enzymes may be another reason for seed germination inhibition [28]. In the present study, the decrease in pollen fertility has been noticed in the mutagenized population. Pollen sterility was maximum in combined mutagen doses than individual mutagen doses. This

may cause meiotic anomalies and leading to the formation of aberrant pollen grains. Reduction in pollen fertility after mutagen treatments have also been reported in lentil [29], [30], cowpea [31], fenugreek [32], faba bean [33]. The pollen sterility percentage was found remarkably less in M_2 than M_1 generation. The survival at maturity was declined with increasing mutagen dose in both the varieties of cowpea. These findings support the earlier reports in *Lathyrus sativus* [34] in *Capsicum annum*, [35]. The reduction of plant survival at maturity may be attributed to the disturbed physiological processes [36], [37] or cytological damages Manjaya [38]. The observations recorded on several bio-physiological and cytological parameters revealed that cowpea var. Pusa-578 was comparatively more sensitive towards the individual and combination doses than Gomati VU-89. The differential response of varieties from the same species towards the mutagens have also been reported earlier in *Cajanas cajan* [39], *Vigna* spp [40], [41], [42], *Phaseolus vulgaris* [43], [44], *Jatropha curcus* [45] and *Lens culinaris* [46]. Sparrow *et al.* [47] suggested that differential genotypic sensitivity towards mutagens may be due to the gene loci mutated by one mutagen in one variety were not necessarily mutated by the second mutagen in another variety. The two cowpea varieties viz., Gomati VU-89 and Pusa-578, used in the present study, were from different races, therefore, it is obvious that the two varieties have substantial variation in their genome and thereby the heritable genetic identity of the two varieties is immensely different. Although the spectrum of variations was more or less similar in both the varieties but the mutagen doses to achieve that were not identical in various parameters studied. It can be concluded from the present bio-physiological that mutagenesis induced a comparable spectrum of apparent variation through random mutation in both the varieties of diverse origin with the variable treatments of gamma rays and SA.

5.2. Chlorophyll mutations

In mutation breeding, the increase in mutation frequency is necessary for crop

improvement. In order to acquire desired mutations at highest rates, the picking of optimum mutagen and appropriate dose is important. Both the varieties of cowpea showed a differential response towards gamma rays, SA and gamma rays+SA treatments. The chlorophyll mutants were identified as per the categorization of Gustafsson [9]. Gamma rays+SA caused maximum frequency of chlorophyll mutations than the gamma rays and SA. Similar trends were reported in chickpea [48], *Oryza sativa* [49], *Vigna mungo* [50] *Zingiber officinale* [51], *Lablab purpureus* [52]. On contrary Pavadai *et al.* [53] in soybean, Barshile *et al.* [54] in chickpea, Kumar *et al.* [55] in mungbean reported the linear increase in chlorophyll mutation frequency with increase in doses of different mutagen treatments. In the present study, albina and chlorina mutants were found more frequently in both the cowpea varieties. SA was found to be more potent in the production of albina and chlorina mutants than gamma rays. The gamma rays+SA treatments have induced maximum frequency of albina and chlorine mutants than gamma rays and SA treatments employed individually. The synergistic effects in the combination treatments are very much evident from the higher frequency of chlorophyll mutations. The possible reason for synergism might be that the mutagen first applied may expose the accessible protected mutable sites to the second mutagen, the repair enzymes may be rendered non functional by the second mutagen which indirectly facilitates mutations fixation [11]. The synergism were earlier documented in *Cicer arietinum* [56], *Phaseolus vulgaris* [57], *Vigna spp.*, [58], *Pisum sativum* [59], *Linum usitatissimum* [60].

5.3. Mutagenic effectiveness and efficiency

Induced mutagenesis has been known as a vital tool for the improvement of wide range of crops. The success of mutation breeding depends on enhancing the efficiency of desired mutation. Further the mutagen choice is imperative to achieve the desired mutation rate [12], [61]. Whereas mutagenic efficiency on the contrary the SA doses were superior among mutagen

doses employed in present study. Similar results have been reported by Balai and Krishna [62] while studying the effectiveness of ethylmethane sulphonate, sodium azide and hydroxylamine on mungbean. Earlier workers such as Kaul and Bhan [63] in rice, Laskar and Khan [64] in lentil, Wani [65] and Shah *et al.* [66] in chickpea, Khan and Tyagi [67] in soybean and Dhulgande *et al.* [68] in pea have also reported the more effectiveness of chemical mutagens over physical mutagens [69] while studying the mutation breeding in cowpea. As per them, in order to attain higher effectiveness and efficiency, the mutation effect was more prominent in the cell that led to meiotic anomalies. Our results reflected a differential effectiveness in two varieties of cowpea which depict the genetic divergence and mutagenic sensitivity towards different mutagen doses. In the present study, the gamma rays+SA were more efficient than individual gamma rays and SA treatments and this may be attributed to the higher chlorophyll mutants. On contrary Zeerak [70] in brinjal reported that gamma rays+ EMS doses were less efficient than the individual doses of gamma rays and EMS. In the present study, lower and medium doses of gamma rays and SA and gamma rays+SA treatments were found to be more efficient in both the varieties of cowpea. The results were in line with the earlier results in *Lens culinaris* [64] *Glycine max* [67], *Vigna mungo* [71]. However, contrary results of linear increase in mutagenic efficiency with the increase in the mutagen concentrations have also been reported in black gram [72] and millet [73]. Makeen *et al.* [74] also reported higher 613 mutagenic efficiency at lower and medium doses in urdbean.

5.4. Morphological mutations

Even though the majority of the induced morphological mutants were uneconomical, however, these mutants can serve as a source of valuable genes in hybridisation programmes. Toker [75] reported that the morphological mutants are useful in gene mapping and phylogenetic studies of crops. Mutagen induced chromosomal anomalies or the pleiotropic effects of mutated genes could be the probable reason for the induction of morphological mutants [76], [77]. Different mutagen doses and duration of treatment may be

attributed to the relative proportion of different mutation types [78].. Khursheed *et al.* [79] reported maximum morphological mutants frequency in gamma rays and lowest with combined treatments while investigating the frequency and spectrum of M₂ morphological faba bean mutants. Previous reports of a high frequency and boarder spectrum of morphological mutations in crops such as *Vigna mungo* [41], *Cicer arietinum* [80], [81], *Lens culinaris* [82], *Phaseolus vulgaris* [83] *Glycine max* [67] also confirm that mutagens are effective in inducing morphological mutations in wide range of crops that may serve as a source of valuable genes for the crop improvement programs. Some morphological mutants (early maturing, dwarf, bushy, bold seeded pods and flower colour), screened in M₂ generation in the present study, can serve as source of valuable genes or used as a parents in cross breeding programmes. Dwarf mutants, as recorded in the present study, have been reported earlier in chickpea [21] barley [84], mungbean spp. [77], lentil [85]. The decreased length of internodes and/or number of internode could be attributed to the induction of dwarf and bushy mutants [86]. In our experimentation dwarf and bushy mutants showed substantial decrease in internode length but number of internodes were similar to that of control plants. The mutated genes that control the altered plant height and growth habits are monogenic recessives [87]. Whereas polygenes govern the semi dwarf trait in mutants in wheat [12] and triticale [88]. However, Qin *et al.* [89] reported a dominant mutation in a single gene in dwarf mutants in rice. Both dwarfness and bushy habit of a genotype contribute to the production and productivity. The bold pod mutants are associated with high yield which may be valuable in increasing seed size that lead to enhanced cowpea yield. Singh [90] reported that gene mutations govern the bold seeded trait in *Vigna mungo*.

Conclusions

Increase in mutagen doses lead to a progressive decline in the values of all bio-physiological parameters. Combination treatments showed an enhanced effect on seed germination, seedling height and pollen fertility compared to individual mutagen treatments. A dose dependant increase in the frequency of chlorophyll mutations was observed in both the varieties. The estimation of coefficient of interaction (k) did not show any additive effects of mutagens. Mutagenic effectiveness was highest at the moderate doses of gamma rays and SA, however, lower doses of gamma rays+SA were most effective and efficient in both the varieties. The effectiveness of mutagens was found as SA> gamma rays + SA> gamma rays. Mp/I and Mp/S based mutagenic efficiency was gamma rays> gamma rays+ SA> SA whereas Mp/Me based order was gamma rays + SA> gamma rays> SA. A diverse range of mutagen induced morphological mutants were obtained in the population of two varieties of cowpea in M₂ generation, many of which are useful from yield improvement prospective. Of all the mutant types, seed and flower mutants followed by growth habit, plant height and pod length were of maximum occurrence in the two varieties. The spectrum of morphological mutations induced by combined mutagens was relatively boarder than gamma rays and SA treatments in both the varieties.

References

1. FAO. Definition and Classification of Commodities: Pulses and Derived Products.1994.
2. DES. Agriculture statistics at a glance. Directorate of Economics and Statistics, Department of Agriculture and Economics, Ministry of Agriculture, Government of India 2017.
3. Summerfield RJ, Huxley PA, Dart PJ, Hughes AP. Some effects of environmental stress on seed yield of cowpea (*Vigna unguiculata* (L.) Walp.) cv. Prima. Plant and Soil. 1976; 44: 527–546.
4. Sprent IJ, Odee WD, Dakora FD. African Legume: a vital but under-utilized resource. J Exp. Bot. 2009; 61: 1257–1265.
5. Fery RL. The cowpea: production, utilization, and research in the United States. Hort. Reviews. 1990; 12: 197–222.

6. Raina A, Khursheed S, Khan S. Optimisation of mutagen doses for gamma rays and sodium azide in cowpea genotypes. *Trends. Biosci.* 2018; 11: 2386–2389
7. FAO. Production Statistics. Food and Agriculture Organization, Rome, Italy 2017.
8. Raina A, Laskar RA, Tantray YR, Khursheed S, Wani MR, Khan S. Characterization of Induced High Yielding Cowpea Mutant Lines Using Physiological, Biochemical and Molecular Markers. *Sci. Rep.* 2020; 10: 1–22.
9. Gustafsson A. The mutation system of the chlorophyll apparatus. Lund Univ. 1940
10. Khan S. Studies on Chemical Mutagenesis in Mungbean (*Vigna radiata* (L.) Wilczek). Ph.D. Thesis. Aligarh Muslim University, Aligarh. 1990.
11. Sharma RP. Increased mutation frequency and wider mutation spectrum in barley induced by combining gamma rays with ethyl methane sulphonate. *Indian J. Genet.* 1970; 30: 180–186.
12. Konzak CF, Nilan RA, Wagner J, Foster RJ. Efficient chemical mutagenesis. *Rad. Bot.* 1965; 5: 49–70.
13. UPOV. Cowpea (*Vigna unguiculata* L. Walp. subsp. *sesquipedalis* (L.) Verdc.). International Union for The Protection of New Varieties of Plants Nairobi, Kenya, 2007; 1–20
14. IBPGR. Cowpea descriptors. IBPGR Secretariat, Rome, Italy, 1983; 1–30.
15. Toker C, Cagiran MI. Spectrum and frequency of induced mutations in chickpea. *Intl. Chickpea Pigeonpea Newslet.* 2004; 11: 8–10.
16. Khan S, Wani MR. Comparison on the effect of chemical mutagens on mungbean. *Advanced Plant Sci.* 2005; 18: 533–535.
17. Lal GM, Toms B, Lal SS (2009) Mutagenic sensitivity in early generation in black gram. *Asian J. Agric. Sci.* 2009; 1: 9–11.
18. Bhat M, Khan S, Kozgar MI. Studies on induced mutations in chickpea (*Cicer arietinum* L.) I. Responses of the mutagenic treatments in M1 biological parameters. *Electronic J. Plant Breed.* 2011; 2: 422–424.
19. Khan H, Laskar RA, Khan S. Comparative estimation of induced cytotoxicity and mutagenic potency of EMS and SA using *Vicia faba* L. as a biological system, In: Krishnan S., Rodrigues BF., editors. *Advances in Plant Sciences and Biotechnology.* Goa University; 2015. pp. 172–186.
20. Laskar RA, Laskar AA, Raina A, Khan S, Younus H. Induced mutation analysis with biochemical and molecular characterization of high yielding lentil mutant lines. *Intl. J. Biol. Macromolecules.* 2018; 109: 167–179.
21. Raina A, Laskar RA, Khursheed S, Khan S, Parveen K, Amin R, Khan S. Induce physical and chemical mutagenesis for improvement of yield attributing traits and their correlation analysis in chickpea. *Intl. Let. Nat. Sci.* 2017; 61: 14–22.
22. Venkateshwarlu S, Singh RM, Singh RB, Singh BD. Radiosensitivity and frequency of chlorophyll mutations in pigeonpea. *Indian J. Genet.* 1978; 38: 90–94.

23. Vandana, Dubey DK. Effect of ethylmethane sulphonate (EMS) and diethyl sulphate (DES) in faba bean. FABIS Newslet. 1988; 32: 18–22.
24. Mensah JK, Obadoni B. Effects of sodium azide on yield parameters of groundnut (*Arachis hypogaea* L.). Afr. J. Biotechnol. 2007; 6: 668–671.
25. Boureima S, Diouf M, Silme RS, Diop T, Van Damme P, Çağırhan Mİ. Radiosensitivity of African sesame cultivars to gamma rays. Turk. J. Field Crops. 2009; 14: 181–190.
26. Verma AK, Banerji BK, Chakrabarty, Datta SK. Studies on Makhana (*Euryale ferox* Salisbury). Current Sci. 2010; 99: 795–800.
27. Chauhan MP, Singh IS. Inheritance of protein content in lentil (*Lens culinaris* Medik.). Legume Res. 1995; 18: 5–8.
28. Usuf KK, Nair PM. Effect of gamma irradiation on the indole acetic acid synthesizing system and its significance in sprout inhibition of potatoes. Rad. Bot. 1974; 14: 251–256.
29. Reddy VRK, Annadurai M. Cytological effects of different mutagens in lentil (*Lens culinaris* Medik.). Cytologia. 1992; 57: 213–216
30. Taziun T, Laskar RA, Amin R, Parveen SK. Effects of dosage and durations of different mutagenic treatment in lentil (*Lens culinaris* Medik.) cultivars Pant L 406 and DPL 62. Legume Res. 2017; 41: 500–509.
31. Singh VV, Krishna RK, Arya RK. Induced chemical mutagenesis in cowpea (*Vigna unguiculata* (L) Walp. Indian J. Genet. 2006; 66: 312–315.
32. Choudhary S, Ansari MYK, Khan Z, Gupta H. Cytotoxic action of lead nitrate on cytomorphology of *Trigonella foenum-graecum* L. Turk. J. Biol. 2012; 36: 267–273.
33. Bhat TA, Wani AA. Mutagenic Effects on Meiosis in Legumes and a Practical Case Study of *Vicia faba* L. In: Bhat TA., Wani AA. editors. Chromosome Structure and Aberrations. Springer; 2007. pp. 219–244.
34. Kumar S, Dubey DK. Effect of separate and simultaneous application of gamma rays and EMS on germination, growth, fertility and yield in cultivars Nirmal and LSD-3 of khesari (*Lathyrus sativus* L.) Var. P-505. J. Phytol. Res. 1998; 11: 165–70.
35. Dhamayanthi KPM, Reddy VRK. Cytogenetic effects of gamma rays and ethyl methane sulphonate in chilli pepper (*Capsicum annuum* L.). Cytologia. 2000; 65: 129–133.
36. Sato M, Gaul H. Effect of ethyl methane sulphonate on the fertility of barley. Rad.Bot. 1967; 7: 7–15.
37. Meherchandani M. Effects of gamma radiation on the dormant seeds of *Avena fatua* L. Rad. Bot. 1975; 15: 439–443.
38. Manjaya JG. Genetic improvement of soybean variety VLS-2 through induced mutations. In: Shu QY. editors. Induced Plant Mutation in the Genomic Era. Food and Agriculture Organization of the United Nations, Rome; 2009. pp. 106–110.
39. Pandey N, Ojha CB, Jha VB, Singh NB. Effect of chemical mutagens on the rate of germination, seedling mortality, and induced sterility in pigeonpea. Int. Chickpea and Pigeonpea Newslet. 1996; 3: 65–67.

40. Khan S, Siddiqui BA. Mutation genetic studies in mungbean. I. Variability components and genetic parameters. Thailand J. Agric. Sci. 1995; 28: 113–124.
41. Goyal S, Khan S. Differential response of single and combined treatment in moist seeds of urdbean. Indian J. Bot. Res. 2010; 6: 183–188.
42. Kozgar MI, Khan S, Goyal S. EMS induced mutational variability in *Vigna radiata* and *Vigna mungo*. Res. J. Bot. 2011; 6: 31–37.
43. Al-Rubeai MAF, Godward MBE. Genetic control of radiosensitivity in *Phaseolus vulgaris* L. Env. Exp. Bot. 1981; 21: 211–216.
44. Svetleva DL, Crino P. Effect of ethyl methanesulfonate (EMS) and n- nitrosen –ethyl urea (ENU) on callus growth of common bean. J. Central European Agric. 2006; 6: 59–64.
45. Dhakshanamoorthy D, Selvaraj R, Chidambaram A. Physical and chemical mutagenesis in *Jatropha curcas* L. to induce variability in seed germination, growth and yield traits. Rom. J. Biol. 2010; 55: 113–125.
46. Gaikwad NB, Kothekar VS. Mutagenic effectiveness and efficiency of ethylmethane sulphate and sodium azide in lentil (*Lens culinaris* Medik.). Indian J. Genet. 2004; 64: 73–74.
47. Sparrow AH, Sparrow RC, Thompson KH, Schairer DA. The use of nuclear and chromosome variables in determining and predicting radiosensitivity. In: Use of Induced Mutation in Plant Breeding. Rep/IAEA, Rome; 1965; 102–132.
48. Khursheed S, Raina A, Laskar RA, Khan S. Effect of gamma radiation and EMS on mutation rate: their effectiveness and efficiency in faba bean (*Vicia faba* L.). Caryologia. 2018; 71: 397–404.
49. Yamaguchi H, Hase Y, Tanaka A, Shikazono N, Degi K, Shimizo A, Morishita T. Mutagenic effects of ion beam irradiation on rice. Breed. Sci. 2009; 59: 169–177.
50. Thilagavathi C, Mullainathan L. Isolation of macro mutants and mutagenic effectiveness, efficiency in Black Gram (*Vigna mungo* (L.) Hepper). Global J. Mol. Sci. 2009; 4: 76–79.
51. Pawar N, Pai S, Nimbalkar M, Kolar F, Dixit G. Induction of chlorophyll mutants in *Zingiber officinale* Roscoe by gamma rays and EMS. Emirates J Food Agric. 2010; 22: 406–411.
52. Srinivas RT, Veerabhadhiran P. Efficiency and effectiveness of physical and chemical mutagens and their combination in inducing chlorophyll mutations in M2 generation of Lablab [*Lablab purpureus* (L.) Sweet var. Typicus.]. Electronic J. Plant Breed. 2010; 1: 752–757.
53. Pavadai P, Giriya M, Dhanavel D. Effectiveness, efficiency and biochemical content of physical and chemical mutagens in soybean (*Glycine max* (L.) Merr.). J. Phytol. 2009; 1: 444–447.
54. Barshile JD, Auti SG, Dalve SC, Apparao BJ. Mutagenic sensitivity studies in chickpea employing SA, EMS and gamma rays. Indian J. Pulses Res. 2006; 19: 43– 46.
55. Kumar A, Parmhansh P, Prasad R. Induced chlorophyll and morphological mutations in

mungbean (*Vigna radiata* (L.) Wilczek). *Legume Res.* 2009; 32: 41–45.

56. Kozgar MI. Induced Mutagenesis: Biophysiological Damages and Cytological Aberrations. In: *Mutation Breeding in Chickpea*. 2014. pp. 13–30.

57. Agarkova SN, Yakovlev AG. Effects of gamma rays and EMS on mutation frequency in *Phaseolus vulgaris*. *Referats Zhurnal*. 1977; 3: 55–56.

58. Singh AK. Mutagenic effectiveness and efficiency of gamma rays and ethylmethane sulphonate in mungbean. *Madras Agric. J.* 2007; 94: 7–13.

59. Mehandjiev A, Kosturkova G, Mihov M. Enrichment of *Pisum sativum* gene resources through combined use of physical and chemical mutagens. *Israel J. Plant Sci.* 2001; 49(4): 280–284.

60. Bhat IA, Pandit UJ, Sheikh IS, Hassan ZU. Physical and chemical mutagenesis in *Linum usitatissimum* L. to induce variability in seed germination, survival and growth rate traits. *Current Bot.* 2016; 7: 28–32.

61. Bhosale SS, Kothekar VS. Mutagenic efficiency and effectiveness in Cluster Bean (*Cyamopsis tetragonoloba* (L.) Taub.). *J. Phytol.* 2010; 2: 21–27.

62. Balai OP, Ram K. Efficiency and effectiveness of chemical mutagens in mungbean. *J. Food Legumes.* 2009; 22: 105–108.

63. Kaul MLH, Bhan AK. Mutagenic effectiveness and efficiency of EMS, DES and gamma rays in rice. *Theor. Appl. Genet.* 1977; 50: 241–246.

64. Laskar RA, Khan S. Mutagenic Effectiveness and Efficiency of Gamma Rays and HZ with Phenotyping of Induced Mutations in Lentil Cultivars. *Intl. Let. Nat. Sci.* 2017; 64: 17–31.

65. Wani AA. Mutagenic effectiveness and efficiency of gamma rays, ethyl methane sulphonate and their combination treatments in chickpea (*Cicer arietinum* L.). *Asian J. Plant Sci.* 2009; 8: 318.

66. Shah TM, Mirza JI, Haq MA, Atta BM. Induced genetic variability in chickpea (*Cicer arietinum* L.) II. Comparative mutagenic effectiveness and efficiency of physical and chemical mutagens. *Pak J. Bot.* 2008; 40(2): 605–613.

67. Khan MH, Tyagi SD. Studies on effectiveness and efficiency of gamma rays, EMS and their combination in soybean (*Glycine max* (L.) Merrill). *J. Plant Breed. Crop Sci.* 2010; 2(3): 55–58.

68. Dhulgande GS, Dhale DA, Pachkore GL, Satpute RA. Mutagenic effectiveness and efficiency of gamma rays and Ethyl methanesulphonate in Pea (*Pisum sativum* L.). *J. Exp. Sci.* 2011; 2(3): 7–8.

69. Giriya M, Dhanavel D. Mutagenic effectiveness and efficiency of gamma rays, ethyl methane sulphonate and their combination treatment in Cowpea (*Vigna unguiculata* L. Walp). *Global J. Mol. Sci.* 2009; 4: 68–75.

70. Zeerak NA. Mutagenic effectiveness and efficiency of gamma rays and ethyl methane sulphonate in brinjal. *J. Nuc. Agric. Biol.* 1992; 21: 84–87.

71. Sharma A, Plaha P, Rathour R, Katoch V, Singh Y, Khalsa GS. Induced mutagenesis for improvement of garden pea. *Intl. J. Vegetable Sci.* 2010; 16: 60–72.
72. Gautam AS, Sood KC, Richaria AK. Mutagenic effectiveness and efficiency of gamma rays, ethylmethane sulphonate and their synergistic effects in black gram (*Vigna mungo* L.). *Cytologia.* 1992; 57: 85–89.
73. Ganapathy S, Nirmalakumari A, Senthil N, Souframanien J, Raveendran TS. Isolation of macromutations and mutagenic effectiveness and efficiency in little millet varieties. *World J. Agric. Sci.* 2008; 4: 483–486.
74. Makeen K, Babu GS. Mutagenic effectiveness and efficiency of gamma rays, sodium azide and their synergistic effects in urd bean (*Vigna mungo* L.). *World J. Agric. Sci.* 2010; 6 (2): 234–237.
75. Toker C. A note on the evolution of kabuli chickpeas as shown by induced mutations in (*Cicer reticulatum* Ladizinsky.) *Genet. Resources Crop Evol.* 2009; 56: 7–12.
76. Gottschalk W. The genetic basis of variation. In: *Improving Vegetatively Propagated Crops*, Academic Press Limited, London, UK; 1987. pp. 317–334.
77. Wani MR, Khan S, Kozgar MI, Goyal S. Induction of morphological mutants in mungbean (*Vigna radiata* (L.) Wilczek) through chemical mutagens. *Nucleus.* 2011; 48: 243–247
78. Nilan RA. Nature of induced mutations in higher plants. Induced mutations and their utilization. In: *Proc Symp Erawim Baur Gedachtnis Orle Singen IV 1966*. Acverlag, Berlin; 1967. pp 5–20
79. Khursheed S, Raina A, Parveen K, Khan S. Induced phenotypic diversity in the mutagenized populations of faba bean using physical and chemical mutagenesis. *J. Saudi Society Agric. Sci.* 2019; 18: 113–119.
80. Khan MR, Qurshi AS, Hussain SA, Ibrahim M. Spectrum of morphological mutations induced by separate and simultaneous application of gamma rays with GA3 in chickpea (*Cicer arietinum* L.). *Pak. J. Biol. Sci.* 2000; 3:1431–5.
81. Laskar RA, Khan S, Khursheed S, Raina A, Amin R. Quantitative analysis of induced phenotypic diversity in chickpea using physical and chemical mutagenesis. *J. Agron.* 2015; 14 (3): 102–111.
82. Solanki IS, Sharma B. Induction and isolation of morphological mutations in different damage groups in lentil (*Lens culinaris* Medik). *Indian J. Genet.* 1999; 59(4): 479–485.
83. Yankulov MT, Isasi EM, Abreu F. Some aspects of the sensitivity and mutability of two French bean varieties under the influence of gamma radiation from ⁶⁰Co and ethyl methanesulphonate (EMS). *Ciencias de la Agricultura.* 1980; 7: 59–64.
84. Sethi GS. Long-penduncled mutant: a new mutant type induced in barley. *Euphytica*, 1974; 23: 237–239.
85. Shahwar D, Ansari MY, Choudhary S. Induction of phenotypic diversity in mutagenized population of lentil (*Lens culinaris* Medik) by using heavy metal. *Heliyon.* 2019; 5(5):e01722.

86. Sjodin J. Induced morphological variations in *Vicia faba* L. *Hereditas*. 1971; 67: 155–180.
87. Reddy VRK, Gupta PK. Induced mutations in hexaploid triticales. Frequency and spectrum of morphological mutations. *Genet. Agric.* 1988; 42: 241–254.
88. Shakoar A, Sadiq MS, Hasan MU, Saleem M. Selection for useful semi dwarf mutants through induced mutation in bread wheat. *Proceed. In: 5th Wheat Genetics Symposium, New Delhi; 1978. pp. 540–546.*
89. Qin R, Qiu Y, Cheng Z, Shan X, Guo X, Zhai H, Wan J. Genetic analysis of a novel dominant rice dwarf mutant 986083D. *Euphytica*. 2008; 160: 379–387.
90. Singh RK. Gamma ray induced bold seeded mutant in *Vigna mungo* (L.) Hepper. *Indian J. Genet.* 1996; 56: 104–108.

Author Contributions

Conceptualization: Aamir Raina, Samiullah Khan.

Data curation: Aamir Raina.

Formal analysis: Aamir Raina.

Investigation: Aamir Raina.

Methodology: Aamir Raina.

Project administration: Aamir Raina.

Resources: Aamir Raina.

Software: Aamir Raina.

Supervision: Samiullah Khan.

Validation: Aamir Raina.

Visualization: Aamir Raina.

Writing - original draft: Aamir Raina.

Writing - review & editing: Aamir Raina.

Competing interests

The authors declare no competing interests.

Gamma rays

SA

Gomati VU-89

Pusa-578