1 **Title:** Intracellular and intercellular gene regulatory networks inference from
2 time-course individual RNA-Seq.

3 **Authors:** Makoto Kashima^{1*}, Yuki Shida², Takashi Yamashiro², Hiromi Hirata¹, and
4 Hiroshi Kurosaka²

5 ¹College of Science and Engineering, Aoyama Gakuin University, 5-10-1, Fuchinobe,
6 Sagamihara Chuo Ku, Kanagawa 252-5258, Japan;

7 ²Department of Orthodontics and Dentofacial Orthopedics, Osaka University,
8 Yamadaoka, Suita, Osaka, 565-0871, Japan.

9 * Corresponding author

10 Contact: kashima@chem.aoyama.ac.jp

11

12 **Abstract**

13 Gene regulatory network (GRN) inference is an effective approach to understand the
14 molecular mechanisms underlying biological events. Generally, GRN inference mainly
15 targets intracellular regulatory relationships such as transcription factors and their
16 associated targets. In multicellular organisms, there are both intracellular and
17 intercellular regulatory mechanisms. Thus, we hypothesize that GRNs inferred from
18 time-course individual (whole embryo) RNA-Seq during development can reveal
19 intercellular regulatory relationships (signaling pathways) underlying the development.
20 Here, we conducted time-course bulk RNA-Seq of individual mouse embryos during
21 early development, followed by pseudo-time analysis and GRN inference. The results
22 demonstrated that GRN inference from RNA-Seq with pseudo-time can be applied for
23 individual bulk RNA-Seq similar to scRNA-Seq. Validation using an
24 experimental-source-based database showed that our approach could significantly infer

GRN for all transcription factors in the database. Furthermore, the inferred ligand-related and receptor-related downstream genes were significantly overlapped. Thus, the inferred GRN based on whole organism could include intercellular regulatory relationships, which cannot be inferred from scRNA-Seq based only on gene expression data. Overall, inferring GRN from time-course bulk RNA-Seq is an effective approach for understanding the regulatory relationships underlying biological events in multicellular organisms.

Introduction

Regulation of gene expression is a fundamental factor that controls cellular events such as proliferation and differentiation. Understanding gene regulatory interconnections is an important step to elucidate the molecular mechanisms underlying cellular events. Recently, gene regulatory network (GRN) inference based on time-course data has attracted much attention in single-cell RNA-Seq (scRNA-Seq). State-of-the-art scRNA-Seq analysis techniques can generate transcriptome information from thousands of cells¹⁻⁵. Transcriptomic heterogeneity of cells due to asynchronous progression of cellular events enables us to infer regulatory relationships of genes. During inference, first, dimensional reduction of scRNA-Seq data provides a trajectory of cellular events such as differentiation and proliferation^{6,7}. Then, assignment of pseudo-time can place cells along the trajectory. Since scRNA-Seq with pseudo-time is a dense time-course observation of cellular events, gene regulatory networks can be inferred by comparing the timing of gene upregulation and downregulation along pseudo-time^{8,9}.

Originally, GRN inference was applied for gene expression data from tissue and pooled cells (bulk samples), generated using DNA microarray and RNA-Seq^{10,11}.

49 Compared to steady-state data, time-course data allows GRN inference by comparing
50 the timing of gene upregulation and downregulation^{12–15}. However, GRN inference from
51 time-course data of bulk samples has not been popular due to the following drawbacks:
52 1) RNA extraction and library preparation of a large number of bulk samples prior to
53 sequencing are too costly and time-consuming¹⁶, and 2) biological variances may result
54 in inconsistencies between real sampling time and transcriptome status¹⁷. Fortunately,
55 recent technical advances related to bulk RNA-Seq have overcome these limitations.
56 Advances in sequencing platforms¹⁸, RNA extraction method^{16,19} and bulk 3' RNA-Seq
57 library preparation methods^{20–22} have enabled a cost-effective time-course individual
58 RNA-Seq²³ (a time-series RNA-Seq targeting a whole embryo or tissue of each
59 individual). Pseudo-time analysis for individual RNA-Seq might capture individual
60 differences in progression speed of biological events such as development. Following
61 assignment of pseudo-time to each individual RNA-Seq data, GRN can be inferred
62 similar to an inference based on scRNA-Seq.

63 Theoretically, GRN inferred from whole-body and tissue RNA-Seq are different
64 from those inferred from scRNA-Seq. scRNA-Seq provides transcriptomic information
65 at the cellular level that enables inference of intracellular GRN involved in the
66 progression of proliferation and differentiation²⁴. In contrast, bulk RNA-Seq of whole
67 body and tissues could contain transcriptomic information at the level of cell
68 populations. Time course individual RNA-Seq during development would enable
69 inference of both intracellular and intercellular GRN (cell-cell communications)
70 involved in the developmental process. For instance, during embryonic development,
71 upregulation of ligand-related genes can be followed by upregulation of downstream
72 genes in cells expressing receptor genes²⁵.

73 We thus hypothesize that GRNs inferred from time-course individual RNA-Seq
74 during embryonic development would include intercellular regulatory relationships
75 between ligand genes and downstream genes of related signaling pathways. To test this
76 hypothesis, we conducted time-course bulk RNA-Seq of individual mouse embryos in
77 early development, followed by pseudo-time analysis and GRN inference.

78

79 **Results**

80 **Pseudo-time analysis of the time-course bulk RNA-Seq for mouse embryos**

81 To determine whether gene regulatory network (GRN) inference based on time-course
82 bulk RNA-Seq is effective, we conducted time-course bulk RNA-Seq for individual
83 mouse embryos. RNA was extracted from each individual whole embryo (n = 10 or 11)
84 at seven time points: E7.5, E8.5, E9.5, E10.5, E11.5, E12.5, and E.13.5, followed by 3'
85 RNA-Seq using the 'Lasy-Seq' method²¹. As a result, we obtained 76 RNA-Seq datasets
86 with an average of 8.5 million reads per sample. The reads were mapped onto the mouse
87 reference sequence, followed by calculation of the read counts of each gene in each
88 sample. We then used 'Seurat' ²⁶, an R package used for single cell omics analysis,
89 normalization of read counts, detection of highly variable genes, and dimension
90 reduction of omics data. On the PC1 and PC2 plane obtained with Seurat, samples at the
91 same stage were close to each other (Fig. 1a). As expected, clusters of each stage were
92 ordered according to the developmental process (Fig. 1b). Using an R package
93 'slingshot'²⁷, we inferred the developmental trajectory of mouse embryos and calculated
94 the pseudo-time for each sample (Fig. 1 a and b). Pseudo-time analysis revealed
95 individual differences among embryos in the speed of their developmental processes.
96 For example, the pseudo-time of a sample at E10.5 and of E11.5 were very close to each

other (Fig. 1 a and b). Using the pseudo-time analysis, the time-course RNA-Seq data of the seven time points could be converted into those of 76 time points. Using pseudo-time as temporal information instead of stage (real sampling time) improved the sum of squared residuals (SSR) between observed and fitted values (Fig. 1c and d). The SSRs along pseudo-time were decreased by 0.745% (all genes) and 3.835% (high variable genes), on average, compared with the SSRs along stage. These results indicate that integrating pseudo-time into the analysis, instead of real sampled stage, could improve capture temporal expression dynamics, by considering individual differences in the progression speed of biological events during the mouse early embryonic development.

Gene regulatory network inference based on individual RNA-Seq of whole mouse embryos

Next, we inferred a GRN from the dataset of time-course individual RNA-Seq of whole mouse embryos. We used SCODE, which solves linear ordinary differential equations to infer GRN⁸. To avoid incorrect emphasis on the technical noise of RNA-Seq, we used a non-scaled normalized gene expression matrix as the input for SCODE. Selection of the D size, a parameter that affects the number of assumed basic patterns of expression dynamics of the dataset, is important for robust GRN inference, since a needlessly large D causes an unstable inferred GRN⁸. In this study, we used $D = 4$ similar to the previous study⁸, with which the SSR was relatively small (Fig. 2a). For 28117 genes whose read counts were greater than zero, SCODE produced A (28117×28117 matrix) corresponding to the inferred gene regulatory network, in which the value of A_{ij} indicates regulatory effects on the downstream gene i from the regulator j . $A_{ij} > 0$ indicates that the regulator j positively regulates gene i , whereas $A_{ij} < 0$ indicates the

121 opposite. Because SCODE optimizes A by random sampling, we optimized A 20 times
 122 to check for reproducibility. Then, Pearson's correlation coefficients between the values
 123 of each A from the 20 optimizations and the mean A were calculated (Fig. 2b). Almost all
 124 optimizations produced similar A with high correlation coefficients (Fig. 2b). In the
 125 subsequent analysis, we used the average values of the top 10 A showing higher
 126 correlations with the mean A from 20 optimizations. We then tried to define the
 127 thresholds for significant regulatory relationships between regulators and downstream
 128 genes. Because the average expression level of downstream genes showed higher
 129 correlation with the absolute values of A compared to the regulators (Fig. 2 c and d), we
 130 independently defined the threshold for each regulator instead of a constant threshold
 131 previously used⁸. For example, absolute values of A indicating regulatory relationships
 132 between *Sox8* and its regulators showed a positive correlation (Pearson's correlation
 133 coefficient = 0.76) (Fig. 2e), whereas absolute values of A showing regulatory
 134 relationships between *Sox8* and its downstream genes showed smaller correlations
 135 (Pearson's correlation coefficient = 0.54) (Fig. 2f). Although most of the inferred values
 136 of A for regulatory relationships between *Sox8* and its downstream genes were around
 137 zero, some values were outliers (Fig. 2g). To define the threshold for the downstream
 138 genes of *Sox8*, we regressed the linear function for the scatter plot of absolute A values
 139 for downstream genes in the decreasing order and defined the threshold (Fig. 2g).
 140 Finally, in the subsequent analysis, genes with larger absolute values of A compared to
 141 the threshold were defined as the inferred genes downstream of *Sox8* (Fig. 2g).

142 **Validation of the inferred network**

143 Next, we evaluated the inferred GRN by comparing the inferred genes downstream of
 144 the TFs with the information in the TF2DNA database, which is an

experimental-source-based database of the binding motifs and downstream genes for 438 mouse TFs^{28–39}. First, we evaluated the effectiveness of target prediction based on the absolute values of A by calculating the area under the curve (AUC) (Fig. 3a), obtaining an average AUC of 0.704, suggesting that the inferred regulatory relationships could reflect the true regulatory network. Also, we calculated AUC for inferred regulatory relationships using dynGENIE3⁴⁰, an algorithm using sampled stage information as temporal information to infer GRN. The average AUC was 0.50 (Fig. 3a), suggesting that advantage of usage of pseudo-time in GRN inference based on time-course individual RNA-Seq. Second, we examined the validity of the defined thresholds by assessing the differences between the validated target gene rate (number of validated target genes / number of all inferred downstream genes) above the defined thresholds and the best validated rate (Fig. 3b). Below the threshold, validated target gene rates for the 438 TFs were 0.69% smaller than the best validated target gene rates on average. The validated target genes were 67.5% of the inferred downstream genes on average (Table S1, Fig. 3c and d). Compared with the background rate (number of known target genes / number of all genes), the validated target gene rates of the inferred downstream genes of all TFs were statistically higher (adjusted p-value < 0.01) (Fig. 3d). These results suggest that our approach could infer the GRN underlying early development in mice.

The inferred network contained regulatory relationships involved in cell-cell interaction

Our GRN inference was based on bulk RNA-Seq containing the information of all cells in the body. We thus hypothesized that the inferred GRN also included the intercellular regulatory network. To examine this possibility, we checked the overlaps of inferred

downstream genes for genes related to the ligands and receptors of nine major signaling pathways (Fig. 4a): Wnt/ β catenin, TNF, TGF- β , Hedgehog, FGF, EGF, Delta/Notch, BMP, and Retinoic acid (RA) signaling pathways (Table S2). As expected, the inferred downstream genes of all pairs of ligand and receptor genes were significantly overlapped (adjusted p-value < 0.01) (Fig. 4a). On average, 94.2% of the inferred downstream genes for the ligand-related and receptor-related genes were overlapped (Fig. 4A). For example, 4510 genes were inferred as the downstream of *Wnt* genes, and 4545 genes were downstream of *Fzd* genes. However, 97.5% of the inferred downstream of *Wnt* genes were also inferred as the downstream of *Fzd* genes (Fig. 4a). Since SCODE can infer whether each downstream gene is positively or negatively regulated⁸, we assessed the overlaps of positively and negatively regulated gene downstream of the representative ligand-related and receptor-related genes of each signaling pathway, i.e., with the largest number of inferred downstream genes among each gene family (Fig. 4 b-d and Supplementary Figure 1,2). For six signaling pathways (TNF β , TGF, FGF, EGF, Delta/Notch, and BMP signaling pathways), which activate the downstream genes only when ligands bind to the receptors⁴¹, the regulatory directions of ligand-related and receptor-related genes for most inferred downstream genes were the same (Fig. 4b and Supplementary Figure 1). In contrast, in case of the RA signaling pathway, wherein RA receptors function as transcriptional repressors without RA binding (Supplementary Figure 2a)⁴², the regulatory directions of a ligand-related gene, *Aldh1a3*, which encodes a protein involved in RA synthesis, and a receptor-related gene, *Rara*, for the downstream genes were opposite (Fig. 4c). In case of the Hedgehog signaling pathway, the regulatory directions of *Ssh2* and *Ptchd4* tended to be opposite (Fig. 4d). The regulatory directions of *Ssh2* and *Gli3* for the downstream genes were

also opposite (Fig. 4d). In the absence of Hedgehog ligands, the full-length Gli family proteins are ubiquitinated and function as repressors (Supplementary Figure 2b)⁴³. With the binding of Hedgehog ligands, PATCHED inhibits the degradation of Gli family proteins and allows them to function as transcriptional activators (Supplementary Figure 2a)⁴³. Thus, the overlap of downstream genes that are positively and negatively regulated by *Ssh2*, *Ptchd4*, and *Gli3* is reasonable. In case of the Wnt/ β catenin signaling pathway, the regulatory directions of *Wnt7a* and *Fzd5* for the downstream were opposite (Supplementary Figure 3a). Our time-course RNA-Seq revealed that among the *Fzd* and *Wnt* genes, *Fzd5* and *Wnt8a*, but not *Wnt7a*, were the only genes expressed dominantly in the early developmental stage among the protein families. (Supplementary Figures 4 and 5). Similar regulatory directions were found for most of the inferred downstream genes of *Wnt8a-Fzd5*, which could be a functional pair in the early developmental stages. Furthermore, the regulatory directions for most of the inferred downstream genes of *Wnt7a-Fzd6* (*Fzd* genes with the second most inferred downstream genes) were also the same (Supplementary Figure 3). In conclusion, our approach allowed successful inference the intercellular regulatory relationships related to the major signaling pathways as well as the intracellular pathways related to TFs.

Discussion

Recently, GRN inference based on the combination of scRNA-Seq and pseudo-time analysis has attracted much attention⁴⁴. However, to our knowledge, this study is the first to report GRN inference based on the combination of individual bulk RNA-Seq and pseudo-time analysis. Herein, pseudo-time explained variances of gene expression during mouse early development better than real sampled stage (Fig. 1 c and d), suggesting that the variance of gene expression observed using our time-course

individual RNA-Seq was not mere stochastic variability in gene expression but individual difference in the progression speed of development. Thus, pseudo-time uses the correlation of gene expression dynamics more effectively than the real temporal information in time-course individual RNA-Seq.

221

Unlike scRNA-Seq, which can provide transcriptomic dynamics in a certain cellular event such as proliferation and differentiation, bulk RNA-Seq could provide a mixture of various transcriptomic dynamics regarding the cellular events occurring in an embryo⁴⁵. Theoretically, GRN inference based on bulk RNA-Seq cannot provide multiple lineage-specific regulatory relationships. This may explain why GRN inference based on bulk RNA-Seq has not been attempted as with scRNA-Seq. In this study, we succeeded in inferring the known regulatory relationships of the TFs in the TF2DNA database with high AUC compared to the GRN inferred from scRNA-Seq⁴⁶ (Fig. 3). This suggests that GRN inference from bulk RNA-Seq provides insights into the regulatory interconnections of genes, even though GRN inferred from bulk RNA-Seq would have limitations in the comprehensiveness of inferred regulatory relationships compared to those from scRNA-Seq. Furthermore, drastic changes in the cell population occur during early developments. It is possible that our GRN inferred from bulk RNA-Seq merely reflect the changes in cell populations but not the interconnections of genes. Considering the results of the inference for *Wnt* and *Fzd* genes (Supplementary Figure 2), these points should be considered when interpreting the inferred GRN based on the time-course bulk RNA-Seq.

Assignment of pseudo-time is an important step in GRN inference from time-course data. In case of scRNA-Seq, the accuracy of pseudo-time assignment is

241 controversial⁴⁷. In contrast, in the case of time-course individual bulk RNA-Seq, the
242 accuracy of pseudo-time assignment could be assured by the actual sampling time,
243 which can be an advantage compared to GRN inference from scRNA-Seq.

244 In this study, we proposed a new strategy for threshold of significant gene
245 regulatory relationships inferred by SCODE (Fig. 2). In the case of *E2f3*, the number of
246 predicted target genes is only ~10% of the ground truth targets under our strategy.
247 Shifting the threshold to approximately 17,000 targets for *E2f3* may still result in a
248 validated target rate of approximately 80% (Fig. 3b). Since decreasing threshold may
249 produce false positive inferred relationships for some genes, the determination of the
250 threshold remains challenging. Considering high AUCs for the TFs in the database (Fig.
251 3a), the threshold should change depending on the study aims.

252 GRN indicates the intracellular interconnections of genes in a narrow sense;
253 intercellular regulation of genes via cell-cell communication is also a key factor to
254 understand the regulatory mechanisms underlying multicellular organisms. Several
255 studies have attempted to systematically identify cell-cell communications based on
256 single cell gene expression profiles and information regarding ligand-receptor pairs^{48–51}.
257 Since these approaches required prior knowledge, they could only be applied for the
258 major model organisms and could not reveal novel signaling pathways. We
259 demonstrated that GRN inference based on time-course individual RNA-Seq could infer
260 intercellular regulatory relationships related to cell-cell communication via cell
261 signaling pathways. This approach only requires time-course RNA-Seq results and is
262 applicable for non-model organisms without ligand-receptor information. Theoretically,
263 the GRN inferred based on scRNA-Seq cannot include intercellular interconnections of
264 genes. Comparison of GRN from scRNA-Seq and bulk RNA-Seq would be useful for

systematic identification of genes involved in cell-cell communications during cellular events of interest. Taken together, our approach is a powerful tool for understanding intracellular and intercellular regulatory relationships of genes, which cannot be achieved using the existing GRN inferences based on scRNA-Seq alone. As discussed above, bulk RNA-Seq is limited in the comprehensiveness of inferred regulatory relationship since multiple lineage-specific regulatory relationships cannot be deduced from GRN inference based on bulk RNA-Seq. A future novel bioinformatic approach that can deconvolute gene expression in each tissue and cell lineage from bulk RNA-Seq will help overcoming this limitation.

Methods

Maintenance of mice

Embryos were collected from pregnant female Institute of Cancer Research (ICR) mice (CLEA, Tokyo, Japan) at each stage (E7.5, E8.5, E9.5, E10.5, E11.5, E12.5, and E13.5). The number of replicates (embryos) were 10 at E7.5 and 11 at the remaining stages. All sacrificed female mice were housed under a 12 h dark-light cycle in which the light phase started from 8 am. All animal experiments were performed in accordance with the guidelines of the Animal Care and Use Committee of Osaka University Graduate School of Dentistry, Osaka, Japan.

RNA extraction

Total RNA extraction was performed using the RNeasy® kit (QIAGEN, Hilden, Germany) according to manufacturer's protocol. The total RNA concentration was measured using Qubit™ RNA HS Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA) and was adjusted to 5 ng/ul and stored at -80 °C until subsequent analysis.

289 **RNA-Seq library preparation and sequencing**

290 Non targeted RNA-Seq were conducted according to the Lasy-Seq ver. 1.1 protocol
 291 (<https://sites.google.com/view/lasy-seq/>)^{21,23}. Briefly, 50 ng of total RNA were reverse
 292 transcribed using an RT primer with index and SuperScript IV reverse transcriptase
 293 (Thermo Fisher Scientific, Waltham, MA, USA). Then, all RT mixtures of the samples
 294 were pooled and purified using an equal volume of AMPure XP beads (Beckman
 295 Coulter, Brea, CA, USA) according to the manufacturer's instructions. Second strand
 296 synthesis was conducted on the pooled samples using RNaseH (5 U/μL, Enzymatics,
 297 Beverly, MA, USA), and DNA polymerase I (10 U/μL, Enzymatics, Beverly, MA,
 298 USA). To avoid the carryover of large amounts of rRNAs, the mixture was subjected to
 299 RNase treatment using RNase T1 (Thermo Fisher Scientific, Waltham, MA, USA).
 300 Then, purification was conducted with a 0.8× volume of AMPure XP beads.
 301 Fragmentation, end-repair, and A-tailing were conducted using 5× WGS Fragmentation
 302 Mix (Enzymatics, Beverly, MA, USA). The Adapter for Lasy-Seq was ligated using 5×
 303 Ligation Mix (Enzymatics, Beverly, MA, USA), and the adapter-ligated DNA was
 304 purified twice with a 0.8× volume of AMPure XP beads. After optimisation of PCR
 305 cycles for library amplification by qPCR using EvaGreen, 20× in water (Biotium,
 306 Fremont, CA, USA) and the QuantStudio5 Real-Time PCR System (Applied
 307 Biosystems, Waltham, MA, USA), the library was amplified using KAPA HiFi HotStart
 308 ReadyMix (KAPA BIOSYSTEMS, Wilmington, MA, USA) on the ProFlex PCR
 309 System (Applied Biosystems, Waltham, MA, USA). The amplified library was purified
 310 with an equal volume of AMPure XP beads. One microliter of the library was then used
 311 for electrophoresis using a Bioanalyzer 2100 with the Agilent High Sensitivity DNA kit
 312 (Agilent Technologies, Santa Clara, CA, USA) to assess quality. Then, sequencing of

150-bp paired-end reads was performed using HiSeq X Ten (Illumina, San Diego, CA, USA).

Mapping and gene quantification

Read 1 reads were processed with fastp (version 0.21.0)⁵² using the following parameters: `--trim_poly_x -w 20 --adapter_sequence=AGATCGGAAGAGCACACGTCTGAACTCCAGTCA --adapter_sequence_r2=AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT -l 31.`

The trimmed reads were then mapped to the mouse reference sequences of Mus_musculus.GRCm38.cdna.all.fa, using BWA mem (version 0.7.17-r1188)⁵³ with the default parameters. The read count for each gene was calculated with salmon using -l IU, which specifies the library type (version v0.12.0)⁵⁴. Then, using R (version 4.0.1)⁵⁵, the sums of read counts per gene were calculated. Genes with read counts greater than zero were used in the subsequent analysis.

Pseudo-time analysis

Read counts were normalized using the “NormalizeData” function with the default parameters in Seurat (version 4.0.0)⁵⁶, which produces natural-log transformed (read per 10,000 + 1). For principal component analysis (PCA), the normalized read counts were centered but not scaled using the “ScaleData” function with the default parameters except for do.scale = F. Then, PCA was conducted using the “RunPCA” function for genes with high dispersion, which were selected using the “FindVariableFeatures” function with default parameters except for selection.method = "mvp". Finally, SingleCellExperiment (version 1.10.1)⁵⁷ and slingshot (version 1.6.1)²⁷ were used to calculate the pseudo-time for each sample.

Evaluation of considering pseudo-time instead of stage

337 The “smooth.spline” function in R (version 4.0.1)⁵⁵ with the default parameters except
 338 for “all.knots = T, lambda = 0.001” was used to obtain smoothed curve for normalized
 339 expression of each gene in the “data” slot of the Seurat object, and stage or pseudo-time.
 340 Then sum of squared residuals (SSR) between observed and fitted values was calculated
 341 for each gene. Mean of SSRs were calculated against all genes and the high variable
 342 genes obtained with the “FindVariableFeatures” function.

343 **Gene regulatory network inference**

344 In the SCODE algorithm⁸, normalized expression data in the “data” slot of the Seurat
 345 object, and pseudo-time were used to infer GRN. A was optimized 20 times with one
 346 hundred iterations and $D = 4$. Pearson’s correlation coefficients between values of each
 347 A from the 20 optimizations and the mean A (the average of each value of A) were
 348 calculated. In the following analysis, we used the average values of the top 10 A
 349 showing higher correlations with the mean A from 20 optimizations. To define the
 350 thresholds for downstream gene selection for each gene, the linear function was
 351 regressed using the “nls” function in R for the scatter plot of absolute values of A for
 352 downstream genes in the decreasing order; the X and Y axes showed the integers from 1
 353 to 28,117, and the absolute values of A , respectively. Genes with larger absolute values
 354 of A compared to the Y values of the regressed line were defined as the inferred gene
 355 downstream of each gene.

356 In the dynGENIE3 algorithm⁴⁰, normalized expression data in the “data” slot of the
 357 Seurat object, and samples stage were used to infer GRN. E7.5, E8.5, E9.5, E10.5,
 358 E.11.5, E12.5, and E.13.5 were converted into 1, 2, 3, 4, 5, 6, and 7, respectively. The all
 359 parameters were used with the default values. weight.matrix inferred by dynGENIE3
 360 was used as inferred regulatory relationships.

361 **Evaluation of inferred GRN by comparing with a TF-downstream gene database**

362 For validating the inferred GRN, information regarding the binding motifs and targets
 363 for 438 mouse TFs in the TF2DNA database was used^{28–39}. AUC values for downstream
 364 prediction based on the absolute values of A were calculated using the “performance”
 365 function in ROCR (version 1.0-11)⁵⁸. Statistical analysis for the enrichment of validated
 366 target genes among the inferred genes was conducted using the “fisher.test” function in
 367 R. Statistical analysis for the overlap of inferred downstream genes for ligand-related
 368 and receptor-related genes was conducted using the “enrichment_test” function in
 369 Rvenn (version 1.1.0)⁵⁹. For all statistical tests, BH correction was performed using the
 370 “p.adjust” function. The upset plots were drawn using the “upset” function in UpSetR
 371 (version 1.4.0)⁶⁰.

372
 373 **Acknowledgments.** This work was supported by Japan Society for the Promotion of
 374 Science [grant number JP19H03858 to H.K.]. We would like to thank Editage
 375 (www.editage.com) for English language editing.

377 **Author contributions**

378 M.K., Y.S., and H.K. conceived and conducted the experiments, and M.K. analyzed the
 379 results and wrote the manuscript. All authors reviewed and approved the final
 380 manuscript.

382 **Competing interests**

383 The authors declare no competing interests.

384

385 **Data availability**

386 All RNA-Seq sequence data is deposited in PRJNA725414 of SRA. The R source code
387 used in the present study is available at <https://github.com/Makoto-Kashima/TIODE>

389 **References**

- 390 1. Sasagawa, Y. *et al.* Quartz-Seq2: A high-throughput single-cell RNA-sequencing
391 method that effectively uses limited sequence reads. *Genome Biol.* **19**, 1–24
392 (2018).
- 393 2. Sasagawa, Y. *et al.* Quartz-Seq: a highly reproducible and sensitive single-cell
394 RNA-Seq reveals non-genetic gene expression heterogeneity. *Genome Biol.* **14**,
395 R31 (2013).
- 396 3. Gao, C., Zhang, M. & Chen, L. The Comparison of Two Single-cell Sequencing
397 Platforms: BD Rhapsody and 10x Genomics Chromium. *Curr. Genomics* **21**,
398 602–609 (2020).
- 399 4. Hayashi, T. *et al.* Single-cell full-length total RNA sequencing uncovers
400 dynamics of recursive splicing and enhancer RNAs. *Nat. Commun.* **9**, 1–16
401 (2018).
- 402 5. Klein, A. M. *et al.* Droplet barcoding for single-cell transcriptomics applied to
403 embryonic stem cells. *Cell* **161**, 1187–1201 (2015).
- 404 6. Treutlein, B. *et al.* Reconstructing lineage hierarchies of the distal lung
405 epithelium using single-cell RNA-seq. *Nature* **509**, 371–375 (2014).
- 406 7. Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. Diffusion
407 pseudotime robustly reconstructs lineage branching. *Nat. Methods* **13**, 845–848
408 (2016).

- 409 8. Matsumoto, H. *et al.* SCODE: an efficient regulatory network inference
410 algorithm from single-cell RNA-Seq during differentiation. *Bioinformatics* **33**,
411 2314–2321 (2017).
- 412 9. Aalto, A., Viitasaari, L., Ilmonen, P., Mombaerts, L. & Gonçalves, J. Gene
413 regulatory network inference from sparsely sampled noisy data. *Nat. Commun.*
414 **11**, 3493 (2020).
- 415 10. Ko, D. K. & Brandizzi, F. Network-based approaches for understanding gene
416 regulation and function in plants. *Plant Journal* vol. 104 302–317 (2020).
- 417 11. Fernandez-Valverde, S. L., Aguilera, F. & Ramos-Díaz, R. A. Inference of
418 developmental gene regulatory networks beyond classical model systems: New
419 approaches in the post-genomic era. *Integr. Comp. Biol.* **58**, 640–653 (2018).
- 420 12. Iglesias-Martinez, L. F., Kolch, W. & Santra, T. BGRMI: A method for inferring
421 gene regulatory networks from time-course gene expression data and its
422 application in breast cancer research. *Sci. Rep.* **6**, 1–12 (2016).
- 423 13. Zhang, J. *et al.* Differential regulatory network-based quantification and
424 prioritization of key genes underlying cancer drug resistance based on
425 time-course RNA-seq data. *PLOS Comput. Biol.* **15**, e1007435 (2019).
- 426 14. Krouk, G., Mirowski, P., LeCun, Y., Shasha, D. E. & Coruzzi, G. M. Predictive
427 network modeling of the high-resolution dynamic plant transcriptome in response
428 to nitrate. *Genome Biol.* **11**, R123 (2010).
- 429 15. Ogami, K. *et al.* Computational gene network analysis reveals TNF-induced
430 angiogenesis. *BMC Syst. Biol.* **6**, 3–8 (2012).
- 431 16. Yoshino, K., Nishijima, R. & Kawakatsu, T. Low-cost RNA extraction method
432 for highly scalable transcriptome studies. *Breed. Sci.* (2020)

- doi:10.1270/jsbbs.19170.
17. Kilfoil, M. L., Lasko, P. & Abouheif, E. Stochastic variation: From single cells to superorganisms. *HFSP J.* **3**, 379–385 (2009).
18. Muir, P. *et al.* The real cost of sequencing: scaling computation to keep pace with data generation. *Genome Biol.* **17**, 53 (2016).
19. Ujibe, K., Nishimura, K., Kashima, M. & Hirata, H. Direct-TRI: High-throughput RNA-extracting method for all stages of zebrafish development. *BIO-PROTOCOL* (2021).
20. Alpern, D. *et al.* BRB-seq: ultra-affordable high-throughput transcriptomics enabled by bulk RNA barcoding and sequencing. *Genome Biol.* **20**, 71 (2019).
21. Kamitani, M., Kashima, M., Tezuka, A. & Nagano, A. J. Lasy-Seq: a high-throughput library preparation method for RNA-Seq and its application in the analysis of plant responses to fluctuating temperatures. *Sci. Rep.* **9**, 7091 (2019).
22. Li, Y. *et al.* Decode-seq: A practical approach to improve differential gene expression analysis. *Genome Biol.* **21**, 66 (2020).
23. Kashima, M., Kamitani, M., Nomura, Y., Hirata, H. & Nagano, A. J. DeLTa-Seq: direct-lysate targeted RNA-Seq from crude tissue lysate. (8/15 words) Running Title: Development of direct-lysate targeted RNA-Seq method Corresponding Author. *bioRxiv* 2020.09.15.299180 (2020) doi:10.1101/2020.09.15.299180.
24. Lam, K. Y., Westrick, Z. M., Müller, C. L., Christiaen, L. & Bonneau, R. Fused Regression for Multi-source Gene Regulatory Network Inference. *PLoS Comput. Biol.* **12**, 1–23 (2016).
25. Basson, M. A. Signaling in Cell Differentiation and Morphogenesis. *Cold Spring*

- 457 *Harb. Perspect. Biol.* **4**, 1–21 (2012).
- 458 26. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**,
459 1888–1902.e21 (2019).
- 460 27. Street, K. *et al.* Slingshot: Cell lineage and pseudotime inference for single-cell
461 transcriptomics. *BMC Genomics* **19**, 477 (2018).
- 462 28. Pujato, M., Kieken, F., Skiles, A. A., Tapinos, N. & Fiser, A. Prediction of DNA
463 binding motifs from 3D models of transcription factors; identifying TLX3
464 regulated genes. *Nucleic Acids Res.* **42**, 13500–13512 (2014).
- 465 29. Badis, G. *et al.* Diversity and complexity in DNA recognition by transcription
466 factors. *Science (80-.).* **324**, 1720–1723 (2009).
- 467 30. Wei, G. H. *et al.* Genome-wide analysis of ETS-family DNA-binding in vitro and
468 in vivo. *EMBO J.* **29**, 2147–2160 (2010).
- 469 31. Weirauch, M. T. *et al.* Evaluation of methods for modeling transcription factor
470 sequence specificity. *Nat. Biotechnol.* **31**, 126–134 (2013).
- 471 32. Berger, M. F. *et al.* Compact, universal DNA microarrays to comprehensively
472 determine transcription-factor binding site specificities. *Nat. Biotechnol.* **24**,
473 1429–1435 (2006).
- 474 33. Mathelier, A. *et al.* JASPAR 2014: An extensively expanded and updated
475 open-access database of transcription factor binding profiles. *Nucleic Acids Res.*
476 **42**, D142–D147 (2014).
- 477 34. Matys, V. *et al.* TRANSFAC and its module TRANSCompel: transcriptional
478 gene regulation in eukaryotes. *Nucleic Acids Res.* **34**, D108 (2006).
- 479 35. Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription
480 factor sequence specificity. *Cell* **158**, 1431–1443 (2014).

- 481 36. Berger, M. F. *et al.* Variation in Homeodomain DNA Binding Revealed by
482 High-Resolution Analysis of Sequence Preferences. *Cell* **133**, 1266–1276 (2008).
- 483 37. Chen, L., Wu, G. & Ji, H. hmChIP: a database and web server for exploring
484 publicly available human and mouse ChIP-seq and ChIP-chip data.
485 *Bioinformatics* **27**, 1447–1448 (2011).
- 486 38. Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell*
487 **152**, 327–339 (2013).
- 488 39. Seb-Pedrs, A. *et al.* Early evolution of the T-box transcription factor family.
489 *Proc. Natl. Acad. Sci. U. S. A.* **110**, 16050–16055 (2013).
- 490 40. Huynh-Thu, V. A. & Geurts, P. DynGENIE3: Dynamical GENIE3 for the
491 inference of gene networks from time series expression data. *Sci. Rep.* **8**, (2018).
- 492 41. Gilbert, S. F. & Barresi, M. J. F. DEVELOPMENTAL BIOLOGY, 11TH
493 EDITION 2016. *Am. J. Med. Genet. Part A* **173**, 1430–1430 (2017).
- 494 42. Glass, C. K. & Rosenfeld, M. G. The coregulator exchange in transcriptional
495 functions of nuclear receptors. *Genes and Development* vol. 14 121–141 (2000).
- 496 43. Skoda, A. M. *et al.* The role of the hedgehog signaling pathway in cancer: A
497 comprehensive review. *Bosnian Journal of Basic Medical Sciences* vol. 18 8–20
498 (2018).
- 499 44. Dai, H., Jin, Q. Q., Li, L. & Chen, L. N. Reconstructing gene regulatory networks
500 in single-cell transcriptomic data analysis. *Zool. Res.* **41**, 599–604 (2020).
- 501 45. Chasman, D. & Roy, S. Inference of cell type specific regulatory networks on
502 mammalian lineages. *Current Opinion in Systems Biology* vol. 2 130–139 (2017).
- 503 46. Chen, S. & Mar, J. C. Evaluating methods of inferring gene regulatory networks
504 highlights their lack of performance for single cell gene expression data. *BMC*

505 *Bioinformatics* **19**, 232 (2018).

506 47. Tritschler, S. *et al.* Concepts and limitations for learning developmental
507 trajectories from single cell genomics. *Development (Cambridge)* vol. 146
508 (2019).

509 48. Kumar, M. P. *et al.* Analysis of Single-Cell RNA-Seq Identifies Cell-Cell
510 Communication Associated with Tumor Characteristics. *Cell Rep.* **25**,
511 1458-1468.e4 (2018).

512 49. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat.
513 *Nat. Commun.* **12**, 1–20 (2021).

514 50. Cabello-Aguilar, S. *et al.* SingleCellSignalR: Inference of intercellular networks
515 from single-cell transcriptomics. *Nucleic Acids Res.* **48**, 55 (2021).

516 51. Wang, S., Karikomi, M., Maclean, A. L. & Nie, Q. Cell lineage and
517 communication network inference via optimization for single-cell transcriptomics.
518 *Nucleic Acids Res.* **47**, 66 (2019).

519 52. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ
520 preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

521 53. Li, H. & Durbin, R. Fast and accurate short read alignment with
522 Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

523 54. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon
524 provides fast and bias-aware quantification of transcript expression. *Nat. Methods*
525 **14**, 417–419 (2017).

526 55. R Core Team. R: A language and environment for statistical computing. R
527 Foundation for Statis-tical Computing, Vienna, Austria.
528 <https://www.r-project.org/> (2017).

- 529 56. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *bioRxiv*
530 2020.10.12.335331 (2020) doi:10.1101/2020.10.12.335331.
- 531 57. Amezquita, R. A. *et al.* Orchestrating single-cell analysis with Bioconductor. *Nat.*
532 *Methods* **17**, 137–145 (2020).
- 533 58. Sing, T., Sander, O., Beerenwinkel, N. & Lengauer, T. ROCR: visualizing
534 classifier performance in R. *Bioinformatics* **21**, 3940–3941 (2005).
- 535 59. Akyol, T. Y. R Venn: An R package for set operations on multiple sets.
536 <https://cran.r-project.org/web/packages/RVenn/vignettes/vignette.html> (2019).
- 537 60. Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the
538 visualization of intersecting sets and their properties. *Bioinformatics* **33**,
539 2938–2940 (2017).

540

541

Figures

Figure 1 Pseudo-time analysis for time-course individual bulk RNA-Seq of mouse embryos in early development

(a) Transcriptomic trajectory of mouse embryos from E7.5 to E13.5. Each point on the PC1-PC2 plane indicates each individual RNA-Seq result. The solid line indicates an inferred trajectory by slingshot. (b) A scatterplot of pseudo-time and corresponding stage for each sample. Each point indicates each individual RNA-Seq result and is jittered along the Y-axis. (c) An example (*Fabp7*) of difference of gene expression

551 dynamics along stage and pseudo-time. The black lines indicate smoothing curves for
 552 each data. Sum of squared residuals (SSR) between observed and fitted values were
 553 calculated. (d) A scatterplot of SSR of normalized gene expression of all genes along
 554 stage and pseudo-time. The red line indicates the same value between stage and
 555 pseudo-time.

556

557

558 **Figure 2. Inference of downstream genes based on the values in A**

559 (a) Selection of a parameter D for SCODE. The sum of squared residuals (SSR) for each
560 D (from 1 to 10 by 0.5) was calculated. We used $D = 4$. (b) Pearson's correlation
561 coefficients between the values of each A from 20 optimizations and the mean A , which
562 is the average of each value from all optimizations. (c) Calculation of Pearson's
563 correlations between the average expression level of each gene and the absolute values
564 in each row and column of A . (d) The violin plots of Pearson's correlations (e, f) Scatter
565 plot of the average expression level of each gene and the absolute values in the column
566 (e) and row (f) of A for *Sox8*. (g) An example of the definition of thresholds for
567 significant regulatory relationship between the regulator and downstream genes. A
568 scatter plot of the values of A for a regulator, *Sox8*, in decreasing order. The solid line

569 indicates a regression for the scatter plot. The dashed lines indicate the defined
570 thresholds.

571

Figure 3. Validation of the inferred regulatory network of transcription factors with the TF2DNA database

(a) Scatter plot of area under the curve (AUC) target selection for each transcription factor (TF) based on the absolute values of A inferred using SCODE and weight.matrix inferred using dyGENIE3. (b) Scatter plots of the validated inferred target gene rate of *E2f3* and *Arid5b* in the TF2DNA database. Downstream genes of TFs were selected based on the absolute values of A in decreasing order. Solid lines indicate genes at the

580 thresholds. (c) Bar graph of validated and non-validated downstream genes of each TF
 581 in the TF2DNA database, in decreasing order of the total number of inferred
 582 downstream genes. Only the top 30 TFs are shown. (d) Scatter plot of the validated
 583 target gene rate of the inferred downstream genes of TF j and the background rate of
 584 target genes in the TF2DNA database. Histograms show the distribution of validated
 585 target gene rates of inferred downstream genes and the background rates of downstream
 586 genes in the TF2DNA database.

587

Figure 4. Overlap of inferred genes downstream of ligand-related and receptor-related genes

(a) Bar plot of the number of inferred downstream genes that are common and unique for each ligand-receptor pair. (b-d) Upset plots of inferred downstream genes that are positively and negatively regulated by the representative ligand-related and receptor-related genes. (b) Retinoic acid signaling pathway. (c) Delta/Notch signaling pathway. (d) Hedgehog signaling pathway.