Host cell amplification of nutritional stress contributes to persistence in *Chlamydia trachomatis*

Nick D. Pokorzynski^{1,2} & Rey A. Carabeo^{1*}

¹Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE, USA

²Current affiliation: Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT, USA

*Correspondence: rey.carabeo@unmc.edu

Persistence, a viable, but non-replicating state has been implicated in diseases caused by *Chlamydia trachomatis*. Multiple nutritional stressors produce a superficially similar “persistent” state, yet no systematic comparison has been made to determine their likeness. We employed host-pathogen dual RNA-sequencing under both iron- and tryptophan-starved conditions to gain insight into chlamydial persistence and identify contributions by the host cell. Analysis of the transcriptome of iron- or tryptophan-starved *Chlamydia* revealed a common “core” component and a stress-specific “accessory” subset. Despite the overall transcriptomic differences of host cells starved for either iron or tryptophan, both stressors induced persistence. A common metabolic consequence of the stressors was a reduction in intracellular GTP levels. Mizoribine inhibition of IMPDH1, which catalyzes the rate-limiting step in *de novo* guanine nucleotide synthesis reproduced to a similar extent GTP depletion, and inhibited chlamydial growth as expected for a pathogen that is auxotrophic for GTP. Thus, the reduction of guanine nucleotide synthesis manifests amplification of either iron or tryptophan starvation contributing to persistence. These findings illustrate that a nutritionally stressed host cell remains effective in arresting growth of *Chlamydia* by targeting metabolic pathways required by the pathogen.

INTRODUCTION

The dynamics of intracellular infection reflect the interaction of the pathogen and the host cell, with the outcome of the battle shaped by the competition between pathogen virulence and host counteractive measures (Prentice et al., 2007). A cytokine that tilts the balance towards the host is interferon-gamma (IFN γ), the effects of which are amplified by the JAK/STAT signaling pathway to induce a varied collection of responsive genes, including several anti-microbial effectors (Shtrichman and Samuel, 2001). In turn, pathogens have evolved to acquire strategies that attenuate or neutralize IFN γ (Finlay and McFadden, 2006). Interestingly, some of these anti-microbial activities could potentially compromise host cells in their ability to deal with infections. For example, differential regulation of iron acquisition and storage mechanisms designed to starve the pathogen of this metal would be expected to starve the host of iron (Abreu et al., 2020; Nairz et al., 2014). A similar scenario applies to tryptophan depletion mediated by the catabolizing enzyme indoleamine-2,3-dioxygenase (IDO1) (Byrne et al., 1986). While it starves intracellular pathogens for tryptophan, it concomitantly deprives the host of this essential amino acid. Here, we interrogate the response of the host cell subjected to models of nutrient starvation (e.g. iron starvation by chelation with 2,2-bipyridyl and tryptophan limitation via growth in tryptophan-depleted medium) typically used to simulate specific IFN γ -responsive anti-microbial effectors. Specifically, we sought to determine if under such growth conditions the host cell is compromised

in clearing the infection, and if the pathogen response reflects this compromised state.

A typical response of *Chlamydia trachomatis* (Ctr) to iron or tryptophan starvation is the establishment of “persistence”. *Chlamydiae* are distinguished by a biphasic developmental cycle that interconverts an infectious, non-replicative elementary body (EB) with a non-infectious, replicative reticulate body (RB) (AbdelRahman and Belland, 2005). *Chlamydiae* can disengage their normal developmental program and enter a persistent state in response to a wide array of stress (Panzzetta et al., 2018; Wyrick, 2010), including antibiotic treatment (Skilton et al., 2009), amino acid starvation (Beatty et al., 1994; Belland et al., 2003; Hatch and Ouellette, 2020) or biometal limitation (Ardisson et al., 2020; Raulston, 1997; Thompson and Carabeo, 2011). Chlamydial persistence shares many hallmarks with conventional bacterial persistence: for instance, persistent *Chlamydiae* are re-activatable (Beatty et al., 1995) and tolerant to bactericidal antibiotics (Phillips-Campbell et al., 2014; Wyrick and Knight, 2004). Yet *Chlamydia* lack most of the suspected genetic determinants (e.g. TA modules, stringent response) implicated in conventional models of bacterial persistence (Eisenreich et al., 2021) and display unique traits that imply a more fundamental process. For example, chlamydial persistence appears homogeneous, rather than stochastic, as all organisms in a given inclusion typically adopt characteristic aberrant bacterial morphology (Beatty et al., 1993; Belland et al., 2003; Matsumoto and Manire, 1970).

The prevailing view is that persistence is the result of the accumulated effects of the stressor on the pathogen.

For example, persistence resulting from iron starvation is thought to arise from the combined action of inactivating iron-dependent enzymes and the dysregulation of the iron-responsive regulon, both of which are expected to have pleiotropic effects on *Chlamydia* (Al-Younes et al., 2001; Raulston, 1997). Tryptophan starvation on the other hand is expected to reduce translation of proteins that are tryptophan-rich; and the resulting skewed proteome disrupts chlamydial growth and development (Ouellette et al., 2016). These are likely to be an oversimplification of the interaction between pathogen and host because the effects of the nutritional stress on the host cell, which itself deploys adaptive responses, is not considered. In other words, the nature of these adaptive responses might inform on the host cell priority, *i.e.* inhibit pathogen growth or survive the side effects of the anti-microbial effectors. It is also possible that these two priorities can coexist, and perhaps cooperate to clear infection effectively.

We therefore studied systematically compared tryptophan- and iron-starved *Ctr*-infected epithelial cells via host-pathogen dual RNA sequencing (RNA-seq). We find that the morphological and developmental fate of *Ctr* is dependent on the nutritional insult applied, but that these outcomes are independent of the treatment regimen, implying that *Ctr* senses and actively initiates a persistent state. RNA-seq analysis of the transcriptomes of both the infected epithelial cell and *Ctr* under these conditions revealed that while the transcriptome of the infected host cell is distinct to each stress, the chlamydial transcriptome demonstrated a high level of conservation between conditions but maintains a subset of stress-specific differentially expressed (DE) “accessory” genes. In addition, we report that the nutrient-deprived host cell appears to enhance its anti-bacterial activity, starving *Ctr* of the essential nucleotide guanosine triphosphate (GTP). This was a common metabolic consequence of both iron- and tryptophan starvation of host cells, regardless of infection. The pathogen responded accordingly by up-regulating genes for GTP-dependent cofactor biosynthesis. Treatment of infected cells with mizoribine, a specific inhibitor of IMPDH1 reproduced both the growth defect and the chlamydial upregulation of riboflavin and folate biosynthesis genes. Together, these data support the view that a nutritionally stressed cell is not compromised in its ability to handle infection by an intracellular pathogen. That GTP depletion is a common metabolic consequence of the two stresses indicate that amplification of the primary stress could be an intrinsic property of host cells. This feature of the host cell should be considered as we build a more detailed picture of host-pathogen interactions using accepted methods of inducing nutritional stress in intracellular pathogens.

RESULTS

Alternate models of chlamydial persistence produce discernable phenotypic differences.

To establish models of chlamydial persistence, we subjected *Ctr*-infected HeLa cells to iron starvation, by treatment with the membrane permeable iron chelator 2,2-bipyridyl (BPD) (Thompson and Carabeo, 2011), or

tryptophan starvation, by culturing in a defined medium lacking tryptophan (TRP) (Pokorzynski et al., 2020, 2019). We applied two treatment regimens (Figure 1A), one which started at the time of infection and continued for 24 hours (h; BPD24, TRP24), reflecting established models of chlamydial persistence, or one that began at 8h post-infection (hpi) and continued for 16h (BPD16, TRP16), thereby allowing *Ctr* to establish a productive infection and differentiate into the replicative RB state prior to nutrient starvation. We then assayed several physiological hallmarks of chlamydial persistence to identify differences between nutrient stress or treatment regimen. We first assayed chlamydial morphology by immunofluorescent confocal microscopy (Figure 1B). In comparison to the untreated control (UTD24), all treatments produced qualitatively smaller inclusions, implying inhibited growth. We observed clear morphological differences between BPD and TRP, with BPD inclusions being occupied by aberrantly enlarged organisms while TRP inclusions displayed an “indiscrete” morphology, obscuring the observation of individual bacteria. We then analyzed genome copy number under each condition and observed that while all treatments significantly reduced genome equivalents compared to UTD24, no differences were statistically distinguishable between BPD and TRP (Figure 1C). In contrast, we found that TRP was more permissive to the generation of infectious progeny (Figure 1D), as BPD reduced recoverable inclusion forming units (IFUs) below the calculated limit of detection. Thus, while a comparable number of genome equivalents, and by extension chlamydial organisms, exist under each treatment condition, whether those chlamydiae can complete their developmental cycle is influenced by the model of persistence employed.

We then analyzed the expression of a panel of chlamydial genes commonly used to indicate developmental dysregulation by reverse transcription quantitative PCR (RT-qPCR): *trpRBA*, *ytgA*, *omcB* and *ompA* (Figure 1E). Consistent with previous reports, we observed that the tryptophan salvage operon, *trpRBA*, was significantly up-regulated in both BPD and TRP, though TRP increased *trpRBA* expression significantly compared to BPD. This is due to dual regulation by the tryptophan-dependent transcriptional repressor TrpR and the iron- and tryptophan-dependent repressor YtgR (Pokorzynski et al., 2020, 2019). In contrast, the expression of the periplasmic iron-binding protein *ytgA*, another gene regulated by YtgR (Thompson et al., 2012), was not significantly altered by any treatment condition, despite prior reports of its iron- and tryptophan-dependent induction (Pokorzynski et al., 2020; Thompson and Carabeo, 2011). Here, we utilized a transcriptome-based normalization method (see Materials and Methods, Figure 1 - Supplementary Text, Figure 1 – Figure Supplement 1), rather than normalizing to genome equivalents. This indicates that while YtgR regulation of *trpRBA* serves to increase the expression of the operon relative to the total transcriptome, YtgR regulation of *ytgA* maintains a constant proportion of transcripts across various developmental or nutritional conditions. We confirm that the expression of *omcB*, a cysteine-rich outer membrane protein associated with differentiation to the EB stage and common

Figure 1. Alternative models of chlamydial persistence are not phenotypically identical. (A) Diagram depicting the experimental design used throughout this study, where 24h treatment conditions (UTD24, BPD24, TRP24) begin at the time of infection, and 16h treatments begin at 8 hours post-infection (BPD16, TRP16). (B) Immunofluorescent confocal microscopic analysis of chlamydial morphology under various models of chlamydial persistence. Micrographs are representative of at least three independent biological replicates (N = 3). Chlamydial organisms were detected by immunostaining against the cytosolic Hsp60 homologs, GroEL_1-GroEL_3. Nuclei were detected by staining with DAPI. Arrowheads indicate aberrantly enlarged bacteria. (C) Determination of genome equivalents by quantitative PCR against the *euo* locus under the various persistence models. (D) Measurement of infectious progeny generation during nutritional stress in a reinfection assay. Dotted line indicates calculated limit of detection for the assay. ND = not detected. (E) Gene expression profiles for various nutritionally- or developmentally-regulated chlamydial genes. All plots represent the mean and standard deviation of three independent biological replicates (N = 3). Statistical significance in all panels was determined by one-way ANOVA followed by Tukey's post-hoc test of honestly significant differences (two-tailed). * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, ns = not significant.

biomarker of persistence, is down-regulated by nutritional stress, as is the expression of *ompA*, the major outer membrane protein in *Ctr*. While some statistically distinguishable differences exist between stress conditions, we find that these are differences in magnitude but not the directionality of expression. We conclude that while both stimuli induce physiological changes characteristic of persistence, iron starvation has a more profound effect on chlamydial physiology than media-defined tryptophan starvation. While more extreme treatment conditions may reveal a higher degree of concordance between the physiological endpoints of various persistence models, our results nevertheless underscore that different persistence models produce varied results.

***Chlamydia* initiates a common transcriptional program in response to distinct nutritional stressors.**

To characterize the chlamydial response systematically and unbiasedly to both tryptophan and iron starvation, we turned to dual RNA-seq to elucidate the persistent chlamydial transcriptome in relation to the transcriptional response of the infected host cell. Our experiment was designed to capture the presumably small fraction of chlamydial transcripts produced in persistently infected HeLa cells, and we accordingly used a multiplicity of infection (MOI) of five for each condition. In agreement with recent reports, we recovered high levels of chlamydial transcripts in each library (Hayward et al., 2021), with no fewer than 7.5×10^6 mapped reads under any condition. We also note that across replicates, no more than 0.01% of the total library mapped to the chlamydial reference genome in any mock-

Figure 2. The persistent chlamydial transcriptome is broadly conserved across conditions. (A) Principal component analysis (PCA) of the chlamydial transcriptomes derived from the various nutrient-deprived conditions. All data are from three independent biological replicates ($N = 3$). (B) Volcano plots of the differentially expressed (DE) genes under each persistence-inducing condition (adjusted p -value < 0.05 , $|\text{Fold change}| > 1.5$). Red = DE, blue = $p < 0.05$, $|\text{FC}| < 1.5$, green = $p > 0.05$, $|\text{FC}| > 1.5$, grey = $p > 0.05$, $|\text{FC}| < 1.5$. (C) Dot plots for enriched pathways identified by clusterProfiler in BPD24 and TRP24. Dot size reflects the number of genes enriched in the pathway and dot color indicates the statistical significance of pathway enrichment. (D) Ridge plots for enriched pathways identified by clusterProfiler in BPD24 and TRP24. Ridges represent normalized density of genes plotted against their $\log_2$ fold change. Ridge color reflects statistical significance of pathway enrichment.

infected sample, indicating a negligible influence of cross-aligned reads in our samples.

Analysis of our RNA-seq datasets revealed that the transcriptional response of *Ctr* is largely conserved across conditions. Complete details of the differential expression analysis can be found in Figure 2 – Supplemental Data 1. Principal component analysis of the variation between chlamydial transcriptomes demonstrated clustering of all treated conditions, albeit with variation between replicates (Figure 2A). Nevertheless, this indicated that the chlamydial transcriptome associated with different stressors and treatment regimens is highly similar. This is further emphasized by surveying the landscape of statistically significant DE genes (adjusted *p*-value < 0.05, |FC| > 1.5), where we observe all conditions produce a more significantly down-regulated than up-regulated set of genes (Figure 2B). Notably, we observe consistently strong down-regulation of virulence-associated genes such as the polymorphic outer membrane protein, *pmpG* (Becker and Hegemann, 2014; Carrasco et al., 2011), and the histone-like, nucleoid condensing genes *hctA* and *hctB* (*hct2*) (Barry III et al., 1993; Belland et al., 2003; Hackstadt et al., 1991). Furthermore, we note that RNA-seq reproduces the substantial up-regulation of *trpRBA* in TRP as observed by RT-qPCR (see Figure 1E). Thus, our RNA-seq data recapitulates expected trends based on previous gene expression studies in *Ctr*.

Next, we subjected the sets of DE genes from each condition to KEGG pathway gene set enrichment analysis (GSEA). Due to the gross similarity between treatment regimens, we limit the analysis here to BPD24 and TRP24 but provide complete details of the analysis for each condition in Figure 2 – Supplemental Data 2. Among the significantly enriched pathways identified, we observed a high level of agreement between treatments, with the pathways “Ribosome” (ctb03010) and “Aminoacyl-tRNA biosynthesis” (ctb00970) significantly enriched in all conditions and “Bacterial secretion system” (ctb03070) significantly enriched in three of four conditions (Figure 2C). Consistent with previous studies, “Ribosome” and “Aminoacyl-tRNA biosynthesis” pathways are activated, while the virulence-associated “Bacterial secretion system” pathway is suppressed (Figure 2D) (Belland et al., 2003). Interestingly, GSEA also revealed the unique enrichment of categories specific to a given treatment. BPD24 significantly activates the “Biosynthesis of cofactors” (ctb01240) pathway, whereas TRP24 results in significant enrichment of the “Biosynthesis of amino acids” (ctb01230) pathway (Figure 2, C and D). While the enrichment of “Biosynthesis of amino acids” is likely driven by the substantial up-regulation of the *trpRBA* operon in TRP24, the identification of functionally unique pathways under both nutrient conditions implies that while the overarching transcriptome may be similar, *Ctr* possess the ability to tailor their transcriptional response to a given stress.

The “core” persistent transcriptome is decorated by differentially expressed “accessory” genes unique to each nutritional condition.

Characterization of the chlamydial transcriptome across multiple nutritional conditions revealed a conserved, “core”

persistent transcriptome in *Ctr*. This core persistent transcriptome consisted of 30.52% of the up-regulated genes (Figure 3A) and 56.7% of the down-regulated genes across conditions (Figure 3B). Network maps were generated from the core DE up- or down-regulated genes obtained from BPD24 and TRP24 (Figure 3, C and D). The full list of core DE genes can be found in Figure 3 – Supplemental Data 1. Within the up-regulated set of genes, a highly interconnected node emerged representing several ribosomal subunit and translation adaptor genes (Figure 3C), consistent with previous reports on the persistent chlamydial transcriptome (Belland et al., 2003). However, it is unclear if this up-regulation leads to increased translational activity (Ouellette et al., 2006). Rather, this may reflect a preparation for reactivation such that translation can rapidly resume once conditions improve. In the down-regulated set of core genes, a more disconnected network was produced, though distinct nodes emerged, including virulence genes (e.g. *tarP*, *hctA*, *pmpE*, *omcB*) and type III secretion system components (e.g. *sctN*, *sctQ*, *sctU*) (Figure 3D). These data collectively confirm previous findings on chlamydial persistence, but provide a more detailed view of the conserved transcriptional stress response in *Ctr*.

Despite the high degree of transcriptional conservation, *Ctr* maintained many DE “accessory” genes that were treatment-specific (Figure 3, A and B). Interestingly, a higher proportion of the up-regulated genes were unique, with 10.33% and 12.21% of all up-regulated genes being unique to BPD24 and TRP24, respectively (Figure 3A), compared to only 4.91% of down-regulated genes for both conditions (Figure 3B). Network plots were generated for the uniquely up-regulated genes for BPD and TRP (Figure 3, E and F). The complete list of up- and down-regulated accessory genes can be found in Figure 3 – Supplemental Data 2. Among the recognized genes, few associations were derived for BPD, though a small node emerged apparently related to energy transduction by ATP synthase (Figure 3E). In contrast, multiple small nodes emerged in TRP, including one for inclusion membrane proteins (Incs; *incC*, *incB*, *ct228/ctl0480*), transcriptional regulators (*nrdR*, *trpR*, *euo*) and a node comprising translation (*pheT*, *rpsT*), transcription (*rpoD*) and genome replication (*gyrB*, *gyrA2*) components (Figure 3F). We selected two accessory genes from each condition to confirm their differential expression by RT-qPCR: for BPD, we selected the peptidoglycan (PG) amidase, *amiA*, and the beta subunit of ATP synthase, *atpB* (*ctl0559*), and for TRP we selected the diaminopimelate aminotransferase *dapL* and the tyrosine/tryptophan amino acid transporter *tyrP* (*tyrP_1*). We confirmed that *amiA* was uniquely up-regulated in BPD24 and that *tyrP* was significantly up-regulated in TRP24 relative to BPD24, however *dapL* was significantly up-regulated in BPD24 compared to TRP24 and *atpB* expression was up-regulated in both conditions compared to UTD24 but not significantly different between conditions (Figure 3 – Figure Supplement 1). Thus, we conclude that PG turnover is uniquely up-regulated by BPD treatment which may explain the distinct bacterial morphology observed under this condition (see Figure 1B) (Brockett and Liechti, 2021).

Figure 3. The core persistent chlamydial transcriptome is decorated by unique accessory genes in each condition. (A) Venn diagram of all differentially up-regulated genes across the various nutritional conditions. Color shading reflects the relative number of genes in each cross-section of the Venn diagram. **(B)** Same as in **(A)** for all differentially down-regulated genes. **(C)** Network plot of the core up-regulated genes in BPD24 and TRP24. Network consists of those genes that were recognized by the STRING-db server. **(D)** Same as in **(C)** for the core down-regulated genes. **(E)** Network plot of the up-regulated accessory genes across BPD conditions. Network consists of those genes recognized by the STRING-db server. **(F)** Same as in **(E)** for the TRP conditions.

The transcriptional response of the infected epithelial cell is dependent on the nutritional condition.

We hypothesized that instances of unique transcriptional responses by *Ctr* could be correlated with specific variations in the transcriptional responses of the infected epithelial cell. Dual RNA-seq produced HeLa transcriptomes with no fewer than 3.3×10^7 mapped reads in any condition. We conducted a similar analysis for the host cell transcriptomes as was performed on the chlamydial transcriptomes. Principal component analysis of the HeLa transcriptomes revealed that, unlike *Ctr*, clustering was dependent on the treatment condition applied, with BPD forming a distinct cluster and TRP clustering with UTD24 (Figure 4A). All infected conditions clustered distinctly compared to the mock-infected control group (Mock). Subsequent analysis of differential gene expression (adjusted *p*-value < 0.05, |FC| > 2) was performed by analyzing UTD24 in reference to Mock, whereas all nutrient-starved samples were analyzed in reference to UTD24 to account for the influence of infection. Complete details of the differential expression analysis can be found in Figure 4 – Supplemental Data 1. We observed that compared to UTD24, all treatment conditions altered the global transcriptional profile, resulting in a more significantly down-regulated portion of the DE genes (Figure 4B). Whereas UTD24 resulted in a down-regulated set of 145 genes compared to Mock, no fewer than 630 genes were down-regulated under any nutrient-deprived condition, when compared to UTD24 (Figure 4 – Supplemental Data 1). We note the significant up-regulation of genes previously identified to respond to acute and persistent chlamydial infection, such as the antiviral protein OASL, in UTD24, TRP16 and TRP24 (Eickhoff et al., 2007). Having accounted for differentially expressed genes due to infection, the remaining differences in gene expression could be assigned confidently to the host response to nutritional stress or unique activities of persistent chlamydiae.

Next, we performed GSEA to identify differentially regulated pathways between conditions. Due to the similarity between treatment regimens (Figure 4A), here we display only the results from the analysis of the 24h conditions but results for all conditions are provided in Figure 4 – Supplemental Data 2. We sorted pathways based on their enrichment score and plotted the results for the 15 most enriched pathways (Figure 4, C and D). We found that both BPD24 and TRP24 shared with UTD24 the enrichment of pathways related to infection, but also displayed uniquely enriched pathways – most notably in BPD24. The most enriched pathways in BPD24 comprised various categories, such as “Focal adhesion” (hsa04510), “PI3K-Akt signaling pathway” (hsa04151) and “Calcium signaling pathway” (hsa04020) (Figure 4C). We next inferred relatedness of the various enriched pathways by generating enrichment network maps for each condition (Figure 4D). We find that both UTD24 and TRP24 produced highly interconnected networks that form nodes related to the response to infection. In contrast, BPD24 produced a disconnected network, indicating functionally disjointed iron-responsive biological processes were induced under this condition. These data suggest that intracellular bacteria such as *Ctr* will encounter highly disparate host cell responses during nutritional stress, which we expect to have varying

influences on the growth and development of resident chlamydiae.

Suppression of host cell purine metabolism is reflected by the induction of GTP-dependent cofactor biosynthesis in persistent chlamydiae.

The distinct transcriptional profiles of the infected host cell prompted further investigation into the relevance of functional pathways downstream of the primary (*i.e.* iron or tryptophan starvation) stresses. We extended our analysis of the HeLa GSEA data to include significantly suppressed pathways, which we found to differ between TRP and BPD. Suppressed pathways uniquely associated with BPD24 (Figure 5A, Figure 4 – Supplemental Data 2), included “Purine metabolism” (hsa00230), which was intriguing given that *Ctr* is auxotrophic for GTP (McClarty and Fan, 1993; Tipples and McClarty, 1993). In addition to DNA replication, transcription, and translation, GTP is also an essential substrate for chlamydial riboflavin and tetrahydrofolate (THF) biosynthesis (Adams et al., 2014). Because these two pathways fall within the broader “Biosynthesis of cofactors” pathway identified as significantly enriched by *Ctr* in BPD24 (Figure 2, C and D), we hypothesized that this enrichment is linked to the suppression of host GTP biosynthesis. We extracted the expression data for both the host and chlamydial pathways from the RNA-seq datasets for the BPD24 and TRP24 conditions and observed that host purine metabolism was markedly down-regulated transcriptionally in BPD24 compared to TRP24, particularly along the biosynthetic arm leading from ribose-5-phosphate to inosine monophosphate (Figure 5 – Figure Supplement 1, A and B).

In Figure 5B and Figure 5C, we depict the chlamydial riboflavin and THF biosynthetic pathways, simplified for clarity, with color-coded expression data for each gene annotated in the pathway. We then qualitatively compared the normalized expression of the two pathways between BPD24 and TRP24 to identify genes that were differentially regulated between the conditions. Genes within the riboflavin and THF biosynthetic pathways are broadly up-regulated in both BPD24 and TRP24, though differences exist in key genes. For example, the promiscuous enzyme *trpC* (*trpF*), which shunts the product of the *ribA* GTP cyclohydrolase towards THF biosynthesis (Adams et al., 2014), appears differentially regulated between the BPD24 and TRP24 conditions (Figure 5C). Thus, the pathway-level identification of suppressed GTP biosynthesis in the host under BPD24 correlated with distinct gene expression patterns in the iron-starved chlamydiae. We therefore hypothesized that the reduction of GTP levels could be “sensed” by *Ctr*, yielding increased expression of genes involved in the GTP-requiring riboflavin and tetrahydrofolate biosynthetic pathways (Figure 5D). We note the intriguing finding, however, that both the riboflavin and THF biosynthetic pathways were generally upregulated in TRP24 (Figure 5, B and C), despite a less pronounced effect on the expression of the host purine metabolism pathway under this condition (Figure 5 – Supplemental Figure 1B). Given that pathway level analysis of gene expression is insufficient to infer actual alterations to metabolism, we addressed this apparent inconsistency with targeted pharmacological and biochemical assays.

Figure 4. The infected host cell transcriptome differs between models of chlamydial persistence. (A) Principal component analysis (PCA) of the HeLa transcriptomes derived from the various experimental conditions. All data are N = 3. **(B)** Volcano plots of the differentially expressed (DE) genes under each persistence-inducing condition (adjusted p -value < 0.05 , $|\text{Fold change}| > 2.0$). Note that UTD24 is relative to the mock-infected condition (Mock) whereas all other conditions are relative to UTD24. Red = DE, blue = $p < 0.05$, $|\text{FC}| < 2.0$, green = $p > 0.05$, $|\text{FC}| > 2.0$, grey = $p > 0.05$, $|\text{FC}| < 2.0$.

Figure 5. Suppressed host cell purine metabolism is reflected by an up-regulation of GTP-dependent biosynthetic pathways in *Chlamydia trachomatis*. (A) Dot plot for the ten most activated or suppressed pathways identified by clusterProfiler in BPD24. Dot size reflects the number of genes enriched in the pathway and dot color indicates the statistical significance of pathway enrichment. (B) Simplified depiction of the chlamydial riboflavin biosynthetic pathway with color-coded normalized gene expression data for each gene represented by adjacent circles (BPD24) or diamonds (TRP24). DARPP = 2,5-diamino-6-(5-phospho-D-ribosylamino)-pyrimidin-4(3H)-one, ArPP = 5-amino-6-(5-phospho-D-ribosylamino)uracil, ArP = 5-amino-6-(5-phospho-D-ribitylamino)uracil, Ru5P = ribulose-5-phosphate, DHPB = 3,4-dihydroxy-2-butanone 4-phosphate, DRL = 6,7-dimethyl-8-ribityllumazine, FAD = flavin adenine dinucleotide. (C) Same as in (B) for the chlamydial tetrahydrofolate biosynthetic pathway. DARPP = 2,5-diamino-6-(5-phospho-D-ribosylamino)-pyrimidin-4(3H)-one, H₂NMP = 7,8-dihydroneopterin 3-phosphate, H₂N = 7,8-dihydroneopterin, HMH₂N = 6-hydroxymethyl-7,8-dihydropterin, H₂Pt = 7,8-dihydro-pterate, DHF = 7,8-dihydrofolate, THF = 5,6,7,8-tetrahydrofolate. (D) Proposed model of reciprocal up-regulation of chlamydial riboflavin and THF biosynthesis in the presence of the iron-dependent suppression of purine metabolism. Created with BioRender.com.

Pharmacological inhibition of host cell GTP biosynthesis by mizoribine induces chlamydial persistence.

Validation of our model that *Ctr* could sense and respond to GTP deprivation required an independent means of specifically depleting the host GTP pool. We therefore turned to mizoribine (MIZ), a selective inhibitor of inosine monophosphate dehydrogenase (IMPDH1/2) and guanosine monophosphate synthase (GMPS) (Turka et al., 1991; Yokota, 2002). MIZ is a potent inhibitor of mammalian GTP biosynthesis (Hoxhaj et al., 2017) and therefore provided a suitable tool to evaluate whether GTP starvation alone influenced chlamydial development and gene regulation. We treated *Ctr*-infected HeLa cells with a two-fold dilution series of MIZ for 24h starting at the time of

infection. Chlamydial morphology and genome replication were acutely sensitive to MIZ, with a perceptible reduction in inclusion size (Figure 6A) and significant decrease in genome copy number (Figure 6B) detectable at the lowest concentration of MIZ tested, 12.5 μ M. However, we sought to determine whether MIZ treatment could induce a persistent state in *Ctr*. We observed that by 100 μ M MIZ treatment, abnormal chlamydiae could be detected by confocal immunofluorescent microscopy (Figure 6A) and furthermore that genome equivalents at this concentration were not statistically distinguishable from either the 50 or 200 μ M treatments (Figure 6B), suggesting that genome replication had been stalled. Therefore, we moved forward with the 100 μ M MIZ treatment (hereafter MIZ24) and determined if *Ctr* retained viability when a reactivation (*i.e.*

Figure 4 cont. (C) Dot plots for the 15 most enriched pathways identified by clusterProfiler in UTD24, BPD24 and TRP24. Dot size reflects the number of genes enriched in the pathway and dot color indicates the statistical significance of pathway enrichment. (D) Enrichment network map for the 15 most enriched pathways identified by clusterProfiler in UTD24, BPD24 and TRP24. Dot size and color are the same as in (C).

Figure 6. Mizoribine treatment induces chlamydial persistence. (A) Immunofluorescent confocal microscopic analysis of chlamydial morphology following a two-fold dilution series of mizoribine (MIZ) beginning at the time of infection. Micrographs are representative of at least three independent biological replicates (N = 3). Chlamydial organisms were detected by immunostaining against the cytosolic Hsp60 homologs, GroEL_1-GroEL_3. Nuclei were detected by staining with DAPI. Arrowheads indicate aberrantly enlarged bacteria. (B) Determination of genome equivalents under the same MIZ dilution series in (A) by quantitative PCR against the *euo* locus. (C) Measurement of infectious progeny generation during 100 μM MIZ (MIZ24) in a reinfection assay. Reactivation was allowed to proceed for 16h by replacement of MIZ-containing media with fresh media. Dotted line indicates calculated limit of detection for the assay. (D) Gene expression profiles for various nutritionally- or developmentally-regulated chlamydial genes during MIZ24 treatment. Statistical significance in all panels was determined by pairwise two-sided unpaired Welch's *t*-test for unequal variance. * = *p* < 0.05, ** = *p* < 0.01, *** = *p* < 0.001. All plots represent the mean and standard deviation of three independent biological replicates (N = 3).

withdrawal of the inhibitor) period of 16h was introduced (Figure 6C). Recoverable IFUs were reduced roughly 2700-fold in MIZ24 relative to UTD24 but recovered to a level only

3-fold lower than UTD24 after reactivation. Thus, *Ctr* adopts an aberrant morphology and stalls genome replication in the

presence of MIZ but retains viability, which are physiological hallmarks of chlamydial persistence.

Our transcriptomic analyses indicated that an analysis of gene expression could inform a more robust categorization of the persistent state induced by MIZ. We returned to the panel of genes analyzed in Figure 1E to compare the gene expression profile of MIZ24 to other well characterized persistence models. Unlike BPD and TRP, MIZ down-regulated the expression of the *trpRBA* operon, but reproduced the transcriptional profile of *ytgA*, *omcB* and *ompA* found under established persistence models (Figure 1E, Figure 6D). It is not surprising that *trpRBA* gene expression profiles were not replicated under this condition because the nutritional co-repressors (*i.e.* iron and tryptophan) were not expected to be negatively affected by MIZ. We additionally analyzed expression of the curated panel of accessory genes (Figure 3 – Figure Supplement 1) in the MIZ24 condition and found that while *atpB* was significantly down-regulated, *amiA*, *dapL* and *tyrP* were all significantly up-regulated (Figure 6 – Figure Supplement 1). The up-regulation of *amiA* and *dapL* implies that the persistent state induced by MIZ24, to a limited extent, may be more comparable to that of BPD24 than TRP24, potentially explaining the comparable bacterial morphology observed between MIZ24 and BPD24 (Figure 1B, Figure 6A).

The depletion of host cell GTP pools in models of persistence elicits specific gene expression profiles in the chlamydial riboflavin and THF biosynthetic pathways.

With the validation of MIZ as a tool for starving *Ctr* of GTP, we next assayed intracellular GTP levels in mock-infected or *Ctr*-infected HeLa cells during nutritional stress. To measure the host GTP pool, we adapted a commercially available kit for assaying GTPase activity (Figure 7A, see Materials and Methods). By comparing mock-infected samples, we observed that all nutrient-depleted conditions resulted in a reduction of GTP levels comparable to that observed with MIZ24 (Figure 7B). Interestingly, we observed that infection alone decreased intracellular GTP levels, which may reflect increased competition for this nucleotide between host and pathogen. However, infection could not further reduce the level of GTP from any nutrient-starved condition, suggesting that GTP was inaccessible to *Ctr*. Whether this is solely the effect of suppressed purine metabolism or if GTP sequestration or depletion occurs is unknown. Therefore, the host cell responds to nutrient limitation by depleting GTP pools, which negatively impacts *Ctr* as they compete for this critical nutrient to sustain replication and development. We note however that only BPD24 was able to reduce total luminescent output from this assay, which reflects the gross suppression of purine

metabolism identified by pathway-level analysis (Figure 7 – Figure Supplement 1, Figure 5A).

Finally, we assayed the expression of a subset of genes from the chlamydial riboflavin and THF biosynthetic pathways under our various nutrient-depleted conditions (Figure 7C), focusing on genes we identified as differentially expressed between BPD24 and TRP24 by RNA-seq (e.g. *ribF*, *trpC* and *ctl0875*, Figure 5, B and C), with *ribA* as the non-differentially regulated control. Indeed, relative to UTD24, the expression of *ribA* was significantly up-regulated in BPD24 and TRP24, but not statistically distinguishable between these groups. However, *ribF*, *trpC* and *ctl0875* all exhibited significantly higher expression in BPD24 compared to TRP24, suggesting that these pathways may respond to other signals specific to iron starvation. In comparison, we find that MIZ24 significantly increased expression of *ribA* and did not alter *trpC* expression (Figure 7D), consistent with the BPD24 condition. However, unlike BPD24, both *ribF* and *ctl0875* were significantly down-regulated by MIZ24, indicating a greater influence of iron starvation than GTP starvation on the expression of these pathways. This finding also implied that GTP starvation, while sufficient in the context of mizoribine treatment, is not the major mechanism by which iron starvation induces persistence. Rather, GTP depletion is one of many contributors to persistence. A broader implication is that the transcriptional signature of a secondary stress within the chlamydial transcriptome is not always readily apparent. Their identification requires co-analysis of the host cell transcriptome and functional validation of candidate secondary stress pathways.

DISCUSSION

Historically, models of nutritional stress during infection have operated under the assumption that downstream phenotypic consequences to the pathogen can be directly attributed to the depletion of the nutrient in question. We challenge this assumption by demonstrating that the nutrient-deprived host cell can deploy an unrelated antibacterial nutritional stress by suppressing GTP biosynthesis, essentially amplifying and diversifying the stress acting upon the pathogen. How this is accomplished is presently unclear. One could argue that chelation of iron by BPD disrupts the iron-sulfur cluster-containing amidophosphoribosyltransferase, PPAT (Itakura and Holmes, 1979). However, this biochemical explanation does not account for the broader transcriptional down-regulation of purine metabolism during BPD24 treatment or the reduction of GTP levels in the TRP24 condition. Thus, a more fundamental process seems to be at play – one that may ultimately benefit the host during persistent infection. In agreement with this is the observation that distantly related eukaryotes, such as the budding yeast *Saccharomyces cerevisiae*, also suppress purine biosynthesis in the absence of iron, despite lacking iron-dependent enzymes in these pathways (Philpott and Protchenko, 2008; Shakoury-Elizeh et al., 2003). Future studies of nutritional stress during intracellular bacterial infection may require a more careful examination of the effect on host cells and the ways in which these effects condition pathogen responses and survival.

The pro-inflammatory cytokine IFN γ has been attributed a major role in the host anti-chlamydial immune response (Finethy and Coers, 2016). Much of this anti-chlamydial activity has been explained by the IFN γ -mediated induction of IDO1, which catabolizes host cell tryptophan pools to starve *Ctr* of this essential amino acid (Beatty et al., 1994; Taylor and Feng, 1991). Whether IDO1 overexpression is sufficient to inhibit chlamydial growth has not been investigated. Instead, IDO1 relevance is based on tryptophan supplementation studies and subsequent rescue of normal growth (Beatty et al., 1994; Byrne et al., 1986). Similarly, studies with IDO1 inhibitors, such as levo-1-methyl-tryptophan (L-1MT) only led to a partial rescue of growth (Ibana et al., 2011). Both published results remain consistent with additional stresses distinct from tryptophan catabolism by IDO1 being involved in IFN γ -mediated growth inhibition. IFN γ has also been shown to suppress purine metabolism by inhibiting eIF4E expression to reduce translation of genes in the purine biosynthetic pathway in primary human macrophages (an effect that depends on IFN γ -mediated IDO1 induction) (Ivashkiv, 2018; Su et al., 2015). This would be analogous to GTP depletion being an outcome of tryptophan starvation. Thus, it may be that reducing purine nucleotide levels, and specifically the GTP pool, is an evolved and redundant immune response of the host cell.

Herein, we provide evidence that both primary and secondary effects of a stress contribute to chlamydial persistence. The character of the chlamydial transcriptome also supports the notion that a stressed host cell induces subsequent stresses. Growth of infected cells in iron- or tryptophan-depleted media yielded transcriptomic profiles consisting of a “core” and an “accessory” component. The latter is specific to the original stress, and could represent remnants of initial chlamydial response, only deteriorating as adverse effects accumulate. On the other hand, the core component could be associated with metabolic consequences that are common to iron and tryptophan starvation, of which reduction in GTP levels is an example. We emphasize that prominent differences remain in the transcriptome components, and a more careful and detailed study is needed to establish their relevance to chlamydial persistence. Based on our collective data, a picture emerges of persistence as a deceptively similar process underpinned by a response that has stress-dependent and -independent components. This response is shaped by the different actions of the primary stress on the pathogen and the host cell, the latter involving the induction of subsequent waves of metabolically oriented stressors that target the pathogen. In other words, despite using a single-stressor experimental model, subsequent stresses with antimicrobial functions are induced, forcing *Chlamydia* to adapt to not just one, but two or more simultaneous stresses. With a limited repertoire of stress adaptation strategies, *Chlamydiae* are likely more sensitive to these simultaneous stressors than other intracellular bacterial pathogens; this does not discount the potential relevance of such combined effects in these experimental systems but instead argues for their careful examination in future studies.

ACKNOWLEDGMENTS

We would like to acknowledge the members of the Carabeo laboratory for their critical feedback on the development of this project. We would also like to acknowledge the input of Dr. John Mishler-Elmore, Technical Services Scientist I at Promega Corporation for their invaluable feedback on the establishment of the GTPase-Glo assay kit to measure intracellular GTP levels. This work was supported by NIAID grant R01-AI132406 to RAC. The University of Nebraska DNA Sequencing Core receives partial support from the National Institute for General Medical Science (NIGMS) INBRE - P20GM103427-19 grant as well as The Fred & Pamela Buffett Cancer Center Support Grant - P30 CA036727. This publication's contents are the sole responsibility of the authors and do not necessarily represent the official views of the NIH or NIGMS.

AUTHOR CONTRIBUTIONS

NDP and RAC designed and analyzed the experiments. NDP performed the experiments. NDP and RAC wrote and edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no conflicts of interest.

REFERENCES

- AbdelRahman, Y.M., Belland, R.J., 2005. The chlamydial developmental cycle. *FEMS Microbiol. Rev.* 29, 949–959. <https://doi.org/10.1016/j.femsre.2005.03.002>
- Abreu, R., Essler, L., Giri, P., Quinn, F., 2020. Interferon-gamma promotes iron export in human macrophages to limit intracellular bacterial replication. *PLoS One* 15, e0240949.
- Adams, N.E., Thiaville, J.J., Proestos, J., Juárez-Vázquez, A.L., McCoy, A.J., Barona-Gómez, F., Iwata-Reuyl, D., de Crécy-Lagard, V., Maurelli, A.T., 2014. Promiscuous and adaptable enzymes fill “Holes” in the tetrahydrofolate pathway in *Chlamydia* species. *MBio* 5, 1–14. <https://doi.org/10.1128/mBio.01378-14>
- Al-Younes, H.M., Rudel, T., Brinkmann, V., Szczepek, A.J., Meyer, T.F., 2001. Low iron availability modulates the course of *Chlamydia pneumoniae* infection. *Cell. Microbiol.* 3, 427–437.
- Anders, S., Pyl, P.T., Huber, W., 2015. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31, 166–169. <https://doi.org/10.1093/bioinformatics/btu638>
- Ardissone, S., Scherler, A., Pilonel, T., Martin, V., Kebbi-Beghdadi, C., Greub, G., 2020. Transcriptional Landscape of *Waddlia chondrophila* Aberrant Bodies Induced by Iron Starvation. *Microorg.* <https://doi.org/10.3390/microorganisms8121848>
- Barry III, C.E., Brickman, T.J., Hackstadt, T., 1993. Hc1-mediated effects on DNA structure: a potential regulator of chlamydial development. *Mol. Microbiol.* 9, 273–283. <https://doi.org/10.1111/j.1365-2958.1993.tb01689.x>
- Beatty, W.L., Belanger, T.A., Desai, A.A., Morrison, R.P., Byrne, G.I., 1994. Tryptophan depletion as a mechanism of gamma interferon-mediated chlamydial persistence. *Infect. Immun.* 62, 3705–3711.
- Beatty, W.L., Byrne, G.I., Morrison, R.P., 1993. Morphologic and antigenic characterization of interferon gamma-mediated persistent *Chlamydia trachomatis* infection in vitro. *Proc. Natl. Acad. Sci. U. S. A.* 90, 3998–4002. <https://doi.org/10.1073/pnas.90.9.3998>
- Beatty, W.L., Morrison, R.P., Byrne, G.I., 1995. Reactivation of persistent *Chlamydia trachomatis* infection in cell culture. *Infect. Immun.* 63, 199–205. <https://doi.org/10.1128/iai.63.1.199-205.1995>
- Becker, E., Hegemann, J.H., 2014. All subtypes of the Pmp adhesin family are implicated in chlamydial virulence and show species-specific function. *Microbiologyopen* 3, 544–556. <https://doi.org/https://doi.org/10.1002/mbo3.186>
- Belland, R.J., Nelson, D.E., Virok, D., Crane, D.D., Hogan, D., Sturdevant, D., Beatty, W.L., Caldwell, H.D., 2003. Transcriptome analysis of chlamydial growth during IFN-gamma-mediated persistence and reactivation. *Proc. Natl. Acad. Sci. U. S. A.* 100, 15971–15976. <https://doi.org/10.1073/pnas.2535394100>
- Blighe, K., Rana, S., Lewis, M., 2021. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling. <https://doi.org/10.18129/B9.bioc.EnhancedVolcano>
- Brinkworth, A.J., Wildung, M.R., Carabeo, R.A., 2018. Genomewide Transcriptional Responses of Iron-Starved *Chlamydia trachomatis* Reveal Prioritization of Metabolic Precursor Synthesis over Protein Translation. *mSystems* 3, e00184-17.
- Brockett, M.R., Liechti, G.W., 2021. Persistence alters the interaction between *Chlamydia trachomatis* and its host cell. *Infect. Immun.* 0, IAI.00685-20. <https://doi.org/10.1128/IAI.00685-20>
- Byrne, G.I., Lehmann, L.K., Landry, G.J., 1986. Induction of tryptophan catabolism is the mechanism for gamma-interferon-mediated inhibition of intracellular *Chlamydia psittaci* replication in T24 cells. *Infect. Immun.* 53, 347–351.
- Caldwell, H.D., Kromhout, J., Schachter, J., 1981. Purification and partial characterization of the major outer membrane protein of *Chlamydia trachomatis*. *Infect. Immun.* 31, 1161–1176.
- Carrasco, J.A., Tan, C., Rank, R.G., Hsia, R., Bavoil, P.M., 2011. Altered developmental expression of polymorphic membrane proteins in penicillin-stressed *Chlamydia trachomatis*. *Cell. Microbiol.* 13, 1014–1025. <https://doi.org/https://doi.org/10.1111/j.1462-5822.2011.01598.x>
- Eickhoff, M., Thalmann, J., Hess, S., Martin, M., Laue, T., Kruppa, J., Brandes, G., Klos, A., 2007. Host Cell Responses to *Chlamydia pneumoniae* in Gamma Interferon-Induced Persistence Overlap Those of Productive Infection and Are Linked to Genes Involved in Apoptosis, Cell Cycle, and Metabolism. *Infect. Immun.* 75, 2853–2863. <https://doi.org/10.1128/IAI.01045-06>
- Eisenreich, W., Rudel, T., Heesemann, J., Goebel, W., 2021. Persistence of Intracellular Bacterial Pathogens—With a Focus on the Metabolic Perspective. *Front. Cell. Infect. Microbiol.*
- Engström, P., Bailey, L., Önskog, T., Bergström, S., Johansson, J., 2010. A comparative study of RNA and DNA as internal gene expression controls early in the developmental cycle of *Chlamydia pneumoniae*. *FEMS Immunol. Med. Microbiol.* 58, 244–253. <https://doi.org/10.1111/j.1574-695X.2009.00631.x>
- Finethy, R., Coers, J., 2016. Sensing the enemy, containing the threat: cell-autonomous immunity to *Chlamydia trachomatis*. *FEMS Microbiol. Rev.* 40, 875–893. <https://doi.org/10.1093/femsre/fuw027>
- Finlay, B.B., McFadden, G., 2006. Anti-Immunology: Evasion of the Host Immune System by Bacterial and Viral Pathogens. *Cell* 124, 767–782. <https://doi.org/https://doi.org/10.1016/j.cell.2006.01.034>
- Hackstadt, T., Baehr, W., Ying, Y., 1991. *Chlamydia trachomatis* developmentally regulated protein is homologous to eukaryotic histone H1. *Proc. Natl. Acad. Sci. U. S. A.* 88,

- 3937–3941. <https://doi.org/10.1073/pnas.88.9.3937>
- Hatch, N.D., Ouellette, S.P., 2020. Inhibition of tRNA Synthetases Induces Persistence in Chlamydia. *Infect. Immun.* 1–15. <https://doi.org/10.1128/iai.00943-19>
- Hayward, R.J., Humphrys, M.S., Huston, W.M., Myers, G.S.A., 2021. Dual RNA-seq analysis of in vitro infection multiplicity and RNA depletion methods in Chlamydia-infected epithelial cells. *Sci. Rep.* 11, 10399. <https://doi.org/10.1038/s41598-021-89921-x>
- Hoxhaj, G., Hughes-Hallett, J., Timson, R.C., Ilagan, E., Yuan, M., Asara, J.M., Ben-Sahra, I., Manning, B.D., 2017. The mTORC1 Signaling Network Senses Changes in Cellular Purine Nucleotide Levels. *Cell Rep.* 21, 1331–1346. <https://doi.org/https://doi.org/10.1016/j.celrep.2017.10.029>
- Ibana, J.A., Belland, R.J., Zea, A.H., Schust, D.J., Nagamatsu, T., AbdelRahman, Y.M., Tate, D.J., Beatty, W.L., Aiyar, A.A., Quayle, A.J., 2011. Inhibition of indoleamine 2,3-dioxygenase activity by levo-1-methyl tryptophan blocks gamma interferon-induced chlamydia trachomatis persistence in human epithelial cells. *Infect. Immun.* 79, 4425–4437. <https://doi.org/10.1128/IAI.05659-11>
- Itakura, M., Holmes, E.W., 1979. Human amidophosphoribosyltransferase. An oxygen-sensitive iron-sulfur protein. *J. Biol. Chem.* 254, 333–338. [https://doi.org/10.1016/s0021-9258\(17\)37922-x](https://doi.org/10.1016/s0021-9258(17)37922-x)
- Ivashkiv, L.B., 2018. IFN γ : signalling, epigenetics and roles in immunity, metabolism, disease and cancer immunotherapy. *Nat. Rev. Immunol.* 18, 545–558. <https://doi.org/10.1038/s41577-018-0029-z>
- Kim, D., Langmead, B., Salzberg, S.L., 2015. HISAT: a fast spliced aligner with low memory requirements. *Nat. Methods* 12, 357–360. <https://doi.org/10.1038/nmeth.3317>
- Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550. <https://doi.org/10.1186/s13059-014-0550-8>
- Luo, W., Brouwer, C., 2013. Pathview: an R/Bioconductor package for pathway-based data integration and visualization. *Bioinformatics* 29, 1830–1831. <https://doi.org/10.1093/bioinformatics/btt285>
- Matsumoto, A., Manire, G.P., 1970. Electron microscopic observations on the effects of penicillin on the morphology of Chlamydia psittaci. *J. Bacteriol.* 101, 278–285. <https://doi.org/10.1128/jb.101.1.278-285.1970>
- McClarty, G., Fan, H., 1993. Purine metabolism by intracellular Chlamydia psittaci. *J. Bacteriol.* 175, 4662–4669. <https://doi.org/10.1128/jb.175.15.4662-4669.1993>
- Nairz, M., Haschka, D., Demetz, E., Weiss, G., 2014. Iron at the interface of immunity and infection. *Front. Pharmacol.* 5. <https://doi.org/10.3389/fphar.2014.00152>
- Ouellette, S.P., Hatch, T.P., AbdelRahman, Y.M., Rose, L.A., Belland, R.J., Byrne, G.I., 2006. Global transcriptional upregulation in the absence of increased translation in Chlamydia during IFN γ -mediated host cell tryptophan starvation. *Mol. Microbiol.* 62, 1387–1401. <https://doi.org/10.1111/j.1365-2958.2006.05465.x>
- Ouellette, S.P., Rueden, K.J., Rucks, E.A., 2016. Tryptophan codon-dependent transcription in chlamydia pneumoniae during gamma interferon-mediated tryptophan limitation. *Infect. Immun.* 84, 2703–2713. <https://doi.org/10.1128/IAI.00377-16>
- Panzetta, M.E., Valdivia, R.H., Saka, H.A., 2018. Chlamydia Persistence: A Survival Strategy to Evade Antimicrobial Effects in-vitro and in-vivo. *Front. Microbiol.* 9, 1–11. <https://doi.org/10.3389/fmicb.2018.03101>
- Phillips-Campbell, R., Kintner, J., Schoborg, R. V., 2014. Induction of the Chlamydia muridarum Stress/Persistence Response Increases Azithromycin Treatment Failure in a Murine Model of Infection. *Antimicrob. Agents Chemother.* 58, 1782–1784. <https://doi.org/10.1128/AAC.02097-13>
- Philpott, C.C., Protchenko, O., 2008. Response to Iron Deprivation in Saccharomyces cerevisiae. *Eukaryot. Cell* 7, 20–27. <https://doi.org/10.1128/EC.00354-07>
- Pokorzynski, N.D., Brinkworth, A.J., Carabeo, R., 2019. A bipartite iron-dependent transcriptional regulation of the tryptophan salvage pathway in Chlamydia trachomatis. *Elife* 8, 1–32.
- Pokorzynski, N.D., Hatch, N.D., Ouellette, S.P., Carabeo, R.A., 2020. The iron-dependent repressor YtgR is a tryptophan-dependent attenuator of the trpRBA operon in Chlamydia trachomatis. *Nat. Commun.* 11, 6430. <https://doi.org/10.1038/s41467-020-20181-5>
- Prentice, A.M., Ghattas, H., Cox, S.E., 2007. Host-Pathogen Interactions: Can Micronutrients Tip the Balance? *J. Nutr.* 137, 1334–1337. <https://doi.org/10.1093/jn/137.5.1334>
- Raulston, J.E., 1997. Response of Chlamydia trachomatis serovar E to iron restriction in vitro and evidence for iron-regulated chlamydial proteins. *Infect. Immun.* 65, 4539–4547.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: An open-source platform for biological-image analysis. *Nat. Methods* 9, 676–682. <https://doi.org/10.1038/nmeth.2019>
- Shakoury-Elizeh, M., Tiedeman, J., Rashford, J., Ferea, T., Demeter, J., Garcia, E., Rolfes, R., Brown, P.O., Botstein, D., Philpott, C.C., 2003. Transcriptional Remodeling in Response to Iron Deprivation in Saccharomyces cerevisiae. *Mol. Biol. Cell* 15, 1233–1243. <https://doi.org/10.1091/mbc.e03-09-0642>
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., Ideker, T., 2003. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res.* 13, 2498–2504. <https://doi.org/10.1101/gr.1239303>
- Shtrichman, R., Samuel, C.E., 2001. The role of gamma interferon in antimicrobial immunity. *Curr. Opin. Microbiol.* 4, 251–259. [https://doi.org/https://doi.org/10.1016/S1369-5274\(00\)00199-5](https://doi.org/https://doi.org/10.1016/S1369-5274(00)00199-5)
- Skilton, R.J., Cutcliffe, L.T., Barlow, D., Wang, Y., Salim, O., Lambden, P.R., Clarke, I.N., 2009. Penicillin Induced Persistence in Chlamydia trachomatis: High Quality Time Lapse Video Analysis of the Developmental Cycle. *PLoS One* 4, e7723.
- Su, X., Yu, Y., Zhong, Y., Giannopoulou, E.G., Hu, X., Liu, H., Cross, J.R., Ratsch, G., Rice, C.M., Ivashkiv, L.B., 2015. Interferon- γ regulates cellular metabolism and mRNA translation to potentiate macrophage activation. *Nat. Immunol.* 16, 838–849. <https://doi.org/10.1038/ni.3205>
- Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva, N.T., Morris, J.H., Bork, P., Jensen, L.J., Mering, C. von, 2019. STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* 47, D607–D613. <https://doi.org/10.1093/nar/gky1131>
- Taylor, M.W., Feng, G.S., 1991. Relationship between interferon-gamma, indoleamine 2,3-dioxygenase, and tryptophan catabolism. *Faseb. J.* 5, 2516–2522.
- Thompson, C.C., Carabeo, R.A., 2011. An optimal method of iron starvation of the obligate intracellular pathogen, Chlamydia trachomatis. *Front. Microbiol.* 2. <https://doi.org/10.3389/fmicb.2011.00020>
- Thompson, C.C., Nicod, S.S., Malcolm, D.S., Grieshaber, S.S., Carabeo, R.A., 2012. Cleavage of a putative metal

- permease in *Chlamydia trachomatis* yields an iron-dependent transcriptional repressor. *Proc. Natl. Acad. Sci. U. S. A.* 109, 10546–51. <https://doi.org/10.1073/pnas.1201398109>
- Tipple, G., McClarty, G., 1993. The obligate intracellular bacterium *Chlamydia trachomatis* is auxotrophic for three of the four ribonucleoside triphosphates. *Mol. Microbiol.* 8, 1105–1114. <https://doi.org/10.1111/j.1365-2958.1993.tb01655.x>
- Turka, L.A., Dayton, J., Sinclair, G., Thompson, C.B., Mitchell, B.S., 1991. Guanine ribonucleotide depletion inhibits T cell activation. Mechanism of action of the immunosuppressive drug mizoribine. *J. Clin. Invest.* 87, 940–948. <https://doi.org/10.1172/JCI115101>
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., Speleman, F., 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* 3, research0034.1. <https://doi.org/10.1186/gb-2002-3-7-research0034>
- Wickham, H., 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
- Wyrick, P.B., 2010. *Chlamydia trachomatis* Persistence In Vitro: An Overview. *J. Infect. Dis.* 201, 88–95. <https://doi.org/10.1086/652394>
- Wyrick, P.B., Knight, S.T., 2004. Pre-exposure of infected human endometrial epithelial cells to penicillin in vitro renders *Chlamydia trachomatis* refractory to azithromycin. *J. Antimicrob. Chemother.* 54, 79–85. <https://doi.org/10.1093/jac/dkh283>
- Yokota, S., 2002. Mizoribine: Mode of action and effects in clinical use. *Pediatr. Int.* 44, 196–198. <https://doi.org/https://doi.org/10.1046/j.1328-8067.2002.01536.x>
- Yu, G., Wang, L.-G., Han, Y., He, Q.-Y., 2012. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. *Omi. A J. Integr. Biol.* 16, 284–287. <https://doi.org/10.1089/omi.2011.0118>

MATERIALS AND METHODS

Data and materials availability

All sequencing data generated in this study have been deposited at the NCBI Gene Expression Omnibus (GEO; Accession number: GSE179003) and are publicly available as of the date of publication. All other source data and original code for the analysis of RNA-sequencing datasets and other experimental data have been deposited at Mendeley Data (DOI: 10.17632/vxvznn6bck.1) and are publicly available as of the date of publication. Microscopy data reported in this paper will be shared by the corresponding author upon request. Any additional information required to reanalyze the data reported in this paper is available from the corresponding author upon request.

Cell lines

Human female cervical epithelial adenocarcinoma HeLa cells (RRID: CVCL_1276) were cultured at 37° C with 5% atmospheric CO₂ in Dulbecco's Modified Eagle Medium (DMEM; Gibco, Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 10 µg/mL gentamicin, 2 mM L-glutamine, and 10% (v/v) filter sterilized fetal bovine serum (FBS). For all experiments, HeLa cells were cultured between passage numbers 3 and 15. HeLa cells were originally authenticated by ATCC via STR profiling and isoenzyme analysis per ATCC specifications.

Bacterial strains

Chlamydia trachomatis serovar L2 (434/Bu) was originally obtained from Dr. Ted Hackstadt (Rocky Mountain National Laboratory, NIAID). Chlamydial EBs were isolated from infected HeLa cells at 36–40 hr post-infection (hpi) and purified by density gradient centrifugation essentially as described (Caldwell et al., 1981). For infections, at 80–90% confluency, cells were first washed with Hanks Buffered Saline Solution (HBSS; Gibco, Thermo Fisher Scientific) and ice-cold inoculum prepared in HBSS at the indicated multiplicity of infection was overlaid onto the cell monolayer. To synchronize the infection, inoculated cells were then centrifuged for 15 min at 500xRCF, 4° C in an Eppendorf 5810 R tabletop centrifuge with an A-4-81 rotor. The inoculum was then aspirated and pre-warmed DMEM (or relevant media with treatment supplementation) was added to the cells. Infected cultures were then returned to the tissue culture incubator until the indicated time post-infection.

Treatment conditions

For iron starvation and media-defined tryptophan starvation, treatment was performed essentially as described previously (Pokorzynski et al., 2020, 2019). In brief, 100 mM 2,2-bipyridyl (Sigma Aldrich, St. Louis, MO, USA; CAS: 366-18-7) prepared in dimethyl sulfoxide (DMSO) was added to complete DMEM (or tryptophan-depleted DMEM-F12, as described below) at a working concentration of 100 µM at the start of infection (BPD24) or at 8 hpi (BPD16). When added after the time of infection, cells were first washed with HBSS prior to bipyridyl treatment. Tryptophan depletion was performed by first washing cells with HBSS and then replacing complete DMEM with tryptophan-depleted DMEM-F12 (U.S. Biological Life Sciences, Salem, MA, USA). Media was replaced either at the time of infection (TRP24) or at 8 hpi (TRP16). Treated cells were then returned to the tissue culture incubator for the remainder of the experimental time course. Mizoribine (Sigma Aldrich, CAS: 50924-49-7) was prepared as a 100 mM stock solution in DMSO, stored at -80° C, and used at the indicated concentrations starting at the time of infection (MIZ24).

Nucleic acid preparation

RNA was harvested from *C. trachomatis*-infected cells by scraping one or two wells of a 6-well tissue culture plate in a total volume of 500 µL Trizol Reagent (Thermo Fisher Scientific). Samples were transferred to RNase-free o-ring capped tubes containing ~100 µL volume of zirconia beads and thoroughly vortexed for 10 min to rupture bacterial cells. Zirconia beads were pelleted by centrifugation at 21,000xg for 10 min at 4° C and supernatant was transferred to an RNase-free tube containing 100 µL chloroform (Sigma Aldrich). Samples were vortexed for 15 s prior to a 10 min RT° C incubation. Phases were then separated by centrifugation at 21,000xg for 15 min at 4° C. The aqueous top layer was transferred to an RNase-free tube containing 250 µL 100% ethanol to precipitate RNA. Samples were briefly vortexed and then applied to an RNA collection column provided in the PureLink™ RNA Mini Kit (Invitrogen, Thermo Fisher Scientific). RNA was isolated as described by the manufacturer with an on-column DNA digestion using the PureLink™ DNase Set (Invitrogen, Thermo Fisher Scientific). RNA was eluted in nuclease-free H₂O and stored at -20° C for short-term storage or -80° C for long-term storage.

Complementary DNA (cDNA) was generated using 1–2 µg of RNA as a template for the SuperScript IV Reverse Transcriptase (RT) VILO master mix (Invitrogen, Thermo Fisher Scientific) with a no-RT control reaction in a half-reaction volume following manufacturer protocols. The no-RT control sample was screened for DNA contamination by qPCR against the *euo* locus (see Data S7 for full list of oligonucleotide primers).

Genomic DNA (gDNA) was harvested from parallel well(s) of a 6-well plate in 200 μ L ice-cold PBS + 10% Proteinase K and processed through the DNeasy Blood and Tissue Kit following manufacture protocols (QIAGEN, Hilden, Germany). gDNA was stored at -20° C for short-term storage or -80° C for long-term storage.

For the preparation of RNA-sequencing libraries, 10 μ g of RNA collected as described above, with an additional round of on-column DNA digestion, was processed in parallel 5 μ g aliquots through the RiboMinus™ Transcriptome Isolation Kit (Invitrogen, Thermo Fisher Scientific) essentially as described in the manufacturer protocol with the exception that the magnetic beads were loaded with 3 μ L of the pan-prokaryotic rRNA probe as well as 4 μ L of the eukaryotic rRNA probe to deplete both host and chlamydial rRNA simultaneously. The resulting rRNA-depleted samples were concentrated in the RNA Clean and Concentrator™ Kit (Zymo Research, Irvine, CA, USA) and submitted to the University of Nebraska DNA Sequencing Core for library preparation and RNA-sequencing.

Library preparation and RNA-sequencing

Submitted RNA samples were determined to be of suitable quality by fragment analysis on an Agilent 2100 Bioanalyzer (Agilent, Santa Clara, CA, USA). TruSeq Stranded Total RNA library preparation kit (Illumina, San Diego, CA, USA) was used to generate RNA-sequencing libraries following manufacturer protocols with a starting amount of 100 ng rRNA-depleted RNA. Depletion of rRNA in the TruSeq kit was performed by the addition of 2.5 μ L each of standard rRNA Removal Mix (RRM) or Prokaryotic RRM. Quality of prepared libraries was determined by concentration and fragment analysis as above. Libraries were sequenced on an Illumina NextSeq NS550 (75 bp single read high output flow cell) or NovaSeq 6000 (75 bp single read SP-100 flow cell). Across replicates, the proportion of sequenced bases with a quality score higher than 30 was at least 95%.

Host-pathogen RNA-sequencing analysis

Transcriptomes were processed using the Galaxy server, version 21.05.1 (usegalaxy.org). Individual sequencing files for each condition within a replicate were concatenated and processed using the fastp application to filter low quality reads, trim reads and cut adapter sequences. Sequences were then aligned to either the *Chlamydia trachomatis* 434/Bu (ASM6858v1) genome assembly or the *Homo sapiens* GRCh38 genome assembly using HISAT2 (Kim et al., 2015). Read counts were generated using htseq-count (Anders et al., 2015) and output files were exported and compiled for differential gene expression analysis by DESeq2 in R (Love et al., 2014). Principal component analysis was performed on the regularized log-transformed count data. Volcano plots were generated using the EnhancedVolcano R package (Blighe et al., 2021). Gene set enrichment analysis for KEGG pathways was conducted using the clusterProfiler R package (Yu et al., 2012). Mapping of gene expression data to KEGG pathways was performed using the Pathview package in R (Luo and Brouwer, 2013). Gene network maps were generated by submitting gene lists to the STRING database (Szklarczyk et al., 2019) and then formatting networks in Cytoscape (Shannon et al., 2003). Note that any chlamydial genes not recognized by STRING were automatically filtered out during analysis.

Quantitative PCR

All quantitative PCR (qPCR) assays were performed using Power Up™ SYBR™ Green Master Mix (Applied Biosystems, Thermo Fisher Scientific) essentially as previously described (Pokorzynski et al., 2020, 2019). In brief, cDNA was diluted 1:5-1:10 and gDNA was diluted 1:50-1:100 in nuclease-free H₂O (dilutions were identical within each experiment). The 2X PCR master mix was diluted to 1X in nuclease-free H₂O with specific primers diluted to 500 nM (see Supplementary Data 2 for complete list of primers). To 79 μ L of the master mix solution, 3.3 μ L of template (cDNA or gDNA) was added and then aliquoted into three 25 μ L technical replicate reactions in a 96-well optical plate. Reactions were analyzed on a QuantStudio™ 3 Real-Time PCR System with standard SYBR cycling conditions. All assays were performed with a melt-curve analysis to ensure specific product amplification across samples. Primer sets (Supplementary File 1) used in qPCR were validated against a standard curve of *C. trachomatis* L2 gDNA diluted from 2 $\times 10^{-3}$ to 2 $\times 10^0$ ng per reaction. C_t values generated from each experimental reaction were then fit to a standard curve and only primer sets with an efficiency of 100% $\pm$ 5% were used.

Genome equivalents (GE) were calculated by first converting the mean C_t of the triplicate technical replicate reactions to a ng quantity of gDNA (ng template) with the linear equation generated from the standard curve of the *euo* primer pair. This value was then normalized to the total ng/ μ L gDNA isolated for each sample as follows:

$$GE = \frac{ng \text{ template}}{\frac{ng}{\mu L} \text{ gDNA}}$$

For the quantification of transcript expression by reverse transcription (RT)-qPCR, a transcriptome-based normalization was used based on the geometric average of multiple control genes, which were empirically determined using the geNorm method (Vandesompele et al., 2002). For more information on the geNorm analysis, see Supplemental Note 1. In brief, all transcript expression data was normalized to the geometric mean of the expression of *groEL_1*, *euo*, *nrdA* and *nrdB*. The $\Delta\Delta C_t$ method was then used to determine relative expression values and the log2-transformed fold change was analyzed to facilitate comparisons between conditions where the magnitude of gene expression changed considerably (e.g. expression of the *trpRBA* operon in iron- or tryptophan-starved conditions). Thus, transcript expression (TE) was calculated as:

$$TE = 2^{-[C_{tExp} - C_{tgeNorm}] - C_{tRef}}$$

Where C_t(Exp) is the C_t value of the experimental gene being analyzed, C_t(geNorm) is the geometric mean of the C_t values for the control genes, and C_t(Ref) is the mean C_t value of the reference condition, in this case UTD24. All C_t values were corrected for dilution prior to the computation of transcript expression.

Immunofluorescent confocal microscopy

To analyze inclusion morphology, HeLa cells were seeded onto acid-washed glass coverslips in 24-well tissue culture plates and infected at MOI = 5. At the indicated times post-infection, coverslips were washed with phosphate-buffered saline (PBS) and cells were fixed with 4% paraformaldehyde in PBS for 15 min at RT° C. Fixation solution was then aspirated and coverslips were either stored at 4° C in PBS or immediately processed for immunofluorescence assays by permeabilizing cells in PBS + 0.2% Triton X-100 (Thermo Scientific™) for 10 min at RT° C with rocking. Permeabilization solution was then decanted and coverslips were washed 3x with PBS. Coverslips were blocked in PBS + 3% bovine serum albumin (BSA) for 1 hr at RT° C with rocking. Coverslips were then washed 3x with PBS prior to being overturned on a 50 μ L droplet of PBS + 3% BSA containing primary antibody diluted 1:1000. To detect chlamydial GroEL, cells were stained with monoclonal mouse anti-cHsp60 (MA3-023, Invitrogen, ThermoFisher Scientific). Coverslips were incubated on primary antibody solution overnight at 4° C in an opaque humidified container. Coverslips were then washed thoroughly by repeated submersion (~50x) in 100 mL PBS before being overturned on a 50 μ L droplet of PBS + 3% BSA + 1:1000 secondary antibody + 2.5 μ g/mL 4',6-diamidino-2-phenylindole (DAPI) to label nuclei. A donkey anti-mouse AlexaFluor-594 secondary antibody (Invitrogen, Thermo Fisher Scientific) was used to label the primary mouse anti-cHsp60. Coverslips were then incubated for at least one hour at RT° C in an opaque humidified container prior to being washed as described above in Milli-Q H₂O and then being mounted on glass microscope slides with 10 μ L Shandon™ Immu-Mount (Thermo Fisher Scientific). Mounting medium was allowed to solidify overnight. Confocal microscopy was performed on a Nikon Ti2 Eclipse spinning-disk confocal microscope. All images were acquired using identical laser power and exposure settings. To enhance visualization of inclusion morphology, contrast and brightness were adjusted as necessary for each condition in Fiji ImageJ (Schindelin et al., 2012). All images are summed Z-projections of Z-stacks spanning the entire depth of the inclusions in the field.

Reinfection assay

At the indicated times post-infection for the relevant treatment conditions, infected cells were scraped into cell culture media and collected in 2 mL microcentrifuge tubes. Cell suspensions were then centrifuged at 21,000xg for 30 min at 4° C to rupture cells. The supernatant was aspirated and the cell pellet was resuspended in 500 µL of sterile-filtered Sucrose-phosphate-glutamate (SPG; 220 mM sucrose, 10 mM Na₂HPO₄, 4 mM KH₂PO₄, 5 mM Glutamic Acid) buffer. The resuspended cell lysate was centrifuged at 200xg for 5 min at 4° C to pellet cell debris. The supernatant was stored at -80° C and used to reinfect a confluent HeLa cell monolayer in one well of a 24-well tissue culture plate in a ten-fold dilution series starting at 10 or 100 µL of inoculum. At 24 hpi, the reinfected cells were fixed and stained as above for DAPI and GroEL and at the appropriate dilution for each condition, inclusions were enumerated per field (total of five fields per replicate) and the number of IFU per mL of inoculum was calculated. For reactivation, media containing mizoribine was removed at 24 hpi and the samples were incubated with fresh media for an additional 16 hours prior to sample collection. The limit of detection was calculated to be one inclusion identified per field at 100 µL of inoculum.

Measurement of intracellular GTP levels

Infected or mock-infected cells under the indicated treatment conditions were collected at 24 hpi by washing cells in 2 mL PBS, aspirating the wash buffer, and then scraping the cells into 250 µL 1% trichloroacetic acid (TCA) solution to precipitate macromolecular complexes. The lysate was centrifuged to collect precipitates and the supernatant was neutralized to pH ~7.5 with 20 µL 1 M Tris-HCl, pH 8.5 prior to storage at -80° C. Intracellular GTP levels were then measured using the GTPase-Glo assay kit (Promega Corporation, Madison, WI, USA), which was adapted to measure GTP from cell lysates (see Fig. 7A). In brief, 60 µL of TCA-precipitated lysate was diluted in 60 µL of GTPase/GAP buffer. A 10 µM stock solution of rGTP, provided by the manufacturer, was used as a positive control for the assay. Each sample was then aliquoted in quadruplicate 25 µL volumes in separate wells of a white polystyrene 96-well plate. Two wells for each sample received GTPase-Glo buffer containing ADP and GTPase-Glo reagent ([ATP]+[GTP]), while the other two wells received GTPase-Glo buffer alone ([ATP]). Samples were then incubated for 30 min at RT° C with shaking. Following incubation, 50 µL of Detection reagent was added to each well and allowed to incubate for another 10 min at RT° C with shaking. Luminescence was then measured on a Tecan Spark® microplate reader (Tecan Group Ltd., Männedorf, Switzerland). GTP levels were calculated by subtracting the baseline [ATP] luminescence reading from the converted [GTP]+[ATP] luminescence reading. All values were normalized to the mean of the untreated, mock-infected control group.

Statistics

All statistical computations were performed in RStudio (version 1.3.1093) using base platform functions and the code is available as indicated above. All plots were made in the ggplot2 base package (version 3.1.0) (Wickham, 2009) and the ggpubr package (version 0.2.3; <https://CRAN.R-project.org/package=ggpubr>) or in Adobe Illustrator (version 24.1.2). All tests are indicated in the figure legends along with the value of N (independent biological replicates). All plots represent the mean and standard deviation of the data. Significance was defined as a *p*-value below 0.05 and a sample size of three was considered satisfactory for estimating normality.

Supplemental Information

Figure 1 - Supplementary Text

The validation of RNA-sequencing data for chlamydial transcriptomes is not straight-forward (Ardisson et al., 2020; Brinkworth et al., 2018), primarily owing to the conventional normalization method utilized for targeted RT-qPCR gene expression data. Two general means of normalizing RT-qPCR data have been proposed: (1) normalization of gene expression to genome equivalents and (2) normalization of gene expression to the expression of an internal control gene. In the case of the former method, gene expression is interpreted on a per-organism basis, reflecting changes in absolute transcript levels. In the case of the latter method, gene expression is theoretically represented as a proportion of the total transcriptome, but the common practice of normalizing to a single control gene carries many assumptions that are often faulty, such as the turnover rate of the control gene under various conditions, leading to erroneous results (Engström et al., 2010). Yet, the nature of genome normalization can produce unclear results depending on the experimental question being asked. During chlamydial persistence, where genome copies are reduced and basal transcriptional activity increases (Ouellette et al., 2006), genome normalization can over-estimate the up-regulation of genes whose abundance as a proportion of the total transcriptome does not change.

We therefore considered an alternative means of RT-qPCR normalization: geometric averaging of multiple control genes by the geNorm method (Vandesompele et al., 2002). This approach relies on the identification of stably expressed groups of control genes, empirically determined by an assessment of their stability (*i.e.* the maintenance of the ratio of their raw expression values) and their pairwise variation (*i.e.* the variation in stability between any two control genes across conditions). The analysis therefore provides a normalization factor that is based on the *stable relationship* of the expression of multiple control genes, rendering it more insensitive to instability or fluctuations in the expression of a single control gene. We used this method to analyze the following seven transcripts: *euo*, *omcB*, *groEL_1*, *ompA*, *nrdA*, *nrdB*, and 16S rRNA. We determined that across our experimental conditions, *euo* and *groEL_1* were the most stably associated genes under the tested conditions (Figure 1 – Figure Supplement 1, A and B) and that the set of *euo*, *groEL_1*, *nrdA* and *nrdB* had the lowest average pairwise variation (Figure 1 – Figure Supplement 1C) and were most suitable for the derivation of a normalization factor. We note that while the suggested cut-off for pairwise variation of a set of control genes is 0.15, we find that all genes analyzed here appear highly stable (with pairwise variation not exceeding 0.027), likely reflecting the strong effect of developmental regulation on the relationship of chlamydial gene expression. This normalization facilitated the confirmation of RNA-sequencing data by RT-qPCR by increasing the sensitivity for down-regulated or unchanged gene expression. This normalization also more accurately reflects the normalization methods utilized during RNA-sequencing, *i.e.* transcriptome-based normalization factors across conditions. We suggest that future gene expression studies in *Chlamydia* carefully consider the most suitable normalization method for the experimental question at hand.

Figure 1 – Figure Supplement 1. Analysis of stable control genes by geNorm. **(A)** Table displaying the step-wise exclusion of genes with the lowest stability (highest M value) following geNorm analysis. **(B)** Plot of average gene stability (M) during the stepwise exclusion of unstable genes. **(C)** Plot of pairwise variation of remaining control genes during the stepwise exclusion of unstable genes.

Figure 3 – Figure Supplement 1. Validation of differential expression for selected accessory genes during nutritional stress. All plots represent the mean and standard deviation of three independent biological replicates (N = 3). Statistical significance in all panels was determined by one-way ANOVA followed by Tukey's post-hoc test of honestly significant differences (two-tailed). * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, ns = not significant.

Figure 6 – Figure Supplement 1. Analysis of accessory gene expression in MIZ24. Statistical significance in all panels was determined by pairwise two-sided unpaired Welch's *t*-test for unequal variance. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$. All plots represent the mean and standard deviation of three independent biological replicates (N = 3).

Figure 7 – Figure Supplement 1. Total luminescent output in unconverted ([ATP]) and converted ([GTP]+[ATP]) reactions for the measurement of intracellular GTP levels. Note that the values in ([GTP]+[ATP]) samples are higher than those in [ATP] alone, reflecting the conversion of the GTP pool. All plots represent the mean and standard deviation of three independent biological replicates (N = 3). Statistical significance in all panels was determined by one-way ANOVA followed by Tukey's post-hoc test of honestly significant differences (two-tailed). * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, ns = not significant.