Title

The OceanDNA MAG catalog contains over 50,000 prokaryotic genomes originated from various marine environments

Authors

Yosuke Nishimura¹, Susumu Yoshizawa^{1, 2, 3}

Affiliations

¹Atmosphere and Ocean Research Institute, The University of Tokyo, Chiba 277-8564, Japan.

²Graduate School of Frontier Sciences, The University of Tokyo, Chiba 277-8563, Japan.

³Collaborative Research Institute for Innovative Microbiology, The University of Tokyo, Tokyo 113-8657, Japan.

corresponding author: Yosuke Nishimura (ynishimura@aori.u-tokyo.ac.jp)

Abstract

Marine microorganisms are immensely diverse and play fundamental roles in global geochemical cycling. Recent metagenome-assembled genome studies, with special attention to large-scale projects such as *Tara* Oceans, have expanded the genomic repertoire of marine microorganisms. However, published marine metagenome data has not been fully explored yet. Here, we collected 2,057 marine metagenomes (>29 Tera bps of sequences) covering various marine environments and developed a new genome reconstruction pipeline. We reconstructed 52,325 qualified genomes composed of 8,466 prokaryotic species-level clusters spanning 59 phyla, including genomes from deep-sea deeper than 1,000 m (n=3,337), low-oxygen zones of <90 $\mu\text{mol O}_2$ per kg water (n=7,884), and polar regions (n=7,752). Novelty evaluation using a genome taxonomy database shows that 6,256 species (73.9%) are novel and include genomes of high taxonomic novelty such as new class candidates. These genomes collectively expanded the known phylogenetic diversity of marine prokaryotes by 34.2% and the species representatives cover 26.5 - 42.0% of prokaryote-enriched metagenomes. This genome resource, thoroughly leveraging accumulated metagenomic data, illuminates uncharacterized marine microbial ‘dark matter’ lineages.

Background & Summary

Marine microorganisms have shaped Earth's environment and played crucial roles in controlling the global climate^{1,2}. Genome-based knowledge is essential to understand microorganisms in various aspects, such as their phylogeny, evolution, metabolism, and physiology. Though difficulty in isolation has limited the genome-based knowledge of marine microorganisms, success of culture independent genome reconstruction techniques such as metagenome-assembled genomes (MAGs) and single-amplified genomes (SAGs) have changed our understanding of microbial ecosystems. Genome information of marine microorganisms supplied by these approaches enabled to uncover new lineages that have been identified as participants in important biogeochemical cycling (e.g., nitrogen fixation³ and carbon fixation^{4,5}), characterize metabolic potentials of uncultured lineages^{6,7,8,9,10}, and reconstruct deep evolutionary trajectories^{11,12}.

Metagenomes of *Tara* Oceans Expeditions^{13,14} have been repeatedly subjected for genome reconstruction^{3,4,10,11,15,16,17}. In contrast, there are many metagenomes from which relatively little effort has been made for genome reconstruction despite large-scale data (e.g., GEOTRACES¹⁸) or from which reported genomes were limited to ones of specific taxa (e.g., metagenomes of the Canada Basin¹⁹). Moreover, genome reconstruction methodologies in many previous studies are considered inefficient (e.g., use of a single binning algorithm and/or coverage profile calculated by a single or only limited samples²⁰). Genome reconstruction using an improved methodology and applying it to a large-scale metagenome dataset is thus promising for expanding our genomic knowledge of marine microorganisms.

We aimed to build an extended genome catalog of marine prokaryotes with taking advantage of accumulated metagenomic data. Practically, two methodological focuses of this study were defined as (1) to compose a large-scale metagenome dataset that covers diverse marine environments including less explored regions such as deep-sea, low-oxygen zones, and polar regions and (2) to develop a new genome reconstruction pipeline to maximize quality of reconstructed genomes. Here, we collected 2,057 published metagenomes originated from diverse marine environments (Fig. 1ab). Then, to improve the quality of genomes, we developed a genome reconstruction pipeline that includes three key processes (Fig. 1c). As a result, we reconstructed 52,325 qualified prokaryotic genomes that were QS (quality score: %-completeness - 5 x %-contamination) ≥ 50 , named as the OceanDNA MAGs. These genomes were reconstructed from various marine environments, including genomes originated from deep-sea deeper than 1,000 m (n=3,337; from 179 metagenomes), low-oxygen zones of <90 $\mu\text{mol O}_2$ per kg water (n=7,884; from 176 metagenomes), and polar regions (n=7,752; from 129 metagenomes) (Fig. 2a).

The OceanDNA MAGs were composed of 8,466 species-level clusters. Genomes were identified as species representatives if the genome quality is the best within each species-cluster

(assessed by ‘QS + ln(N50)’). The median genome completeness and contamination of the OceanDNA MAGs were estimated as >80% and <2%, respectively (Fig 2b). The species representatives were originated from various metagenomic projects (divisions), and not dominated by ones from *Tara* Oceans (Fig. 2c). Taxonomic classification based on the genome taxonomy database (GTDB) release 05-RS95²¹ showed that the OceanDNA MAGs covered various marine prokaryotic lineages spanning 59 phyla (Fig. 2d). As taxonomic novelty assessment according to GTDB, 11 species representatives were not assigned to any existing class, suggesting that these species potentially belong to new classes. Likewise, 44 species representatives were suggested to belong to new orders, 290 were to new families, and 1,395 were to new genera (Fig. 2e). Overall, A large part of representatives (n=6,256; 73.9%) was not assigned to existing species in the database.

Novelty of the OceanDNA MAGs were further evaluated by a collection of published marine prokaryotic genomes (n=29,292; QS ≥ 50). Among the 8,466 species representatives, a large part (80.1%) was not overlapped with the published genomes at species level (56.8%) or was overlapped but of superior genome quality (here assessed by ‘QS + ln(N50)’ to the published genomes (23.3%) (Fig. 2f). The OceanDNA MAGs expanded the known phylogenetic diversity of marine prokaryotes by 34.2% (34.8% for bacteria and 29.4% for archaea) that was evaluated by the sum of branch length of bacterial and archaeal phylogenomic trees (Fig. 2g). The species representative genomes collectively covered 26.5 - 42.0% of metagenomic reads of prokaryote-enriched metagenomes at ≥95% nucleotide identity (Fig 3a). The OceanDNA MAG catalog is available as an unprecedented-scale genome resource of marine prokaryotes that enable to characterize microbial ‘dark matter’ lineages and to elucidate yet unsolved questions of marine microbial ecosystems.

Methods

Collection of metagenomes. We composed a dataset of marine metagenomic samples originated from a broad range of geographic regions (Fig 1ab). These metagenomes were reported by various research groups, and we organized these into 24 divisions for operational purpose, considering various factors such as related publications, research groups, and geographic regions (Table 1). These metagenomes include ones originated from long-distance cruises (e.g., *Tara* Oceans^{22,23,24}, GEOTRACES¹⁸, and Malaspina²⁵) and from time-series and/or transect sampling in a specific marine region (e.g., the Mediterranean Sea^{26,27}, the Baltic Sea²⁸, the Saanich Inlet²⁹, Station ALOHA³⁰, and the San Pedro Channel³¹). Associated metadata such as location, date, depth, oxygen concentration was collected from original publication and the BioSample database (Supplementary File S1). The metagenomic samples were originated from pole-to-pole (76.96°S - 85.02°N), sea surface to deep-sea (0 - 10,899 m below sea level), oxic to anoxic zones, coastal to pelagic seas (Fig. 1ab). The samples contain ones from aphotic zones

(179 metagenomes from deeper than 1,000 m; 200 metagenomes from 200 - 1,000 m) and low-oxygen zones (73 dysoxic (20 - 90 $\mu\text{mol/kg}$), 86 suboxic (1 - 20 $\mu\text{mol/kg}$), and 17 anoxic (<1 $\mu\text{mol/kg}$) metagenomes³²; Fig 1b). Most samples were originated from prokaryote-enriched fractions (here defined as sea water pass through a prefilter of 0.45 - 5 μm pore and collected on a filter of 0.1 - 0.45 μm pore; $n=732$), prokaryote- and eukaryote-enriched fractions (pass through a prefilter of 20 μm pore or no prefilter and collected on a filter of 0.2 - 0.8 μm pore; $n=832$), and virus-enriched fractions (pass through a prefilter of 0.2 - 0.22 μm pore; $n=312$; Fig 1b). In addition to water samples, metagenomes originated from sediment traps^{33, 34} ($n=63$) and in situ formation of biofilms³⁵ ($n=104$) were collected. Overall, these metagenomes cover various marine environments.

Sequence assemblies and metagenome binning. Metagenomic sequence data in a paired-end layout was downloaded from NCBI SRA and quality controlled by using Trimmomatic³⁶ v0.35, with an option ‘LEADING:20 TRAILING:20 MINLEN:60’. If one side of the pair was discarded due to its low quality, the other was retained when it passed the quality control. The qualified reads were assembled in a sample-by-sample manner (i.e., all qualified reads from one sample were used in one assembly; statistics are described in Supplementary File S1) using MEGAHIT³⁷ v1.1.4. Resulting contigs were retained if the length is no less than 1 kbps.

We then calculated a coverage profile for each metagenome using all metagenomes belong to the same division for better binning performance (Table 1; see also ‘Technical Validation’). An exception was applied to the division of GEOTRACES, which includes many metagenomes ($n=610$). This division is split into six subdivisions and the coverage profiles were calculated within each subdivision (Supplementary File S1). Read mapping was performed by bowtie2³⁸ v2.3.5.1 using qualified paired-end reads. Mapping result was sorted by samtools (http://www.htslib.org/) v1.9 and coverage was calculated by jgi_summarize_bam_contig_depths that is bundled in MetaBAT2³⁹, customizing a parameter ‘-percentIdentity’ set to 90. We then performed metagenome binning using three algorithms, MetaBAT2³⁹ v2.12.1, MaxBin2⁴⁰ v2.2.6, and CONCOCT⁴¹ v1.0.0. These algorithms were run with default settings, but for MetaBAT2, the ‘--minContig’ parameter was set to 1,500 following the software instruction, which states this value should not be less than 1,500. The resulting three sets of bins were then dereplicated and merged using the bin_refinement module of MetaWRAP⁴² v1.2.1 with minimum completion is set to 50. The quality score (QS) was defined as ‘%-completeness - 5 x %-contamination’ and genomes of $\text{QS} \geq 50$ were retained. Completeness and contamination of genome bins was estimated by taxon specific sets of single-copy marker genes through the lineage-specific workflow of CheckM v1.0.13⁴³. After removal of genomes likely derived from internal standard ($n=63$; *Thermus thermophilus* and *Blautia producta*⁴⁴), 54,614 genomes were obtained.

Post-refinement of genome bins. For quality improvement of the reconstructed genome bins, we developed a post-refinement module to decontaminate potential misassigned contigs for each genome bin (Fig 1c; see also ‘Technical Validation’). This module consists of three independent decontamination filters: (1) taxonomic filter, (2) mobile element filter, and (3) outlier filter. First, the taxonomic filter was designed to detect taxonomically inconsistent contigs with each genome. Coding regions were predicted with prodigal⁴⁵ v2.6.3 and resulting proteins were used as input of CAT and BAT⁴⁶ v5.0.3 to assign taxonomy for contigs and genomes, respectively. CAT and BAT was run with the default setting using NCBI Taxonomy downloaded in January 2020. Then, predicted taxonomy was quality controlled to remove less reliable assignment. Namely, predicted taxonomy was recursively trimmed from the low level until either of the following three types of assignment are not detected: (a) ‘suggestive’ taxonomic assignment that is less confident, indicated by stars in the BAT and CAT output, (b) very low-level assignment equal to or lower than species-level, and (c) some ambiguous assignment (i.e., classified as ‘environmental samples’ or classifications start with ‘unclassified’). For each pair of a genome and its contig, the pair was recognized as taxonomically consistent only if the lowest common ancestor of the genome and the contig was the same as either of these. For example, suppose taxonomy of a genome is ‘A; B; C’, a contig is taxonomically consistent if taxonomy of a contig is ‘A; B’ or ‘A; B; C; D’, and inconsistent if ‘A; B; E’ or ‘A; F’.

Second, the mobile element filter was designed to remove possible contamination of viral and plasmid contigs within genome bins. As genome bins are likely contaminated with viral and plasmid contigs that have similar coverage and nucleotide composition to the genome²⁰, we adopted a conservative approach to remove possible mobile elements, though these contigs are possibly true parts of the genome (e.g., as a provirus). First, circular contigs were identified as potential viral and plasmid contigs by detecting terminal redundancy through ccfind (<https://github.com/yosuken/ccfind>)⁴⁷. Second, viral contigs were detected using additional two types of methods. VirSorter⁴⁸ v1.0.6 was used to detect viral contigs of ≥ 3 kb. The prediction result of category 1-6 was considered as viral, but for category 4-6 (predicted as provirus), only if length of viral region was $\geq 50\%$ of the total length, the contig was considered as viral. To supplement the detective power for short contigs (1kb to 10kb), we additionally scanned for *terL* genes that are one of the hallmark genes of prokaryotic viruses, by following steps. We prepared 11 *terL* HMMs (Supplementary File S2) that were constructed from *terL* protein sequences obtained from previously identified aquatic viral MAGs (EVGs: circularly assembled environmental viral genomes)⁴⁷. We searched for *terL* candidates using hmmsearch (HMMER⁴⁹ v3.2.1; $\text{evalue} < 1\text{e-}10$) with the 11 HMMs as query. We validated sequence homology of the candidates with known *terL* genes using

pipeline_for_high_sensitive_domain_search
(https://github.com/yosuken/pipeline_for_high_sensitive_domain_search), which utilizes jackhmmer (HMMER⁴⁹ v3.2.1) to build a protein HMM of each gene and hhsearch⁵⁰ (HH-suite⁵⁰ v3.2.0) to identify homology between the built HMMs and *terL* HMMs included in pfam 32.0. The candidates were identified as *terL* if the best hit is one of *terL* domains (i.e., Terminase_1, Terminase_3, Terminase_6, Terminase_GpA, DNA_pack_N, Terminase_3C, and Terminase_6C) among all the pfam domain and if probability of the HHsearch hit is >97%. We used proteins encoded in EVGs as a database of jackhmmer (jackhmmer parameters: ‘-N 5 --incE 0.001 --incdomE 0.001’).

Third, the outlier filter was designed to detect outlier contigs in terms of coverage and tetranucleotide frequency (<-2.5 or >2.5 s.d. within each genome bin). Principal component analysis was performed using the prcomp function of R v3.6.2 (with default parameters) and the first primary component was evaluated. As a coverage profile, a part (related to contigs of the bin) of a coverage profile that was used for binning was extracted and normalized within each sample. Contigs identified as outliers were removed from the genome bin. Overall, after the detection and removal of possible contamination using these three filters, completeness and contamination of each genome bin was again estimated with the lineage-specific workflow of CheckM.

Finally, 52,325 genomes of QS ≥ 50 were obtained and here named as the OceanDNA MAGs (Data Citation 1; Table S2). The OceanDNA MAGs reconstructed from various marine environments and size-fractions (Fig 2a), including deep-sea deeper than 1,000m (3,337 genomes from 176 samples), low-oxygen zones of <90 $\mu\text{mol O}_2$ per kg water (7,884 genomes from 176 samples), polar regions (7,752 genomes from 129 samples), viral enriched fraction (pass through a filter of 0.2 or 0.22 μm pore; 5,998 genomes from 312 samples). Basic statistics of genome assemblies were evaluated with QUAST⁵¹ v5.0.2 (Supplementary File S3). Ribosomal RNAs and transfer RNAs were identified using Barrnap v0.9 (<https://github.com/tseemann/Barrnap>) and tRNAscan-SE⁵² v2.0.5, respectively.

Taxonomic assignment and their novelty evaluation using GTDB. We performed species-level clustering and identified species representatives of the OceanDNA MAGs through the following two rounds. First, for each of the 24 divisions, species-level clustering was performed using dRep⁵³ v2.2.2 with a cutoff value of average nucleotide identity set as 95% and aligned fraction as 30%. We identified genomes of species representatives if ‘QS + ln(N50)’ was the highest within each species-level cluster. From the 24 divisions, 13,357 species representatives were identified at this round. Then, the secondary clustering was performed among these representatives using dRep, and 8,466 species-level clusters were obtained. The representatives of the 8,466 species-level clusters (Data Citation 2) were identified using the same criteria. The

median genome completeness and contamination of both the species representatives and the other genomes (n=43,859; Data Citation 3) were estimated as >80% and <2%, respectively (Fig 2b). The species representatives were originated from various metagenomic projects and not dominated by ones from *Tara* Oceans (Fig. 2c).

The OceanDNA MAGs were taxonomically classified using GTDB (Genome Taxonomy DataBase) release 05-RS95²¹ through the classify workflow of GTDB-Tk⁵⁴ v1.3.0. As classification based on GTDB, the species representatives spanned 59 phyla (Fig. 2d). Of these, 11 species representatives were not assigned to any existing class, suggesting that these species potentially belong to new classes. Likewise, it was suggested that 44 species representatives belong to new orders, 290 belong to new families, and 1,395 belong to new genera and 4,516 belong to new species (Fig. 2e). Overall, most of the species representatives (n=6,256; 73.9%) were not assigned to existing species in the database.

Novelty evaluation using published marine genomes. For further novelty assessment of the OceanDNA MAGs, we comprehensively collected published genomes of marine prokaryotes. First, genomes contained in MarDB and MarRef⁵⁵ v5.0, which are curated genome collection of marine prokaryotes originated from isolates/SAGs/MAGs, were downloaded (n=14,209). Second, to supplement these with very recently published genomes and/or genomes that are not stored in NCBI, we collected genomes (n=26,946; SAGs and MAGs) of marine origin from 15 research articles^{3,5,6,10,24,33,35,56,57,58,59,60,61,62,63} (Supplementary File S4). After selection of qualified genomes (QS $\geq$ 50), 29,292 genomes were retained in total (11,985 from marRef/MarDB and 17,307 genomes from the 15 articles; Supplementary File S5). We then organized a unified genome catalog of marine prokaryotes (UGCMP; n=81,617), composed of the 29,292 published genomes and the 52,325 OceanDNA MAGs (Fig. 2f). We identified species representatives of UGCMP by following two steps. Species-level clusters (n=13,669) and the representatives were identified separately for MarDB/MarRef and for each publication, using the same criteria as the OceanDNA MAGs. After unifying the species representatives of OceanDNA MAGs (n=8,466) and published marine genomes (n=13,669) into one set, the second-round species-level clustering was performed with the same conditions. We finally identified 16,141 species representatives of UGCMP using the same criteria (Supplementary File S6). The OceanDNA MAGs exclusively composed 4,806 species-level clusters (56.8% of the species representatives of the OceanDNA MAGs) and selected as species representatives in 1,971 non-exclusive species-level clusters (23.3% of the species representatives of OceanDNA MAGs) based on the better genome quality evaluated by 'QS + ln(N50)'. Overall, a large part (80.1%; n=6,777) of the species representatives of the OceanDNA MAGs was still species representatives in UGCMP.

We then assessed phylogenomic diversity of UGCMP for bacteria (n=74,214) and archaea (n=7,403). For domain and phylum-level classification, taxonomic assignment of UGCMP genomes were performed using GTDB release 05-RS95 and GTDB-Tk v1.3. Phylogenomic trees of bacteria and archaea were reconstructed with FastTree v2.1.11 (option: '-wag -gamma') using alignments that were built by GTDB-Tk (Fig 2g). The alignments included 5,040 sites of high phylogenetic signal from 120 single copy marker genes for bacteria, and 5,124 sites from 122 genes for archaea as well. After midpoint rooting using gotree (<https://github.com/evolbioinfo/gotree>) v0.4.0, sum of branch length was calculated for two categories: (1) branches that were represented only by the OceanDNA MAGs (2) branches that were other than (1). The expanded phylogenetic diversity by the OceanDNA MAGs was 34.2% (34.8% for bacteria and 29.4% for archaea), estimated from a ratio of (1) to (2).

Back mapping of metagenomic reads. We assessed the fraction of metagenomic reads recruited onto the OceanDNA MAGs. Sequence reads of the 2,057 metagenomes, which were used for genome reconstruction, were back mapped onto the 8,466 species representatives of the OceanDNA MAGs. For cases that one sample has multiple sequencing runs, only the biggest run was used. Read mapping was performed with bowtie2³⁸ v2.3.5.1 with the default setting using the quality controlled paired-end reads of each run, but if the run was bigger than 5 Gbps, a subset of 5 Gbps sequences that was randomly sampled using seqtk (<https://github.com/lh3/seqtk>) v1.3 was used for read mapping. Then, the mapping result was sorted using samtools (<http://www.htslib.org/>) v1.9, and only mapping of $\geq 95\%$ identity, ≥ 80 bp, and $\geq 80\%$ aligned fraction of the read length was extracted using msamtools (<https://github.com/arumugamlab/msamtools>) that are bundled in MOCAT2⁶⁴ v2.1.3. Finally, the mapped reads were counted using featureCounts⁶⁵ that were bundled in Subread v2.0.0. The species representatives collectively cover 10.4 - 35.0% (the 25th to 75th percentile) of metagenome reads of the 2,057 metagenomes (Fig 3a). Especially, where only prokaryotes-enriched metagenomes (n=731) were considered, 26.5 - 42.0% of metagenomic reads were mapped onto the species representatives.

Next, we evaluated mapped read fractions onto species representatives of UGCMP, the OceanDNA MAGs, and four sets of marine prokaryotic genomes from large-scale genome reconstruction studies^{3,5,16, 62} (Fig 3b). Read mapping was performed using only species representatives of qualified genomes (i.e., QS ≥ 50) for all these genome collections. In terms of the medians of mapped read fractions, the OceanDNA MAGs was the highest among the previously reported genome collections, and UGCMP was about 10% higher than the OceanDNA MAGs.

Data Records

Genome sequences of the OceanDNA MAGs (Data Citation 1) were available at figshare (<https://figshare.com/s/e2aa3456d68aa51e617c>) and submitted to DDBJ/ENA/GenBank under BioProject accession no. PRJDB11811. Genome sequences of the 8,466 species representatives (Data Citation 2) were submitted as WGS entries, and sequences of the other genomes (n=43,859) were submitted as DDBJ analysis entries (Data Citation 3; available via DDBJ). Supplementary files (listed below) are available at figshare (<https://figshare.com/s/e2aa3456d68aa51e617c>). Related information of the OceanDNA MAGs can be accessed at <https://OceanDNA-MAGs.aori.u-tokyo.ac.jp>.

Supplementary File S1. A list of metagenomes used in this study, with various information used for generating figures.

Supplementary File S2. Eleven multiple alignments and HMMs of *terL* protein sequences obtained from aquatic viral MAGs.

Supplementary File S3. A list of the OceanDNA MAGs with basic statistics, functional RNAs, genome quality, and genome-based taxonomy.

Supplementary File S4. A list of 15 publications of marine SAGs and MAGs

Supplementary File S5. A custom collection of published marine prokaryotic genomes of QS ≥ 50

Supplementary File S6. A list of species representatives of UGCMP

Technical Validation

For maximization of the genome quality, our genome reconstruction pipeline was carefully designed, including three key processes (Fig. 1c): (1) high-resolution coverage profiles were calculated using all metagenomes within each division, (2) metagenome binning was performed using three algorithms and subsequently dereplicated, (3) an automated post-refinement process for detection of possible contaminations, including ones could be missed by prokaryotic single-copy marker gene-based assessment. Here we assessed efficiency of these processes.

First, binning algorithms are primarily based on a coverage profile among multiple metagenomes and *k*-mer (e.g., tetranucleotide) composition of metagenomic contigs^{66,67}. It was shown that if a coverage profile was calculated using only a few metagenomes, it would result in underperformance of a binning algorithm (e.g., CONCOCT)⁴¹. Here, to assess the effect of the number of metagenomes in coverage profile, we selected 20 *Tara* Oceans metagenomes included in the “*Tara* prok” division (Table 1) of which geographic region and water depth was widely distributed. We performed metagenome binning of the selected metagenomes with different coverage profiles. The coverage profiles were calculated with all metagenomes within the same division (n=139) or with randomly sampled 10, 25, and 50 metagenomes with three

replicates out of the 139 metagenomes. If multiple sequencing runs were available from one metagenome, only the largest run was used for coverage profiles. Then, binning was performed same as the OceanDNA MAGs but except for the post-refinement part, and the resulting number of bins of $QS \geq 50$ was compared (Fig 4a). As a result, coverage profiles of all metagenomes reconstructed the greater number of qualified bins (i.e., $QS \geq 50$) than coverage profiles of subsampled metagenomes. The result suggests the superiority of the ‘high-resolution’ coverage profiles calculated with many metagenomes.

Second, using the same 20 metagenomes of the “*Tara prok*” division, binning result of single algorithm (MetaBAT2, CONCOCT, MaxBin2) and dereplicated result of the three algorithms using the `bin_refinement` module of MetaWRAP were compared (Fig 4b). Dereplication of bins generated from three algorithms significantly increased the number of qualified (i.e., $QS \geq 50$) bins.

Third, we designed an automated post-refinement process to remove possible contamination from each MAG using three filters that are independent of prokaryotic single-copy marker genes: (1) taxonomic filter, (2) mobile element filter, and (3) outlier filter. Similar strategies were applied in previous studies (e.g., MAGpurify⁶⁸). The aim of this refinement process is to remove possible contamination for genome quality improvement. Especially, contamination over the domain (i.e., eukaryotic and viral contigs included in prokaryotic genomes) could not be detected through analysis of prokaryotic single-copy marker genes. Genomes from *Tara* Oceans MAG studies were predicted to contain viral contigs (in a few cases, more than 50) within a single genome⁶⁹. Viral contigs could be contamination of viral genome fragments that have similar coverage profiles and k-mer compositions to the prokaryotic genome²⁰. Though removing viral and plasmid sequences possibly results in the exclusion of true element of the genome (e.g., provirus) and identification of viral and plasmid contigs could contain false positives, we set a priority on removing those as possible contamination, not retaining those as true genomic fragments. The three filters of the post-refinement module identified 561,804, 39,289, and 436,143 potential misassigned contigs, respectively. Overall, from 54,614 qualified genome bins, 1,000,417 contigs were filtered out (18.3 contigs per genome bin on average) and 2,289 genome bins were discarded, due to the reduction of genome completeness as a result of the decontamination process. Code for the post-refinement process is available at GitHub (<https://github.com/yosuken/MAGRE>).

Usage Notes

We carefully designed the genome reconstruction pipeline for genome quality improvement, including the automated post-refinement process. Nevertheless, due to difficulty of perfect decontamination, misassigned contigs might be still included in the genomes. Manual quality control is recommended before use of the genomes, as is the case for MAGs from other studies.

We collected metagenome samples covering various marine environments. Nevertheless, note that some marine environments (e.g., hydrothermal vents, sediments, coral reefs, and oil spills) were not included in the dataset of this study.

Genome completeness evaluated by CheckM are likely underestimated for genomes of specific taxa that have experienced extreme genome reduction and may have a symbiotic lifestyle (e.g., lineages of the phylum Patescibacteria, also known as Candidate Phyla Radiation). Ribosomal RNA operons are difficult regions to reconstruct due to co-existence of closely related sequences that confuse de Bruijn graph-based assemblers²⁰. 5S, 16S, 23S ribosomal RNAs were identified in 24.2%, 6.8%, 3.8% of the OceanDNA MAGs, respectively (including full sequences or >25% fragments of the whole length).

SAR11 and Prochlorococcus are two of the most abundant lineages in the ocean. However, despite their high abundance, not so many genomes of these lineages were reconstructed in this study. This shortfall is probably attributable to coexisting closely related strains of these lineages that cause difficulty for genome reconstruction²⁰. Among the OceanDNA MAGs, 780 genomes were reconstructed from 85 species-level clusters of ‘o__Pelagibacterales’ (SAR11) and 157 genomes were reconstructed from 8 species-level clusters of ‘g__Prochlorococcus’. For these lineages, SAGs could supplement genomic information. For example, recently reported SAGs that were reconstructed from the tropical and subtropical euphotic ocean⁵ includes 2,108 genomes consisted of 1,215 species-level clusters of ‘o__Pelagibacterales’ and 327 genomes consisted of 155 species-level clusters of ‘g__Prochlorococcus’, where genomes are limited to those of $QS \geq 50$ (Supplementary File S5).

Code Availability

Code of the post-refinement module is available at GitHub as MAGRE (<https://github.com/yosuken/MAGRE>).

The options and parameters of all tools used for the analysis are described in the main text.

Acknowledgements

We thank all persons who contributed to generation of the metagenome sequence data, as well as all persons who developed the software and databases used in this study. This study is partly supported by JSPS KAKENHI Grant Number 18K19224, 18H04136, and 21K19134 (S.Y.). Computation time was provided by the Super Computer System, Institute for Chemical Research, Kyoto University.

Author contributions

Y.N. conceived the study, designed and implemented the pipeline, performed analysis, and wrote a draft. S.Y. reviewed and edited a draft.

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. Overview of the study. (a) Geographic distribution of the 2,057 metagenomes analysed in this study (shown by black points). The map was drawn using marmap⁷⁰ and ggplot2 (<https://ggplot2.tidyverse.org/>). (b) Origin of the metagenome samples. Types of the fraction were described in the main text. (c) Schematic representation of the developed pipeline for MAG reconstruction. Three key processes were highlighted by brown stars. Source data of (a) and (b) was provided in Supplementary File S1.

Figure 2. Origin, quality, and novelty of the OceanDNA MAGs. (a) Origin of the OceanDNA MAGs. Types of the fraction were described in the main text. (b) Genome completeness and contamination evaluated by CheckM. (c) Origin of metagenome divisions of the 8,466 species representatives. (d) Phyla of the species representatives assigned by GTDB-Tk. (e) Potential novelty of the species representatives assessed using GTDB-Tk. (f) Origins and compositions of the unified catalog UGCMP and the species representatives. (g) Bacterial (left) and archaeal (right) phylogenetic trees of the species representatives of UGCMP. The trees were midpoint rooted for visualization purpose. The number of species representatives and %-expanded phylogenetic diversity were described for individual phyla of which the number of species was at least 100 for bacteria and 10 for archaea. These phyla were highlighted in the trees with the corresponding colours. If a phylum was not monophyletic in the trees, only the largest monophyletic unit was highlighted (three phyla represented by asterisks in the legend). Note that %-expanded phylogenetic diversity was estimated using all the genomes of UGCMP (not limited to the species representatives).

Figure 3. Recruitment of metagenomic reads. The fraction of mapped reads of 2,057 metagenomes were evaluated at $\geq 95\%$ nucleotide identity. (a) Recruitment onto the species representatives of the OceanDNA MAGs. X-axis shows types of metagenome fractions. P: prokaryote-enriched metagenomes, P + E: prokaryote- and eukaryote-enriched metagenomes, V: virus enriched metagenomes. (b) Recruitment of prokaryote-enriched metagenome reads. X-axis shows genome collections. Note that all these genome collections include only species

representatives of qualified genomes (i.e., $QS \geq 50$). UGCMP and OceanDNA MAGs include genomes reconstructed in this study. Nayfach+, 2021⁶², Pachiadaki+, 2019⁵, Tully+, 2018¹⁶, and Delmont+, 2018³ are reported genome collections. For Nayfach+, 2021, genomes are limited to ones of which ‘ecosystem type’ is marine.

Figure 4. Assessment of the genome reconstruction pipeline. Using selected 20 *Tara* Oceans metagenomes included in the “*Tara* prok” division, the impact of high-resolution coverage profiles (a) and use of multiple binning algorithms (b) were assessed. The number of qualified genome bins ($QS \geq 50$) was compared between (a) coverage profiles calculated with all metagenomes within the same division (n=139) or with randomly sampled 10, 25, and 50 metagenomes with three replicates, and (b) different algorithms: MaxBin2, CONCOCT, MetaBAT2, and dereplicated results of the three algorithms using the bin_refinement module of MetaWRAP.

Tables

Table 1. metagenome divisions

division name	related publication (selected)	samples	QCed read (Gbp)	MAGs
<i>Tara</i> prok	Sunagawa et al., 2015 ²²	139	4,935	8,624
Saanich Inlet	Hawley et al., 2017 ²⁹	85	1,041	5,087
NS polar	Cao et al., 2020 ⁵⁸	59	847	3,511
<i>Tara</i> virus	Gregory et al., 2019 ²³	131	3,887	3,271
Monterey bloom	Nowinski et al., 2019 ⁴⁴	84	681	3,223
biofilm	Zhang et al., 2019 ³⁵	130	2,577	3,209
GEOTRACES	Biller et al., 2018 ¹⁸	610	4,998	3,063
North Sea	Kruger et al., 2019 ⁵⁶	38	832	3,019
<i>Tara</i> polar	Salazar et al., 2019 ²⁴	41	1,416	2,762
<i>Tara</i> girus	Sunagawa et al., 2015 ²²	59	1,612	2,757
Baltic Sea	Alneberg et al., 2018 ²⁸	81	566	2,335
Mediterranean	Lopez-Perez et al., 2017 ⁷¹ , Haro-Moreno et al., 2019 ⁷² , Martin-Cuadrado et al., 2015 ⁷³	37	599	2,292
HOT	Mende et al., 2017 ³⁰	85	1,000	2,109
Malaspina	Acinas et al., 2021 ²⁵ , Gregory et al., 2019 ²³	72	209	1,320
Med. coastal	Galand et al., 2018 ²⁷	40	276	1,243
Canada Basin	Colatriano et al., 2018 ¹⁹	12	362	1,083
Hawaii bloom	Wilson et al., 2017 ⁷⁴	88	530	641
San Pedro Channel	Sieradzki et al., 2019 ³¹ , Ignacio-Espinoza et al., 2019 ⁷⁵	65	1,527	554
sediment trap	Poff et al., 2021 ³⁴	63	470	506
low oxygen	Thrash et al., 2017 ⁶ , Tsementzi et al., 2016 ⁷⁶ , Glass et al., 2015 ⁷⁷	26	123	476
Atlantic	Bergauer et al., 2018 ⁷⁸	7	180	451
Red Sea	Haroon et al., 2016 ⁷⁹	45	83	319
NW Pacific	Saw et al., 2020 ¹⁰ , Li et al., 2018 ⁸⁰	35	96	248
Baltic Sea virus	Nilsson et al., 2019 ⁸¹	25	261	222

References

1. Falkowski, P. G., Fenchel, T. & DeLong, E. F. The microbial engines that drive Earth's biogeochemical cycles. *Science* **320**, 1034–1039 (2008).
2. Falkowski, P. Ocean Science: The power of plankton. *Nature* **483**, S17–20 (2012).
3. Delmont, T. O. *et al.* Nitrogen-fixing populations of Planctomycetes and Proteobacteria are abundant in surface ocean metagenomes. *Nat Microbiol* **3**, 804–813 (2018).
4. Graham, E. D., Heidelberg, J. F. & Tully, B. J. Potential for primary productivity in a globally-distributed bacterial phototroph. *ISME J* **12**, 1861–1866 (2018).
5. Pachiadaki, M. G. *et al.* Charting the Complexity of the Marine Microbiome through Single-Cell Genomics. *Cell* **179**, 1623–1635.e11 (2019).
6. Thrash, J. C. *et al.* Metabolic Roles of Uncultivated Bacterioplankton Lineages in the Northern Gulf of Mexico "Dead Zone". *MBio* **8**, e01017–17 (2017).
7. Haro-Moreno, J. M., Rodriguez-Valera, F., López-García, P., Moreira, D. & Martin-Cuadrado, A.-B. New insights into marine group III Euryarchaeota, from dark to light. *ISME J* **11**, 1102–1117 (2017).
8. Rinke, C. *et al.* A phylogenomic and ecological analysis of the globally abundant Marine Group II archaea (Ca. Poseidoniales ord. nov.). *ISME J* **13**, 663–675 (2019).
9. Tully, B. J. Metabolic diversity within the globally abundant Marine Group II Euryarchaea offers insight into ecological patterns. *Nat Commun* **10**, 271 (2019).
10. Saw, J. H. W. *et al.* Pangenomics Analysis Reveals Diversification of Enzyme Families and Niche Specialization in Globally Abundant SAR202 Bacteria. *MBio* **11**, 93 (2020).
11. Martijn, J., Vosseberg, J., Guy, L., Offre, P. & Ettema, T. J. G. Deep mitochondrial origin outside the sampled alphaproteobacteria. *Nature* **557**, 101–105 (2018).
12. Getz, E. W., Tithi, S. S., Zhang, L. & Aylward, F. O. Parallel Evolution of Genome Streamlining and Cellular Bioenergetics across the Marine Radiation of a Bacterial Phylum. *MBio* **9**, e01089–18 (2018).
13. Karsenti, E. *et al.* A holistic approach to marine eco-systems biology. *PLoS Biol* **9**, e1001177 (2011).
14. Sunagawa, S. *et al.* Tara Oceans: towards global ocean ecosystems biology. *Nat. Rev. Microbiol.* **18**, 428–445 (2020).
15. Tully, B. J., Sachdeva, R., Graham, E. D. & Heidelberg, J. F. 290 metagenome-assembled genomes from the Mediterranean Sea: a resource for marine microbiology. *PeerJ* **5**, e3558 (2017).
16. Tully, B. J., Graham, E. D. & Heidelberg, J. F. The reconstruction of 2,631 draft metagenome-assembled genomes from the global oceans. *Sci Data* **5**, 170203 (2018).
17. Parks, D. H. *et al.* Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nat Microbiol* **2**, 1533–1542 (2017).
18. Biller, S. J. *et al.* Marine microbial metagenomes sampled across space and time. *Sci Data* **5**, 180176 (2018).
19. Colatriano, D. *et al.* Genomic evidence for the degradation of terrestrial organic matter by pelagic Arctic Ocean Chloroflexi bacteria. *Commun Biol* **1**, 90 (2018).
20. Chen, L.-X., Anantharaman, K., Shaiber, A., Eren, A. M. & Banfield, J. F. Accurate and complete genomes from metagenomes. *Genome Res* **30**, 315–333 (2020).
21. Parks, D. H. *et al.* A complete domain-to-species taxonomy for Bacteria and Archaea. *Nat Biotechnol* **38**, 1079–1086 (2020).
22. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).
23. Gregory, A. C. *et al.* Marine DNA Viral Macro- and Microdiversity from Pole to Pole. *Cell* **177**, 1109–1123.e14 (2019).
24. Salazar, G. *et al.* Gene Expression Changes and Community Turnover Differentially Shape the Global Ocean Metatranscriptome. *Cell* **179**, 1068–1083.e21 (2019).
25. Acinas, S. G. *et al.* Deep ocean metagenomes provide insight into the metabolic architecture of bathypelagic microbial communities. *Commun Biol* **4**, 604 (2021).
26. Haro-Moreno, J. M. *et al.* Fine metagenomic profile of the Mediterranean stratified and mixed water columns revealed by assembly and recruitment. *Microbiome* **6**, 128 (2018).
27. Galand, P. E., Pereira, O., Hochart, C., Auguet, J.-C. & Debroas, D. A strong link between marine microbial community composition and function challenges the idea of functional redundancy. *ISME J* **12**, 2470–2478 (2018).
28. Alneberg, J. *et al.* BARM and BalticMicrobeDB, a reference metagenome and interface to

513 meta-omic data for the Baltic Sea. *Sci Data* **5**, 180146 (2018).

514 29. Hawley, A. K. *et al.* A compendium of multi-omic sequence information from the Saanich Inlet
515 water column. *Sci Data* **4**, 170160 (2017).

516 30. Mende, D. R. *et al.* Environmental drivers of a microbial genomic transition zone in the ocean's
517 interior. *Nat Microbiol* **2**, 1367–1373 (2017).

518 31. Sieradzki, E. T., Ignacio-Espinoza, J. C., Needham, D. M., Fichot, E. B. & Fuhrman, J. A.
519 Dynamic marine viral infections and major contribution to photosynthetic processes shown by
520 spatiotemporal picoplankton metatranscriptomes. *Nat Commun* **10**, 1169 (2019).

521 32. Wright, J. J., Konwar, K. M. & Hallam, S. J. Microbial ecology of expanding oxygen minimum
522 zones. *Nat. Rev. Microbiol.* **10**, 381–394 (2012).

523 33. Boeuf, D. *et al.* Biological composition and microbial dynamics of sinking particulate organic
524 matter at abyssal depths in the oligotrophic open ocean. *Proc Natl Acad Sci USA* **116**, 11824–
525 11832 (2019).

526 34. Poff, K. E., Leu, A. O., Eppley, J. M., Karl, D. M. & DeLong, E. F. Microbial dynamics of
527 elevated carbon flux in the open ocean's abyss. *Proc Natl Acad Sci USA* **118**, (2021).

528 35. Zhang, W. *et al.* Marine biofilms constitute a bank of hidden microbial diversity and functional
529 potential. *Nat Commun* **10**, 200–10 (2019).

530 36. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
531 data. *Bioinformatics* **30**, 2114–2120 (2014).

532 37. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node
533 solution for large and complex metagenomics assembly via succinct de Bruijn graph.
534 *Bioinformatics* **31**, 1674–1676 (2015).

535 38. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
536 357–359 (2012).

537 39. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome
538 reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

539 40. Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning algorithm to
540 recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605–607 (2016).

541 41. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat Methods*
542 **11**, 1144–1146 (2014).

543 42. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP-a flexible pipeline for genome-
544 resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).

545 43. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
546 assessing the quality of microbial genomes recovered from isolates, single cells, and
547 metagenomes. *Genome Res* **25**, 1043–1055 (2015).

548 44. Nowinski, B. *et al.* Microbial metagenomes and metatranscriptomes during a coastal
549 phytoplankton bloom. *Sci Data* **6**, 129 (2019).

550 45. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
551 identification. *BMC Bioinformatics* **11**, 119 (2010).

552 46. Meijnenfeldt, von, F. A. B., Arkhipova, K., Cambuy, D. D., Coutinho, F. H. & Dutilh, B. E.
553 Robust taxonomic classification of uncharted microbial sequences and bins with CAT and
554 BAT. *Genome Biol* **20**, 707–14 (2019).

555 47. Nishimura, Y. *et al.* Environmental Viral Genomes Shed New Light on Virus-Host Interactions
556 in the Ocean. *mSphere* **2**, e00359–16 (2017).

557 48. Roux, S., Enault, F., Hurwitz, B. L. & Sullivan, M. B. VirSorter: mining viral signal from
558 microbial genomic data. *PeerJ* **3**, e985 (2015).

559 49. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**, e1002195 (2011).

560 50. Söding, J. Protein homology detection by HMM-HMM comparison. *Bioinformatics* **21**, 951–
561 960 (2005).

562 51. Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D. & Gurevich, A. Versatile genome
563 assembly evaluation with QUAST-LG. *Bioinformatics* **34**, i142–i150 (2018).

564 52. Chan, P. P. & Lowe, T. M. tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences.
565 *Methods Mol Biol* **1962**, 1–14 (2019).

566 53. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate
567 genomic comparisons that enables improved genome recovery from metagenomes through de-
568 replication. *ISME J* **11**, 2864–2868 (2017).

569 54. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
570 genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2020).

571 55. Klemetsen, T. *et al.* The MAR databases: development and implementation of databases
572 specific for marine metagenomics. *Nucleic Acids Res* **46**, D692–D699 (2018).

573 56. Krüger, K. *et al.* In marine Bacteroidetes the bulk of glycan degradation during algae blooms is

- 574 mediated by few clades using a restricted set of genes. *ISME J* **13**, 2800–2816 (2019).
- 575 57. Thrash, J. C. *et al.* Metagenomic Assembly and Prokaryotic Metagenome-Assembled Genome
576 Sequences from the Northern Gulf of Mexico "Dead Zone". *Microbiol Resour Announc* **7**,
577 e01033–18 (2018).
- 578 58. Cao, S. *et al.* Structure and function of the Arctic and Antarctic marine microbiota as revealed
579 by metagenomics. *Microbiome* **8**, 47 (2020).
- 580 59. Sun, X. *et al.* Uncultured Nitrospina-like species are major nitrite oxidizing bacteria in oxygen
581 minimum zones. *ISME J* **13**, 2391–2402 (2019).
- 582 60. Aylward, F. O. & Santoro, A. E. Heterotrophic Thaumarchaea with Small Genomes Are
583 Widespread in the Dark Ocean. *mSystems* **5**, e00415–20 (2020).
- 584 61. Alneberg, J. *et al.* Ecosystem-wide metagenomic binning enables prediction of ecological
585 niches from genomes. *Commun Biol* **3**, 415–10 (2020).
- 586 62. Nayfach, S. *et al.* A genomic catalog of Earth's microbiomes. *Nat Biotechnol* **39**, 499–509
587 (2021).
- 588 63. Pachiadaki, M. G. *et al.* Major role of nitrite-oxidizing bacteria in dark ocean carbon fixation.
589 *Science* **358**, 1046–1051 (2017).
- 590 64. Kultima, J. R. *et al.* MOCAT2: a metagenomic assembly, annotation and profiling framework.
591 *Bioinformatics* **32**, 2520–2523 (2016).
- 592 65. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
593 assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).
- 594 66. Kang, D. D., Froula, J., Egan, R. & Wang, Z. MetaBAT, an efficient tool for accurately
595 reconstructing single genomes from complex microbial communities. *PeerJ* **3**, e1165 (2015).
- 596 67. Yue, Y. *et al.* Evaluating metagenomics tools for genome binning with real metagenomic
597 datasets and CAMI datasets. *BMC Bioinformatics* **21**, 334 (2020).
- 598 68. Nayfach, S., Shi, Z. J., Seshadri, R., Pollard, K. S. & Kyrpides, N. C. New insights from
599 uncultivated genomes of the global human gut microbiome. *Nature* **568**, 505–510 (2019).
- 600 69. Tominaga, K., Morimoto, D., Nishimura, Y., Ogata, H. & Yoshida, T. In silico Prediction of
601 Virus-Host Interactions for Marine Bacteroidetes With the Use of Metagenome-Assembled
602 Genomes. *Front Microbiol* **11**, 738 (2020).
- 603 70. Pante, E. & Simon-Bouhet, B. marmap: A package for importing, plotting and analyzing
604 bathymetric and topographic data in R. *PLoS ONE* **8**, e73051 (2013).
- 605 71. López-Pérez, M., Haro-Moreno, J. M., Gonzalez-Serrano, R., Parras-Moltó, M. & Rodriguez-
606 Valera, F. Genome diversity of marine phages recovered from Mediterranean metagenomes:
607 Size matters. *PLoS Genet* **13**, e1007018 (2017).
- 608 72. Haro-Moreno, J. M., Rodriguez-Valera, F. & López-Pérez, M. Prokaryotic Population
609 Dynamics and Viral Predation in a Marine Succession Experiment Using Metagenomics. *Front*
610 *Microbiol* **10**, 2926 (2019).
- 611 73. López-Pérez, M., Haro-Moreno, J. M., Coutinho, F. H., Martinez-Garcia, M. & Rodriguez-
612 Valera, F. The Evolutionary Success of the Marine Bacterium SAR11 Analyzed through a
613 Metagenomic Perspective. *mSystems* **5**, e00605–20 (2020).
- 614 74. Wilson, S. T. *et al.* Coordinated regulation of growth, activity and transcription in natural
615 populations of the unicellular nitrogen-fixing cyanobacterium Crocosphaera. *Nat Microbiol* **2**,
616 17118 (2017).
- 617 75. Ignacio-Espinoza, J. C., Ahlgren, N. A. & Fuhrman, J. A. Long-term stability and Red Queen-
618 like strain dynamics in marine viruses. *Nat Microbiol* **5**, 265–271 (2020).
- 619 76. Tsementzi, D. *et al.* SAR11 bacteria linked to ocean anoxia and nitrogen loss. *Nature* **536**, 179–
620 183 (2016).
- 621 77. Glass, J. B. *et al.* Meta-omic signatures of microbial metal and nitrogen cycling in marine
622 oxygen minimum zones. *Front Microbiol* **6**, 998 (2015).
- 623 78. Bergauer, K. *et al.* Organic matter processing by microbial communities throughout the
624 Atlantic water column as revealed by metaproteomics. *Proc Natl Acad Sci USA* **115**, E400–
625 E408 (2018).
- 626 79. Haroon, M. F., Thompson, L. R., Parks, D. H., Hugenholtz, P. & Stingl, U. A catalogue of 136
627 microbial draft genomes from Red Sea metagenomes. *Sci Data* **3**, 160050 (2016).
- 628 80. Li, Y. *et al.* Metagenomic Insights Into the Microbial Community and Nutrient Cycling in the
629 Western Subarctic Pacific Ocean. *Front Microbiol* **9**, 623 (2018).
- 630 81. Nilsson, E. *et al.* Genomic and Seasonal Variations among Aquatic Phages Infecting the Baltic
631 Sea Gammaproteobacterium Rheinheimera sp. Strain BAL341. *Appl. Environ. Microbiol.* **85**,
632 e01003–19 (2019).
- 633

a

bioRxiv preprint doi: <https://doi.org/10.1101/2021.08.18.456858>; this version posted August 18, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

2,057 metagenomes

b

c

a

bioRxiv preprint doi: <https://doi.org/10.1101/2021.08.18.456858>; this version posted August 18, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

b **c** **d** **e** **f** **g**

a **b**