NR4A3 is oxidative stress responsible transcription factor through HMOX1, and also controls cell cycle through CCNE1 and CDK2 in pancreatic islet derived 1.1B4 cells

Motoko Nakayama,¹ Etsuko Ueta,² Mitsuru Yoshida,¹ Yuri Shimizu,^{1,3} Reiko Oguchi,¹ Atsuko Tokuda,¹ Yasuko Sone,^{1,4} Yuri Nomi,^{1,5} and Yuzuru Otsuka^{1,*}

¹*Department of Life Science, Graduate School, Ochanomizu University, Bunkyo-ku, Tokyo, 112-8610, Japan,*

²*School of Health Science, Faculty of Medicine, Tottori University, Yonago, Tottori 683-8503, Japan,*

³*Department of Food Business, Nihon University, Fujisawa, Kanagawa 252-0880, Japan,*

⁴*Department of Health and Nutrition, Takasaki University of Health and Welfare, Takasaki, Gunma 370-0033, Japan*

⁵*Faculty of Applied Life Sciences, Niigata University of Pharmacy and Applied Life Sciences, Niigata 956-8603, Japan*

* To whom correspondence should be addressed. Tel: +81-466-21-8149; Fax: +81-466-21-8149; Email: otsuka.yuzuru@ocha.ac.jp

Running title "NR4A3 is oxidative transcription factor with HMOX1"

Key words: NR4A3, oxidative stress, transcription factor, microarray, HMOX1, cell cycle, CDK2

ABSTRACT

The mechanism of antioxidant defense system is still controversial. As islet β -cell is weak in oxidative condition, that causes diabetes mellitus, therefore, antioxidant defense system of human pancreatic islet derived 1.1B4 cell was analyzed. Cells were exposed to H_2O_2 and comprehensive gene expression was analyzed by Agilent human microarray. *HMOX1* and *NR4A3*, member of orphan receptor, were up-regulated. Therefore, *NR4A3* was knocked down with siRNA, then analyzed gene expression by microarray, and found that the knocked down cells were weak in oxidative stress. *HMOX1* expression was strongly inhibited by siRNA of *NR4A3*, and *NR4A3* responsible sequence of aaggtca was found near the *HMOX1* gene, suggesting *NR4A3* is oxidative stress responsible transcription factor through *HMOX1* expression. The expression of *CCNE1* and *CDK2* was also inhibited by knocked down of *NR4A3*, it is suggested *NR4A3* is also important transcription factor for cell growth regulation.

[Supplemental material is available for this article.]

INTRODUCTION

The relentless decline in β -cell function frequently observed in type 2 diabetic patients and loss of functional β -cell mass is a hallmark of type 1 and type 2 diabetes, and methods for restoring these cells are needed (Tessem et al. 2014). However, despite optimal drug management, loss of β -cell is not possible to stop, and has variously been attributed to glucose toxicity and lipo-toxicity. The former theory posits hyperglycemia, that elevated glucose concentrations increase levels of reactive oxygen species (ROS) in β -cell (Robertson 2004), which takes place within multiple mitochondrial and non-mitochondrial pathways. For example, high concentrations of glucose in vitro increase intracellular peroxide levels in islets, and decrease insulin expression by decreasing *PDX-1* and *MafA* expression (Robertson and Harmon 2006). In their paper, six

biochemical pathways along which glucose metabolism suggested to form ROS. One of pathway is glycation that is hyperglycemia produces glycation end product (AGE) that increases ROS, as AGE-BSA is known to increase production of ROS, and its apoptogenic effect was blocked by the antioxidant N-acetylcysteine (Weinberg et al. 2014). In physiologic concentrations, endogenous ROS help to maintain homeostasis, however, when ROS accumulate in excess for prolonged periods of time, they cause chronic oxidative stress and adverse effects. Thus, increased glucose produces ROS, and pancreatic β -cells are more sensitive to oxidative stress than other cell types because the expressions of antioxidant enzymes are low levels in pancreatic β -cell (Robertson 2004). Therefore, oxidative stress leads cellular damage to β -cell dysfunction. The effect of oxidative stress, however, on the β -cell is not understood well. Nitric oxide, peroxynitrite, hydrogen peroxide and other oxygen-reactive species might be involved in β -cell destruction during diabetes development (Mathis et al. 2001). Oxidative stress induces gene expression in heme oxygenase 1 (HMOX1), and the HMOX1 gene (*HMOX1*) is frequently activated under a variety of cellular stress conditions, by four pathways including heat-shock factor, nuclear factor-kappa B, nuclear factor E2-related factor 2 (NRF2), and activator protein-1 families, are arguably the most important regulators of the cellular stress response in vertebrates (Alam and Cook 2007). One of them, the Kelch-Like ECH-Associated Protein 1 (KEAP1)-NRF2 pathway in which pathway, NRF2 transcription factor directly bind to antioxidant responsive element (ARE) to induce antioxidant enzymes (Nguyen et al. 2009). However, another unknown pathway may be involved. For example, chronically excessive glucose and ROS levels can cause decreased insulin gene expression via loss of the transcription factors pancreatic and duodenal homeobox 1 (PDX1) and v-maf avian musculoaponeurotic fibrosarcoma oncogene homolog A (MAFA) and can also accelerate rates of apoptosis (Robertson 2004).

High glucose and oxidative stress induce cell damage not only to β -cell but also to other cells. There are many transcription factors and sensing proteins. Oxidants such as H_2O_2 can damage

proteins, regulate transcription factors, and sometimes induces cell apoptosis (Marinho et al. 2014). In his review, the regulatory mechanisms by which H₂O₂ modulates the activity of transcription factors in bacteria (*OxyR* and *PerR*), lower eukaryotes (*Yap1*, *Mafl*, *Hsf1* and *Msn2/4*) and mammalian cells (*AP-1*, *NRF2*, *CREB*, *HSF1*, *HIF-1*, *TP53*, *NF-κB*, *NOTCH*, *SP1* and *SCREB-1*) are summarized. Furthermore, Klamt *et al.* (Klamt et al. 2009) found that the actin-binding protein cofilin is a key target of oxidation. When oxidation of this single regulatory protein is prevented, oxidant-induced apoptosis is inhibited.

As described above, high glucose increases ROS, and antioxidants repairs β-cells undergoing damage by oxidative stress, however, molecular mechanism of ROS on the islet β-cell is not well understood. Therefore, in this paper we focus on the mechanism of gene expression by ROS using H₂O₂ in human pancreatic β-cells derived hybrid cell. Because, transformed cell used for in vitro experiment, the cells sometime lost their characteristic especially apoptosis pathway, otherwise cell will die. Furthermore, human normal islet cells are not available easily, 1.1B4 cells, the hybrid cell of normal islet cell with β-cell derived transformed PANC-1 cell was used here (McCluskey et al. 2011). In this paper we analyzed the effect of H₂O₂ on the gene expression of 1.1B4 with Agilent microarray. Then key protein was knock down by siRNA method and analyzed cell viability under the H₂O₂ addition. And gene expression of siRNA treated cells was analyzed by microarray again to find how this knock down of the key protein effect gene expression.

RESULTS

Effect of H₂O₂ on 1.1B4 cells growth and gene expression analyzed by DNA microarray

Human pancreatic islet derived 1.1B4 cells were exposed to 10, 20, 30, 40, 50, 100, 200 μM H₂O₂, and the cell viability was measured by MTT assay (Fig. 1). We found that there was no toxicity of H₂O₂ up to 50 μM, and the cell growth was inhibited at 100 μM of H₂O₂ and completely suppressed at 200 μM. From this result, it was suggested that even the 1.1B4 cells express antioxidant enzymes to survive from H₂O₂ stress but at higher concentration above 50 μM of H₂O₂, cells were not able to grow well because of not enough antioxidant defense system. Therefore, to find antioxidant system of these cells, comprehensive analysis of gene expression of H₂O₂ treated 1.1B4 cells was analyzed by DNA microarray at 100 μM H₂O₂. The 1.1B4 cells

Fig.1. Effect of H_2O_2 concentration on the cell growth of 1.1B4 islet derived cell. Islet derived 1.1B4 cell was seeded and cultured in RPMI-1640 medium, and 24 hours later cell was added H_2O_2 to 10, 20, 30, 40, 50, 100, 200 μM and cell viability was analyzed at 24, 48 and 72 hours by MTT assay. (n=5)

Table 1. Comparison of expression of genes measured by microarray and real time PCR between H₂O₂ treated cells and control cells.

Average of 2 microarray data of H₂O₂ treated cells was compared with 1 microarray data of control cells after normalization. For comparison with real time PCR, 12 RNA from 12 cell culture dishes in 2 groups obtained for microarray were used, and the result of real time PCR was expressed by average of GAPDH and β -Actin. * p<0.05

Gene Symbol	Gene Name	Expression ratio by Microarray (H ₂ O ₂ /Controll)	Expression ratio by Real Time PCR(H ₂ O ₂ /Controll)
<i>HMOX1</i>	<i>heme oxygenase (decycling) 1</i>	4.73	2.06*
<i>SOD1</i>	<i>superoxide dismutase 1, soluble</i>	1.22	1.05
<i>SOD2</i>	<i>superoxide dismutase 2, mitochondrial</i>	1.17	1.36*
<i>SOD3</i>	<i>superoxide dismutase 3, extracellular</i>		1.89*
<i>CAT</i>	<i>Catalase</i>	0.84	1.08
<i>GPX1</i>	<i>glutathione peroxidase 1</i>	1.86	1.21*
<i>GCLC</i>	<i>glutamate-cysteine ligase, catalytic subunit</i>	2.28	2.27*
<i>FOSB</i>	<i>FBJ murine osteosarcoma viral oncogene homolog B</i>	906.21	71.81*
<i>JUNB</i>	<i>jun B proto-oncogene</i>	7	3.65*
<i>KEAP1</i>	<i>kelch-like ECH-associated protein 1</i>	0.99	1.35
<i>MAFF</i>	<i>v-maf musculoaponeurotic fibrosarcoma oncogene homolog F</i>	15.59	9.18*
<i>CYCS</i>	<i>cytochrome c, somatic</i>	2.63	1.37*
<i>CIDEA</i>	<i>cell death-inducing DFFA-like effector a</i>	8.4	1.40*
<i>OSGIN1</i>	<i>oxidative stress induced growth inhibitor 1</i>	12.37	6.38*
<i>REL</i>	<i>v-rel reticuloendotheliosis viral oncogene homolog</i>	1.88	1.67*
<i>NR4A3</i>	<i>nuclear receptor subfamily 4, group A members 3</i>	136.75	24.9*
<i>MAOA</i>	<i>monoamine oxidase A</i>	2.6	2.16*
<i>NFE2L2</i>	<i>nuclear factor (erythroid-derived 2)-like 2 (NRF2)</i>	1.65	1.85*
<i>NR4A1</i>	<i>nuclear receptor subfamily 4, group A, member 1</i>	7.79	6.02*
<i>CXCL3</i>	<i>chemokine (C-X-C motif) ligand 3</i>	14.46	10.4*
<i>GPR68</i>	<i>G protein-coupled receptor 68</i>	6.42	2.08*
<i>NFKB1A</i>	<i>nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha</i>	4.35	3.02*
<i>TP63</i>	<i>tumor protein p63</i>	34.72	1.86*
<i>SEPP1</i>	<i>selenoprotein P, plasma, 1</i>	10.55	2.46*
<i>PMAIP1</i>	<i>phorbol-12-myristate-13-acetate-induced protein 1</i>	10.11	5.44*
<i>BTK</i>	<i>Bruton agammaglobulinemia tyrosine kinase</i>	9.51	4.35*
<i>IL8</i>	<i>interleukin 8</i>	7.51	3.70*
<i>EGR1</i>	<i>early growth response 1</i>	79.85	25.96*
<i>NQO1</i>	<i>NAD(P)H dehydrogenase, quinone 1</i>	0.86	1.39*

were treated with 100 μ M H₂O₂ for 4 hours, then total RNA was extracted and labeled with Cy-3 then analyzed by Agilent human whole genome microarray. Results were submitted to NCBI GEO data base with accession number GSE83369. (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83369>). Total 2903 genes were up-regulated more than two times of untreated cells, and 2283 genes were down regulated less than 1/2. Most of the gene were classified unknown or others, the genes classified to nucleotide metabolism were major group (Table 1, Supplemental Fig. S1, S2). Some of gene expression was confirmed by RT- real time PCR (Table 1). Antioxidant enzyme *HMOX1*, was up-regulated, but *SOD1* and *CAT* were not changed. On the other hand, *GPXI* and *GCLC*, a glutathione synthesis and antioxidant enzymes, were increased. Among the transcription factors that are up-regulated by H₂O₂ described by Marinho *et al.* (Marinho et al. 2014), *FOSB* and *JUNB* were strongly up-regulated, suggesting AP-1 pathway was up-regulated. Several apoptosis related enzymes were also up-regulated. Among the antioxidant pathway (Marinho et al. 2014), *KEAP1* was not changed. There were many genes that function was unknown or slightly known, up-regulated or down-regulated. Among them, *NR4A3*, kind of orphan receptor, was up-regulated significantly, therefore, the effect of H₂O₂ concentration on the expression of this gene was measured (Fig. 2A). The expression of *NR4A3* was increased by H₂O₂ concentration until 100 μ M at 4 hours significantly. Therefore, the expression of *NR4A3* after H₂O₂ treatment was measured up to 24 hours (Fig. 2B). It is interesting that *NR4A3* expression was increased rapidly and decreased to original level by 24 hours, suggesting that the *NR4A3* is early responsible gene to H₂O₂ stress, and it is suggested that NR4A3 might have important role in antioxidant defense system in 1.1B4 cell against H₂O₂ stress.

Knock down of *NR4A3* mRNA by RNA interference

To investigate the roll of *NR4A3* in 1.1B4 cell, *NR4A3* mRNA was knocked down with siRNA. The 1.1B4 cells were incubated with 5 nM siRNA of *NR4A3* for 48 hours, the mRNA of *NR4A3* was decreased to 29.6% of control cell (Fig. 3A). The 1.1B4 cells knocked down with siRNA of *NR4A3* were then treated with 100 μ M H₂O₂, and analyzed cell viability by MTT assay (Fig. 3B). After 72 hours, the viability of siNR4A3 treated 1.1B4 cell was significantly decreased, and when H₂O₂ was added to culture medium, the growth was completely stopped.

A **B**

*p<0.05, **p<0.01

Fig. 2. Effect of H₂O₂ concentration on *NR4A3* gene expression and time course of expression.

A, Human 1.1B4 cell (1.0×10^5 cells/ml) was incubated in 2ml culture medium and 24 hours later medium was changed to individual concentration of H₂O₂ containing medium, then 4 or 24 hours later total RNA was extracted with Isogen. Expression of *NR4A3* was measured by real time PCR with condition B. n=3. **B**, After addition of 100 μM of H₂O₂, cell was incubated individual time and RNA was extracted. n=3. β-Actin was used for control.

A

Fig. 3.

Fig. 3. Effect of *NR4A3* knocked down on the expression of *NR4A3* and cell viability. **A**, After knocked down of *NR4A3* gene in 1.1B4 cell with siRNA, total RNA was extracted and expression of *NR4A3* and β -actin was measured by real time PCR with Cyber green method with condition B. $n=3$ * $p<0.05$. **B**, After knocked down of *NR4A3* gene of 1.1B4 cell, cell was incubated with 100 μM of H_2O_2 , and cell viability was measured by MTT assay. $n=5$. NC: negative control; NFW: nuclease free water added cell; siNR4A3: siRNA of *NR4A3* treated cell.

Comprehensive analysis of gene expression in 1.1B4 cells by DNA microarray after the knock down of *NR4A3* with siRNA

To analyze the role of *NR4A3* in pancreatic islet cell further, we performed comprehensive analysis of gene expression of 1.1B4 cells by Agilent human DNA microarray after the knocked down of *NR4A3* mRNA. Data were submitted to NCBI GEO data base with accession number GSE86924. (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE86924>). We found 1,044 genes were significantly increased over 1.5 folds and 859 genes were significantly decreased under 0.67 folds (Supplemental Fig. S3A, B). Using those genes, pathway analysis was performed by KEGG pathway database. (Supplemental Table S2). Many genes in PI3K-Akt signaling pathway were up-regulated, and Rap1 signaling pathway genes were down-regulated. Changes in gene expression measured by microarray of transcription factor, related to H₂O₂ addition (Marinho et al. 2014), antioxidant genes and major cell growth related genes are also listed in Table 2. Some of these results were confirmed by real time PCR (Fig. 4A - N, Supplemental Fig. S4A - H). *NR4A3* expression was down regulated (Fig. 4A) but this group of orphan receptor, *NR4A1* (Fig. 4B) was not changed and *NR4A2* (Fig. 4C) was up-regulated. Among the genes of antioxidant enzymes, the expression of *HMOX1* was decreased when measured by microarray, and confirmed by real time PCR (Fig. 4E), the expression of *HMOX1* was reduced to 68%. However, *GCLC* (Supplemental Fig. S4E) was only 0.88 times of control, and *SOD1* (Supplemental Fig. S4F) was not changed significantly. We found interesting gene expression of antioxidant enzyme of *GLRX* (Fig. 4F). After the knocked down the *NR4A3*, *GLRX* expression was increased, different from *HMOX1*. *SOD3* was also increased by *NR4A3* knock down (Fig.4G). Among the transcription factors related to H₂O₂ oxidation, *MAFA* was slightly down regulated measured by microarray but was not changed measured by real time PCR (Supplemental Fig. S4A - D). Other redox sensitive transcription factors were not down regulated.

Gizard *et al.* (Gizard et al. 2011) reported that NOR1 (NR4A3 or Nur77) is recruited to a nerve growth factor-induced clone B response element (NBRE, aaaggaca or aaaggtca) site (Philips et al. 1997). Therefore, whether the *HOMX1* gene has NBRE sequence, or not, was investigated. The *HMOX1* gene structure was obtained from NCBI database (AY460337). There are two NBRE like sequence of aaggtca, tentative NR4A3 responsible element, at second intron (Fig. 5). Transcription factor responsible element in second intron might works when H₂O₂ is present in 1.1B4 cell culture medium.

Table 2. Changes of gene expression after siRNA of NR4A3 treatment measured by microarray.

Average of 3 microarray data of siRNA of NR4A3 treated cells after normalization was compared with average of 3 microarray data of control cells.

Gene symbol	Gene name	Fold changes (SiRNA vs Cont)	Gene symbol	Gene name	Fold changes (SiRNA vs Cont)
Orphan receptor			Cell cycle		
NR4A3	nuclear receptor subfamily 4, group A members 3	0.85	E2F1	E2F transcription factor 2	0.66688
NR4A2	nuclear receptor subfamily 4, group A, member 2	1.06	CDK1	cyclin-dependent kinase 1	0.7342
			CDK2	cyclin-dependent kinase 2	0.34304
Antioxidant responsible Transcription Factor			CDK4	cyclin-dependent kinase 4	0.87803
MAFA	v-maf musculoaponeurotic fibrosarcoma oncogene homolog A (avian)	0.75	CCNA1	cyclin A1	0.84123
FOSB	FBJ murine osteosarcoma viral oncogene homolog B	0.96	CCNB1	cyclin B1	0.72389
FOS	FBJ murine osteosarcoma viral oncogene homolog	2.5	CCND1	cyclin D1	0.99547
JUNB	jun B proto-oncogene	1.02	CCNE1	cyclin E1	0.44222
JUN	jun proto-oncogene	0.76	CCNY	cyclin Y	0.62658
KEAP1	kelch-like ECH-associated protein 1	0.93	CDKN1B	cyclin-dependent kinase inhibitor 1B (p27, Kip1)	1.66505
NFE2L2	nuclear factor (erythroid-derived 2)-like 2, NRF2	No data	CDKN1B	cyclin-dependent kinase inhibitor 1B (p27, Kip1)	1.66505
CREB5	cAMP responsive element binding protein 5	2			
TP53	tumor protein p53	0.83	Others		
NOTCH4	notch 4	1.36	RAB5B	member RAS oncogene family (RAB5B)	0.54527
NFKB1	nuclear factor of kappa light polypeptide gene enhancer in B-cells 1	0.95	RAP1B	RAP1B, member of RAS oncogene family	0.58507
NFKB2	nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (p49/p100)	0.77	TGFB3	transforming growth factor, beta 3	0.50684
SP1	Sp1 transcription factor	0.8	PIK3CA	phosphoinositide-3-kinase, catalytic, alpha polypeptide	1.24095
HIF1A	hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)	1.02	MRAS	muscle RAS oncogene homolog	0.4872
SREBF1	sterol regulatory element binding transcription factor 1	0.76	RAP1B	RAP1B, member of RAS oncogene family	0.59426
HSF1	heat shock transcription factor 1	0.84	SREBF1	sterol regulatory element binding transcription factor 1 (A_33_P3222139)	0.85885
			HIF1A	hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)	0.91634
Antioxidant Responsible Enzyme			BAD	BCL2-associated agonist of cell death	0.65799
HMOX1	heme oxygenase (decycling) 1	0.48585	AKT1	v-akt murine thymoma viral oncogene homolog 1	0.85096
GCLC	glutamate-cysteine ligase, catalytic subunit	0.88148	BAD	BCL2-associated agonist of cell death	2.09416
GPX1	glutathione peroxidase 1(A_33_P3239849)	0.81867	FOXO1	Homo sapiens forkhead box O1 (FOXO1), mRNA [NM_002015]	1.0298
GPX1	glutathione peroxidase 1(A_33_P3354322)	0.81248	FOXO3	Homo sapiens forkhead box O3 (FOXO3), transcript variant 1, mRNA	1.08933
CAT	catalase	1.59628	GAPDH	glyceraldehyde-3-phosphate dehydrogenase	0.73695
SOD1	superoxide dismutase 1, soluble	0.87111	TBP	TATA box binding protein	0.93576
SOD2	superoxide dismutase 2, mitochondrial	0.6454	ACTB	actin, beta	0.89854
SOD3	superoxide dismutase 3, extracellular	0.65966			
GLRX	glutaredoxin (thioltransferase)	1.97201			
GLRX2	glutaredoxin 2	0.83005			
TXN	thioredoxin	0.82374			
PMEL	premelanosome protein	1.6221

Fig. 4.

Fig. 4.

Fig. 4.

M $p=0.00$

N $p=0.00$

Fig. 4. Changes of Various Gene expression in 1.1B4 Cell after the knocked down of *NR4A3* mRNA measured by real time PCR.

(A) Human 1.1B4 cells were incubated 24hours, half of them (n=9) were added siRNA (siNR4A3) and another half (n=9) were added nuclease free water (NFW), and after 52 hours, mRNA was purified with RNeasy Plus Mini Kit. Expression of *NR4A3* was measured by real time PCR with Taqman method using condition C. TATA binding protein (TBP) mRNA was used for internal control. (B) The *NR4A1* expression, (C) *NR4A2* expression. (D) - (E) *NFKB1* and *HMOX1* respectively. (F) *GLRX* expression (n=8). (G) Human 1.1B4 cells were incubated 24hours, half of them (n=3) were added siRNA (siNR4A3) and another half (n=3) were added NFW, and after 52 hours, mRNA was purified with Isogen reagent and treated with RNase-free DNase. Expression of *SOD3* was measured by real time PCR with Cyber green method using condition B. b-Actin was used for control. (H) Expression of *GPX1* using same method of (G). (I) *E2F1* using same method of (A) (n=8). (J) - (M) *p27Kip1*, *CDK2*, *CCNE1* and *PMEL*, respectively, using same method of (A). (N) *RAB5B* using same method of (A) (n=8).

Fig. 5. Gene structure of *HMOX1*.

Human *HMOX1* gene was obtained from PubMed gene (ACCESSION AY460337). NR4A3 binding motif of aaggtca were searched and abstract of the gene structure was displayed. NFkB shows NFkB binding motif. FOSJUN shows FOS and JUN binding motif. EX1 etc. shows exon 1 etc.

As described before, knock down cells decreased cells growth, therefore, cell cycle regulation genes that expression was modified by siRNA of *NR4A3*, were investigated. *CCNE1* (*Cyclin E*) and *CDK2* expressions were decreased remarkably (Table 2). Then those down-regulations were confirmed by RT-real time PCR (Fig. 4K, L). *CDK2* and *CCNE1* expression were decreased strongly compared with un-treated cell. To investigate the mechanism, gene structure of *CDK2* and *CCNE1* were analyzed (Supplemental Fig. S5A, B and S6). There was no aaggtca at up-stream of *CDK2*, however, down-stream of *CDK2*, there is aaggtca sequence. This area is also up-stream of the gene of *RAB5B*, therefore, the effect of siRNA of *NR4A3* on this *RAB5B* expression was investigated. There is also *PMEL* is near this area, the expression of *PMEL* was also measured. As shown in Fig 4M and N, those two genes expression were also decreased or increased by siRNA of *NR4A3*. It is, therefore, suggested that *NR4A3* promotes proliferation through activating *Rap1* signaling pathway, therefore, it is also suggested that the *NR4A3* contribute to homeostasis against extracellular stress such as oxidative stress.

DISCUSSION

It is known that pancreatic islet β -cell weakly express antioxidant enzymes, therefore, oxidative stress enhances diabetes mellitus. As shown in this experiment of microarray of H_2O_2 treated 1.1B4 cell, among the antioxidant enzymes, *HMOX1*, *GPX1* and *GCLC* were up-regulated (Table 1). Other major antioxidant enzymes were not up-regulated, for example, *SOD1* and *CAT* were not induced by H_2O_2 addition, therefore, those *HMOX1*, *GPX1* and *GCLC* enzymes are responsible for antioxidant defense system in pancreatic cell, and those are very important enzymes for defending against oxidative stress at diabetes. Alam and Cook (Alam and Cook 2007) described that there are 4 pathways that regulates *HMOX1* gene expression at oxidative stress. But here, not only those 4 pathway, we showed that *HMOX1* and *NR4A3* were up-regulated when H_2O_2 was added, and the expression of *HMOX1* was down-regulated when *NR4A3* was knocked down by siRNA resulting loss of antioxidative resistance. Furthermore, the second intron of *HMOX1* has NBRE like sequence of aaggtca. Therefore, we concluded that the *NR4A3* is antioxidant responsible transcription factor and regulates *HMOX1* expression at oxidative stress. Not only 1.1B4 cell, *NR4A3* expression was also increased in HUCF2 cell at oxidative stress (Shimizu et al. 2015). Therefore, we propose that the *NR4A3* is new oxidative stress responsible transcription factor not listed before (Marinho et al. 2014), and this pathway is major antioxidant pathway in 1.1B4 cell (Graphical abstract). *FOSB* is also up-regulated by

Graphical abstract

Hydrogen peroxide induces NR4A3 and binds to aaggtca sequence of HMOX1, and increased transcription of HMOX1. Resulting heme oxygenase produces biliverdin, antioxidants, from heme. NR4A3 also bind to aaggtca sequence of CDK2 and CCNE1, resulting CDK2 and Cyclin E. CDK2 bind to cyclin E and cell goes from G1 to S phase.

H₂O₂ in this experiment, therefore, FOS pathway might be also another major antioxidant pathway.

There are two NBRE like sequence of aaggtca in *HMOX1*. NBRE contains aggtca which is typically recognized by retinoic acid receptor/retinoid X receptor (RAR/RXR) subfamily, and it also includes two aa residues preceding this hexanucleotide (Philips et al. 1997). On the other hand, NR4A1 and NR4A2 forms heterodimer with RXR and binds to direct repeat of aggtca (Safe et al. 2016). Therefore, we propose aaggtca is NR4A3 responsible element.

Micro-array analysis has shown that knock down of *NR4A3* decreased *CDK2* and *CCNE1* (*cyclin E*) expression (Table 2). This result confirmed by real time PCR and promotor region analysis (Fig. 4K and 4L, Supplemental Fig. S5A, B and S6). These results show that *NR4A3* is key transcription factor not only for antioxidant system but also for cell cycle control in 1.1B4 cell. Tessem *et al.* (Tessem et al. 2014) demonstrate that *E2F1* and *CCNE1* are key cell cycle inducers. Our result of *cyclin E* consistent with Tessem results, but our results showed *E2F1* expression was only slightly increased in *NR4A3* knock down cell (Fig. 4I). Therefore, it is suggested that NR4A3 directly controls *cyclin E* and *CDK2*, not via *E2F1*.

Robertson (Robertson 2010) described that main stays of therapy for type 2 diabetes involve drugs that are insulin-centric, i.e., they are designed to increase insulin secretion and decrease insulin resistance. The mechanism for this unrelenting deterioration of β -cell function is related to chronic oxidative stress. This suggests that drug discovery should not exclusively focus on insulin-centric targets, but also include glucose-centric strategies, such as antioxidant protection of the β -cells. This may facilitate repair of β -cells undergoing damage by oxidative stress secondary to chronic hyperglycemia. Furthermore, Gao *et al.* (Gao et al. 2014) have shown that over expression of *NR4A3* results in down-regulation of insulin gene transcription and insulin secretion. It is suggested that oxidative stress up-regulated *NR4A3* expression resulting down-regulation of insulin secretion. Therefore, antioxidant material might recover from islet damage. For this aim, many antioxidants were used for diabetes therapy. For example, Cinnamtannin D-1, one of the main A-type procyanidin oligomers in *C. tamala*, was discovered to dose-dependently reduce palmitic acid- or H₂O₂-induced apoptosis and oxidative stress in INS-1 cells, MIN6 cells, and primary cultured murine islets (Wang et al. 2014), however, antioxidant drugs were not in use for human diabetes. Therefore, we screened new antioxidant from food stuff for

potential therapy use, and recently found strong antioxidant Zeylaniin A from edible vegetable (Nomi et al. 2012). This polyphenol may be used for diabetes treatment.

For microarray experiment, Agilent array was used and confirmed by real time PCR. Some times other type of microarray shows different data from real time PCR and shows false positive data, but Agilent array used here shows very co-related data to real time PCR as far as we measured in this experiment. And three microarray data of same conditions showed very similar data. Use of internal control gene for real time PCR was not easy, because many house keeping genes are known to change their transcription depend on the conditions. Therefore, average of two house keeping genes of *GAPDH* and *β-actin* were used for H₂O₂ experiment. *TBP* was also used for siRNA treated cell because *TBP* was not changed by siNR4A3 treatment measured by microarray (Table 2).

METHODS

Reagents

RPMI-1640 medium (GIBCO, Tokyo or Sigma, Tokyo), FBS (CELLect FBS, MP Bomedical Japan, Tokyo Japan, LOT#7997K or Biowest, Nuaillé, France, Lot.No:S05831S1820), Penicillin (50 IU/ml) - Streptomycine (50 micro g/ml) (ICN Biomedicals, Irvine, California, United States), 96 well plate (IWAKI, Tokyo, Japan), NR4A3 Silencer Select siRNA (Applied Biosystems - Thermo Fisher Scientific Japan, Tokyo, siRNA ID : s15542), Silencer Select Negative Control #1 siRNA, Nuclease Free Water and Lipofectamine 2000 (Applied Biosystems - Thermo Fisher Scientific Japan, Tokyo). RNeasy Plus Mini Kit (Qiagen, Tokyo, Japan), Bio-analyzer (Agilent Technology, Palo Alto, CA, USA). Isogen reagent and RNase-free DNase (Nippon gene, Toyama, Japan). Agilent Whole Human Genome Oligo Microarray 4x44K Ver. 2.0 (Agilent Technology, Palo Alto, CA, USA) Reagents for microarray (Agilent Technology, Palo Alto, CA, USA). Agilent G2565BA Microarray Scanner System (Agilent Technology, Palo Alto, CA, USA). The scanned images were analyzed with Feature Extraction Software 9.5.1.1 (Agilent Technology, Palo Alto, CA, USA) using default parameters (protocol GE1-v5_95_feb07 and Grid: 014850_D_F_20101031). Spot Fire software (TIBCO, NTTCom, Tokyo, Japan) and the GeneSpringGX10 v 7.3.1 (Agilent Technology, Palo Alto, CA, USA). DNaseI-treated total RNA with M-MLV reverse transcriptase (Invitrogen, Carlsbad, CA, USA)

and Oligo (dT) 15 primer (Promega, Madison, WI, USA). STEP ONE PLUS Real Time PCR system (Applied Biosystems Japan, Tokyo, Japan), SYBR GREEN PCR Master Mix (Applied Biosystems Japan, Tokyo, Japan). Takara Premix Ex Taq (Probe qPCR) (TaKaRa, Kyoto, Japan), Designs of PCR probes: Universal Probe Library Assay Design Center at Roche (<http://www.universalprobelibrary.com>). ABI PRISM 7900HT Sequence Detection System (Applied Biosystems, Tokyo, Japan).

Biological Resources: Human pancreatic islet derived cell, 1.1B4

Islet derived hybrid cell of 1.1B4 (ECACC No. 10012801) formed by the electrofusion of a primary culture of human pancreatic islets with PANC-1, a human pancreatic ductal carcinoma cell line (ECACC catalogue number 87092802), was obtained from DS Pharma Biomedical Co., Ltd., Osaka, Japan, and cultured in RPMI-1640 medium (GIBCO, Tokyo or Sigma, Tokyo, Japan) supplemented with 10% FBS (CELLelect FBS, MP Bomedical Japan, Tokyo Japan, LOT#7997K or Biowest, Nuaille, France, Lot.No:S05831S1820) with Penicilin (50 IU/ml) - Streptomycine (50 µg/ml) (ICN Biomedicals, Irvine, CA, USA) in humidified air at 37 °C with 5% CO₂.

Cell toxicity of H₂O₂ and the effect of H₂O₂ on gene expression of 1.1B4

The cells of 1.1B4 were cultured in 96 well plate (IWAKI, Tokyo, Japan) at 6.0×10^3 cells / well with RPMI-1640 medium for 24 hours, then 0, 10, 20, 30, 40, 50, 100, 200 µM H₂O₂ was added to medium, and cell viability was analyzed at 24, 48 and 72 hours by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) colorimetric assay (Mosmann 1983). We found that there was no toxicity up to 150 µM H₂O₂. Next, we examined the effect of H₂O₂ on gene expression by real time PCR. Human 1.1B4 cell (1.0×10^5 cells/ml) was incubated in 2ml culture medium and 24 hours later medium was changed to 25, 50 and 100 µM H₂O₂ containing medium, then 4 or 24 hours later total RNA was extracted with Isogen. Expression was measured by real time PCR with condition B. Cells of 1.1B4 were also treated with 100 µM H₂O₂ and we analyzed gene expression after the exposure to H₂O₂ for 2, 6, 12, 24 hours.

Knock down of NR4A3 mRNA by RNA interference

NR4A3 Silencer Select siRNA (ABI, siRNA ID: s15542) was used for knocked down the *NR4A3* mRNA, and Silencer Select Negative Control #1 siRNA (ABI) and Nuclease Free Water (ABI) were used for negative control. *NR4A3* mRNA of 1.1B4 cells was knocked down by incubating with 5 nM siRNA and 0.1 % of Lipofectamine 2000 for 48 h. Total RNA was isolated by Isogen then mRNA was measured as described before. The 1.1B4-siNR4A3 treated cells or control cells were treated with or without 100 μ M H₂O₂, and cell viability was assayed by MTT assay.

Extraction and purification of mRNA

For micro-array, total RNA was isolated using the RNeasy Plus Mini Kit (Qiagen, Tokyo, Japan). The purity of RNA was assessed by a Bio-analyzer (Agilent Technology, Palo Alto, CA, USA) before microarray analysis. The RNA thus obtained was also used for real time PCR. For only real time PCR use, total RNA was extracted with Isogen reagent (Nippon gene, Toyama Japan) as described previous paper (Oba et al. 2006) and treated with RNase-free DNase (Nippon gene, Toyama, Japan).

Global analysis of gene expression by DNA microarray in 1.1B4 cells

(Experiment A) Pancreatic derived 1.1B4 cell was seeded as 1.5×10^5 cells / ml to RPMI-1640 medium (GIBCO, Tokyo, Japan) in 6 well plate, and 24 hours later, cell was incubated with 100 μ M H₂O₂ 4 hours, then total RNA was extracted with RNeasy Mini Kit (Qiagen, Tokyo) (n=6 for control and treated respectively). The quality of all 12 RNA samples was checked by Agilent RNA 6000 Nano Reagents Part1 (Agilent Technologies, Tokyo, Japan), and 3 samples of RNA were combined to one microarray sample, then analyzed by Agilent human microarray by the protocol of manufacture. In brief, Cy3-labeled cRNA was generated from 200 ng input total RNA using Agilent's Low Input Quick Amp Labeling Kit (Agilent Technologies, Tokyo, Japan). cDNA was generated with the primer containing a T7 polymerase promoter, and then was transcribed into cRNA in accompany with the dye labeling. For every type of cell, 1.65 μ g cRNA from each labeling reaction was hybridized to the Agilent Whole Human Genome Oligo Microarray (Agilent Technologies, Tokyo, Japan). The Whole Human Genome Oligo Microarray is in a 4 x 44k slide format and each block represents more than 41,000 unique

human genes and transcripts. After hybridization, the slides were washed and then scanned with the Agilent G2565BA Microarray Scanner System (Agilent Technologies, Tokyo, Japan). The scanned images were analyzed with Feature Extraction Software 9.5.1.1 (Agilent Technologies, Tokyo, Japan) using default parameters (protocol GE1-v5_95_feb07 and Grid: 014850_D_F_20101031) to obtain background subtracted and spatially detrended Processed Signal intensities. Features flagged in Feature Extraction as Feature Non-uniform outliers were excluded. Data were further normalized using Gene Spring GX10 using default parameters of recommended protocol with Median shift normalization to the 75th percentile and baseline transformed to the median of all samples. Data normalization and filtering was performed using Spot Fire software (TIBCO, NTTCom, Tokyo, Japan) and the Gene Spring v 7.3.1 (Agilent Technologies, Tokyo, Japan). After the reading, one of microarray for control sample was scratched, therefore, the gene expression was compared average of two treated array samples vs one control sample. These 12 RNA samples were also used for real time PCR analysis.

(Experiment B) The microarray experiment was designed as follows. Human 1.1B4 cells (n=24) were incubated in RPMI-1640 medium 24hours, half of them (n=12) were added siRNA (siNR4A3) and another half (n=12) were added nuclease free water (NFW), and 48 hours later 3 cells of each group were added H₂O₂ to 100 mM. Those 6 cells samples were stored but not used for this paper. After 52 hours, mRNA of remaining 18 cells was purified with RNeasy Plus Mini Kit. The quality of mRNA was checked by Bio-analyzer, 3 mRNA samples were combined to one microarray sample (3 knocked down microarray samples and 3 control microarray samples), and applied on microarray analysis with Agilent Whole Human Genome Oligo Microarray 4 x 44k. The expression was analyzed by Gene-spring software and Spot fire software. Average of 3 array samples treated by H₂O₂ was compared to average of 3 array samples without treated by H₂O₂. For real time PCR, 18 RNA samples were used individually. Real time PCR with Taqman method was used (condition C). *TBP* mRNA was used for internal control. All the sample-labeling, hybridization, washing, and scanning steps were conducted following the manufacturer's specifications as described before.

Real-time RT-PCR

The cDNAs were synthesized from DNaseI-treated total RNA with M-MLV reverse transcriptase (Invitrogen, Carlsbad, CA, USA) and Oligo (dT) 15 primer (Promega, Madison, WI, USA) according to the manufacturer's instructions. Analysis of gene expression was

performed with a STEP ONE PLUS Real Time PCR system (Applied Biosystems Japan, Tokyo, Japan) using SYBR GREEN PCR Master Mix (Applied Biosystems Japan, Tokyo, Japan) and the primers listed in Supplemental Table S1. **Condition A**, 0.2 µl of cDNA was mixed with 0.1 µl forward primer (50 pmol / ml), 0.1 µl reverse primer (50 pmol / ml), 4.7 µl water and 5 µl 2×SYBR Green PCR Master MIX. Reactive condition was first incubated at 50 °C for 2 min and 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 sec and 60 °C for 1 min, then final of 95 °C for 15 sec, 60 °C for 15 sec and 95 °C for 15 sec. Normalization of the data was achieved by quantitating the cycle time at an arbitrary fluorescence intensity in the linear exponential phase using Step One Plus Real-Time system Software (Applied Biosystems Japan, Tokyo, Japan) by calculating the ratio of the relative concentration of each mRNA relative to that of average of *GAPDH* and *β-actin* or *β-actin* only. To confirm amplification specificity, the PCR products from each primer pair were subjected to a melting curve analysis. The relative quantification of gene expression was computed by using the comparative Ct (threshold cycle) method. **Condition B**, 0.2 µl of cDNA was mixed with 1 µl forward primer (50 µM), 1 µl reverse primer (50 µM), 2.8 µl water and 5 ml 2×SYBR Green PCR Master MIX with same reaction condition.

Condition C was performed with a TaqMan method. Briefly, the synthesized cDNA products were subjected to real-time PCR in a reaction mixture (10 µl) containing Takara Premix Ex Taq (Probe qPCR) (TaKaRa, Kyoto, Japan), 200 nM hydrolysis probes and 200 nM PCR primers. Two µl of cDNA was mixed with 5 µl of Premix Ex Taq (2×conc.), 0.2 µl of ROX Reference Dye (50X), 0.2 µl of forward primer (10 µM), 0.2 µl of reverse primer (10 µM) and 0.2 µl of TaqMan Probe (10 µM). Designs of PCR probes and primers were obtained from the Universal Probe Library Assay Design Center at Roche (<http://www.universalprobelibrary.com>) (Supplemental Table S1). Real-time amplifications were performed on the ABI PRISM 7900HT Sequence Detection System (Applied Biosystems, Tokyo, Japan). The settings for the thermal profile were an initial denaturation (30 s at 95°C) followed by 40 amplification cycles: denaturation for 5 s at 95°C; annealing for 30 s at 60°C. Gene-specific standard curves were generated using 2-fold serial dilutions of cDNA. The amount of target mRNA was expressed as the ratio to *TBP* mRNA. Primers are listed in Table S1.

Classification of genes

The genes up-regulated or down-regulated significantly were categorized by panther process name of the gene prepared by Agilent data for human microarray.

Pathway analysis and statistical analysis

The genes up-regulated significantly were submitted to the KEGG pathway analysis at Kyoto University. Statistical analysis was performed with R (R Core Team; R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2017 URL <https://www.R-project.org/>). For comparisons between groups, analysis of covariance (ANNOV method and t test for post hoc test) was used to assess the statistical significance of mean differences between groups, and pairwise t test was used for post hoc testing. The significance levels were set at $Pr(>F)$ 0.05 for anova, and $p < 0.05$ and $p < 0.01$ for post hoc testing.

DATA ACCESS

Microarray data of H₂O₂ treated cells is available in NCBI GEO data base with accession number GSE83369. (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83369>).

Microarray data of siRNA of NR4A3 treated cells is available in NCBI GEO data base with accession number GSE86924. (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE86924>).

COMPETING INTEREST STATEMENT

CONFLICT OF INTEREST

No potential conflicts of interest relevant to this article were reported. Declarations of interest: none. The datasets generated during and/or analyzed during the current study are available from the corresponding author or author of E.U. upon reasonable request. The datasets are also available from NCBI GEO data base. Disclosure Statement: The authors have nothing to disclose.

1

2 **ACKNOWLEDGEMENTS**

3 This work was supported by Japan Society for the Promotion of Science (JSPS) Grant-in-Aid
4 for Scientific Research (A, B and C) [grant numbers 17500533, 20300243, 24240096,
5 25282020], and Grant-in-Aid for Challenging Exploratory Research [grant number 21650196].
6 Founders had no role in study design, data collection and analysis, decision to publish, or
7 preparation of the manuscript.

8

9 **SUPPLEMENTARY DATA**

10 Supplementary Data are available.

11

12 **REFERENCES**

- 13 Alam J, Cook JL. 2007. How many transcription factors does it take to turn on the heme
14 oxygenase-1 gene? *Am J Respir Cell Mol Biol* **36**: 166-174.
- 15 Gao W, Fu Y, Yu C, Wang S, Zhang Y, Zong C, Xu T, Liu Y, Li X, Wang X. 2014. Elevation
16 of NR4A3 expression and its possible role in modulating insulin expression in the pancreatic
17 beta cell. *PLoS One* **9**: e91462.
- 18 Gizard F, Zhao Y, Findeisen HM, Qing H, Cohn D, Heywood EB, Jones KL, Nomiya T,
19 Bruemmer D. 2011. Transcriptional regulation of S phase kinase-associated protein 2 by NR4A
20 orphan nuclear receptor NOR1 in vascular smooth muscle cells. *J Biol Chem* **286**: 35485-35493.
- 21 Klamt F, Zdanov S, Levine RL, Pariser A, Zhang Y, Zhang B, Yu LR, Veenstra TD, Shacter E.
22 2009. Oxidant-induced apoptosis is mediated by oxidation of the actin-regulatory protein cofilin.
23 *Nat Cell Biol* **11**: 1241-1246.
- 24 Marinho HS, Real C, Cyrne L, Soares H, Antunes F. 2014. Hydrogen peroxide sensing,
25 signaling and regulation of transcription factors. *Redox Biol* **2**: 535-562.
- 26 Mathis D, Vence L, Benoist C. 2001 β -Cell death during progression to diabetes. *Nature* **414**:
27 792-798.
- 28 McCluskey JT, Hamid M, Guo-Parke H, McClenaghan NH, Gomis R, Flatt PR. 2011.
29 Development and functional characterization of insulin-releasing human pancreatic beta cell
30 lines produced by electrofusion. *J Biol Chem* **286**: 21982-21992.

- 1 Mosmann T. 1983. Rapid colorimetric assay for cellular growth and survival: application to
2 proliferation and cytotoxicity assays. *J Immunol Methods* **65**: 55-63.
- 3 Nomi Y, Shimizu S, Sone Y, Mai TT, Tien PG, Kamiyama M, Shibamoto T, Shindo K, Otsuka
4 Y. (2012) Isolation and antioxidant activity of zeylaniin A, a new macrocyclic ellagitannin from
5 *Syzygium zeylanicum* leaves. *J Agric Food Chem* **60**: 10263-10269.
- 6 Nguyen T, Nioi P, Pickett CB. 2009. The Nrf2-antioxidant response element signaling pathway
7 and its activation by oxidative stress. *J Biol Chem* **284**: 13291-13295.
- 8 Oba R, Kudo Y, Sato N, Noda R, Otsuka Y. 2006. A new method of competitive reverse
9 transcription polymerase chain reaction with SYBR Gold staining for quantitative analysis of
10 mRNA. *Electrophoresis* **27**: 2865-2868.
- 11 Philips A, Lesage S, Gingras R, Maira MH, Gauthier Y, Hugo P, Drouin J. 1997. Novel
12 dimeric Nur77 signaling mechanism in endocrine and lymphoid cells. *Mol Cell Biol* **17**: 5946-
13 5951.
- 14 Robertson RP. 2004 Chronic oxidative stress as a central mechanism for glucose toxicity in
15 pancreatic islet beta cells in diabetes. *J Biol Chem* **279**: 42351-42354.
- 16 Robertson RP, Harmon JS. 2006. Diabetes, glucose toxicity, and oxidative stress: A case of
17 double jeopardy for the pancreatic islet beta cell. *Free Radic Biol Med* **41**:177-184.
- 18 Robertson RP. 2010. Antioxidant drugs for treating beta-cell oxidative stress in type 2 diabetes:
19 glucose-centric versus insulin-centric therapy. *Discov Med* **9**: 132-137.
- 20 Safe S, Jin UH, Morpurgo B, Abudayyeh A, Singh M, Tjalkens RB. 2016. Nuclear receptor 4A
21 (NR4A) family - orphans no more. *J Steroid Biochem Mol Biol* **157**: 48-60.
- 22 Shimizu Y, Miyakura R, Otsuka Y. 2015. Nuclear receptor subfamily 4, group A, member 1
23 inhibits extrinsic apoptosis and reduces caspase-8 activity in H₂O₂-induced human HUC-F2
24 fibroblasts. *Redox Rep* **20**: 81-88.
- 25 Tessem JS, Moss LG, Chao LC, Arlotto M, Lu D, Jensen MV, Stephens SB, Tontonoz P,
26 Hohmeier HE, Newgard CB. 2014. Nkx6.1 regulates islet beta-cell proliferation via Nr4a1 and
27 Nr4a3 nuclear receptors. *Proc Natl Acad Sci U S A* **111**: 5242-5247. doi:
- 28 Wang T, Sun P, Chen L, Huang Q, Chen K, Jia Q, Li Y, Wang H. 2014. Cinnamtannin D-1
29 Protects Pancreatic β -Cells from Palmitic Acid-Induced Apoptosis by Attenuating Oxidative
30 Stress. *J Agric Food Chem* **62**: 5038-5045.
- 31 Weinberg E, Maymon T, Weinreb M. 2014. AGEs induce caspase-mediated apoptosis of rat
32 BMSCs via TNF α production and oxidative stress. *J Mol Endocrinol* **52**: 67-76.

33 34 TABLE AND FIGURES LEGENDS

Table 1. Comparison of expression of genes measured by microarray and real time PCR between H₂O₂ treated cells and control cells.

Average of 2 microarray data of H₂O₂ treated cells was compared with 1 microarray data of control cells after normalization. For comparison with real time PCR, 12 RNA from 12 cell culture dishes in 2 groups obtained for microarray were used, and the result of real time PCR was expressed by the average of GAPDH and β -actin. * p<0.05

Table 2. Changes of gene expression after siRNA of NR4A3 treatment measured by microarray.

Average of 3 microarray data of siRNA of NR4A3 treated cells after normalization was compared with average of 3 microarray data of control cells.

Fig.1. Effect of H₂O₂ concentration on the cell growth of 1.1B4 islet derived cell.

Islet derived 1.1B4 cell was seeded and cultured in RPMI-1640 medium, and 24 hours later cell was added H₂O₂ to 10, 20, 30, 40, 50, 100, 200 μ M and cell viability was analyzed at 24, 48 and 72 hours by MTT assay. (n=5)

Fig. 2. Effect of H₂O₂ concentration on NR4A3 gene expression and time course of expression.

A, Human 1.1B4 cell (1.0×10^5 cells/ml) was incubated in 2ml culture medium and 24 hours later medium was changed to individual concentration of H₂O₂ containing medium, then 4 or 24 hours later total RNA was extracted with Isogen. Expression of NR4A3 was measured by real time PCR with condition B. n=3. **B**, After addition of 100 μ M of H₂O₂, cell was incubated individual time and RNA was extracted. n=3. β -Actin was used for control.

Fig. 3. Effect of NR4A3 knocked down on the expression of NR4A3 and cell viability. **A**, After knocked down of NR4A3 gene in 1.1B4 cell with siRNA, total RNA was extracted and expression of NR4A3 and β -actin was measured by real time PCR with Cyber green method with condition B. n=3 *p<0.05. **B**, After knocked down of NR4A3 gene of 1.1B4 cell, cell was incubated with 100 μ M of H₂O₂, and cell viability was measured by MTT assay. n=5. NC: negative control; NFW: nuclease free water added cell; siNR4A3: siRNA of NR4A3 treated cell.

Fig. 4. Changes of Various Gene expression in 1.1B4 Cell after the knocked down of *NR4A3* mRNA measured by real time PCR.

(A) Human 1.1B4 cells were incubated 24hours, half of them (n=9) were added siRNA (siNR4A3) and another half (n=9) were added nuclease free water (NFW), and after 52 hours, mRNA was purified with RNeasy Plus Mini Kit. Expression of *NR4A3* was measured by real time PCR with Taqman method using condition C. TATA binding protein (TBP) mRNA was used for internal control. (B) The *NR4A1* expression, (C) *NR4A2* expression. (D) - (E) *NFKB1* and *HMOX1* respectively. (F) *GLRX* expression (n=8). (G) Human 1.1B4 cells were incubated 24hours, half of them (n=3) were added siRNA (siNR4A3) and another half (n=3) were added NFW, and after 52 hours, mRNA was purified with Isogen reagent and treated with RNase-free DNase. Expression of *SOD3* was measured by real time PCR with Cyber green method using condition B. β -Actin was used for control. (H) Expression of *GPX1* using same method of (G). (I) *E2F1* using same method of (A) (n=8). (J) - (M) *p27Kip1*, *CDK2*, *CCNE1* and *PMEL*, respectively, using same method of (A). (N) *RAB5B* using same method of (A) (n=8).

Fig. 5. Gene structure of *HMOX1*.

Human *HMOX1* gene was obtained from PubMed gene (ACCESSION AY460337). NR4A3 binding motif of aagggtca were searched and abstract of the gene structure was displayed. NFkB shows NFkB binding motif. FOSJUN shows FOS and JUN binding motif. EX1 etc. shows exon 1 etc.