Cell Cycle Delay, Pro-metaphase Arrest and C-metaphase Inducing Effects of Petroleum Ether Fraction of Leaf Aqueous Extract of *Clerodendrum viscosum* Vent.

Sujit Roy¹, Lalit Mohan Kundu^{1,2}, Gobinda Chandra Roy^{1,3}, Manabendu Barman¹ and Sanjib Ray^{1*}

¹Molecular Biology and Genetics Unit, Department of Zoology, The University of Burdwan, Golapbag, Purba Bardhaman-713104, West Bengal, India.

²Department of Zoology, Kalna College. Kalna, West Bengal, India.

³Department of Zoology, Dinhat College, Dinhat, Cooch Behar, West Bengal, India.

*Corresponding author, e-mail: sray@zoo.buruniv.ac.in

Summary

Clerodendrum viscosum is a traditionally used medicinal plant. The present study aimed to analyze cell cycle delay, pro-metaphase arrest, and c-metaphase inducing effects of the petroleum ether fraction (AQPEF) of leaf aqueous extract of *C. Viscosum* (LAECV). The LAECV was fractionated with petroleum ether and its metaphase arrest, cell cycle delay, and c-metaphase inducing activities were tested on *A. cepa* root tip cells. The AQPEF induced cell cycle delay, and colchicine like metaphase, c-metaphase, in *A. cepa* root tip cells. Thus, the present study explores AQPEF as an active fraction of LAECV having metaphase arresting, cell cycle delay, and c-metaphase inducing potentials.

Keywords: Diterpenoids; Mitotic abnormality; *Allium cepa*; Metaphase arrest.

Clerodendrum viscosum (Family: Lamiaceae) is widely distributed throughout Asia, Africa, Australia, and America. It is used in traditional, Ayurvedic, Homeopathic, and Unani medicine (Nandi and Lyndem 2015). The whole plant juice is used against worm infection, cough, itching, leprosy, scorpion sting, asthma, bronchitis, fever, etc. (Kirtikar and Basu 1991, Bhattacharjee 2011). The plant is well known for its effectiveness against rheumatism in Unani medicine (Singh *et al.* 1997). It is prescribed to treat postnatal complications, diarrhea, and fresh wounds in the Indian Homeopathic system (Hamilton 1997, Nadkarni and Nadkarni 2002). Its bark juice is used to relieve indigestion and abdominal pain. The different parts of this plant are used as a remedy for asthma, malaria, cataract, diseases of the skin, blood, and lung by the Indian Tribals of Chotanagpur plateau (Kirtikar and Basu 1991).

The plant's ethanolic extracts have antioxidant, antimicrobial, hepatoprotective, wound healing, and antidiarrheal activities (Bhattacharjee *et al.* 2011). The anthelmintic activity was reported in leaf ethanolic, methanolic, and aqueous extracts against *Pheretima posthuma* (Islam *et al.* 2013). The leaves and roots have great potential against different microbial and fungal strains. Acetone and chloroform extracts of *C. viscosum* leaves have an inhibitory effect on the growth of *Shigella sp.*, *Vibrio cholerae*, *Klebsiella pneumonia*, and *Pseudomonas aeruginosa*, etc. (Lobo *et al.* 2010) while ethanolic fraction has shown antifungal activity against *Aspergillus niger*, *A. flavus*, and *Candida albicans* (Modi *et al.* 2010). A saponin isolated

from petroleum ether extract of leaves has analgesic activity (Sannigrahi *et al.* 2009). In vivo, antinociceptive activity of the methanolic extract is comparable to diclofenac sodium drug (Rahman *et al.* 1970). Reduction in CCl₄ induced hepatotoxicity on rats after treatment with methanolic leaf extract reveals its hepatoprotective activity which is further supported by biochemical blood parameters (Sannigrahi *et al.* 2009). Successive ethanolic extract of *C. infortunatum* shows the highest antioxidant activity compared to petroleum ether and chloroform extracts, whereas significant wound healing activity was exhibited by petroleum ether and ethanol extract. These pharmacological effects were also correlated with the total phenolic content of the plant (Gouthamchandra *et al.* 2010). Leaf methanolic extract exerts a restoring effect on blood glucose levels after streptozotocin treatment (Arvind *et al.* 2002, Das *et al.* 2011). Allelochemicals from leaf aqueous extract have been found to harm the growth and germination of weeds in agro-ecosystem (Qasem and Foy 2001, Devi *et al.* 2013). A promising positive correlation is established between the plant's parts and their insect repellent and insecticidal activity (Muh *et al.* 2014).

Acute toxicity test reveals that these plant parts are safe up to 2000 mg kg⁻¹ body weight (Gupta *et al.* 2012). The crude leaf extracts contain phenolics viz. fumaric acids, acetoside, methyl esters of caffeic acids, terpenoids like clerodin, flavonoids such as apigenin, acacetin, scutellarein, quercetin, hispidulin, steroids such as clerodone, clerodolone, clerodol, clerosterol and some fix oils

containing linolenic acid, oleic acids, stearic acid and lignoceric acid (Singhmura 2016).

In our previous study, we have reported that treatment with leaf aqueous extract of *C. viscosum* (LAECV) on root apical meristem cells of wheat and onion gave an increased metaphase frequency along with a reduction in mitotic index, antiproliferative, and apoptosis-inducing effects (Ray *et al.* 2012). The metaphase arrest and cell cycle delay-inducing effects were somewhat comparable to colchicine's action (Ray *et al.* 2013, Kundu and Ray 2016, Roy and Ray 2017).

Colchicine inhibits spindle formation in cells which leads to generating signals delaying the transition from metaphase to anaphase (Salmon *et al.* 1984). Later on, when the concentration of colchicine decreases in the environment, the chromatids separate abnormally and the plant cells become polyploid (Caperta *et al.* 2006). Colchicine treatment to onion root apical meristem cells results in root growth inhibition, root's swelling, haphazardly arranged condensed chromosomes, and increased frequency in metaphase (Ray *et al.* 2013). In another comparative study, colchicine and LAECV treatments on *A. cepa* root tip cells revealed similar cytotoxic effects (Kundu and Ray 2016). The metaphase arrest and cell cycle delay-inducing effects of LAECV raised the key question about its active principle(s) (Ray *et al.* 2013). Therefore, this study aims to fractionate the bioactive molecule(s) of LAECV with non-polar petroleum ether solvent and to test its antiproliferative and metaphase arresting activities. The LAECV was fractionated with petroleum ether and the extract fraction was tested for cell cycle delay, pro-metaphase arrest, and c-metaphase inducing effects on *A. cepa* root tip cells.

Materials and methods

Plant collection and extraction

The fresh *C. viscosum* leaves were collected from Burdwan University campus, West Bengal, India, and it was taxonomically identified by Professor A. Mukherjee, Department of Botany, The University of Burdwan. A voucher specimen (No. BUTBSR011) is maintained in the Department of Zoology, B.U., for future reference.

Fresh leaves were washed in tap water, dried in shade, ground by Philips Mixer Grinder HL1605, and the obtained powder was stored in a tightly sealed container for further use. 100 g of this pulverized material was extracted in 2.5 L of boiling distilled water for 2-3 h and the extract was filtered with filter paper and the extract coded as leaf aqueous extract of *C. viscosum* (LAECV). The LAECV was fractionated by petroleum ether with the help of a magnetic stirrer for 10-12 h. After collection by siphoning, the solvent fraction (AQPEF) was concentrated by rotary vacuum evaporator and stored in a glass container.

Culture and treatment for root growth retardation

1% sodium hypochlorite mediated surface-sterilized *A. cepa* bulbs were placed in 6-well plates containing distilled water and kept in the environmental test chamber for germination (25-27°C, humidity 50%). The 48h aged similar-sized *A. cepa* roots (2-3 cm root length) were treated with 12.5, 25, 50, 100, and 150 $\mu\text{g mL}^{-1}$ concentrations of AQPEF in 1% DMSO continuously for 24, 48, and 72 h. The experiments were performed in triplicate.

Cell cycle delay and c-metaphase inducing effects analysis

The 48h aged similar-sized *A. cepa* roots (2-3 cm) were treated with 50, 100, 150, 200 $\mu\text{g mL}^{-1}$ of AQPEF and 150 $\mu\text{g mL}^{-1}$ of colchicine for 2 and 4 h. The stock solution of the Petroleum ether fraction was prepared in DMSO for treatment. After 2 and 4 h exposure, 8-10 roots were fixed and processed for squash preparation following the standard procedure (Chaudhuri and Ray 2015). The remaining roots were allowed to grow further for another 16 h in distilled water (4h T+16h RS) and subsequently, root tips were fixed. The control group, which had not received any treatment, was maintained in distilled water simultaneously with the treatment groups. The treated and untreated root tips were fixed in aceto-methanol (3 parts methanol: 1part glacial acetic acid) for 24 h and then hydrolyzed for 10 min in 1 M HCl at 60°C. The roots were stained with 2% aceto-orcin and finally squashed in 45% acetic acid (Sharma and Sharma 1999, Ray *et al.* 2013). The well-spread areas of squashed roots were focused under the bright field light microscope for observation and scoring the cell division phase and c- metaphase frequencies.

Scoring and statistical analysis

In the case of squash preparation of *A. cepa* root apical meristem cells, at least three randomly coded slides were observed under the light microscope. Calculation of the mitotic index was done by counting the number of dividing cells per total cells scored for each concentration. Different cell phase frequencies, mitotic index, and c-metaphase cells were analyzed by 2X2 contingency χ^2 -test. Pearson correlation was analyzed using GraphPad Prism 8.4.3.

Results and discussion

The focus of the present investigation was to analyze cell cycle delay, metaphase arrest, and c-metaphase inducing effects of AQPEF in *A. cepa* root apical meristem cells. Many authors suggested that antiproliferative and cytotoxic effects of plant extracts or chemical substances can be evaluated using *A. cepa* root tip cells (Levan 1938, Bakare *et al.* 2000, 2012, 2013, Frescura *et al.* 2012). The root growth

Table 1: Root growth retardation effect of AQPEF in *A. cepa*.

AQPEF ($\mu\text{g mL}^{-1}$)	Root length (cm) (Root length retardation %)			Growth rate (cm/24 h) Growth rate retardation (%)	
	24 h	48 h	72 h	48 h	72 h
0	3.08 $\pm$ 0.10	4.32 $\pm$ 0.10	5.51 $\pm$ 0.11	1.24 $\pm$ 0.10	1.19 $\pm$ 0.03
12.5	2.57 $\pm$ 0.07 (16.51 $\pm$ 0.39)	3.37 $\pm$ 0.04 ^b (22.00 $\pm$ 0.83)	3.73 $\pm$ 0.10 ^a (32.92 $\pm$ 1.01 ^a)	0.79 $\pm$ 0.06 ^b (35.62 $\pm$ 2.62 ^b)	0.36 $\pm$ 0.11 ^a (69.87 $\pm$ 5.05 ^a)
25	2.38 $\pm$ 0.08 ^b (22.68 $\pm$ 0.87 ^b)	2.80 $\pm$ 0.08 ^a (35.08 $\pm$ 1.06 ^a)	3.09 $\pm$ 0.12 ^a (43.92 $\pm$ 1.32 ^a)	0.42 $\pm$ 0.04 ^a (65.48 $\pm$ 3.84 ^a)	0.28 $\pm$ 0.06 ^a (75.98 $\pm$ 2.95 ^a)
50	2.17 $\pm$ 0.07 ^a (29.59 $\pm$ 1.52 ^a)	2.45 $\pm$ 0.09 ^a (43.31 $\pm$ 1.97 ^a)	2.59 $\pm$ 0.18 ^a (52.94 $\pm$ 1.87 ^a)	0.28 $\pm$ 0.16 ^a (76.55 $\pm$ 8.87 ^a)	0.14 $\pm$ 0.03 ^a (87.91 $\pm$ 1.58 ^a)
100	1.80 $\pm$ 0.08 ^a (41.64 $\pm$ 1.33 ^a)	1.99 $\pm$ 0.07 ^a (53.91 $\pm$ 0.74 ^a)	2.13 $\pm$ 0.07 ^a (61.38 $\pm$ 0.78 ^a)	0.19 $\pm$ 0.03 ^a (84.17 $\pm$ 2.43 ^a)	0.13 $\pm$ 0.02 ^a (88.50 $\pm$ 1.05 ^a)
150	1.50 $\pm$ 0.06 ^a (51.33 $\pm$ 1.31 ^a)	1.63 $\pm$ 0.05 ^a (62.31 $\pm$ 0.44 ^a)	1.73 $\pm$ 0.05 ^a (68.51 $\pm$ 0.56 ^a)	0.13 $\pm$ 0.03 ^a (89.32 $\pm$ 1.93 ^a)	0.10 $\pm$ 0.03 ^a (91.01 $\pm$ 1.51 ^a)

Allium cepa root growth retardation effect of the AQPEF was performed in triplicate and data analyzed with Student's t-test. ^a significant at $p < 0.001$, ^b significant at $p < 0.01$ and ^c significant at $p < 0.05$ as compared to their respective control by Student's t-test. Data represented as Mean $\pm$ SEM.

Table 2: Metaphase arrest and c-metaphase inducing effects of AQPEF.

h	Conc. ($\mu\text{g mL}^{-1}$)	Tc	Tdc	MI%	Pro%	Meta%	Ana%	Telo%	C-Meta%
2	0	1959	111	5.67 $\pm$ 0.30	21.95 $\pm$ 4.90	36.86 $\pm$ 2.97	25.87 $\pm$ 3.32	15.13 $\pm$ 0.70	1.66 $\pm$ 1.66
	50	2181	144	6.63 $\pm$ 0.41	16.67 $\pm$ 1.27	42.41 $\pm$ 1.35	25.64 $\pm$ 4.45	15.25 $\pm$ 2.48	22.18 $\pm$ 1.52 ^a
	100	2445	183	7.58 $\pm$ 0.5 ^c	15.93 $\pm$ 2.63	56.52 $\pm$ 1.96	18.48 $\pm$ 3.42	11.09 $\pm$ 0.94	39.2 $\pm$ 4.05 ^a
	150	2404	234	9.76 $\pm$ 0.26 ^a	16.32 $\pm$ 1.01	63.67 $\pm$ 3.27 ^c	14.95 $\pm$ 1.28	5.05 $\pm$ 1.76 ^b	55.62 $\pm$ 2.03 ^a
	200	2389	251	10.51 $\pm$ 0.80 ^a	17.74 $\pm$ 1.90	53.67 $\pm$ 2.05	17.96 $\pm$ 2.23	10.63 $\pm$ 1.13	45.15 $\pm$ 3.86 ^a
	150©	1987	248	12.55 $\pm$ 1.44 ^a	11.91 $\pm$ 1.41	79.75 $\pm$ 4.46 ^a	5.98 $\pm$ 2.74 ^a	2.32 $\pm$ 0.49 ^a	77.95 $\pm$ 5.63 ^a
4	0	2914	207	7.09 $\pm$ 0.62	19.02 $\pm$ 0.97	37.72 $\pm$ 2.60	30.77 $\pm$ 2.07	12.45 $\pm$ 0.51	0.00 $\pm$ 0.00
	50	3423	352	10.3 $\pm$ 0.56 ^a	8.72 $\pm$ 2.61 ^b	66.47 $\pm$ 5.70 ^a	14.54 $\pm$ 1.98 ^a	10.24 $\pm$ 2.27	49.38 $\pm$ 5.71 ^a
	100	2162	222	10.3 $\pm$ 0.29 ^a	10.04 $\pm$ 1.61 ^c	81.44 $\pm$ 1.61 ^a	6.67 $\pm$ 1.04 ^a	1.83 $\pm$ 0.52 ^a	80.08 $\pm$ 1.65 ^a
	150	3035	403	13.25 $\pm$ 0.72 ^a	9.34 $\pm$ 1.11 ^c	78.48 $\pm$ 2.41 ^a	8.76 $\pm$ 2.53 ^a	3.39 $\pm$ 0.76 ^a	70.86 $\pm$ 2.74 ^a
	200	3525	352	9.95 $\pm$ 0.80 ^a	14.33 $\pm$ 2.74	76.43 $\pm$ 2.61 ^a	8.00 $\pm$ 0.27 ^a	2.26 $\pm$ 0.48 ^a	68.88 $\pm$ 1.67 ^a
	150©	3286	301	9.17 $\pm$ 0.56 ^b	9.35 $\pm$ 1.92 ^b	84.88 $\pm$ 1.26 ^a	4.70 $\pm$ 0.63 ^a	1.00 $\pm$ 0.07 ^a	83.00 $\pm$ 1.93 ^a
4+16	0	2449	241	10.06 $\pm$ 0.92	24.63 $\pm$ 2.84	38.95 $\pm$ 0.78	23.27 $\pm$ 0.93	13.11 $\pm$ 2.99	0.00 $\pm$ 0.00
	50	3946	276	7.00 $\pm$ 0.29 ^a	14.83 $\pm$ 0.34 ^c	45.32 $\pm$ 0.66	24.58 $\pm$ 0.94	15.23 $\pm$ 0.43	0.36 $\pm$ 0.36 ^a
	100	2130	177	8.45 $\pm$ 1.64	12.77 $\pm$ 3.18 ^c	65.69 $\pm$ 3.87 ^b	12.23 $\pm$ 0.72 ^c	9.28 $\pm$ 1.71	43.05 $\pm$ 13.28 ^a
	150	2068	136	6.51 $\pm$ 0.77 ^a	20.08 $\pm$ 0.92	52.42 $\pm$ 6.02	18.64 $\pm$ 2.49	8.82 $\pm$ 3.99	33.73 $\pm$ 1.91 ^a
	200	2162	161	7.43 $\pm$ 0.38 ^b	25.00 $\pm$ 2.85	35.78 $\pm$ 2.46	29.68 $\pm$ 2.72	9.51 $\pm$ 2.22	24.81 $\pm$ 5.00 ^b
	150©	1922	69	3.55 $\pm$ 1.10 ^a	23.35 $\pm$ 3.40	38.81 $\pm$ 1.66	11.60 $\pm$ 2.05	26.20 $\pm$ 3.57	9.96 $\pm$ 3.40 ^a

^a Significant at $p < 0.0001$, ^b significant at $p < 0.001$ and ^c significant at $p < 0.05$ as compared with their respective control by 2x2 Contingency χ^2 - test with respective $df = 1$. h; Hours, Comp; Compound, Conc; Concentration, Tc; Total no of cells, Tdc; Total no of dividing cells, MI; Mitotic

index, Pro; Prophase, Meta; Metaphase, LAECV; Leaf aqueous extract of *C. viscosum*, h; hour, Anaphase, Ana; Telo; Telophase, C-Meta; Colchicine like metaphase or C- metaphase, AQPEF; Petroleum ether fraction of LAECV, ©; Colchicine. Data represented as Mean $\pm$ SEM.

Fig. 1: Photomicrograph showing the squash preparation of A, Untreated; B, AQPEF (petroleum ether fraction); C, Colchicine induced increased C-metaphase frequency in root apical cells of *A. cepa* at 4 h continuous treatment.

Fig. 2: Graph showing the effect of AQPEF and Colchicine on A, metaphase and anaphase frequency; B, metaphase and C-metaphase frequency. Data represented as mean $\pm$ sem. ^a significant at $p < 0.0001$ and ^b significant at $p < 0.001$, ^c significant at $p < 0.05$ as compared to their respective control by 2x2 Contingency χ^2 -test with respective $df = 1$. Pearson Correlation and simple linear regression analysis of Colchicine ($150 \mu\text{g mL}^{-1}$) and AQPEF ($150 \mu\text{g mL}^{-1}$) between MI% (Mitotic Index%), META% (Metaphase%), ANA% (Anaphase%), and cM% (C-metaphase%) considering the change in their relative percentages at 2 h, 4 h and 4 h T+16 h RS. C, MI% and Metaphase%; D, MI% and Anaphase%; E, Metaphase% and cM%.

retardation depends on the antiproliferative potentials of the treated substances. Data indicate that AQPEF induced a concentration-dependent *A. cepa* root growth retardation effect. The highest root length retardation percentage ($68.51 \pm 0.56\%$, $p < 0.001$) in $150 \mu\text{g mL}^{-1}$ at 72 h of AQPEF treatment and IC_{50} value of the root growth

retardation was $23.68 \pm 5.5 \mu\text{g mL}^{-1}$ at 48 h treatment (Table 1). Recently, Barman *et al.* (2020) and Das *et al.* (2021) reported that plant extract induced root growth retardation and the levels of root growth retardation were increased with the increasing extract concentration and suppression of cell division in *A. cepa* root tips (Murthy *et al.* 2011, Ray *et al.* 2013). For the determination of

antiproliferative and cytotoxic potentials of synthetic or biochemicals including plant extracts, the *A. cepa* test model is widely used (Frescura *et al.* 2013, Khanna *et al.* 2013).

The mitotic index percentage in root apical meristem cells varies with the growing periods and the used AQPEF concentrations. Here, the highest MI (13.25 ± 0.72 %) was scored at 4 h of AQPEF ($150 \mu\text{g mL}^{-1}$) treatment and the lowest MI (3.55 ± 1.10 %) was found in the case of colchicine ($150 \mu\text{g mL}^{-1}$) at 4 h followed by 16 h recovery samples. At 2 h, MI% values were gradually increased- 5.67 ± 0.30 , 6.63 ± 0.41 , 7.58 ± 0.5 , 9.76 ± 0.26 , and 10.51 ± 0.80 % respectively for 0, 50, 100, 150, and $200 \mu\text{g mL}^{-1}$. The MI% also increased in the case of 4 h treated samples - 7.09 ± 0.62 , 10.3 ± 0.56 , 10.3 ± 0.29 , 13.25 ± 0.72 , and 9.95 ± 0.80 % respectively for 0, 50, 100, 150, and $200 \mu\text{g mL}^{-1}$ of AQPEF. In the case of 16 h recovery samples (4 h T + 16 h RS), MI% values were 10.06 ± 0.92 , 7.00 ± 0.29 , 8.45 ± 1.64 , 6.51 ± 0.77 , and 7.43 ± 0.38 % respectively for 0, 50, 100, 150 and $200 \mu\text{g mL}^{-1}$ of AQPEF. In the case of $150 \mu\text{g mL}^{-1}$ colchicine treatment, data indicate the MI% decreasing tendencies at 2 h (12.55 ± 1.44 %), 4 h (9.17 ± 0.56 %) and 4 h followed by 16 h recovery samples (3.55 ± 1.10 %). Plant extracts having mitostatic effects can be regarded as cytogenotoxic substance (Frescura *et al.* 2013, Khanna *et al.* 2013) and the reduction in the mitotic index is considered as an important parameter to examine the antimitotic as well as cytotoxic effects of biochemicals. The reduction in MI may be the result of their cytotoxic effects like blockage during DNA synthesis or G₂ phase or insufficient synthesis of ATP during spindle fiber formation and elongation (Sreeranjini and Siril 2011).

The investigation by squash preparation of *A. cepa* root apical meristem cells revealed that AQPEF induced increased metaphase frequencies and decreased frequencies of prophase, anaphase, and telophase in a concentration-dependent manner at both 2 and 4 h treated samples indicate that AQPEF has definite metaphase arresting activity. Subsequently, in recovery root tip cells (4 h T + 16 h RS), the frequencies of prophase, anaphase, and telophase showed an increasing tendency as compared to 2 and 4 h treated cells (Table 2). The highest metaphase cells percentage (81.44 ± 1.61 %) was scored from 4 h AQPEF treated ($100 \mu\text{g mL}^{-1}$) samples and that was followed by the concentration of $150 \mu\text{g mL}^{-1}$ (78.48 ± 2.41 %). However, in the case of 16 h recovery samples, the metaphase frequency showed a reducing tendency, and the metaphase frequencies were scored as 65.69 ± 3.87 % and 52.42 ± 6.02 % respectively for 100 and $150 \mu\text{g mL}^{-1}$ of AQPEF. Similarly. Colchicine treatment for 2 and 4 h notably induced increased metaphase frequency, 79.75 ± 4.46 %, and 84.88 ± 1.26 % respectively. In the case of 16 h recovery after 4 h colchicine treatment, the metaphase frequency (38.81 ± 1.66 %) was comparable to untreated control (Table 2). The mitotic index elevation in 2 and 4 h treated

cells may be due to an increase in metaphase frequencies. A similar colchicine-induced change in mitotic index and metaphase frequencies was observed earlier (Davidson *et al.* 1966).

Data indicate that the C-metaphase is the most frequent type of abnormalities induced by AQPEF in *A. cepa* root apical meristem cells. The AQPEF and colchicine treatments for 2 and 4 h induced a significant ($p < 0.001$ & 0.0001) increase in the frequency of C-metaphase in *A. cepa* root apical meristem cells in a concentration-dependent manner and decreased in recovery treatments (Table 2, Fig. 1, 2). In the case of 2 h treatment, $150 \mu\text{g mL}^{-1}$ of AQPEF showed the highest C-metaphase frequency (55.62%) but in the case of 4 h treatment, $100 \mu\text{g mL}^{-1}$ of AQPEF induced the highest (80.08 %) C-metaphase. Similarly, C-metaphase frequency was also increased in colchicine ($150 \mu\text{g mL}^{-1}$) treated samples as 77.95 ± 5.63 % and 83.00 ± 1.93 % respectively for 2 and 4 h. In the case of 16 h recovery samples (4 h T + 16 h RS), c-metaphase frequency (9.96 ± 3.40 %) was decreased. The formation of C-metaphase is directly correlated with the microtubule disruption and indicates that phytochemicals present in AQPEF may have colchicine-like microtubule destabilizing activity (Fiskesjö 1985, Bonciu *et al.* 2018, Shahin and El-Amoodi 1991). These data also correlate with the occurrence of C-metaphase in LAECV and colchicine-treated *A. cepa* root tip cells (Kundu and Ray 2016). Recently, Barman *et al.* (2021) reported colchicine like *C. inermis* leaf aqueous extract, LAECI, induced haphazardly arranged condensed chromosomes, C-metaphase, in *A. cepa*. Correlation between the formation of C-metaphase, vagrant chromosome, and polyploidy was evident by many investigators in *A. cepa* root tip cells (Carvalho *et al.* 2019). Based on this correlative evidence, we can postulate that AQPEF has a similar effect to LAECV on *A. cepa* root tip cells.

From Pearson correlation and simple linear regression analysis of colchicine ($150 \mu\text{g mL}^{-1}$) and AQPEF ($150 \mu\text{g mL}^{-1}$) between MI% (Mitotic Index%), META% (Metaphase%), ANA% (Anaphase%), and cM% (C-metaphase%) considering the change in their relative percentages at 2, 4, and 4h T+16h RS, a positive correlation was drawn between MI % and META % whereas a negative correlation was found between MI % and ANA % both in colchicine ($r = 0.885$ in META% and $r = -0.849$ in ANA%) and AQPEF ($r = 0.998$ in META% and $r = -0.992$ in ANA%). Positive correlation was also drawn between META% and cM% in AQPEF ($r = 0.983$) and colchicine (0.999). The above observations indicate that an increase in metaphase frequency was correlated with the decrease in anaphase frequency of both colchicine and AQPEF treatment. It can be extrapolated as the metaphase arresting nature of colchicine and AQPEF, leads to an increase in metaphase frequency and a simultaneous decrease in anaphase frequency, a possible mechanism for the cell cycle delay. A positive correlation between META% and

cM% indicate that both compounds possess metaphase arresting activity; therefore, AQPEF may have colchicine-like microtubule destabilizing attributes (Fig.2). The LAECV and its petroleum ether fraction, AQPEF, has colchicine-like metaphase arrest and mitotic abnormalities inducing potentials (Ray *et al.* 2012, Kundu and Ray 2016, Roy *et al.* 2021) and this study indicates AQPEF has cell cycle delay, metaphase arrest, c-metaphase inducing potentials in *A. cepa* root apical meristem cells. A further detailed investigation is required for purification of active principles of *C. viscosum* and to test their relative cell cycle delay, pro-metaphase arrest, and mitotic abnormality-inducing potentials.

Disclosure statement

No conflict of interest was declared.

Acknowledgments

The authors acknowledge Prof. A. Mukherjee for plant species authentication and the financial support of UGC-SRF (FC(Sc)/RS/UGC/ZOO/2018-19/129, w.e.f. 07.04.2018, dated: 04.02.2019), and the DST-PURSE, DST-FIST, and UGC-DRS-MRP-sponsored facilities in the Department of Zoology.

References

- Arvind, K., Pradeep, R., Deepa, R. and Mohan, V. 2002. Diabetes and coronary artery diseases. *Indian J. Med. Res.* **116**: 163-176.
- Bakare, A.A., Adeyemi, A. O., Adeyemi, A., Alabi, O. A. and Osibanjo, O. 2012. Cytogenotoxic effects of electronic waste leachate in *Allium cepa*. *Caryologia*. **65**: 94-100.
- Bakare, A. A., Alabi, O. A., Gbadebo, A. M., Ogunsuyi, O. I. and Alimba, C. G. 2013. In vivo cytogenotoxicity and oxidative stress induced by electronic waste leachate and contaminated well water. *Challenges*. **4**:169–187.
- Bakare, A. A., Mosuro, A. A. and Osibanjo, O. 2000. Effect of simulated leachate on chromosomes and mitosis in roots of *Allium cepa* (L). *J. Environ. Biol.* **21**: 263–271.
- Barman, M., Roy, S. and Ray, S. 2020. Colchicine like metaphase and cell cycle delay inducing effects of leaf aqueous extract of *Clerodendrum inerme* L. Gaertn. in *Allium cepa* root apical meristem cells. *Cytologia*. **85**: 197-201.
- Barman, M., Roy, S. and Ray, S. 2020. Mitotic abnormality inducing effects of leaf aqueous extract of *Clerodendrum inerme* Gaertn. on *Allium cepa* root apical meristem cells. *Cytologia*. **86**: 113–118.
- Bhattacharjee, D., Das, A., Das, S.K. and Chakraborty, G.S. 2011. *Clerodendrum infortunatum* L.: A review. *J. Adv. Pharm. Healthcare Res.* **1**: 82-5.
- Bonciu, E., Firbas, P., Fontanetti, C. S., Wusheng, J., Karaismailoğlu, M. C., Liu, D., Menicucci, F., Pesnya, D. S., Popescu, A., Romanovsky, A. V. and Schiff, S. 2018. An evaluation for the standardization of the *Allium cepa* test as cytotoxicity and genotoxicity assay. *Caryologia*. **71**: 191-209.
- Caperta, A. D., Delgado, M., Ressurreicao, F., Meister, A., Jones, R. N., Viegas, W. and Houben, A. 2006. Colchicine-induced polyploidization depends on tubulin polymerization in c-metaphase cells. *Protoplasma*. **227**: 147-153.
- Carvalho, M. S., Andrade-Vieira, L. F., dos Santos, F. E., Correa, F. F., das Graças Cardoso, M. and Vilela, L. R. 2019. Allelopathic potential and phytochemical screening of ethanolic extracts from five species of *Amaranthus* spp. in the plant model *Lactuca sativa*. *Sci. Hortic.* **9**: 90-8.
- Chaudhuri, A. and Ray, S. 2015. Antiproliferative activity of phytochemicals present in aerial parts aqueous extract of *Ampelocissus latifolia* (Roxb.) Planch. on apical meristem cells. *Int. J. Pharm. Bio. Sci.* **6**: 99-108.
- Das, R., Goswami, S., Barman, A., Barman, M. and Ray, S. 2021. Antimitotic and cytotoxic effects of leaf aqueous extract of *Scadoxus multiflorus* (Martyn) Raf. in *Allium cepa* root apical meristem cells. *Cytologia*. **86**:
- Das, S., Bhattacharya, S., Prasanna, A., Kumar, RBS., Pramanik, G. and Halder, PK. 2011. Preclinical evaluation of antihyperglycemic activity of *Clerodendrum infortunatum* leaf against streptozotocin-induced diabetic rats. *Diabetes ther.* **2**: 92-100.
- Davidson, D., MacLeod, R. and O'Riordan, M. 1966. Changes in mitotic index induced by colchicine. *Nature*. **212**: 1541–1542.
- Devi, O. I., Dutta, B. K. and Choudhury, P. 2013. Allelopathy effect of aqueous extract of *Clerodendrum viscosum* Vent, *Ageratum conyzoides* and *Parthenium hysterophorus* on the seed germination and seedling vigour of chickpea seeds (*Cicer arietinum* L) in vitro. *J. Nat. Appl. Sci.* **5**: 37-40.
- Fiskesjö, G. 1985. The *Allium* tests as a standard in environmental monitoring. *Hereditas*. **102**: 99–112.
- Frescura, V. D., Laughinghouse, H. D. 4th. and Tedesco SB. 2012. Antiproliferative effect of the tree and medicinal species *Luehea divaricata* on the *Allium cepa* cell cycle. *Caryologia*. **65**: 27-33.
- Frescura, V. D., Kuhn, A. W., Laughinghouse, H. D. 4th., Paranhos, J. T. and Tedesco, S. B. 2013. Post-treatment with plant extracts used in Brazilian folk medicine caused a partial reversal of the antiproliferative effect of glyphosate in the *Allium cepa* test. *Biocell*. **37**: 23-28.

- Gouthamchandra, K., Mahmood, R. and Manjunatha H. 2010. Free radical scavenging, antioxidant enzymes and wound healing activities of leaves extracts from *Clerodendrum infortunatum* L. *Environ. Toxicol. Pharmacol.* **30**: 11-18.
- Gupta, R. and Singh, H. K. 2012. Nootropic potential of *Alternanthera sessilis* and *Clerodendrum infortunatum* leaves on mice. *Asian Pac. J. Trop. Dis.* **2**: 465-470.
- Hamilton, F. 1997. The Flora Homoeopathica. B Jain (New Delhi).
- Islam, Md. S., Moghal, Md. M.R., Ahamed, S.K., Ahmed, J. and Islam, Md. A. 2013. A study on cytotoxic and anthelmintic activities of crude extracts of leaves of *Clerodendrum viscosum*. *Int. Res. J. Pharm.* **4**: 99-102.
- Khanna, N. and Sharma, S. 2013. *Allium cepa* root chromosomal aberration assay: a review. *Indian J. Pharm. Biol. Res.* **1**: 105-119.
- Kirtikar, K.R. and Basu, B. D. 1991. Indian Medicinal Plants. III. 2nd ed. Bishen Singh and Mahendra Pal Singh. Dehradun.
- Kundu, L. M. and Ray, S. 2016. Mitotic abnormalities and micronuclei inducing potentials of colchicine and leaf aqueous extracts of *Clerodendrum viscosum* Vent. in *Allium cepa* root apical meristem cells. *Caryologia.* **70**: 7-14
- Levan, A. 1938. The effect of colchicine on root mitoses in *Allium*. *Hereditas.* **24**: 471-86.
- Lobo, R., Chandrshakar, K. S., Jaykumar, B. and Mamatha B. 2010. In vitro antimicrobial activity of *Clerodendrum viscosum* Vent. *Der Pharm. Lett.* **2**: 257-260.
- Modi, A. J., Khadabadi, S. S., Farooqui, I. A. and Ghorpade, D.S. 2010. Studies on antimicrobial activity of *Clerodendrum infortunatum*, *Argyrea nervosa* and *Vitex negundo*: A comparison. *Der Pharm. Lett.* **2**: 102-105.
- Muh, T., Waliullah, A., Yeasmin, A. M., Wahedul, I. M. and Parvez, H. 2014. Insecticidal and repellent activity of *Clerodendrum viscosum* Vent. (Verbenaceae) against *Tribolium castaneum* (Herbst) (Coleoptera: tenebrionoidea). *Acad. J. Entomol.* **7**: 63-69.
- Murthy, G.S., Francis, T.P., Singh, C.R., Nagendra, H.G. and Naik, C. 2011. An assay for screening anti-mitotic activity of herbal extracts. *Curr. Sci.* **100**: 1399-1404.
- Nadkarni, K. M. and Nadkarni, A. K. 2002. Indian Materia Medica. Bombay.
- Nandi, S. and Lyndem, L. M. 2015. *Clerodendrum viscosum*: traditional uses, pharmacological activities and phytochemical constituents. *Nat. Prod. Res.* **30**: 497-506.
- Qasem, J. R. and Foy, C. L. 2001. Weed allelopathy, its ecological impacts and future prospects. *J. Crop. Prod.* **4**: 43-119.
- Ray, S., Kundu, L. M., Goswami, S. and Chakrabarti, C. S. 2012. Antiproliferative and apoptosis inducing activity of allelochemicals present in leaf aqueous extract of traditionally used antitumor medicinal plant, *Clerodendrum viscosum* Vent. *Int. J. Pharm. Res. Dev.* **4**: 332-345.
- Ray, S., Kundu, L. M., Goswami, S., Roy, G. C., Chatterjee, S., Dutta, S., Chaudhuri, A. and Chakrabarti, C. S. 2013. Metaphase arrest and delay in cell cycle kinetics of root apical meristems and mouse bone marrow cells treated with leaf aqueous extract of *Clerodendrum viscosum* Vent. *Cell Prolif.* **46**: 109-117.
- Roy, G. C. and Ray, S. 2017. Antiproliferative and apoptosis inducing activities of Leaf organic solvent extract fractions of *Clerodendrum viscosum* Vent. *Int. J. Pharma. Bio Sci.* **8**: 58-66.
- Roy, S., Kundu, L.M., Roy, G.C., Barman, M. and Ray S. 2021. Cytotoxic and micronuclei inducing effects of petroleum ether fraction of leaf aqueous extract of *Clerodendrum viscosum* Vent. in *Allium cepa* root tip cells. *Cytologia.* **86**: 1-6.
- Salmon, E. D., Mickseel, M. and Hays, T. 1984. Rapid rate of tubulin dissociation from microtubule in the mitotic spindle *in vivo* measured by blocking polymerization with colchicine. *JCB.* **99**: 1066-1075.
- Sannigrahi, S., Mazumder, U. K., Pal, D. K. and Mishra, S. L. 2009. Hepatoprotective potential of methanol extract of *Clerodendrum infortunatum* Linn. against CCl₄ induced hepatotoxicity in rats. *Pharmacogn. Mag.* **5**: 394-399.
- Shahin, S. A. and El-Amoodi, K. H. H. 1991. Induction of numerical chromosomal aberrations during DNA synthesis using the fungicides nimrod and rumigan-4 in root tips of *Vicia faba* L. *Mutat. Res.* **261**: 169-176.
- Sharma, A. K., and Sharma, A. 1999. Plant Chromosomes: Analysis, Manipulation and Engineering. *Hardwood Academic Publishers.*
- Singh, V. K., Ali, Z. A. and Siddiqui, M. K. 1997. Medicinal plants used by the forest ethnics of Gorakhpur district (Uttar Pradesh), India. *Int. J. Pharm.* **35**: 194-206.
- Singhmura, S. 2016. A comprehensive overview of a traditional medicinal herb: *Clerodendrum infortunatum* Linn. *J. pharm. sci.* **5**: 80-84.
- Sreeranjini, S. and Siril, E. A. 2011. Evaluation of anti-genotoxicity of the leaf extracts of *Morinda citrifolia* L. *Plant. Soil. Environ.* **57**: 222-227.