1 Viral spillover risk in High Arctic increases with 2 melting glaciers

3 **Audrée Lemieux¹, Graham A. Colby¹, Alexandre J. Poulain¹, Stéphane**
4 **Aris-Brosou^{1,2,✉}**

5 ¹Department of Biology, University of Ottawa, Ottawa, ON K1N 6N5, Canada

6 ²Department of Mathematics and Statistics, University of Ottawa, Ottawa, ON K1N
7 6N5, Canada

8 ✉e-mail: sarisbro@uottawa.ca

9 Abstract

10 While many viruses have a single natural host, host restriction can be incomplete, hereby
 11 leading to spillovers to other host species, potentially causing significant diseases as it is
 12 the case with the Influenza A, Ebola, or the SARS-CoV-2 viruses. However, such spillover
 13 risks are difficult to quantify. As climate change is rapidly transforming environments,
 14 it is becoming critical to quantify the potential for spillovers, in an unbiased manner.
 15 For this, we resorted to a metagenomics approach, and focused on two environments in
 16 the High Arctic, soil and lake sediments from Lake Hazen. We used DNA and RNA
 17 sequencing to reconstruct the lake's virosphere and its range of eukaryotic hosts, and
 18 show that spillover risk is higher in lake sediments than in soil and increased with runoff
 19 from glacier melt - a proxy for climate change. Should climate change also shift species
 20 range of potential vectors northwards, the High Arctic could become fertile ground for
 21 emerging pandemics.

Viruses are ubiquitous and are often described as the most abundant replicators on Earth (Dávila-Ramos et al., 2019; Koonin et al., 2015; Zhang et al., 2018). In spite of having highly diverse genomes, viruses are not independent “organisms” or replicators (Dawkins, 1989), as they need to infect a host’s cell in order to replicate. These virus/host relationships seem relatively stable within superkingdoms, so that viruses can be classified as *archaeal*, *bacterial* (also known as *bacteriophages*), and *eukaryotic* (Malik et al., 2017; Nasir et al., 2014, 2017), but host changes are not uncommon below this rank. Indeed, many species causing significant diseases can infect more than one host, as it is the case with the Influenza A (Long et al., 2019), Ebola (Rewar & Mirdha, 2014), and SARS-CoV-2 (Zhou & Shi, 2021) viruses. While these three viruses have non-human wild animal reservoirs as natural hosts, others have a broader host range, or their reservoir is more challenging to identify. Such host restrictions (or alternatively spillover capacities or risks) are to date poorly defined pragmatically and hence, difficult to assess without resorting to expert opinion (Grange et al., 2021).

Numerous factors are thought to influence such a viral spillover risk. For instance, viral particles need to attach themselves to specific receptors on their host’s cell to invade it (Longdon et al., 2014; Maginnis, 2018; Parrish et al., 2008). The conservation of those receptors across multiple species allows these hosts to be more predisposed to becoming infected by the same virus (Parrish et al., 2008; Woolhouse et al., 2005). Indeed, from an evolutionary standpoint, viruses are more prone to infecting hosts that are phylogenetically close to their natural host (Longdon et al., 2014, 2018), potentially because it is easier for them to infect and colonize species that are genetically similar (Geoghegan et al., 2017). Alternatively, but not exclusively, high mutation rates might explain why RNA viruses change host more often than other viruses (Longdon et al., 2014), as most lack proofreading mechanisms (Parrish et al., 2008).

To further investigate the role of host relatedness in viral host switch events, multiple studies have focused on the cophylogenetic relationships between viruses and their hosts (Bellec et al., 2014; Bennett et al., 2020; Jackson & Charleston, 2004; Madinda et al., 2016; Olival et al., 2017). Conceptually, if both viruses and their hosts cospeciate, the topologies of their respective phylogenetic trees should be identical or *congruent*. On the other hand, the occurrence of spillovers would result in incongruent virus/host phylogenies.

To date only few studies have characterized the communities and genomes of viruses in extreme environments (Dávila-Ramos et al., 2019), and even fewer, if any, describe their spillover risk. The High Arctic is of special interest as it is particularly affected by climate change, warming faster than the rest of the world (Colby et al., 2020; Lehnherr et al., 2018; Ruuskanen et al., 2020). Warming climate and rapid transitions of the environment increase the risks of spillover events by varying the global distributions and dynamics of viruses, and their reservoirs and vectors (Parkinson et al., 2014; Waits et al., 2018), as shown for arboviruses (Ciota & Keyel, 2019) and the Hendra virus (Martin et al., 2018). Furthermore, as the climate changes, the metabolic activity of the Arctic's microbiosphere also shifts, which in turns affects numerous ecosystem processes such as the emergence of new pathogens (Messan et al., 2020). It has now become critical to quantify the risk of these spillovers.

To address these outstanding questions, we resorted to a combination of metagenomics and of cophylogenetic modeling to quantify viral spillover risk by sampling in an unbiased manner both the virosphere and its range of hosts (Zhang et al., 2018). Focusing on eukaryotic hosts, which are critically affected by viral spillovers (Carrasco-Hernandez et al., 2017), we contrasted two local environments, lake sediments and soil samples of a High Arctic lake, Lake Hazen, to test how viral spillover risk is affected by glacier runoff, and hence potentially by global warming, which is expected to increase runoff with increasing

glacier melt at Lake Hazen (Colby et al., 2020; Lehnher et al., 2018). We show here that the risk increases with warming climate, but is likely to remain low in the absence of “bridge vectors.”

Results and Discussion

Plant and fungal viruses are overrepresented. Based on our most sensitive annotation pipeline (Supplementary Methods), viruses represented less than 1% of all contigs (See Supplementary Methods). RNA viruses of all kinds (*i.e.*, dsRNA, +ssRNA, and -ssRNA viruses) were found to be significantly more abundant than DNA viruses in all samples ($> 70.5\%$; binomial tests, $P < 2.48 \times 10^{-7}$; Figure 1, Table 1), with 7.3% to 36.5% of contigs *per* sampling site that had no annotations (binomial tests, $P < 3.82 \times 10^{-5}$; Table S2). This dominance of RNA viruses is not unexpected, as fungi biomass for instance surpasses that of bacteria in Arctic environments by 1-2 orders of magnitude (Schmidt & Bölter, 2002), and eukaryotes are known to be the main targets of RNA viruses (Koonin et al., 2015; Malik et al., 2017; Nasir et al., 2014, 2017). Our results are however difficult to compare with previous studies in the High Arctic, as those were based solely on DNA metagenomics sequencing (Aguirre de Cárcer et al., 2015; Gregory et al., 2019; Labbé et al., 2020), probably because RNA viruses are thought to be unstable (Yau & Seth-Pasricha, 2019), or due to inadequate sampling strategies to extract RNA viruses (Wang, 2020). Our results show nonetheless that it is possible to recover RNA viruses from RNA-seq metagenomics.

All viral types confounded, those infecting plants and/or fungi were overrepresented compared to those infecting animals and protists ($> 68.8\%$, binomial tests, $P < 2.48 \times 10^{-7}$; Table 1). This overrepresentation might reflect a preservation bias, due to the

constitutive defenses found in plants and fungi offered by their waxy epidermal cuticles and cell walls (Kozieł et al., 2021), even if most plant viruses lack a protective lipoprotein envelope as found in animal viruses (Stavolone & Lionetti, 2017). But irrespective of such a preservation bias, this imbalance could imply a high spillover potential among plants and fungi in the High Arctic for two reasons. First, RNA viruses are the most likely pathogens to switch hosts, due to their high rates of evolution (Aris-Brosou et al., 2019; Longdon et al., 2014). Second, plant biomass has been increasing over the past two decades in the High Arctic due to regional warming (Hudson & Henry, 2009), and is likely to keep doing as warming continues.

Spillover risk increases with glacier runoff. Given these viral and eukaryotic host representations, can spillover risk be assessed in these environments? To address this question, we resorted to the novel global-fit model Random TaPas. The stability of its results was assessed by running this algorithm three times, and by combining the results for the normalized Gini coefficients ($G^* \in [0, 1]$), a direct measure of spillover risk.

In all samples, when the runoff volume was negligible (the C sites; Figure 2a), spillover risk was high, with median G^* exceeding 0.675 and thus the 2/3 threshold, and was significantly higher in soil than in lake sediments for both GD and PACo (Dunn test, Benjamini-Hochberg [BH] correction, $P < 0.001$). However, in the presence of a low runoff volume (the L sites), spillover risk was higher in lake sediments than in soil for GD, but lower for PACo (Dunn test, BH correction, $P < 0.001$; Figure 2b). Finally, in the high runoff regime (the H sites), for both GD and PACo, spillover risk was higher in lake sediments than in soil (Dunn test, BH correction, $P < 0.001$; Figure 2c). Altogether, these results show that as runoff volume increases from almost non-existent to high, spillover risk shifted from higher in soil, to higher in lake sediments.

119 This pattern is consistent with the predictions of the Coevolution Effect hypothesis
 120 (Zohdy et al., 2019), and provides us with a mechanism explaining the observed increase
 121 in spillover risk with runoff. Lake Hazen was recently found to have undergone a dramatic
 122 change in sedimentation rates since 2007 compared to the previous 300 years: an increase
 123 in glacial runoff drives sediment delivery to the lake, leading to increased turbidity that
 124 perturbs anoxic bottom water known from the historical record (Lehnherr et al., 2018).
 125 Not only this, but turbidity also varies within the water column throughout the season
 126 (St. Pierre et al., 2019), hence fragmenting the lake habitat every year, and more so
 127 since 2007. This fragmentation of the aquatic habitat creates conditions that are, under
 128 the Coevolution Effect, favorable to spillover. Fragmentation creates barriers to gene
 129 flow, that increases genetic drift within finite populations, accelerating the coevolution of
 130 viruses and of their hosts. This acceleration leads to viral diversification which, should it
 131 be combined with “bridge vectors” (such as mosquitoes in terrestrial systems), increases
 132 spillover risk (Zohdy et al., 2019). Lake sediments are environmental archives; over time,
 133 they can preserve genetic material from aquatic organisms but also, and probably to a
 134 lesser extent, genetic material from its drainage basin. The co-evolutionary signal detected
 135 in lake sediments reflects interactions that may have happened in the fragmented aquatic
 136 habitat but also elsewhere in the drainage basin. Regardless of where the interaction
 137 occurred, our results show that spillover risk increases with runoff, a proxy of climate
 138 warming (Figure 2).

139 **Spillovers might already be happening.** To go one step further and identify the
 140 viruses most at risk of spillover, we focused on the model predictions made by Random
 141 TaPas. Under the null model, the occurrence of each virus/host association is evenly
 142 distributed on their cophylogeny (when sub-cophylogenies are drawn randomly, from a

uniform law). Departures from an even distribution are measured by the residuals of the linear fit. Positive residuals indicate a more frequent association than expected, that is pairs of host/virus species that contribute the most to the cophylogenetic signal. On the other hand, negative residuals indicate a less frequent association than expected, and hence pairs of host/virus species that contribute little to the cophylogenetic signal, because they tend to create incongruent phylogenies, a signature of spillover risk.

For both soil and lake sediments, the magnitude of the largest residuals tended either to decrease (Soil; Figure 3a) or to stay the same (Sediment; Figure 3b). This means that with increasing runoff, the strength of the cophylogenetic signal may remain steady, or may even weaken. On the other hand, the magnitude of the most negative residuals either remained globally unchanged (Soil; Figure 3a), or tended to become more negative (Sediment; Figure 3b). This latter pattern indicates that as runoff increases, the strength of the cophylogenetic signal deteriorates, potentially implying a higher spillover risk in lake sediments.

With this, Random TaPas can identify the viruses driving the spillover signal. Our results show that certain nucleocytoplasmic large DNA viruses (NCLDV), such as the cowpox, horsepox, monkeypox, and vaccinia viruses, infect more than one species when the runoff is high (H sites), including humans (Figure 4). This indicates that spillovers are already occurring with these viral species. Among those, giant viruses (*e.g.*, *Megavirus chiliensis* and *Acanthamoeba polyphaga mimivirus*) are known to infect a wide range of eukaryotes from protists to animals (Schulz et al., 2020), which could impact the eukaryotic communities in the High Arctic (Figure 4, S8). As plant viruses represented $\sim 44\%$ of viruses infecting more than one host, this indicates that plants could also be affected.

To our knowledge, this is the first attempt to assess the complete virosphere of both DNA and RNA viruses, and their spillover capacity in any given environments, leading us

168 to show that increased glacier runoff, a direct consequence of climate change, is expected
169 to increase viral spillover risk. However, as this is the first study applying the Random
170 TaPas algorithm, we do not have yet any comparators in order to gauge the efficacy of
171 G^* in assessing spillover capacity, both qualitatively and quantitatively.

172 Altogether, we provided here a proof of concept for a novel and unbiased approach to
173 assessing spillover risk. This is not the same as predicting spillovers or even pandemics,
174 because as long as “bridge vectors” (Zohdy et al., 2019) are not present in the environment,
175 the likelihood of dramatic events probably remains low. But as climate change leads to
176 shifts in species ranges and distributions, new interactions can emerge (Wallingford et al.,
177 2020), bringing in vectors that can mediate viral spillovers (Rocklöv & Dubrow, 2020).
178 This twofold effect of climate change, both increasing spillover risk and leading to a
179 northward shift in species ranges (Osland et al., 2021), could have dramatic effect in the
180 High Arctic. Disentangling this risk from actual spillovers and pandemics will be a critical
181 endeavor to pursue in parallel with surveillance activities, in order to mitigate the impact
182 of future major outbreaks.

183 Materials and Methods

184 **Data acquisition.** Between May 10 and June 10, 2017, sediments and soil cores were
185 collected from Lake Hazen (82°N, 71°W; Quttinirpaaq National Park, northern Ellesmere
186 Island, Nunavut, Canada), the largest High Arctic lake by volume in the world (Lehnherr
187 et al., 2018). Sampling took place as the lake was still completely covered in ice (Table S1),
188 as detailed in Colby et al. Colby et al. (2020). The sediment accumulation at the bottom
189 of the Lake is caused by both allochthonous and autochthonous processes. The former
190 are characterized by meltwaters that flow between late June and the end of August, and

run from the outlet glaciers along the northwestern shoreline through poorly consolidated river valleys, while the latter refer to the sedimentation process within the lake.

To contrast soil and sediment sites, core samples were paired, whenever possible, between these two environments. Soil samples were taken at three locations (Figure S2; C-Soil, L-Soil, and H-Soil) in the dried streambeds of the tributaries, on the northern shore, upstream of the lake and its sediments. The corresponding paired lake sediment samples were also cored at three locations, separated into hydrological regimes by seasonal runoff volume: negligible, low, and high runoff (Figure S2; C-Sed, L-Sed, and H-Sed). Specifically, the C (for *Control*) sites were both far from the direct influence of glacial inflows, while L sites were at a variable distance from Blister Creek, a small glacial inflow, and the H sites were located adjacent to several larger glacial inflows (Abbé River and Snow Goose). The water depth at L-Sed and H-Sed was respectively 50 m and 21 m, and the overlying water depth for site C-Sed was 50 m.

Before sample collection, all equipment was sterilized with 10% bleach and 90% ethanol, and non-powdered latex gloves were worn to minimize contamination. Three cores of ~ 30 cm length were sampled at each location, and the top 5 and 10 cm of each sediment and soil core, respectively, were then collected and homogenized for genetic analysis as detailed in Colby et al. Colby et al. (2020). DNA was extracted on each core using the DNeasy PowerSoil Pro Kit, and RNA with the RNeasy PowerSoil Total RNA Kit (MO BIO Laboratories Inc, Carlsbad, CA, USA), following the kit guidelines, except that the elution volume was 30 μ L. DNA and RNA were thereby extracted three times per sampling site, and elution volumes were combined for a total volume of 90 μ L instead of 100 μ L.

To sequence both DNA and RNA, a total of 12 metagenomic libraries were prepared ($n = 6$ for DNA, $n = 6$ for RNA), two for each sampling site, and run on an Illumina

216 HiSeq 2500 platform (Illumina, San Diego, CA, USA) at Génome Québec, using Illu-
217 mina’s TruSeq LT adapters (forward: AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC, and back-
218 ward: AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT) in a paired-end 125 bp configuration. Each
219 library was replicated ($n = 2$ for DNA, $n = 3$ for RNA) for each sample.

220 **Data preprocessing and taxonomic assignments.** A first quality assessment of the
221 raw sequencing data was made using FastQC v0.11.8 (Andrews et al., 2010). Trimmomatic
222 v0.36 (Bolger et al., 2014) was then employed to trim adapters and low-quality reads and
223 bases using the following parameters: `phred33, ILLUMINACLIP:adapters/TruSeq3-PE-2.`
224 `fa:3:26:10, LEADING:3, TRAILING:3, SLIDINGWINDOW:4:20, CROP:105, HEADCROP:15,`
225 `AVGQUAL:20, MINLEN:36`. A second round of quality check was performed with FastQC
226 to ensure that Illumina’s adapter sequences and unpaired reads were properly removed.
227 Reads assembly into contigs was done *de novo* with both SPAdes v3.13.1 (Prjibelski et
228 al., 2020) and metaSPAdes v3.13.1 (Nurk et al., 2017) for DNA, and with Trinity v2.9.0
229 (Grabherr et al., 2011), rnaSPAdes v3.13.1 (Bushmanova et al., 2019), and metaSPAdes
230 for RNA (See Supplementary text). In all cases, the pipelines were used with their default
231 settings.

232 Once assembled, a high-level (superkingdom) taxonomic assignment was determined
233 based on BLASTn v2.10.0 (Zhang et al., 2000) searches. Those were performed at a
234 stringent 10^{-19} *E*-value threshold against the partially non-redundant nucleotide (nr/nt)
235 database from NCBI v5 (NCBI, 1988) (`ftp.ncbi.nlm.nih.gov/blast/db/nt*tar.gz`; down-
236 loaded on June 17, 2020). The proportions of taxonomic annotations (“Archaea,” “Bacte-
237 ria,” “Eukaryota,” or “Viruses”) were calculated, and a 95% consensus was taken to assign
238 a superkingdom rank for each contig. When no such 95% consensus could be determined,
239 the contigs were classified as “Other.”

To refine the taxonomic assignment of “viruses,” GenBank’s viral nucleotide sequences v238.0 (Sayers et al., 2019) were retrieved (`ftp.ncbi.nlm.nih.gov/genbank/gbvrl*seq.gz`; downloaded on July 23, 2020), concatenated, converted into FASTA with `seqret` v6.6.0 (Rice et al., 2000), and used to create a local database for BLASTn alignments. For each sampling location, after combining the DNA and RNA contigs classified as viral in the previous step, BLASTn searches were again conducted at the same stringent 10^{-19} *E*-value threshold, and the accession numbers of all the High-scoring Segment Pairs (HSPs) were used to retrieve their corresponding taxonomy identifiers (IDs) and their full taxonomic lineages with the R package `taxonomizr` v0.5.3 (Sherrill-Mix, 2019). The viral contigs were also mapped with Bowtie2 v2.3.5.1 (Langmead & Salzberg, 2012) using default settings to compare BLASTn and Bowtie2 efficiencies in refining these taxonomic annotations. As searches were found to be more sensitive with BLASTn than with Bowtie2 (see Supplementary text), only BLASTn results are shown hereafter, as our goal was to find as many similar sequences as possible in more than one species to eventually infer the virosphere from the virome.

Eukaryotic contigs were processed as above, based off the nr/nt database. Results were filtered by keeping a maximum of 12 HSPs whose *E*-value $< 10^{-100}$ *per* contig, for which lineages were obtained.

All samples were filtered to remove non-eukaryotic and uncultured hosts as well as viral and eukaryotic sequences with no taxonomy information. The ViralZone (Hulo et al., 2011) and International Committee on Taxonomy of Viruses (ICTV) (Lefkowitz et al., 2018) databases were consulted to obtain host range information on each found viruses. These taxonomic assignments were then used to retrieve their phylogenetic placements according to the Tree of Life (ToL) (`tolweb.org`), hence generating two trees: one for viruses and one for eukaryotes. For this, we used the `classification` and `class2tree`

functions from the R package `taxize` v0.9.99 (S. Chamberlain et al., 2020; S. A. Chamberlain & Szocs, 2013). In each environment, vertices of the viral and eukaryotic trees were then put in relation with each other according to the Virus-Host DB (downloaded on Sep. 29, 2020) (Mihara et al., 2016). These relations were saved in a binary association matrix (0: no infection; 1: infection), one for each environment. To simplify downstream computations without losing any information, only eukaryotic hosts associated to at least one virus were kept in the non-viral tree.

Spillover quantification. To quantify viral spillovers based on the viral and eukaryotic hosts identified, we employed the Random Tanglegram Partitions algorithm (Random TaPas) (Balbuena et al., 2020). This algorithm computes the cophylogenetic signal or congruence between two phylogenetic trees, the viral and the host trees. When congruence is large, or “perfect,” the two trees are identical and hence, there is strong cophylogenetic signal – and absence of spillover. On the other hand, weak congruence is evidence for the existence of spillovers. Random TaPas quantifies congruence in two ways: a geodesic distance (GD) (Schardl et al., 2008), or a Procrustes distance (Procrustes Approach to Cophylogeny: PACo) (Balbuena et al., 2013), the latter measuring the distance between two trees geometrically transformed to make them as identical as possible. To partially account for phylogenetic non-independence when measuring congruence, Random TaPas further implements a resampling scheme where $N = 10^4$ subtrees of about 20% of the total number of virus/hosts links are randomly selected. This selection is used to generate a distribution of the empirical frequency of each association, measured by either GD or PACo.

Each empirical frequency is then regressed against a uniform distribution, and the residuals are used in two ways: (i) to quantify co-speciation, which is inversely propor-

289 tional to spillover risk; and (ii) to identify those virus/host pairs that contributed the least
 290 to the cophylogenetic signal, *i.e.*, the most to spillover risk. This risk is finally quantified
 291 by the shape of the distribution of residuals (for GD or PACo), with a normalized Gini
 292 coefficient (G^*) that takes its values between 0 (perfect congruence, no spillover) to 1
 293 (maximal spillover risk), with a defined threshold of 2/3 indicating a “large” value of G^* .
 294 To account for phylogenetic uncertainty, the process is repeated $n = 1,000$ times, each
 295 replicate being a random resolution of the multifurcating virus/host trees of life into a
 296 fully bifurcating tree.

297 **Data availability.** The raw data used in this study can be found at [www.ncbi.nlm.nih](http://www.ncbi.nlm.nih.gov/bioproject/556841)
 298 [.gov/bioproject/556841](http://www.ncbi.nlm.nih.gov/bioproject/556841) (DNA-Seq) and at www.ncbi.nlm.nih.gov/bioproject/PRJNA746497/
 299 (RNA-Seq). The code developed for this work is available from [github.com/sarisbro/](https://github.com/sarisbro/data)
 300 [data](https://github.com/sarisbro/data).

301 **Acknowledgements.** We thank Frances Pick for helpful comments on an early version
 302 of this work. This study was financially supported by the Natural Sciences Research
 303 Council of Canada (A.J.P., S.A.B.) and by the University of Ottawa (A.L., G.A.C.).

304 **Author contributions.** S.A.B. and A.J.P. designed research; G.A.C. collected and
 305 processed the samples; A.L. performed all analyses and wrote the original draft; A.L. and
 306 S.A.B. wrote the manuscript with contributions and suggestions from G.A.C. and A.J.P.;
 307 and S.A.B. and A.J.P. supervised this study and acquired funding.

308 **Competing interests**

309 The authors declare no competing interests.

References

- Aguirre de Cárcer, D., López-Bueno, A., Pearce, D. A., & Alcamí, A. (2015). Biodiversity and distribution of polar freshwater DNA viruses. *Sci. Adv.*, 1(5). doi: 10.1126/sciadv.1400127
- Andrews, S., Krueger, F., Segonds-Pichon, A., Biggins, L., Krueger, C., & Wingett, S. (2010). *FastQC*. Retrieved from <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Aris-Brosou, S., Parent, L., & Ibeh, N. (2019). Viral long-term evolutionary strategies favor stability over proliferation. *Viruses*, 11(8). doi: 10.3390/v11080677
- Balbuena, J. A., Míguez-Lozano, R., & Blasco-Costa, I. (2013). PACo: a novel procrustes application to cophylogenetic analysis. *PLoS ONE*, 8(4), e61048. doi: 10.1371/journal.pone.0061048
- Balbuena, J. A., Pérez-Escobar, Ó. A., Llopis-Belenguer, C., & Blasco-Costa, I. (2020). Random tanglegram partitions (Random TaPas): an alexandrian approach to the cophylogenetic Gordian knot. *Syst. Bio.*, 69(6), 1212–1230. doi: 10.1093/sysbio/syaa033
- Bellec, L., Clerissi, C., Edern, R., Foulon, E., Simon, N., Grimsley, N., & Desdevises, Y. (2014). Cophylogenetic interactions between marine viruses and eukaryotic picophytoplankton. *BMC Evol. Biol.*, 14(1), 59. doi: 10.1186/1471-2148-14-59
- Bennett, A. J., Paskey, A. C., Kuhn, J. H., Bishop-Lilly, K. A., & Goldberg, T. L. (2020). Diversity, transmission, and cophylogeny of Ledanteviruses (*Rhabdoviridae*: *Ledantevirus*) and Nycteribiid bat flies parasitizing Angolan soft-furred fruit bats in Bundibugyo District, Uganda. *Microorganisms*, 8(5), 750. doi: 10.3390/microorganisms8050750
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), 2114–2120. doi: 10.1093/bioinformatics/btu170
- Bushmanova, E., Antipov, D., Lapidus, A., & Prjibelski, A. D. (2019). rnaSPAdes: a de novo transcriptome assembler and its application to RNA-Seq data. *GigaScience*, 8(9), 1–13. doi: 10.1093/gigascience/giz100
- Carrasco-Hernandez, R., Jácome, R., López Vidal, Y., & Ponce de León, S. (2017). Are RNA viruses candidate agents for the next global pandemic? A review. *ILAR J.*, 58(3), 343–358. doi: 10.1093/ilar/ilx026

- 342 Chamberlain, S., Szocs, E., Foster, Z., Arendsee, Z., Boettiger, C., Ram, K., ... Grenié,
343 M. (2020). *taxize: Taxonomic information from around the web*. Retrieved from
344 <https://github.com/ropensci/taxize>
- 345 Chamberlain, S. A., & Szocs, E. (2013). *taxize: taxonomic search and retrieval in R*.
346 *F1000Research*, 2(191). doi: 10.12688/f1000research.2-191.v2
- 347 Ciota, A. T., & Keyel, A. C. (2019). The role of temperature in transmission of zoonotic
348 arboviruses. *Viruses*, 11(11). doi: 10.3390/v11111013
- 349 Colby, G. A., Ruuskanen, M. O., St.Pierre, K. A., St.Louis, V. L., Poulain, A. J., & Aris-
350 Brosou, S. (2020). Warming climate is reducing the diversity of dominant microbes
351 in the largest High Arctic lake. *Front. Microbiol.*, 11, 2316. doi: 10.3389/fmicb.2020
352 .561194
- 353 Dávila-Ramos, S., Castelán-Sánchez, H. G., Martínez-ávila, L., Sánchez-Carbente,
354 M. D. R., Peralta, R., Hernández-Mendoza, A., ... Batista-García, R. A. (2019).
355 A review on viral metagenomics in extreme environments. *Front. Microbiol.*, 10, 2403.
356 doi: 10.3389/fmicb.2019.02403
- 357 Dawkins, R. (1989). *The selfish gene*. Oxford University Press.
- 358 Geoghegan, J. L., Duchêne, S., & Holmes, E. C. (2017). Comparative analysis estimates
359 the relative frequencies of co-divergence and cross-species transmission within viral
360 families. *PLoS Pathog.*, 13(2), 1–17. doi: 10.1371/journal.ppat.1006215
- 361 Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., ...
362 Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a
363 reference genome. *Nat. Biotechnol.*, 29(7), 644–652. doi: 10.1038/nbt.1883
- 364 Grange, Z. L., Goldstein, T., Johnson, C. K., Anthony, S., Gilardi, K., Daszak, P., ...
365 Mazet, J. A. K. (2021). Ranking the risk of animal-to-human spillover for newly
366 discovered viruses. *P. Natl. Acad. Sci.*, 118(15). doi: 10.1073/pnas.2002324118
- 367 Gregory, A. C., Zayed, A. A., Conceição-Neto, N., Temperton, B., Bolduc, B., Alberti,
368 A., ... Sullivan, M. B. (2019). Marine DNA viral macro- and microdiversity from pole
369 to pole. *Cell*, 177(5), 1109–1123.e14. doi: 10.1016/j.cell.2019.03.040
- 370 Hudson, J. M. G., & Henry, G. H. R. (2009). Increased plant biomass in a High Arctic
371 heath community from 1981 to 2008. *Ecology*, 90(10), 2657–2663. doi: 10.1890/
372 09-0102.1
- 373 Hulo, C., de Castro, E., Masson, P., Bougueleret, L., Bairoch, A., Xenarios, I., & Le
374 Mercier, P. (2011). ViralZone: a knowledge resource to understand virus diversity.
375 *Nucleic Acids Res.*, 39(Database issue), D576–D582. doi: 10.1093/nar/gkq901

- 376 Jackson, A. P., & Charleston, M. A. (2004). A cophylogenetic perspective of RNA–virus
377 evolution. *Mol. Biol. Evol.*, *21*(1), 45–57. doi: 10.1093/molbev/msg232
- 378 Koonin, E. V., Dolja, V. V., & Krupovic, M. (2015). Origins and evolution of viruses
379 of eukaryotes: the ultimate modularity. *Virology*, *479–480*, 2–25. doi: 10.1016/j.virol
380 .2015.02.039
- 381 Koziel, E., Otulak-Koziel, K., & Bujarski, J. J. (2021). Plant cell wall as a key player
382 during resistant and susceptible plant-virus interactions. *Front. Microbiol.*, *12*, 495.
383 doi: 10.3389/fmicb.2021.656809
- 384 Labbé, M., Girard, C., Vincent, W. F., & Culley, A. I. (2020). Extreme viral partitioning
385 in a marine-derived High Arctic lake. *mSphere*, *5*(3), e00334–20. doi: 10.1128/mSphere
386 .00334-20
- 387 Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat.*
388 *Methods*, *9*, 357–359. doi: 10.1038/nmeth.1923
- 389 Lefkowitz, E. J., Dempsey, D. M., Hendrickson, R. C., Orton, R. J., Siddell, S. G., &
390 Smith, D. B. (2018). Virus taxonomy: the database of the International Committee on
391 Taxonomy of Viruses (ICTV). *Nucleic Acids Res.*, *46*(D1), D708–D717. doi: 10.1093/
392 nar/gkx932
- 393 Lehnherr, I., St. Louis, V. L., Sharp, M., Gardner, A. S., Smol, J. P., Schiff, S. L., ...
394 Talbot, C. H. (2018). The world’s largest High Arctic lake responds rapidly to climate
395 warming. *Nat. Commun.*, *9*(1), 1290. doi: 10.1038/s41467-018-03685-z
- 396 Long, J. S., Mistry, B., Haslam, S. M., & Barclay, W. S. (2019). Host and viral determi-
397 nants of Influenza A virus species specificity. *Nat. Rev. Microbiol.*, *17*(2), 67–81. doi:
398 10.1038/s41579-018-0115-z
- 399 Longdon, B., Brockhurst, M. A., Russell, C. A., Welch, J. J., & Jiggins, F. M. (2014). The
400 evolution and genetics of virus host shifts. *PLoS Pathog.*, *10*(11), 1–8. doi: 10.1371/
401 journal.ppat.1004395
- 402 Longdon, B., Day, J. P., Alves, J. M., Smith, S. C. L., Houslay, T. M., McGonigle, J. E., ...
403 Jiggins, F. M. (2018). Host shifts result in parallel genetic changes when viruses evolve in
404 closely related species. *PLoS Pathog.*, *14*(4), 1–14. doi: 10.1371/journal.ppat.1006951
- 405 Madinda, N. F., Ehlers, B., Wertheim, J. O., Akoua-Koffi, C., Bergl, R. A., Boesch, C.,
406 ... Calvignac-Spencer, S. (2016). Assessing host-virus codivergence for close relatives
407 of Merkel cell Polyomavirus infecting African great apes. *J. Virol.*, *90*(19), 8531–8541.
408 doi: 10.1128/JVI.00247-16

- 409 Maginnis, M. S. (2018). Virus–receptor interactions: the key to cellular invasion. *J. Mol.*
410 *Biol.*, *430*(17), 2590–2611. doi: 10.1016/j.jmb.2018.06.024
- 411 Malik, S. S., Azem-e Zahra, S., Kim, K. M., Caetano-Anollés, G., & Nasir, A. (2017). Do
412 viruses exchange genes across superkingdoms of life? *Front. Microbiol.*, *8*, 2110. doi:
413 10.3389/fmicb.2017.02110
- 414 Martin, G., Yanez-Arenas, C., Chen, C., Plowright, R. K., Webb, R. J., & Skerratt, L. F.
415 (2018). Climate change could increase the geographic extent of Hendra Virus spillover
416 risk. *Ecohealth*, *15*(3), 509–525. doi: 10.1007/s10393-018-1322-9
- 417 Messan, K. S., Jones, R. M., Doherty, S. J., Foley, K., Douglas, T. A., & Barbato, R. A.
418 (2020). The role of changing temperature in microbial metabolic processes during
419 permafrost thaw. *PLoS ONE*, *15*(4), e0232169. doi: 10.1371/journal.pone.0232169
- 420 Mihara, T., Nishimura, Y., Shimizu, Y., Nishiyama, H., Yoshikawa, G., Uehara, H., ...
421 Ogata, H. (2016). Linking virus genomes with host taxonomy. *Viruses*, *8*(3), 66. doi:
422 10.3390/v8030066
- 423 Nasir, A., Forterre, P., Kim, K. M., & Caetano-Anollés, G. (2014). The distribution and
424 impact of viral lineages in domains of life. *Front. Microbiol.*, *5*, 194. doi: 10.3389/
425 fmicb.2014.00194
- 426 Nasir, A., Kim, K. M., & Caetano-Anollés, G. (2017). Long-term evolution of viruses: a
427 Janus-faced balance. *BioEssays*, *39*(8), 1700026. doi: 10.1002/bies.201700026
- 428 Nucleotide [Internet]. Bethesda (MD): National Library of Medicine (US), National Center
429 for Biotechnology Information; [1988] - [cited 2021 Jan 20]. Available from [https://](https://www.ncbi.nlm.nih.gov/nucleotide/)
430 www.ncbi.nlm.nih.gov/nucleotide/. (n.d.).
- 431 Nurk, S., Meleshko, D., Korobeynikov, A., & Pevzner, P. A. (2017). metaSPAdes: a
432 new versatile metagenomic assembler. *Genome Res.*, *27*(5), 824-834. doi: 10.1101/
433 gr.213959.116
- 434 Olival, K. J., Hosseini, P. R., Zambrana-Torrel, C., Ross, N., Bogich, T. L., & Daszak,
435 P. (2017). Host and viral traits predict zoonotic spillover from mammals. *Nature*,
436 *546*(7660), 646–650. doi: 10.1038/nature22975
- 437 Osland, M. J., Stevens, P. W., Lamont, M. M., Brusca, R. C., Hart, K. M., Waddle, J. H.,
438 ... Seminoff, J. A. (2021). Tropicalization of temperate ecosystems in north america:
439 The northward range expansion of tropical organisms in response to warming winter
440 temperatures. *Glob. Change Biol.*, *27*(13), 3009–3034. doi: 10.1111/gcb.15563

- 441 Parkinson, A. J., Evengard, B., Semenza, J. C., Ogden, N., Børresen, M. L., Berner, J., ...
442 Albiñ, A. (2014). Climate change and infectious diseases in the Arctic: establishment
443 of a circumpolar working group. *Int. J. Circumpol. Heal.*, 73, 25163. doi: 10.3402/
444 ijch.v73.25163
- 445 Parrish, C. R., Holmes, E. C., Morens, D. M., Park, E.-C., Burke, D. S., Calisher, C. H., ...
446 Daszak, P. (2008). Cross-species virus transmission and the emergence of new epidemic
447 diseases. *Microbiol. Mol. Biol. R.*, 72(3), 457–470. doi: 10.1128/MMBR.00004-08
- 448 Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., & Korobeynikov, A. (2020). Using
449 spades de novo assembler. *Curr. Protoc. Bioinformatics*. doi: 10.1002/cpbi.102
- 450 Rewar, S., & Mirdha, D. (2014). Transmission of Ebola virus disease: an overview. *Ann.*
451 *Glob. Health*, 80(6), 444–451. doi: 10.1016/j.aogh.2015.02.005
- 452 Rice, P., Longden, L., & Bleasby, A. (2000). EMBOSS: The european molecular biology
453 open software suite. *Trends Genet.*, 16(6), 276–277. doi: 10.1016/S0168-9525(00)02024
454 -2
- 455 Rocklöv, J., & Dubrow, R. (2020). Climate change: an enduring challenge for vector-
456 borne disease prevention and control. *Nat. Immunol.*, 21(5), 479–483. doi: 10.1038/
457 s41590-020-0648-y
- 458 Ruuskanen, M. O., Colby, G., St.Pierre, K. A., St.Louis, V. L., Aris-Brosou, S., & Poulain,
459 A. J. (2020). Microbial genomes retrieved from High Arctic lake sediments encode for
460 adaptation to cold and oligotrophic environments. *Limnol. Oceanogr.*, 65(S1), S233–
461 S247. doi: 10.1002/lno.11334
- 462 Sayers, E. W., Cavanaugh, M., Clark, K., Ostell, J., Pruitt, K. D., & Karsch-Mizrachi, I.
463 (2019). GenBank. *Nucleic Acids Res.*, 47(D1), D94–D99. doi: 10.1093/nar/gky989
- 464 Schardl, C. L., Craven, K. D., Speakman, S., Stromberg, A., Lindstrom, A., & Yoshida,
465 R. (2008). A novel test for host-symbiont codivergence indicates ancient origin of fungal
466 endophytes in grasses. *Syst. Bio.*, 57(3), 483–498. doi: 10.1080/10635150802172184
- 467 Schmidt, N., & Bölter, M. (2002). Fungal and bacterial biomass in tundra soils along an
468 arctic transect from Taimyr Peninsula, central Siberia. *Polar Biol.*, 25(12), 871–877.
469 doi: 10.1007/s00300-002-0422-7
- 470 Schulz, F., Roux, S., Paez-Espino, D., Jungbluth, S., Walsh, D. A., Denef, V. J., ...
471 Woyke, T. (2020). Giant virus diversity and host interactions through global metage-
472 nomics. *Nature*, 578(7795), 432–436. doi: 10.1038/s41586-020-1957-x
- 473 Sherrill-Mix, S. (2019). *taxonomizr: functions to work with NCBI accessions and taxon-*
474 *omy*. Retrieved from <https://cran.r-project.org/package=taxonomizr>

- 475 Stavolone, L., & Lionetti, V. (2017). Extracellular matrix in plants and animals: Hooks
476 and locks for viruses. *Front. Microbiol.*, 8, 1760. doi: 10.3389/fmicb.2017.01760
- 477 St. Pierre, K. A., St. Louis, V. L., Lehnher, I., Schiff, S. L., Muir, D. C. G., Poulain,
478 A. J., ... Gardner, A. S. (2019). Contemporary limnology of the rapidly changing
479 glacierized watershed of the world's largest High Arctic lake. *Sci. Rep.*, 9(1), 4447. doi:
480 10.1038/s41598-019-39918-4
- 481 Waits, A., Emelyanova, A., Oksanen, A., Abass, K., & Rautio, A. (2018). Human
482 infectious diseases and the changing climate in the Arctic. *Environ. Int.*, 121, 703–713.
483 doi: 10.1016/j.envint.2018.09.042
- 484 Wallingford, P. D., Morelli, T. L., Allen, J. M., Beaur, E. M., Blumenthal, D. M.,
485 Bradley, B. A., ... Sorte, C. J. B. (2020). Adjusting the lens of invasion biology to
486 focus on the impacts of climate-driven range shifts. *Nat. Clim. Change*, 10(5), 398–405.
487 doi: 10.1038/s41558-020-0768-2
- 488 Wang, D. (2020). 5 challenges in understanding the role of the virome in health and
489 disease. *PLoS Pathog.*, 16(3), 1–6. doi: 10.1371/journal.ppat.1008318
- 490 Woolhouse, M. E. J., Haydon, D. T., & Antia, R. (2005). Emerging pathogens: the
491 epidemiology and evolution of species jumps. *Trends Ecol. Evol.*, 20(5), 238–244. doi:
492 10.1016/j.tree.2005.02.009
- 493 Yau, S., & Seth-Pasricha, M. (2019). Viruses of polar aquatic environments. *Viruses*,
494 11(2). doi: 10.3390/v11020189
- 495 Zhang, Y. Z., Shi, M., & Holmes, E. C. (2018). Using metagenomics to characterize an
496 expanding virosphere. *Cell*, 172(6), 1168–1172. doi: 10.1016/j.cell.2018.02.043
- 497 Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for aligning
498 DNA sequences. *J. Comput. Biol.*, 7(1-2), 203–214. doi: 10.1089/10665270050081478
- 499 Zhou, P., & Shi, Z.-L. (2021). SARS-CoV-2 spillover events. *Science*, 371(6525), 120–122.
500 doi: 10.1126/science.abf6097
- 501 Zohdy, S., Schwartz, T. S., & Oaks, J. R. (2019). The Coevolution Effect as a driver of
502 spillover. *Trends Parasitol.*, 35(6), 399–408. doi: 10.1016/j.pt.2019.03.010

503 Tables

Table 1. Abundance of the viral families of all the viral HSPs. The host range information was obtained from the ViralZone and International Committee on Taxonomy of Viruses (ICTV) databases. Viruses with a missing family were excluded from this table.

Viral family	Known eukaryotic host range	Count by site					
		Control		Low runoff		High runoff	
		C-Soil	C	L-Soil	L1	H-Soil	H1
dsDNA viruses							
<i>Alloherpesviridae</i>	Fish						1
<i>Ascoviridae</i>	Insects: mainly Noctuids						4
	SfAV: <i>Spodoptera</i> species only						
<i>Asfarviridae</i>	Pigs warthogs bushpigs						2
	Vector: Argasid ticks						
<i>Baculoviridae</i>	Arthropods: Lepidoptora, Hymenoptera, Diptera		3				
	Crustacean: Decapoda (Shrimps)						
<i>Herpesviridae</i>	Vertebrates			2	4	2	
<i>Iridoviridae</i>	Insects		11		13		159
<i>Lavidaviridae</i>	Protists infected by <i>Mimivirus</i>				1		8
<i>Marseilleviridae</i>	Amoeba						3
<i>Mimiviridae</i>	Amoeba	86	105	60	167	82	886
<i>Papillomaviridae</i>	Vertebrates						4
<i>Phycodnaviridae</i>	Alga	1	69	1	108	37	1,858
<i>Pithoviridae</i>	Amoeba	1	2		4		2
<i>Poxviridae</i>	Human, vertebrates, and arthropods		2		18	20	173
dsRNA viruses							
<i>Chrysoviridae</i>	Fungi	12		9		116	69
<i>Endornaviridae</i>	Plants, fungi, and oomycetes	33	20	115	21	4	212
<i>Hypoviridae</i>	Fungi			2			4
<i>Megabirnaviridae</i>	Fungi	1		3			
<i>Partitiviridae</i>	Fungi and plants	206	384	112	304	352	1,140
<i>Picobirnaviridae</i>	Vertebrates and invertebrates		3		3		36
<i>Reoviridae</i>	Vertebrates, invertebrates, plants, and fungi		1		4		106
<i>Totiviridae</i>	<i>Totivirus</i> : Fungi	4	44	45	124	19	957

<i>Victorivirus</i> : Fungi							
-ssRNA viruses							
<i>Fimoviridae</i>	Plant: European mountain ash	6					
<i>Mymonaviridae</i>	<i>Sclerotinia sclerotiorum</i> fungi	1		3		6	
<i>Phasmaviridae</i>	Insects (mosquitos, cockroaches, water striders, psyllids, odonates, and drosophilids)	3					2
<i>Phenuiviridae</i>	RVFV: ruminants, camels, and humans Vector: Mosquitoes	3		8		7	
+ssRNA viruses							
<i>Alphaflexiviridae</i>	Plants and fungi	20			8		50
<i>Barnaviridae</i>	Cultivated mushroom (<i>Agaricus bisporus</i>)	1					
<i>Betaflexiviridae</i>	Plants and fungi	11		103	1,716		4,631
<i>Botourmiaviridae</i>	Plants and fungi	52		7		142	
<i>Bromoviridae</i>	Plants			2	4		8
<i>Caliciviridae</i>	Vertebrates		6		7	2	7
<i>Closteroviridae</i>	Plants						2
<i>Deltaflexiviridae</i>	Fungi and plants	2	30			27	
<i>Dicistroviridae</i>	Invertebrates				31		6
<i>Iflaviridae</i>	Insects	2	3		3		3
<i>Luteoviridae</i>	Plants		2				
<i>Marnaviridae</i>	Phytoplankton <i>Heterosigma akashiwo</i>		7		23		5
<i>Mitoviridae</i>	Fungi	66	4		8	26	37
<i>Narnaviridae</i>	Fungi	59	3	26		207	3
<i>Nodaviridae</i>	Vertebrates and invertebrates		3		4		
<i>Picornaviridae</i>	Vertebrates				8		9
<i>Potyviridae</i>	Plants				12		33
<i>Solemoviridae</i>	Plants (few species of Gramineae)	114		54			26
<i>Tombusviridae</i>	Plants		21		9	8	40
<i>Virgaviridae</i>	Plants					4	16
Total		684	723	552	2,604	1,061	10,502

504 **Figures**

Figure 2. Normalized Gini coefficients (G^*) obtained with Random TaPas ($n = 3$ runs). The values are separated by runoff volume: (A) control; (B) low runoff; and (C) high runoff. The two global-fit models used were GD (geodesic distances in tree space) and PACo (Procrustes Approach to Cophylogeny). Significant results (Dunn test, BH correction) are marked with letters from *a* to *j* ($\alpha = 0.05$). Blue represents the soil and yellow, the lake sediments.

Figure 3. Largest and smallest residuals per sampling site for (A) soil and (B) lake sediments samples. Residuals were computed by Random TaPas ($n = 3$ runs) using GD (geodesic distances in tree space). Significant results (Dunn test, BH correction) are marked with an asterisk (*) ($\alpha = 0.05$). Red represents the largest and blue, the smallest residuals. Figure S6 further shows these results to be robust to the distance used to compare trees.