

A dietary vitamin B12 deficiency impairs motor function and changes neuronal survival and choline metabolism after ischemic stroke in middle aged male and female mice

Gyllian B. Yahn BA¹, Brandi Wasek², Teodoro Bottiglieri PhD², Olga Malysheva PhD³, Marie A. Caudill PhD³, Nafisa M. Jadavji PhD^{1,4}

Affiliations

¹ Department of Biomedical Sciences, Midwestern University, Glendale, USA

² Center of Metabolomics, Institute of Metabolic Disease, Baylor Scott & White Research Institute, Dallas, TX, USA

³ Division of Nutritional Sciences, Cornell University, Ithaca, USA

⁴ Department of Neuroscience, Carleton University, Ottawa, Canada

Sources of support: Midwestern University Start-up Funds

Conflict of Interest and Funding Disclosure:

Corresponding Author:

Nafisa M. Jadavji, PhD

Department of Biomedical Sciences

College of Graduate Studies

Midwestern University

19555 N 59th Ave

Glendale, AZ 85308, USA

Phone: 623-572-3665

Fax : 623-572-3647

E-mail: njadav@midwestern.edu

Word Count: 3,634

Number of Figures: 6

Number of Tables: 2

Supplementary data submitted: None

Running Title: Vitamin B12 deficiency and stroke

Abbreviations: Total homocysteine; tHcy, neuronal nuclei; NeuN, 4',6-diamidino-2-phenylindole; DAPI

Abstract

Background

Nutrition is a modifiable risk factor for ischemic stroke. As people age their ability to absorb some nutrients decreases, a primary example is vitamin B12. Older individuals with a vitamin B12 deficiency are at a higher risk for ischemic stroke and have worse outcome after stroke. However the mechanisms through which these occur remain unknown.

Objective

The aim of the study was to investigate the role of vitamin B12 deficiency in ischemic stroke outcome and mechanistic changes in a mouse model.

Methods

At 10-weeks of age male and female mice were put on control or vitamin B12 deficient diets for 4-weeks prior to and after ischemic stroke to the sensorimotor cortex. At 18 weeks of age, we assessed motor outcome using the accelerating rotarod and forepaw placement tasks. At the end of experiments, tissues were collected to assess potential mechanisms.

Results

All animals maintained on the vitamin B12 deficient diet had increased levels of total homocysteine in plasma and liver tissue. After ischemic stroke, male and female mice maintained on a vitamin B12 deficient diet had impaired motor function compared to control animals. In ischemic brain tissue no difference between groups in lesion volume. However, there was an increase in total apoptosis within the ischemic region of brain tissue in male vitamin B12 deficient animals. More neuronal survival was present in ischemic brain tissue of the vitamin B12 deficient group compared to controls. Additionally, there were changes in choline metabolites in ischemic brain tissue because of a vitamin B12 deficiency.

Conclusions

The data presented in this study confirms that a vitamin B12 deficiency impacts motor function in older adult male and female mice after ischemic stroke. The mechanisms driving this change may be a result of neuronal survival and compensation in choline metabolism within the damaged brain tissue.

Keywords

Vitamin B12; stroke, sex differences; motor function; apoptosis; choline metabolism

Introduction

A large complication in public health today is overall health and diet. It is known that eating healthy is important for good health and can reduce the risk of diseases such as stroke (1). It is becoming increasingly popular to partake in a vegetarian or vegan diet, for many reasons. These kinds of diets eliminate meat, which is one of the main sources of our daily required B vitamins and are extremely important for maintaining good health and everyday functioning. Vitamin B12 is also referred to as cobalamin and is a water-soluble vitamin found naturally in foods such as red meat, poultry, eggs, dairy products, seafood, and fortified cereals (2). Vitamin B12 has very important roles in the human body as it is necessary for proper red blood cell formation, neurological function, myelin synthesis and repair, and DNA synthesis. A vitamin B12 deficiency is a serious and common complication that is defined as low plasma and tissue levels of vitamin B12 (3). It can affect all age ranges, however it has a much higher prevalence within the elderly population (1,4).

In elderly patients, a vitamin B12 deficiency is commonly caused by malabsorption, decreased acid secretion, and reduced intrinsic factor production (4). Having a deficiency in vitamin B12 results in a multitude of problems and can exacerbate outstanding conditions or developing conditions. The recommended daily amount of vitamin B12 is 2.4 μg and the average daily diet contains a range of 3-30 μg (5). Interestingly, only 50% of patients who are vitamin B12 deficient present with low levels of serum vitamin B12, resulting in a substantial number of undiagnosed patients (2). Clinical manifestations can be set back due to the high amounts of hepatic storage of vitamin B12 (5).

Adequate levels of vitamin B12 are needed for successful aging (6). The presence of atrial fibrillation, vitamin B12 deficiency, and resultant elevated levels of plasma total homocysteine

(tHcy) increases with age and are a risk factor for stroke (7,8). Clinical data strongly suggest that low levels of vitamin B12 are a risk factor for ischemic stroke and also impacts stroke outcome (9,10). There is a gap in the understanding of the mechanisms in which a vitamin deficiency leads to increased risk of stroke and worse stroke outcome. This study aims to assess the effects of a vitamin B12 deficiency on stroke outcome and related mechanisms.

Methods

Animals

All experiments in animals were approved by the Midwestern University IACUC committee. Female and male C57/BL6J mice were obtained for Jackson laboratories for this study. A total of 44 mice were obtained, 22 males and 22 females. The mice 10-months-old upon arrival and were habituated for 1 week prior to the start of experiments.

Experimental design

An overview of all experimental manipulations is outlined in Figure 1. After mice were habituated to the Midwestern University animal facility, mice were randomly assigned to control or vitamin B12 deficient groups and placed on diets for 4 weeks. At 11 months of age, ischemic stroke was induced in animals using the photothrombosis model (11–15) this corresponds to middle aged in humans (16). The ischemic stroke damaged the sensorimotor cortex, which allows for motor function assessment. Four weeks after damage motor function was measured in animals using the forepaw placement and accelerating rotarod tasks. After the completion of behavioral testing animals were euthanized and tissue was collected, including brain, liver, and blood tissue to study mechanisms.




Figure 1. Timeline of experimental manipulation (A). Male and female C57Bl/6J mice arrived from Jackson Laboratories 10-months-old, and after one week of acclimation, animals were placed on a control or vitamin B12 deficient diet for four weeks. Following the four weeks, ischemic stroke was induced using the Photothrombosis model in the sensorimotor cortex. After stroke animals were maintained on respective diets for 4 additional weeks, after which motor function of the mice was measured using the accelerating rotarod and forepaw placement tasks. At the completion of *in vivo* experiments animals were euthanized, and brain and liver tissue, as well as plasma was collected for further analysis. Weekly body weights of animals maintained on control and vitamin B12 deficient diets (B).

Diet

The mice were placed on a vitamin B12 deficient (0 mg/kg) or a control (0.025 mg/kg vitamin B12) diet four weeks prior to photothrombosis damage and four weeks post photothrombosis damage. The diets were formulated by and purchased from Envigo. The control diet (TD. 190790) contains the recommended dietary amount of nutrients for mice (17). The vitamin B12 deficient diet (TD. 190791) was formulated based on a previous study in mice that has shown it to be safe and have no negative side effects (18). The mice had *ad libitum* access to food and water throughout the experiment. Body weights of each animal were recorded weekly.

Photothrombosis model

Using the photothrombosis model of ischemic stroke damage, mice were anesthetized with 4-5% isoflurane in oxygen. After anesthetization the mice had the top of their heads shaved and disinfected. Tear gel was used to prevent their eyes from drying out during while anesthetized. 0.03 mg/kg of Buprenorphine and one mL of saline were administered subcutaneously. Mice were then transferred to a stereotaxic apparatus (Harvard Apparatus) and maintained at 2-2.5% isoflurane. The mice were placed on a heating pad and a probe was rectally inserted to maintain a body temperature of 37°C. Prior to laser exposure, mice were intraperitoneally injected with 10 mg/kg of photoactive Rose Bengal (Sigma) followed by a 5- minute delay to allow the dye to enter circulation. Skin at the top of the head was surgically cut to expose the skull and then the sensorimotor cortex was targeted using stereotaxic coordinates (3 cm above, mediolateral + 0.24 mm from Bregma). The skull of the mice was exposed to a laser (Beta Electronics, wavelength: 532 nm) for 15 minutes. For recovery of post-operative pain, Buprenorphine was administered to all animals prior to ischemic damage.

Behavioral Testing

Accelerating Rotarod

A standard accelerating rotarod apparatus (Harvard Apparatus) was used to measure walking movements and balance previously described (13,19,20). Thirty centimeters above the ground, mice were placed on a rotating rod 3 cm in diameter and 6 cm wide in which the speed gradually increases from 4 to 60 revolutions per minute over 8 minutes. When mice fall off the rotarod, a digital sensor recorded the latency, in seconds, to fall off the accelerating rod. An average of three trials per mouse was taken with an inter trial interval of five minutes.

Forepaw placement task

To measure spontaneous forepaw usage, mice were placed in a 19 cm high, 14 cm diameter cylinder, and the placement of their forepaws on the cylinder wall during natural exploratory rearing behaviors was recorded using a digital camera for frame-by-frame analysis (13,21). During a rear, the first forepaw placement on the wall was recorded as impaired, non-impaired, or both.

Total homocysteine levels

At the time of euthanization, blood was collected by cardiac puncture in EDTA coated tubes, centrifuged at 7000g for 7 minutes at 4C to obtain plasma. Liver tissue was also removed at the same time and samples were stored at -80C, until time of analysis. Total homocysteine (tHcy) in plasma and liver were measured by liquid chromatography tandem mass spectrometry (LC-MS/MS) as previously described ((22)).

Brain tissue processing

Brain tissue was sectioned using a cryostat at 30 μ m and slide mounted in serial order. There were six slides full of brain tissue sections of the damaged area per mouse and each animal had a minimum of four sections that were used for quantification. ImageJ (NIH) was used to quantify ischemic damage volume by measuring the area of damaged tissue (23).

Immunofluorescence experiments

Brain tissue was used in immunofluorescence analysis to assess molecular mechanisms and brain tissue staining was performed to investigate potential mechanisms. Primary antibodies included, active caspase-3 (1:100, Cell Signaling Technologies) to measure apoptosis and phospho-AKT (protein kinase B) (1:100, Cell Signaling Technologies). All brain sections were

stained with a marker for neuronal nuclei, NeuN (1:200, AbCam). Primary antibodies were diluted in 0.5% Triton X and incubated with brain tissue overnight at 4°C. The next day, brain sections were incubated in Alexa Fluor 488 or 555 (Cell Signaling Technologies) and secondary antibodies were then incubated at room temperature for 2 hours and stained with 4', 6-diamidino-2-phenylindole (DAPI) (1:1000, Thermo Fisher Scientific). The stains were analyzed using a microscope (Zeiss) and all images were collected at the magnification of 40X.

In brain tissue within the ischemic region, co-localization of active caspase-3 or phospho-AKT with NeuN labelled neurons were counted and averaged per animal. A positive cell was indicated by co-localization of the antibodies of interest located within a defined cell. Cells were distinguished from debris by identifying a clear cell shape and intact nuclei (indicated by DAPI and NeuN) under the microscope. All cell counts were conducted by two individuals blinded to treatment groups. The number of positive cells were counted in three brain sections per animal. For each section, three fields were analyzed. The number of positive cells were averaged for each animal.

Choline metabolites measurements

Frozen brain tissue from ischemic and non-ischemic cortex was measured for acetylcholine, betaine, choline, glycerophosphocholine, phosphocholine, phosphatidylcholine, and sphingomyelin levels using the LC-MS method as previously reported (24).

Data analysis and statistics

All data were analyzed by two individuals that were blinded to experimental treatment groups. GraphPad Prism 6.0 was used to analyze all data from the study. Two-way ANOVA analysis was performed when comparing the mean measurement of both sex and dietary group for

behavioral testing, plasma tHcy measurements, lesion volume, immunofluorescence staining, and choline measurements. Significant main effects of two-way ANOVAs were followed up with Tukey's post-hoc test to adjust for multiple comparisons. All data are presented as mean \pm standard error of the mean (SEM). Statistical tests were performed using a significance level of 0.05.

Results

Body weight

To assess health animals were weighed weekly. There was difference between male and females in body weight (Figure 1B ($t_{1,38}$) = 4.63, p = 0.001). There was no change in body weight between groups prior to or after photothrombosis damage (Figure 1B, $F_{(1,33)}$ = 45.5, p = 0.27).

Behavioral Testing

Accelerating Rotarod

Four weeks after ischemic damage to the sensorimotor cortex, balance and coordination were measured via the accelerating rotarod task (19). We measured the latency to fall, there was no difference between diet groups (**Figure 2A**; $F_{(1,32)}$ = 3.06, p = 0.09). Female mice stayed on the rotarod longer compared to male mice (**Figure 2A**; $F_{(1,32)}$ = 5.52, p = 0.025).

The revolutions per minute reached by each animal were also measured during the accelerating rotarod task. Control diet mice achieved higher levels of speed compared to vitamin B12 deficient mice (Figure 2B; $F_{(1,32)}$ = 7.44, p = 0.010). Male control mice were able to achieve a higher RPM compared to vitamin B12 deficient animals (Tukey's pairwise, p = 0.021). Female mice achieved higher speeds compared to male mice (Figure 2B; $F_{(1,32)}$ = 9.33, p = 0.0045).

Forepaw Placement

The forepaw placement task evaluated forepaw usage through natural rearing behaviors of rodents (Theoret et al., 2016). Four weeks after ischemic damage, there were differences in impaired forepaw usage scores between control and vitamin B12 deficient mice (Figure 2C; $F_{(1,38)} = 6.27$, $p = 0.017$). Female control mice used their impaired forepaw more than vitamin B12 deficient animals (Tukey's pairwise, $p = 0.015$). There was no sex difference ($F_{(1,38)} = 1.86$, $p = 0.18$).

There was a diet difference in non-impaired forepaw usage scores between control and vitamin B12 deficient mice (Figure 2D; $F_{(1,38)} = 11.41$, $p = 0.002$, diet effect) and sex difference ($F_{(1,38)} = 6.136$, $p = 0.018$). Female control mice used their non-impaired forepaw more than vitamin B12 deficient animals (Tukey's pairwise comparison, $p = 0.012$).





Figure 2. Impact of dietary vitamin B12 deficiency and ischemic stroke on motor function. Coordination and balance were assessed using the accelerating rotarod behavioral task. The latency to fall (A) and revolutions per minute (B) were measured. The forepaw placement task was used to measure forepaw usage. The number of impaired (A) and non-impaired (B) forepaw placements

were recorded. Depicted are means of \pm SEM of 10 to 11 mice per group. Tukey's pairwise comparison * $p < 0.01$ between indicated groups.

Total homocysteine levels

To confirm vitamin B12 deficiency, total homocysteine (tHcy) levels in the plasma and liver were measured. In plasma, vitamin B12 deficient mice, plasma homocysteine levels were elevated compared to control diet mice (Figure 3A; $F_{(1,26)} = 15.05$, $p = 0.0006$). Female vitamin B12 deficient mice had elevated levels of homocysteine compared to control animals (Tukey's pairwise comparison, $p = 0.048$). There was no difference between male and female mice (Figure 3A; $F_{(1,26)} = 0.20$, $p = 0.66$).

In vitamin B12 deficient mice, liver tHcy levels were elevated compared to control diet mice (Figure 3B; $F_{(1,19)} = 47.86$, $p < 0.0001$) and there was also a sex effect (Figure 3B; $F_{(1,19)} =$



10.30, $p = 0.005$). Female vitamin B12 deficient mice had elevated levels of tHcy compared to control animals (Tukey's pairwise comparison, $p = 0.001$).

Figure 3. Impact of dietary vitamin B12 deficiency and ischemic stroke on total homocysteine levels. Plasma (A) and liver (B) homocysteine levels (B). Depicted are means of \pm SEM of 7 to 8 mice per group. *** $p < 0.01$ between indicated groups.

Ischemic Damage Volume Quantification

Ischemic brain damage to the sensorimotor cortex was measured. Representative images of brain

tissue are shown in **Figure 4A**. There was no diet effect (Figure 4B; $F_{(1,14)} = 3.65$, $p = 0.077$) or sex effect ($F_{(1,14)} = 1.27$, $p = 0.28$).




Figure 4. Impact of dietary vitamin B12 deficiency and ischemic stroke on damage volume. Representative cresyl violet image (A) and ischemic damage volume quantification (B). Depicted are means of \pm SEM of 4 to 5 mice per group.

Brain Tissue Immunofluorescence Staining

Neuronal Apoptosis

Neuronal apoptosis was assessed within the damage area using active caspase-3 and neuronal nuclei (NeuN), a marker of neuronal nuclei immunofluorescence experiments. Representative images are shown in Figure 5A. Quantification of neuronal apoptosis revealed that there was no diet (Figure 5B; $F_{(1,10)} = 4.60$, $p = 0.056$) or sex ($F_{(1,10)} = 2.81$, $p = 0.13$) effects.

Evaluation of total active caspase-3 within the damage area was compiled by counting the number of antibody active caspase 3 positive profiles. Representative images are shown in Figure 5C. There was a main effect of diet (Figure 5D; $F(1, 10) = 8.28$, $p = 0.017$), but no effect of sex ($F(1, 10) = 0.070$, $p = 0.80$). Male vitamin B12 deficient animals had more total active caspase-3 cells within the damage area (Tukey's pairwise comparison, $p = 0.0055$).




Figure 5. Impact of dietary vitamin B12 deficiency and ischemic stroke on neuronal active caspase-3 cell counts. Representative images of immunofluorescence staining with positive semi quantitative spatial co-localization of active caspase-3 with neuronal nuclei (NeuN) and 4',6-diamidino-2-phenylindole (DAPI) (A). Quantification of active caspase-3, NeuN, and DAPI cell counts (B). Depicted are means of \pm SEM of 3 to 4 mice per group. The scale bar = 50 μm .

Neuronal Cell Survival and Proliferation

Neuronal survival and proliferation were measured using colocalization of pAKT and NeuN. Representative images are shown in Figure 6A. Vitamin B12 deficient animals had more positive neuronal pAKT (Figure 6B, $F_{(1,9)} = 16.6$, $p = 0.0028$), but no difference between male and females ($F_{(1,9)} = 0.03$, $p = 0.856$).




Figure 6. Impact of dietary vitamin B12 deficiency and ischemic stroke on phospho-AKT (pAKT) cell counts. Representative images of immunofluorescence staining with positive semi quantitative spatial co-localization of active caspase-3 with neuronal nuclei (NeuN) and 4',6-diamidino-2-phenylindole (DAPI) (A). Quantification of pAKT, NeuN, and DAPI cell counts (B). Depicted are means of \pm SEM of 3 to 4 mice per group. The scale bar = 50 μ m.

Choline Metabolite Measurements

A dietary vitamin B12 deficiency has been linked to changes in choline metabolism (25,26). Offspring choline metabolites were measured in brain tissue with ischemic (Table 1) and non-ischemic (Table 2) brain tissue.

Table 1. Choline metabolite concentration in cortical ischemic hemisphere of male and female mice maintained on control and vitamin B12 diets.

Metabolites (nmol/g)	Control diet		Vitamin B12 Deficient diet		Effect	p-value
	Female	Male	Female	Male		
Methionine	62.91 ± 13.12	85.66 ± 5.83	97.00 ± 2.03	103.31 ± 11.79	Diet	0.0232
Acetylcholine	3.90 ± 0.92	4.92 ± 5.08	5.64 ± 1.59	3.95 ± 0.34		
Betaine	20.62 ± 1.10	13.15 ± 0.94	15.35 ± 1.63	11.61 ± 1.63	Sex and diet	0.0366; 0.0033
Choline	67.47 ± 4.11	51.44 ± 2.29	55.67 ± 6.89	64.05 ± 7.02		
GPC	679.61 ± 18.52	756.71 ± 60.35	676.34 ± 24.74	756.71 ± 60.35		
Phosphocholine	381.49 ± 33.30	333.18 ± 2.58	338.49 ± 20.54	329.26 ± 13.17		
Phosphatidylcholine	34162.62 ± 566.20	35640.89 ± 566.20	34706.88 ± 651.01	35626.43 ± 227.88		
Sphingomyelin	4269.26 ± 186.93	4075.13 ± 202.93	4268.67 ± 68.54	3999.57 ± 23.03		

Table 2. Choline metabolite concentration in cortical non-ischemic hemisphere of male and female mice maintained on control and vitamin B12 diets.

Metabolites (nmol/g)	Control diet		Vitamin B12 Deficient diet		Effect	p-value
	Female	Male	Female	Male		
Methionine	77.51 ± 12.43	79.19 ± 9.36	90.77 ± 5.94	107.07 ± 9.84	Diet	0.029
Acetylcholine	3.64 ± 0.36	4.17 ± 0.012	5.03 ± 0.59	4.79 ± 0.33		
Betaine	25.53 ± 2.91	46.47 ± 1.45	15.14 ± 0.83	15.24 ± 4.93	Sex	0.0151
Choline	67.92 ± 6.39	46.47 ± 6.90	67.92 ± 6.39	82.09 ± 4.95		
GPC	635.66 ± 5.68	611.81 ± 40.95	649.40 ± 10.85	692.71 ± 2.11		
Phosphocholine	353.28 ± 9.50	385.74 ± 16.73	353.28 ± 9.50	309.66 ± 14.10		
Phosphatidylcholine	34546.56 ± 1271.33	32335.41 ± 1178.83	35283.09 ± 497.43	34727.34 ± 262.79		
Sphingomyelin	4342.97 ± 148.72	3653.92 ± 41.67	4518.94 ± 39.26	3766.14 ± 116.91	Sex	0.0001

Measurements within the ischemic cortex showed that methionine levels were increased in vitamin B12 tissue compared to controls (Table 1; diet, $F_{(1,8)} = 7.84$, $p = 0.023$). Betaine levels were reduced in male and female animals on a vitamin B12 deficient diet compared to control diet animals (Table 1; sex, $F_{(1,8)} = 17.0$, $p = 0.0033$; diet, $F_{(1,8)} = 6.28$, $p = 0.036$).

In the non-ischemic cortex acetylcholine levels were increased between vitamin B12 and control diet animals (Table 2, diet, $F_{(1,8)} = 7.84$, $p = 0.023$). Phosphocholine levels were decreased in vitamin B12 deficient animals compared to control animals (Table 2; diet, $F_{(1,8)} = 0.95$, $p = 0.0151$). Sphingomyelin levels were also increased in vitamin B12 deficient animals compared to controls (Table 2; diet, $F_{(1,8)} = 9.50$, $p = 0.015$).

Discussion

Nutrition is a modifiable risk factor for stroke (27). The elderly population is at higher risk of stroke as well as a vitamin B12 deficiency (4). Worse stroke outcome has been reported in patients with a vitamin B12 deficiency (9,10). The aim of the study was to investigate the role of vitamin B12 deficiency in ischemic stroke outcome and mechanistic changes in a mouse model. Our

findings showed that vitamin B12 deficient mice had more impairments in coordination and balance compared to control diet mice after ischemic damage. Forepaw usage was decreased in female mice maintained on the vitamin B12 diet. Male mice on a vitamin B12 deficient diet showed impaired coordination and balance. In both plasma and liver there was an increase in plasma and liver tHcy because of dietary vitamin B12 deficiency. There was no difference in the damage size and neuronal apoptosis induced by the ischemic stroke between groups. There was an increase in overall apoptosis within the ischemic cortex. We also report increased neuronal survival in vitamin B12 deficient male and female mice, along with changes in choline metabolism in brain tissue.

Lacunar stroke patients with vitamin B12 deficiency reported significantly worse stroke outcome, including fatigue and depressive-like symptoms (28). A common stroke outcome is impaired motor function, and the two most common motor deficits from stroke are spasticity and paresis, resulting in decreased usage of that limb. We measured forepaw usage and saw that female mice had significantly reduced impaired and non-impaired forepaw usage. Male vitamin B12 deficient mice showed impairments in coordination and balance using the rotarod task. Our preclinical study results are consistent with clinical observations in terms of worse outcome after ischemic stroke when patients are vitamin B12 deficient (10).

In the present study, we did not observe any differences in damage volume between dietary groups; however, this is a gross measurement. In our immunofluorescence experiments we report a difference in in apoptosis between dietary groups. In our neuronal apoptosis data, there was a decrease in neuronal death in vitamin B12 deficient mice. This may be due to increased neuronal plasticity in the damaged area (29). However, total levels of apoptosis showed an increase in overall positive active caspase-3 levels in male vitamin B12 deficient mice. Increased levels of

apoptosis have been reported in clinical studies, for example, in an autopsy cohort consisting of 13 cases of fatal ischemic stroke, TUNEL-labelled cells with apoptotic morphology were disproportionately more frequent in the ischemic core, compared percentage of the cells in the non-damaged hemisphere (30). Furthermore, in post-ischemic stroke there is elevated levels of neuronal death due to apoptosis in the damaged area of the brain (31). A potential reason there is not an increase in apoptosis in female mice is the neuroprotective effects of estrogen and progesterone (32). This finding could also reflect an increased vulnerability of glial cells, the other type of cell in the brain. Overall, these findings support clinical data being reported (10).

One-carbon metabolism incorporates several vitamins and nutrients. Other studies have shown that vitamin B12 status affects choline status in animals (25,26). We measured levels of choline in brain tissue in ischemic and non-ischemic cortical tissue. Our results show that ischemic brain tissue has higher methionine levels in vitamin B12 deficient mice. The increase in methionine levels might be in compensation to the dietary vitamin B12 deficiency, since the brain is sensitive to methionine (33). We also report lower levels of betaine in ischemic brain tissue of vitamin B12 deficient mice, which might reflect what's happening in liver and plasma, and availability for uptake by brain.

Part of this study investigated sex differences in response to ischemic stroke between dietary groups. This is an important component of preclinical studies as there have been a low number of studies using female mice (34,35). In our study, male and female mice maintained on vitamin B12 deficiency had impaired stroke outcome, although they had impairments of different motor tasks. In tissue, male mice showed more total apoptosis, however in our present study our data shows both males and females had increased neuronal survival within damage region in brain tissue. This is an interesting finding considering research has shown that men are more susceptible

to a vitamin B12 deficiency (36) . The results of our study emphasizes the importance of studying of sex differences in preclinical stroke experiments and the possibility of individualized medicine (37).

Conclusion

The data presented in this study provides evidence that a vitamin B12 deficiency impacts motor function in older adult male and female mice after ischemic stroke. The mechanisms driving this change may be a result of increased apoptosis, changes in neuronal survival, and compensation in choline metabolism within the damaged brain tissue.

Acknowledgments

The authors would like to acknowledge the Chelsea Adamson for her assistance in photothrombosis surgeries.

Statement of authors' contributions to manuscript

Conceptualization, NMJ; Data curation, GBY, BW, TB, OM, MAC, NMJ; Formal analysis, GBY, TB, MAC, NMJ; Funding acquisition, NMJ; Investigation; Methodology, GBY, BW, TB, OM, MAC, NMJ; Supervision, NMJ; Visualization; Roles/Writing - original draft, GBY and NMJ; Writing - review & editing, GBY, TB, MAC, NMJ. N.M.J. had primary responsibility for final content. All authors read and approved the final manuscript.

References

1. Spence JD. Nutrition and risk of stroke. *Nutrients* 2019;11.
2. Langan RC, Goodbred AJ. Vitamin B12 Deficiency: Recognition and Management. *Am Fam Physician* 2017;96:384–9.
3. Spence JD. Metabolic B12 deficiency: A missed opportunity to prevent dementia and stroke. *Nutrition Research Elsevier Inc.*; 2016;36:109–16.
4. Andrès E, Loukili NH, Noel E, Kaltenbach G, Abdelgheni MB, Perrin AE, Noblet-Dick M, Maloisel F, Schlienger J-L, Blicklé J-F. Vitamin B12 (cobalamin) deficiency in elderly patients. *CMAJ* 2004;171:251–9.
5. Moridani M, Ben-Poorat S. Laboratory Investigation of Vitamin B12 Deficiency. *Laboratory Medicine* 2006;37:166–74.
6. Vogiatzoglou A, Refsum H, Johnston C, Smith SM, Bradley KM, de Jager C, Budge MM, Smith AD. Vitamin B12 status and rate of brain volume loss in community-dwelling elderly. *Neurology* 2008;71:826–32.
7. Spence JD. Homocysteine-lowering therapy: a role in stroke prevention? *Lancet neurology* 2007;6:830–8.
8. Spence JD, Stampfer MJ. Understanding the complexity of homocysteine lowering with vitamins: The potential role of subgroup analyses. *JAMA - Journal of the American Medical Association* 2011;306:2610–1.
9. Ahmed S, Bogiatzi C, Hackam DG, Rutledge AC, Sposato LA, Khaw A, Mandzia J, Azarpazhoo MR, Hachinski V, Spence JD. Vitamin B 12 deficiency and hyperhomocysteinaemia in outpatients with stroke or transient ischaemic attack: A cohort study at an academic medical centre. *BMJ Open* 2019;9:1–8.
10. Yahn G, Abato J, Jadavji N. Role of vitamin B12 deficiency in ischemic stroke risk and outcome. *Neural Regeneration Research* 2021;16.
11. Abato JE, Mofteh M, Cron GO, Smith PD, Jadavji NM. Methylene tetrahydrofolate reductase deficiency alters cellular response after ischemic stroke in male mice. *Nutritional Neuroscience [Internet]* Taylor and Francis Ltd.; 2020; Available from: <https://pubmed.ncbi.nlm.nih.gov/32448097/>
12. Jadavji N, Emmerson JT, Shanmugalingam U, Willmore WG, Macfarlane AJ, Smith PD. A genetic deficiency in folic acid metabolism impairs recovery after ischemic stroke. *Experimental neurology* 2018;309:14–22.

13. Jadavji NM, Emmerson J, Willmore WG, MacFarlane AJ, Smith P. B-vitamin and choline supplementation increases neuroplasticity and recovery after stroke. *Neurobiology of Disease* 2017;103:89–100.
14. Labat-gest V, Tomasi S. Photothrombotic ischemia: a minimally invasive and reproducible photochemical cortical lesion model for mouse stroke studies. *Journal of visualized experiments : JoVE* 2013;e50370–e50370.
15. Lee J-K, Kim J-E, Sivula M, Strittmatter SM. Nogo receptor antagonism promotes stroke recovery by enhancing axonal plasticity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2004;24:6209–17.
16. Flurkey K, Curren JM, Harrison DE. Mouse Models in Aging Research. In: Fox J, Barthold S, Davisson M, Newcomer C, Quimby F, Smith A, editors. *The-Mouse-in-Biomedical-Research, Normative Biology, Husbandry, and Models*. [Internet] 2nd ed. Elsevier.; 2007. p. 637–72. Available from: https://mouseion.jax.org/stfb2000_2009/1685
17. Reeves PG. Components of the AIN-93 Diets as Improvements in the AIN-76A Diet. *The Journal of Nutrition* 1997;127:838S-841S.
18. Bernard DJ, Pangilinan FJ, Cheng J, Molloy AM, Brody LC. Mice lacking the transcobalamin-vitamin B12 receptor, CD320, suffer from anemia and reproductive deficits when fed vitamin B12-deficient diet. *Human Molecular Genetics* 2018;27:3627–40.
19. Balkaya M, Kröber J, Gertz K, Peruzzaro S, Endres M. Characterization of long-term functional outcome in a murine model of mild brain ischemia. *Journal of neuroscience methods* 2013;213:179–87.
20. Jadavji NM, Farr T, Khalil A, Boehm-Sturm P, Foddis M, Harms C, Füchtmeier M, Dirnagl U. Elevated levels of plasma homocysteine, deficiencies in dietary folic acid and uracil-DNA glycosylase impair learning in a mouse model of vascular cognitive impairment. *Behavioural brain research* 2015;283:215–26.
21. Theoret JK, Jadavji NM, Zhang M, Smith PD. Granulocyte macrophage colony-stimulating factor treatment results in recovery of motor function after white matter damage in mice. *European Journal of Neuroscience* 2016;43.
22. Ducros V, Belva-Besnet H, Casetta B, Favier A. A robust liquid chromatography tandem mass spectrometry method for total plasma homocysteine determination in clinical practice. *Clinical Chemistry and Laboratory Medicine (CCLM)* 2006;44:987–90.
23. Jensen EC. Quantitative Analysis of Histological Staining and Fluorescence Using ImageJ. *The Anatomical Record John Wiley & Sons, Ltd*; 2013;296:378–81.
24. Koc H, Mar MH, Ranasinghe A, Swenberg JA, Zeisel SH. Quantitation of choline and its metabolites in tissues and foods by liquid chromatography/electrospray ionization-isotope dilution mass spectrometry. *Analytical Chemistry* 2002;74:4734–40.

25. Compher C, Kinosian B, Stoner N, Lentine D, Buzby G. Choline and vitamin B12 deficiencies are interrelated in folate-replete long-term total parenteral nutrition patients. *JPEN J Parenter Enteral Nutr* 2002;26:57–62.
26. Hegsted DM, Roach AM, McCombs HL. Effect of Vitamin B12 in Choline Deficiency in the Rat. *The Journal of Nutrition* 1967;92:403–11.
27. Spence JD. Stroke Nutrition in Stroke Prevention. 2017;45–51.
28. Huijts M, Duits A, Staals J, van Oostenbrugge RJ. Association of vitamin B12 deficiency with fatigue and depression after lacunar stroke. *PLoS One* 2012;7:e30519.
29. Cirillo C, Brihmat N, Castel-Lacanal E, Le Fricc A, Barbieux-Guillot M, Raposo N, Pariente J, Viguier A, Simonetta-Moreau M, Albucher J-F, et al. Post-stroke remodeling processes in animal models and humans. *J Cereb Blood Flow Metab* 2020;40:3–22.
30. Sairanen T, Karjalainen-Lindsberg M-L, Paetau A, Ijäs P, Lindsberg PJ. Apoptosis dominant in the periinfarct area of human ischaemic stroke--a possible target of antiapoptotic treatments. *Brain* 2006;129:189–99.
31. Uzdensky AB. Photothrombotic Stroke as a Model of Ischemic Stroke. *Translational Stroke Research Springer US*; 2018;9:437–51.
32. Kim T-H, Vemuganti R. Effect of sex and age interactions on functional outcome after stroke. *CNS Neurosci Ther* 2015;21:327–36.
33. Selhub J, Bagley LC, Miller J, Rosenberg IH. B vitamins, homocysteine, and neurocognitive function in the elderly. *The American journal of clinical nutrition* 2000;71:614S-620S.
34. Persky RW, Turtzo LC, McCullough LD. Stroke in women: Disparities and outcomes. *Current Cardiology Reports* 2010;12:6–13.
35. Singh D, Wasan H, Reeta KH. Preclinical Stroke Research and Translational Failure: A Bird's Eye View on Preventable Variables. *Cell Mol Neurobiol* 2021;
36. Margalit I, Cohen E, Goldberg E, Krause I. Vitamin B12 Deficiency and the Role of Gender: A Cross-Sectional Study of a Large Cohort. *Ann Nutr Metab* 2018;72:265–71.
37. Bushnell CD, Chaturvedi S, Gage KR, Herson PS, Hurn PD, Jiménez MC, Kittner SJ, Madsen TE, McCullough LD, McDermott M, et al. Sex differences in stroke: Challenges and opportunities. *J Cereb Blood Flow Metab* 2018;38:2179–91.