

SUPPORTING INFORMATION

A Useful Guide to Lectin Binding: Machine-Learning Directed Annotation of 57 Unique Lectin Specificities

Daniel Bojar¹✉, Lawrence Meche²✉, Guanmin Meng³✉, William Eng², David F. Smith⁴, and Richard D. Cummings⁵, Lara K. Mahal^{2,3*}

1. Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, Sweden. Wallenberg Centre for Molecular and Translational Medicine, Gothenburg, SWEDEN
2. Biomedical Chemistry Institute, New York University Department of Chemistry, 100 Washington Square East, Room 1001, New York, NY, USA, 10003
3. Department of Chemistry, University of Alberta, Edmonton, CANADA, T6G 2S2;
4. Department of Biochemistry, Glycomics Center, Emory University, School of Medicine, Atlanta, GA, USA, 30322.
5. Department of Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA, 02115.




Figure S1. Overview of *N*- and *O*-glycans. **A.** *N*-glycosylation pathway. High mannose is trimmed Man_5 and then elaborated to hybrid, biantennary, triantennary and tetraantennary *N*-glycans. **B.** An example of a more complex biantennary *N*-glycan. The biantennary core is highlighted in grey. **C.** *O*-glycan cores. **D.** An example of a more complex core 1 *O*-glycan. The core is highlighted in grey.

Tables S1-5: Excel sheets.