1 Update on the mosquito fauna (Diptera: Culicidae) distribution in Cabo 2 Verde: occurrence of the species complexes *Anopheles gambiae* and *Culex* 3 *pipiens* (*pipiens*, *quinquefasciatus* and their hybrids)

4 Authors and affiliation

5 Silvânia Da Veiga Leal ^{1,*}, Isaias Baptista Fernandes Varela ¹, Davidson Daniel Sousa
6 Monteiro ¹, Celivianne Marisia Ramos de Sousa ¹, Maria da Luz Lima Mendonça ¹,
7 Adilson José De Pina ² Aderitow Augusto Lopes Gonçalves ¹, and Hugo Costa Osório ^{3,4}

8 ¹ Laboratório de Entomologia Médica, Instituto Nacional de Saúde Pública, Largo do
9 Desastre da Assistência, Chã de Areia, Praia 719, Cabo Verde;

10 ² Programa de Eliminação do Paludismo, CCS-SIDA, Ministério da Saúde e da Segurança
11 Social, Varzea, Praia 855, Cabo Verde;

12 ³ Instituto Nacional de Saúde Doutor Ricardo Jorge, Centro de Estudos de Vectores e
13 Doenças Infecciosas, Avenida da Liberdade 5, 2965-575 Águas de Moura, Portugal;

14 ⁴ Instituto de Saúde Ambiental, Faculdade de Medicina da Universidade de Lisboa, Av.
15 Prof. Egas Moniz, Ed. Egas Moniz, Piso 0, Ala C, 1649-028 Lisboa, Portugal

16 * Correspondence: silvania.leal@insp.gov.cv

17 Abstract

18 In this study, we aimed to update the mosquito species composition and distribution based
19 on a national entomological survey in all municipalities of Cabo Verde. This includes the
20 sibling species of the *Culex pipiens* complex, namely *Cx. pipiens*, *Cx. quinquefasciatus*
21 and their hybrids, in locations where information is not available. The entomological
22 survey took place from October 2017 to September 2018, in all municipalities of Cabo

Mosquito larvae and pupae were collected in breeding sites and samples were sent to the Laboratory of Medical Entomology of the National Institute of Public Health for the morphological identification of the species. The mosquitoes morphologically identified in *Anopheles gambiae* and *Culex pipiens* complexes of species were further molecular analysed to species confirmation. A total of 814 breeding sites were surveyed and 10 mosquito species of five genera were identified. The greatest number of mosquito species was reported in the island of Santiago. The most widespread species in the country were *Aedes aegypti* and *Culex quinquefasciatus*. *Anopheles arabiensis* was the only species identified in the *Gambiae* complex of species. The results of this study will assist decision makers in important health policies to control mosquitoes and vector-borne diseases towards a strategic approach by timely detection of changes in species diversity.

Introduction

Vector-borne diseases represent a public health problem worldwide. At least 80% of the world population is at risk of infection by one or more diseases [1]. A large proportion of these diseases, such as malaria, dengue, chikungunya and Zika their infectious agents are transmitted by mosquitoes, which represent a major threat to human health, with millions of deaths annually [2,3].

In Cabo Verde some of the mosquito species, namely *Aedes aegypti* and *Anopheles arabiensis*, are vectors of pathogens, increasing the risk of diseases outbreaks. Moreover, the flow of people and goods between Cabo Verde and other countries poses a risk for the introduction of new mosquito species into the archipelago, as well as mosquito-borne pathogens. In the last 10 years, Cabo Verde has been affected by three mosquito-borne diseases outbreaks: dengue in 2009, Zika in 2015 and malaria in 2017 [3-7]. To evaluate

the risk of transmission of vector-borne diseases in a given region, a continuous updating of the geographic distribution of insect vectors is required [8].

The first studies on the mosquito fauna in Cabo Verde date from the last century, after the identification of *Anopheles gambiae* s.l. in 1909 by Sant'Anna as described by Ribeiro and collaborators [9]. In 1977, an extensive study was carried out on all nine inhabited islands, resulting in the first and unique dichotomous key of the mosquitoes of Cabo Verde, in which eight species were described in the archipelago [10]. In 1984, the species *Lutzia tigripes* (formerly *Culex tigripes*) was first reported in the country [11]. The last update of Cabo Verde's mosquito fauna was in 2007, based on bibliographic research, wherein entomological survey was carried out only in Maio, Santiago, Fogo and Brava islands. In that study, the species *Culex perexiguus*, a member of the *Culex univittatus* species complex was reported for the first time [12, 13]. Later, the species *Culex tritaeniorhynchus* was identified, raising to a total of 11 known species of mosquitoes recognized in the archipelago [14].

Culex pipiens pipiens and *Culex pipiens quinquefasciatus*, hereinafter referred as *Culex pipiens* and *Culex quinquefasciatus*, are vectors responsible for the transmission of lymphatic filariasis and neurotropic arboviruses to humans, namely West Nile virus [15,16]. These sibling species of the *Culex pipiens* complex were described in Cabo Verde in 1950 and 1980, respectively [10]. The nominal species of the complex, *Culex pipiens*, is found primarily in temperate zones, while *Cx. quinquefasciatus* occurs in warmer tropical and subtropical zones with a higher degree of humidity [17]. However, Cabo Verde is a region where both species coexist in sympatry and where hybrids of the two species were reported from Fogo and Maio locations [12,18]. Here, high levels of hybridization rates between the two species were detected, together with second-generation hybrids identified. The presence of these hybrid forms may locally potentiate

the transmission of arboviruses to humans, such as West Nile virus, and therefore it is so important to have current information about its distribution.

In this study, we aimed to update the mosquito species composition and distribution based on a national entomological survey in all municipalities of Cabo Verde. This includes the sibling species of the *Culex pipiens* complex, namely *Cx. pipiens*, *Cx. quinquefasciatus* and their hybrids, in locations where information is not available. Due to the air and maritime transporting of people and goods in the archipelago, dispersion of species from one island to another, may have occurred after the last mosquito fauna survey in Cabo Verde in 2007. This could result in changes of composition, distribution and abundance of species, which potentially affects the pattern of disease occurrence and transmission. Our results will assist decision makers in important health policies to control mosquitoes and vector-borne diseases towards a strategic approach.

Methodology

Study area

Cabo Verde is a volcanic archipelago with an area of 4.033 km² located about 550 km off the coast of Senegal in West Africa (Figure 1). The archipelago consists of ten islands, nine of which are inhabited with about 537.660 inhabitants. It has a warm and dry, arid and semi-arid climate, with an average annual temperature around 25°C, low rainfall, with two identified seasons: the dry season, from December to June, and the rainy season, from August to October [19].

Mosquito collection

The entomological survey took place from October 2017 to April 2018 in all municipalities of Cabo Verde, except in the island of Santo Antão, in which the survey took place in September 2018. Mosquito larvae and pupae were collected in breeding

sites that included domestic containers and others. The samples were sent to the Laboratory of Medical Entomology (LME) of the National Institute of Public Health in 100 ml pots of water from the breeding site. Larvae L1 and L2 were kept in the insectary and fed with flocculated food for fish (Tropical mix flakes; Ref: F042.1), for later assembly of the L3/L4 stages in slides, prior species identification.

Morphological identification

Larvae in development stage L3 and L4 and adults were used in the morphological identification of mosquito species. The larvae were mounted on slides with 2% glycerinated Hoyer's medium. Morphological identification of the species was performed with the aid of a stereomicroscope using dichotomous keys according to Ribeiro et al. (1980) [10], Ribeiro & Ramos (1995) [9] and auxiliary taxonomic keys according to Dehghan et al. (2016) [20] and Azari-Hamidian & Harbach (2009) [21].

Molecular analyses

The mosquitoes morphologically identified as *Anopheles gambiae* s.l. and *Culex pipiens* s.l. were submitted to single total DNA extraction using the NZY Tissue gDNA isolation kit. Sampled mosquitoes were individually grinded in Lysis Buffer using glass pearls in 2ml Eppendorf and DNA extraction was performed using the prepared lysate suspensions with the NZY Tissue gDNA isolation kit according to manufacturer's recommendations. The Identification of *An. gambiae* complex species were performed by PCR-RFLP following the protocol described by Scott et al. (1993) [22] and Identification of *Cx. pipiens* species was performed according to Smith & Fonseca (2004) [23].

Results

Identification and geographical distribution of mosquito species

All the municipalities of the islands of Cabo Verde (N = 22) were inspected for mosquito larvae and pupae in a total of 814 breeding sites. Five genera and 10 mosquito species were identified (**Error! Reference source not found.**).

The greatest number of mosquito species was reported in the islands of Santiago and Santo Antão (N = 8); followed by Maio (N=7); Boavista, São Vicente, and São Nicolau (N = 6); Fogo (N=5), Sal and Brava (N = 4) (Figure 1).

Figure 1: Geographic distribution of mosquito species in Cabo Verde (*ArcGIS* 10.5).

The most widespread species in the country were *Aedes aegypti* and *Culex quinquefasciatus*, present in all islands of the archipelago (N= 09), followed by *Anopheles pretoriensis* (N = 08) and *Culiseta longiareolata* (N = 06). All the other species were circumscribed to less than five Islands. *Culex tritaeniorhynchus* was the most restricted species found only on the island of Santiago, in the municipality of Tarrafal (Table 1).

A total of 156 mosquitoes identified in the *An. gambiae* complex were molecularly identified as *An. arabiensis* (N = 156; 100%).

In the *Culex pipiens* complex the species the *Cx. quinquefasciatus* was the most identified species (N = 195; 83%) found in every island (Table 2). *Culex pipiens* was found in two islands, Maio and Santo Antão and in lower frequencies. Hybrid forms were identified in four islands (Maio, Fogo, Santo Antão and São Nicolau). Higher frequencies of hybrids were detected in Maio (N = 15; 50%). The species *Culex perexiguus* was not detected.

Discussion

In this study, we present the most recent data on mosquito species distribution in Cabo Verde based on an entomological survey carried in all municipalities of the country. From the 10 species reported, four major mosquito vectors with medical importance were identified, namely *Anopheles arabiensis*, *Aedes aegypti*, *Culex pipiens* and *Cx. quinquefasciatus*.

The only malaria vector found in Cabo Verde was *An. arabiensis*, and it was identified for the first time in Maio and São Vicente in our survey. We corroborate previous studies reporting this species in Santiago, Fogo and Boavista [12, 24,25], however we found that its current distribution is wider. The exclusive presence of *An. arabiensis* in Cabo Verde can be explain by the Sahelian conditions of the archipelago, since this member of the *An. gambiae* complex is the most tolerant to aridity [26,27].

Aedes aegypti was identified in all 22 municipalities in the country. This was the second species of mosquito identified in Cabo Verde in 1930 in São Vicente. Half a century later, this species was identified on all inhabited islands except in Maio [10]. Although *Ae. aegypti* was not found in the island of Maio in the last entomological survey performed in 2007 [12], there was evidence of its presence, since the occurrence of cases of dengue

and Zika in 2009 and 2015-2016, respectively [5,28]. In this survey, *Ae. aegypti* was reported in all the municipalities, and at municipality of Maio for the first time [29].

Culex quinquefasciatus and *Cx. pipiens* are the most ubiquitous mosquitoes in the tropical and temperate regions of the globe, respectively. In Cabo Verde, these species and their hybrids have been described since 1950 [10]. Entomological surveys have reported the presence of *Cx. pipiens* s.l. in all inhabited islands [10,12] which are corroborated in this study, in which we report its presence in all 22 municipalities. Although *Cx. pipiens* s.s. was identified in Maio and Santo Antão, which corroborates previous data [10], the presence of hybrid forms in Fogo and São Nicolau can suggest its presence in these islands. This brings up to four the locations of the archipelago that are more likely to present adequate environmental conditions for a contact zone between this species and *Cx. quinquefasciatus*.

Culex quinquefasciatus was identified in all inhabited islands of Cabo Verde. When in sympatry with *Cx. pipiens*, the former was always the most abundant form (N = 195; 83.3%).

Hybrid forms of *Cx. quinquefasciatus* and *Cx. pipiens* were found for the first time in Santo Antão. The subtropical location of the archipelago is likely to present adequate environmental conditions for a contact zone between the two species, as documented in previous studies [30]. Interestingly, there is no known hybrid zone between the two sibling species in mainland Africa, which probably reflects the effect of the Sahara Desert as a geographic barrier to the distribution of both species.

A recent study conducted on Santiago island showed a high human blood index (HBI) in engorged populations of *Cx. pipiens* s.l., followed by domestic dog and chicken blood meals (personal communication). Although depending on biotic and abiotic factors,

including host-insect interaction factors, mosquito genetic traits, host availability and host density, it is generally assumed that each *taxa* has a particular biting behaviour, with *Culex pipiens* showing essentially an ornithophilic preference and *Culex quinquefasciatus* anthropophilic behaviour [31-43]. Hybrid forms can have intermediary host feeding patterns which can lead to a different role in pathogen transmission [35, 44-46].

Aedes caspius was identified in Sal, Boavista, Maio, and for the first time in São Vicente. *Aedes caspius* was found for the first time by Meira and collaborators in 1952 on the island of Sal [10,12].

Our results show *An. pretoriensis* widespread in Cabo Verde, except in Sal, where it was never reported (*Entomological Surveillance Bulletins: https://insp.gov.cv/index.php/pilar-02-laboratorio-nacional-de-saude-publica*, accessed in 27/03/2020). First detection of this species was in São Nicolau, in 1947. Later it was also reported in the islands of Boavista, São Vicente, Santo Antão, Santiago, Maio, Fogo and Brava [10].

Culex bitaeniorhynchus (formerly *Cx. ethiopicus*) was identified on the islands of Santo Antão and Santiago, where it has always been recorded, since its first identification, in 1977 [10,12]. And the *Culex tritaeniorhynchus* was found only on the island of Santiago in municipality of Tarrafal. This species was first described in the archipelago in 2011, in the municipality of Santa Cruz [14], which indicates dissemination of the species.

Culiseta longiareolata was recorded in Santo Antão, São Nicolau, Sal, Santiago, Fogo and Brava. This species was first described in 1947, in São Nicolau and Boavista, and in 1977 it was registered on all islands except Sal [8], while *Lutzia tigripes* was identified in Santo Antão, São Vicente, São Nicolau and Boavista. In Cabo Verde, this species was first reported in Maio [11], and later in Sal, Santiago and Fogo [12].

In this survey, the species *Culex perexiguus* was not detected. This species was reported for the first time in Santiago in 2010 [12], and since then no more reports of this species occurred. However, the confirmation of this species, included in the *Cx. univitattus* complex, in Cabo Verde is important, since this is a competent vector of West Nile, Sindbis, and Rift Valley fever viruses [47,48].

Conclusion

Mosquitoes are the most important group of vectors in public health, because some species are highly competent for the transmission of pathogens to humans. The knowledge of the mosquito fauna and its geographic distribution provide important data to assess the potential transmission risk and outbreak occurrence.

With this work, 10 species of mosquitoes were identified in Cabo Verde, including members of two species complexes, four of which are important vectors of human pathogens, namely: *Anopheles arabiensis*, *Aedes aegypti*, *Culex pipiens* and *Cx. quinquefasciatus*.

The only malaria vector found in Cabo Verde was *An. arabiensis* and we report for the first time its presence in Maio and São Vicente. *Aedes aegypti* was spread across the archipelago and it was reported for the first time in Maio and *Aedes caspius* was reported for the first time in São Vicente.

Hybrids forms of *Cx. quinquefasciatus* and *Cx. pipiens* were found in four islands, bringing up to four the locations of the archipelago that are more likely to these species occur in sympatry. In this survey *Cx. quinquefasciatus* was the most abundant species of the *Cx. pipiens* complex.

Timely detection of changes in species diversity provide valuable knowledge to health authorities, the scientific community and entities which may take control measures of vector populations reducing their impact on public health.

Acknowledgements

We acknowledge the National Institute of Public Health of Cabo Verde for supporting the study, and the National Program for vector control to all the Health facilities. Also, we thank the health delegates, control vector agents, drivers, and administrators that provided their support during the entomological survey. We acknowledge Jonas Antonio Lopes Gomes, from the National Institute of Public Health of Cabo Verde, for the map drawing. Special acknowledge Tomás Alves de Só Valdez for trust and support.

Authors' contributions

Design of the study, S.D.V.L. and I.B.F.V.; Collections and laboratory work, D.D.S.M., C.M.R.d.S., A.A.L.G., and I.B.F.V.; Data analyses, I.B.F.V., A.A.L.G., H.C.O., and S.D.V.L. Original Draft Preparation, I.B.F.V., A.A.L.G., H.C.O., D.D.S.M., and S.D.V.L.; Writing-Review and Editing, I.B.F.V., A.A.L.G., M., A.J.D.P., H.C.O., and S.D.V.L.; Revision and Supervision, A.J.D.P., and M.d.L.L.M.

Conflict of interest

All authors read and approved the final manuscript.

Financing

The study was made possible through the Program Investing to achieve elimination for Malaria and impact against TB and HIV in Cape Verde, Grant CPZ-Z-CCSSIDA, supported by The Global Fund to Fight AIDS, Tuberculosis and Malaria and implemented by the Coordination Committee of the Fight against AIDS of Cape Verde (CCS-SIDA).

References:

1. WHO. Global vector control response 2017–2030. Geneva: World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO. 2017.
2. WHO Study Group. Malaria vector control and personal protection. World Health Organization technical report series, 2006. 936, 1.
3. World Health Organization. World malaria report 2017. Geneva. Licence: CC BY-NC-SA 3.0 IGO.
4. Dia I, Diagne CT, Ba Y, Diallo D, Konate L, and Diallo M. Insecticide susceptibility of *Aedes aegypti* populations from Senegal and Cape Verde Archipelago. *Parasites & vectors*. 2012; 5 (1): 238.
5. Lourenço J, de Lourdes MM, Valdez T, Monteiro JR, Pybus O, and Faria RN. Epidemiology of the Zika Virus Outbreak in the Cabo Verde Islands, West Africa. *PLoS Curr*. 2018; 15:10.
6. DePina AJ, Niang EH, Andrade AJ, Dia AK, Moreira A, Faye O, et al. Achievement of malaria pre-elimination in Cape Verde according to the data collected from 2010 to 2016. *Mal J*. 2018; 17: 236.
7. DePina AJ, Andrade AJ, Dia AK, Moreira AL, Furtado UD, Baptista H, et al. Spatiotemporal Characterization and risk factor analysis of Malaria Outbreak in Cabo Verde in 2017. *Tropical Medicine and Health*. 2019; 47:3.
8. Weaver SC and Reisen WK. Present and future arboviral threats. *Antiviral research*. 2010; 85 (2): 328–345.
9. Ribeiro H, and da Cunha RH. *Guia ilustrado para a identificação dos mosquitos de Angola (Diptera: Culicidae)*. Sociedade Portuguesa de Entomologia. 1995.

10. Ribeiro H, Ramos HD, Capela RA, and Pires CA. Os mosquitos de Cabo Verde (Diptera: Culicidae): Sistemática, distribuição, bioecologia e importância medica. Junta Invest. Cient. Ultramar Estud. Ens. Dot. 1980; 135: 141.
11. Cambournac FJ, Oliveira MC, Correia A, Coutinho MA, Torinho J and Soares, AB. *Culex* (Lutzia) *tigripes* (Grandepré); Mais uma espécie nova Para Cabo Verde. Anais Instituto de Higiene e Medicina Tropical. 1984; 10(1/4): 41-46.
12. Alves J, Gomes B, Rodrigues R, Silva J, Arez AP, Pinto J, et al. Mosquito fauna on the Cape Verde Islands (West Africa): an update on species distribution and a new finding. Journal of Vector Ecology. 2010;35 (2): 307-312.
13. Harbach RE, Culverwell CL, and Kitching IJ. Phylogeny of the nominotypical subgenus of *Culex* (Diptera: Culicidae): insights from analyses of anatomical data into interspecific relationships and species groups in an unresolved tree. Systematics and Biodiversity. 2016; 15 (4): 296-306.
14. Alves J, DePina AJ, Diallo M, and Dia I. First report of *Culex* (*Culex*) *tritaeniorhynchus* Giles, 1901 (Diptera: Culicidae) in the Cape Verde Islands. Zoologia Caboverdiana. 2014; 5(1): 4-19.
15. Lourens GB, and Ferrell DK. Lymphatic Filariasis. The Nursing clinics of North America. 2019; 54(2): 181–192.
16. Sule WF, Oluwayelu DO, Hernández-Triana LM, Fooks AR, Venter M, and Johnson N. Epidemiology and ecology of West Nile virus in sub-Saharan Africa. Parasites & vectors. 2018; 11(1): 414.
17. Fonseca DM, Smith JL, Wilkerson RC, and Fleischer RC. Pathways of expansion and multiple introductions illustrated by large genetic differentiation among worldwide populations of the southern house mosquito. The American journal of tropical medicine and hygiene. 2006; 74(2): 284–289.

18. Gomes B, Alves J, Sousa CA, Santa-Ana M, Vieira I, Silva T L, et al. Hybridization and population structure of the *Culex pipiens* complex in the islands of Macaronesia. *Ecology and evolution*. 2012; 2(8): 1889-1902.
19. Instituto Nacional da Estatística de Cabo Verde. *Projeção Demográficas de Cabo Verde, 2010 – 2030*. 2013.
20. Dehghan H, Sadraei J, Moosa-Kazemi SH, Abolghasemi E, Solimani H, Nodoshan AJ, et al. A pictorial key for *Culex pipiens* complex (Diptera: Culicidae) in Iran. *Journal of arthropod-borne diseases*. 2016; 10(3): 291.
21. Azari-Hamidian S, and Harbach RE. Keys to the adult females and fourth-instar larvae of the mosquitoes of Iran (Diptera: Culicidae). *Zootaxa*. 2009; 2078(1): 1-33.
22. Scott JA, Brogdon WG, and Collins FH. Identification of single specimens of the *Anopheles gambiae* complex by the polymerase chain reaction. *The American journal of tropical medicine and hygiene*. 1993; 49(4): 520-529.
23. Smith JL, and Fonseca DM. Rapid assays for identification of members of the *Culex* (*Culex*) *pipiens* complex, their hybrids, and other sibling species (Diptera: culicidae). *The American journal of tropical medicine and hygiene*. 2004; 70(4): 339–345.
24. Cambournac FJ, Petrarca V, and Coluzzi M. *Anopheles arabiensis* in the Cape Verde archipelago. *Parassitologia*. 1982; 24(2-3): 265.
25. DePina AJ, Namountougou M, Leal SV, Varela IB, Monteiro DD, de Sousa CM, et al. Susceptibility of *Anopheles gambiae* Sl to the Insecticides in Praia, Cape Verde: A Country in the Pre-Elimination of Malaria. *Vector Biol J*. 2018; 3:2.
26. Coetzee M, Craig M, and le Sueur D. Distribution of African malaria mosquitoes belonging to the *Anopheles gambiae* complex. *Parasitol. Today*. 2000; 16: 74-77.

27. Coluzzi M, Sabatini A, della Torre A, Di Deco MA, and Petrarca V. A polytene chromosome analysis of the *Anopheles gambiae* species complex. *Science*. 2002; 298: 1415-1418.
28. DePina AJ, Sangare MB, Dia AK, Manyangadze T, Moreira AL, Seck I, et al. Clinical and Epidemiological Characterization of Dengue Outbreak in Cabo Verde in 2009-2010. *J Trop Dis* 2019; 7 (296): 2.
29. Leal S, Varela IB, Gonçalves AL, Monteiro DD, de Sousa CM, Mendonça, ML, et al. Abundance and Updated Distribution of *Aedes aegypti* (Diptera: Culicidae) in Cabo Verde Archipelago: A Neglected Threat to Public Health. *International journal of environmental research and public health*. 2020; 17(4): 1291.
30. Kothera L, Zimmerman EM, Richards CM, and Savage HM. Microsatellite characterization of subspecies and their hybrids in *Culex pipiens* complex (Diptera: Culicidae) mosquitoes along a north-south transect in the central United States. *J. Med. Entomol*. 2009; 46: 236-248.
31. Farajollahi A, Fonseca DM, Kramer LD, and Kilpatrick AM. “Bird biting” mosquitoes and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology. *Infection, genetics and evolution*. 2011; 11(7): 1577-1585.
32. Gomes AC, Silva NN, Marques GR, and Brito M. Host-feeding patterns of potential human disease vectors in the Paraíba Valley region, State of São Paulo, Brazil. *Journal of Vector Ecology*. 2003; 28: 74-78.
33. Hamer GL, Kitron UD, Brawn JD, Loss SR, Ruiz MO, Goldberg TL, et al. *Culex pipiens* (Diptera: Culicidae): a bridge vector of West Nile virus to humans. *Journal of medical entomology*. 2008; 45(1): 125-128.

34. Kilpatrick AM, Kramer LD, Jones MJ, Marra PP, and Daszak P. West Nile virus epidemics in North America are driven by shifts in mosquito feeding behavior. PLoS Biology. 2006; 4(4):e82.
35. Kilpatrick AM, Kramer LD, Jones MJ, Marra PP, Daszak P, and Fonseca DM. Genetic influences on mosquito feeding behavior and the emergence of zoonotic pathogens. The American journal of tropical medicine and hygiene. 2007; 77(4): 667-671.
36. Lehane MJ. The biology of blood-sucking in insects. Cambridge University Press. 2005.
37. Kilpatrick AM, Daszak P, Jones MJ, Marra PP, and Kramer LD. Host heterogeneity dominates West Nile virus transmission. Proc Biol Sci. 2006; 273(1599): 2327-2333.
38. Molaei G, Andreadis TG, Armstrong PM, Bueno JR, Dennett JA, Real SV et al. Host feeding pattern of Culex quinquefasciatus (Diptera: Culicidae) and its role in transmission of West Nile virus in Harris County, Texas. The American journal of tropical medicine and hygiene. 2007; 77(1): 73-81.
39. Kay BH, Boreham PFL, and Williams GM. Host preferences and feeding patterns of mosquitoes (Diptera: Culicidae) at Kowanyama, Cape York Peninsula, northern Queensland. Bulletin of Entomological Research. 1979;69(3): 441-457.
40. Rizzoli A, Bolzoni L, Chadwick EA, Capelli G, Montarsi F, Grisenti M, et al. Understanding West Nile virus ecology in Europe: Culex pipiens host feeding preference in a hotspot of virus emergence. Parasites & vectors. 2015; 8(1): 213.
41. Thiemann TC, and Reisen WK. Evaluating sampling method bias in Culex tarsalis and Culex quinquefasciatus (Diptera: Culicidae) bloodmeal identification studies. Journal of medical entomology. 2012; 49(1): 143-149.

42. Thiemann TC, Lemenager DA, Klueh S, Carroll BD, Lothrop HD, and Reisen WK. Spatial variation in host feeding patterns of *Culex tarsalis* and the *Culex pipiens* complex (Diptera: Culicidae) in California. *Journal of medical entomology*. 2012; 49(4): 903-916.
43. Turell MJ, Dohm DJ, Sardelis MR, O'guinn ML, Andreadis TG, and Blow JA. An update on the potential of North American mosquitoes (Diptera: Culicidae) to transmit West Nile virus. *Journal of medical entomology*. 2005; 42(1): 57-62.
44. Fonseca DM, Keyghobadi N, Malcolm CA, Mehmet C, Schaffner F, Mogi M, et al. Emerging vectors in the *Culex pipiens* complex. *Science*. 2004; 303(5663): 1535-1538.
45. Huang S, Hamer GL, Molaei G, Walker ED, Goldberg TL, Kitron UD, et al. Genetic variation associated with mammalian feeding in *Culex pipiens* from a West Nile virus epidemic region in Chicago, Illinois. *Vector-Borne and Zoonotic Diseases*. 2009; 9(6): 637-642.
46. Vogels CB, Fros JJ, Göertz GP, Pijlman GP, and Koenraadt CJ. Vector competence of northern European *Culex pipiens* biotypes and hybrids for West Nile virus is differentially affected by temperature. *Parasites & vectors*. 2016; 9(1): 393.
47. Jupp PG, McIntosh BM, and Blackburn NK. Experimental assessment of the vector competence of *Culex (Culex) neavei* Theobald with West Nile and Sindbis viruses in South Africa. *Trans. R. Soc. Trop. Med. Hyg.* 1986; 80: 226-230.
48. Turell MJ, Presley SM, Gad AM, Cope SE, Dohm DJ, Morrill JC, et al. Vector competence of Egyptian mosquitoes for Rift Valley fever virus. *Am. J. Trop. Med. Hyg.* 1996; 54: 136-139.

Figure 2: Geographic distribution of mosquito species in Cabo Verde (*ArcGIS* 10.5).

405 Table 1: Species composition in the islands of Cabo Verde.

Island	Species										
	<i>Ae. aegypti</i>	<i>Ae. caspius</i>	<i>An. arabiensis</i>	<i>An. pretoriensis</i>	<i>Cx. bitaeniorhynchus</i>	<i>Cx. pipiens s.s.</i>	<i>Cx. quinquefasciatus</i>	<i>Culex pipiens hybrids</i>	<i>Cx. tritaeniorhynchus</i>	<i>Cs. longiareolata</i>	<i>Lz. tigripes</i>
Santo Antão	X			X	X	x	x	x		X	X
São Vicente	X	X	X	X			x				X
São Nicolau	X			X			x	x		X	X
Sal	X	X					x			X	
Boavista	X	X	X	X			x				X
Maio	X	X	X	X		x	x	x			
Santiago	X		X	X	X		x		X	X	X
Fogo	X			X			x	x		X	
Brava	X			X			x			X	

407 Table 2. Frequencies of *Culex pipiens*, *Cx. quinquefasciatus* and their hybrids in Cabo Verde determined by the molecular assay ACE-2.
 408

Species	Localities									
	N	Maio	Boavista	Brava	Fogo	Sal	Santiago	S. Antão	S. Nicolau	S. Vicente
<i>Cx. pipiens</i>	9 (3.8)	5 (16.7)						4 (13.3)		
<i>Cx. quinquefasciatus</i>	195 (83.3)	10 (33.3)	23 (100)	30 (100)	2 (66.7)	29 (100)	30 (100)	15 (50.0)	26 (86.7)	30 (100)
Hybrids	31 (13.2)	15 (50.0)			1 (33.3)			11 (36.7)	4 (13.3)	
Total	235	30	23	30	3	29	30	30	30	30

409 *(% in brackets)

Elementos do Mapa

- Cabo Verde*
- ▲ *Aedes aegypti*
- *Aedes caspius*
- *Anopheles arabiensis*
- ◆ *Anopheles pretoriensis*
- *Culex bitaeniorhynchus*
- ✦ *Culex* hybrid forms
- *Culex pipiens* s.s.
- ▼ *Culex quinquefasciatus*
- ✦ *Culex tritaeniorhynchus*
- *Culiseta longiareolata*
- *Lutzia tigripes*

0 10 20 40 60 80 Km

Figure

Table 1: Species composition in the islands of Cabo Verde.

Island	Species										
	<i>Ae. aegypti</i>	<i>Ae. caspius</i>	<i>An. arabiensis</i>	<i>An. pretoriensis</i>	<i>Cx. bitaeniorhynchus</i>	<i>Cx. pipiens s.s.</i>	<i>Cx. quinquefasciatus</i>	<i>Culex pipiens hybrids</i>	<i>Cx. tritaeniorhynchus</i>	<i>Cs. longiareolata</i>	<i>Lz. tigripes</i>
Santo Antão	X			X	X	x	x	x		X	X
São Vicente	X	X	X	X			x				X
São Nicolau	X			X			x	x		X	X
Sal	X	X					x			X	
Boavista	X	X	X	X			x				X
Maio	X	X	X	X		x	x	x			
Santiago	X		X	X	X		x		X	X	X
Fogo	X			X			x	x		X	
Brava	X			X			x			X	

1 Table 2. Frequencies of *Culex pipiens*, *Cx. quinquefasciatus* and their hybrids in Cabo Verde determined by the molecular assay ACE-2.

Species	N	Localities								
		Maio	B. Vist a	Brav a	Fog o	Sal	Santi ago	S. Antão	S. Nicola u	S. Vicent e
<i>Cx. pipiens</i>	9 (3.8)	5 (16.7)						4 (13.3)		
<i>Cx. quinquefasciatus</i>	195 (83.3)	10 (33.3)	23 (100)	30 (100)	2 (66.7)	29 (100)	30 (100)	15 (50.0)	26 (86.7)	30 (100)
Hybrids	31 (13.2)	15 (50.0)			1 (33.3)			11 (36.7)	4 (13.3)	
Total	235	30	23	30	3	29	30	30	30	30

2 *(% in brackets)

3