1 Genomic evidence for a chemical link 2 between redox conditions and 3 microbial community composition

4 Jeffrey M. Dick^{1,2*} and Jingqiang Tan^{1*}

*For correspondence:

jeff@chnosz.net (JMD);

tanjingqiang@csu.edu.cn (JT)

5 ¹Key Laboratory of Metallogenic Prediction of Nonferrous Metals and
6 Geological Environment Monitoring, Ministry of Education, School of
7 Geosciences and Info-Physics, Central South University, Changsha 410083,
8 China; ²State Key Laboratory of Organic Geochemistry, Guangzhou Institute of
9 Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

11 **Abstract** Environmental influences on community structure are often assessed through
12 multivariate analyses in order to relate microbial abundances to separately measured
13 physicochemical variables. However, genes and proteins are themselves chemical entities; in
14 combination with genome databases, differences in microbial abundances directly encode for
15 chemical variability. We predicted that the carbon oxidation state of inferred community
16 proteomes, obtained by combining taxonomic abundances from published 16S rRNA gene
17 sequencing datasets with predicted microbial proteomes from the NCBI Reference Sequence
18 (RefSeq) database, would reflect environmental oxidation-reduction conditions in various
19 natural and engineered settings including shale gas wells. Our analysis confirms the
20 geobiochemical predictions for environmental redox gradients within and between
21 hydrothermal systems and stratified lakes and marine environments. Where they are present,
22 a common set of taxonomic groups (Gamma- and Deltaproteobacteria and Clostridia) act as
23 drivers of the community-level differences in oxidation state, whereas Flavobacteria most often
24 oppose the overall changes. The geobiochemical signal is largest for the steep redox gradients
25 associated with hydrothermal systems and between surface water and produced fluids from
26 shale gas wells, demonstrating the ability to determine the magnitude of redox effects on
27 microbial communities from 16S sequencing alone.

29 Introduction

30 How environment affects community structure is a major question for microbial ecology.
 31 Physicochemical conditions affect the energetics of metabolic reactions (e.g. *Lin et al., 2016*)
 32 and consequently the distribution of organisms with specific metabolic capabilities. In some
 33 environments such as sediments, there are well-known links between inorganic redox couples
 34 and metabolic types. However, at a global scale, salinity is thought to be the major driver of
 35 community structure (*Lozupone and Knight, 2007; Thompson et al., 2017*).

36 The elucidation of environmental drivers of community structure depends on two types
 37 of data: environmental DNA sequencing (e.g. 16S amplicon or shotgun metagenomes) and
 38 physicochemical measurements. Many studies have used multivariate techniques, particularly
 39 ordination, to relate microbial abundances determined from sequence data to measured
 40 physicochemical variables (*Ramette, 2007*). Despite the sophistication of these techniques,
 41 there are some drawbacks: the environmental factors are represented by dataset-dependent
 42 synthetic variables that can be difficult to interpret (*Ramette, 2007*), and the analyses are
 43 inherently correlative (*James and McCulloch, 1990*). A complementary approach would
 44 be to use the particular biomolecular compositions of different organisms to derive chemical
 45 metrics, which have a stable definition that enables comparisons across datasets. Such a
 46 chemical representation is also a precursor to thermodynamic models that yield quantitative
 47 predictions of the effects of the physicochemical environment on the relative abundances of
 48 different microbial organisms (*Dick and Shock, 2013*).

49 The analysis of environmental DNA sequences is most often regarded as a taxonomic
 50 classification problem: What organisms are there? However, from a geobiochemical perspective,
 51 microbial ecosystems can also be viewed as chemical systems, which prompts a different
 52 question: What are the organisms made of? One approach to answer this question is to
 53 obtain the elemental compositions of proteins coded by the genomes of different organisms.
 54 The methodological advance in this study is to combine proteins coded by whole-genome
 55 reference sequences with taxonomic profiles from 16S rRNA gene sequences in order to infer
 56 community proteomes. These inferred proteomes have a definite elemental composition and
 57 therefore provide a novel source of information about biochemical differences between microbial
 58 communities.

59 Our guiding hypothesis is that physicochemical gradients shape the chemical composition

of microbial communities. The specific predictions we make are that where an environment is more reducing, the community proteomes would have a lower average oxidation state of carbon (Z_C), and where an environment is more saline (which is a condition associated with lower water activity), the community proteomes would have a lower stoichiometric hydration state (n_{H_2O}). This choice of variables reflects the anticipation that oxidation-reduction and hydration-dehydration reactions are major types of biochemical transformations in metabolism and, by extension, at the ecosystem scale. We previously described the rationale and derivation of Z_C and n_{H_2O} applied to shotgun metagenomic data for redox gradients and salinity gradients (*Dick et al., 2019, 2020*). Here we deploy these concepts in an analysis of 16S rRNA data to test the predictions in a wider range of environments.

Results

We used data from the NCBI Reference Sequence (RefSeq) database separately and in combination with 16S-based community profiles. By processing these data (see Materials and Methods) we obtained *predicted proteomes*, which refer to proteomes for individual taxonomic groups in the RefSeq database that are based on automatic translation of genomic sequences, and *inferred community proteomes*, which represent the combination of predicted RefSeq proteomes with 16S-based microbial abundance profiles to estimate the amino acid compositions of proteomes for entire communities.

We analyzed the amino acid compositions of proteomes using two chemical metrics. Carbon oxidation state (Z_C) of an organic molecule denotes the average charge on carbon atoms required for electroneutrality given formal charges of all other atoms. Also known by other names such as nominal oxidation state of carbon (NOSC), Z_C is an established metric for describing chemical transformations of natural organic matter (*Kroll et al., 2011; Arndt et al., 2013; Visconti and de Paz, 2021*), and has also been applied to analysis of metagenomic sequences (*Fones et al., 2019; Lecoivre et al., 2021*). The equation for Z_C of proteins only requires elemental abundances (*Dick, 2014; Dick et al., 2020*).

The second chemical metric, stoichiometric hydration state (n_{H_2O}), is computed from the number of water molecules in theoretical reactions to form proteins from a set of thermodynamic components. These components are also referred to as basis species, and provide a way of projecting elemental composition into chemical species such as O_2 and H_2O . Other than the requirement that they are the minimum number of species that can be linearly combined to represent all possible elemental compositions of the primary sequence of proteins, there is no

Figure 1. Distinct chemical parameters of predicted proteomes for major taxonomic groups. (A) Archaeal, bacterial, and viral phyla with more than 500 members at all lower levels (identified by unique taxonomic IDs in the NCBI taxonomy database); (B) archaeal and bacterial phyla with more than 60 lower-level members; (C) proteobacterial classes. Large symbols are for high-level taxa (phyla in A and B; classes in C) and small outlined symbols represent lower-level taxa (classes in A and B; orders in C). Points labeled 1, 2, 3, 4, and 5 are for the euryarchaeotal classes Methanococci, Archaeoglobi, Thermococci, Halobacteria, and Nanohaloarchaea, respectively, and the point labeled 6 is for the class Clostridia in the phylum Firmicutes. The taxonomic names are taken from the current NCBI taxonomy, but the Epsilonproteobacteria have recently been reclassified as Campylobacterota (phyl. nov.) (Waite et al., 2018).

Figure 1—source data 1. Computed mean values of Z_C and n_{H_2O} and names of all taxonomic groups shown in the plot, and numbers of individual taxa used to compute the means for each group.

thermodynamic constraint on the choice of basis species. A particular set of basis species – glutamine, glutamic acid, cysteine, O_2 , and H_2O (abbreviated QEC) – was chosen to maximize the covariation between the independently computed Z_C of proteins and the number of O_2 in their theoretical formation reactions (which are both measures of oxidation state) and to reduce that between Z_C and n_{H_2O} (Dick et al., 2020). These basis species have been used to identify distinct signatures of oxidation-reduction and hydration-dehydration reactions in both metagenomic sequences and clinical proteomic data (Dick et al., 2020; Dick, 2021b).

To provide a taxonomic context for the analysis of 16S gene sequence datasets, the presentation of results begins with a visualization of the chemical metrics of the predicted proteomes of taxonomic groups in the RefSeq database.

Chemical differences among taxonomic groups

Chemical metrics of predicted proteomes for phyla with the greatest number of representative lower-level taxa in the RefSeq database are plotted in *Figure 1*. The first panel includes viruses and archaeal and bacterial phyla, and reveals that proteins in many viruses have a lower stoichiometric hydration state than most cellular organisms except for Bacteroidetes. The next panel excludes viruses to consider cellular organisms. The proteomes of organisms affiliated to Bacteroidetes have the lowest overall n_{H_2O} , and those for Crenarchaeota, Fusobacteria, Actinobacteria, and Euryarchaeota, except for the classes Haloarchaea and Nanohaloarchaea, have relatively high n_{H_2O} . In terms of Z_C , Actinobacteria, Planctomycetes, and Haloarchaea and Nanohaloarchaea within the Euryarchaeota are the groups with the most oxidized proteomes, whereas Crenarchaeota, Thermotogae, Fusobacteria, and Tenericutes have the most reduced proteomes. In the third panel, the stoichiometric hydration state distinguishes the main classes of Proteobacteria, with most orders of Alphaproteobacteria and Gammaproteobacteria at high and low n_{H_2O} , respectively, although some orders of the Alphaproteobacteria are considerably less hydrated. Notably, the proteobacterial class with the most reduced proteins is Epsilonproteobacteria. Members of this class are often identified in hydrothermal vent communities (Nakagawa *et al.*, 2005; Campbell *et al.*, 2006), and in a proposed reclassification this class has been reassigned as a new phylum named Campylobacterota (Waite *et al.*, 2018).

The chemical representation of microbial proteomes generates many hypotheses about the effects of oxidation-reduction and hydration-dehydration reactions on biochemical evolution. For instance, the highly reducing environmental conditions of sediments and hydrothermal fluids (Lovley and Klug, 1986; Lin *et al.*, 2016) may have shaped the evolutionary processes that resulted in the low Z_C of proteins in the Methanococci, Archaeoglobi, and Thermococci, which are the most reduced classes in the Euryarchaeota and among the most reduced of all microbial classes shown in *Figure 1*. In an even broader evolutionary context, the lower hydration state of viral proteomes than those of cellular organisms may not be surprising given the absence of a cytoplasmic compartment in viruses; in particular, the water content per gram of dry matter of geometrical viruses (those without a viral envelope) is considerably lower than that of *E. coli* cells (Matthews, 1975). The low n_{H_2O} of Bacteroidetes, which are often enriched in suspended particles in seawater (Fernández-Gómez *et al.*, 2013), aligns with our previous finding of lower n_{H_2O} of proteins coded by metagenomes of particle-associated compared to free-living communities (Dick *et al.*, 2020).

Some natural clustering is evident in *Figure 1*, which indicates that the chemical com-

Table 1. Sources of data used in this study.

Description	BioProject	Description	BioProject
<i>Natural environment datasets</i>			
Guerrero Negro mat	N/A (a,b)	Black Sea water	PRJNA423140
O ₂ gradient (in mat)	<i>Harris et al. (2013)</i>	O ₂ gradient	<i>Sollai et al. (2019)</i>
Yellowstone hot springs	PRJNA207095	Swiss lakes	PRJEB27579 (b)
Archaea and Bacteria	<i>Bowen De León et al. (2013)</i>	O ₂ gradient	<i>Mayr et al. (2020)</i>
ETNP water	PRJNA263621	Sansha Yongle Blue Hole	PRJNA503500
O ₂ gradient	<i>Ganesh et al. (2015)</i>	O ₂ gradient	<i>He et al. (2020)</i>
Baltic Sea water	PRJEB1245 (b,c)	<i>Shale gas-affected streams</i>	
Salinity gradient	<i>Herlemann et al. (2016)</i>	NW Pennsylvania	PRJNA394724
Lake Fryxell mat	PRJNA291280	Water and sediment	<i>Ulrich et al. (2018)</i>
O ₂ gradient (in water)	<i>Jungblut et al. (2016)</i>	Penn. State Forests (PASF)	PRJNA544240
Tibetan Plateau lakes	PRJNA294836	Water (spring and fall)	<i>Mumford et al. (2020)</i>
Salinity gradient	<i>Zhong et al. (2016)</i>	<i>Source water, fracturing fluid, and produced water</i>	
Manus Basin vents	PRJEB15554 (b)	Marcellus Shale	PRJNA229085
O ₂ and T gradient	<i>Meier et al. (2017)</i>	Pennsylvania, USA	<i>Cluff et al. (2014)</i>
Qarhan Salt Lake	PRJNA388250 (d)	Denver–Julesburg Basin	PRJNA438710
Saline and normal soils	<i>Xie et al. (2017)</i>	Colorado, USA	<i>Hull et al. (2018)</i>
Ursu Lake water	PRJNA395513	Duvernay Formation	PRJNA407226
O ₂ and salinity gradient	<i>Baricz et al. (2021)</i>	Alberta, Canada	<i>Zhong et al. (2019)</i>

a. 16S rRNA gene sequences were obtained from GenBank (JN427016 to JN539989) using the `read.GenBank()` function from R package ape (*Paradis and Schliep, 2019*). For all other datasets, sequences were downloaded from the NCBI Sequence Read Archive (*Leinonen et al., 2011*) with the BioProject accessions listed in the table. b. Only bacterial 16S primers were used in these studies; other studies include both archaeal and bacterial sequences. c. To minimize the effects of seasonality and depth, we selected samples taken in the summer from depths of not more than 20 m. d. The analysis includes samples from both surface (0–10 cm) and subsurface (15–30 cm) layers. Normal (low salinity) soil samples are from outside the Qarhan Salt Lake (Water Park, Tianjin, China).

position of proteomes of classes within a phylum tend to be more similar to each other than to classes in other phyla. An interesting exception is the much higher Z_C and lower n_{H_2O} of Haloarchaea and Nanohaloarchaea than for other euryarchaeotal classes. The lower water activity associated with hypersaline environments may be one reason for these groups to have evolved proteomes with lower n_{H_2O} . However, the high Z_C of these groups is most likely not an adaptation to more oxidizing conditions; indeed, the solubility of O₂ decreases at higher salinity (*Sherwood et al., 1991*). Instead, the proteomes of many halophiles have greater numbers of acidic residues that stabilize the three-dimensional structure of proteins in high-salt conditions; because of the oxygen atoms contained in carboxylic acid groups, this adaptation also results in higher average oxidation state of carbon of the proteins (*Dick et al., 2020*).

Microbial communities encode redox and salinity gradients

Published datasets from studies of hydrothermal systems, stratified water bodies, and microbial mats were selected to represent both local (within datasets) and global (across datasets) redox and salinity gradients. Accession numbers and literature references for the datasets analyzed here are listed in *Table 1*.

Figure 2. Inferred community proteomes from different environments have distinct chemical signatures. n_{H_2O} and Z_C were calculated for inferred community proteomes using 16S rRNA gene sequencing datasets for hydrothermal systems, seawater, hypersaline environments, and microbial mats. Sources of data are listed in *Table 1*. The plot for each dataset shows individual samples as points and the convex hull containing all the samples. The convex hulls for individual datasets are assembled in the center index plot. The gray lines here have a slope corresponding to that of the regression between n_{H_2O} and Z_C for amino acids, and therefore represent the background covariation between these metrics when n_{H_2O} is calculated using the QEC basis species (glutamine, glutamic acid, cysteine, O_2 , and H_2O) (*Dick et al., 2020*). The practical salinity values reported for the Baltic Sea (supplementary information of *Herlemann et al., 2016*) have no units.

Figure 2-source data 1. Computed values of Z_C and n_{H_2O} , sample names, and SRA Run accessions used to make the plots.

As visualized in *Figure 2*, local redox gradients represented by stratified water bodies including the Black Sea (*Sollai et al., 2019*) and Lake Fryxell in Antarctica (*Jungblut et al., 2016*), submarine vents in the Manus Basin (*Meier et al., 2017*), and within the Guerrero Negro microbial mat (*Harris et al., 2013*) all provide evidence that carbon oxidation state is aligned with environmental redox conditions. Z_C is locally lower in the deep euxinic water of the Black Sea and anoxic water of Lake Fryxell, lower in the hotter water samples for the Manus Basin (which have greater input of reduced hydrothermal fluids), and lower just below the surface of the hypersaline Guerrero Negro microbial mat, which experiences a sharp oxygen gradient during the day (*Harris et al., 2013*). The Lake Fryxell dataset is also for mat samples, but the environmental gradient here is between relatively oxygenated shallow water and anoxic deeper water (*Jungblut et al., 2016*).

Inferred proteomes of archaeal sequences from alkaline hot spring communities in Yellowstone National Park (*Bowen De León et al., 2013*) have higher n_{H_2O} compared to bacterial sequences. This is aligned with expected phylogenetic differences, since the proteomes of major archaeal groups detected in these samples, including Crenarchaeota and Euryarchaeota (*Bowen De León et al., 2013*), generally have higher n_{H_2O} than bacterial proteomes (*Figure 1B*). Similarly, fauna samples in the Manus Basin dataset have lower n_{H_2O} than the majority of fluid and rock samples, but in this case no archaeal sequences were reported, so differences in bacterial abundances are the reason for the observed chemical differences.

The Yellowstone and Manus Basin datasets represent hydrothermal systems that emit highly reduced fluids. In the index plot in the center of *Figure 2*, it can be seen that these datasets are distributed toward lower Z_C than for the lake and seawater environments, thereby demonstrating that Z_C provides a signature of oxidation-reduction conditions on a global scale.

Both the Baltic Sea (*Herlemann et al., 2016*) and saline soils in the Qarhan salt lake (*Xie et al., 2017*) exhibit decreasing n_{H_2O} with greater salinity; we previously found a similar trend for proteins inferred from shotgun metagenomic sequences for various marine and freshwater environments (*Dick et al., 2020*). However, the trends for the Qarhan salt lake soils and Tibetan Plateau lake datasets (*Zhong et al., 2016*) are more strongly dominated by increasing Z_C in hypersaline conditions.

Many water bodies around the world develop vertical redox gradients as a result of microbial respiration of organic matter derived from the photic zone that leads to oxygen depletion with depth. Besides the Black Sea, we analyzed data for permanently stratified lakes in Switzerland (*Mayr et al., 2020*), the oxygen minimum zone of the Eastern Tropical North Pacific (ETNP)

Figure 3. Lower carbon oxidation state is tied to oxygen depletion in water columns. All sites except for ETNP and station C4 outside the Blue Hole are permanently stratified. Z_C values were calculated for inferred community proteomes using published microbial 16S rRNA gene sequences (see Table 1). Oxygen concentrations were taken from the same publications, except for locations inside and outside the Blue Hole (Xie *et al.*, 2019). For the Blue Hole, ratios of nitrate to nitrite (NO_3^- / NO_2^-) are also plotted based on NO_3^- and NO_2^- concentrations reported by Xie *et al.* (2019). No Z_C value is shown for 1 m depth in Ursu Lake in April 2016 because fewer than 200 sequences remained for this sample after all sequence processing steps.

Figure 3—source data 1. Computed values of Z_C , chemical measurements taken from the literature, sample names and depths, and SRA Run accessions used to make the plots.

(*Ganesh et al., 2015*), the Sansha Yongle Blue Hole in the South China Sea (*He et al., 2020*), and Ursu Lake in Central Romania (*Baricz et al., 2021*). At each location, the Z_C of the inferred community proteomes generally decreases with depth (*Figure 3*). At the ETNP, the Z_C decreases strongly with depth in the free-living communities (0.2–1.6 μm size fraction), but to a lesser extent in particle-associated communities (1.6–30 μm size fraction). This might reflect environmental microniches and cell-cell interactions that to some extent reduce the sensitivity of these communities to external redox conditions.

Although Z_C trends are dominated by oxygen gradients, a closer look suggests that other elemental redox systems may also contribute to the observed trends. There is a moderate rise in Z_C in the deepest (euxinic) layers of the Black Sea. Likewise, the low- to high-sulfide layers of the Guerrero Negro mat (below 3 mm) (*Harris et al., 2013*) have higher Z_C than the minimum at 3 mm (*Figure 2*). Therefore, despite the absence of oxygen, Z_C appears to rise locally in some sulfide-rich environments. At a non-stratified location outside the Sansha Yongle Blue Hole, the water remains oxygenated at depth. In the upper 100 meters, the profile of Z_C has a “C” shape, but maintains relatively high values at greater depths. In this case, the overall higher Z_C with depth appears to be more closely associated with the ratio of nitrate to nitrate than with O_2 concentration.

Data from a study of Ursu Lake in Central Romania from summer 2015 to spring 2016 (*Baricz et al., 2021*) provide an opportunity to examine the temporal variability of oxygen concentration and chemical metrics of the microbial communities. At all times, both O_2 and Z_C show an overall decline with depth. The profile of Z_C exhibits a broad maximum at around 2 m depth in November that becomes narrower and deeper through the winter and spring. The development of the Z_C maximum precedes that of an O_2 maximum in February, and the two profiles exhibit a remarkable meter-scale correspondence in April.

Physicochemical signal is strongest at high taxonomic levels

Not only the abundances but also the proteomic composition of individual taxonomic groups contribute to the overall chemical differences between microbial communities. To assess the chemical differences along redox and salinity gradients in more detail, the abundances and calculated Z_C or $n_{\text{H}_2\text{O}}$ of major taxonomic groups at the domain, phylum, class, and genus levels are plotted in *Figure 4* for the Manus Basin and Baltic Sea. Because the lowest-level taxonomic classification for each sequence was used for mapping to the NCBI taxonomy to generate the inferred microbial proteomes (see Materials and Methods), within-group variation of Z_C and $n_{\text{H}_2\text{O}}$ is possible above the genus level and is represented in *Figure 4* by sloping

Figure 4. Changes of abundance and chemical composition at different taxonomic levels. Symbols represent taxon abundance-weighted mean values of Z_C for subsets of samples with relatively high and low temperature as a proxy for reducing and oxidizing conditions ($N = 4$ and 28 , respectively) from the Manus Basin or n_{H_2O} for subsets of samples with high and low salinity from the Baltic Sea ($N = 19$ and 18); environmental criteria used to allocate samples to subsets are shown in the plot. The leftmost plots represent all sequences classified by the RDP Classifier and mapped to the NCBI taxonomy at the domain level. In subsequent plots, sequences classified at lower taxonomic levels were combined to calculate the percentage abundance and Z_C or n_{H_2O} for each taxonomic group whose sequences make up at least 2% (for phylum or class) or 1% (for genus) of the total number of sequences from all samples. Percentages in the plot titles indicate the fraction of the whole community represented by groups shown in the plot. Arrows connect the same taxonomic group in different sample subsets and point to samples with higher temperature (Manus Basin) or lower salinity (Baltic Sea). Abundance-weighted means for all taxonomic groups shown in each plot are indicated by dashed lines for high temperature or low salinity samples and dotted lines for low temperature or high salinity samples. All taxonomic names are taken from the output of the RDP Classifier. Campilobacterota is probably a misspelling of the phylum name Campylobacterota (Waite et al., 2018).

lines.

At the phylum level, the high-temperature samples in the Manus Basin are associated with greater numbers of Aquificae and Campylobacterota (formerly Epsilonproteobacteria) and fewer Proteobacteria. These groups have relatively low and high Z_C , respectively, which to a large extent explains the chemical difference at the whole-community level (leftmost plot). However, the campylobacterotal sequences themselves are affiliated with organisms whose proteomes have lower Z_C in the higher-temperature samples. Therefore, the whole-community chemical differences are due to both differential taxonomic abundances at the phylum level, which have the largest effect, as well as differential abundances within phyla. A similar finding applies to the major classes; at this level it is apparent that the proteobacterial contribution is mainly due to lower numbers of Gammaproteobacteria. The differential abundances of the identified genera yield a small Z_C difference between hotter and cooler fluids in the same direction as the whole-community trend.

Analogous reasoning can be used to interpret the trends in the Baltic Sea. The relatively high n_{H_2O} in low-salinity samples is mostly controlled by an increase in Actinobacteria. In contrast, Proteobacteria become less abundant at lower salinity, which to some extent counteracts the n_{H_2O} rise, but the within-group variation of Alphaproteobacteria, Betaproteobacteria, and Gammaproteobacteria is toward higher n_{H_2O} at lower salinity. The genus-level assignments suggest an opposite trend (higher n_{H_2O} at higher salinity), but this is less likely to represent the actual differences because the low classification rate to the genus level (37%; see *Table S1*) together with the 1% abundance cutoff for genera in *Figure 4* results in a low fraction of assignments represented at this level (23%).

Application to datasets for shale gas operations

During hydrocarbon extraction from unconventional reservoirs (shale gas), hydraulic fracturing fluid is injected into shale formations to create extensive horizontal fracture networks that improve hydrocarbon recovery. The injected fluid mixes and reacts with natural formation waters and the fractured rock surfaces. Water that returns to the surface is initially referred to as flowback and later as produced water during the hydrocarbon production stage of the well (*Khan et al., 2021*). Chemical oxidants are commonly added to the injected fracturing fluid; these additives enhance mineral dissolution and can also have biocidal effects. Even without such additives, fracturing fluids generally consist of large amounts of water from surface sources, which makes them highly oxidized compared to the reducing conditions in organic-rich shale.

Changes in the chemistry and biology of streams that are in proximity to or may be affected

249 by shale gas operations are an important issue for environmental assessments of shale gas
250 operations. The chemistry of affected surface streams is thought to reflect the possible input
251 of methane and/or chemical additives in fracking fluid from nearby wells, though the overall
252 strength of these associations has been debated (*Mumford et al., 2020*).

253 Other authors have noted that little quantitative information is available about the changes
254 in redox conditions of flowback and produced fluid over time (*Mouser et al., 2016; Liden
255 et al., 2019*). In one case, oxygen was found to be rapidly depleted in flowback and produced
256 waters in the Duvernay Formation in Alberta, Canada (*Zhong et al., 2019*). In a study on
257 the Marcellus Shale in Pennsylvania, USA, the abundances of S-bearing organic compounds
258 determined from FT-ICR-MS measurements exhibited a decrease in carbon oxidation state
259 compared to injected fluids (*Luek et al., 2019*). Furthermore, oxidation-reduction potential
260 (ORP) measured using a multiprobe is lower (more reducing) in groundwater samples with
261 higher concentrations of CH₄ (*LeDoux et al., 2016*). Therefore, we predicted that inferred
262 community proteomes in produced water would be shifted toward a more reduced state
263 compared to source water. A related prediction is that putative hydrocarbon input to surface
264 streams has a similar reducing effect, but the effect may be smaller because of the greater
265 extent of dilution.

266 Analysis of 16S sequence data for water from streams in Northwestern Pennsylvania (*Ulrich
267 et al., 2018*) shows that both Z_C and n_{H_2O} are lower in streams affected by Marcellus Shale
268 activity compared to unaffected streams (*Figure 5A*). Smaller differences, but in the same
269 direction, are found for stream sediments in the same study and for stream water in another
270 study (*Mumford et al., 2020*) (*Figure 5B*). Much larger shifts, toward lower Z_C and
271 higher n_{H_2O} , occur in flowback and produced waters compared to source waters and injected
272 fracturing fluids. This trend is evident for not only the Marcellus Shale (*Cluff et al., 2014*)
273 but also the Denver–Julesburg Basin in Colorado, USA (*Hull et al., 2018*) and the Duvernay
274 Formation (*Zhong et al., 2019*) (*Figure 5C* and *Figure 5D*). Notably, the magnitudes of
275 the Z_C differences in the Denver–Julesburg Basin and Duvernay Formation are the largest
276 among all datasets considered in this study, and the Marcellus Shale has a larger negative
277 ΔZ_C compared to all datasets except for the Manus Basin hydrothermal vents (*Figure 6*).
278 Therefore, highly reducing conditions associated with subsurface or hydrothermal systems are
279 manifested as great differences in carbon oxidation state of inferred community proteomes,
280 whereas oxygen gradients in water bodies exhibit a smaller ΔZ_C .

Figure 5. Decreased carbon oxidation state of inferred proteomes for communities affected by shale gas extraction. (A) Water samples from streams affected by Marcellus Shale activity (MSA+) and non-affected streams (MSA-) in Northwestern Pennsylvania (Ulrich *et al.*, 2018); star-shaped symbols represent group means. **(B)** Mean values for sample groups in various studies of streams in Pennsylvania: Northwestern Pennsylvania (water and sediment samples) (Ulrich *et al.*, 2018) and Pennsylvania State Forests (PASF; water samples in spring and fall) (Mumford *et al.*, 2020). **(C)** Injected fluids and produced water from a hydraulically fractured well in the Marcellus Shale (Cluff *et al.*, 2014); star-shaped symbols represent group means. **(D)** Mean values for samples of produced water compared to injected fluids or source water for hydraulically fractured wells in the Marcellus Shale (Cluff *et al.*, 2014), Denver-Julesburg Basin (Hull *et al.*, 2018), and Duvernay Formation (Zhong *et al.*, 2019). Numbers of runs used for calculating group means are shown next to points in (B) and (D); for the Marcellus Shale, there are two runs for each biological sample. In this figure, data for the samples collected in 2014 for NW PA water and sediment (Ulrich *et al.*, 2018) are used; in Figure 6 water samples from all available years (2012 to 2016) are used. **Figure 5-source data 1.** Computed values of Z_c and n_{H_2O} , SRA Run accessions, and sample names and types (e.g. MSA+ or MSA-, or source or produced water) used for calculating group means.

Discussion

The main finding of this study is that the oxidation state of inferred community proteomes decreases in more reducing conditions at global and local scales. Closer examination of selected datasets (*Figure 4*) indicates that the whole-community chemical differences are mainly associated with changes in abundances of particular phyla and that within-group variation of particular classes is often in the same direction, indicating that physicochemical conditions shape microbial communities at multiple taxonomic levels.

The results here are consistent with our earlier analysis of shotgun metagenomic data that showed lower Z_C of proteins in Yellowstone hot springs and the Lost City hydrothermal vents compared to ambient seawater (*Dick et al., 2019*). Similarly, *Lecoeuvre et al. (2021)* analyzed shotgun metagenome sequences from the newly discovered Old City hydrothermal field. The mean value they reported for Z_C of proteins (-0.168) lies within the range for hydrothermal systems obtained here (*Figure 2*) and is lower than all but the most reduced community proteomes inferred for anoxic waters (*Figure 3*). Therefore, Z_C values calculated using inferred community proteomes (this study) or from shotgun metagenomic sequences (*Dick et al., 2019; Lecoeuvre et al., 2021*) appear to be commensurate.

In our previous analysis of shotgun metagenomic data, we found conflicting trends for Z_C in oxygen minimum zones (*Dick et al., 2019*). We speculated that preferential degradation of low-GC content extracellular DNA by heterotrophs could leave behind resistant genes that are more likely to undergo horizontal gene transfer; the putative enrichment of AT-rich genes would be manifested by higher Z_C of the inferred proteins (*Dick et al., 2019*). It has also been proposed that horizontal gene transfer could hinder the use of shotgun sequences for accurate taxonomic identification (*Tessler et al., 2017*), although we note that our previous analysis of Z_C of shotgun sequences was not based on taxonomic assignments. In contrast to the results for shotgun metagenomes, the present study uncovered a strong signal of decreasing Z_C with depth in multiple 16S rRNA datasets for stratified water bodies.

The highly saline produced waters from many shales converge toward a common profile dominated by the halophilic and anaerobic *Halanaerobium* (*Mouser et al., 2016*), but in the Denver–Julesburg Basin *Thermoanaerobacter*, which has similar metabolic capabilities, is present instead (*Hull et al., 2018*). The predicted RefSeq proteomes of these groups have Z_C values of -0.199 and -0.227, respectively; the very low oxidation states of these abundant groups, which are both members of Clostridia (see *Figure 1B*) accounts for much of the decrease in Z_C in produced fluids. Notably, hypersaline conditions in other environments

including lakes and soils are characterized by relatively high Z_C (this study and *Dick et al.*, 2020), so the finding of an opposite trend for shale gas wells strengthens the interpretation that reducing conditions are a primary driver of community structure in produced water.

More work is needed to derive a chemical metric that captures the relationship between community composition and salinity. Because of osmotic forces, higher salinity should have a dehydrating effect, but this prediction is not supported by the increase of n_{H_2O} of inferred community proteomes that is observed for produced fluids. Higher n_{H_2O} may be intrinsically linked to lower Z_C as a result of the background correlation between these metrics when n_{H_2O} is calculated using the QEC basis species (gray lines in *Figure 5A* and *Figure 5C*; see also *Dick et al.*, 2020). However, the relations between Z_C and n_{H_2O} are not universally dictated by the basis species, as shown by the positive slopes between datasets for stream communities in *Figure 5B*.

At this point it is not possible to predict with this method which taxa are actually present in a community. Nevertheless, analysis of the contribution of individual classes to the differences in Z_C within datasets reveals some provocative patterns for these taxa that are present (*Figure 6*). In all datasets where they make a major contribution to the oxidation-reduction differences, the Gammaproteobacteria, Deltaproteobacteria, and Clostridia contribute to lower the overall Z_C of the communities in more reducing environments, due to either changes in abundance, within-group Z_C , or both (see *Figure 6–Figure Supplement 1* for a depiction of the values used for these calculations). These groups can therefore be regarded as the microbial drivers of the environmental shaping by redox gradients. In contrast, the change in abundance of Flavobacteriia opposes the community-level ΔZ_C for many datasets; a similar trend is also apparent for Cytophagia, which is another group within the Bacteroidetes, in some other datasets. These classes have relatively reduced proteomes (i.e. more negative Z_C), but they tend to be more abundant in more oxidizing environments, which explains their opposing contribution (*Figure 6–Figure Supplement 1*). Therefore, these classes can be described as detracting from the shaping of various communities by local redox gradients. To what extent the phenotypic traits of Bacteroidetes, known for degradation of high molecular weight organic matter (*Fernández-Gómez et al.*, 2013), may underlie the unique behavior of these groups remains unknown; whether these trends could be related to the very low n_{H_2O} of their proteomes (*Figure 1*) is also an open question.

Our findings for shale gas systems suggest that in addition to salinity, the steep redox gradient between the surface (oxidizing) and subsurface (reducing) is a primary factor that

Figure 6. Contributions of taxonomic classes to the difference in carbon oxidation state of inferred community proteomes between relatively oxidized and reduced communities.

Samples in each dataset were categorized into oxidized and reduced subsets as indicated in the dataset descriptions; numbers of samples in each category are given in parentheses. For each dataset, colored boxes indicate classes that are at least 2% abundant across all included samples. Inferred community proteomes for the oxidized and reduced subsets of samples were aggregated and used to calculate values of ΔZ_C . The color scale represents the contribution of each class to the overall ΔZ_C , calculated using Equation 1. Positive percentages mean that the change in abundance and/or Z_C of the class itself is in the same direction as the overall ΔZ_C (i.e. more negative in reducing conditions). Negative percentages indicate contributions by classes that are in the opposite direction. The color scale is limited to percentages in the range [-100, 100]; values > 100 are indicated by triangles inside the boxes. The sum of percentages for each dataset is 100. Datasets representing redox gradients were selected from Table 1 and are ordered as in the table. Datasets for NW Pennsylvania sediment and PASF Streams (fall) were omitted because they have a very small ΔZ_C at the class level. Classes are grouped by phylum but their ordering is otherwise arbitrary. Abbreviations: FW – flowback water; IF – injected fluids; PW – produced water; SW – source water. MSA+, MSA-, highest, and lowest are terms used by previous authors to describe the level of disturbance of streams from unconventional hydrocarbon extraction (Ulrich et al., 2018; Mumford et al., 2020). Because of the different sample depths and O_2 profiles of the Swiss lakes (Figure 3), the deepest and shallowest samples from both Lake Zug and Lake Lugano were selected.

Figure 6–Figure supplement 1. Plots of abundance and Z_C for taxonomic classes in oxidized and reduced sample subsets.

347 shapes the structure of microbial communities. Other authors have commented on the dearth
348 of redox and oxygen measurements in samples collected from black shale well sites (*Mouser*
349 *et al.*, 2016), and a lack of such measurements is also apparent in the USGS National
350 Produced Waters Geochemical Database (*Blondes et al.*, 2018). Of the 114943 records in
351 the database, there are only 66 with measurements of dissolved oxygen (O₂) or ORP, and these
352 are only for conventional or geothermal wells. The multivariate analyses performed in previous
353 studies (*Cluff et al.*, 2014; *Hull et al.*, 2018; *Zhong et al.*, 2019) identified variables
354 including dissolved inorganic and organic carbon, salinity, pH, and temperature as major
355 environmental drivers of the differences in microbial community composition. Because oxygen
356 or redox measurements of produced waters from unconventional wells were not available, those
357 analyses could not have been used to identify an association between microbial communities
358 and oxidation-reduction state of the fluids. However, such an association is exactly what is
359 predicted by the chemical analysis in this study.

360 In spite of the oxygenation of injected fluids, subsurface conditions are likely to be anoxic
361 (*Zhong et al.*, 2019), so other types of measurements should be considered for monitoring
362 the redox state of produced water. For instance, the USGS database cites a study from
363 2009 that gives both NO₃⁻ and NO₂⁻ measurements in injected and produced water from
364 the Marcellus Shale (*Hayes*, 2009). Monitoring the dynamics of N species in conjunction
365 with biological sampling would give further insight into the extent of nitrate reduction in the
366 subsurface (*Orcutt et al.*, 2011; *Cluff et al.*, 2014) and could also serve as a proxy for
367 *in situ* redox conditions through a metric such as the NO₃⁻/NO₂⁻ ratio (*Duchazeau et al.*,
368 2014). ORP is another measure of redox conditions; however, they are generally considered
369 more difficult to interpret because the electroactivity of chemical species in the sample is
370 affected by kinetic barriers that affect the potential reading but are not readily quantified.
371 Nevertheless, ORP measurements can yield information about electrochemical reactions that
372 are relevant to microbial growth, especially if the measurements are made continuously in time
373 (*Markelova et al.*, 2017).

374 By leveraging the chemical information contained in genomic sequences, it is possible to
375 achieve a broader view of the coupling between inorganic and organic oxidation-reduction
376 reactions that is essential for all ecosystems (*Burgin et al.*, 2011; *Orcutt et al.*, 2011).
377 Analysis of inferred community proteomes indicates that redox gradients have the strongest
378 effect on microbial communities in produced fluids, followed by hydrothermal environments
379 and oxygen gradients in stratified water bodies. The results suggest that more comprehensive

monitoring of dissolved oxygen concentrations or other redox indicators should be used to better characterize the responses of microbial communities in produced water and streams affected by unconventional hydrocarbon extraction, and ecosystems in general.

Materials and Methods

Data sources, processing, and classification

16S rRNA gene sequences were downloaded from the NCBI Sequence Read Archive (SRA) except for the Guerrero Negro microbial mat sequences (*Harris et al., 2013*), which were obtained from GenBank. The processing pipeline consisted of merging of paired-end reads, length and quality filtering, removal of singletons, subsampling, chimera removal, and taxonomic classification. VSEARCH version 2.15.0 (*Rognes et al., 2016*) was used to merge Illumina paired-end reads; for some datasets with low-quality reverse reads, only the forward reads were used as in previous studies (*Ulrich et al., 2018*). For Illumina datasets, quality and length filtering were done with the options `-fastq_maxee_rate 0.005` (i.e. maximum one expected error for every 200 bases) and `-fastq_truncflen` with a length value depending on the specific dataset. For 454 datasets, where reads are generally longer but have more variable length, quality and length filtering were done with `-fastq_truncqual 15 -fastq_minlen 200 -fastq_maxlen 600`; also, the option `-fastq_stripleft 18` was used to remove adapter sequences. Sequence processing statistics and additional details are given in *Table S1*.

After filtering, the remaining sequences for all samples in each dataset were pooled and singletons (sequences that appear exactly once, but not including subsequences of other sequences) were removed. Then, samples were subsampled to a depth of 10000 sequences; samples with fewer than 10000 sequences were not affected. The subsampling reduces the processing time for chimera detection, which is the longest step in the pipeline, and retains enough sequences for classifying the major taxonomic groups in the communities. Reference-based chimera detection was performed using the VSEARCH command `-uchime_ref` with the SILVA 138.1 SSURef NR99 database (*Quast et al., 2012*). Sequences identified as chimeras or borderline chimeras were removed.

The remaining sequences were processed with the Ribosomal Database Project (RDP) (*Cole et al., 2014*) Classifier version 2.13 (*Wang et al., 2007*) with the provided training set (RDP 16S rRNA training set No. 18 07/2020). The sequence classifications for all samples in each dataset were merged using the RDP Classifier command `merge-count`.

411 RefSeq proteomes

412 Predicted protein sequences were obtained from the RefSeq database release 206 (2021-05-21)
 413 (*O’Leary et al., 2016*) for all 49448 bacterial, archaeal and viral taxa identified by a unique
 414 taxonomic ID (taxid). For each taxid, the available taxonomic names at ranks of superkingdom,
 415 phylum, class, order, family, genus, and species were parsed from the current (as of the RefSeq
 416 release date) NCBI taxonomy files, and the amino acid compositions of all available protein
 417 sequences were summed and divided by the number of proteins to generate the average amino
 418 acid composition of the predicted proteome. To calculate the amino acid compositions for
 419 taxonomic groups at genus and higher levels, those for each taxon within this group, including
 420 lower levels, were summed then divided by the number of taxa. The number of proteomes,
 421 corresponding to the number of taxonomic groups at each level, is 4788 (genus), 763 (family),
 422 303 (order), 140 (class), 78 (phylum), and 3 (superkingdom).

423 Taxonomy mapping

424 To infer the amino acid compositions of communities from 16S sequencing data, RDP classifi-
 425 cations at only the root or domain level were first omitted because they provide very little
 426 taxonomic resolution. Sequences assigned to RDP class- and family-level name Chloroplast or
 427 genus-level names Chlorophyta and Bacillariophyta were also discarded because they do not
 428 fall within the archaeal and bacterial taxonomy used by NCBI. All remaining classifications
 429 were retained for attempted mapping to the NCBI taxonomy. The lowest-level classification
 430 returned by the RDP classifier for each sequence was used; the classification rate to genus
 431 level varies widely among datasets (*Table S1*) so higher-level classifications were used where
 432 needed.

433 In general, mapping from the RDP Classifier to the NCBI taxonomy was performed
 434 by text matching of both the taxonomic rank and name. Some particular mappings were
 435 used to improve the representation of common taxa in the datasets. The RDP phylum
 436 Cyanobacteria/Chloroplast, class Planctomycetacia, and genus *Escherichia/Shigella* were
 437 mapped to the NCBI phylum Cyanobacteria, class Planctomycetia, and genus *Escherichia*,
 438 respectively. The RDP order Rhizobiales was mapped to the NCBI order Hyphomicrobiales
 439 (*Hördt et al., 2020*). The RDP taxon Spartobacteria genera *incertae sedis*, which is
 440 relatively abundant in the Baltic Sea (*Herlemann et al., 2016*), was mapped to NCBI class
 441 Spartobacteria. The RDP taxon Subdivision3 genera *incertae sedis*, which was identified here in
 442 some stream datasets (*Mumford et al., 2020*) was mapped to NCBI family Verrucomicrobia

subdivision 3. The RDP taxon Marinimicrobia genera *incertae sedis*, which was identified in this study in some deep ocean datasets (*Ganesh et al., 2015; Meier et al., 2017*), was mapped to NCBI species Candidatus Marinimicrobia bacterium, which is the only representative of the Candidatus Marinimicrobia phylum in the RefSeq database. Among Acidobacteria, which are fairly abundant in river water and sediment, the RDP genus-level classifications Gp1 and Gp6 were mapped to NCBI genera *Acidobacterium* and *Luteitalea*, respectively, which are members of Acidobacteria subdivisions 1 and 6 (*Eichorst et al., 2018*). The RDP genus-level cyanobacterial groups GpI, GpIIa, and GpVI were mapped to the NCBI genera *Nostoc*, *Synechococcus*, and *Pseudanabaena*, and the RDP taxon Family II was mapped to the family Synechococcaceae; these mappings are based on names of members of these groups given in Bergey's Manual (*Castenholz et al., 2001*), although the mappings are necessarily imperfect because of inconsistencies with the NCBI taxonomy.

Any other RDP classifications whose rank and name could not be matched to the NCBI taxonomy were removed from the subsequent calculations. Note that sequences were mapped using the lowest available taxonomic level in the output of the RDP Classifier; this is why the phylum Campilobacterota (the name output by the RDP Classifier) is represented in *Figure 4* although no explicit mapping to the NCBI name Campylobacterota was done. Across all datasets, a median of 96.2% of RDP classifications at all levels from genus to phylum were mapped to the NCBI taxonomy (*Table S1*). The lowest classification rate is for a dataset for produced water from shale gas wells (*Hull et al., 2018*), in which the genus *Cavicella* makes up 27% of the RDP classifications but has no counterpart in the available RefSeq proteomes (*Table S2*).

Analysis of chemical composition

Values of n_C , Z_C and n_{H_2O} were computed for each taxonomic group in RefSeq whose amino acid compositions were obtained as described above, using the `ZCAA()` and `H2OAA()` functions in the canprot package (*Dick, 2021b*) and a modified function to calculate n_C (number of carbon atoms). For individual or aggregated samples, the RDP counts for each mapped taxon were multiplied by n_{H_2O} of that taxon, summed, and divided by the total counts to obtain the abundance-weighted mean n_{H_2O} for the inferred community proteome. For Z_C , the mean was additionally weighted by carbon number as described previously (*Dick et al., 2020*): $Z_C = \sum n_i n_{C,i} Z_{C,i} / \sum n_i n_{C,i}$, where n_i , $n_{C,i}$ and $Z_{C,i}$ are the count, carbon number, and carbon oxidation state of the i th taxon. Because Z_C is a per-carbon quantity and n_{H_2O} values are per-residue, comparisons can be made between proteomes with different average protein length.

To compare subsets of samples within datasets, RDP classifications for all samples in each subset were first aggregated, then Z_C and n_{H_2O} were calculated from the aggregated taxon counts. The percent contribution of the i th taxon ($\Delta Z_C\%$) to the difference in Z_C between sample subsets “o” and “r” (for oxidized and reduced) was calculated from

$$\Delta Z_C\% = 100 \times \left[\frac{n_{i,r}}{\sum n_{i,r}} (Z_{C,i,r} - Z_{C,o}) - \frac{n_{i,o}}{\sum n_{i,o}} (Z_{C,i,o} - Z_{C,o}) \right] \quad (1)$$

where $n_{i,o}$ and $n_{i,r}$ are the abundances of the i th taxon in the oxidized and reduced subsets, $Z_{C,i,o}$ and $Z_{C,i,r}$ are the carbon oxidation states of the i th taxon in the oxidized and reduced subsets, and $Z_{C,o}$ is the carbon oxidation state of the oxidized subset. The value of $Z_{C,i}$ was assumed to be constant for any taxon that was present in one subset but not the other. Values of $\Delta Z_C\%$ were visualized using the R package `plot.matrix` (*Klinke, 2021*).

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

JMD: Conceptualization, Software, Formal analysis, Writing; JT: Conceptualization, Funding acquisition.

Funding

This research was supported by the National Natural Science Foundation of China (grant nos. 72088101 and 41872151 to JT) and the State Key Laboratory of Organic Geochemistry (grant no. SKLOG-201928 to JMD).

Data Availability Statement

All data analyzed in this study were obtained from public databases: NCBI RefSeq (*O’Leary et al., 2016*) and SRA (*Leinonen et al., 2011*); the BioProject accession numbers used to download the SRA data are listed in Table 1. The JMDplots package version 1.2.8 deposited at Zenodo (*Dick, 2021a*) has the scripts used for processing RefSeq proteins and 16S rRNA gene sequences (in the directories `extdata/refseq` and `extdata/chem16S`, respectively) and processed data files including amino acid compositions of taxa compiled from

the RefSeq database (`extdata/refseq/protein_refseq.csv.xz`), chemical metrics calculated for taxonomic groups in RefSeq (`extdata/chem16S/taxon_metrics.csv`), and RDP Classifier results (`extdata/geo16S/RDP`). All figures were made using R (*R Core Team, 2021*) with data files and code provided in the JMDplots package; the `geo16S.Rmd` vignette in the package runs the functions to make each of the figures and the Source Data files. The compiled vignette is available in an HTML file in the package and can also be viewed at <https://chnosz.net/JMDplots/doc/geo16S.html> (accessed on 2021-09-02).

References

- Arndt, S., Jørgensen, B. B., LaRowe, D. E., Middelburg, J. J., Pancost, R. D., and Regnier, P. (2013). Quantifying the degradation of organic matter in marine sediments: A review and synthesis. *Earth-Science Reviews*, 123:53–86.
- Baricz, A., Chiriac, C. M., Andrei, A.-Ş., Bulzu, P.-A., Levei, E. A., Cadar, O., Battes, K. P., Cîmpean, M., Şenilă, M., Cristea, A., Muntean, V., Alexe, M., Coman, C., Szekeres, E. K., Sicora, C. I., Ionescu, A., Blain, D., O'Neill, W. K., Edwards, J., Hallsworth, J. E., and Banciu, H. L. (2021). Spatio-temporal insights into microbiology of the freshwater-to-hypersaline, oxic-hypoxic-euxinic waters of Ursu Lake. *Environmental Microbiology*, 23(7):3523–3540.
- Blondes, M. S., Gans, K. D., Engle, M. A., Kharaka, Y. K., Reidy, M. E., Saraswathula, V., Thordsen, J., Rowan, E. L., and Morrissey, E. A. (2018). U.S. Geological Survey National Produced Waters Geochemical Database (ver. 2.3, January 2018). Data Release, U. S. Geological Survey.
- Bowen De León, K., Gerlach, R., Peyton, B. M., and Fields, M. W. (2013). Archaeal and bacterial communities in three alkaline hot springs in Heart Lake Geyser Basin, Yellowstone National Park. *Frontiers in Microbiology*, 4:330.
- Burgin, A. J., Yang, W. H., Hamilton, S. K., and Silver, W. L. (2011). Beyond carbon and nitrogen: How the microbial energy economy couples elemental cycles in diverse ecosystems. *Frontiers in Ecology and the Environment*, 9(1):44–52.
- Campbell, B. J., Engel, A. S., Porter, M. L., and Takai, K. (2006). The versatile ϵ -proteobacteria: Key players in sulphidic habitats. *Nature Reviews Microbiology*, 4(6):458–468.
- Castenholz, R. W., Wilmotte, A., Herdman, M., Rippka, R., Waterbury, J. B., Itean, I., and Hoffmann, L. (2001). Phylum bx. cyanobacteria. In Boone, D. R., Castenholz, R. W., and Garrity, G. M., editors, *Bergey's Manual® of Systematic Bacteriology: Volume One: The Archaea and the Deeply Branching and Phototrophic Bacteria*, pages 473–599. Springer.

533 Cluff, M. A., Hartsock, A., MacRae, J. D., Carter, K., and Mouser, P. J. (2014). Temporal changes in
534 microbial ecology and geochemistry in produced water from hydraulically fractured Marcellus Shale
535 gas wells. *Environmental Science & Technology*, 48(11):6508–6517.

536 Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., Brown, C. T., Porras-Alfaro,
537 A., Kuske, C. R., and Tiedje, J. M. (2014). Ribosomal Database Project: Data and tools for high
538 throughput rRNA analysis. *Nucleic Acids Research*, 42(D1):D633–D642.

539 Dick, J. M. (2014). Average oxidation state of carbon in proteins. *Journal of the Royal Society,*
540 *Interface*, 11:20131095.

541 Dick, J. M. (2021a). JMDplots 1.2.8. Zenodo. <https://doi.org/10.5281/zenodo.5375747>.

542 Dick, J. M. (2021b). Water as a reactant in the differential expression of proteins in cancer. *Computa-*
543 *tional and Systems Oncology*, 1(1):e1007.

544 Dick, J. M. and Shock, E. L. (2013). A metastable equilibrium model for the relative abundances of
545 microbial phyla in a hot spring. *PLOS One*, 8(9):e72395.

546 Dick, J. M., Yu, M., and Tan, J. (2020). Uncovering chemical signatures of salinity gradients through
547 compositional analysis of protein sequences. *Biogeosciences*, 17(23):6145–6162.

548 Dick, J. M., Yu, M., Tan, J., and Lu, A. (2019). Changes in carbon oxidation state of metagenomes
549 along geochemical redox gradients. *Frontiers in Microbiology*, 10:120.

550 Ducluzeau, A.-L., Schoepp-Cothenet, B., van Lis, R., Baymann, F., Russell, M. J., and Nitschke, W.
551 (2014). The evolution of respiratory O₂/NO reductases: An out-of-the-phylogenetic-box perspective.
552 *Journal of the Royal Society, Interface*, 11(98):20140196.

553 Eichorst, S. A., Trojan, D., Roux, S., Herbold, C., Rattei, T., and Woebken, D. (2018). Genomic insights
554 into the *Acidobacteria* reveal strategies for their success in terrestrial environments. *Environmental*
555 *Microbiology*, 20(3):1041–1063.

556 Fernández-Gómez, B., Richter, M., Schüler, M., Pinhassi, J., Acinas, S. G., González, J. M., and
557 Pedrós-Alió, C. (2013). Ecology of marine Bacteroidetes: A comparative genomics approach. *ISME*
558 *Journal*, 7(5):1026–1037.

559 Fones, E. M., Colman, D. R., Kraus, E. A., Nothaft, D. B., Poudel, S., Rempfert, K. R., Spear, J. R.,
560 Templeton, A. S., and Boyd, E. S. (2019). Physiological adaptations to serpentinization in the
561 Samail Ophiolite, Oman. *ISME Journal*, 13(7):1750–1762.

562 Ganesh, S., Bristow, L. A., Larsen, M., Sarode, N., Thamdrup, B., and Stewart, F. J. (2015). Size-
563 fraction partitioning of community gene transcription and nitrogen metabolism in a marine oxygen
564 minimum zone. *ISME Journal*, 9(12):2682–2696.

Harris, J. K., Caporaso, J. G., Walker, J. J., Spear, J. R., Gold, N. J., Robertson, C. E., Hugenholtz, P., Goodrich, J., McDonald, D., Knights, D., Marshall, P., Tufo, H., Knight, R., and Pace, N. R. (2013). Phylogenetic stratigraphy in the Guerrero Negro hypersaline microbial mat. *ISME Journal*, 7(1):50–60.

Hayes, T. (2009). Sampling and Analysis of Water Streams Associated with the Development of Marcellus Shale Gas. Final Report, Marcellus Shale Coalition. <https://www.water-research.net/naturalgasPA/pdffiles/MSCCommission-Report.pdf>, Accessed on 2021-05-27.

He, P., Xie, L., Zhang, X., Li, J., Lin, X., Pu, X., Yuan, C., Tian, Z., and Li, J. (2020). Microbial diversity and metabolic potential in the stratified Sansha Yongle Blue Hole in the South China Sea. *Scientific Reports*, 10(1):5949.

Herlemann, D. P. R., Lundin, D., Andersson, A. F., Labrenz, M., and Jürgens, K. (2016). Phylogenetic signals of salinity and season in bacterial community composition across the salinity gradient of the Baltic Sea. *Frontiers in Microbiology*, 7:1883.

Hördt, A., López, M. G., Meier-Kolthoff, J. P., Schleuning, M., Weinhold, L.-M., Tindall, B. J., Gronow, S., Kyrpides, N. C., Woyke, T., and Göker, M. (2020). Analysis of 1,000+ type-strain genomes substantially improves taxonomic classification of *Alphaproteobacteria*. *Frontiers in Microbiology*, 11:468.

Hull, N. M., Rosenblum, J. S., Robertson, C. E., Harris, J. K., and Linden, K. G. (2018). Succession of toxicity and microbiota in hydraulic fracturing flowback and produced water in the Denver–Julesburg Basin. *Science of The Total Environment*, 644:183–192.

James, F. C. and McCulloch, C. E. (1990). Multivariate analysis in ecology and systematics: Panacea or Pandora’s box? *Annual Review of Ecology and Systematics*, 21(1):129–166.

Jungblut, A. D., Hawes, I., Mackey, T. J., Krusor, M., Doran, P. T., Sumner, D. Y., Eisen, J. A., Hillman, C., and Goroncy, A. K. (2016). Microbial mat communities along an oxygen gradient in a perennially ice-covered Antarctic lake. *Applied and Environmental Microbiology*, 82(2):620–630.

Khan, H. J., Spielman-Sun, E., Jew, A. D., Bargar, J., Kavscek, A., and Druhan, J. L. (2021). A critical review of the physicochemical impacts of water chemistry on shale in hydraulic fracturing systems. *Environmental Science & Technology*, 55(3):1377–1394.

Klinke, S. (2021). *plot.matrix: Visualizes a Matrix as Heatmap*. R package version 1.6.

Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R., Wilson, K. R., Altieri, K. E., Mazzoleni, L. R., Wozniak, A. S., Bluhm, H., Mysak, E. R., Smith, J. D., Kolb, C. E., and Worsnop, D. R. (2011). Carbon oxidation state as a metric for describing the chemistry of atmospheric organic aerosol. *Nature Chemistry*, 3(2):133–139.

- 598 Lecoivre, A., Ménez, B., Cannat, M., Chavagnac, V., and Gérard, E. (2021). Microbial ecology of the
599 newly discovered serpentinite-hosted Old City hydrothermal field (southwest Indian ridge). *ISME*
600 *Journal*, 15(3):818–832.
- 601 LeDoux, S. T. M., Szykiewicz, A., Faiia, A. M., Mayes, M. A., McKinney, M. L., and Dean, W. G.
602 (2016). Chemical and isotope compositions of shallow groundwater in areas impacted by hydraulic
603 fracturing and surface mining in the Central Appalachian Basin, Eastern United States. *Applied*
604 *Geochemistry*, 71:73–85.
- 605 Leinonen, R., Sugawara, H., and Shumway, M. (2011). The Sequence Read Archive. *Nucleic Acids*
606 *Research*, 39(suppl_1):D19–D21.
- 607 Liden, T., Santos, I. C., Hildenbrand, Z. L., and Schug, K. A. (2019). Analytical methods for the
608 comprehensive characterization of produced water. In Ahuja, S., editor, *Evaluating Water Quality to*
609 *Prevent Future Disasters*, volume 11 of *Separation Science and Technology*, pages 199–217. Academic
610 Press.
- 611 Lin, T. J., Ver Eecke, H. C., Breves, E. A., Dyar, M. D., Jamieson, J. W., Hannington, M. D., Dahle,
612 H., Bishop, J. L., Lane, M. D., Butterfield, D. A., Kelley, D. S., Lilley, M. D., Baross, J. A., and
613 Holden, J. F. (2016). Linkages between mineralogy, fluid chemistry, and microbial communities
614 within hydrothermal chimneys from the Endeavour Segment, Juan de Fuca Ridge. *Geochemistry,*
615 *Geophysics, Geosystems*, 17(2):300–323.
- 616 Lovley, D. R. and Klug, M. J. (1986). Model for the distribution of sulfate reduction and methanogenesis
617 in freshwater sediments. *Geochimica et Cosmochimica Acta*, 50(1):11–18.
- 618 Lozupone, C. A. and Knight, R. (2007). Global patterns in bacterial diversity. *Proceedings of the*
619 *National Academy of Sciences*, 104(27):11436–11440.
- 620 Luek, J. L., Harir, M., Schmitt-Kopplin, P., Mouser, P. J., and Gonsior, M. (2019). Organic
621 sulfur fingerprint indicates continued injection fluid signature 10 months after hydraulic fracturing.
622 *Environmental Science: Processes & Impacts*, 21(2):206–213.
- 623 Markelova, E., Parsons, C. T., Couture, R.-M., Smeaton, C. M., Madé, B., Charlet, L., and Van Cap-
624 pellen, P. (2017). Deconstructing the redox cascade: What role do microbial exudates (flavins) play?
625 *Environmental Chemistry*, 14(8):515–524.
- 626 Matthews, R. E. F. (1975). A classification of virus groups based on the size of the particle in relation
627 to genome size. *Journal of General Virology*, 27(2):135–149.
- 628 Mayr, M. J., Zimmermann, M., Guggenheim, C., Brand, A., and Bürgmann, H. (2020). Niche
629 partitioning of methane-oxidizing bacteria along the oxygen-methane counter gradient of stratified
630 lakes. *ISME Journal*, 14(1):274–287.

631 Meier, D. V., Pjevac, P., Bach, W., Hourdez, S., Girguis, P. R., Vidoudez, C., Amann, R., and
632 Meyerdierks, A. (2017). Niche partitioning of diverse sulfur-oxidizing bacteria at hydrothermal
633 vents. *ISME Journal*, 11:1545–1558.

634 Mouser, P. J., Borton, M., Darrah, T. H., Hartsock, A., and Wrighton, K. C. (2016). Hydraulic
635 fracturing offers view of microbial life in the deep terrestrial subsurface. *FEMS Microbiology Ecology*,
636 92(11):fiw166.

637 Mumford, A. C., Maloney, K. O., Akob, D. M., Nettemann, S., Proctor, A., Ditty, J., Ulsamer, L.,
638 Lookenbill, J., and Cozzarelli, I. M. (2020). Shale gas development has limited effects on stream
639 biology and geochemistry in a gradient-based, multiparameter study in Pennsylvania. *Proceedings*
640 *of the National Academy of Sciences*, 117(7):3670–3677.

641 Nakagawa, S., Takai, K., Inagaki, F., Hirayama, H., Nunoura, T., Horikoshi, K., and Sako, Y. (2005).
642 Distribution, phylogenetic diversity and physiological characteristics of epsilon-*Proteobacteria* in a
643 deep-sea hydrothermal field. *Environmental Microbiology*, 7(10):1619–1632.

644 O’Leary, N. A., Wright, M. W., Brister, J. R., Ciufu, S., Haddad, D., McVeigh, R., Rajput, B.,
645 Robbertse, B., Smith-White, B., Ako-Adjei, D., Astashyn, A., Badretdin, A., Bao, Y., Blinkova,
646 O., Brover, V., Chetvernin, V., Choi, J., Cox, E., Ermolaeva, O., Farrell, C. M., Goldfarb, T.,
647 Gupta, T., Haft, D., Hatcher, E., Hlavina, W., Joardar, V. S., Kodali, V. K., Li, W., Maglott, D.,
648 Masterson, P., McGarvey, K. M., Murphy, M. R., O’Neill, K., Pujar, S., Rangwala, S. H., Rausch,
649 D., Riddick, L. D., Schoch, C., Shkeda, A., Storz, S. S., Sun, H., Thibaud-Nissen, F., Tolstoy, I.,
650 Tully, R. E., Vatsan, A. R., Wallin, C., Webb, D., Wu, W., Landrum, M. J., Kimchi, A., Tatusova,
651 T., DiCuccio, M., Kitts, P., Murphy, T. D., and Pruitt, K. D. (2016). Reference sequence (RefSeq)
652 database at NCBI: Current status, taxonomic expansion, and functional annotation. *Nucleic Acids*
653 *Research*, 44(D1):D733–D745.

654 Orcutt, B. N., Sylvan, J. B., Knab, N. J., and Edwards, K. J. (2011). Microbial ecology of the dark
655 ocean above, at, and below the seafloor. *Microbiology and Molecular Biology Reviews*, 75(2):361–422.

656 Paradis, E. and Schliep, K. (2019). ape 5.0: An environment for modern phylogenetics and evolutionary
657 analyses in R. *Bioinformatics*, 35(3):526–528.

658 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.
659 (2012). The SILVA ribosomal RNA gene database project: Improved data processing and web-based
660 tools. *Nucleic Acids Research*, 41(D1):D590–D596.

661 R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for
662 Statistical Computing, Vienna, Austria.

663 Ramette, A. (2007). Multivariate analyses in microbial ecology. *FEMS Microbiology Ecology*, 62(2):142–
664 160.

665 Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. (2016). VSEARCH: A versatile open
666 source tool for metagenomics. *PeerJ*, 4:e2584.

667 Sherwood, J. E., Stagnitti, F., Kokkinn, M. J., and Williams, W. D. (1991). Dissolved oxygen
668 concentrations in hypersaline waters. *Limnology and Oceanography*, 36(2):235–250.

669 Sollai, M., Villanueva, L., Hopmans, E. C., Reichart, G.-J., and Sinninghe Damsté, J. S. (2019). A
670 combined lipidomic and 16S rRNA gene amplicon sequencing approach reveals archaeal sources of
671 intact polar lipids in the stratified Black Sea water column. *Geobiology*, 17(1):91–109.

672 Tessler, M., Neumann, J. S., Afshinnekoo, E., Pineda, M., Hersch, R., Velho, L. F. M., Segovia, B. T.,
673 Lansac-Toha, F. A., Lemke, M., DeSalle, R., Mason, C. E., and Brugler, M. R. (2017). Large-
674 scale differences in microbial biodiversity discovery between 16S amplicon and shotgun sequencing.
675 *Scientific Reports*, 7(1):6589.

676 Thompson, L. R., Sanders, J. G., McDonald, D., Amir, A., Ladau, J., Locey, K. J., Prill, R. J., Tripathi,
677 A., Gibbons, S. M., Ackermann, G., Navas-Molina, J. A., Janssen, S., Kopylova, E., Vázquez-Baeza,
678 Y., González, A., Morton, J. T., Mirarab, S., Zech Xu, Z., Jiang, L., Haroon, M. F., Kanbar, J.,
679 Zhu, Q., Jin Song, S., Kosciulek, T., Bokulich, N. A., Lefler, J., Brislawn, C. J., Humphrey, G.,
680 Owens, S. M., Hampton-Marcell, J., Berg-Lyons, D., McKenzie, V., Fierer, N., Fuhrman, J. A.,
681 Clauset, A., Stevens, R. L., Shade, A., Pollard, K. S., Goodwin, K. D., Jansson, J. K., Gilbert,
682 J. A., Knight, R., and The Earth Microbiome Project Consortium (2017). A communal catalogue
683 reveals Earth’s multiscale microbial diversity. *Nature*, 551(7681):457–463.

684 Ulrich, N., Kirchner, V., Drucker, R., Wright, J. R., McLimans, C. J., Hazen, T. C., Campa, M. F.,
685 Grant, C. J., and Lamendella, R. (2018). Response of aquatic bacterial communities to hydraulic
686 fracturing in northwestern Pennsylvania: A five-year study. *Scientific Reports*, 8(1):5683.

687 Visconti, F. and de Paz, J. M. (2021). Estimation of the carbon valence from its average formal oxidation
688 state in the soil organic matter. *European Journal of Soil Science*. <https://doi.org/10.1111/ejss.13122>.

689 Waite, D. W., Vanwonterghem, I., Rinke, C., Parks, D. H., Zhang, Y., Takai, K., Sievert, S. M., Simon,
690 J., Campbell, B. J., Hanson, T. E., Woyke, T., Klotz, M. G., and Hugenholtz, P. (2018). Addendum:
691 Comparative genomic analysis of the class *Epsilonproteobacteria* and proposed reclassification to
692 *Epsilonbacteraeota* (phyl. nov.). *Frontiers in Microbiology*, 9:772.

693 Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naïve Bayesian classifier for rapid
694 assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental*
695 *Microbiology*, 73(16):5261–5267.

696 Xie, K. H., Deng, Y., Zhang, S. C., Zhang, W. H., Liu, J. R., Xie, Y. L., Zhang, X. Z., and Huang, H.
697 (2017). Prokaryotic community distribution along an ecological gradient of salinity in surface and
698 subsurface saline soils. *Scientific Reports*, 7:13332.

699 Xie, L., Wang, B., Pu, X., Xin, M., He, P., Li, C., Wei, Q., Zhang, X., and Li, T. (2019). Hydrochemical
700 properties and chemocline of the Sansha Yongle Blue Hole in the South China Sea. *Science of The*
701 *Total Environment*, 649:1281–1292.

702 Zhong, C., Li, J., Flynn, S. L., Nesbø, C. L., Sun, C., von Gunten, K., Lanoil, B. D., Goss, G. G.,
703 Martin, J. W., and Alessi, D. S. (2019). Temporal changes in microbial community composition
704 and geochemistry in flowback and produced water from the Duvernay formation. *ACS Earth and*
705 *Space Chemistry*, 3(6):1047–1057.

706 Zhong, Z.-P., Liu, Y., Miao, L.-L., Wang, F., Chu, L.-M., Wang, J.-L., and Liu, Z.-P. (2016).
707 Prokaryotic community structure driven by salinity and ionic concentrations in plateau lakes of the
708 Tibetan Plateau. *Applied and Environmental Microbiology*, 82(6):1846–1858.

709

Appendix 1

710

Appendix 1 Table S1. Sequence processing statistics.

Study key	BioProject	Samples	Reads per sample (RPS) (a)	Filtered RPS (b)	Filtered read length (c)	Single-ton % (d)	Chim-era % (e)	RPS used for classification (f)	Classification Map to to genus level (%) (g)	NCBI taxonomy (%) (h)
Natural Environment Datasets										
HCW+13	N/A (GenBank)	10	11297	11297	(g)	(g)	18.1	9249	30	86.9
BGPF13	PRJNA207095	14	36386	17886	(h)	60.2	2.0	6120	20	84.7
GBL+15	PRJNA263621	15	20371	18123	250	6.1	7.0	8882	29	97.4
HLA+16	PRJEB1245	105	1878	1464	(h)	(i)	0.4	1458	37	98.0
JHM+16	PRJNA291280	8	1165067	420093 (j)	230	16.1	10.4	8960	17	96.6
ZLM+16	PRJNA294836	46	59934	48973	250	14.2	5.4	9456	33	97.6
MPB+17	PRJEB15554	54	165298	88931	450	54.4	7.7	8359	48	97.9
XDZ+17	PRJNA388250	29	59876	54360	250	8.9	2.4	9757	46	95.7
BCA+21	PRJNA395513	36	280205	95996	450	34.9	9.9	8638	43	98.0
SVH+19	PRJNA423140	15	6631	5718	(h)	11.6	5.9	4753	17	97.6
MZG+20	PRJEB27579	134 (k)	47294	32779	450	59.3	4.0	9151	54	95.6
HXZ+20	PRJNA503500	21 (l)	92727	75235	440	34.3	2.3	9773	38	95.8
Shale Gas Datasets										
UKD+18 .sediment	PRJNA394724	93	129499	115420 (m)	100	10.4	1.4	9799	22	85.7
UKD+18 .water	PRJNA394724	80	87546	73960 (m)	100	14.2	1.3	9635	21	93.1
MMA+20	PRJNA544240	138	154859	114533	250	17.8	4.3	9496	38	85.5
CHM+14	PRJNA229085	46	5852	4169	(h)	25.4	35.2	2014	96	97.4
HRR+18	PRJNA438710	9	388424	175069 (n)	300	44.4	23.5	7655	99	72.7
ZLF+19	PRJNA407226	6	64644	32638	290	16.8	0.6	9938	66	98.0

a. Paired forward and reverse reads are counted as one. Paired-end reads were merged with “vsearch -fastq_mergepairs” with default options; reads that failed merging were excluded from the subsequent analysis. All “reads per sample” columns are averages of all samples.

b. Filtering was done with “vsearch -fastq_filter” with the options “-fastq_maxee_rate 0.005” and “-fastq_truncLen length” with length value given in next column.

c. Reads in all samples (runs) were pooled and singletons (sequences appearing exactly once) were identified with “vsearch -derep_fulllength” with the option “-maxuniquesize 1”. The singletons were removed from the pooled file using “seqtk subseq”; the remaining sequences for each run were extracted from the pooled file using an awk script. After removing the singletons, 10000 sequences were subsampled from each run for the subsequent analysis using “vsearch -fastx_subsample” with the options “-sample_size 10000” and “-randseed 1234”; subsampling was not done on runs with fewer than 10000 sequences remaining after filtering and singleton removal.

d. After subsampling, runs were pooled again and chimeras were removed using “vsearch -uchime_ref” with the option

“-nonchimeras” to output non-chimeric sequences (i.e. those not classified as either chimeras or borderline chimeras). The remaining sequences for each run were extracted from the output and used for taxonomic classification.

e. Classification using RDP Classifier. Any samples with < 200 classified sequences at all taxonomic levels were excluded from downstream analysis.

f. This shows the percentage of successful mappings (exact matches of rank and taxonomic name, with exceptions noted in the text) between RDP and the NCBI taxonomy for all sequences classified at phylum and lower levels.

g. FASTA sequences were downloaded from GenBank. No quality filtering or singleton removal was done.

h. For these 454 sequencing experiments, filtering was done with “vsearch -fastq_filter” with options “-fastq_strip 18” to remove the primer sequence and “-fastq_minlen 200 -fastq_maxlen 600 -fastq_truncqual 15” for read length and quality filtering.

i. Because of the low number of sequences, singletons were not removed.

j. After merging, reads were filtered only on length, not quality scores.

k. Library “c” was chosen, representing 134 out of 536 total runs, because it has more sequences per run than the other available libraries (a, b, and d).

l. Sequences from Station C4 were used for samples outside the Blue Hole.

m. Only forward reads were used, as indicated by the previous authors.

n. Only forward reads were used because a large majority of merge attempts failed for one or more runs (likely because of low-quality reverse reads).

Appendix 1 Table S2. Most abundant unmapped taxonomic groups.

Study key	Site	Most abundant unmapped groups
HCW+13	Guerrero Negro mat	genus_Potamolinea (2.27%), genus_Saccharicenans (1.54%)
BGPF13	Yellowstone hot springs	genus_Armatimonadetes_gp7 (4.05%), genus_Diapherotrites Incertae Sedis AR10 (3.91%)
GBL+15	ETNP water	phylum_Woearchaeota (0.95%), genus_Pacearchaeota Incertae Sedis AR13 (0.47%)
HLA+16	Baltic Sea water	family_Thalassobaculaceae (0.92%), genus_Cryptomonadaceae (0.48%)
JHM+16	Lake Fryxell mat	genus_Acidibacter (0.43%), phylum_Woearchaeota (0.4%)
ZLM+16	Tibetan Plateau lakes	genus_Cryptomonadaceae (1.1%), phylum_Woearchaeota (0.42%)
MPB+17	Manus Basin vents	genus_Parcubacteria_genera_incertae_sedis (0.56%), genus_Thiopropundum (0.28%)
XDZ+17	Qarhan Salt Lake soils	genus_Candidatus_Nanosalina (1.59%), genus_Gp4 (0.46%) (a)
BCA+21	Ursu Lake	genus_Desulfonatrobacter (1.53%), class_Subdivision3 (b) (0.24%)
SVH+19	Black Sea water	genus_Pacearchaeota Incertae Sedis AR13 (0.52%), genus_Cryptomonadaceae (0.36%)
MZG+20	Swiss lakes	genus_Cryptomonadaceae (1.12%), genus_GpXI (1.07%) (c)
HXZ+20	Sansha Yongle Blue Hole	family_Arcobacteraceae (1.6%), genus_Parcubacteria_genera_incertae_sedis (0.84%)
UKD+18.sediment	Pennsylvania Streams	genus_Gp2 (5.52%) (a), genus_Gp3 (1.79%) (a)
UKD+18.water	Pennsylvania Streams	genus_Gp6 (0.91%) (a), genus_GpVIII (0.66%) (c)
MMA+20	Penn. Streams Water	genus_Gp3 (1.64%) (a), genus_Gp2 (1.57%) (a)
CHM+14	Marcellus Shale	family_Arcobacteraceae (2.11%), order_Clostridiales (0.44%)
HRR+18	Denver-Julesburg Basin	genus_Cavicella (27.02%), genus_Gelria (0.13%)
ZLF+19	Duvernay Formation	genus_Armatimonas/Armatimonadetes_gp1 (1.14%), genus_Subdivision3_genera_incertae_sedis (0.53%) (c)

a. Acidobacteria. b. Verrucomicrobia. c. Cyanobacteria.

Figure 6—Figure supplement 1. Calculations were performed as described in Figure 4. Arrows are drawn between subsets of samples classified as oxidized and reduced according to the descriptions in Figure 6.