Reduced H⁺ channel activity disrupts pH homeostasis and calcification in coccolithophores at low ocean pH

Dorothee Kottmeier^{1,3}, Abdul Chrachri¹, Gerald Langer¹, Katherine Helliwell^{1,4}, Glen L. Wheeler^{1*}, Colin Brownlee^{1,2*}

¹Marine Biological Association, The Laboratory, Citadel Hill, Plymouth. PL1 2PB

²School of Ocean and Earth Science, University of Southampton, Southampton, SO14 3ZH, UK.

³MARUM Center for Marine Environmental Sciences, University of Bremen, 28334 Bremen, Germany

⁴Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, EX4 4QD, UK

*Correspondence: cbr@mba.ac.uk and glw@mba.ac.uk

Key words: Calcification, coccolithophore, *Coccolithus braarudii*, pH regulation, H⁺ channel

Abstract

Coccolithophores produce the bulk of ocean biogenic calcium carbonate but this process is predicted to be negatively affected by future ocean acidification scenarios. Since coccolithophores calcify intracellularly, the mechanisms through which changes in seawater carbonate chemistry affect calcification remain unclear. Here we show that voltage-gated H^+ channels in the plasma membrane of *Coccolithus braarudii* serve to regulate pH and maintain calcification under normal conditions, but have greatly reduced activity in cells acclimated to low pH. This disrupts intracellular pH homeostasis and impairs the ability of *C. braarudii* to remove H^+ generated by the calcification process, leading to specific coccolith malformations. These coccolith malformations can be reproduced by pharmacological inhibition of H^+ channels. Heavily-calcified coccolithophore species such as *C. braarudii*, which make the major contribution to carbonate export to the deep ocean, have a large intracellular H^+ load and are likely to be most vulnerable to future decreases in ocean pH.

Introduction

Anthropogenic CO₂ emissions and the subsequent dissolution of CO₂ in seawater have resulted in substantial changes in ocean carbonate chemistry¹. The resultant decrease in seawater pH, termed ocean acidification, is predicted to be particularly detrimental for calcifying organisms². Mean global surface ocean pH is predicted to fall as low as 7.7 by 2100³ and is likely to continue to fall further in the following centuries. Present day marine organisms can experience significant fluctuations in seawater pH, particularly in coastal and upwelling regions^{4,5}. Ocean acidification is therefore predicted to have an important influence not only on mean surface ocean pH, but also on the extremes of pH experienced by marine organisms^{6,7}. Coccolithophores, characterised by their covering of intricately-formed calcite scales (coccoliths), account for the bulk of global biological calcification and around 20% of ocean productivity, making major contributions to global biogeochemical cycles, including the long-term export of both inorganic and organic carbon from the ocean photic zone to deep waters^{8,9}. Unlike the vast majority of calcifying organisms, coccolithophore calcification occurs in an intracellular compartment, the Golgi-derived coccolith vesicle (CV), effectively isolating the calcification process from direct changes in seawater carbonate chemistry. Nevertheless, extensive laboratory observations indicate that ocean acidification may negatively impact coccolithophore calcification, albeit with significant variability of responses between species and strains¹⁰⁻¹⁴. The negative impact on calcification rates occurs at calcite saturation states (Ω) >1, indicating it results primarily from impaired cellular production rather than dissolution^{10,15}. However, prediction of how natural coccolithophore populations may respond to future changes in ocean pH are hampered by lack of mechanistic understanding of pH impacts at the cellular level¹⁰.

As calcification occurs intracellularly using external HCO₃⁻ as the primary dissolved inorganic carbon (DIC) source¹⁶⁻¹⁸, coccolith formation is not directly dependent on external CO₃²⁻ concentrations. However, the uptake of HCO₃⁻ as a substrate for calcification results in the equimolar production of CaCO₃ and H⁺ in the CV¹⁸. In order to maintain saturation conditions for calcite formation, H⁺ produced by the calcification process must be rapidly removed from the CV, placing extraordinary demands for cellular pH regulation to prevent cellular acidosis¹⁸.

Lower calcification rates under ocean acidification conditions appear to be primarily due to decreased pH rather than other aspects of carbonate chemistry^{10,19,20}. Coccolithophores exhibit highly unusual membrane physiology, including the presence of voltage-gated H⁺

channels in the plasma membrane²¹ and a high sensitivity of cytosolic pH (pH_{cyt}) to changes in external pH (pH_0)^{21, 22}. Voltage-gated H^+ channels are associated with rapid H^+ efflux in a number of specialised animal cell types²³ and contribute to effective pH regulation in coccolithophores²¹. As H^+ channel function is dependent on the electrochemical gradient of H^+ across the plasma membrane, this mechanism could be impaired under lower seawater pH. However, it remains unknown whether H^+ channels play a direct role in removal of calcification-derived H^+ or contribute to the sensitivity of coccolithophores to ocean acidification.

Coccolithophores exhibit significant diversity in their extent of calcification (Supplementary Fig. 1). The abundant bloom forming species *Emiliania huxleyi* is moderately calcified (ratio of particulate inorganic carbon to particulate organic carbon (PIC/POC) of around 1) and has been the focus of the vast majority of the studies into the effects of environmental change in coccolithophores¹³. Coastal species belonging to the Pleurochrysidaceae and Hymenomonadaceae are lightly calcified, commonly exhibiting a PIC/POC of less than 0.5²⁴⁻²⁷. Species such as *Coccolithus braarudii*, *Calcidiscus leptoporus* and *Helicosphaera carteri* exhibit much higher PIC/POC ratios and contribute the majority of carbonate export to the deep ocean in many areas²⁸⁻³⁰. The physiological response of heavily-calcified coccolithophores to ocean acidification is therefore of considerable biogeochemical significance. Growth and calcification rates in *C. leptoporus* and *C. braarudii* are sensitive to pH values predicted to prevail on a future decadal timescale^{10, 15, 31, 32}. However, a mechanistic understanding of the different sensitivity of coccolithophore species to changing ocean carbonate chemistry is lacking.

The net H^+ load in a cell is determined by the combination of metabolic processes that consume or produce H^+ . H^+ fluxes in coccolithophores will be primarily determined by the balance of H^+ consumed by photosynthesis and H^+ generated by calcification, with uptake of different carbon sources a particularly important consideration (Fig. 1a). CO_2 uptake for photosynthesis results in no net production or consumption of H^+ , whereas uptake of HCO_3^- requires the equimolar consumption of H^+ in order to generate CO_2 . Growth at elevated CO_2 causes a switch from HCO_3^- uptake to predominately CO_2 uptake in *E. huxleyi*^{33, 34}. The associated net decrease in H^+ consumption will therefore increase the H^+ load in coccolithophores grown at elevated CO_2 , which may exacerbate the potential for cytosolic acidosis caused by lower seawater pH.

In this study we set out to better understand the cellular mechanisms underlying the sensitivity of coccolithophore calcification to lower pH. We subjected the heavily-calcified species *C. braarudii*, which is commonly found in temperate upwelling regions^{35, 36}, to conditions that reflect the range of pH values it may experience in current and future oceans. We show that acclimation to low pH leads to loss of H⁺ channel function and disruption of cellular pH regulation in *C. braarudii*. These effects are coincident with very specific defects in coccolith morphology that can be reproduced by direct inhibition of H⁺ channels. We conclude that H⁺ efflux through H⁺ channels is essential for maintaining both calcification rate and coccolith morphology. By providing a mechanistic insight into pH regulation during the calcification process, our results indicate that disruption of coccolithophore calcification in a future acidified ocean is likely to be most severe in heavily calcified species.

Results

Cellular H⁺ load varies with DIC use for calcification and photosynthesis.

To examine more closely how the balance of photosynthesis and calcification may influence the cellular H⁺ load, we measured physiological parameters in *C. braarudii* cells acclimated to a broad range of carbonate chemistries (Supplementary Table 1). *C. braarudii* exhibited pH optima for growth, PIC and POC production of 8.32 ± 0.01 , 8.20 ± 0.03 and 8.24 ± 0.02 respectively (pH_{NBS}, n=3, $\pm$ SE), with sharp declines in these parameters exhibited by cells grown at pH 7.85 and 7.55 (Fig. 1b-d). PIC production decreased more strongly than POC production in acidified conditions, leading to lower PIC:POC ratios. These trends are in close agreement with other laboratory studies examining the response of *C. braarudii* to changing carbonate chemistries^{10, 15, 31, 32, 37}. Calculation of the H⁺ load from values of PIC and POC production indicated that H⁺ production by calcification exceeded H⁺ consumption by photosynthesis under all scenarios, being highest at optimal PIC:POC ratios (Fig. 1e). Although the large decrease in calcification rates at seawater pH 7.55 result in lower H⁺ production, the net H⁺ load could still be substantial due to a likely increase in CO₂ uptake under these conditions (Fig. 1e)³⁴. The results illustrate that changes in the relative rates of photosynthesis and calcification, as well as in the carbon source used for photosynthesis, will have a major impact on the cellular H⁺ budget in *C. braarudii*, although in all cases there is a resultant requirement for net H⁺ efflux.

Growth at low pH results in specific defects in coccolith morphology

Morphological defects in coccoliths are widely reported in coccolithophores grown under simulated ocean acidification conditions^{37, 38}, although there is little information on the specific nature of these malformations to aid mechanistic understanding of the impacts of low seawater pH on the calcification process. Scanning electron microscopy (SEM) analysis of coccolith morphology revealed that only 19.0 ± 5.0 % and 30.1 ± 2.7 % of coccoliths exhibited normal morphology at pH 7.55 and pH 7.85 respectively (n=3, $\pm$ SE) (Fig. 2a,b). Moreover, by performing a detailed categorisation of each morphological defect, we found a very distinct ‘type-pH’ malformation at low pH, in which the shield elements are malformed and greatly reduced in length so that the coccolith appears as a ring of calcite rather than a fully formed shield (Fig. 2b). This differs from an immature coccolith, in which the individual elements are reduced in length but correctly formed (Supplementary Fig. 2). Cells grown at low pH exhibited a large increase in the number of collapsed coccospheres observed by SEM analysis

(Fig. 2c-d), indicating that the extensive malformations result in an inability to maintain the structural integrity of the coccosphere. Although the type-pH malformation has not been explicitly described previously, it can clearly be observed in other studies where *C. braarudii* has been grown at low pH³⁷. Importantly, type-pH malformations are distinct from coccolith malformations induced by other stressors, such as phosphate limitation or the Si analogue Ge^{39, 40}.

H⁺ channel function is greatly reduced following acclimation to lower pH

To investigate how these defects in coccolith morphology could arise, we examined the physiology of *C. braarudii* cells grown at low pH. *C. braarudii* exhibits an unusual large outwardly rectifying H⁺ current at membrane potentials positive of the H⁺ equilibrium potential (E_{H+}), due to the activity of voltage-gated H⁺ channels in the plasma membrane²¹. Our previous studies showed that the H⁺ channel activation potential tracked E_{H+} across the plasma membrane. Calculation of the proton motive force (pmf) across the *C. braarudii* plasma membrane (at a resting membrane potential of -46 mV²¹) indicates that there is a small net outward pmf at a seawater pH of 8.15 (Fig 3a). A decrease in pH_{cyt} results in a shift in the activation potential of the H⁺ current to more negative membrane potential values and increases the outwardly-directed pmf. These combined effects result in net H⁺ efflux and allow restoration of resting pH_{cyt} (Supplementary Fig. 3). However, at pH_o 7.55 the activation potential shifts to more positive values and the calculated pmf is no longer outward, so channel mediated H⁺ efflux would only occur following depolarisation of the membrane potential and/or further reductions in pH_{cyt} (Supplementary Figure 3).

In order to determine the impact of growth at unfavourable seawater pH on H⁺ channel function, we monitored H⁺ currents using patch clamp recordings in *C. braarudii* cells previously acclimated to pH_o 7.55, 8.15 or 8.75. The mean amplitude of the outward H⁺ current, when measured at pH_o 8.15, was greatly reduced in cells that had been acclimated to pH 7.55 (Fig. 3b-d). 52.9% of cells acclimated to pH 7.55 exhibited either greatly reduced or undetectable outward current (Fig. 3d), although these cells still displayed inward Cl⁻ currents typical of healthy *C. braarudii* cells⁴¹ (Supplementary Fig.4). The outward currents exhibited a reversal potential (E_{rev}) close to E_{H+}, indicating that H⁺ remained as the primary charge carrier in all cases (Fig. 3e). The results suggest that acclimation of *C. braarudii* to a low seawater pH unsuited to the operation of H⁺ channels results in the loss of H⁺ channel function.

A unique family of H⁺ channels in calcifying coccolithophores

Homologues of the mammalian voltage-gated H⁺ channel, Hv1, are present in coccolithophores and a range of other phytoplankton, although the large outward H⁺ currents typical of *C. braarudii* have not been reported in other algal cells, suggesting that H⁺ channels are utilised for alternative roles in non-calcifying cells (e.g. in supporting NADPH oxidase activity⁴² or in dinoflagellate bioluminescence^{43, 44}). We previously characterised Hv1 channels from *E. huxleyi* and *C. braarudii*²¹. Further analysis of haptophyte transcriptomes⁴⁵ revealed that coccolithophores possess an additional H⁺ channel homologue (Hv2) that was not found in non-calcifying haptophytes (Supplementary Fig.5). *C. braarudii* Hv2 exhibited robust H⁺ currents when expressed in a heterologous system (Supplementary Fig.5). The presence of this additional homologue suggests that coccolithophore H⁺ channels have undergone functional specialisation related to calcification. In support of a specific role in calcification, we found that *HV1* and *HV2* were only expressed in the heavily calcified heterococcolith-bearing diploid life cycle phase of *C. braarudii* and were not detected in the lightly calcified holococcolith-bearing haploid life cycle phase (Supplementary Fig.6). However, we did not find any significant change in the expression of either *HV1* or *HV2* in diploid cells acclimated to low pH (Supplementary Fig.6). This suggests that the greatly reduced H⁺ conductance in these cells results from post-transcriptional or post-translational regulation of H⁺ channels.

pH homeostasis is disrupted at low seawater pH

To determine whether the reduced H⁺ channel activity in cells acclimated to low pH led to disrupted cellular pH homeostasis, we examined resting pH_{cyt}. Cells acclimated to pH 7.55 exhibited a significantly lower mean pH_{cyt} values than cells acclimated to pH 8.15 or pH 8.75 (Fig. 4a, Supplementary Fig.7). Cells acclimated to pH 8.15 or 8.75 retained the ability to adjust intracellular pH rapidly within seconds when exposed to a higher or lower pH^{21, 22}, but this response was greatly reduced in cells acclimated to pH 7.55 (Fig. 4b-d). To determine the capacity for H⁺ efflux, we transiently exposed cells to pH 6.5 and examined their ability to restore pH_{cyt} on transfer to pH 8.15. Nearly all cells acclimated to pH 8.15 or 8.75 showed a substantial decrease in cytosolic [H⁺] on transfer from pH 6.5 to 8.15 (Fig. 4e). However, many cells acclimated to pH 7.55 showed little or no capacity to lower cellular [H⁺] on transfer from pH 6.5 to 8.5, indicating the presence of distinct populations of responsive and unresponsive cells. (Fig. 4e-f). Thus, a significant proportion of cells acclimated to pH 7.55 exhibit a defect

in H⁺ efflux, which likely reflects the highly reduced H⁺ channel activity in these cells (Fig. 3d).

Pharmacological inhibition of H⁺ channel function disrupts coccolith morphology

Our results suggest that loss of H⁺ channel function and subsequent disruption of pH homeostasis is directly responsible for the defects in calcification exhibited by *C. braarudii* grown at low pH. To directly test this hypothesis, we treated cells with two inhibitors of Hv channels, Zn²⁺²³ and 2-guanidinobenzimidazole (2-GBI)⁴⁶. Cells grown in 35 μM Zn²⁺, which inhibits the outward H⁺ conductance in *C. braarudii* by approximately 60%²¹, showed only a small reduction in growth rate (control 0.54 ± 0.01 d⁻¹ compared to Zn²⁺-treated 0.47 ± 0.01 d⁻¹, n=3, se) (Fig. 5a). However, SEM examination of Zn²⁺-treated cells after 5 d revealed severe disruptions of coccolith morphology (Fig. 5b-d). Importantly, Zn-treated cells exhibited the unique type-pH coccolith malformations, which were completely absent from control cells. Growth of cells in 15 μM 2-GBI for 5 d also resulted in the presence of type-pH coccolith malformations (Supplementary Fig.8), suggesting that this calcification phenotype is specifically associated with impaired H⁺ channel function. Our results show that pharmacological inhibition of the H⁺ current leads to highly specific malformations in coccolith morphology (type-pH) that have only previously been observed in cells grown at low pH.

Discussion

We have shown that voltage-gated H^+ channels play a critical role in pH homeostasis during coccolith formation. Hv channels are regulated by the plasma membrane H^+ electrochemical gradient and are primed to respond to decreases in intracellular pH, allowing rapid H^+ efflux to restore intracellular pH²¹. However, as extracellular pH decreases to values predicted in future ocean acidification scenarios, H^+ efflux diminishes to the extent where this mechanism is no longer effective. Under such conditions, H^+ channel function may only be maintained by reduced pH_{cyt} or depolarisation of the membrane potential, with likely pronounced impacts on other physiological processes. Indeed, the loss of capacity to generate outward H^+ currents shown here in cells acclimated to lower pH suggests a physiological need to switch off this mechanism of pH_{cyt} control. Our results also indicate that alternative mechanisms to maintain cytosolic pH homeostasis, such as energised forms of H^+ transport (e.g. H^+ -ATPases, Na^+/H^+ exchangers), are incapable of dealing with the exceptional H^+ load generated by intracellular calcification (Supplementary Table 2). Loss of H^+ channel function will therefore lead to cytoplasmic acidosis unless calcification rate is reduced. The reliance on H^+ channels for pH homeostasis may constrain the ability of coccolithophores to adapt to lower pH environments.

The inactivation of the outward H^+ current at low seawater pH most likely involves changes in protein translation or post-translational modifications that modify channel activation. Elucidating these cellular mechanisms will be key in determining whether the loss of H^+ channel function is reversible. Long-term experiments examining whether coccolithophores may eventually adapt to ocean acidification conditions have yielded complex results, but support a trend of decreased calcification rates^{47, 48}. Whilst physiological adaptations to low seawater pH are possible, such as recruiting alternative mechanisms for pH regulation or reducing calcification rates to lower the H^+ load, these would either incur increased energetic costs or reduce the overall degree of calcification in future populations (Supplementary Table 2). The substantial heterogeneity in individual cell responses to low pH observed in the present study is also likely to be significant in determining how selection may operate on natural populations.

The sensitivity of *C. braarudii* calcification to low pH may not only be relevant to future ocean scenarios, but may also contribute to its distribution in current oceans. *C. braarudii* is commonly found in temperate regions, including the Iberian and Benguela upwelling systems that are associated with significant variability in seawater pH^{35, 36, 49}. It is

notable that *C. braarudii* populations in the Iberian Peninsula are predominately associated with frontal systems at the periphery of the upwelled waters³⁵, rather than directly within the upwelling regions that experience the greatest extremes of seawater pH.

Calculated cellular H⁺ budgets differ considerably between coccolithophore species. In heavily-calcified species, where calcification can occur at twice the rate of photosynthesis¹⁰, rapid removal of excess H⁺ is essential. However, in lightly-calcified species H⁺ production by calcification may be balanced by H⁺ consumption during photosynthesis, resulting in a much lower dependence on functional H⁺ channels. Calcification status may therefore be an important determinant in the sensitivity of coccolithophores to ocean acidification. A recent meta-analysis of multiple species revealed that the sensitivity of calcification rate to elevated seawater CO₂ showed a strong positive correlation to PIC/POC ratio¹¹. Moreover, heavily-calcified species such as *Calcidiscus leptoporus* and *Gephyrocapsa oceanica* show highly malformed coccoliths under future ocean acidification conditions, whereas coccolith morphology in lightly-calcified species, such as *Syracosphaera pulchra*, *Chrysotila carterae* and *Ochrosphaera neopolitana* is less sensitive^{15, 25, 27, 47, 50}. Indeed, evidence from boron isotope approaches indicated that *O. neopolitana* is able to maintain a constant pH in the coccolith vesicle over a range of seawater carbonate chemistries, although the pH range examined was relatively narrow (pH 8.05-8.35)²⁵. Our data provide mechanistic insight into the differential sensitivity of coccolithophore species, suggesting that H⁺ load is likely to be the key determinant of their sensitivity to ocean acidification. This conclusion is seemingly at odds with observations of over-calcified morphotypes of *E. huxleyi* at higher CO₂ levels in the Southern Ocean⁵¹, and millennial-scale trends indicating a correlation between increasing prevalence of ‘over-calcified’ morphotypes of *E. huxleyi* with increased atmospheric CO₂ over approximately the past 150 ka⁵². However, laboratory analyses of ‘over-calcified’ *E. huxleyi* morphotypes suggest that this phenotype relates primarily to coccolith morphology rather than calcification rate, as their PIC/POC ratios are not higher than those with normal coccolith morphology⁵³.

While non-specific defects in coccolith morphology reflecting reduced calcification in response to ocean acidification have been observed in many studies^{15, 37}, the unique malformations observed here in *C. braarudii* now provide a mechanistic link between seawater pH, the ability to regulate pH_{cyt} and coccolith morphology. The highly specific nature of the *C. braarudii* malformations may facilitate the identification of low pH stress in environmental populations and aid the characterisation of past ocean acidification events in the fossil record

³⁷. Modelled reconstructions indicate that, apart from the last ca 25 Ma, surface ocean pH has been lower than at present over much of the 200 Ma since the emergence of calcifying coccolithophores ^{54, 55}. This suggests that coccolithophores possess some capacity to adapt to the lowering of seawater pH over geological timescales. However, the very rapid predicted decline in surface ocean pH driven by anthropogenic CO₂ emissions may limit the degree to which coccolithophores can adapt their physiology. Recent evidence indicates that the mass extinction event at the (K-Pg) boundary (66 MYA), which led to the loss of 90% of coccolithophore species, was associated with rapid ocean acidification ⁵⁶. It is notable that many of the coccolithophore species that survived the K-Pg Cretaceous–Paleogene mass extinction event were coastal species ⁵⁷⁻⁵⁹, which may have been better suited to variable seawater pH and therefore less sensitive to ocean acidification.

Multiple environmental parameters in addition to carbonate chemistry are predicted to change in future oceans, including temperature, nutrient availability and ecosystem scale changes in the abundance of predators, pathogens and competitors ⁶⁰. Predicting the response of coccolithophore populations to future environmental change is therefore highly complex. Our incomplete understanding of the haplo-diplontic life cycle of coccolithophores further limits our ability to predict how natural populations may respond to unfavourable conditions ⁶¹. However, our results show that the physiology of heavily-calcified species such as *C. braarudii* is best suited to a constant seawater pH and that calcification is likely to be severely affected by ocean acidification. The ability of coccolithophores to calcify intracellularly, which has facilitated the evolution of remarkably diverse coccolith architecture, required the development of specialised physiological mechanisms for pH homeostasis that ultimately may constrain the ability of certain species to adapt to rapid changes in ocean pH.

Methods

Cell culturing

Cultures of *Coccolithus braarudii* (PLY182g) (formerly *Coccolithus pelagicus* ssp. *braarudii*) were grown in sterile-filtered seawater containing additions of nitrate, phosphate, trace metals and vitamins according to standard F/2 medium as described previously⁴¹. Silicon, selenium and nickel were also supplemented in concentrations of 10 μM , 0.0025 μM and 0.0022 μM , respectively. Dilute-batch cultures were maintained at 15°C under an irradiance of 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$ with a 16:8 h light:dark cycle. Cells were cultured in autoclaved borosilicate bottles with minimal headspace and gas-tight lids to avoid in- and outgassing of CO_2 (Duran Group, Mainz, Germany).

Acclimation to various seawater pH

Cultures were pre-acclimated for 4 d in a range of seawater pH/carbonate chemistry conditions and then used to inoculate test cultures (Supplementary Table 1). Triplicate cultures were used for all analyses, except for the pH_{cyt} measurements where 5 replicate cultures were grown. Growth rates and coccolith morphology were determined after 5 d in test conditions (i.e. a minimum of 9 d acclimation). Physiological measurements (pH_{cyt} , patch clamp) were performed between 5-10 d after inoculation into test conditions. Adjustment of seawater pH/carbonate chemistry was performed by modulating total alkalinity (TA) with amounts of HCl or NaOH at constant DIC in sealed containers. Cell density was kept between 500 and 4000 cells mL^{-1} to minimise carbonate chemistry drifts. Carbonate chemistry was measured immediately after cell inoculation and at the end of the acclimation period measuring pH_{NBS} and total alkalinity TA with a pH meter (Mettler Toledo, UK) and alkalinity titrator (TitroLine alpha plus, Schott Instruments, Germany). TA measurements were corrected with certified reference materials (CRM; provided by A. Dickson; Scripps Institution of Oceanography, U.S.A.). Data was accuracy-corrected with certified reference materials supplied by A. Dickson (Scripps Institution of Oceanography, US). Calculations were made with CO2SYS⁶².

Phenotypic changes in physiology

Cell growth was assessed by daily cell counting with Sedgewick Rafter counting chamber (Graticule Optics, UK) using a Leica DM 1000 LED light microscope. Specific growth rates (μ) were calculated from daily increments in cell concentrations counted every 24 or 48 h⁶³.

Cellular POC content was estimated by measuring the area of decalcified cells microscopically. The area was converted to volume, assuming cells were spherical. The mean volume [μm^3] of at least 20 – 50 cells per culture was converted into POC quota using the equation:

$$\text{POC} [\text{pg cell}^{-1}] = a \times V^b$$

where a and b are constants (0.216 and 0.939, respectively) for non-diatom protists⁶⁴. The cellular PIC contents were also estimated microscopically, using the volume of the coccosphere. To obtain the cellular PIC quota, the volume of the coccolith is required. The following equation was used.

$$V_c = k_s \times L^3$$

Here, V_c is the volume of coccoliths and can be estimated using coccolith length L and the shape constant k_s ⁶⁵ which is 0.06 for *C. braarudii*. The cellular PIC quota is calculated from the following equation:

$$\text{PIC} [\text{pg cell}^{-1}] = n \times V_c \times \rho \times (M_c / M_{\text{CaCO}_3})$$

where n is the total number of coccoliths per cell including the discarded coccoliths; ρ is the calcite density of $2.7 \text{ pg } \mu\text{m}^{-3}$ assuming coccoliths are pure calcite; M_c / M_{CaCO_3} is the molar mass ratio of C and CaCO_3 .

Time points for sampling cell volumes ($t = 10 \text{ h}$ after the onset of the light phase) were chosen in order present daily means (according to a modified version of the model provided in⁶³). Production rates of POC and PIC ($\text{pmol cell}^{-1} \text{ d}^{-1}$) were approximated as cellular POC content [pmol cell^{-1}] $\times \mu \times [\text{d}^{-1}]$ or cellular PIC content [pmol cell^{-1}] $\times \mu \times [\text{d}^{-1}]$, respectively. Determination of pH optima was performed by determining the vertex of a polynomial fit (second order) of three independent experiments (each experiment consists of triplicate cultures acclimated to the five different seawater pH).

Calculation of the proton motive force

The proton motive force (pmf) at the different seawater pH was estimated as

$$\text{pmf} = (zF V_m + (RT \ln([H^+]_i / [H^+]_o)) / F$$

where z is the electrical charge of H^+ , F is the Faraday constant [$\text{J V}^{-1} \text{ mol}^{-1}$], V_m is resting membrane potential [V], R is the gas constant [$\text{J mol}^{-1} \text{ K}^{-1}$] and T is temperature [K]. We used a value of -46 mV for V_m , which represents a mean of previously measured values in *C.*

*braarudii*²¹. To show how changes in pH_{cyt} and V_m may influence pmf during changes in pH_o, we calculated pmf using two values for pH_{cyt} (6.8 and 7.1) and resting V_m (-46 and -28 mV).

Calculation of H⁺ production rates

H⁺ production and consumption during photosynthesis and calcification were calculated based on POC and PIC production rates. To determine the possible range for net H⁺ load, we estimated maximal and minimal values based on a cell taking up only CO₂ (no H⁺ consumed per C fixed during photosynthesis, 2 H⁺ generated per C precipitated during calcification) or taking up only HCO₃⁻ (1 H⁺ consumed per C fixed during photosynthesis, 1 H⁺ generated per C precipitated during calcification).

Scanning electron microscopy analysis of coccolith morphology

Samples for SEM analysis were filtered on polycarbonate filters (0.8 µm pore-size), dried in a drying cabinet at 50°C for 24 h, then sputter-coated with gold-palladium using an Emitech K550 sputter coater at the Plymouth Electron Microscopy Centre (PEMC). Scanning electron micrographs were produced with both Jeol JSM-6610LV and Jeol JSM-7001F at PEMC. The following categories were used to describe coccolith morphology of *C. braarudii*: normal; minor malformation (element malformation that does not impair interlocking); major malformation (shield malformation that impairs interlocking); malformation type-R (gap in both shields so that the shield elements do not form a closed oval shape); rhomb-like malformation (elements severely malformed displaying rhomb-like crystal morphology); incomplete (closed oval shape, but with incompletely grown shield elements that do not exhibit malformations); type-pH (closed oval shape, but with short shield elements exhibiting malformations) (Fig. 2). Despite a superficial similarity between the categories “incomplete” and “type-pH” that can make them difficult to distinguish in the light microscope, they are easily distinguished in SEM. An incomplete coccolith indicates that coccolith growth was stopped prematurely, but it does not indicate a malfunctioning of the morphogenetic machinery. Therefore, the label “incomplete” should only be applied to coccoliths that do not feature any malformations^{66,67}. A malformed coccolith contains individual elements that have a disrupted morphology, rather than just an abnormal size. An average of ~350 coccoliths was analysed per replicate culture flask, with triplicate cultures examined for each treatment⁶⁸. Coccolith categorization and counting employed standard methodologies as described in detail in⁶⁶.

***C. braarudii* Patch-Clamp Recording and Analysis**

C. braarudii, cells were decalcified by washing cells with Ca^{2+} -free ASW containing 25 mM EGTA^{21, 41}. The recording chamber volume was 1.5 cm³, and solutions exchanged using gravity-fed input and suction output at a rate of 5 cm³ min⁻¹. Patch electrodes were pulled from filamented borosilicate glass (1.5 mm OD, 0.86mm ID) using a P-97 puller (Sutter Instruments, Novato, CA, USA) to resistances of 3-6 MΩ. All external and pipette solutions are described in Supplementary Table 3. Sorbitol was added to pipette solutions to adjust the osmolarity to 1,200 mOsmol kg⁻¹. Liquid junction potentials were calculated using the junction potential tool in Clampex (Molecular Devices, Sunnyvale, CA) and corrected off-line. Whole cell capacitance and seal resistance (leak) were periodically monitored during experiments by applying a <5 mV test pulse. Currents were linear leak subtracted in Clampfit (Molecular Devices, Sunnyvale, CA) using the pre-test seal resistance. Current voltage relations were determined on leak subtracted families by measuring the maximum steady state amplitude (averaging between 10 and 50 ms of the current trace). Reversal potentials were determined by manually measuring the peak tail currents of leak subtracted families of traces and calculating a linear regression versus test voltage. Series resistance was monitored throughout the experiments and whole cell currents were analysed only from recordings in which series resistance varied by less than 15%.

Cloning of *C. braarudii* HV2 into mammalian expression vector

HV2 (CAMPEP_0183380698) from *C. braarudii* was identified by sequence similarity searches of the *C. braarudii* transcriptome (MMETSP0164)⁴⁵ using *C. braarudii* HV1 as a query (ADM25825.1). The predicted coding sequence for HV2 was synthesised (GenScript, Piscataway, NJ) after being codon-optimised for expression in human cells, and sub-cloned into pcDNA3.1-C-eGFP using *HindIII* and *BamHI*. A 6 bp Kozak sequence (GCCACC) was included upstream of the ATG, and the stop codon removed.

Culturing and transfection of HEK293 Cells

HEK293 cells (ATCC CRL-1573) were cultured at 37°C in a humidified atmosphere containing 95% CO₂ in a Dulbecco's modified Eagle's medium (DMEM, Gibco, 12800-017) containing 10% fetal bovine serum (Gibco™ Fetal Bovine Serum, Qualified, Cat. 26140095), 2 mM glutamine, penicillin 100 U mL⁻¹ and streptomycin 100 µg mL⁻¹. Cells were passaged every 3 to 4 d at 1:6 or 1:12 dilutions (cell mm⁻²). HEK293 cells were plated for transfection

onto 35 mm poly-L-lysine coated glass-bottom dishes (35-mm) (www.ibidi.com). Transfections of HEK293 were performed with 1.0 µg of expression vector using Lipofectamine 2000 (ThermoFisher). After 12 to 30 h of incubation, cells were rinsed and maintained with fresh growth media until used for electrophysiological experiments. Cells exhibiting GFP fluorescence were subsequently selected for electrophysiological analysis. HEK293 cells transfected with pCDNA3.1-eGFP alone were used as a control.

HEK293 Whole Cell Patch-Clamp Electrophysiology

Standard whole-cell patch recordings were performed at room temperature with a Multiclamp 700B amplifier (Molecular Devices, Sunnyvale, California) under the control of pClamp10 software (Molecular Devices, Sunnyvale, California). Patch electrodes were pulled from filamented borosilicate glass (1.5 mm OD, 0.86mm ID) using a P-97 puller (Sutter Instruments, Novato, CA, USA) to resistances of 3-6 MΩ. Voltage errors incurred from the liquid junction potentials (LJPs) and series resistance (recorded from the amplifier) were corrected by subtraction post hoc. These corrected voltages were used to plot IV curves and in all subsequent investigations.

Intracellular pH Measurements

For pH_{cyt} measurements, *C. braarudii* cells were loaded with the cell-permeant acetoxymethyl ester for of the pH sensitive fluorescent dye, carboxy SNARF-1. Cells were incubated with SNARF-1 (5 µM) for 20-40 minutes, before being washed with ASW and placed in a poly-lysine coated imaging dish. Cells were imaged using a Nikon Ti Eclipse fluorescence (TIRF) system, equipped with a Photometrics Evolve EM-CCD camera and a Photometrics DV2 beamsplitter. SNARF-1 was excited between 540-560 nm and fluorescence emission was captured at 580 nm (570-590nm) and 630 nm (620-640 nm). Images were recorded at the rate of 3.3 frames s⁻¹ (300 ms exposure). Background fluorescence was minimal and was therefore not subtracted.

pH_{cyt} values at acclimation conditions were derived by measuring SNARF-1 fluorescence in ASW with identical carbonate chemistry to that used for acclimation. For each treatment, pH_{cyt} was measured on a minimum of three independent days. To measure the response of cells to changes in external pH (pH_o), acclimated cells were loaded with SNARF-1 in control ASW at pH 8.15. pH_o was then changed by consecutively perfusing the cells with ASW of pH 6.55, 7.55 and 8.75 for 5 minute time intervals (flow-through approximately 3 mL min⁻¹ in 0.5 mL total volume). In a final step, cells were washed with ASW of pH 8.15 to determine the pH_{cyt}

drift between the beginning and the end of the experiment. If the pH_{cyt} offset was $> 4\%$, measurements were discarded from analysis.

For image processing, the mean fluorescence emission ratio (F_{630}/F_{580}) was determined using a region of interest encompassing the whole cell. We were unable to achieve a satisfactory *in vivo* calibration for SNARF-loaded cells using the nigericin technique, as we found that dye fluorescence was not stable after the addition of this protonophore. We therefore used an *in vitro* calibration, measuring the fluorescence emission ratio (F_{630}/F_{580}) of SNARF-1 (40 μM) in buffer (130 mM KCl, 1 mM MgCl_2 , 15 mM HEPES) of a range of pH (pH 6.75 - 7.5). From the calibration curve, the following relation was obtained ($R^2=0.86$):

$$\text{pH}_{\text{cyt}} = (0.8205 \times \ln(F_{630}/F_{580})) + 7.1101$$

qPCR analysis of gene expression

Quantitative reverse-transcriptase polymerase chain reaction (qPCR) was performed for *HVI* and *HV2* in cultures acclimated to pH 7.55, 8.15 and 8.75 (in triplicates). The expression of two endogenous reference genes (ERGs), *EFL* and *RPS1*, was measured alongside expression of the two target genes. Primers were designed to amplify products approximately 150 bp in length. Primer quality was tested by performing efficiency curves for serial dilutions (1:5) of each primer pair (efficiencies were $> 98\%$, R^2 values > 0.96). RNA was extracted from *C. braarudii* cells using the Isolate II RNA Mini Kit (Bioline) with on column DNA digestion. 30 ml of exponential growth phase culture (approximately 4,000 cells mL^{-1}) was centrifuged at 3800 g for 5 min at 4°C . Quality and quantity of extracted RNA were tested using a Nanodrop 1000 (Thermo Fisher Scientific) (A_{260}/A_{280} ratios > 1.80). cDNA was synthesised from 50 ng RNA using a SensiFAST cDNA Synthesis Kit (Bioline), with a combination of random hexamers and oligo dT primers. No Reverse-Transcriptase Controls (NRTCs) were generated to ensure no DNA contamination had occurred. qPCR runs were performed using a Rotorgene 6000 cycler (Qiagen, USA). Each reaction (20 μL) consisted of 1 μL of cDNA substrate and 19 μL of a SensiFAST No-ROX Kit Master Mix (Bioline, UK). Following primer optimisation, 0.4 μM primer were used for all genes. PCR cycles were run with Rotorgene Q series software, comprising an initial 95°C 2 min hold period, 40 cycles of 95°C denaturing for 5 s, 62°C annealing for 10 s and 72°C extension step (acquisition at end of extension step) for 20 s. A high resolution melt (HRM) curve, $72 - 95^\circ\text{C}$ with 1°C ramp was conducted after amplification to ensure all amplicons had comparable melting temperatures.

For each sample, 1 μ L of cDNA were analysed in technical triplicates (target genes) or duplicates (reference genes). One qPCR plate contained all *HV1* or *HV2* primer reactions (or the NRTCs), as well as the ERG reactions (*EFL* and *RPS1*), control reactions (= *HV1* expressed in the pH 8.15 acclimation), no template controls (NTC) and two positive controls. Stability of the ERG was tested using geNorm⁶⁹. qPCR data were analysed using a efficiency corrected DDCt method, normalizing to the geometric mean of the two ERGs⁶⁹. Expression of *EFL* in all NRTC was at least 10 Ct smaller than the sample.

Phylogenetic Analyses

Previously identified Hv1 sequences from coccolithophores²¹ were used as queries for sequence similarity searches of the available haptophyte transcriptomes within the Marine Microbial Eukaryote Sequencing Project^{45, 70}; reassembled reads NCBI accession PRJEB37117). Further Hv sequences from other representative protist lineages were obtained from the Joint Genome Institute (<https://phycocosm.jgi.doe.gov/phycocosm/home>). Protist Hv sequences possess an extended extracellular loop between transmembrane domains S1 and S2^{21, 44}, enabling the generation of a longer multiple sequence alignment and improved resolution of the haptophyte Hv sequences. Hv sequences from other lineages (e.g. animals) lack the extended S1-S2 loop, although phylogenetic trees constructed with a wider range of eukaryotes exhibited a similar topology. Potential Hv sequences identified by sequence similarity searches were manually inspected using a multiple sequence alignment to assess the presence of conserved residues essential for H⁺ channel function²³. The multiple sequence alignments were then refined using GBLOCKS 0.91b to remove poorly aligned residues⁷¹. Phylogenetic analysis was performed using the maximum likelihood method within MEGAX software⁷² after prior determination of the best substitution model (WAG with gamma and invariant).

Statistical analyses

For coccolith morphologies mean and SEM values were calculated from experimental replicates with a minimum of 350 coccoliths scored for each replicate. For electrophysiology means and SEM values were calculated from individual replicate cells from each treatment, with n numbers given in each Fig. For intracellular pH measurements, differences in pH between treatments were tested with 1-way ANOVA Holm-Sidak post hoc tests. Differences in distribution of pH values between treatments were assessed with 2-sample Kolmogorov-Smirnov tests.

References

1. Doney, S.C., Fabry, V.J., Feely, R.A. & Kleypas, J.A. Ocean acidification: the other CO₂ problem. *Ann Rev Mar Sci* **1**, 169-192 (2009).
2. Rost, B., Zondervan, I. & Wolf-Gladrow, D. Sensitivity of phytoplankton to future changes in ocean carbonate chemistry: current knowledge, contradictions and research directions. *Mar Ecol-Prog Ser* **373**, 227-237 (2008).
3. Jiang, L.Q., Carter, B.R., Feely, R.A., Lauvset, S.K. & Olsen, A. Surface ocean pH and buffer capacity: past, present and future. *Scientific reports* **9**, 18624 (2019).
4. Hofmann, G.E. *et al.* High-frequency dynamics of ocean pH: a multi-ecosystem comparison. *PLoS One* **6**, e28983 (2011).
5. Wootton, J.T., Pfister, C.A. & Forester, J.D. Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset. *Proc Natl Acad Sci U S A* **105**, 18848-18853 (2008).
6. Capone, D. & Hutchins, D. Microbial biogeochemistry of coastal upwelling regimes in a changing ocean. *Nat Geosci* **6**, 711-717 (2013).
7. Waldbusser, G.G. & Salisbury, J.E. Ocean acidification in the coastal zone from an organism's perspective: multiple system parameters, frequency domains, and habitats. *Ann Rev Mar Sci* **6**, 221-247 (2014).
8. Balch, W.M. The Ecology, Biogeochemistry, and Optical Properties of Coccolithophores. *Annual Review of Marine Science*, Vol 10 **10**, 71-98 (2018).
9. Balch, W.M. *et al.* Coccolithophore distributions of the North and South Atlantic Ocean. *Deep-Sea Res Pt I* **151** (2019).
10. Bach, L.T., Riebesell, U., Gutowska, M.A., Federwisch, L. & Schulz, K.G. A unifying concept of coccolithophore sensitivity to changing carbonate chemistry embedded in an ecological framework. *Prog Oceanogr* **135**, 125-138 (2015).
11. Gafar, N.A., Eyre, B.D. & Schulz, K.G. Particulate inorganic to organic carbon production as a predictor for coccolithophorid sensitivity to ongoing ocean acidification. *Limnol Oceanogr Lett* **4**, 62-70 (2019).
12. Kroeker, K.J. *et al.* Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biol* **19**, 1884-1896 (2013).
13. Meyer, J. & Riebesell, U. Reviews and Syntheses: Responses of coccolithophores to ocean acidification: a meta-analysis. *Biogeosciences* **12**, 1671-1682 (2015).
14. Raven, J.A. & Crawford, K. Environmental controls on coccolithophore calcification. *Mar Ecol Prog Ser* **470**, 137-166 (2012).
15. Langer, G. *et al.* Species-specific responses of calcifying algae to changing seawater carbonate chemistry. *Geochem Geophys Geosy* **7** (2006).
16. Bach, L.T. *et al.* Dissecting the impact of CO₂ and pH on the mechanisms of photosynthesis and calcification in the coccolithophore *Emiliana huxleyi*. *New Phytologist* **199**, 121-134 (2013).
17. Sikes, C.S., Roer, R.D. & Wilbur, K.M. Photosynthesis and Coccolith Formation - Inorganic Carbon-Sources and Net Inorganic Reaction of Deposition. *Limnology and Oceanography* **25**, 248-261 (1980).
18. Taylor, A.R., Brownlee, C. & Wheeler, G. Coccolithophore Cell Biology: Chalking Up Progress. *Annu Rev Mar Sci* **9**, 283-310 (2017).
19. Bach, L.T., Bauke, C., Meier, K.J.S., Riebesell, U. & Schulz, K.G. Influence of changing carbonate chemistry on morphology and weight of coccoliths formed by *Emiliana huxleyi*. *Biogeosciences* **9**, 3449-3463 (2012).
20. Bach, L.T., Riebesell, U. & Schulz, K.G. Distinguishing between the effects of ocean acidification and ocean carbonation in the coccolithophore *Emiliana huxleyi*. *Limnology and Oceanography* **56**, 2040-2050 (2011).

21. Taylor, A.R., Chrachri, A., Wheeler, G., Goddard, H. & Brownlee, C. A voltage-gated H⁺ channel underlying pH homeostasis in calcifying coccolithophores. *PLoS Biol* **9**, e1001085 (2011).
22. Suffrian, K., Schulz, K.G., Gutowska, M.A., Riebesell, U. & Bleich, M. Cellular pH measurements in *Emiliania huxleyi* reveal pronounced membrane proton permeability. *New Phytol* **190**, 595-608 (2011).
23. DeCoursey, T.E. Voltage-gated proton channels: molecular biology, physiology, and pathophysiology of the H(V) family. *Physiol Rev* **93**, 599-652 (2013).
24. Fresnel, J. & Probert, I. The ultrastructure and life cycle of the coastal coccolithophorid *Ochrosphaera neapolitana* (Prymnesiophyceae). *European Journal of Phycology* **40**, 105-122 (2005).
25. Liu, Y.W., Eagle, R.A., Aciego, S.M., Gilmore, R.E. & Ries, J.B. A coastal coccolithophore maintains pH homeostasis and switches carbon sources in response to ocean acidification. *Nature communications* **9** (2018).
26. Saez, A.G., Zaldivar-Riveron, A. & Medlin, L.K. Molecular systematics of the Pleurochrysidaceae, a family of coastal coccolithophores (Haptophyta). *Journal of Plankton Research* **30**, 559-566 (2008).
27. White, M.M. *et al.* Calcification of an estuarine coccolithophore increases with ocean acidification when subjected to diurnally fluctuating carbonate chemistry. *Mar Ecol Prog Ser* **601**, 59-76 (2018).
28. Daniels, C.J. *et al.* Species-specific calcite production reveals *Coccolithus pelagicus* as the key calcifier in the Arctic Ocean. *Mar Ecol Prog Ser* **555**, 29-47 (2016).
29. Daniels, C.J., Sheward, R.M. & Poulton, A.J. Biogeochemical implications of comparative growth rates of *Emiliania huxleyi* and *Coccolithus* species. *Biogeosciences* **11**, 6915-6925 (2014).
30. Hernandez, A.S.R. *et al.* Coccolithophore biodiversity controls carbonate export in the Southern Ocean. *Biogeosciences* **17**, 245-263 (2020).
31. Krug, S.A., Schulz, K.G. & Riebesell, U. Effects of changes in carbonate chemistry speciation on *Coccolithus braarudii*: a discussion of coccolithophorid sensitivities. *Biogeosciences* **8**, 771-777 (2011).
32. Muller, M.N. *et al.* Response of the coccolithophores *Emiliania huxleyi* and *Coccolithus braarudii* to changing seawater Mg²⁺ and Ca²⁺ concentrations: Mg/Ca, Sr/Ca ratios and delta Ca-44/40, delta Mg-26/24 of coccolith calcite. *Geochimica Et Cosmochimica Acta* **75**, 2088-2102 (2011).
33. Kottmeier, D.M., Rokitta, S.D. & Rost, B. H⁺-driven increase in CO₂ uptake and decrease in HCO₃⁻ uptake explain coccolithophores' acclimation responses to ocean acidification. *Limnology and Oceanography* **61**, 2045-2057 (2016).
34. Kottmeier, D.M., Rokitta, S.D., Tortell, P.D. & Rost, B. Strong shift from HCO₃⁻ to CO₂ uptake in *Emiliania huxleyi* with acidification: new approach unravels acclimation versus short-term pH effects. *Photosynthesis Research* **121**, 265-275 (2014).
35. Cachao, M. & Moita, M.T. *Coccolithus pelagicus*, a productivity proxy related to moderate fronts off Western Iberia. *Mar Micropaleontol* **39**, 131-155 (2000).
36. Giraudeau, J., Monteiro, P.M.S. & Nikodemus, K. Distribution and malformation of living coccolithophores in the northern Benguela upwelling system off Namibia. *Mar Micropaleontol* **22**, 93-110 (1993).
37. Faucher, G., Riebesell, U. & Bach, L.T. Can morphological features of coccolithophores serve as a reliable proxy to reconstruct environmental conditions of the past? *Clim Past* **16**, 1007-1025 (2020).
38. Riebesell, U. *et al.* Reduced calcification of marine plankton in response to increased atmospheric CO₂. *Nature* **407**, 364-367 (2000).

39. Durak, G.M. *et al.* A role for diatom-like silicon transporters in calcifying coccolithophores. *Nature communications* **7**, 10543 (2016).
40. Gerecht, A.C., Supraha, L., Edvardsen, B., Langer, G. & Henderiks, J. Phosphorus availability modifies carbon production in *Coccolithus pelagicus* (Haptophyta). *J Exp Mar Biol Ecol* **472**, 24-31 (2015).
41. Taylor, A.R. & Brownlee, C. A novel Cl⁻ inward-rectifying current in the plasma membrane of the calcifying marine phytoplankton *Coccolithus pelagicus*. *Plant Physiology* **131**, 1391-1400 (2003).
42. Taylor, A.R., Brownlee, C. & Wheeler, G.L. Proton channels in algae: reasons to be excited. *Trends Plant Sci* **17**, 675-684 (2012).
43. Kigundu, G., Cooper, J.L. & Smith, S.M.E. Hv 1 Proton Channels in Dinoflagellates: Not Just for Bioluminescence? *J Eukaryot Microbiol* **65**, 928-933 (2018).
44. Rodriguez, J.D. *et al.* Identification of a vacuolar proton channel that triggers the bioluminescent flash in dinoflagellates. *PLoS One* **12**, e0171594 (2017).
45. Keeling, P.J. *et al.* The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through Transcriptome Sequencing. *PLoS Biol* **12**, e1001889 (2014).
46. Hong, L., Kim, I.H. & Tombola, F. Molecular determinants of Hv1 proton channel inhibition by guanidine derivatives. *Proc Natl Acad Sci U S A* **111**, 9971-9976 (2014).
47. Fiorini, S., Middelburg, J.J. & Gattuso, J.P. Effects of elevated CO₂ partial pressure and temperature on the coccolithophore *Syracosphaera pulchra*. *Aquatic Microbial Ecology* **64**, 221-232 (2011).
48. Tong, S.Y., Gao, K.S. & Hutchins, D.A. Adaptive evolution in the coccolithophore *Gephyrocapsa oceanica* following 1,000 generations of selection under elevated CO₂. *Global Change Biol* **24**, 3055-3064 (2018).
49. Padin, X.A., Velo, A. & Pérez, F.F. ARIOS: a database for ocean acidification assessment in the Iberian upwelling system (1976–2018). *Earth Syst. Sci. Data* **12**, 2647–2663 (2020).
50. Hermoso, M. & Minoletti, F. Mass and Fine-Scale Morphological Changes Induced by Changing Seawater pH in the Coccolith *Gephyrocapsa oceanica*. *J Geophys Res-Biogeosci* **123**, 2761-2774 (2018).
51. Rigual-Hernandez, A.S. *et al.* Full annual monitoring of Subantarctic *Emiliania huxleyi* populations reveals highly calcified morphotypes in high-CO₂ winter conditions. *Scientific reports* **10** (2020).
52. McClelland, H.L. *et al.* Calcification response of a key phytoplankton family to millennial-scale environmental change. *Scientific reports* **6**, 34263 (2016).
53. von Dassow, P. *et al.* Over-calcified forms of the coccolithophore *Emiliania huxleyi* in high-CO₂ waters are not preadapted to ocean acidification. *Biogeosciences* **15**, 1515-1534 (2018).
54. Honisch, B. *et al.* The geological record of ocean acidification. *Science* **335**, 1058-1063 (2012).
55. Monteiro, F.M. *et al.* Why marine phytoplankton calcify. *Sci Adv* **2**, e1501822 (2016).
56. Henahan, M.J. *et al.* Rapid ocean acidification and protracted Earth system recovery followed the end-Cretaceous Chicxulub impact. *Proc Natl Acad Sci U S A* **116**, 22500-22504 (2019).
57. Bown, P.R., Lees, J.A. & Young, J.R. Calcareous nannoplankton evolution and diversity through time. *Coccolithophores: From Molecular Processes to Global Impact*, 481-508 (2004).
58. Gibbs, S.J. *et al.* Algal plankton turn to hunting to survive and recover from end-Cretaceous impact darkness. *Science Advances* **6** (2020).

59. Hagino, K. *et al.* Re-discovery of a "living fossil" coccolithophore from the coastal waters of Japan and Croatia. *Mar Micropaleontol* **116**, 28-37 (2015).
60. Van de Waal, D.B. & Litchman, E. Multiple global change stressor effects on phytoplankton nutrient acquisition in a future ocean. *Philos Trans R Soc Lond B Biol Sci* **375**, 20190706 (2020).
61. Frada, M.J., Bendif, E.M., Keuter, S. & Probert, I. The private life of coccolithophores. *Perspectives in Phycology* **6**, 11-30 (2018).
62. Xu, Y.Y., Pierrot, D. & Cai, W.J. Ocean carbonate system computation for anoxic waters using an updated CO2SYS program. *Mar Chem* **195**, 90-93 (2017).
63. Kottmeier, D.M., Terbruggen, A., Wolf-Gladrow, D.A. & Thoms, S. Diel variations in cell division and biomass production of *Emiliana huxleyi*-Consequences for the calculation of physiological cell parameters. *Limnology and Oceanography* **65**, 1781-1800 (2020).
64. Menden-Deuer, S. & Lessard, E.J. Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. *Limnology and Oceanography* **45**, 569-579 (2000).
65. Young, J.R. & Ziveri, P. Calculation of coccolith volume and its use in calibration of carbonate flux estimates. *Deep-Sea Research Part 2. Topical Studies in Oceanography* **47**, 9-11 (2000).
66. Langer, G., Oetjen, K. & Brenneis, T. On culture artefacts in coccolith morphology. *Helgoland Mar Res* **67**, 359-369 (2013).
67. Young, J.R. & Westbroek, P. Genotypic Variation in the Coccolithophorid Species *Emiliana-huxleyi*. *Mar Micropaleontol* **18**, 5-23 (1991).
68. Langer, G. & Benner, I. Effect of elevated nitrate concentration on calcification in *Emiliana huxleyi*. *Journal of Nannoplankton Research* **30**, 77-80 (2009).
69. Vandesompele, J. *et al.* Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* **3**, RESEARCH0034 (2002).
70. Johnson, L.K., Alexander, H. & Brown, C.T. Re-assembly, quality evaluation, and annotation of 678 microbial eukaryotic reference transcriptomes. *Gigascience* **8** (2019).
71. Talavera, G. & Castresana, J. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Systematic biology* **56**, 564-577 (2007).
72. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549 (2018).
73. Langer, G. *et al.* Role of silicon in the development of complex crystal shapes in coccolithophores. *New Phytol* (2021).
74. Blanco-Ameijeiras, S. *et al.* Phenotypic Variability in the Coccolithophore *Emiliana huxleyi*. *Plos. One* **11**, e0157697 (2016).
75. Bretherton, L. *et al.* Day length as a key factor moderating the response of coccolithophore growth to elevated pCO₂. *Limnol. Oceanogr.* **64**, 1284-1296 (2019).
76. Langer, G., Nehrke, G., Probert, I., Ly, J. & Ziveri, P. Strain-specific responses of *Emiliana huxleyi* to changing seawater carbonate chemistry. *Biogeosci.* **6**, 2637-2646 (2009).
77. Sett, S. *et al.* Temperature Modulates Coccolithophorid Sensitivity of Growth, Photosynthesis and Calcification to Increasing Seawater pCO₂. *Plos. One* **9**, e88308 (2014).

78. Supraha, L., Gerecht, A.C., Probert, I. & Henderiks, J. Eco-physiological adaptation shapes the response of calcifying algae to nutrient limitation. *Sci. Reports* **5**, 16499 (2015).
79. Zondervan, I., Rost, B. & Riebesell, U. Effect of CO₂ concentration on the PIC/POC ratio in the coccolithophore *Emiliania huxleyi* grown under light-limiting conditions and different daylengths. *J Exp Mar Biol Ecol* **272**, 55-70 (2002).

Data availability

All data is deposited and archived on our institutional servers or relevant genomics databases, according to institutional and funders' requirements. Data will made available from the corresponding authors on request following publication.

Acknowledgements

The work was supported by an ERC Advanced Grant to CB (ERC-ADG-670390) and a NERC award to GLW (NE/N011708/1). Electron microscopy analyses were performed at the PEMC (Plymouth University, UK).

Author contributions

DK, GLW and CB conceived the study. DK was responsible for the majority of experimental analyses, with AC performing electrophysiology and GL performing the SEM analysis of coccolith morphology. KEH performed cloning of *HV2*. DK, AC and GL analysed the data. DK, GLW and CB wrote the manuscript. All data will be made available on request following publication.

The authors declare that they have no competing interests.

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. The data can be provided by the corresponding authors pending scientific review and a completed material transfer agreement. Requests for data should be submitted to the corresponding authors.

Fig. 1. Physiology and H⁺ fluxes of *Coccolithus braarudii* cells grown at different seawater pH. **a** Schematic indicating H⁺ fluxes associated with photosynthesis and calcification in a coccolithophore cell. Whilst many metabolic processes may contribute to the cellular H⁺ budget, these two processes are likely to be the major contributors. In a cell taking up HCO₃⁻, the H⁺ budget is balanced between H⁺ consumed during photosynthesis and H⁺ generated during calcification. In a cell taking up CO₂, 2 H⁺ are produced for each molecule of CaCO₃ produced and H⁺ are no longer consumed during photosynthesis. Excess H⁺ may be removed from the cell by H⁺ transporters in the plasma membrane, such as voltage-gated H⁺ channels

(Hv). Coccolithophores take up both HCO_3^- and CO_2 across the plasma membrane, with increasing proportions of DIC taken up as CO_2 as seawater CO_2 increases³⁴. **b** Growth rate of *C. braarudii* cells acclimated to different seawater pH. n=3 replicates per treatment, line represents polynomial fit to mean. **c** Cellular production of particulate organic carbon (POC) through photosynthesis and particulate inorganic carbon (PIC) through calcification. The optima for both processes are close to the control conditions (pH 8.15). **d** As a consequence of the unequal changes in cellular POC and PIC production across the applied pH values, cellular PIC:POC ratios are minimal at pH 7.55 (≈ 1.0) and maximal at pH 8.45 (≈ 1.8). **e** Calculated net H^+ budgets under the different pH regimes, based on rates of photosynthesis and calcification shown in (c) (See Methods). The concentration of CO_2 in seawater is also shown (dashed line). Estimates are shown for cells using taking up only HCO_3^- or only CO_2 . As *C. braarudii* cells will likely take up a mixture of both DIC species, with a shift towards greater CO_2 usage at elevated CO_2 , the shaded area represents the potential range of H^+ production. Regardless of DIC species used *C. braarudii* produces excess H^+ at all applied pH values, but H^+ production is much lower at pH 7.55 due to the decrease in calcification.

Fig. 2. A unique defect in coccolith morphology occurs at low seawater pH. a Representative scanning electron micrographs of cells acclimated to pH 7.55, 7.85, 8.15 and 8.75. The majority of cells grown at pH ≥ 8.15 had intact coccospheres without crystal or coccolith malformation. The majority of cells grown under acidified conditions produced malformed coccoliths, resulting in abnormal or collapsed coccospheres. Bar = 5 μ m **b** Quantification of coccolith malformations reveal an increasing proportion of defective

coccoliths with seawater acidification. Coccoliths were grouped into morphological categories (see Methods). The morphological categories representing rhomb-like, R-type, major and minor malformations are commonly observed *C. braarudii* cells grown under various stressors⁷³. However, the distinct ‘type-pH’ was only observed in this study and appears unique to low pH (high CO₂) conditions. The counts represent the mean of three independent replicates. A minimum of 350 coccoliths were counted for each replicate. **c** An example of *C. braarudii* cells grown at pH 7.55 exhibiting a high proportion of the distinctive ‘type-pH’ malformations. As the shield elements are not properly formed, the coccoliths are unable to interlock in the normal manner, resulting in the collapse of the coccospheres during preparation for SEM imaging. Bar = 10 μm. **d** The proportion of collapsed coccospheres increases at lower seawater pH. n= 3 replicates per treatment. Error bars represent SE.

Fig. 3: A reduced outward H⁺ current in *C. braarudii* cells acclimated to low seawater pH.

a Estimation of the impact of changes in seawater pH on the proton motive force (PMF) across the plasma membrane. Models of PMF based on measured maximal or minimal pH_{cyt} (pH 6.8 and 7.1: See Fig. 4) in combination with measured minimal and maximal ΔE (-46mV and -28mV, ²¹) suggest that PMF is close to zero at a seawater pH of approximately pH 7.55.

829 Therefore passive H^+ efflux via voltage-gated H^+ channels becomes unfavourable, unless
 830 mediated by excursions of pH_{cyt} (lower cytosolic pH) or V_m (depolarisation). **b**
 831 Electrophysiological measurements of whole cell currents in response to incremental 1 s 10
 832 mV depolarisations from -80 to +40 mV performed in artificial seawater buffered to pH 8.15.
 833 The large outward-directed ion current is predominately carried by H^+ . The maximal current is
 834 much smaller in cells acclimated to pH 7.55. Example of currents from a single cell are shown
 835 for each pH treatment. **c** Mean whole cell currents (plotted as pA/pF vs mV) for acclimated
 836 cells. Outward currents are observed when the plasma membrane is depolarised to potentials
 837 more positive than the equilibrium potential of H^+ (E_{H^+} ; arrow). Cells acclimated to pH 7.55
 838 exhibit a greatly reduced outward current. Values in parentheses represent n, bars represent SE.
 839 **d** The proportion of cells that do not show an outward current (defined as an outward current
 840 <2.5 pA/pF at +45 mV) is greatly increased in cultures acclimated to a seawater pH of 7.55
 841 compared to cultures acclimated to pH 8.15 or 8.75. n=18. **e** Tail current analysis indicating
 842 that reversal potentials (E_{rev}) are close to E_{H^+} and more positive than the E_{K^+} and E_{Cl^-} in all
 843 treatments, suggesting that the observed currents in all treatments are predominately carried by
 844 H^+ .

Fig. 4: Changes in intracellular pH (pH_{cyt}) in response to seawater acidification and alkalisation. **a** Intracellular pH (pH_{cyt}) measured at acclimation pH conditions in *C. braarudii* cells loaded with the pH-responsive fluorescent dye SNARF-AM. Cells acclimated to pH 7.55 show a lower mean pH_{cyt} to those acclimated to pH 8.15 and pH 8.75. $n = 63, 61$ and 36 cells for pH 7.55, 8.15 and 8.75 respectively. 1-way ANOVA, Holm-Sidak post-hoc test, *

p<0.05, ** p<0.01. Box plots indicate mean (open square), median, 25-75th percentiles (box) and 10-90th percentiles (whiskers). **b** pH_{cyt} regulation following rapid changes in external pH. Acclimated cells were perfused with seawater at pH 6.55, 7.55, 8.15 and 8.75 for 2 minutes each to examine their ability to regulate pH_{cyt}. Cells acclimated to pH 8.15 and 8.75 show the rapid adjustment of pH_{cyt} typical of coccolithophore cells ^{21,22}. However, cells acclimated to pH 7.55 show a much lower change in pH_{cyt} (n=25, 61 and 36 respectively for pH 7.55, 8.15 and 8.75). Shaded areas represent SE. **c** Example of rapid changes in intracellular pH (pH_{cyt}) in cells acclimated to pH 8.15. Cells were perfused with ASW pH 8.15 and pHi was monitored as the perfusion was switched to a higher or lower pH. Two representative cells are shown. **d** Example of cells acclimated to pH 7.55 exhibiting little change in pH_{cyt} following changes in external pH. Note that the time course of the perfusion differs slightly from that shown in (C). Two representative cells are shown. **e** Detailed examination of pH_{cyt} recovery during a transition from seawater pH 6.5 to seawater pH 8.15. The frequency histogram indicates the change in pH_{cyt} (shown as $\Delta[H^+]$) in individual cells acclimated to different pH regimes. Whilst nearly all cells acclimated to pH 8.15 and 8.75 exhibit a substantial decrease in $[H^+]$ on transfer from pH 6.5 to higher pH, many cells acclimated to pH 7.55 are unable to respond, indicative of a defect in H⁺ efflux. n=55, 61 and 36 cells). Cells acclimated to pH 7.55 exhibit a significantly different distribution to pH 8.15 or 8.75 (2-sample Kolmogorov-Smirnov test, p<0.05). **f** Proportions of cells exhibiting defective H⁺ efflux in the experiment described in (E). Defective H⁺ efflux was defined as a $\Delta[H^+]$ less than 20 pM.

Fig. 5: Effects of Hv inhibitors on coccolith morphology in *C. braarudii*. **a** Cell growth in the presence of the H⁺ channel inhibitor ZnCl₂ (35 μM) at seawater pH of 8.15. n=3. Error bars = SE. Application of a similar concentration of Zn (30 μM) results in a decrease of approximately 50% in the amplitude of the outward H⁺ current²¹. **b** SEM image of *C. braarudii* cells treated with ZnCl₂ (35 μM) for 5 days showing the presence of many distinctive 'type-pH' coccolith malformations. Note also the collapse of the coccospheres due to the inability of the coccoliths to interlock. Bar = 10 μm. **c** Quantitative analysis of coccolith morphology. Coccoliths were categorised into morphological categories (see Methods). The counts represent the mean of three independent replicate treatments. A minimum of 350 coccoliths were counted for each replicate. Cells exposed to 35 μM Zn exhibit a substantial increase in the proportion of the distinctive type-pH coccolith malformations. **d** Higher resolution SEM images of type-pH morphological defects found in cells exposed to low pH (pH 7.55 from experiment described in Fig. 2) and Zn (35 μM). Bar = 2 μm.