1 **Do forest over- and understory respond to the same environmental variables when viewed at the**
2 **taxonomic and trait level?**

3

4 Kenny Helsen¹, Yeng-Chen Shen¹, Tsung-Yi Lin¹, Chien-Fan Chen^{1,2}, Chu-Mei Huang³,
5 Ching-Feng Li^{4†} & David Zelený¹

6

7 ¹ Institute of Ecology and Evolutionary Biology, National Taiwan University, Taipei, Taiwan

8 ² Division of Botanical Garden, Taiwan Forestry Research Institute, Taipei, Taiwan

9 ³ Division of Silviculture, Taiwan Forestry Research Institute, Taipei, Taiwan

10 ⁴ School of Forestry and Resource Conservation, National Taiwan University, Taipei, Taiwan

11 [†] Deceased 29 November 2019

12

13 Contact author: Kenny Helsen

14 Corresponding author: David Zelený

15

16 Running head: filtering of forest over- and understory

17 **Abstract**

18 While the relative importance of climate filtering is known to be higher for woody species
 19 assemblages than herbaceous assemblage, it remains largely unexplored whether this pattern
 20 is also reflected between the woody overstory and herbaceous understory of forests. While
 21 climatic variation will be more buffered by the tree layer, the understory might also respond
 22 more to small-scale soil variation, next to experiencing additional environmental filtering due
 23 to the overstory's effects on light and litter quality. For (sub)tropical forests, the understory
 24 often contains a high proportion of fern and lycophyte species, for which environmental
 25 filtering is even less well understood. We explored the proportional importance of climate
 26 proxies and soil variation on the species, functional trait and (functional) diversity patterns of
 27 both the forest overstory and fern and lycophyte understory along an elevational gradient
 28 from 850 to 2100 m a.s.l. in northern Taiwan. We selected nine functional traits expected to
 29 respond to soil nutrient or climatic stress for this study and furthermore verified whether they
 30 were positively related across vegetation layers, as expected when driven by similar
 31 environmental drivers. We found that climate was a proportionally more important predictor
 32 than soil for the species composition of both vegetation layers and trait composition of the
 33 understory. The stronger than expected proportional effect of climate for the understory was
 34 likely due to fern and lycophytes' higher vulnerability to drought, while the high importance
 35 of soil for the overstory seemed driven by deciduous species. The environmental drivers
 36 affected different response traits in both vegetation layers, however, which together with
 37 additional overstory effects on understory traits, resulted in a strong disconnection of
 38 community-level trait values across layers. Interestingly, species and functional diversity
 39 patterns could be almost exclusively explained by climate effects for both vegetational layers,
 40 with the exception of understory species richness. This study illustrates that environmental
 41 filtering can differentially affect species, trait and diversity patterns and can be highly

divergent for forest overstory and understory vegetation, and should consequently not be extrapolated across vegetation layers or between composition and diversity patterns.

Key words

Trait-environment relationship, ferns, lycophytes, forest overstory, forest understory, functional traits, functional diversity, species richness, subtropical montane cloud forest

Introduction

Although the effects of environmental or abiotic filtering on plant communities is often reflected in their species composition and richness, it is believed that this filtering mainly acts on the plants' functional traits, rather than directly on the species' identities (Lavorel and Garnier 2002, Kraft et al. 2015). Many studies have consequently observed strong trait – environment relationships across ecosystems (Wright et al. 2005, Ordoñez et al. 2009, Bruelheide et al. 2018). Not only functional trait composition, but also functional diversity can be affected by environmental filtering (Aros-Mualin et al. 2021). Functional diversity is, more specifically, expected to be reduced under environmentally stressful conditions, since only a limited number of functionally similar species will be able to establish (cf. trait underdispersion) (Weiher and Keddy 1995). The spatial extent at which environmental filtering occurs furthermore seems to differ among different environmental factors (Mokany and Roxburgh 2010, Bruelheide et al. 2018). While climatic factors mainly drive differences in species and trait composition across relatively large spatial scales, at smaller spatial scales, community (trait) composition is mainly structured by local-scale factors, such as variation in soil conditions (Bruelheide et al. 2018).

A recent study focusing on large-scale trait-environment patterns, suggested that the relative importance of different drivers even differs between woody and herbaceous species

67 assemblages, with climatic variation more strongly impacting woody plant communities
68 (Šímová et al. 2018). Consequently, within multi-layered forest ecosystems, environmental
69 filtering might also differentially affect the woody overstory and herbaceous understory. For
70 example, climate or (micro)climatic-related topography might more strongly impact the
71 overstory, because the overstory is fully exposed to climatic variation, while the understory
72 experiences buffered climatic variation under the protection of the forest canopy (Šímová et
73 al. 2018, De Frenne et al. 2019). The species composition of the overstory might also be more
74 likely to be filtered by more coarse-scale soil variation compared to that of the herbaceous
75 understory, whose roots will be much more localized. The understory might additionally
76 experience filtering due to small-scale environmental variation caused directly by variation in
77 the overstory. Several studies have, for example, shown the impact of overstory related light
78 availability and leaf litter on understory species composition (Komiya et al. 2001, Wang et
79 al. 2019, Majasalmi and Rautiainen 2020), trait composition (Maes et al. 2020) and
80 functional diversity (Chabrierie et al. 2010).

81 Surprisingly little studies have, however, tried to quantify the similarities in
82 environmental filtering between the over- and understory layers of forests (however see
83 Ruokolainen et al. 2007, Rogers et al. 2008, Salazar et al. 2012). While this comparison is
84 complicated for many temperate forest types, due to the often limited overstory species
85 diversity, (sub)tropical forests offer an ideal study system to explore these relationships. In
86 this study we focus on the proportional impact of several climate proxies (topography and
87 ground fog frequency) and soil variation on the over- and the understory of the subtropical
88 montane forests of northern Taiwan, along an elevation gradient ranging from 870 to 2130 m
89 a.s.l.

90 Interestingly, the understory of subtropical montane forests contains a high diversity
91 of fern and lycophyte species, next to angiosperms. For this reason, we focus specifically on

the understory fern and lycophyte species in this study, while excluding angiosperms. Some studies have observed similar leaf trait-trait (Karst and Lechowicz 2007, Lin et al. 2020) and trait-environment relationships (Kessler et al. 2007, Kluge and Kessler 2007, Zhu et al. 2016, Campany et al. 2019) for ferns as for angiosperms, suggesting that functional patterns are similarly structured and thus comparable across both phylogenetic groups. Environmental filtering and trait-environment relationships nevertheless remain less well understood for fern and lycophyte communities, compared to angiosperm communities (Kessler et al. 2016). Effects of environmental filtering on understory fern community functional diversity has, for example, been observed in a few studies (Tanaka and Sato 2015, Zhang et al. 2017, Sessa et al. 2018), but not in others (Kluge and Kessler 2011, Aros-Mualin et al. 2021).

To allow optimal trait comparisons across both vegetation layers, we measured the same nine functional leaf traits for both overstory woody species and understory fern and lycophyte species. These nine traits were specifically chosen for their expected link to soil nutrient and/or climatic stress. Using this dataset, we addressed the following research questions:

- Do climate proxies and soil factors explain equal proportions of variation in the over- and understory community-level species and functional trait composition along the elevation gradient?
- Are these potential climate proxy and soil filtering processes also reflected in species and functional diversity along the elevation gradient?
- Can we find additional filtering of the understory species and trait composition due to variation in the overstory?

Methods

Study design

The study was performed in the Wulai district, New Taipei City, northern Taiwan, along an elevational transect ranging from Mt. Meilu (870 m a.s.l., 24.85°N 121.53°E) to Mt. Taman (2130 m a.s.l., 24.71°N 121.45°E, Fig. 1). The geological substrates mainly consist of argillite, shale, slate, sandstone and phyllite (Central Geological Survey, MOEA), with soils of low pH and high soil organic matter content. The study region is characterized by a humid subtropical climate ('Cfa' climate sensu the Köppen-Geiger system), with an average annual temperature of 16.1°C and average annual precipitation of 2070 mm (Lalashan weather station, 1374 m a.s.l., 24.68°N 121.40°E). Most precipitation falls during the summer, although the region is also affected by the north-eastern winter monsoon. The forest vegetation along the gradient varies from lower elevation *Pyrenaria-Machilus* subtropical winter monsoon forest, across mid-elevation *Quercus* montane evergreen broad-leaved cloud forest to higher elevation *Chamaecyparis* montane mixed cloud forest (Li et al. 2013).

Along the transect, six elevation zones were delimited at 850, 1100, 1350, 1600, 1850 and 2100 m a.s.l. $\pm$ 50 m (Fig. 1). At each elevation zone, ten 10 m $\times$ 10 m plots were established across a secondary gradient in aspect and topography, ranging from the northeast facing (windward) to the southwest facing slope (leeward) across the ridge. Plots were positioned at least 50 m apart. Due to logistic constraints, only 9 plots were established at the 1850 m elevation zone, resulting in a total of 59 plots. For each plot, we recorded the presence of all woody species (angiosperm and gymnosperm shrubs and trees) taller than 2 m and with diameter in breast height (DBH) $\geq$ 1 cm (i.e. the 'overstory' vegetation). We also recorded the presence of all terrestrial (non-epiphytic) fern and lycophyte species, which together make up 66% of the plot-level herbaceous understory species richness in our study (i.e. the 'understory' vegetation).

140

141 Climate proxies

142 We measured two topographic variables (i.e. elevation and heat load) that are known to relate
 143 to (micro)climatic conditions. The exact elevation of each plot was measured using GPS
 144 (GPSMAP 64st, Garmin, USA). To calculate heat load, we first measured the slope of each
 145 plot with a clinometer (SUUNTO PM-5/360 PC Clinometer, SUUNTO, Finland) from the
 146 upper plot edge, and the aspect of each plot with a compass (SILVA, Sweden). Aspect was
 147 then transformed into folded aspect, which was defined as the azimuth angle difference
 148 between the aspect and 45° (McCune and Keon 2002). We calculated heat load based on
 149 folded aspect and slope using equation 2 of McCune and Keon (2002). We additionally
 150 extracted average annual ground fog frequency for each plot from the ground fog frequency
 151 raster map for Taiwan (250 m per pixel resolution), developed by Schulz et al. (2017), based
 152 on MODIS satellite data. Ground fog frequency is expected to be an important environmental
 153 factor impacting cloud forest vegetation through effects on temperature, light availability,
 154 evapotranspiration and water availability (Fahey et al. 2016).

155

156 Soil variables

157 For each plot, four soil samples of the top 0-10 cm were collected and pooled together for soil
 158 analysis. Each pooled soil sample was analyzed for pH, carbon:nitrogen ratio, total nitrogen,
 159 and phosphorous, potassium, magnesium, zinc, calcium, manganese, copper and iron content.
 160 See Appendix S1 for details of the soil chemical analysis.

161 To prevent collinearity among soil predictor variables, we performed a principal
 162 component analysis (PCA) on all measured soil chemical variables (after logarithmic
 163 transformation of Ca, C:N ratio, Mg, Mn, P and Zn and subsequent standardization to zero
 164 mean and unit standard deviation for all variables, Appendix S2). The three retained soil PC

axes together explained 81.4% of the total variation. The first axis (termed ‘soil NPK’ in the text) reflected a gradient in nitrogen, phosphorous and potassium, next to several other micronutrients. The second axis (‘soil pH’) reflected an increase in pH and soil manganese, while the third axis (‘soil Cu’) was most strongly related to soil copper (positively) and calcium and iron (negatively) (Appendix S2).

We additionally measured soil depth at four positions in each plot using a 30 cm long soil depth meter (diameter 0.6 cm) and averaged values per plot. Soil more than 30 cm deep was recorded as 35 cm (16.9 % of the plots). Soil rockiness (the percentage content of rocks in the top 0-10 cm of the soil) was also estimated.

Functional traits

Nine leaf traits were measured for 91 overstory woody species (sampled during 10/2014 and 12/2016-09/2018) and 48 understory fern and lycophyte species (sampled during 5/2017-10/2018), including all common species present in the plots and covering 74.0% and 63.2% of our over- and understory species pools, respectively (Appendix S3). The nine measured traits consisted of specific leaf area (SLA, mm²/mg), leaf dry matter content (LDMC, mg/g), area-based leaf chlorophyll content (SPAD units), leaf nitrogen content (leaf N, mg/g), leaf area (cm²), leaf thickness (Lth, mm), equivalent water thickness (EWT, mg/mm²), leaf ¹³C/¹²C stable isotope ratio (δ¹³C, ‰) and leaf ¹⁵N/¹⁴N stable isotope ratio (δ¹⁵N, ‰). Note that the first four traits are related to the leaf economics spectrum (LES) (cf. Wilson et al. 1999, Wright et al. 2004). Trait measurements largely followed standard protocols (Pérez-Harguindeguy et al. 2013), with some modifications for fern and lycophyte species. See Appendix S1 for trait measurement details.

These traits are expected to vary along climatic and soil variation gradients, because of their expected links to either soil nutrient stress (LES traits, δ¹⁵N, Lth; Wright and Cannon

2001, Hodgson et al. 2011), temperature or climatic stress (LES traits, leaf area; Wright et al. 2005, Dong et al. 2020) or drought (leaf area, Lth, EWT, $\delta^{13}\text{C}$; Medeiros et al. 2019, Maréchaux et al. 2020). While high values of $\delta^{13}\text{C}$ are known to reflect high long-term water-use efficiency, and thus drought tolerance (Farquhar et al. 1982, Pérez-Harguindeguy et al. 2013), $\delta^{15}\text{N}$ relates to a plant's nitrogen acquisition strategy (Craine et al. 2015). More specifically, $\delta^{15}\text{N}$ values around 0 ‰ usually indicate nitrogen fixation, while values around -2, -3 and -6 ‰ indicate plant nitrogen acquisition through arbuscular, ericoid and ectomycorrhiza, respectively (Craine et al. 2015). Note that EWT expresses the water mass content of a fresh leaf per unit leaf area, and is sometimes also called 'succulence' (Mantovani 1999, Féret et al. 2019).

All leaf level trait values were averaged at the species level after exclusion of leaf-level outliers (Z-score > 2.5 at the species level), as we assumed these values to most likely occur from measurement errors. This resulted in the exclusion of 0.61 % and 0.16 % of the trait values from the full leaf $\times$ trait matrix for the over- and understory, respectively. For leaf area we did not remove 'outliers', since all trait values could be verified for measurement errors. Missing trait values for nine overstory species were replaced by mean trait values across all overstory species, prior to further data analysis.

Data analysis

We calculated the plot-level community mean (CM) values for each trait, as the average trait across all species present in the plot. Since only presence-absence data was collected, CM trait values were not weighted by species abundance. Species richness was calculated as the number of species present in the plot. We calculated two measures of functional diversity for each plot. Scheiner et al. (2017) has recently proposed to express functional diversity by the two parameters trait dispersion (M') and trait 'evenness' or

equability (${}^1E(T)$), next to species richness. While M' quantifies ‘magnitude’, i.e. the amount of difference in trait values among species in a community, ${}^1E(T)$ quantifies ‘variability’, i.e. the extent to which species are equally different from each other in trait values. Both measures are based on pairwise trait dissimilarities. Unlike the more traditionally used functional diversity parameters (i.e. functional richness, evenness and divergence), M' and ${}^1E(T)$ are independent from species richness and evenness, and thus solely reflect trait magnitude and variability, respectively (Scheiner 2019, Kosman et al. 2021). M' and ${}^1E(T)$ were calculated with the R script provided by Malavasi et al. (2018), based on the formulas of Scheiner et al. (2017), using Gower dissimilarity on the species $\times$ trait matrix (with traits standardized to Z-scores) to construct the trait dissimilarity matrix. Leaf area was logarithmically transformed prior to CM and functional diversity calculation. CM $\delta^{15}N$ was additionally transformed using the equation $(x + 4)^2$ for the overstory dataset and $\log(x + 2.5)$ for the understory dataset to obtain symmetrical distribution.

Before statistical analysis, we calculated variance inflation factors (VIF) to identify potential collinearity issues, separately among the different climate proxies and among soil variables. Collinearity was identified for folded aspect and heat load ($VIF > 5$), and thus we excluded folded aspect from all statistical models. We consequently retained three climate proxies (elevation, heat load and ground fog frequency) and five soil variables (soil depth, soil rockiness and three soil PC axes, namely soil NPK, soil pH and soil Cu). Heat load was squared and soil rockiness was square root transformed prior to statistical analyses to improve symmetrical distribution of their values.

To test the effects of climate proxies and soil variables on the over- and understory species composition, we performed separate redundancy analyses (RDA) on the respective plot $\times$ species matrices. These RDA models were repeated on the plot $\times$ species matrices including only species for which traits were measured. Similar RDA models were performed

for the over- and understory plot $\times$ CM trait matrices, with CM of traits standardized to Z-scores (CWM-RDA, Nygaard and Ejrnæs 2004). The overstory trait RDA model was also repeated after excluding all deciduous species, to assess the potential effect of deciduous species on the trait patterns.

After assuring global significance of each of the global RDA models containing all environmental predictors, we performed variation partitioning on each model to assess the proportions of variation in each plot $\times$ species or plot $\times$ CM trait matrix explained by either climate proxies or soil variables, expressed by adjusted R^2 . Next, we performed forward model selection among all environmental predictors, based on adjusted R^2 -values (conditional effects) and their significance assessed using Monte Carlo permutation tests (9999 permutations). This model selection allowed us to assess which environmental variables were most important predictors for each species and CM trait dataset. We are aware that the CWM-RDA method is prone to inflated Type I error rate when using the standard Monte Carlo permutation test (Šmilauer and Lepš 2014, Zelený 2018). Since there is no published solution to this problem, however, we nonetheless use these tests. Before forward model selection, we again used VIFs to ensure that no collinearity occurred among any of the combined climate and soil variables. All multivariate analyses were performed with the ‘vegan’ R package (Oksanen et al. 2017).

We additionally performed partial redundancy analysis for the understory plot $\times$ species and plot $\times$ CM trait datasets, to assess the potential additional effects of the overstory. To achieve these models, we first performed two PCA’s, one on the overstory plot $\times$ species matrix and one on the overstory plot $\times$ CM trait matrix. For both models, we only retained PCA axes for which the eigenvalue was larger than the average inertia (Kaiser-Guttman criterion, Ibanez 1973). Hence, we retained 15 and 3 axes for the overstory species and CM trait PCA’s, respectively. Next, we performed a partial RDA (pRDA) on the understory plot

× species matrix with all retained overstory species PCA axes as explanatory variables, while partialling out the effects of all measured environmental variables, and assessed the model’s overall significance. Similarly, we assessed the effect (including overall significance) of the retained overstory trait PCA axes on the understory plot × trait matrix while partialling out all climate proxies and soil variables.

The effects of climate proxies and soil variables on species and functional diversity were assessed using generalized linear models. For species richness (a count variable) we used a Poisson probability distribution and log link function. For M' and ${}^1E(T)$ we used a gamma probability distribution with inverse link function, since these variables consist of positive, continuous, right skewed data. Due to the nonlinear relationship between elevation and understory species richness, and fog frequency and overstory ${}^1E(T)$, we included the quadratic term for elevation and fog frequency to the respective models. We performed variation partitioning on the full diversity models to quantify the proportion of explained deviance by climate proxies and soil variables. Each full model was then reduced using a AIC-based comparison of all predictor-subset combinations of the full model, using the ‘dredge’ function in the ‘MuMIn’ R package (Barton 2019). Model assumptions were checked using the ‘DHARMA’ R package (Hartig 2021).

If traits and diversity are shaped by similar environmental drivers for the under- and overstory along our gradient, we expect them to be positively correlated between vegetation layers. To verify this, we performed simple linear regressions between each understory CM trait (response) and overstory CM trait (predictor). Based on the scatterplots for these regressions, we also included a quadratic term for the CM SLA model (i.e. understory CM $SLA \sim \text{overstory CM SLA} + (\text{overstory CM SLA})^2$). These regressions were also performed for overstory CM traits excluding deciduous species. Similar regressions were additionally constructed for S , M' and ${}^1E(T)$ between under- and overstory. All p-values of the pairwise

regressions were corrected for type-I error inflation using the false discovery rate method (Benjamini and Hochberg 1995) with the ‘p.adjust’ function in the ‘stats’ R package. All analyses were performed with R version 4.0.5.

Results

Species composition of both over- and understory species was significantly affected by climate proxy and soil variation, together explaining 29.9% and 30.7% of the total variation (i.e. adjusted R^2), respectively (Fig. 2A&B). Variation partitioning indicated that for both over- and understory species composition, climate proxies explained a higher relative proportion of this variation (overstory: 80.4%, understory: 85.0%) than soil (overstory: 54.9%, understory: 54.2%). However, for both vegetation layers, around one third of the explained variation was shared by climate proxies and soil (Fig. 2A&B). The final RDA models after forward model selection retained all climate proxy variables and soil pH for both the overstory and understory species composition. For the overstory species composition, soil NPK and soil Cu were additionally retained (Table 1, Fig. 3A&B). RDA and variation partitioning results were barely affected when RDA was performed on the plot $\times$ species matrices including only species for which traits were measured (Appendices S4 & S5). The pRDA indicated that an additional 7.1% ($p < 0.001$) of the total variation in understory species composition was explained purely by overstory species composition.

CM trait composition variation was also related to climate proxies and soil for both over- (adjusted $R^2 = 49.3\%$) and understory species (adjusted $R^2 = 42.2\%$) (Fig. 2C&D). Variation partitioning showed that climate proxies were the most important predictors of CM trait variation for understory species (77.4% of the total explained variation for climate proxies vs. 58.5% for soil), while soil explained most variation for overstory CM traits (58.4% for climate proxies vs. 80.5% for soil). For both vegetation layers’ trait composition,

around one third of the explained variation was shared by climate proxies and soil (Fig. 2C&D). After forward model selection, elevation, ground fog frequency, soil pH and soil Cu were retained for both CM trait RDAs, while soil NPK was additionally retained for the overstory CM trait RDA (Fig. 3C&D). The RDA results were largely similar for overstory traits excluding deciduous species. However, climate proxies became more important than soil after variance partitioning (Appendices S4 & S5).

The results of the CM trait RDAs furthermore suggest an increase in CM LDMC with elevation for both vegetation layers and a decrease in CM of leaf area and $\delta^{15}\text{N}$ with elevation for the overstory. High ground fog frequency seems to result in lower CM EWT for both vegetation layers, while high soil pH seems to result in high CM of SLA and leaf N and low CM Lth for the overstory, and high CM leaf N for the understory (Fig. 3C&D). Note that these patterns can be inferred from Fig. 3 because the involved environmental variables and traits were well represented by the first two RDA axes (high axis loadings). Evaluation of the third RDA axis loadings for the understory furthermore suggests that high soil pH is also related to high CM leaf N for the understory (results not shown). The pRDA showed that the overstory trait composition could explain an additional 4.5% ($p = 0.011$) of the variation in the understory trait composition after partialling out the effects of climate proxies and soil variables.

CM of LDMC and leaf N were positively related, while CM of Lth and $\delta^{15}\text{N}$ were negatively related between both vegetation layers. CM SLA showed a parabolic relationship, with the highest values for the understory at intermediate values for the overstory species. CM of leaf area, chlorophyll content, $\delta^{13}\text{C}$ and EWT, on the other hand, were not significantly related between both vegetation layers (Fig. 4, Appendix S6). Excluding deciduous species for the overstory weakened, but did not change the direction of all trait -

trait relationships, except for leaf thickness, for which the negative relationship strengthened (Appendices S6 & S7).

Compared to soil, climate proxy variables explained more of the variation in species richness, functional divergence and functional equability for both vegetation layers (Table 1, Appendix S8). Soil variation nonetheless contributed a small additional proportion to the total explained variation (< 20%) in the understory species richness and functional equability for both vegetation layers (Table 1, Appendix S8). While elevation was the strongest climate proxy predictor for all diversity measures of both the over- and understory, the direction of these relationship differed between both vegetation layers. While overstory species richness declined with elevation, understory species richness was highest at intermediate elevation (Fig. 5A). Functional dispersion increased with elevation for both vegetation layers (Fig. 5C), while functional equability increased with elevation for the overstory, and decreased for the understory (Fig. 5E). Soil pH was the strongest soil predictor for understory species richness (positive relationship) (Fig. 5B) and functional equability (negative relationship) (Fig. 5F).

Species richness and functional equability were not significantly related for both vegetation layers, while functional dispersion was significantly positively related between the over- and understory (Appendices S6 & S9).

Discussion

Species and trait composition

Climate proxies were more strongly related than soil to the species composition of the overstory. This result was expected, since, unlike the understory, the overstory is fully exposed to climatic variation (Šímová et al. 2018, De Frenne et al. 2019). Surprisingly, however, also the species and trait composition of the fern and lycophyte understory were mainly shaped by climate proxies. This nonetheless agrees with the literature, where climate

often strongly impacts fern trait composition (e.g. Kessler et al. 2007, Kluge and Kessler 2007, Sessa and Givnish 2014). It has been suggested that the limited potential for controlling evaporation makes ferns more sensitive to drought than angiosperms (Brodribb and Holbrook 2004, Zhang et al. 2014). This could thus explain why elevation (temperature) and ground fog frequency (relative humidity) rather than soil variation were the main drivers of fern species and trait composition in our study.

The high relative importance of elevation is not surprising, considering the quite steep elevational gradient of 1260 m in our dataset. The trait responses followed expectations of more resource conservative (e.g. high LMDC) and smaller leaves with increasing elevation for both vegetation layers (Wright et al. 2005, Dong et al. 2020). The high ground fog frequency likely increased water availability, as suggested by drought-related trait states (high EWT for both vegetation layers and Lth for understory) in low fog plots in our study (cf. Medeiros et al. 2019, Maréchaux et al. 2020). Frequent fog can, however, additionally reduce light availability (up to 10-50%) and local temperature (up to 3-6°C) (Lai et al. 2006, Reinhardt and Smith 2008) and negatively impact photosynthesis by preventing leaf transpiration and promoting the growth of epiphyllous lichens and algae. While these conditions are highly suitable for fern species, which often seem adapted to low light and high water availability (Sessa and Givnish 2014, Hernández-Rojas et al. 2020), they likely present less suitable growing conditions for most tree species by hampering photosynthesis (Fahey et al. 2016). These potential different responses of both vegetation layers were, however, not reflected in their respective trait composition. More detailed future work using field-based fog or relative humidity measurements should be used to further explore these potential effects.

Soil did nevertheless still impact the species and trait composition of both vegetation layers and was, unexpectedly, more important than climate proxies in structuring overstory

trait composition. The higher importance of soil for overstory traits compared to species composition was not due to differences in the species included in both models, since the species RDA model including only species for which traits were measured gave similar results. The ‘soil pH’ ordination axis was the most important soil driver, and likely better reflects plant nutrient availability than soil NPK in the typically highly acidic soils with low decomposition rates and high soil organic matter of cloud forests (Fahey et al. 2016). The high nutrient levels following standard soil analysis of these soils likely reflect nutrients trapped in undecomposed organic matter, rather than plant available nutrients. This could explain the presence of acquisitive leaf traits (high SLA and leaf N) in less acidic soils for both vegetation layers (Wright and Cannon 2001). The lower relative importance of soil variables compared to climate proxies for the understory, on the other hand, might be because understory fern species respond to more small-scale soil variation than that measured at the plot-level in this study.

Interestingly, if deciduous species are excluded from the overstory, climate proxies become the most important driver of overstory trait composition, due to reduced importance of soil pH. Deciduous woody species are more common on steep wind-exposed slopes in northern Taiwan, probably because of their ability to avoid environmental stress during winter by shedding their leaves. These steep slopes usually have higher soil pH in comparison with less steep or flat ridges at the same elevation, perhaps due to surface erosion removing more acidic soil and litter and increasing availability of more cation-rich weathered parental rock material. This contradiction between seemingly stress-adapted niches and more acquisitive LES traits (cf. low leaf longevity, Wright et al. 2004) of deciduous species, makes it difficult to assess if these soil pH patterns are caused by LES nutrient-availability or wind-exposure and slope. This shows how soil, topography and climate can interact in their impact

on species- and trait composition, as also illustrated by the high overlap in explained variation by climate proxies and soil for all variation partitioning analyses.

We found that the overstory can act as an additional filter on the understory's species and trait composition, through potential light availability or leaf litter effects, as observed in previous studies (Komiyama et al. 2001, Wang et al. 2019, Maes et al. 2020, Majasalmi and Rautiainen 2020). The observational nature of our study does, however, not allow us to assess the causality of this effect. Alternatively, the relationship between overstory and understory could be caused by unmeasured environmental conditions affecting species and trait composition of both layers simultaneously.

Despite some similarities in the environmental responses of both vegetation layers, only two of the nine measured traits showed a significant positive relationship between over- and understory. This illustrates that environmental filtering differs substantially for trees and ferns and is most clearly illustrated by Lth. For ferns, Lth seemed to respond to drought (cf. Kluge and Kessler 2007), but was linked to low nutrient availability in trees (cf. Read et al. 2006), resulting in a negative correlation between both. Direct impact of the overstory on the understory could also have shaped unexpected trait relationships. The negative correlation for $\delta^{15}\text{N}$ might, for example, reflect niche differentiation among vegetation layers to prevent competition for different nitrogen sources. The quadratic relationship for SLA could also be due to overstory impact on the understory. While nutrient availability mainly structured SLA for the overstory, for the understory, SLA might be affected by a combination of nutrient limitation under low overstory SLA (cf. Kessler et al. 2007) and more strong light competition due to shading under high overstory SLA.

Species and functional diversity

Species richness was most strongly affected by elevation for both vegetation layers. Both the decrease in overstory species richness and humped-shaped relationship for understory fern species along elevation are consistent with previous studies in (sub)tropical forests (Kluge and Kessler 2011, Qian and Ricklefs 2016, Hernández-Rojas et al. 2020). At lower elevation, lower water availability is expected to reduce diversity of drought-sensitive lifeforms such as ferns and lycophytes (Kessler et al. 2011, Weigand et al. 2020). At higher elevations, on the other hand, species richness of both ferns and angiosperms will be reduced by the stronger climatic stress associated with lower temperatures (Kessler et al. 2011). Interestingly, soil explained a much higher proportion of variation in understory than overstory species richness, thus showing the opposite pattern as observed for species and trait composition. The impact of soil productivity (soil pH), next to climate on fern species richness is nonetheless in agreement with previous work (Tuomisto et al. 2014, Weigand et al. 2020). Not surprisingly, species richness was not correlated between the two vegetation layers, further illustrating that species richness is shaped by different environmental drivers for overstory trees and understory ferns and lycophytes.

Functional diversity patterns suggested that for the overstory, the community trait composition is not experiencing increased trait convergence among species with elevation, as expected under increased environmental filtering (Weiher and Keddy 1995). On the contrary, species trait overlap seems to be reduced, resulting in higher average trait distances among species (M'), combined with high equal spacing of species in the trait space ($^1E(T)$), a pattern usually attributed to increased importance of competition among species (Kraft et al. 2008). Functional equability was nonetheless highest at intermediate ground fog frequency levels, potentially reflecting trait clustering (environmental filtering) due to drought stress and high air humidity at each respective end of the ground fog frequency gradient.

For the understory, trait composition seemingly clustered in separate distinct trait sets, potentially reflecting (environmental) filtering of a few alternative trait combinations for ferns at high elevation (increasing M' , decreasing $^1E(T)$). A similar pattern was also observed for terrestrial fern communities along four tropical elevation gradients (Aros-Mualin et al. 2021). Elevation related most strongly to functional dispersion in both vegetation layers and functional equability for the overstory. This again mirrors the results of Aros-Mualin et al. (2021), who found that temperature was the most important predictor of fern functional diversity. Functional equability of the understory, on the other hand, was almost equally strongly affected by soil variation and climate proxies. This soil (pH)-driven environmental trait filtering for the fern understory (cf. Zhang et al. 2017, Sessa et al. 2018), but not the overstory, thus mirrors the impact of soil on the species richness patterns in our study.

Conclusions

Both climate proxies and soil were important predictors of species and trait composition of both vegetation layers. The stronger effects of climate proxies for understory ferns and lycophytes compared to overstory trees is likely due to their higher vulnerability to drought. The environmental drivers furthermore seem to affect very different response traits in both vegetation layers, which together with additional overstory effects on understory traits, results in a disconnection of community-level trait values across layers. Interestingly, the relative importance of soil and climate proxies on species or trait composition cannot be extrapolated to species or trait diversity, which showed very different patterns. This study illustrates that environmental filtering can differentially affect species, trait and diversity patterns and can be highly divergent for forest overstory and understory vegetation.

Acknowledgements

We would like to thank to all volunteers who contributed to the field work and lab trait measurements. Hsin-Yen Teng and Kun-Sung Wu helped with the logistics of the field work, specimen determination and dataset preparation. This study was supported by Ministry of Science and Technology, Taiwan (106-2621-B-002-003-MY3 and 109-2811-B-002-644).

Literature cited

- Aros-Mualin, D., S. Noben, D. N. Karger, C. I. Carvajal-Hernández, L. Salazar, A. Hernández-Rojas, J. Kluge, M. A. Sundue, M. Lehnert, D. Quandt, and M. Kessler. 2021. Functional diversity in ferns is driven by species richness rather than by environmental constraints. *Frontiers in Plant Science* 11:615723.
- Barton, K. 2019. MuMIn: Multi-Model Inference. R package version 1.43.6. <https://CRAN.R-project.org/package=MumIn>.
- Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B* 57:289–300.
- Brodribb, T. J., and N. M. Holbrook. 2004. Stomatal protection against hydraulic failure: a comparison of coexisting ferns and angiosperms. *New Phytologist* 162:663–670.
- Bruehlheide, H., J. Dengler, O. Purschke, J. Lenoir, B. Jiménez-Alfaro, S. M. Hennekens, Z. Botta-Dukát, M. Chytrý, R. Field, et al. 2018. Global trait–environment relationships of plant communities. *Nature Ecology and Evolution* 2:1906–1917.
- Campany, C. E., L. Martin, and J. E. Watkins Jr. 2019. Convergence of ecophysiological traits drives floristic composition of early lineage vascular plants in a tropical forest floor. *Annals of Botany* 123:793–803.

509 Chabrierie, O., J. Loinard, S. Perrin, R. Saguez, and G. Decocq. 2010. Impact of *Prunus*
510 *serotina* invasion on understory functional diversity in a European temperate forest.
511 Biological Invasions 12:1891–1907.

512 Craine, J. M., E. N. J. Brookshire, M. D. Cramer, N. J. Hasselquist, K. Koba, E. Marin-
513 Spiotta, and L. Wang. 2015. Ecological interpretations of nitrogen isotope ratios of
514 terrestrial plants and soils. Plant and Soil 396:1–26.

515 Dong, N., I. C. Prentice, I. J. Wright, B. J. Evans, H. F. Togashi, S. Caddy-retalic, F. A.
516 Mcinerney, B. Sparrow, E. Leitch, and A. J. Lowe. 2020. Components of leaf-trait
517 variation along environmental gradients. New Phytologist 228:82–94.

518 Fahey, T. J., R. E. Sherman, and E. V. J. Tanner. 2016. Tropical montane cloud forest:
519 environmental drivers of vegetation structure and ecosystem function. Journal of
520 Tropical Ecology 32:355–367.

521 Farquhar, G. D., M. H. O’Leary, and J. A. Berry. 1982. On the relationship between carbon
522 isotope discrimination and the intercellular carbon dioxide concentration in leaves.
523 Australian Journal of Plant Physiology 9:121–137.

524 Féret, J.-B., G. le Maire, S. Jay, D. Berveiller, R. Bendoula, G. Hmimina, A. Cheraiet, J. C.
525 Oliveira, F. J. Ponzoni, et al. 2019. Estimating leaf mass per area and equivalent water
526 thickness based on leaf optical properties: Potential and limitations of physical modeling
527 and machine learning. Remote Sensing of Environment 231:110959.

528 De Frenne, P., F. Zellweger, F. Rodríguez-Sánchez, B. R. Scheffers, K. Hylander, M. Luoto,
529 M. Vellend, K. Verheyen, and J. Lenoir. 2019. Global buffering of temperatures under
530 forest canopies. Nature Ecology and Evolution 3:744–749.

531 Hartig, F. 2021. DHARMA: residual diagnostics for hierarchical (multi-level/mixed)
532 regression models. R package version 0.4.1.

533 Hernández-Rojas, A. C., J. Kluge, T. Krömer, C. Carvajal-Henández, L. Silva-Mijangos, G.

534 Miehe, M. Lehnert, A. Weigand, and M. Kessler. 2020. Latitudinal patterns of species
535 richness and range size of ferns along elevational gradients at the transition from tropics
536 to subtropics. *Journal of Biogeography* 47:1383–1397.

537 Hodgson, J. G., G. Montserrat-Martí, M. Charles, G. Jones, P. Wilson, B. Shipley, M. Sharafi,
538 B. E. L. Cerabolini, J. H. C. Cornelissen, et al. 2011. Is leaf dry matter content a better
539 predictor of soil fertility than specific leaf area? *Annals of Botany* 108:1337–1345.

540 Ibanez, F. 1973. Méthode d’analyse spatio-temporelle du processus d’échantillonnage en
541 planctologie, son influence dans l’interprétation des données par l’analyse en
542 composantes principales. *Annales de l’Institut océanographique (Paris)* 49:83–111.

543 Karst, A. L., and M. J. Lechowicz. 2007. Are correlations among foliar traits in ferns
544 consistent with those in the seed plants? *New Phytologist* 173:306–312.

545 Kessler, M., D. N. Karger, and J. Kluge. 2016. Elevational diversity patterns as an example
546 for evolutionary and ecological dynamics in ferns and lycophytes. *Journal of*
547 *Systematics and Evolution* 54:617–625.

548 Kessler, M., J. Kluge, A. Hemp, and R. Ohlemüller. 2011. A global comparative analysis of
549 elevational species richness patterns of ferns. *Global Ecology and Biogeography*
550 20:868–880.

551 Kessler, M., Y. Siorak, M. Wunderlich, and C. Wegner. 2007. Patterns of morphological leaf
552 traits among pteridophytes along humidity and temperature gradients in the Bolivian
553 Andes. *Functional Plant Biology* 34:963–971.

554 Kluge, J., and M. Kessler. 2007. Morphological characteristics of fern assemblages along an
555 elevational gradient: patterns and causes. *Ecotropica* 13:27–43.

556 Kluge, J., and M. Kessler. 2011. Phylogenetic diversity, trait diversity and niches: species
557 assembly of ferns along a tropical elevational gradient. *Journal of Biogeography*
558 38:394–405.

- 559 Komiyama, A., S. Kato, and M. Teranishi. 2001. Differential overstory leaf flushing
560 contributes to the formation of a patchy understory. *Journal of Forest Research* 6:163–
561 171.
- 562 Kosman, E., S. M. Scheiner, and H. R. Gregorius. 2021. Severe limitations of the FEve
563 metric of functional evenness and some alternative metrics. *Ecology and Evolution*
564 11:123–132.
- 565 Kraft, N. J. B., P. B. Adler, O. Godoy, E. C. James, S. Fuller, and J. M. Levine. 2015.
566 Community assembly, coexistence and the environmental filtering metaphor. *Functional*
567 *Ecology* 29:592–599.
- 568 Kraft, N. J. B., R. Valencia, and D. D. Ackerly. 2008. Functional traits and niche-based tree
569 community assembly in an Amazonian forest. *Science* 322:580–583.
- 570 Lai, I.-L., S.-C. Chang, P.-H. Lin, C.-H. Chou, and J.-T. Wu. 2006. Climatic characteristics
571 of the subtropical mountainous cloud forest at the Yuanyang lake long-term ecological
572 research site, Taiwan. *Taiwania* 51:317–329.
- 573 Lavorel, S., and E. Garnier. 2002. Predicting changes in community composition and
574 ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional Ecology*
575 16:545–556.
- 576 Li, C. F., M. Chytrý, D. Zelený, M. Y. Chen, T. Y. Chen, C. R. Chiou, Y. J. Hsia, H. Y. Liu,
577 S. Z. Yang, C. L. Yeh, J. C. Wang, C. F. Yu, Y. J. Lai, W. C. Chao, and C. F. Hsieh.
578 2013. Classification of Taiwan forest vegetation. *Applied Vegetation Science* 16:698–
579 719.
- 580 Lin, D., S. Yang, P. Dou, H. Wang, F. Wang, S. Qian, G. Yang, L. Zhao, Y. Yang, and N.
581 Fanin. 2020. A plant economics spectrum of litter decomposition among coexisting fern
582 species in a sub-tropical forest. *Annals of Botany* 125:145–155.
- 583 Maes, S. L., M. P. Perring, L. Depauw, M. Bernhardt-Römermann, H. Blondeel, G. Brūmelis,

584 J. Brunet, G. Decocq, J. den Ouden, et al. 2020. Plant functional trait response to
585 environmental drivers across European temperate forest understorey communities. *Plant*
586 *Biology* 22:410–424.

587 Majasalmi, T., and M. Rautiainen. 2020. The impact of tree canopy structure on understory
588 variation in a boreal forest. *Forest Ecology and Management* 466:118100.

589 Malavasi, M., V. Bartak, M. L. Carranza, P. Simova, and A. T. R. Acosta. 2018. Landscape
590 pattern and plant biodiversity in Mediterranean coastal dune ecosystems: Do habitat loss
591 and fragmentation really matter? *Journal of Biogeography* 45:1367–1377.

592 Mantovani, A. 1999. A method to improve leaf succulence quantification. *Brazilian Archives*
593 *of Biology and Technology* 42:9–14.

594 Maréchaux, I., L. Saint-André, M. K. Bartlett, L. Sack, and J. Chave. 2020. Leaf drought
595 tolerance cannot be inferred from classic leaf traits in a tropical rainforest. *Journal of*
596 *Ecology* 108:1030–1045.

597 McCune, B., and D. Keon. 2002. Equations for potential annual direct incident radiation and
598 heat load. *Journal of Vegetation Science* 13:603–606.

599 Medeiros, C. D., C. Scoffoni, G. P. John, M. K. Bartlett, F. Inman-Narahari, R. Ostertag, S.
600 Cordell, C. Giardina, and L. Sack. 2019. An extensive suite of functional traits
601 distinguishes Hawaiian wet and dry forests and enables prediction of species vital rates.
602 *Functional Ecology* 33:712–734.

603 Mokany, K., and S. H. Roxburgh. 2010. The importance of spatial scale for trait-abundance
604 relations. *Oikos* 119:1504–1514.

605 Nygaard, B., and R. Ejrnæs. 2004. A new approach to functional interpretation of vegetation
606 data. *Journal of Vegetation Science* 15:49–56.

607 Oksanen, A. J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R.
608 Minchin, R. B. O. Hara, G. L. Simpson, et al. 2017. *vegan: community ecology package.*

609 R package version 2.4-2.

610 Ordoñez, J. C., P. M. van Bodegom, J.-P. M. Witte, I. J. Wright, P. B. Reich, and R. Aerts.

611 2009. A global study of relationships between leaf traits, climate and soil measures of

612 nutrient fertility. *Global Ecology and Biogeography* 18:137–149.

613 Pérez-Harguindeguy, N., S. Diaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S.

614 Bret-Harte, W. K. Cornwell, J. M. Craine, et al. 2013. New handbook for standardised

615 measurement of plant functional traits worldwide. *Australian Journal of Botany* 61:167–

616 234.

617 Qian, H., and R. E. Ricklefs. 2016. Out of the tropical lowlands: latitude versus elevation.

618 *Trends in Ecology and Evolution* 31:738–741.

619 Read, J., G. D. Sanson, M. de Garine-Wichatitsky, and T. Jaffré. 2006. Sclerophylly in two

620 contrasting tropical environments: low nutrients vs. low rainfall. *American Journal of*

621 *Botany* 93:1601–1614.

622 Reinhardt, K., and W. K. Smith. 2008. Impacts of cloud immersion on microclimate,

623 photosynthesis and water relations of *Abies fraseri* (Pursh.) Poiret in a temperate

624 mountain cloud forest. *Oecologia* 158:229–238.

625 Rogers, D. A., T. P. Rooney, D. Olson, and D. M. Waller. 2008. Shifts in Southern

626 Wisconsin forest canopy and understory richness, composition, and heterogeneity.

627 *Ecology* 89:2482–2492.

628 Ruokolainen, K., H. Tuomisto, M. J. Macía, M. A. Higgins, and M. Yli-halla. 2007. Are

629 floristic and edaphic patterns in Amazonian rain forests congruent for trees,

630 pteridophytes and Melastomataceae? *Journal of Tropical Ecology* 23:13–25.

631 Salazar, L., J. Homeier, C. Leuschner, M. Kessler, and J. Kluge. 2012. Altitudinal change in

632 biomass, productivity and leaf functional traits in the Ecuadorian Andes: Comparing

633 terrestrial ferns with trees. Pages 89–111 *Unraveling the causal links between ecosystem*

634 productivity measures and species richness using terrestrial ferns in Ecuador. PhD thesis.

635 Scheiner, S. M. 2019. A compilation of and typology for abundance-, phylogenetic- and

636 functional-based diversity metrics. bioRxiv:1–29.

637 Scheiner, S. M., E. Kosman, S. J. Presley, and M. R. Willig. 2017. Decomposing functional

638 diversity. *Methods in Ecology and Evolution* 8:809–820.

639 Schulz, H. M., C.-F. Li, B. Thies, S.-C. Chang, and J. Bendix. 2017. Mapping the montane

640 cloud forest of Taiwan using 12 year MODIS-derived ground fog frequency data. *PLOS*

641 *ONE* 12:e0172663.

642 Sessa, E. B., S. M. Chambers, D. Li, L. Trotta, L. Endara, J. G. Burleigh, and B. Baiser. 2018.

643 Community assembly of the ferns of Florida. *American Journal of Botany* 105:549–564.

644 Sessa, E. B., and T. J. Givnish. 2014. Leaf form and photosynthetic physiology of *Dryopteris*

645 species distributed along light gradients in eastern North America. *Functional Ecology*

646 28:108–123.

647 Šímová, I., C. Violle, J.-C. Svenning, J. Kattge, K. Engemann, B. Sandel, R. K. Peet, S. K.

648 Wiser, B. Blonder, et al. 2018. Spatial patterns and climate relationships of major plant

649 traits in the New World differ between woody and herbaceous species. *Journal of*

650 *Biogeography* 45:895–916.

651 Šmilauer, P., and J. Lepš. 2014. *Multivariate Analysis of Ecological Data using CANOCO 5*.

652 Cambridge University Press, Cambridge.

653 Tanaka, T., and T. Sato. 2015. Taxonomic, phylogenetic and functional diversities of ferns

654 and lycophytes along an elevational gradient depend on taxonomic scales. *Plant Ecology*

655 216:1597–1609.

656 Tuomisto, H., G. Zuquim, and G. Cárdenas. 2014. Species richness and diversity along

657 edaphic and climatic gradients in Amazonia. *Ecography* 37:1034–1046.

658 Wang, H., M. Zhang, and H. Nan. 2019. Abiotic and biotic drivers of species diversity in

understory layers of cold temperate coniferous forests in North China. *Journal of Forestry Research* 30:2213–2225.

Weigand, A., S. Abrahamczyk, I. Aubin, C. Bitá-Nicolae, H. Bruelheide, C. I. Carvajal-Hernández, D. Cicuzza, L. E. Nascimento da Costa, J. Csiky, et al. 2020. Global fern and lycophyte richness explained: How regional and local factors shape plot richness. *Journal of Biogeography* 47:59–71.

Weiher, E., and P. A. Keddy. 1995. Assembly rules, null models, and trait dispersion: new questions from old patterns. *Oikos* 74:159–164.

Wilson, P. J., K. Thompson, and J. G. Hodgson. 1999. Specific leaf area and leaf dry matter content as alternative predictors of plant strategies. *New Phytologist* 143:155–162.

Wright, I. J., and K. Cannon. 2001. Relationships between leaf lifespan and structural defences in a low-nutrient, sclerophyll flora. *Functional Ecology* 15:351–359.

Wright, I. J., P. B. Reich, J. H. C. Cornelissen, D. S. Falster, P. K. Groom, K. Hikosaka, W. Lee, C. H. Lusk, Ü. Niinemets, J. Oleksyn, N. Osada, H. Poorter, D. I. Warton, and M. Westoby. 2005. Modulation of leaf economic traits and trait relationships by climate. *Global Ecology and Biogeography* 14:411–421.

Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares, T. Chapin, J. H. C. Cornelissen, et al. 2004. The worldwide leaf economics spectrum. *Nature* 428:821–827.

Zelený, D. 2018. Which results of the standard test for community-weighted mean approach are too optimistic? *Journal of Vegetation Science* 29:953–966.

Zhang, H., S. Zhu, R. John, R. Li, H. Liu, and Q. Ye. 2017. Habitat filtering and exclusion of weak competitors jointly explain fern species assemblage along a light and water gradient. *Scientific Reports* 7:298.

Zhang, S.-B., M. Sun, K.-F. Cao, H. Hu, and J.-L. Zhang. 2014. Leaf photosynthetic rate of

684 tropical ferns is evolutionarily linked to water transport capacity. PLOS ONE 9:e84682.
685 Zhu, S.-D., R.-H. Li, J. Song, P.-C. He, H. Liu, F. Berninger, and Q. Ye. 2016. Different leaf
686 cost–benefit strategies of ferns distributed in contrasting light habitats of sub-tropical
687 forests. Annals of Botany 117:497–506.
688

Tables

Table 1. Parameter estimates of the reduced redundancy analyses (RDA) for species and trait composition and the reduced generalized linear models for species richness (S), functional divergence (M') and functional equability ($^1E(T)$), for the overstory and understory datasets separately. Test statistic (F) for each retained predictor and full model adjusted R^2 provided. For soil principal components 'soil NPK', 'soil pH' and 'soil Cu', see Table 1. $^{(*)}0.10 \geq p\text{-value} > 0.05$; $^*0.05 \geq p\text{-value} > 0.01$; $^{**}0.01 \geq p\text{-value} > 0.001$; $^{***}0.001 \geq p\text{-value}$. sqrt = square root transformation, sq = squared transformation of environmental factor. El = elevation, fog = ground fog frequency.

climate proxies				soil					R ²
elevation	fog	heat load ^{sq}	soil depth	soil NPK	soil pH	soil Cu	soil rockiness ^{sqrt}		
overstory									
species composition	10.4 ^{***}	4.7 ^{***}	1.6 ^(*)	-	1.7 [*]	3.2 ^{***}	1.6 ^(*)	-	29.2
CM trait composition	10.0 ^{***}	4.2 ^{**}	-	-	4.6 ^{**}	13.3 ^{***}	7.7 ^{***}	-	49.3
S	18.1 ^{***}	3.1 ^(*)	10.0 ^{**}	-	-	-	-	-	36.5
M'	97.5 ^{***}	-	2.5	-	-	-	-	-	64.2
¹ E(T)	5.3 [*]	fog: 12.1 ^{**} fog ² : 9.2 ^{**}	5.0 [*]	-	3.6 ^(*)	-	-	-	48.5
understory									
species composition	10.4 ^{***}	7.3 ^{***}	1.9 [*]	-	-	3.9 ^{***}	-	-	29.8
CM trait composition	21.0 ^{***}	5.4 ^{***}	-	-	-	3.9 ^{**}	7.2 ^{***}	-	42.2
S	el: 11.4 ^{**} el ² : 69.7 ^{***}	-	-	-	5.1 [*]	6.4 ^{*sq}	-	4.3 [*]	65.7
M'	14.4 ^{***}	4.0 ^(*)	-	-	-	-	-	-	31.7
¹ E(T)	8.2 ^{**}	-	7.5 ^{**}	3.2 ^(*)	-	9.8 ^{**}	4.6 [*]	-	46.5

Figure legends

Figure 1. Location of the six elevation zones at which vegetation plots were established in northern Taiwan.

Figure 2. Venn diagrams visualizing the variation partitioning between climate proxy and soil variable effects on A. the overstory plot × species matrix, B. the understory plot × species matrix, C. the overstory plot × community mean (CM) trait matrix, D. the understory plot × CM trait matrix, using redundancy analysis (RDA). Numbers in the Venn diagrams correspond to the relative proportions of the total explained variation. The total explained variance (adjusted R^2) is also presented.

Figure 3. Triplots for the redundancy analyses (RDA) after forward model selection, exploring the effect of environment (climate proxies and soil) on A. the overstory plot × species matrix, B. the understory plot × species matrix, C. the overstory plot × community mean (CM) trait matrix, D. the understory plot × CM trait matrix. Plots visualized as points, with colors indicating plot elevation, 40% most common species with 20% best fit to the RDA axes are visualized as codes (see Appendix S3), CM trait vectors visualized as vector tips with names in italics, environmental variables visualized as vectors. Note that ‘soil NPK’, ‘soil pH’ and ‘soil Cu’ refer to the first three soil PCA axes, respectively (Table 1). Chl = leaf chlorophyll content, $\delta^{13}\text{C}$ = the leaf $^{13}\text{C}/^{12}\text{C}$ stable isotope ratio, $\delta^{15}\text{N}$ = the leaf $^{15}\text{N}/^{14}\text{N}$ stable isotope ratio, EWT = equivalent water thickness, Lth = leaf thickness.

Figure 4. Scatterplots for pairwise regressions between plot-level overstory and understory community mean (CM) trait values. Solid regression line + SE presented for

725 significant regressions, dashed line for marginally significant regression (see Appendix S6).
 726 Each datapoint corresponds to one vegetation plot, with colors indicating plot elevation. $\delta^{13}\text{C}$
 727 = the leaf $^{13}\text{C}/^{12}\text{C}$ stable isotope ratio, $\delta^{15}\text{N}$ = the leaf $^{15}\text{N}/^{14}\text{N}$ stable isotope ratio, EWT =
 728 equivalent water thickness, Lth = leaf thickness.

729

730 **Figure 5. Scatterplots between three species and functional diversity measures, on the**
 731 **one hand and elevation and the second soil PCA axis (soil pH), on the other hand. Blue**
 732 circles and lines = overstory species, green triangles and lines = understory species. Solid
 733 regression line + SE presented for significant regressions (Table 1).

734 **Figures**

735
736 **Figure 1.**

Figure 2.

Figure 4.

Figure 5.