Generation and diversification of recombinant monoclonal antibodies for studying mitosis

Keith F. DeLuca^{1*}, Jeanne E. Mick^{1*}, Amy L. Hodges¹, Wanessa C. Lima², Lori Sherman³, Kristin L. Schaller⁴, Steven M. Anderson^{3,5}, Ning Zhao¹, Timothy J. Stasevich^{1,6}, Dileep Varma⁷, Gary J. Gorbisky⁸, Jakob Nilsson⁹, and Jennifer G. DeLuca^{1#}

¹Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins, CO, USA

²Geneva Antibody Facility, Faculty of Medicine, University of Geneva, Geneva, Switzerland

³CU Cancer Center Cell Technologies Shared Resource, University of Colorado Cancer Center, Anschutz Medical Campus, Aurora, CO

⁴Department of Pediatric Hematology, Oncology and Bone Marrow Transplant, University of Colorado Anschutz Medical Campus, Aurora, CO

⁵Department of Pathology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.

⁶Cell Biology Center and World Research Hub Initiative, Tokyo Institute of Technology, Yokohama, Japan

⁷Department of Cell and Developmental Biology, Feinberg School of Medicine, Northwestern University, Chicago, USA

⁸Cell Cycle and Cancer Biology Research Program, Oklahoma Medical Research Foundation, Oklahoma City, OK, USA

⁹The Novo Nordisk Foundation Center for Protein Research, University of Copenhagen, Faculty of Health and Medical Sciences, Copenhagen, Denmark

*Authors contributed equally

#For correspondence: Jennifer G. DeLuca, jdeluca@colostate.edu

Abstract

Antibodies are indispensable tools used for a large number of applications in both foundational and translational bioscience research; however, there are drawbacks to using traditional antibodies generated in animals. These include a lack of standardization leading to problems with reproducibility, high costs of antibodies purchased from commercial sources, and ethical concerns regarding the large number of animals used to generate antibodies. To address these issues, we have developed practical methodologies and tools for generating low-cost, high-yield preparations of recombinant monoclonal antibodies and antibody fragments directed to protein epitopes from primary sequences. We describe these methods here, as well as approaches to diversify monoclonal antibodies, including customization of antibody species specificity, generation of genetically encoded small antibody fragments, and conversion of single chain antibody fragments (e.g. scFv) into full-length, bivalent antibodies. This study focuses on antibodies directed to epitopes important for mitotic cell division; however, the methods and reagents described here are applicable to antibodies and antibody fragments for use in any field.

41 **Introduction**

42 Antibodies are used in a diverse array of applications in the biomedical sciences including detection of
 43 biomolecules in cells, tissues, and biological fluids; protein purification; functional depletion of proteins
 44 from cells and cell extracts; medical diagnostics, and therapeutic medicine. While these reagents are
 45 essential for almost all areas of research in the biosciences, there are drawbacks to using traditional
 46 antibodies generated in animals. First, there are growing concerns regarding reproducibility, and this is
 47 in part due to a lack of standardized and thoroughly defined immunological reagents. In many cases,
 48 antibodies are incompletely characterized, not well understood at the molecular level, and variable in
 49 performance across lots (Bradbury and Plückthun, 2015; Bordeaux et al., 2010; Bradbury et al., 2018;
 50 Baker, 2015; Weller, 2016). Second, the continued availability of traditionally generated antibodies is not
 51 guaranteed, as the existence of such reagents depends on active maintenance and storage, or continued
 52 production in animals (Cosson and Hartley, 2016). Third, traditional, commercially available antibodies
 53 are expensive. For many researchers these costs are prohibitive and in turn, significantly limit productivity
 54 and research innovation. Finally, a large number of vertebrate animals are used for the generation of
 55 traditional antibodies for biomedical research, which presents ethical concerns (Gray et al., 2020; Gray
 56 et al., 2016; Leenaars et al., 1998).

57
 58 In recent years, it has been possible to sequence monoclonal antibodies from purified antibody samples
 59 and from hybridoma cell lines such that their primary amino acid composition is explicitly identified (Lima
 60 et al., 2020; Cosson and Hartley, 2016; Vazquez-Lombardi et al., 2016). In addition, techniques have
 61 been developed in which antibodies to nearly any antigen can be isolated through clonal selection of
 62 sequence-defined antibody fragments (Gavilondo and Larrick, 2000; Saeed et al., 2017; Alfaleh et al.,
 63 2020; Almargo et al., 2019; Laustsen et al., 2021). Using these approaches, the generation of sequence-
 64 defined recombinant antibodies and antibody fragments is feasible, which circumvents many of the
 65 problems listed above regarding traditionally generated antibodies. First, using recombinant antibodies
 66 generated from an invariant primary sequence increases reagent reproducibility. Second, after a primary
 67 sequence is determined, recombinant antibodies and their derivatives are accessible in perpetuity. Third,
 68 recombinant antibodies can be produced in large quantities using low-cost expression and purification
 69 systems, such that researchers can produce large-scale yields of recombinant antibodies for a fraction
 70 of the cost of antibodies purchased from commercial sources. In addition, plasmids are easily distributed
 71 for direct expression in cell line of choice. Finally, the use of recombinant antibodies significantly reduces
 72 the number of animals required for antibody production.

73
 74 An additional advantage of recombinant antibodies is the potential for increased reagent versatility. With
 75 the primary amino acid sequence of an antibody in-hand, researchers can diversify the original reagent
 76 and create derivative tools such as antibody fragments, that can be genetically fused to other protein

molecules, such as fluorophores, to generate custom tools with diverse functionalities. Here we describe methods and tools for generating low-cost, high-yield preparations of recombinant monoclonal antibodies and antibody fragments from mammalian cell culture, benchmarking the approach with mitotic epitopes. Furthermore, we describe straightforward and accessible approaches to diversify immunological reagents including customization of antibody species specificity, generation of genetically encoded small antibody fragments, and conversion of single chain antibody fragments (e.g. scFv) into full-length, bivalent antibodies. While this study focuses on antibodies relevant to cell division and mitosis, these approaches are widely applicable for antibodies and antibody fragments across fields.

Results

Generation of recombinant monoclonal antibodies to mitotic targets

In the process of mitosis, chromosomes must properly segregate into two daughter cells in order to maintain genomic integrity. Kinetochores are structures built at the primary constriction of mitotic chromosomes which mediate attachments to spindle microtubules and are largely responsible for both powering and regulating chromosome congression and segregation (Musacchio and Desai, 2017). The primary factor that connects kinetochores to microtubules is the kinetochore associated NDC80 complex, and within this complex the Ndc80/Hec1 subunit serves as the direct link to microtubules (DeLuca and Musacchio, 2011; Varma and Salmon, 2013; Wimbish and DeLuca, 2020). Research in many labs utilizes the Hec1 monoclonal antibody “9G3”, which was originally generated in mice to a purified protein fragment encompassing amino acids 56-632 of the human Hec1 protein (Chen et al., 1997). The specific epitope was later mapped using peptide array analysis to amino acids 200-215, a region which resides within the well-ordered calponin homology domain of the protein (DeLuca et al., 2006). While this antibody is commercially available, the quality of the antibody is variable between lots, which is not uncommon in the case of commercially produced immunological reagents (Pozner-Moulis, 2007; Garg and Loring, 2017; Katzman et al., 2017; Bradbury and Plückthun, 2015; Bordeaux et al., 2010; Bradbury et al., 2018; Baker, 2015; Weller, 2016).

To ensure its continued consistency, we sequenced the 9G3 Hec1 mouse monoclonal antibody (Rapid Novor, Kitchener, Ontario, Canada) using tandem mass spectrometry with W-ion isoleucine and leucine determination (Johnson et al., 1987; Zhokhov et al., 2017). A 100 µg sample of purified monoclonal antibody was used as the source material for sequencing. Figure 1A shows the antibody sequence and annotations for the hyper-variable regions (HVR, also known as complementarity determining regions or CDRs) in green, the framework regions (FR) within the variable regions in blue, as well as the constant regions (CR1, CR2, and CR3 in the heavy chain; CR in the light chain) in orange. The heavy chain was identified as class IgG2a and the light chain as kappa (Figure 1A, B). We next generated geneblocks encoding for both heavy chain and light chain sequences optimized for expression in human cells. The

Figure 1. Generation of a recombinant Hec1 monoclonal antibody. (A) Sequence data obtained for the Hec1 monoclonal antibody is annotated for the heavy chain (HC) and light chain (LC) variable regions (HC=light blue; LC=dark blue), hypervariable regions (green), constant regions (HC=light orange; LC=dark orange), and the flexible hinge (dark gray). (B) rMAb-Hec1^{ms} antibody structure and nomenclature. (C) Cloning, transfection, and purification scheme. Heavy and light chain coding regions were cloned into separate plasmids for transfection into human Expi293F cells. Cell media containing secreted antibodies was collected and applied to a Protein A Sepharose column. After column washing, antibodies were eluted using a low pH buffer. (D) Purified rMAb-Hec1^{ms} antibody was serially diluted and run on a 12% SDS- polyacrylamide gel. The prominent band that runs above the 245 kDa molecular mass marker is likely a population of non-denatured antibody. The bands running at ~63 kDa and ~50 kDa are glycosylated and non-glycosylated heavy chains, respectively. The antibody light chain runs at ~25 kDa. (E) HeLa cells immunostained with the purified rMAb-Hec1^{ms} antibody. Cells were also stained with DAPI to detect chromosomes. (F) Immunoblot of control and Hec1 siRNA-depleted HeLa cell lysates. Increasing amounts of lysates are shown, and the blot is probed with the purified rMAb-Hec1^{ms} antibody. (G) HeLa cell treated with Hec1 siRNA and stained with the rMAb-Hec1^{ms} antibody. Cell is also stained with DAPI to detect chromosomes. Scale bars are 10 µm.

127
 128 geneblocks were cloned separately into GFP-N1 vectors with the GFP removed. Signal peptide
 129 sequences for each chain (Burton, 1994; Yu et al., 2006) were cloned N-terminal to the heavy chain and
 130 light chain sequences to direct the expressed antibody for secretion into the cell media (Figure 1C). The
 131 heavy and light chain-containing expression vectors were co-transfected at a ratio of 2:3 (heavy
 132 chain:light chain) into 30 mL cultures of human HEK293 cells grown in suspension (Expi293F cells) using
 133 PEI transfection reagent (Figure 1C). The cell supernatant was collected five days post-transfection, and
 134 the antibody was purified on a Protein A Sepharose column (Figure 1C). From 30 mL cell suspension
 135 cultures, purification yields of the recombinant Hec1 antibody (rMAb-Hec1^{ms}) ranged from 0.5-1.4 mg of
 136 purified antibody (Figure 1D). To test the quality and specificity of rMAb-Hec1^{ms}, we carried out
 137 immunofluorescence staining and found that in HeLa cells, rMAb-Hec1^{ms} recognized kinetochores during
 138 all phases of mitosis, as expected (Figure 1E). We also analyzed mitotic cell lysates from control cells
 139 and cells treated with Hec1 siRNA by immunoblotting, and as shown in Figure 1F, rMAb-Hec1^{ms}
 140 recognized a single band at ~72 kDa (corresponding to the predicted mass of the 642 amino acid protein,
 141 73.9 kDa) in control lysates, but not in lysates depleted of Hec1. Consistently, rMAb-Hec1^{ms} antibodies
 142 did not recognize kinetochores in cells treated with Hec1 siRNA (Figure 1G).

143
 144 We implemented a similar strategy for generating a recombinant phospho-specific antibody to a
 145 conserved, repeating four amino acid motif ("MELT" motif) in the kinetochore scaffolding protein KNL1
 146 whose phosphorylation at the threonine (T) residue by the mitotic kinase Mps1 is required to recruit a
 147 suite of spindle assembly checkpoint proteins to kinetochores (Shepherd et al., 2012; Yamagishi et al.,
 148 2012; London et al., 2012), and a recombinant antibody to CENP-C, an inner kinetochore protein
 149 required for kinetochore assembly (Kixmoeller et al., 2020; Hara and Fukagawa, 2020; Navarro and
 150 Cheeseman, 2021). The KNL1 pMELT monoclonal antibody (Fisher Scientific) was generated in rabbit
 151 against a peptide containing phosphorylated Thr943 and phosphorylated Thr1155 (Nijenhuis et al.,
 152 2014). Sequence results revealed that the pMELT rabbit antibody belongs to class IgG with a kappa-
 153 class light chain (Supplemental Figure 1A). The CENP-C monoclonal antibody was generated in mice
 154 (Abcam), and sequencing identified the heavy chain as IgG2b, and the light chain as belonging to the
 155 kappa class (Supplemental Figure 1B). Both the pMELT and CENP-C antibody heavy and light chains
 156 were cloned into expression vectors as described for the rMAb-Hec1^{ms} antibody. Expression plasmids
 157 were transfected into human Expi293F cells, and the antibodies were purified on Protein A Sepharose
 158 columns. Immunofluorescence experiments revealed that the rMAb-pMELT^{rb} antibody recognized
 159 kinetochores, and as expected, staining was high at kinetochores in early mitosis and decreased as
 160 chromosomes aligned at the spindle equator (Figure 2A) (Nijenhuis et al., 2014; Vleugel et al., 2015).
 161 To test the phospho-specificity of the rMAb-pMELT^{rb} antibody, we treated cells with 10 μM reversine,
 162 an Mps1 kinase inhibitor (Santaguida et al., 2010), and found the antibody reactivity was significantly

Figure 2. Characterization of recombinant antibodies to KNL1 pMELT, CENP-C, BubR1, 3F3/2 antigen, and Mad2-C. (A) HeLa cells immunostained with rMab-pMELT^{rb} antibodies. (B) HeLa cell treated with 10 μ M reversine and immunostained with rMab-pMELT^{rb} antibodies. (C) Immunoblots of lysates generated from untreated or reversine-treated HeLa cells expressing a 100 kDa fragment of KNL1 containing multiple pMELT domains, and probed with rMab-pMELT^{rb} antibodies (left) and tubulin antibodies as a loading control (right). (D) HeLa cells immunostained with rMab-CENP-C^{ms} antibodies. (E) HeLa cells immunostained with rMab-BubR1^{ms} antibodies. (F) Left: HeLa cell treated with BubR1 siRNA and immunostained with rMab-BubR1^{ms} antibodies. Right: Immunoblots of lysates generated from control or BubR1 siRNA-treated HeLa cells and probed with rMab-BubR1^{ms} antibodies (left) and tubulin antibodies as a loading control (right). (G) HeLa cells immunostained with rMab-3F3/2^{ms} antibodies. (H) HeLa cells immunostained with rMab-Mad2-C^{ms} antibodies. (I) HeLa cells (+/- Mad2 siRNA) pre-treated with 500 nM nocodazole for 12 hr and immunostained with rMab-Mad2-C^{ms} antibodies. In all immunofluorescence images, cells were stained with DAPI to detect chromosomes. Scale bars are 10 μ m.

174 reduced upon Mps1 inhibition, as detected by immunofluorescence and immunoblotting (Figure 2B, C).
 175 We found that the rMAb-CENP-C^{ms} antibody also recognized kinetochores during mitosis; however, the
 176 antibody exhibited low levels of cross-reactivity with rabbit secondary antibodies (not shown). We
 177 therefore cloned a new CENP-C antibody by combining the rMAb-Hec1^{ms} antibody constant regions with
 178 the sequenced CENP-C antibody variable regions. In this case, immunofluorescence experiments
 179 revealed that the new rMAb-CENP-C^{ms} antibody recognized kinetochores at all phases in mitosis as
 180 expected (Hara and Fukagawa, 2020) (Figure 2D), and there was no cross-reactivity with rabbit
 181 secondary antibodies (not shown).

182

183 We next wanted to obtain primary sequences and generate recombinant antibodies from existing mouse
 184 hybridoma cell lines producing monoclonal antibodies to key mitotic epitopes. To this end, we used cell
 185 lines expressing antibodies to the following targets: (1) the kinetochore-associated and spindle assembly
 186 checkpoint protein BubR1 (Chan et al., 1999; Zhang et al., 2015; Lischetti and Nilsson, 2015); (2) the
 187 3F3/2 phospho-epitope which recognize kinetochores that are lacking tension from microtubule pulling
 188 forces (Gorbsky and Ricketts, 1993; Nicklas et al., 1995; Waters et al., 1998); and (3) the active form of
 189 the kinetochore-associated and spindle assembly checkpoint protein Mad2, which recognize the “closed”
 190 conformation of Mad2 molecules found in Mitotic Checkpoint Complexes or bound to Mad1 (Sedgwick et
 191 al., 2016; De Antoni et al., 2005; Mapelli et al., 2007). For each hybridoma cell line, the mRNA
 192 transcriptome was obtained and used to generate a cDNA library from which the antibody sequences
 193 were identified through whole transcriptome shotgun sequencing (Absolute Antibody; Boston, MA).
 194 Based on the obtained sequences, the antibodies were classified as follows: BubR1: IgG2b / kappa light
 195 chain; 3F3/2: IgG2a / kappa light chain; and Mad2-Closed (Mad2-C): IgG1 / kappa light chain
 196 (Supplemental Figure 2A-C). Through this approach, we were able to additionally obtain the sequences
 197 of the native N-terminal signal peptides (also referred to as leader peptides) for both heavy and light
 198 chains for all three antibodies (Supplemental Figure 2A-C). Heavy and light chain sequences were cloned
 199 into expression vectors as described for rMAb-Hec1^{ms} above, with the exception that the native signal
 200 peptides were used. Expression plasmids were transfected into human Expi293F cells, and the
 201 antibodies were purified on Protein A Sepharose columns. Immunofluorescence experiments revealed
 202 that all three antibodies recognized kinetochores during mitosis (Figure 2E, G, and H). As expected,
 203 staining for all antibodies was high at kinetochores in early mitosis and decreased as chromosomes
 204 aligned at the spindle equator (Skoufias et al., 2001; Hoffman et al., 2001; Campbell and Gorbsky, 1995)
 205 (Figure 2E, G, and H). To test the specificity of rMAb-BubR1^{ms}, HeLa cells were treated with an siRNA
 206 targeted to BubR1 and processed for immunofluorescence or immunoblotting. As shown in Figure 2F,
 207 cells depleted of BubR1 exhibited significantly reduced reactivity with rMAb-BubR1^{ms} antibodies at
 208 kinetochores (left), or in cell lysates (right). Similarly, HeLa cells treated with Mad2 siRNA exhibited no
 209 reactivity at kinetochores with rMAb-Mad2-C^{ms} antibodies (Figure 2I). In this case, cells were treated with

nocodazole to enrich for kinetochore-association of Mad2, which is rapidly evicted from kinetochores in the presence of spindle microtubules in untreated cells.

Modification of antibody species specificity

For indirect immunofluorescence experiments, a commonly used cell biology approach, a fluorescently tagged secondary antibody recognizes a primary antibody and provides signal detection and amplification. In many cases, commercial primary antibodies directed to target antigens are only available in a single animal specificity, meaning the antibodies were generated in a particular species and contain heavy and light chain constant regions unique to that species. In practice, this limits the number and combinations of targets that can be probed for and detected in such experiments. A solution to this problem is to directly conjugate primary antibodies with fluorescent dyes, obviating the need for secondary antibodies, and allowing for the simultaneous use of two primary antibodies generated in the same host species. However, this approach requires a substantial amount of primary antibody, which may be cost prohibitive if the antibodies are purchased from a commercial source. Furthermore, this approach eliminates secondary antibody-mediated amplification of primary antibody signals. We therefore set out to alter the species specificity of our recombinant monoclonal antibodies to expand our

Figure 3. Generation of recombinant antibodies with modified species specificity. (A) Schematic of “species swap” approach. (B-F) HeLa cells stained with antibodies to rMAb-Hec1^{rb}, panel B; rMAb-Hec1^{hu}, panel C; rMAb-CENP-C^{hu}, panel D; rMAb-BubR1^{hu}, panel E; and rMAb-Mad2-C^{hu}, panel F. For all panels B-F, cells were probed with the species-appropriate secondary antibody and also with secondary antibodies specific for the original species. Cells were also stained with DAPI to detect chromosomes. Scale bars are 10 μm.

toolbox of available reagents. To this end, we generated a new Hec1 primary antibody, termed rMAb-Hec1^{rb}, such that it is recognized by a rabbit secondary antibody. We generated the new sequence by removing the constant regions from both the heavy and light chains of the rMAb-Hec1^{ms} antibody and replacing them with the constant regions from a rabbit IgG antibody (Figure 3A). The new heavy and light chain expression plasmids were transfected into human Expi293F cells and purified on a Protein A Sepharose column as described above. We tested the antibody in immunofluorescence assays and determined that the rMAb-Hec1^{rb} localized to kinetochores in mitotic cells, and importantly, was recognized by rabbit secondary antibodies, but not by mouse secondary antibodies. (Figure 3B). We diversified the Hec1 antibody even further by generating a version that is recognized by a human secondary antibody. In this case, we acquired human IgG amino acid sequences from the UniProt Knowledgebase (UniProt # P01857 and P01834) and generated geneblocks for the heavy and light chain constant regions. We then combined the variable regions of both the heavy and light chains from the rMAb-Hec1^{ms} IgG2a sequence with the human IgG antibody constant sequences to generate plasmids encoding for rMAb-Hec1^{hu} (Figure 3C). We purified and tested this antibody in immunofluorescence assays and determined that the rMAb-Hec1^{hu} localized to kinetochores in mitotic cells, and was recognized by human, but not mouse secondary antibodies (Figure 3C). We went on to use this approach to generate a number of additional species variants, including rMAb-Cenp-C^{hu}, rMAb-Mad2-C^{hu}, and rMAb-BubR1^{hu} antibodies (Figure 3D-F).

Generation of recombinant antibody fragments

For some cell biological and biomedical applications, antibody fragments, generated by either proteolysis or genetic engineering, offer advantages over the use of intact, bivalent antibodies. For example, their smaller size allows more efficient penetration of tissue samples and may provide better access to “buried” epitopes; they are able to bind targets without inducing cross-linking; they reduce steric effects (compared with intact antibodies) when monitoring an antigen in living cells, and they have reduced immunogenicity, which may be desirable for therapeutic applications (Hayashi-Takanaka et al., 2011; Cheloha et al., 2020; Zhao et al., 2019; Xenaki et al., 2017; Berland et al., 2021; Ries et al., 2012; Morisaki et al., 2016; Yan et al., 2016; Wang et al., 2016; Stasevich et al., 2014). Another advantage is that single chain antibody fragments can be fused to a fluorescent protein and genetically encoded for expression in living cells for the purpose of real-time antigen tracking. This ability becomes particularly important when an antibody recognizes a post-translational protein modification or specific protein conformation, which cannot be tracked simply by expressing a fluorescently-tagged version of a protein of interest (Sato et al., 2013; Kimura et al., 2015). Finally, the use of antibody fragments is becoming increasingly important in super-resolution microscopy approaches, where the achieved spatial resolution depends not only on the imaging method, but on the size of the probes used (Ries et al., 2012; Traenkle and Rothbauer, 2017; Mikhaylova et al., 2015; Zhao et al., 2019).

To further diversify our recombinant antibodies and to capitalize on the advantages listed above, we set out to generate three types of antibody fragments: (1) scFvC (single chain variable plus a truncated constant region), (2) scFv (single chain variable fragment), and (3) Fab (antigen binding fragment). We first used the antibody sequences to Hec1 and Mad2-C to generate recombinantly expressed and purified

Figure 4. Generation of antibody fragments. (A) Schematic illustrating the generation of three types of fragments: scFvC (left), scFv (center), and Fab (right). (B and C) Hec1 and Mad2-C scFvC fragments were purified on Protein A Sepharose columns and bead-loaded into HeLa cells. Cells were fixed and stained with anti-rabbit secondary antibodies and DAPI to detect chromosomes. (D and E) HeLa cells were transfected with the scFvC-Hec1 plasmid or the scFvC-Mad2-C plasmid. Cells were fixed and stained with anti-rabbit secondary antibodies and DAPI to detect chromosomes. (F) HeLa cells were transfected with the scFv-pMELT-GFP plasmid and time-lapse imaged using confocal microscopy, and a representative cell is shown. At time=0 min, many kinetochores are positive for scFv-pMELT-GFP, and by 18 min, when the cell has reached metaphase, no kinetochore-associated pMELT signals are detected. Arrows in the 8 min timepoint image point to the kinetochores which retain detectable MELT phosphorylation in late prometaphase. (G) Hec1 Fab were generated through proteolysis and directly labeled with an Alexa 647 fluorophore. HeLa cells were bead-loaded with Hec1 Fab⁶⁴⁷, stained with DAPI to detect chromosomes, and imaged. (H) Hec1 Fab were generated through proteolysis and directly labeled with both Alexa 405 and Alexa 647 fluorophores. HeLa cells were fixed, permeabilized, incubated with Hec1 Fab^{405/647}, and subjected to both wide-field (left) and STORM imaging (right). Scale bars are 10 μm.

scFvC fragments, which contain the variable regions of the heavy and light chains connected by a flexible linker, attached to the rabbit IgG-specific heavy chain constant regions (CH2 + CH3) in a single polypeptide chain and totaling ~50 kDa in mass (Figure 4A, left panel). Single plasmids encoding for the Hec1 and Mad2-C scFvC fragments (scFvC-Hec1^{rb} and scFvC-Mad2-C^{rb}), which include signal peptides (Lima and Cosson, 2019), were transfected into human Expi293F cells and purified on Protein A Sepharose columns. We expressed and were able to purify the scFvC fragments on our affinity columns, since they contain a portion of the heavy chain constant region that is recognized by Protein A. The purified Hec1 and Mad2-C scFvC fragments were introduced into cells by bead-loading (McNeil and Warder, 1987; Cialek et al., 2021), and the cells were subsequently fixed and incubated with anti-rabbit secondary antibodies. As shown in Figure 4B and 4C, the fragments recognized kinetochores in mitotic cells, and the results were similar to those obtained with the intact, bivalent antibodies. Next, we generated genetically encoded scFvC fragments using both the Hec1 and Mad2-C antibody sequences. For these constructs, we removed the signal peptide sequence and transiently expressed a single plasmid encoding for the heavy and light chain variable regions connected by the flexible linker, and the rabbit IgG-specific heavy chain constant regions (CH2 + CH3). HeLa cells were transfected with the single plasmids, prepared for immunofluorescence and stained with an anti-rabbit secondary antibody, and as shown in Figure 4D and E, the genetically encoded Hec1 and Mad2-C scFvC fragments both recognized kinetochores in mitotic cells.

We next used the rMAb-pMELT^{rb} sequence to generate an scFv, comprised of the variable regions of the heavy and light chains connected by a flexible linker and totaling ~25 kDa in mass (Figure 4A, middle panel). In this case, the scFv does not contain the Fc (fragment crystallizable) region (comprised of the heavy chain constant regions CR2 and CR3), which is recognized by Protein A. We therefore did not express and purify the antibody for this experiment. However, we added a GFP tag to the C-terminus of the polypeptide chain, omitted the signal peptide, and expressed the fluorescently tagged scFv in HeLa cells. We collected time-lapse images, and as shown in Figure 4F, the genetically encoded pMELT scFv-GFP recognizes kinetochores in early mitosis when phosphorylation of the KNL1 MELT repeats is high, but much less so in late mitosis, when phosphorylation of the MELT repeats is low. These results demonstrate that small immunological probes can be generated from primary monoclonal antibody sequences and successfully expressed in mitotic cells to detect mitotic antigens, and importantly, post-translational protein modifications of mitotic targets in living cells.

Finally, we used the recombinant, purified rMAb-Hec1^{ms} antibody to generate antigen binding fragments (Fab), containing a single constant region of each chain and the variable regions from both the heavy and light chains (Figure 4A, right panel). The purified rMAb-Hec1^{ms} antibody was enzymatically digested with papain protease, the digestion reaction was centrifuged through a Protein A spin column, and the

antigen binding fragments, which do not bind the Protein A resin, were collected in the flow through. We directly labeled the rMAb-Hec1^{ms} Fab with an Alexa 647 fluorescent dye and bead-loaded the fragment into HeLa cells, and as shown in Figure 4G, the labeled Fab recognizes kinetochores in mitotic cells. We next tested if the labeled rMAb-Hec1^{ms} Fab is appropriate for use in super-resolution STORM imaging. For this experiment, we double-labeled the rMAb-Hec1^{ms} Fab with Alexa 647 and Alexa 405 dyes at a ratio of 1:1. HeLa cells were then fixed and stained with the dually-labeled rMAb-Hec1^{ms} Fab^{405/647} fragment (Figure 4H). Samples were excited with 405 and 640 nm lasers, and images were collected on a Nikon N-STORM imaging system. The Fab robustly recognized kinetochores and the resulting STORM images demonstrate that our recombinantly expressed antibody fragments are compatible with super-resolution imaging approaches (Figure 4H). Importantly, these super-resolution techniques benefit from the use of small, directly labeled probes by preventing loss of spatial resolution that occurs with the use of larger, intact bivalent primary and secondary antibodies (Ries et al., 2012; Traenkle and Rothbauer, 2017; Mikhaylova et al., 2015).

Reverse engineering antibody fragments into full-length, bivalent antibodies

Advances in genetic engineering techniques such as antibody phage display and hyper-variable domain grafting have provided new routes to generating antibody fragments including scFv (single-chain variable fragments) and single-domain antibodies (sdAbs, also known as nanobodies) (Panikar et al., 2021; Laustsen et al., 2021; Valldorf et al., 2021; Shim, 2017; Shukra et al., 2014; Zhao et al., 2019). As a result, there are growing numbers of published antibody fragment sequences. As discussed above, there are numerous advantages to using such antibody fragments; however, there are also cases in which a full-length, bivalent antibody is preferred. For example, in standard indirect immunofluorescence experiments, multiple secondary antibodies bind to the constant regions of a full-length, bivalent primary antibody, which results in signal amplification and increased sensitivity. Similar arguments can be made for the use of full-length, bivalent antibodies in immunoblotting and immunoprecipitation experiments. Given the rise in availability of antibody fragment sequences, we wanted to use our approach to convert scFv sequences into full-length, bivalent antibodies. For this reason, we modified an HA (hemagglutinin)-tag scFv (also known as the HA-tag “frankenbody”) sequence (Zhao et al., 2019) into a full-length, bivalent antibody. Specifically, we cloned the heavy chain variable regions of the HA-tag scFv onto the heavy chain constant regions from a rabbit IgG antibody (Figure 5A). Similarly, we cloned the light chain variable region of the HA-tag scFv onto the rabbit IgG light chain constant region (Figure 5A). In addition, signal peptides were cloned into both heavy and light chain plasmids. Both plasmids were transfected into human Expi293F cells as described above and full-length, bivalent antibody secreted into the cell media was purified using a Protein A Sepharose column. To test the antibody in cells and to confirm specificity, we expressed the following versions of Hec1 in HeLa cells: HA-tagged (Figure 5B); HA- and GFP-tagged (Figure 5C), and GFP-tagged (Figure 5D). Cells were prepared for immunofluorescence and

stained with the rMAb-HA^{rb} antibody and a fluorescently-labeled rabbit secondary. As shown in Figure 5 B-D, the rMAb-HA^{rb} antibody recognized Hec1 at kinetochores in cells expressing Hec1-HA and Hec1-HA-GFP, but not in cells expressing Hec1-GFP.

Figure 5. Generation of full-length, bivalent antibodies from fragment sequences. (A) Schematic illustrating the generation of a reverse engineered, full-length HA-tag antibody from an scFv. (B-D) HeLa cells were transfected with Hec1-HA (B), Hec1-HA-GFP (C), or Hec1-GFP (C), and immunostained with rMAb-HA^{rb} antibodies. In panels C and D, GFP was imaged. (E) Immunoblot of HeLa lysate probed with reverse engineered, full-length tubulin antibody, rMAb-Tub^{rb}. (F) HeLa cells immunostained with rMAb-Hec1^{ms} and rMAb-Tub^{rb}. In all immunofluorescence images, cells were stained with DAPI to detect chromosomes. Scale bars are 10 μm.

Finally, we carried out a similar reverse engineering approach to generate a full-length, bivalent IgG tubulin antibody using the sequence of an alpha tubulin scFvC antibody fragment (Lima and Cosson, 2019). Similar to the approach used for the HA-tag scFv described above, we cloned the heavy and light chain variable regions from the tubulin scFvC onto the heavy and light chain constant regions from a rabbit IgG antibody. Additionally, signal peptides were cloned into both the heavy and light chain plasmids. The full-length antibody was expressed and purified from human Expi293F cells. We confirmed that the new, bivalent rMAb-Tub^{rb} antibody recognized a ~55 kDa protein in HeLa cell lysate, which corresponds to the molecular weight of α and β tubulin monomers (Figure 5E). In addition, the antibody

recognizes microtubules in cells, and as expected, the primary antibody is recognized by rabbit but not mouse secondary antibodies (Figure 5F).

Discussion

Here we describe a set of tools and protocols that can be used to generate high-yield, low-cost antibodies and antibody fragments from primary amino acid sequences. While most of the antibodies and antibody derivatives described here are directed to antigens involved in mitotic cell division and kinetochore function, the plasmids and protocols are applicable to any monoclonal antibody sequence. Using relatively small volumes (~30 mL) of human Expi293F cells grown in suspension, we demonstrate production of up to 1.4 mg of protein per preparation (rMAb-Hec1^{ms}). We note, however, that the yields for individual antibody preparations varied, and our lowest yields were in the range of 0.1 mg of protein from a 30 mL suspension culture. We have not yet defined the source of yield variation, but we are working to normalize these values by optimizing the cloning, expression, and purification conditions for the lower-yield reagents.

In some cases, the major cost incurred for the approach described here is the protein sequencing itself. Currently, the approximate costs of sequencing a purified monoclonal antibody ranges from ~\$5000-\$12,000. However, obtaining sequence information from a monoclonal hybridoma cell line is much less expensive (~\$500-\$1500), and is offered through many companies and non-profit facilities as a stand-alone service, or as an “add-on” cost for custom monoclonal antibody production services. We propose that if individual research labs submit existing hybridoma cell lines for antibody sequencing and share this information publicly, this would greatly benefit the entire research community by providing a means for individual labs to produce low-cost antibodies and generate customized reagents for use in their own research. Importantly, there are growing numbers of available monoclonal antibody sequences available through databases, obtained through protein sequencing or from atomic structures. These include the ABCD (Antibodies Chemically Defined) database at the Geneva Antibody Facility (Lima et al, 2000); <https://web.expasy.org/abcb>; the ABG (Antibody Group) directory (<http://pt7mdv.ceinge.unam.mx/vir/structure/structures.html>); the Kabat database (Johnson and Wu, 2000); <http://bigd.big.ac.cn/databasecommons/database/id/5425>; SAbDab (<http://opig.stats.ox.ac.uk/webapps/newsabdab/sabdab/>); abYbank (www.abYbank.org); and the IMGT (International Immunogenetics Information System) monoclonal antibodies database (<http://www.imgt.org/mAb-DB/doc>). The approaches described are intended to facilitate the expression and purification of recombinant antibodies and antibody fragments in individual labs using accessible, non-specialized equipment and reagents; however, we note that many non-profit, institutional core facilities also offer these services at relatively low costs, which provide a convenient and accessible alternative to expressing and purifying antibodies in-house. We also note that costs can be reduced by

ordering geneblocks for only the variable regions of a newly sequenced antibody and combining these sequences with previously-validated constant regions of choice (e.g. those described here for rabbit, mouse, and human IgG antibodies). Finally, it is important to consider downstream usage of data generated from antibodies purchased from commercial sources, as different companies may have unique sets of terms and conditions.

In addition to providing low-cost methods for in-house monoclonal antibody production, the approaches described here allow for diversification of sequenced monoclonal antibodies. One practical example of such diversification is altering the species specificity by “swapping” constant regions. Researchers are typically limited to the combination of antigens they wish to detect in indirect immunofluorescence or immunohistochemical experiments by species specificity of the antibodies. Once a monoclonal antibody sequence is in hand, it is simple and straightforward to produce a new antibody containing the constant regions specific to a different species.

Another advantage of the methodology described here is the ability to generate antibody fragments from a primary sequence. Advantages of such fragments include more efficient penetration of tissue and cell samples; access to less-accessible epitopes; the ability to bind targets without inducing cross-linking; and reduced steric effects compared with full-length bivalent antibodies. Additionally, antibody fragments can be used for tracking protein modifications and specific protein conformations in living cells. To study protein dynamics in cells, fluorescent proteins such as GFP are typically fused to a protein of interest, expressed, and monitored in living cells. However, such fluorescent protein tags cannot track or identify protein modifications (e.g. phosphorylation), nor can they detect specific protein conformations, both of which may play critical roles in a protein’s cellular function. In recent years, fluorescently labeled antibody fragments such as Fab, scFv, scFvC, and nanobodies have been used for the purpose of tracking post-translational modifications and specific protein conformations (Kimura et al., 2015; Lyon and Stasevich, 2017; Sachs et al., 1972). Antibody fragments may be preferable to intact bivalent antibodies for tracking modifications and specific conformations in living cells since their smaller size minimizes steric effects of the probes themselves, which could lead to interference with protein function. Such antibody fragments can be purified, directly labeled with a fluorescent dye, and introduced into cells by microinjection, electroporation, or bead-loading, and visualized in living cells. Alternatively, single chain antibody fragments (e.g. scFv, scFvC) fused to a fluorescent protein such as GFP can be genetically encoded for expression and monitoring in living cells. Here, we describe tools to generate both types of labeled antibody fragments from a primary monoclonal antibody sequence, which will allow for investigations of processes in cells that have previously not been feasible. For example, during mitotic cell division the regulation of kinetochore-microtubule attachments is driven in large part by reversible and dynamic phosphorylation events, yet these events have not yet been monitored in real-time during mitotic

progression in living cells. The methodologies described here will enable the generation of reagents for real-time tracking of phosphorylation events on key kinetochore proteins during chromosome congression and spindle assembly checkpoint inactivation, which will provide critical insight into the mechanisms that regulate kinetochore function during mitosis.

It is important to note that while we were successful in generating a recombinant scFv directed to the KNL1 pMELT domain using the native sequence, we were not successful in generating scFv fragments to Hec1, BubR1, or Mad2-C. Why this is the case is not clear; however, there are other methods to obtain an scFv from the primary sequence that we have not yet explored. For example, it has been demonstrated that the hyper-variable domains, obtained from the primary sequence of a monoclonal antibody, can be grafted onto optimized scFv scaffolds which are known to function in cells (Zhao et al., 2019). This approach has led to the generation of genetically encoded single chain antibody fragments referred to as “frankenbodies,” which are capable of recognizing and tracking specific epitopes in cells (Zhao et al., 2019).

In this study, we also demonstrated that our recombinant antibody fragments can be directly labeled with fluorescent dyes for use in super-resolution light microscopy approaches. The use of small, directly-labeled probes is desirable for super-resolution imaging approaches, such as PALM/STORM, where the goal is precise molecular mapping of antigens on the nanometer scale. The size of the probe is directly relevant since larger probes, such as intact, bivalent primary and secondary antibodies, will limit the achievable resolution (Ries et al., 2012; Traenkle and Rothbauer, 2017; Mikhaylova et al., 2015). For example, an intact bivalent antibody is approximately 10-15 nm long, and when using an imaging technique with a spatial resolution in the range of 20-30 nm (e.g. STED; PALM/STORM), such a large labeling probe reduces the resolution by approximately 2-fold.

In summary, we describe methodologies to generate and purify recombinant versions of a suite of antibodies directed to kinetochore and mitotic proteins. We also present a set of molecular biological methods to expand the versatility of these antibodies by altering species specificity and generating antibody fragments that can be either genetically encoded for tracking antigens in cells or recombinantly expressed and purified. As tools such as these become more widely shared, access to low-cost, sequence-defined antibodies will increase, benefiting all fields that utilize antibody and antibody-based tools.

Materials and Methods

Antibody sequencing. Purified samples (100 µg each) of Hec1 (Genetex), KNL1 pMELT (Fisher Scientific), and CENP-C (Abcam) monoclonal antibodies were sequenced by Rapid Novor (Kitchener,

Ontario, Canada) using tandem mass spectrometry. Data obtained from these antibodies presented in the current study are for non-commercial purposes only, in accordance with the terms and conditions of the companies from which the CENP-C, Hec1 9G3, and KNL1 pMELT antibodies were purchased. Cell samples (10^6 cells each) of Mad2, BubR1, and 3F3/2 hybridoma cell lines were submitted to Absolute Antibody (Boston, MA, USA) for sequencing. For each cell sample, the mRNA transcriptome was obtained through whole transcriptome shotgun sequencing, the resulting reads were assembled into contigs, and antibody transcripts were identified based on homology.

Plasmid generation. For full-length Hec1, KNL1 pMELT, CENP-C, Mad2-C, BubR1, and 3F3/2 antibodies, the protein sequence for each was used to design DNA geneblocks optimized for expression in human cells using the IDT (Integrated DNA Technologies) codon optimization tool. For full-length Hec1, KNL1 pMELT, and CENP-C antibodies, an N-terminal signal peptide sequence was added to the geneblock (Burton, 1994; Yu et al., 2006). For Mad2-C, BubR1, and 3F3/2 antibodies, the native signal peptides were included in the geneblock design. The resulting DNA fragments/geneblocks were cloned using the Gibson assembly method into the pEGFP-N1 vector (Clontech) with the GFP removed by Sac1/Not1 digestion, henceforth referred to as the rMAbParent plasmid. For each full-length antibody, a heavy chain (HC) and light chain (LC) plasmid was generated for co-expression in HEK293 suspension culture cells (Expi293F cells) (Fisher Scientific). For species specificity swapping experiments, geneblocks corresponding only to the variable regions of the HCs and LCs were designed and ordered. PCR fragments were generated corresponding to the target species constant regions for both HCs and LCs. DNA fragments from the geneblocks for the variable regions were combined with the PCR fragments for the constant regions and cloned into the rMAbParent plasmid using the Gibson assembly method. For construction of each of the Mad2 and Hec1 scFvC plasmids, the following PCR fragments were generated: (1) HC variable region (for expression in Expi293F cells, a signal peptide was also included; for genetic encoding, the signal peptide was not included) (Sasada et al., 1988), (2) flexible linker, (3) LC variable region, and (4) rabbit IgG HC constant regions (CH2 and CH3). PCR fragments were cloned by the Gibson assembly method into the rMAbParent plasmid resulting in a final single scFvC plasmid. For construction of the KNL1 pMELT scFv, the following PCR fragments were generated: (1) HC variable region (for expression in Expi293F cells a signal peptide was also included; for genetic encoding, the signal peptide was not included) (Sasada et al., 1988), (2) flexible linker, and (3) LC variable region. PCR fragments were cloned by the Gibson assembly method into the rMAbParent plasmid (containing the sequence for GFP), resulting in a final single scFv plasmid. To generate the full-length bivalent HA-tag antibody, PCR fragments were generated corresponding to the HC and LC variable regions of the HA-tag scFv, which were generated by grafting the HA-tag HC and LC hyper-variable regions (a.k.a. complementarity determining regions or CDRs) into the 15F11 scaffold (Zhao et al., 2019). The PCR fragments containing the HC and LC variable regions also included N-terminal signal peptide sequences

(Burton, 1994; Yu et al., 2006). PCR fragments corresponding to the rabbit IgG-specific LC and HC constant regions were generated and, together with the HC and LC variable regions, were cloned into the rMAbParent plasmid using the Gibson assembly method. To generate the full-length bivalent tubulin antibody, geneblocks corresponding to the HC and LC variable regions of a tubulin scFv-C (AA345) (Lima and Cosson, 2019) were designed. N-terminal signal peptide sequences were added to both the HC and LC variable regions (Burton, 1994; Yu et al., 2006). PCR fragments corresponding to the rabbit IgG-specific LC and HC constant regions were generated and, together with the HC and LC variable regions, were cloned into the rMAbParent plasmid using the Gibson assembly method. Descriptions of all plasmids generated in this study are listed in Table 1.

Cell culture. Human HEK293 suspension culture cells (Expi293F) were cultured in Expi293F expression media (Fisher Scientific) and maintained at 37°C in 8% CO₂ in 125 mL spinner flasks on an orbital shaker rotating at 125 rpm. HeLa cells were cultured in DMEM supplemented with 10% FBS and 1% antibiotic/antimycotic solution and maintained at 37°C in 5% CO₂.

Expi293F cell transfection. HEK293 Expi293F cells were grown to 30 mL volumes in 125 mL spinner flasks. For generation of full-length antibodies, cells were treated as follows: 24 hr prior to transfection, cells were seeded at 0.9×10^6 cells/mL (27×10^6 cells total) with a viability no less than 90% (typically 94-99% viability). On the day of transfection, cells were counted and 30×10^6 cells (at >90% viability) were pelleted by centrifugation at 2000 x g for 15 min at 4°C. The resulting supernatant was aspirated off and the pellet was gently resuspended in 15 mL fresh Expi293F media. Spinner flasks were returned to the incubator while transfection reactions were prepared. For transfection, 100 µl of polyethylenimine (PEI) (Polysciences Inc.) at 1 mg/mL was added to 1 mL Optimem media (Gibco). In a second tube, 50 µg LC plasmid and 35 µg HC plasmid were mixed with 1 mL Optimem media. The two separate tubes were incubated for 5 min at room temperature with occasional flicking. After 5 min, the contents of the two tubes were combined and mixed by flicking every 3 min for a total of 20 min at room temperature. After 20 min, the mixture was added to the 15 mL Expi293F culture and incubated overnight, rotating at 125 rpm. Approximately 15 hr later, 15 mL fresh Expi293F media was added to the original 15 mL culture, as well as 300 µl of 220 mM valproic acid (Sigma). Cells were subsequently cultured for an additional 4 days, after which cells were pelleted by centrifugation at 2000 x g for 15 min at 4°C. The supernatant was harvested and filtered through a 0.2 µm filter (GenClone). The volume of the filtered supernatant was measured, and 1.0 M Tris-HCL, pH 7.4 was added to bring the pH to between 7.4-7.7. The filtered supernatant was stored at 4°C until purification, which was optimally within 1 day of harvesting.

Antibody purification. The Protein A slurry was prepared by washing 1.5 g Protein A Sepharose (Sigma) 4 times in 1X Tris Buffered Saline (1X TBS: 50 mM Tris-Cl, 150 mM NaCl, pH 7.5), and raised in a final

volume of 40 mL 1X TBS, pH 7.5. Three mL of the washed slurry was added to the 30 mL of filtered Expi293F cell supernatant and gently inverted for 12 hr at 4°C. Following the 12 hr incubation, the antibody-containing cell supernatant plus Protein A Sepharose mixture was added to a 9 cm high, 2 mL bed volume (0.8 x 4 cm) empty polypropylene column (BioRad) fitted with a two-way stopcock up to the fill line. A funnel reservoir was attached to the top of the column and the remaining cell supernatant/Protein A slurry was poured into the funnel. The column and the reservoir were transferred to a 4°C cooler, covered loosely, and left to settle for 1 hr. After 1 hr, the stopcock was opened and the flow-through was collected at a flow rate of 1 mL/min. Once the cell supernatant approached the top of compacted Protein A slurry (with ~ 1 mL remaining), the column flow was stopped and the flow-through was added back to the column/funnel for a second round of binding. The column was left to settle for 15 min, the stopcock was opened and the flow-through was again collected at 1 mL/min. After the flow-through approached the compacted slurry, 3 mL of 1X TBS (pH 8.0) was added to the column and incubated for 15 min. After 15 min, the flow-through was collected at 1 mL/min. To elute the purified antibody, 9 mL of low pH elution buffer (0.15 M NaCl, 0.1 M glycine, pH 2.95) was added to the column, which was then connected to a peristaltic pump (BioRad). The flow rate was adjusted to 5 mL/min, and the eluate was collected in a 15 mL conical tube containing 0.9 mL 1 M Tris-HCl, pH 8.0. The eluate was then dialyzed and concentrated by transferring it to pre-soaked dialysis membrane (SpectrumLabs), which was then placed into 1 L of 1X phosphate buffered saline (PBS) buffer (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄, 1.47 mM KH₂PO₄, pH 7.4) and gently stirred at 4°C for 4 hr. After 4 hr, the 1X PBS was replaced and incubated overnight at 4°C. Purified, dialyzed antibodies were retrieved from the dialysis tubing and subsequently concentrated in a 10,000 kDa cutoff concentrator (Millipore) to final volume of between 100-200 µL. The protein concentration was calculated, and glycerol was added to a final concentration of 15%. Purified, concentrated antibodies were stored in 2-10 µL aliquots at -20°C.

Cell treatments. For live-cell imaging experiments, HeLa cells were seeded and imaged in 35-mm glass-bottomed dishes (constructed in-house). For fixed-cell analysis, cells were grown on sterile, acid-washed coverslips in six-well plates. siRNAs were transfected into HeLa cells using 6 µL Oligofectamine (Fisher Scientific) and 160 nM of the appropriate siRNA: siHec1 (5'-CCCUGGGUCGUGUCAGGAA-3'); siBubR1 (5'-AAGGAGACAACUAAACUGCAA-3'); siMad2 (5'-CUGAAAGUAACUCAUAUAUCUA -3') (Qiagen). For transfection of the scFv or scFvC plasmids, 2.5 µL of Lipofectamine 3000 (Fisher Scientific) and 0.5-1 µg of DNA were used. All siRNA and DNA transfections were incubated for 24-30 hr before the cells were either imaged, fixed for immunofluorescence or harvested for immunoblot analysis. For some experiments, cells were incubated with 500 nM nocodazole (Tocris) for 15 hr prior to fixation for immunofluorescence or harvesting for immunoblot analysis. To inhibit Mps1 kinase, cells were treated with 10 µM reversine (Adooq Biosciences) for 1 hr prior to fixation for immunofluorescence or harvesting for immunoblot analysis. For bead loading experiments, directly labeled Fab-Hec1⁶⁴⁷ (see below) was

bead loaded into HeLa cells 1 hr prior to fixing (Cialek et al., 2021). Briefly, 10 μ l of Fab-Hec1⁶⁴⁷ were placed directly atop growing HeLa cells in a 35-mm glass-bottomed dish. Glass beads were then sprinkled atop the cells and the dish was agitated by sharply striking against the countertop. Fresh cell media was added to the dish, which was returned to the incubator for 1 hr prior to fixation.

Immunofluorescence. Cells were rinsed in 37°C PHEM buffer (60 mM PIPES, 25 mM HEPES, 10 mM EGTA, and 4 mM MgSO₄, pH 7.0) and then lysed for 5 min in freshly prepared lysis buffer (PHEM buffer + 0.5% Triton X-100). Cells were subsequently fixed for 20 min at room temperature in freshly prepared 4% paraformaldehyde in PHEM buffer (37°C). After fixation, cells were washed 5 X 3 min in PHEM-T (PHEM buffer + 0.1% Triton X-100) and then blocked in 10% boiled donkey serum (BDS) in PHEM for 1 hr at room temperature. Primary antibodies diluted in 5% BDS were added to coverslips and allowed to incubate for 1 hr at room temperature. The following primary antibody and antibody fragment concentrations were used for immunofluorescence: rMAb-Hec1^{ms} at 1.5 μ g/mL, rMAb-Hec1^{rb} at 0.2 μ g/mL, rMAb-Hec1^{hu} at 1 μ g/mL, rMAb-pMELT^{rb} at 1.9 μ g/mL, rMAb-CENP-C^{ms} at 0.66 μ g/mL, rMAb-CENP-C^{hu} at 0.6 μ g/mL, rMAb-BubR1^{ms} at 2.1 μ g/mL, rMAb-BubR1^{hu} at 0.85 μ g/mL, rMAb-3F3/2^{ms} at 0.84 μ g/mL, rMAb-Mad2-C^{ms} at 1.6 μ g/mL, rMAb-Mad2-C^{hu} at 1.12 μ g/mL, scFvC-Hec1^{rb} at 0.5 μ g/mL, scFvC-Mad2-C^{rb} at 1.0 μ g/mL, rMAb-tubulin^{rb} at 1.2 μ g/mL, and rMAb-HA^{rb} at 1.2 μ g/mL. After primary antibody incubation, cells were rinsed 5 X 3 min in PHEM-T and then incubated for 45 min at room temperature with secondary antibodies conjugated to either Alexa 647 or Cy3 (Jackson ImmunoResearch Laboratories, Inc.) at 1.5 μ g/mL diluted in 5% BDS. Cells were rinsed 5 X 3 min in PHEM-T, incubated in a solution of 2 ng/ml DAPI diluted in PHEM, rinsed 5 X 3 min, quick-rinsed in PHEM, and then mounted onto glass slides in an antifade solution (90% glycerol + 0.5% *N*-propyl gallate). Coverslips were sealed with nail polish and stored at 4°C.

Imaging. All fixed cell images were acquired on an Inverted Olympus microscope incorporated into a GE Ultra imaging system (GE Healthcare) with SoftWoRx software (GE Healthcare) using a 60X 1.42 NA differential interference contrast Plan Apochromat oil immersion lens (Olympus) with a final magnification of 107.6 nm/pixel at the camera sensor (edge4.2, PCO Inc.). For live-cell imaging experiments, cells were imaged in 35-mm glass-bottomed dishes (constructed in-house) and imaged in Leibovitz's L-15 media (Invitrogen) supplemented with 10% FBS, 7 mM HEPES, 4.5 g/liter glucose, pH 7.0. Images were captured on a Nikon Ti-E microscope equipped with a Piezo Z-control (Physik Instrumente), stage top incubation system (Okolab), and spinning disk confocal scanner unit (CSUX1; Yokogawa), using a 60X, 1.49NA objective and an iXon DU888 EM-CCD camera (Andor). Five z-planes at 0.75 μ m steps, were acquired every 3 min for the duration of filming using the 488 nm laser.

Fluorophore conjugations. For direct labeling of the Fab fragment, full length rMab-Hec1^{ms} was first digested using a Pierce Fab Preparation Kit (Fisher Scientific) according to the manufacturer's instructions. Purified Hec1 Fab was then directly conjugated with Alexa 647 (Invitrogen) according to the following procedure: 0.065 mg of Fab-Hec1^{ms} was incubated with 6 µl 1M NaHCO₃ and 1 µl Alexa 647 (final reaction volume 50 µL). The tube containing the reaction mix was wrapped in foil, rotated for 30 min at room temperature, and then diluted with an additional 140 µl of 1X PBS. This solution was then added to the center of a Nap-5 gel filtration column and allowed to enter the column via gravity. 500 µl of 1X PBS was added to the column and the fastest-eluting fluorescent band containing the fluorescently-conjugated protein was collected in a fresh tube. Hec1 Fab⁶⁴⁷ concentration and labeling ratio was calculated.

STORM imaging. Cells were seeded and fixed as above in 35-mm glass-bottomed dishes. The Hec1 Fab used for STORM was directly labeled as described above, but in addition to Alexa 647, the Fab was also conjugated with Alexa 405 by adding 3 µl of the dye to the same reaction volume. The directly labeled Hec1 Fab^{405/647} was used at 1:500 following the above immunofluorescence procedure. For imaging, STORM imaging buffer was made immediately before use according to Nikon STORM protocol: 24 µl glucose oxidase, 280 µl 1M MEA 2480 µl (50 mM Tris-HCl + 10 mM NaCl + 10% glucose). 2 mL of the imaging buffer was added to the dish and the cells were imaged on a Nikon Ti-Eclipse microscope using a 1.49NA 100X Plan Apo TIRF lens equipped with an iXon3 DU897 EM-CCD (Andor). STORM images were acquired using 405 nm and 640 nm lasers on N-STORM software, version 3.30.

Immunoblotting. Samples were run on 12% SDS-polyacrylamide gels and subsequently transferred to 0.2 µm polyvinylidene difluoride membrane (PVDF) (Millipore). Membranes were washed with 1X TBS and incubated with a 5% solution of BSA for 1 hr at room temperature. Primary antibodies were added to membranes in quick-seal bags, and incubated, rocking for 1 hr at room temperature. The following primary antibody and antibody fragment concentrations were used for immunoblotting: rMab-Hec1^{ms} at 1.5 µg/mL, rMab-BubR1^{ms} at 2.1 µg/mL, rMab-pMELT^{rb} at 1.9 µg/mL and rMab-Tubulin^{rb} at 1.2 µg/mL. Membranes were washed with TBS-T (1 X TBS + 0.05% Tween-20) and incubated with HRP-tagged secondary antibodies for 1 hr at room temperature. Membranes were washed with TBS-T and scanned on a chemiluminescence imager IA600 (GE Healthcare). For the KNL1 pMELT immunoblots, a truncated version of KNL1 with a molecular mass of ~100 kDa and containing multiple MELT motifs was transiently expressed in HeLa cells 24 hr prior to processing for SDS-PAGE and immunoblot analysis.

Plasmid name	Description
pDL001_Hec1-ms_IgG_HC	Hec1 heavy chain variable and constant regions (mouse) + exogenous N-term signal peptide
pDL002_Hec1-ms_IgG_LC	Hec1 light chain variable and constant region (mouse) + exogenous N-term signal peptide
pDL003_Hec1-rb_IgG_HC	Hec1 heavy chain variable region + rabbit heavy chain constant regions + exogenous N-term signal peptide
pDL004_Hec1-rb_IgG_LC	Hec1 light chain variable region + rabbit light chain constant region + exogenous N-term signal peptide
pDL005_Hec1-hu_IgG_HC	Hec1 heavy chain variable region + human heavy chain constant regions (UniProt P01857) + exogenous N-term signal peptide
pDL006_Hec1-hu_IgG_LC	Hec1 light chain variable region + human light chain constant region (UniProt P01834) + exogenous N-term signal peptide
pDL007_pMELT-rb_IgG_HC	KNL1 pMELT heavy chain variable and constant regions (rabbit) + exogenous N-term signal peptide
pDL008_pMELT-rb_IgG_LC	KNL1 pMELT light chain variable and constant region (rabbit) + exogenous N-term signal peptide
pDL009_CenpC-ms_IgG_HC	CENP-C heavy chain variable region + mouse heavy chain constant regions (from Hec1 sequence) + exogenous N-term signal peptide
pDL010_CenpC-ms_IgG_LC	CENP-C light chain variable region + mouse light chain constant region (from Hec1 sequence) + exogenous N-term signal peptide
pDL011_CenpC-hu_IgG_HC	Cenp-C heavy chain variable region + human heavy chain constant regions (UniProt P01857) + exogenous N-term signal peptide
pDL012_CenpC-hu_IgG_LC	Cenp-C light chain variable region + human light chain constant region (UniProt P01834) + exogenous N-term signal peptide
pDL013_BubR1-ms_IgG_HC	BubR1 heavy chain variable and constant regions (mouse); (contains endogenous N-term signal peptide)
pDL014_BubR1-ms_IgG_LC	BubR1 light chain variable and constant region (mouse); (contains endogenous N-term signal peptide)
pDL015_BubR1-hu_IgG_HC	BubR1 heavy chain variable region + human heavy chain constant regions (UniProt P01857) (contains endogenous N-term signal peptide)
pDL016_BubR1-hu_IgG_LC	BubR1 light chain variable region + human light chain constant region (UniProt P01834) (contains endogenous N-term signal peptide)
pDL017_Mad2C-ms_IgG_HC	Mad2-C heavy chain variable and constant regions (mouse) (contains endogenous N-term signal peptide)
pDL018_Mad2C-ms_IgG_LC	Mad2-C light chain variable and constant region (mouse) (contains endogenous N-term signal peptide)
pDL019_Mad2C-hu_IgG_HC	Mad2-C heavy chain variable region + human heavy chain constant regions (UniProt P01857) (contains endogenous N-term signal peptide)
pDL020_Mad2C-hu_IgG_LC	Mad2-C light chain variable region + human light chain constant region (UniProt P01834) (contains endogenous N-term signal peptide)
pDL021_3F3/2-ms_IgG_HC	3F3/2 heavy chain variable and constant regions (mouse) (contains endogenous N-term signal peptide)
pDL022_3F3/2-ms_IgG_LC	3F3/2 light chain variable and constant region (mouse) (contains endogenous N-term signal peptide)
pDL023_Tub-rb_IgG_HC	α -tubulin heavy chain variable region + rabbit heavy chain constant regions + exogenous N-term signal peptide
pDL024_Tub-rb_IgG_LC	α -tubulin light chain variable region + rabbit light chain constant region + exogenous N-term signal peptide
pDL025_HA-rb_IgG_HC_15F11	HA-tag heavy chain hypervariable regions grafted into framework 15F11 scaffold (Zhao et al., 2019) + rabbit heavy chain constant regions + exogenous N-term signal peptide
pDL026_HA-rb_IgG_LC_15F11	HA-tag light chain hypervariable regions grafted into framework 15F11 scaffold (Zhao et al., 2019) + rabbit light chain constant regions + exogenous N-term signal peptide
pDL027_scFvC_Hec1-rb	Hec1 heavy and light chain variable regions connected by linker + rabbit heavy chain constant regions 2 and 3 + exogenous N-term signal peptide
pDL028_scFvC_Mad2C-rb	Mad2-C heavy and light chain variable regions connected by linker + rabbit heavy chain constant regions 2 and 3 + exogenous N-term signal peptide
pDL029_scFv_pMELT_GFP_NoSP	KNL1 pMELT heavy and light chain variable regions connected by linker + GFP (no signal peptide included)
pDL030_scFvC_Hec1-rb_NoSP	Hec1 heavy and light chain variable regions connected by linker + rabbit heavy chain constant regions 2 and 3 (no signal peptide included)
pDL031_scFvC_Mad2C-rb_NoSP	Mad2-C heavy and light chain variable regions connected by linker + rabbit heavy chain constant regions 2 and 3 (no signal peptide included)

Table 1. Descriptions of antibody-related plasmids generated in this study. Plasmids will be made available by contacting the corresponding author. Use of the plasmids and sequence information is for non-commercial purposes only.

669

670 **Acknowledgements**

671 The authors thank members of the DeLuca lab and Dr. Steven Markus for helpful advice on the project.
672 We also thank Drs. Martha Cyert and Marc Kirschner for providing the 3F3/2 hybridoma cell line to the
673 Gorbisky lab. This work was supported by grants from the National Institutes of Health, National Institute
674 of General Medical Sciences to JGD (R35GM130365), TJS (R35GM119728), NZ (K99GM141453), GJG
675 (R35GM126980), and DV (R01GM135391); and a grant from the National Science Foundation to TJS
676 (MCB-1845761). SMA, LS and the Cell Technology Shared Resource are supported by Cancer Center
677 Support Grant (P30CA046934).

678

679 **Competing Interests**

680
681 The authors have no competing financial interests to declare.
682

683 **References**

- 684 Alfaleh, M.A., H.O. Alsaab, A.B. Mahmoud, A.A. Alkayyal, M.L. Jones, S.M. Mahler, and A.M. Hashem.
685 2020. Phage Display Derived Monoclonal Antibodies: From Bench to Bedside. *Front Immunol.* 11:1986.
- 686 Almagro, J.C., M. Pedraza-Escalona, H.I. Arrieta, and S.M. Pérez-Tapia. 2019. Phage Display Libraries
687 for Antibody Therapeutic Discovery and Development. *Antibodies (Basel)*. 8.
- 688 Baker, M. 2015. Reproducibility crisis: Blame it on the antibodies. *Nature*. 521:274-276.
- 689 Berland, L., L. Kim, O. Abousaway, A. Mines, S. Mishra, L. Clark, P. Hofman, and M. Rashidian. 2021.
690 Nanobodies for Medical Imaging: About Ready for Prime Time? *Biomolecules*. 11.
- 691 Bordeaux, J., A. Welsh, S. Agarwal, E. Killiam, M. Baquero, J. Hanna, V. Anagnostou, and D. Rimm.
692 2010. Antibody validation. *Biotechniques*. 48:197-209.
- 693 Bradbury, A., and A. Plückthun. 2015. Reproducibility: Standardize antibodies used in research. *Nature*.
694 518:27-29.
- 695 Bradbury, A.R.M., N.D. Trinklein, H. Thie, I.C. Wilkinson, A.K. Tandon, S. Anderson, C.L. Bladen, B.
696 Jones, S.F. Aldred, M. Bestagno, O. Burrone, J. Maynard, F. Ferrara, J.S. Trimmer, J. Görnemann, J.
697 Glanville, P. Wolf, A. Frenzel, J. Wong, X.Y. Koh, H.Y. Eng, D. Lane, M.P. Lefranc, M. Clark, and S.
698 Dübel. 2018. When monoclonal antibodies are not monospecific: Hybridomas frequently express
699 additional functional variable regions. *MAbs*. 10:539-546.
- 700 Burton, D.R., J. Pyati, R. Koduri, S.J. Sharp, G.B. Thornton, P.W. Parren, L.S. Sawyer, R.M. Hendry, N.
701 Dunlop, P.L. Nara, and et al. 1994. Efficient neutralization of primary isolates of HIV-1 by a recombinant
702 human monoclonal antibody. *Science*. 266:1024-1027.
- 703 Campbell, M.S., and G.J. Gorbisky. 1995. Microinjection of mitotic cells with the 3F3/2 anti-
704 phosphoepitope antibody delays the onset of anaphase. *J Cell Biol.* 129:1195-1204.
- 705 Chan, G.K., S.A. Jablonski, V. Sudakin, J.C. Hittle, and T.J. Yen. 1999. Human BUBR1 is a mitotic
706 checkpoint kinase that monitors CENP-E functions at kinetochores and binds the cyclosome/APC. *J Cell*
707 *Biol.* 146:941-954.

708 Cheloha, R.W., T.J. Harmand, C. Wijne, T.U. Schwartz, and H.L. Ploegh. 2020. Exploring cellular
709 biochemistry with nanobodies. *J Biol Chem.* 295:15307-15327.

710 Chen, Y., D.J. Riley, P.L. Chen, and W.H. Lee. 1997. HEC, a novel nuclear protein rich in leucine heptad
711 repeats specifically involved in mitosis. *Mol Cell Biol.* 17:6049-6056.

712 Cialek, C.A., G. Galindo, A.L. Koch, M.N. Saxton, and T.J. Stasevich. 2021. Bead Loading Proteins and
713 Nucleic Acids into Adherent Human Cells. *J Vis Exp.*

714 Cosson, P., and O. Hartley. 2016. Recombinant Antibodies for Academia: A Practical Approach. *Chimia*
715 *(Aarau).* 70:893-897.

716 De Antoni, A., C.G. Pearson, D. Cimini, J.C. Canman, V. Sala, L. Nezi, M. Mapelli, L. Sironi, M. Faretta,
717 E.D. Salmon, and A. Musacchio. 2005. The Mad1/Mad2 complex as a template for Mad2 activation in
718 the spindle assembly checkpoint. *Curr Biol.* 15:214-225.

719 DeLuca, J.G., W.E. Gall, C. Ciferri, D. Cimini, A. Musacchio, and E.D. Salmon. 2006. Kinetochore
720 microtubule dynamics and attachment stability are regulated by Hec1. *Cell.* 127:969-982.

721 DeLuca, J.G., and A. Musacchio. 2012. Structural organization of the kinetochore-microtubule
722 interface. *Curr Opin Cell Biol.* 24:48-56.

723 Garg, B.K., and R.H. Loring. 2017. Evaluating Commercially Available Antibodies for Rat $\alpha 7$ Nicotinic
724 Acetylcholine Receptors. *J Histochem Cytochem.* 65:499-512.

725 Gavalondo, J.V., and J.W. Larrick. 2000. Antibody engineering at the millennium. *Biotechniques.* 29:128-
726 132, 134-126, 138 passim.

727 Gorbsky, G.J., and W.A. Ricketts. 1993. Differential expression of a phosphoepitope at the kinetochores
728 of moving chromosomes. *J Cell Biol.* 122:1311-1321.

729 Gray, A., A.R.M. Bradbury, A. Knappik, A. Plückthun, C.A.K. Borrebaeck, and S. Dübel. 2020. Animal-
730 free alternatives and the antibody iceberg. *Nat Biotechnol.* 38:1234-1239.

731 Gray, A.C., S.S. Sidhu, P.C. Chandrasekera, C.F.M. Hendriksen, and C.A.K. Borrebaeck. 2016. Animal-
732 Friendly Affinity Reagents: Replacing the Needless in the Haystack. *Trends Biotechnol.* 34:960-969.

733 Hara, M., and T. Fukagawa. 2020. Dynamics of kinetochore structure and its regulations during mitotic
734 progression. *Cell Mol Life Sci.* 77:2981-2995.

735 Hayashi-Takanaka, Y., K. Yamagata, T. Wakayama, T.J. Stasevich, T. Kainuma, T. Tsurimoto, M.
736 Tachibana, Y. Shinkai, H. Kurumizaka, N. Nozaki, and H. Kimura. 2011. Tracking epigenetic histone
737 modifications in single cells using Fab-based live endogenous modification labeling. *Nucleic Acids Res.*
738 39:6475-6488.

739 Hoffman, D.B., C.G. Pearson, T.J. Yen, B.J. Howell, and E.D. Salmon. 2001. Microtubule-dependent
740 changes in assembly of microtubule motor proteins and mitotic spindle checkpoint proteins at PtK1
741 kinetochores. *Mol Biol Cell.* 12:1995-2009.

742 Johnson, G., and T.T. Wu. 2000. Kabat database and its applications: 30 years after the first variability
743 plot. *Nucleic Acids Res.* 28:214-218.

744 Johnson, R.S., S.A. Martin, K. Biemann, J.T. Stults, and J.T. Watson. 1987. Novel fragmentation process
745 of peptides by collision-induced decomposition in a tandem mass spectrometer: differentiation of leucine
746 and isoleucine. *Anal Chem.* 59:2621-2625.

747 Katzman, B.M., K.M. Ness, and A. Algeciras-Schimmich. 2017. Evaluation of the CLSI EP26-A protocol
748 for detection of reagent lot-to-lot differences. *Clin Biochem.* 50:768-771.

749 Kimura, H., Y. Hayashi-Takanaka, T.J. Stasevich, and Y. Sato. 2015. Visualizing posttranslational and
750 epigenetic modifications of endogenous proteins in vivo. *Histochem Cell Biol.* 144:101-109.

751 Kixmoeller, K., P.K. Allu, and B.E. Black. 2020. The centromere comes into focus: from CENP-A
752 nucleosomes to kinetochore connections with the spindle. *Open Biol.* 10:200051.

753 Laustsen, A.H., V. Greiff, A. Karatt-Vellatt, S. Muyldermans, and T.P. Jenkins. 2021. Animal
754 Immunization, in Vitro Display Technologies, and Machine Learning for Antibody Discovery. *Trends*
755 *Biotechnol.*

756 Leenaars, P.P., M.A. Koedam, P.W. Wester, V. Baumans, E. Claassen, and C.F. Hendriksen. 1998.
757 Assessment of side effects induced by injection of different adjuvant/antigen combinations in rabbits and
758 mice. *Lab Anim.* 32:387-406.

759 Lima, W.C. and Cosson, P. 2019. The AA344 and AA345 antibodies detect human tubulin by
760 immunofluorescence in HeLa cells. *Antib. Rep.* 2(4):e108.

761 Lima, W.C., E. Gasteiger, P. Marcatili, P. Duek, A. Bairoch, and P. Cosson. 2020. The ABCD database:
762 a repository for chemically defined antibodies. *Nucleic Acids Res.* 48:D261-d264.

763 Lischetti, T., and J. Nilsson. 2015. Regulation of mitotic progression by the spindle assembly
764 checkpoint. *Mol Cell Oncol.* 2:e970484.

765 London, N., S. Ceto, J.A. Ranish, and S. Biggins. 2012. Phosphoregulation of Spc105 by Mps1 and PP1
766 regulates Bub1 localization to kinetochores. *Curr Biol.* 22:900-906.

767 Lyon, K., and T.J. Stasevich. 2017. Imaging Translational and Post-Translational Gene Regulatory
768 Dynamics in Living Cells with Antibody-Based Probes. *Trends Genet.* 33:322-335.

769 Mapelli, M., L. Massimiliano, S. Santaguida, and A. Musacchio. 2007. The Mad2 conformational dimer:
770 structure and implications for the spindle assembly checkpoint. *Cell.* 131:730-743.

771 McNeil, P.L., and E. Warder. 1987. Glass beads load macromolecules into living cells. *J Cell Sci.* 88 (Pt
772 5):669-678.

773 Mikhaylova, M., B.M. Cloin, K. Finan, R. van den Berg, J. Teeuw, M.M. Kijanka, M. Sokolowski, E.A.
774 Katrukha, M. Maidorn, F. Opazo, S. Moutel, M. Vantard, F. Perez, P.M. van Bergen en Henegouwen,
775 C.C. Hoogenraad, H. Ewers, and L.C. Kapitein. 2015. Resolving bundled microtubules using anti-tubulin
776 nanobodies. *Nat Commun.* 6:7933.

777 Morisaki, T., K. Lyon, K.F. DeLuca, J.G. DeLuca, B.P. English, Z. Zhang, L.D. Lavis, J.B. Grimm, S.
778 Viswanathan, L.L. Looger, T. Lionnet, and T.J. Stasevich. 2016. Real-time quantification of single RNA
779 translation dynamics in living cells. *Science.* 352:1425-1429.

780 Musacchio, A., and A. Desai. 2017. A Molecular View of Kinetochore Assembly and Function. *Biology*
781 *(Basel)*. 6.

782 Navarro, A.P., and I.M. Cheeseman. 2021. Kinetochore assembly throughout the cell cycle. *Semin Cell*
783 *Dev Biol*.

784 Nicklas, R.B., S.C. Ward, and G.J. Gorbsky. 1995. Kinetochore chemistry is sensitive to tension and may
785 link mitotic forces to a cell cycle checkpoint. *J Cell Biol*. 130:929-939.

786 Nijenhuis, W., Vallardi, G., Teixeira, A., Kops, G.J., and A. T. Saurin. 2014. Negative feedback at
787 kinetochores underlies a responsive spindle checkpoint signal. *Nat Cell Biol*. 16:1257-64.

788 Panikar, S.S., N. Banu, J. Haramati, S. Del Toro-Arreola, A. Riera Leal, and P. Salas. 2021. Nanobodies
789 as efficient drug-carriers: Progress and trends in chemotherapy. *J Control Release*. 334:389-412.

790 Pozner-Moulis, S., M. Cregger, R.L. Camp, and D.L. Rimm. 2007. Antibody validation by quantitative
791 analysis of protein expression using expression of Met in breast cancer as a model. *Lab Invest*. 87:251-
792 260.

793 Ries, J., C. Kaplan, E. Platonova, H. Eghlidi, and H. Ewers. 2012. A simple, versatile method for GFP-
794 based super-resolution microscopy via nanobodies. *Nat Methods*. 9:582-584.

795 Sachs, D.H., A.N. Schechter, A. Eastlake, and C.B. Anfinsen. 1972. An immunologic approach to the
796 conformational equilibria of polypeptides. *Proc Natl Acad Sci U S A*. 69:3790-3794.

797 Saeed, A.F., R. Wang, S. Ling, and S. Wang. 2017. Antibody Engineering for Pursuing a Healthier
798 Future. *Front Microbiol*. 8:495.

799 Santaguida, S., A. Tighe, A.M. D'Alise, S.S. Taylor, and A. Musacchio. 2010. Dissecting the role of MPS1
800 in chromosome biorientation and the spindle checkpoint through the small molecule inhibitor reversine. *J*
801 *Cell Biol*. 190:73-87.

802 Sasada, R., Marumoto, R., and Igarashi, K. 1988. Secretion of human EGF and IgE in mammalian cells
803 by recombinant DNA techniques; use of a IL-2 leader sequence. *Cell Struct Funct*. 13:129-41.

804 Sato, Y., M. Mukai, J. Ueda, M. Muraki, T.J. Stasevich, N. Horikoshi, T. Kujirai, H. Kita, T. Kimura, S.
805 Hira, Y. Okada, Y. Hayashi-Takanaka, C. Obuse, H. Kurumizaka, A. Kawahara, K. Yamagata, N. Nozaki,
806 and H. Kimura. 2013. Genetically encoded system to track histone modification in vivo. *Sci Rep*. 3:2436.

807 Sedgwick, G.G., M.S. Larsen, T. Lischetti, W. Streicher, R.R. Jersie-Christensen, J.V. Olsen, and J.
808 Nilsson. 2016. Conformation-specific anti-Mad2 monoclonal antibodies for the dissection of checkpoint
809 signaling. *MAbs*. 8:689-697.

810 Shepperd, L.A., J.C. Meadows, A.M. Sochaj, T.C. Lancaster, J. Zou, G.J. Buttrick, J. Rappsilber, K.G.
811 Hardwick, and J.B. Millar. 2012. Phosphodependent recruitment of Bub1 and Bub3 to Spc7/KNL1 by
812 Mph1 kinase maintains the spindle checkpoint. *Curr Biol*. 22:891-899.

813 Shim, H. 2017. Antibody Phage Display. *Adv Exp Med Biol*. 1053:21-34.

814 Shukra, A.M., N.V. Sridevi, C. Dev, and M. Kapil. 2014. Production of recombinant antibodies using
815 bacteriophages. *Eur J Microbiol Immunol (Bp)*. 4:91-98.

816 Skoufias, D.A., P.R. Andreassen, F.B. Lacroix, L. Wilson, and R.L. Margolis. 2001. Mammalian mad2
817 and bub1/bubR1 recognize distinct spindle-attachment and kinetochore-tension checkpoints. *Proc Natl*
818 *Acad Sci U S A*. 98:4492-4497.

819 Stasevich, T.J., Y. Hayashi-Takanaka, Y. Sato, K. Maehara, Y. Ohkawa, K. Sakata-Sogawa, M.
820 Tokunaga, T. Nagase, N. Nozaki, J.G. McNally, and H. Kimura. 2014. Regulation of RNA polymerase II
821 activation by histone acetylation in single living cells. *Nature*. 516:272-275.

822 Traenkle, B., and U. Rothbauer. 2017. Under the Microscope: Single-Domain Antibodies for Live-Cell
823 Imaging and Super-Resolution Microscopy. *Front Immunol*. 8:1030.

824 Valldorf, B., S.C. Hinz, G. Russo, L. Pekar, L. Mohr, J. Klemm, A. Doerner, S. Krah, M. Hust, and S.
825 Zielonka. 2021. Antibody display technologies: selecting the cream of the crop. *Biol Chem*.

826 Varma, D., and E.D. Salmon. 2012. The KMN protein network--chief conductors of the kinetochore
827 orchestra. *J Cell Sci*. 125:5927-5936.

828 Vazquez-Lombardi, R., D. Nevoltris, A. Luthra, P. Schofield, C. Zimmermann, and D. Christ. 2018.
829 Transient expression of human antibodies in mammalian cells. *Nat Protoc*. 13:99-117.

830 Vleugel, M., M. Omerzu, V. Groenewold, M.A. Hadders, S.M.A. Lens, and G. Kops. 2015. Sequential
831 multisite phospho-regulation of KNL1-BUB3 interfaces at mitotic kinetochores. *Mol Cell*. 57:824-835.

832 Wang, C., B. Han, R. Zhou, and X. Zhuang. 2016. Real-Time Imaging of Translation on Single mRNA
833 Transcripts in Live Cells. *Cell*. 165:990-1001.

834 Waters, J.C., R.H. Chen, A.W. Murray, and E.D. Salmon. 1998. Localization of Mad2 to kinetochores
835 depends on microtubule attachment, not tension. *J Cell Biol*. 141:1181-1191.

836 Weller, M.G. 2016. Quality Issues of Research Antibodies. *Anal Chem Insights*. 11:21-27.

837 Wimbish, R.T., and J.G. DeLuca. 2020. Hec1/Ndc80 Tail Domain Function at the Kinetochore-
838 Microtubule Interface. *Front Cell Dev Biol*. 8:43.

839 Xenaki, K.T., S. Oliveira, and P.M.P. van Bergen En Henegouwen. 2017. Antibody or Antibody
840 Fragments: Implications for Molecular Imaging and Targeted Therapy of Solid Tumors. *Front Immunol*.
841 8:1287.

842 Yamagishi, Y., C.H. Yang, Y. Tanno, and Y. Watanabe. 2012. MPS1/Mph1 phosphorylates the
843 kinetochore protein KNL1/Spc7 to recruit SAC components. *Nat Cell Biol*. 14:746-752.

844 Yan, X., T.A. Hoek, R.D. Vale, and M.E. Tanenbaum. 2016. Dynamics of Translation of Single mRNA
845 Molecules In Vivo. *Cell*. 165:976-989.

846 Yu, X., D.H. Gilden, A.M. Ritchie, M.P. Burgoon, K.M. Keays, and G.P. Owens. 2006. Specificity of
847 recombinant antibodies generated from multiple sclerosis cerebrospinal fluid probed with a random
848 peptide library. *J Neuroimmunol*. 172:121-131.

849 Zhang, G., T. Lischetti, D.G. Hayward, and J. Nilsson. 2015. Distinct domains in Bub1 localize RZZ and
850 BubR1 to kinetochores to regulate the checkpoint. *Nat Commun*. 6:7162.

851 Zhao, N., K. Kamijo, P.D. Fox, H. Oda, T. Morisaki, Y. Sato, H. Kimura, and T.J. Stasevich. 2019. A
852 genetically encoded probe for imaging nascent and mature HA-tagged proteins in vivo. *Nat Commun.*
853 10:2947.

854 Zhokhov, S.S., S.V. Kovalyov, T.Y. Samgina, and A.T. Lebedev. 2017. An EThcD-Based Method for
855 Discrimination of Leucine and Isoleucine Residues in Tryptic Peptides. *J Am Soc Mass Spectrom.*
856 28:1600-1611.

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

Supplemental Figure 2. BubR1, 3F3/2, and Mad2-C antibody classification and domain architecture. (A) Sequence data and domains for the BubR1 antibody are annotated for the heavy chain (HC) and light chain (LC) variable regions (HC=light gray; LC=dark gray), hypervariable regions (green), constant regions (HC=light orange; LC=dark orange), and the flexible hinge (dark gray). (B) Sequence data and domains for the 3F3/2 antibody are annotated for the heavy chain (HC) and light chain (LC) variable regions (HC=light turquoise; LC=dark turquoise), hypervariable regions (green), constant regions (HC=light orange; LC=dark orange), and the flexible hinge (dark gray). (C) Sequence data and domains for the Mad2-C antibody are annotated for the heavy chain (HC) and light chain (LC) variable regions (HC=light purple; LC=dark purple), hypervariable regions (green), constant regions (HC=light orange; LC=dark orange), and the flexible hinge (dark gray).