Population dynamics of Amazonian floodplain forest species support spatial variation on genetic diversity but not range expansions through time.

Gregory Thom^{1,2,*}; Camila C. Ribas³; Eduardo Shultz³; Alexandre Aleixo⁴; Cristina Y. Miyaki²

¹Department of Ornithology, American Museum of Natural History, New York, NY 10024, USA.

²Departamento de Genética e Biologia Evolutiva, Universidade de São Paulo, Rua do Matão, 277, Cidade Universitária, São Paulo, SP 05508-090, Brazil.

³Instituto Nacional de Pesquisas da Amazônia (INPA), Av. André Araújo 2936, Manaus, AM 69060-001, Brazil.

⁴Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland.

*Corresponding author. Email: gthomesilva@amnh.org

Abstract

Aim: We tested if historical demographic changes of populations occurring on the floodplains of a major Amazon Basin tributary could be associated with range expansions from upper and middle sections of the river, following the establishment of widespread river-created environments during the Late Pleistocene and Holocene.

Location: Solimoes River, Western Amazon, South America

Taxon: *Myrmoborus lugubris*, *Thamnophilus cryptoleucus* and *Myrmotherula assimilis*

Methods: We analyzed thousands of UltraConserved Elements to explore spatial patterns of genetic diversity and connectivity between individuals. Range expansions were tested with alternative methods. We quantified habitat preference for the analyzed species in order to test if the occupation of dynamic habitats could predict spatial patterns of genetic diversity.

Results: Our study did not support shared population range expansions related to historical regionalized changes in habitat availability. We found considerable variation in the spatial distribution of the genetic diversity between studied taxa, and that species with higher levels of specialization to dynamic environments have a more heterogeneous distribution of genetic diversity and reduced levels of gene flow across space.

Main conclusions: Our results suggest that demographic expansions along the Solimões River might be linked to geographic homogeneous oscillation in the distribution of floodplain environments, promoting effective population size changes but not range expansion. We found that habitat specificity might be a good predictor of population connectivity along the Amazonian floodplains.

Introduction

Understanding the factors that shape the spatial distribution of species is a long-standing challenge in comparative biology that is attributable to the complex interaction of environmental and species trait variables. Over evolutionary time scales the geographic distribution of populations might be affected by physiographic processes (Silva et al., 2019; Peter et al., 2020), climatic oscillations (Hewitt, 2000; Raposo do Amaral et al., 2018; Musher et al., 2020), biotic interactions (Araújo & Luoto, 2007; Wisz et al., 2013), and trait evolution (Alves et al., 2019), driving dispersion, extinction, and cladogenesis. Although processes such as the appearance of novel traits, interspecific competition, and colonization of new environments might lead to a single burst of range size change, in many cases species distributions are dynamic over time, oscillating between periods of range contraction followed by expansion (Davis & Shaw, 2001; Taberlet & Cheddadi, 2002; Carnaval & Moritz, 2008). Range expansions are expected to be accompanied by a rapid increase in the number of individuals, and strong spatial directionality (Excoffier et al., 2009). At the expansion front, successive founder events lead to a fast differentiation from the source, producing a characteristic demographic signature that is distinguishable from non-spatial population growth (Excoffier et al., 2009; François et al., 2010; Peter & Slatkin, 2013; Alvarado-Serrano & Hickerson, 2018). By tracking the spatial dynamics of populations one can locate zones of climatic stability (Potter et al., 2016), track sources of population invasions (Fischer et al., 2017), and understand how genetic diversity was affected by past environmental changes (Reid et al., 2019), allowing for more robust predictions to explore classical diversification hypotheses.

One of the main processes used by phylogeographic approaches to explain population size changes through time is the variation on available habitat distribution due to vegetation response to historical climate change, as described by the refugia hypothesis (Haffer, 1969, 1997;

Hewitt, 2004). This mechanism suggests that populations tend to be isolated in "islands" of optimum habitats during glacial (e.g., lowland forest taxa) or interglacial (e.g., montane forest taxa) periods followed by population expansion and secondary contact as habitats expand. In the Neotropics, the region for which the refugia hypothesis was initially proposed (Haffer, 1969), the effects of past climatic oscillations have been used to explain patterns of genetic diversity in multiple environments, including the Atlantic Forest (Carnaval et al., 2009; Amaral et al., 2018; Silva et al., 2019), Amazonia (Thom et al., 2020b), Caatinga (Gehara et al., 2017), the Chaco dry forests (Camps et al., 2018) and the Andes/Patagonia (Cosacov et al., 2010). However, few studies have sought to understand if demographic patterns are linked to range size change (Camargo et al., 2013; Guarnizo et al., 2016; Baranzelli et al., 2017). By testing models including range expansions, one can explicitly explore alternative hypotheses regarding the location of putative stable habitats through time, providing a better understanding of the historical dynamic of specific environments (Potter et al., 2016; He et al., 2017; Reid et al., 2019).

In the Amazon floodplains, historical variation in precipitation and global sea-level affected sedimentation dynamics of the main rivers (Latrubesse & Franzinelli, 2005; Irion et al., 2009; Soares et al., 2010; Pupim et al., 2019), potentially shaping the distribution and connectivity of populations occurring on river-created environments (Aleixo, 2006; Choueri et al., 2017; Thom et al., 2020b). Phylogeographic studies have reported demographic expansions for populations of birds specialized in river islands, suggesting widespread habitat alterations across the entire Amazon river system (Aleixo, 2006; Thom et al., 2020b). Historical oscillations in the availability of floodplain environments tend to be more pronounced in regions where large tributaries with distinct sediment loads meet (Gualtieri et al., 2018). For instance, the confluence of the Negro (low sediment load) and Solimoes rivers (high sediment load) leads to an increase in the water discharge and dilution of sediment concentration of the main Amazon river channel, promoting erosion and sediment bypass (Thom et al., 2020b). The current dynamics between Negro and Solimoes rivers prevent the building of long-lasting substrates for flooded habitats downstream of the confluence for approximately 150 km, until the high sediment input of the Madeira River, which restores suitable conditions for the development of large and stable floodplains in the main channel of the Amazon river (Filizola & Guyot, 2009). Recent

phylogeographic studies suggest that the central portion of the Amazon Basin, between the Negro and Madeira rivers, is indeed a suture zone, partially or completely isolating recently diverged populations, mostly restricted to Solimões, Negro, Madeira, and lower Amazonas rivers (Choueri et al., 2017; Thom et al., 2018, 2020b). This spatial congruence indicates that fluvial sedimentation dynamics may directly influence the connectivity of species occurring in river-created environments along Amazonian floodplains through time.

The historical interaction between the sediment-poor Negro River and the sediment-rich Solimões-Amazonas main channel, mediated by past oscillations in precipitation and sea level, may have been dynamic over time, shaping the availability of flooded habitats in central Amazonia. For example, the Negro River is currently impacted by a damming effect caused by the larger water input from the Solimões-Amazonas River, maintaining a ria lake at its lower section, which prevents island formation and creates a barrier for the biota associated with river-created habitats (Choueri et al., 2017). Similarly, past increases in the Negro River discharge could have periodically affected the flooding pulse of the Solimões (Irion et al., 2009; Passos et al., 2020), causing suppression of seasonally flooded habitats from its lower and middle sections (Latrubesse & Franzinelli, 2005; Soares et al., 2010; Passos et al., 2020). Under this scenario, we expect seasonally flooded habitats in the upper Solimões River to be historically more stable than areas closer to the Negro River confluence (lower Solimões River). Hence, the demographic expansions detected by previous studies for populations of birds (Thom et al., 2020b) would be linked to recent range expansions from the upper/middle towards the lower sections of the Solimões River. This demographic process should lead to higher genetic diversity on upriver populations, and if this was a pervasive mechanism, we expect to detect the same pattern in co-occurring populations that occupy similar habitats. Alternatively, population demographic expansions could be related to geographic homogeneous oscillation in habitat availability (e.g., oscillations in the carrying capacity of the environments). Under this scenario, we expect the genetic diversity to be mostly explained by the geographic distance between individuals (Isolation by distance), with a relatively homogeneous distribution of the genetic diversity across the whole extension of the Solimões River, or governed by species-specific characteristics. In the latter, species would have variable patterns across space, potentially associated with habitat specificity.

The Amazon floodplains are an intricate mosaic of microhabitats that roughly segregate with the sedimentary dynamic of the rivers and the geographic distance from the main channel (Junk et al., 2012, 2015). Areas next to the margins or on islands of rivers with high sedimentary budgets as the Solimões, tend to be more dynamic over time, keeping vegetation in early successional stages of regeneration due to the constant erosion and sedimentation. Given the high diversity of microhabitats along the floodplains, we might expect that species with higher dependence on more dynamic environments should have been more affected by historical changes on the floodplains, having a more heterogeneous spatial distribution of genetic diversity. In this case, higher association to more dynamic habitats should predict spatial levels of genetic diversity.

We explored the levels of connectivity and the spatial distribution of the genetic diversity of three antbird species, *Myrmoborus lugubris*, *Thamnophilus cryptoleucus*, and *Myrmotherula assimilis* with populations restricted to the Solimões River, testing if the population demographic expansions reported in a previous study (Thom et al., 2020b) are related to range expansions following a period of isolation in the upper/middle course of this river. We reanalyzed the data published by Thom and collaborators (2020b) using a set of methods designed to track population range expansions and explore how genetic diversity is distributed across space. Our results refute the idea of congruent range shifts related to historical regionalized changes in habitat availability. Instead, our data suggest that demographic expansions along the Solimões floodplains might be associated with variation in habitat availability across the entire range of the species, promoting effective population size changes but not range expansion. We found that species with higher levels of specialization to dynamic environments have a more heterogeneous distribution of genetic diversity and reduced levels of gene flow across space, indicating that habitat specificity might be a good predictor of population connectivity along the Amazonian floodplains.

Material and Methods

We selected samples from the Solimões River populations of *Myrmoborus lugubris* (N = 31), *Thamnophilus cryptoleucus* (N = 20), and *Myrmotherula assimilis* (N = 16) that were previously analyzed by Thom et. al (2020b; Figure 1; Table S1). Genome-wide variation was

obtained through the sequence capture of Ultraconserved Elements (UCEs) using a probe set targeting 2,312 UCE loci and 97 additional probes targeting exons commonly used in Avian phylogenomic studies (Hackett et al., 2008; Kimball et al., 2009; Faircloth et al., 2012). Raw sequences were downloaded from the Sequence Read Archive (SRA; BioProject ID: PRJNA595086; SubmissionID: SUB6206537; www.ncbi.nlm.nih.gov/bioproject/595086). We used Trinity 2.4 (Grabherr et al., 2011) to assemble *de novo* reads into contigs. The contigs were matched to the probes with LASTZ (<http://www.bx.psu.edu/~rsharris/lastz/>) using “match_contigs_to_probes.py” from PHYLUCE (Faircloth et al., 2012; Faircloth, 2016). Contigs that did not align to probe sequences and those that matched multiple loci were removed. The resulting fasta files from the different individuals were aligned in MAFFT v7.475 (Katoh & Standley, 2013), allowing for missing individuals and without trimming long ragged-ends. These long ragged-ends were then trimmed by applying a threshold of 50% of missing sequences among individuals with TrimAl v1.4.rev15 (Capella-Gutierrez et al., 2009). The longest sequence without indels of each locus was selected as a reference for the following steps. First, reads of each individual were aligned to the reference allowing 4 mismatches per read using BWA v0.7.17-r1188 (Li & Durbin, 2009). The sam files obtained were converted to bam format with Samtools v1.10 (Li et al., 2009). Reads were trimmed to match the reference using CleanSam.jar, reassigned to groups with AddOrReplaceReadGroups.jar, and duplicated reads were identified with Markduplicates.jar from PICARD v.2.0.1 (Broad Institute, Cambridge, MA; <http://broadinstitute.github.io/picard/>). We merged individual bam files into a single bam file of all samples with MergeSamFiles.jar from PICARD. With GATK v3.6 (McKenna et al., 2010) we realigned all reads, identified indels (RealignerTargetCreator; IndelRealigner), and called SNPs hard-masking low-quality bases (< Q30) with UnifiedGenotyper and VariantAnnotator. We obtained raw vcf files for each species that were filtered for a minimum read depth of > 8 using VCFTOOLS v0.1.15 (Danecek et al., 2011). Loci with any site with heterozygosity higher than 0.75 were excluded. Finally, we randomly selected one SNP per locus, excluding sites with missing data, resulting in a complete SNP matrix.

Genetic Structure

We tested the best-fit number of ancestral populations (k) for each species complex and

clustered individuals to populations in sNMF v1.2 (Frichot et al., 2014) by applying a sparse non-negative matrix factorization to compute least-square estimates of ancestry coefficients. We explored values of k between 1 and 8 and performed 100 replicates for each value with an alpha regularization parameter of 100. Given that populations occurring in linear and continuous environments such as rivers are more subjected to the effects of the geographic distance among individuals, and that IBD can generate discrete genetic clusters (Thomaz et al., 2016), we tested genetic structure controlling for the effects of IBD in the R package conStruct v1.0.3 (Bradburd et al., 2018). ConStruct simultaneously estimate continuous and discrete patterns of population structure by assuming a rate of decay in the relatedness among individuals as a function of the geographic distance. Thus, discrete groups are only assumed when genetic variation significantly deviates from an IBD scenario. We ran models assuming the same number of layers as the number of ancestral populations obtained in the best sNMF model and calculated the relative layer contribution with the function “calculate.layer.contribution”, in order to observe if the genetic structure inferred by sNMF could be explained exclusively by IBD. For each conStruct run, we performed 50,000 iterations discarding the first 50% as burn-in. We calculated the geographic distance matrices for each species following river connectivity.

Isolation by distance, effective diversity, and gene flow

We identified spatial variation in gene flow among localities and tested deviations from isolation by distance (IBD) scenario for each species with EEMS (Petkova et al., 2016). EEMS estimates the effective migration surface among drawn demes mapping genetic differentiation based on a spatially explicit approach by using a Markov Chain Monte Carlo (MCMC) to estimate effective demographic parameters given the observed genetic dissimilarity between individuals. Euclidian genetic dissimilarity matrices between individuals for each species complex were generated in ADEGENET v2.1.3 (Jombart & Ahmed, 2011). Habitat polygons were produced based on the geographic distribution of each species complex and 300 demes were distributed over the habitat area. Each MCMC run was performed for 30×10^6 generations with the first 5×10^6 generations excluded as burn-in. Maps were generated using additional features of the EEMS R package. We calculated the inbreeding coefficient (F) per individual in VCFTOOLS and interpolated the obtained values across the species distribution in QGIS

(<https://www.qgis.org/>). Additionally, to test IBD we performed a Mantel correlation test between genetic and geographic distances in ADEGENET with 999 replicates.

Geographic range expansion

We tested for population range expansion using two alternative approaches, first estimating the directionality index using rangeExpansion v0.0.0.9 in R (Peter & Slatkin, 2013, 2015) and second, estimating the geographic spectrum of shared alleles in the GSSA v0.0 program (Alvarado-Serrano & Hickerson, 2018). The rangeExpansion applies a founder effect algorithm assuming a stepping stone model of population expansion from a single location, testing the strength of the spatial expansion and the most likely origin of the founder effect, measuring the effective founder distance. The effective founder distance represents the size of the deme in which the effective population size is reduced by 1% during the founder event occurring in the expansion front. Thus, low values for the effective founder distance suggest a strong founder effect. For this approach, in order to calculate the derived state of each SNP, we assumed as outgroup one individual from a closely related population (Thom et al., 2020b).

The GSSA (Alvarado-Serrano & Hickerson, 2018) captures the range expansion signal by summarizing the shared co-ancestry between sampled localities. This statistic consists of estimating the absolute number of minor alleles shared between sampled localities, taking into account the geographic distance between localities. Given that the amount of genetic drift increases with the distance to the founder event (source of the expansion), the obtained per locality histograms contain information regarding the position of a given locality with respect to the source of the geographic expansion. Some advantages of the GSSA are that no outgroup needs to be used and it can handle less geographic dense sampling when compared to the rangeExpansion method. GSSA implementation allows for the user to provide a geographic distance matrix, instead of assuming linear distances from geographic coordinates. Given the intimate relationship of the studied species with the floodplains, we calculated the geographic distance matrixes for each species following the river's connectivity. The information contained in the shape of the GSSA histogram was summarized using Harpending's raggedness index (Harpending, 1994) and the obtained values per locality were interpolated across the distribution of each species in QGIS. We expect that locations further away from the expansion source share

more genetic variants with nearby locations that were colonized through the same expansion route, and thus have higher values for the Harpending's raggedness index. For both analyses, we selected a single individual per locality with the lowest amount of missing data.

Habitat occupancy and river dynamism

To quantify differences in habitat occupation between the three species that could predict the spatial patterns of genetic diversity we used a metric of river dynamism extracted from the Occurrence Change Intensity Map available at <https://global-surface-water.appspot.com/> (Pekel et al., 2016). This layer contains information on whether the water surface increased (e.g. erosion) or decreased (e.g. deposition of sediments) around bodies of water, and its intensity based on both the intra and inter-annual variability and changes, extracted from Landsat satellite images between 1984 and 2019. On this layer, values range from -100 to 100, where negative values represent a loss of water surface, positive values represent an increase of water surface, and zero represent stable water surfaces. To calculate the dynamicity of a radius around occurrence records for the studied species, we: 1) converted negative values to positive, so both gain and loss represent how much a given pixel has changed over time; 2) drew a buffer of 2km around every stable body of water, removing areas that were permanently water across the analyzed time section; 3) used random drawn points and occurrence records for the three species to calculate average values of dynamicity using a 1000 meters radius around the coordinates. To create the random distribution we draw 1000 points across the floodplains of the Solimoes Basin. We plotted the values for dynamicity and calculated pairwise Wilcoxon tests in R, to explore if values extracted from random points and each species statistically differ. Spatial layers were processed using the raster v3.4 in R (Hijmans & van Etten, 2013). Occurrence records were obtained from video, photos, and sound recordings from Macaulay Library (www.macaulaylibrary.org) and Xeno-canto (www.xeno-canto.org), and deposited specimens on scientific collections (Table S2). Each record was individually checked, and records with a generic locality description or low-resolution coordinates were excluded. Occurrences at the same locality and date were counted only once. Geographic coordinates outside the 2km buffer were moved to the closest pixel within the buffer.

Results

Our bioinformatics pipeline produced raw vcf files with 5,602 (16.8 mean depth; 2.8 sites per loci), 10,624 (28.4 mean depth; 4.7 sites per loci), and 10,283 (19.9 mean depth; 7.1 sites per loci) variant sites for *M. lugubris*, *T. cryptoleucus*, and *M. assimilis*, respectively. After randomly selecting one SNP per UCE loci we obtained matrices with 1,476, 1,805, and 1,776 SNPs for *M. lugubris*, *T. cryptoleucus*, and *M. assimilis*, respectively. The number of genetic clusters obtained with sNMF was partially concordant between species. For *M. lugubris* and *T. cryptoleucus*, the best value of masked cross-entropy was achieved with K=2 while for *M. assimilis* the best value suggested K=1 (Figure 2, S1). However, the results for K=2 for *M. assimilis* recovered a similar geographic structure as the ones observed in the other two species, with a gradual transition over the central portion of the Solimões basin (Figure 2). For *T. cryptoleucus* and *M. assimilis*, this transition in the ancestry coefficients occurred between Tefé and Codajas (localities 9 to 11 in Figure 1) while for *M. lugubris* it occurred more to the west, at Santo Antonio do Ica (locality 6 in Figure 1). When accounting for geographic distance with conStruct, assuming 2 layers, the relative contribution of the second layer was less than 1% and without geographic correspondence in all three species. These results suggest that the genetic structure obtained with sNMF is better explained by isolation by distance (IBD).

The EEMS detected slightly lower effective migration than the average in the eastern Solimões basin, suggesting progressively more dissimilar individuals towards the east in all three species (Figure 2). However, the estimated effective diversity produced contrasting results between species. While for *M. lugubris* we detected lower diversity in eastern localities, for *T. cryptoleucus* there was considerably lower diversity in western localities, and for *M. assimilis* we found a relatively homogeneous distribution of genetic diversity (Figures S2, S3). The results for a Mantel test were congruent with EEMS analyses, significantly supporting an IBD scenario in all three species (p-value < 0.01; Mantel r statistics of 0.59, 0.45, and 0.74 for *M. lugubris*, *T. cryptoleucus*, and *M. assimilis*, respectively). The inbreeding coefficient indicated a similar scenario as reported by EEMS effective diversity results for *M. lugubris* and *M. assimilis*, and a homogeneous distribution for *T. cryptoleucus* (Figure S4).

The geographic expansion model was strongly supported over an equilibrium/IBD model in the rangeExpansion analysis only for *T. cryptoleucus* (p-value < 0.001; q = 0.0004; Table 1;

Figure S5) indicating a geographic expansion from western Solimões. For *M. lugubris* and *M. assimilis* the equilibrium/IBD model could not be rejected (p-value of 36.17 and 0.86 for *M. lugubris* and *M. assimilis*, respectively). The GSSA estimations support a geographic directionality for the raggedness index in *M. lugubris*, with lower values in the eastern and higher values in the western Solimões as suggested by the rangeExpansion (despite the non-significant p-value; Figure 3). For *T. cryptoleucus*, despite the significant results in the rangeExpansion, there is no geographic signal supporting a range expansion using the GSSA.

Data processing of occurrence records produced 60, 92, and 44 points for *M. lugubris*, *T. cryptoleucus*, and *M. assimilis*, respectively. The pairwise Wilcoxon tests indicated that all three species occupy areas that are more dynamic than randomly drawn points, with *M. lugubris* and *T. cryptoleucus* occurring in areas significantly more dynamic than *M. assimilis* (Figure 4).

Discussion

Demographic histories are not linked to a shared pattern of range expansion

We used the genetic variation within thousands of Ultraconserved Elements to explore spatial patterns of genetic diversity in three species of antbirds that inhabit river-created environments along the Solimões River. Previous studies suggested that populations of these three species experienced synchronic demographic expansions in the last 0.2 million years ago (Thom et al., 2020b). Here we tested if demographic expansions were linked to geographic range expansions associated with historical alteration in the distribution of river-created environments (Latrubesse & Franzinelli, 2005; Irion et al., 2009; Soares et al., 2010; Govin et al., 2013; Pupim et al., 2019). Our results supported that only one species, *T. cryptoleucus*, had a significant range expansion with geographic origin in the western Solimões River. The spatial distribution of the genetic diversity reported here refutes historical events triggering common regionalized changes in population ranges along the Solimões River. Our results suggest that despite the high dynamism of floodplain environments over time, significant regional suppression of seasonally flooded habitats was probably restricted to specific portions of the floodplains. For example, the central portion of the Amazon, where the sedimentary basin of the Solimões River flows into the eastern Amazon basin, delimits the distribution of many floodplain lineages and is a suture zone splitting populations and species with distinct levels of genetic structure and gene flow (Farias &

Hrbek, 2008; Albernaz et al., 2012; Zimmer & Isler, 2016; Thom et al., 2020b). Hence, the demographic expansions reported by Thom and collaborators (2020) might be related to homogeneous oscillations in habitat availability across the entire geographic distribution of the populations, promoting population size changes but not directional range contractions and expansions.

Genetic diversity is heterogeneously distributed across the Solimões floodplains

We observed species-specific patterns for the spatial distribution of genetic diversity along the Solimões River, potentially associated with variation in the occupancy of dynamic habitats (Figures 2, S2, S3, S4). The antbird species studied here are intimately related to the river edge forests that are mostly distributed over islands of large rivers (Remsen & Parker, 1983; Junk et al., 2011). Given the low dispersal capacity of the studied species (Zimmer & Isler, 2016) and the historical dynamism of river islands, one might expect a shared pattern of genetic diversity between species. However, our data do not support this scenario, with the three species having different patterns of the spatial distribution of their genetic diversity. Despite occupying similar environments, our data suggest that species differ in their levels of association to dynamic habitats. We found that species occurring in significantly more dynamic areas have higher inbreeding coefficients, more heterogeneous spatial distribution of genetic diversity, and more variable levels of effective gene flow across space. These results indicated that habitat specificity might predict the spatial connectivity of floodplain species. Despite the congruency in historical demography, reported in previous studies (Thom et al., 2020b), our results suggest a pronounced effect of intrinsic ecological traits leading to distinct dispersal capacities affecting the spatial distribution of the genetic diversity within populations.

Heterogeneous patterns of genetic diversity have been reported along the Amazonian floodplains (Choueri et al., 2017; Jardim de Queiroz et al., 2017; dos Anjos Oliveira et al., 2019; Farias et al., 2019). Choueri and collaborators (2017) studying landscape genetics of river island specialist birds found that genetic diversity over Negro River archipelagos is not homogeneously distributed and that the temporal dynamics of formation and disappearance of islands shaped the genetic structure and gene flow between populations. Additionally, these authors suggest that the increase in sediment accumulation and island formation during the Holocene expanded

connectivity between archipelagos (Latrubesse & Franzinelli, 2005; Latrubesse & Stevaux, 2015; Choueri et al., 2017). However, in contrast to the demography of Solimões River populations, only slight signals of expansion were detected for the Negro River populations, corroborating the idea that rivers with black and clear water (sediment-poor) are less dynamic over time than white-water rivers (sediment-rich rivers), such as the Solimões (Thom et al., 2020b).

The patterns of genetic structure we detected with sNMF ($K=2$) were likely a product of the strong effects of IBD in linear distributions, as observed in the conStruct analyses, which suggested a minor contribution of a second layer (K) to explain the genetic structure. These results are in agreement with simulation-based, and empirical studies reporting the high effect of IBD in the linear distribution of river systems, including the formation of differentiated lineages (Selkoe et al., 2015; Thomaz et al., 2016; Farias et al., 2019). However, studies have suggested that the riverway distance plays a minor role in structuring genetic diversity of floodplain populations across the entire Amazonian Basin, suggesting the more pronounced effect of other environmental features (Hrbek et al., 2005; dos Anjos Oliveira et al., 2019; Thom et al., 2020b). Our results reinforce the idea that IBD has to be taken into account when testing for genetic structure in continuous populations on the risk of assuming spurious discrete populations in downstream population genetics analyses.

Future directions to explore spatial patterns of genetic diversity across the Amazonian floodplains

Our study provided new insights on the spatial dynamics of floodplain forest environments over time, refuting a pattern of common range expansion for co-occurring species. However, our data is limited in the sense that more sampled localities could improve our power in detecting range size changes. Although studies have reported robust results with the rangeExpansion method (Peter & Slatkin, 2013) sampling a single diploid individual for 10 localities and 1,500 SNPs, which is a data set similar to ours, the ideal data set lies around 20 diploid individuals (1 per locality) and 7,000 SNPs (Peter & Slatkin, 2013, 2015; Potter et al., 2016). An important factor that might hinder our ability to detect if a range expansion occurred is the time since populations reached a stationary condition (Alvarado-Serrano & Hickerson, 2018). All three species we analyzed have extensive distributions across the Solimões River, and might

already have reached isolation by distance equilibrium, eroding the signal for directional range expansions. Recent evidence has supported that secondary contact and gene flow are pervasive processes across the floodplains, although controlled by varying degrees of connectivity through time (Thom et al., 2018, 2020b). The methods we applied do not incorporate the effects of gene flow from unsampled populations, which could inflate genetic diversity in eastern Solimões localities, masking the genomic signal of range expansions. The inferred range expansion in *T. cryptoleucus* pointed to an interesting scenario, since this taxon is currently in contact with *T. nigrocinereus tschudi* from the Madeira River, suggesting diversification in allopatry followed by range expansion and recent secondary contact. These results suggest that the approach we used might be robust to some level of gene flow from distinct populations. Future studies aiming to explore the spatial demographic dynamics of floodplain populations should target widespread geographic distribution of samples (e.g. one individual per locality and many localities), and more genetic markers, obtained with reduced representation techniques such as RADseq (Andrews et al., 2016), or whole-genome sequencing (Ellegren, 2014). Additionally, spatial explicit simulations of demographic histories coupled with supervised machine learning classification algorithms could improve our ability to test range size changes through time with a reduced number of samples (Battey et al. 2020; Reid et al. 2019).

Ecological niche modeling has been used to track the current and paleo distribution of various Amazonian organisms (Prates et al., 2016; dos Anjos Oliveira et al., 2019; Silva et al., 2019). However, the intimate relationship of floodplain species with microhabitats linked to the sedimentation dynamics of the rivers, probably cannot be captured by interpolated bioclimatic variables (e.g. Worldclim). Meaningful variables that could describe the sedimentation process and detailed distribution of floodplain habitats with a good resolution for the entire Basin are becoming available (Pekel et al., 2016; Fassoni-Andrade & Paiva, 2019) and might provide new insights on how the current sediment budget of rivers and vegetation dynamics shape patterns of genetic diversity. Amazonia is currently at the center of the debate about the impact of hydroelectric dams on biodiversity and the sedimentary budget of rivers (Latrubesse et al., 2017). Understanding how the genetic diversity (dos Anjos Oliveira et al., 2019) and species richness (Laranjeiras et al., 2021) might be associated with river dynamism and productivity is paramount to predict future impacts caused by anthropic activities.

Acknowledgments

We thank the curatorial staff of the Museu Paraense Emílio Goeldi (MPEG), Instituto Nacional de Pesquisas da Amazônia (INPA), Laboratório de Genética e Evolução Molecular de Aves (LGEMA) for loaning tissue samples under their care. We thank members of the Smith lab, Hickerson lab, and Carnaval lab: Brian Smith, Kaiya L. Provost, Lucas R. Moreira, Vivien Chua, Jon Merwin, Ana C. Carnaval, Connor French, Andrea Paz, Kathryn Mercier, Rilquer Mascarenhas, Lidia Martins, Michael Hickerson, Isaac Overcast, Alexander Xue. We thank L. E. Araujo-Silva for the partnership during field expeditions. We thank Oscar Johnson and Rob Brumfield for sharing UCE sequences. We thank the Research Center on Biodiversity and Computing (BioComp) of the Universidade de São Paulo (USP), supported by the USP Provost's Office for Research. This study was co-funded by FAPESP (BIOTA, 2012/50260-6 and 2013/50297-0), NSF (DOB 1343578), NASA, CNPq (310593/2009-3, 574008/2008-0, 563236/2010-8, and 471342/2011-4), PEER/USAID (AID-OAA-A-11-00012), and FAPESPA (ICAAF 023/2011). G.T. was granted by CAPES and then FAPESP scholarships (2014/00113-2, 2015/12551-7, 2018/17869-3, and 2017/25720-7). A.A., C.C.R., and C.M. are supported by CNPq research productivity fellowships (310880/2012-2, 308927/2016-8, and 3062204/2019-3).

Biosketch

Gregory Thom is broadly interested in comparative genomics, testing the major factors impacting population differentiation over historical timescales. His overriding goal is to unearth the mechanisms driving Neotropical bird evolution by using genomics and advanced computational biology.

Author contributions

G.T. conceived the study and performed bioinformatics and phylogeographic analyses. G.T., C.C.R., and C.M. wrote the paper with input from A.A. and ES. GT and ES performed the river dynamicity analyses. All authors reviewed and approved the final version of the manuscript before submission.

References

- Albernaz, A.L., Pressey, R.L., Costa, L.R.F., Moreira, M.P., Ramos, J.F., Assunção, P.A., & Franciscon, C.H. (2012) Tree species compositional change and conservation implications in the white-water flooded forests of the Brazilian Amazon: Compositional patterns of floodplain tree species. *Journal of Biogeography*, **39**, 869–883.
- Aleixo, A. (2006) Historical diversification of floodplain forest specialist species in the Amazon: a case study with two species of the avian genus *Xiphorhynchus* (Aves: Dendrocolaptidae). *Biological journal of the Linnean Society. Linnean Society of London*, **89**, 383–395.
- Alvarado-Serrano, D.F. & Hickerson, M.J. (2018) Detecting spatial dynamics of range expansions with geo-referenced genomewide SNP data and the geographic spectrum of shared alleles. *bioRxiv*, .
- Alves, J.A., Gunnarsson, T.G., Sutherland, W.J., Potts, P.M., & Gill, J.A. (2019) Linking warming effects on phenology, demography, and range expansion in a migratory bird population. *Ecology and Evolution*, **9**, 2365–2375.
- Amaral, F.R., Alvarado-Serrano, D.F., Maldonado-Coelho, M., Pellegrino, K.C.M., Miyaki, C.Y., Montesanti, J.A.C., Lima-Ribeiro, M.S., Hickerson, M.J., & Thom, G. (2018) Climate explains recent population divergence, introgression and persistence in tropical mountains: phylogenomic evidence from Atlantic Forest warbling finches. *bioRxiv*, 439265.
- Andrews, K.R., Good, J.M., Miller, M.R., Luikart, G., & Hohenlohe, P.A. (2016) Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews. Genetics*, **17**, 81–92.
- dos Anjos Oliveira, J., Farias, I.P., Costa, G.C., & Werneck, F.P. (2019) Model-based riverscape genetics: disentangling the roles of local and connectivity factors in shaping spatial genetic patterns of two Amazonian turtles with different dispersal abilities. *Evolutionary Ecology*, **33**, 273–298.
- Araújo, M.B. & Luoto, M. (2007) The importance of biotic interactions for modelling species distributions under climate change. *Global Ecology and Biogeography: a Journal of Macroecology*, **16**, 743–753.
- Baranzelli, M.C., Cosacov, A., Ferreiro, G., Johnson, L.A., & Sérsic, A.N. (2017) Travelling to the south: Phylogeographic spatial diffusion model in *Monttea aphylla* (Plantaginaceae), an endemic plant of the Monte Desert. *PloS One*, **12**, e0178827.
- Battey, C.J., Ralph, P.L., & Kern, A.D. (2020) Space is the place: effects of continuous spatial structure on analysis of population genetic data. *Genetics*, **215**, 193–214.
- Bradburd, G.S., Coop, G.M., & Ralph, P.L. (2018) Inferring continuous and discrete population

485 genetic structure across space. *Genetics*, **210**, 33–52.

486 Camargo, A., Werneck, F.P., Morando, M., Sites, J.W., & Avila, L.J. (2013) Quaternary range
487 and demographic expansion of *Liolaemus darwini* (Squamata: Liolaemidae) in the Monte
488 Desert of Central Argentina using Bayesian phylogeography and ecological niche
489 modelling. *Molecular Ecology*, **22**, 4038–4054.

490 Camps, G.A., Martínez-Meyer, E., Verga, A.R., Sérsic, A.N., & Cosacov, A. (2018) Genetic and
491 climatic approaches reveal effects of Pleistocene refugia and climatic stability in an old
492 giant of the Neotropical Dry Forest. *Biological Journal of the Linnean Society*, **125**, 401–
493 420.

494 Capella-Gutierrez, S., Silla-Martinez, J.M., & Gabaldon, T. (2009) trimAl: a tool for automated
495 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, **25**, 1972–1973.

496 Carnaval, A.C., Hickerson, M.J., Haddad, C.F.B., Rodrigues, M.T., & Moritz, C. (2009) Stability
497 predicts genetic diversity in the Brazilian Atlantic forest hotspot. *Science*, **323**, 785–789.

498 Carnaval, A.C. & Moritz, C. (2008) Historical climate modelling predicts patterns of current
499 biodiversity in the Brazilian Atlantic forest. *Journal of Biogeography*, **35**, 1187–1201.

500 Choueri, É.L., Gubili, C., Borges, S.H., Thom, G., Sawakuchi, A.O., Soares, E.A.A., & Ribas,
501 C.C. (2017) Phylogeography and population dynamics of Antbirds (Thamnophilidae) from
502 Amazonian fluvial islands. *Journal of Biogeography*, **44**, 2284–2294.

503 Cosacov, A., Sérsic, A.N., Sosa, V., Johnson, L.A., & Cocucci, A.A. (2010) Multiple periglacial
504 refugia in the Patagonian steppe and post-glacial colonization of the Andes: the
505 phylogeography of *Calceolaria polyrhiza*: Phylogeography of *Calceolaria polyrhiza*.
506 *Journal of Biogeography*, no–no.

507 Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
508 Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., & 1000 genomes project
509 analysis group (2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–
510 2158.

511 Davis, M.B. & Shaw, R.G. (2001) Range shifts and adaptive responses to Quaternary climate
512 change. *Science*, **292**, 673–679.

513 Ellegren, H. (2014) Genome sequencing and population genomics in non-model organisms.
514 *Trends in Ecology & Evolution*, **29**, 51–63.

515 Excoffier, L., Foll, M., & Petit, R.J. (2009) Genetic consequences of range expansions. *Annual
516 Review of Ecology, Evolution, and Systematics*, **40**, 481–501.

517 Faircloth, B.C. (2016) PHYLUCE is a software package for the analysis of conserved genomic
518 loci. *Bioinformatics*, **32**, 786–788.

- 519 Faircloth, B.C., McCormack, J.E., Crawford, N.G., Harvey, M.G., Brumfield, R.T., & Glenn,
520 T.C. (2012) Ultraconserved elements anchor thousands of genetic markers spanning
521 multiple evolutionary timescales. *Systematic Biology*, **61**, 717–726.
- 522 Farias, I.P. & Hrbek, T. (2008) Patterns of diversification in the discus fishes (*Symphysodon* spp.
523 Cichlidae) of the Amazon basin. *Molecular Phylogenetics and Evolution*, **49**, 32–43.
- 524 Farias, I.P., Willis, S., Leão, A., Verba, J.T., Crossa, M., Foresti, F., Porto-Foresti, F., Sampaio,
525 I., & Hrbek, T. (2019) The largest fish in the world’s biggest river: Genetic connectivity and
526 conservation of *Arapaima gigas* in the Amazon and Araguaia-Tocantins drainages. *PLOS*
527 *ONE*, **14**, e0220882.
- 528 Fassoni-Andrade, A.C. & Paiva, R.C.D. de (2019) Mapping spatial-temporal sediment dynamics
529 of river-floodplains in the Amazon. *Remote Sensing of Environment*, **221**, 94–107.
- 530 Filizola, N. & Guyot, J.L. (2009) Suspended sediment yields in the Amazon basin: an assessment
531 using the Brazilian national data set. *Hydrological Processes*, **23**, 3207–3215.
- 532 Fischer, M.L., Salgado, I., Beninde, J., Klein, R., Frantz, A.C., Heddergott, M., Cullingham, C.I.,
533 Kyle, C.J., & Hochkirch, A. (2017) Multiple founder effects are followed by range
534 expansion and admixture during the invasion process of the raccoon (*Procyon lotor*) in
535 Europe. *Diversity & Distributions*, **23**, 409–420.
- 536 François, O., Currat, M., Ray, N., Han, E., Excoffier, L., & Novembre, J. (2010) Principal
537 component analysis under population genetic models of range expansion and admixture.
538 *Molecular Biology and Evolution*, **27**, 1257–1268.
- 539 Frichot, E., Mathieu, F., Trouillon, T., Bouchard, G., & François, O. (2014) Fast and efficient
540 estimation of individual ancestry coefficients. *Genetics*, **196**, 973–983.
- 541 Gehara, M., Garda, A.A., Werneck, F.P., Oliveira, E.F., da Fonseca, E.M., Camurugi, F.,
542 Magalhães, F. de M., Lanna, F.M., Sites, J.W., Jr, Marques, R., Silveira-Filho, R., São
543 Pedro, V.A., Colli, G.R., Costa, G.C., & Burbrink, F.T. (2017) Estimating synchronous
544 demographic changes across populations using hABC and its application for a
545 herpetological community from northeastern Brazil. *Molecular Ecology*, **26**, 4756–4771.
- 546 Govin, A., Chiessi, C.M., Zabel, M., Sawakuchi, A.O., Heslop, D., Hörner, T., Zhang, Y., &
547 Mulitza, S. (2013) Terrigenous input off northern South America driven by changes in
548 Amazonian climate and the North Brazil Current retroflexion during the last 250 ka.
549 *Climate of the Past Discussions*, **9**, 5855–5898.
- 550 Grabherr, M.G., Haas, B.J., Yassour, M., et al. (2011) Full-length transcriptome assembly from
551 RNA-Seq data without a reference genome. *Nature Biotechnology*, **29**, 644–652.
- 552 Gualtieri, C., Filizola, N., de Oliveira, M., Santos, A.M., & Ianniruberto, M. (2018) A field study
553 of the confluence between Negro and Solimões Rivers. Part 1: Hydrodynamics and

554 sediment transport. *Comptes Rendus Geoscience*, **350**, 31–42.

555 Guarnizo, C.E., Werneck, F.P., Giugliano, L.G., Santos, M.G., Fenker, J., Sousa, L.,
556 D'Angiolella, A.B., Dos Santos, A.R., Strüssmann, C., Rodrigues, M.T., Dorado-Rodrigues,
557 T.F., Gamble, T., & Colli, G.R. (2016) Cryptic lineages and diversification of an endemic
558 anole lizard (Squamata, Dactyloidae) of the Cerrado hotspot. *Molecular Phylogenetics and*
559 *Evolution*, **94**, 279–289.

560 Hackett, S.J., Kimball, R.T., Reddy, S., Bowie, R.C.K., Braun, E.L., Braun, M.J., Chojnowski,
561 J.L., Cox, W.A., Han, K.-L., Harshman, J., Huddleston, C.J., Marks, B.D., Miglia, K.J.,
562 Moore, W.S., Sheldon, F.H., Steadman, D.W., Witt, C.C., & Yuri, T. (2008) A
563 phylogenomic study of birds reveals their evolutionary history. *Science*, **320**, 1763–1768.

564 Haffer, J. (1969) Speciation in amazonian forest birds. *Science*, **165**, 131–137.

565 Haffer, J. (1997) Alternative models of vertebrate speciation in Amazonia: an overview.
566 *Biodiversity & Conservation*, **6**, 451–476.

567 Harpending, H.C. (1994) Signature of ancient population growth in a low-resolution
568 mitochondrial DNA mismatch distribution. *Human Biology*, **66**, 591–600.

569 He, Q., Prado, J.R., & Knowles, L.L. (2017) Inferring the geographic origin of a range
570 expansion: Latitudinal and longitudinal coordinates inferred from genomic data in an ABC
571 framework with the program x-origin. *Molecular Ecology*, **26**, 6908–6920.

572 Hewitt, G. (2000) The genetic legacy of the Quaternary ice ages. *Nature*, **405**, 907–913.

573 Hewitt, G.M. (2004) Biodiversity: A climate for colonization. *Heredity*, **92**, 1–2.

574 Hijmans, R.J. & van Etten, J. (2013) Raster: raster: Geographic data analysis and modeling. *R*
575 *package version*, 2–3.

576 Hrbek, T., Farias, I.P., Crossa, M., Sampaio, I., Porto, J.I.R., & Meyer, A. (2005) Population
577 genetic analysis of *Arapaima gigas*, one of the largest freshwater fishes of the Amazon
578 basin: implications for its conservation. *Animal Conservation*, **8**, 297–308.

579 Irion, G., Müller, J., Morais, J.O., Keim, G., de Mello, J.N., & Junk, W.J. (2009) The impact of
580 Quaternary sea level changes on the evolution of the Amazonian lowland. *Hydrological*
581 *Processes*, **23**, 3168–3172.

582 Jardim de Queiroz, L., Torrente-Vilara, G., Quilodran, C., Rodrigues da Costa Doria, C., &
583 Montoya-Burgos, J.I. (2017) Multifactorial genetic divergence processes drive the onset of
584 speciation in an Amazonian fish. *PloS One*, **12**, e0189349.

585 Jombart, T. & Ahmed, I. (2011) adegenet 1.3-1: new tools for the analysis of genome-wide SNP
586 data. *Bioinformatics*, **27**, 3070–3071.

587 Junk, W.J., Piedade, M.T.F., Schöngart, J., Cohn-Haft, M., Adeney, J.M., & Wittmann, F. (2011)
588 A classification of major naturally-occurring Amazonian lowland wetlands. *Wetlands*, **31**,
589 623–640.

590 Junk, W.J., Piedade, M.T.F., Schöngart, J., & Wittmann, F. (2012) A classification of major
591 natural habitats of Amazonian white-water river floodplains (várzeas). *Wetlands Ecology*
592 *and Management*, **20**, 461–475.

593 Junk, W.J., Wittmann, F., Schöngart, J., & Piedade, M.T.F. (2015) A classification of the major
594 habitats of Amazonian black-water river floodplains and a comparison with their white-
595 water counterparts. *Wetlands Ecology and Management*, **23**, 677–693.

596 Katoh, K. & Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7:
597 improvements in performance and usability. *Molecular Biology and Evolution*, **30**, 772–
598 780.

599 Kimball, R.T., Braun, E.L., Barker, F.K., Bowie, R.C.K., Braun, M.J., Chojnowski, J.L.,
600 Hackett, S.J., Han, K.-L., Harshman, J., Heimer-Torres, V., Holznagel, W., Huddleston,
601 C.J., Marks, B.D., Miglia, K.J., Moore, W.S., Reddy, S., Sheldon, F.H., Smith, J.V., Witt,
602 C.C., & Yuri, T. (2009) A well-tested set of primers to amplify regions spread across the
603 avian genome. *Molecular Phylogenetics and Evolution*, **50**, 654–660.

604 Laranjeiras, T.O., Naka, L.N., Leite, G.A., & Cohn Haft, M. (2021) Effects of a major
605 Amazonian river confluence on the distribution of floodplain forest avifauna. *Journal of*
606 *Biogeography*, **48**, 847-860.

607 Latrubesse, E.M., Arima, E.Y., Dunne, T., Park, E., Baker, V.R., d’Horta, F.M., Wight, C.,
608 Wittmann, F., Zuanon, J., Baker, P.A., Ribas, C.C., Norgaard, R.B., Filizola, N., Ansar, A.,
609 Flyvbjerg, B., & Stevaux, J.C. (2017) Damming the rivers of the Amazon basin. *Nature*,
610 **546**, 363–369.

611 Latrubesse, E.M. & Franzinelli, E. (2005) The late Quaternary evolution of the Negro River,
612 Amazon, Brazil: Implications for island and floodplain formation in large anabranching
613 tropical systems. *Geomorphology*, **70**, 372–397.

614 Latrubesse, E.M. & Stevaux, J.C. (2015) The Anavilhanas and Mariuá Archipelagos: Fluvial
615 Wonders from the Negro River, Amazon Basin. *World Geomorphological Landscapes*,
616 157–169.

617 Li, H. & Durbin, R. (2009) Fast and accurate short read alignment with Burrows–Wheeler
618 transform. *Bioinformatics*, **25**, 1754–1760.

619 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
620 Durbin, R., & 1000 Genome Project Data Processing Subgroup (2009) The Sequence
621 Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.

- 622 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernysky, A., Garimella,
623 K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M.A. (2010) The Genome Analysis
624 Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
625 *Genome Research*, **20**, 1297–1303.
- 626 Musher, L.J., Galante, P.J., Thom, G., Huntley, J.W., & Blair, M.E. (2020) Shifting ecosystem
627 connectivity during the Pleistocene drove diversification and gene flow in a species
628 complex of Neotropical birds (Tityridae: *Pachyramphus*). *Journal of Biogeography*, **47**,
629 1714–1726.
- 630 Passos, M.S., Soares, E.A.A., Tatumi, S.H., Yee, M., Mittani, J.C.R., Hayakawa, E.H., &
631 Salazar, C.A. (2020) Pleistocene-Holocene sedimentary deposits of the Solimões-Amazonas
632 fluvial system, Western Amazonia. *Journal of South American Earth Sciences*, **98**, 102455.
- 633 Pekel, J.-F., Cottam, A., Gorelick, N., & Belward, A.S. (2016) High-resolution mapping of
634 global surface water and its long-term changes. *Nature*, **540**, 418–422.
- 635 Peter, B.M., Petkova, D., & Novembre, J. (2020) Genetic Landscapes Reveal How Human
636 Genetic Diversity Aligns with Geography. *Molecular Biology and Evolution*, **37**, 943–951.
- 637 Peter, B.M. & Slatkin, M. (2013) Detecting range expansions from genetic data. *Evolution;*
638 *International Journal of Organic Evolution*, **67**, 3274–3289.
- 639 Peter, B.M. & Slatkin, M. (2015) The effective founder effect in a spatially expanding
640 population. *Evolution; International Journal of Organic Evolution*, **69**, 721–734.
- 641 Petkova, D., Novembre, J., & Stephens, M. (2016) Visualizing spatial population structure with
642 estimated effective migration surfaces. *Nature Genetics*, **48**, 94–100.
- 643 Potter, S., Bragg, J.G., Peter, B.M., Bi, K., & Moritz, C. (2016) Phylogenomics at the tips:
644 inferring lineages and their demographic history in a tropical lizard, *Carlia amax*.
645 *Molecular Ecology*, **25**, 1367–1380.
- 646 Prates, I., Xue, A.T., Brown, J.L., Alvarado-Serrano, D.F., Rodrigues, M.T., Hickerson, M.J., &
647 Carnaval, A.C. (2016) Inferring responses to climate dynamics from historical demography
648 in neotropical forest lizards. *Proceedings of the National Academy of Sciences of the United*
649 *States of America*, **113**, 7978–7985.
- 650 Pupim, F.N., Sawakuchi, A.O., Almeida, R.P., Ribas, C.C., Kern, A.K., Hartmann, G.A.,
651 Chiessi, C.M., Tamura, L.N., Mineli, T.D., Savian, J.F., Grohmann, C.H., Bertassoli, D.J.,
652 Stern, A.G., Cruz, F.W., & Cracraft, J. (2019) Chronology of Terra Firme formation in
653 Amazonian lowlands reveals a dynamic Quaternary landscape. *Quaternary Science*
654 *Reviews*, **210**, 154–163.
- 655 Raposo do Amaral, F., Maldonado-Coelho, M., Aleixo, A., Luna, L.W., Rêgo, P.S. do, Araripe,
656 J., Souza, T.O., Silva, W.A.G., & Thom, G. (2018) Recent chapters of Neotropical history

- 657 overlooked in phylogeography: Shallow divergence explains phenotype and genotype
658 uncoupling in *Antilophia manakins*. *Molecular Ecology*, **27**, 4108–4120.
- 659 Reid, B.N., Kass, J.M., Wollney, S., Jensen, E.L., Russello, M.A., Viola, E.M., Pantophlet, J.,
660 Iverson, J.B., Peery, M.Z., Raxworthy, C.J., & Naro-Maciel, E. (2019) Disentangling the
661 genetic effects of refugial isolation and range expansion in a trans-continentially distributed
662 species. *Heredity*, **122**, 441–457.
- 663 Remsen, J.V., Jr & Parker, T.A., III (1983) Contribution of river-created habitats to bird species
664 richness in Amazonia. *Biotropica*, **15**, 223–231.
- 665 Selkoe, K.A., Scribner, K.T., & Galindo, H.M. (2015) Waterscape genetics--applications of
666 landscape genetics to rivers, lakes, and seas. *Landscape Genetics*, **13**, 220–246.
- 667 Silva, S.M., Peterson, A.T., Carneiro, L., et al. (2019) A dynamic continental moisture gradient
668 drove Amazonian bird diversification. *Science Advances*, **5**, eaat5752.
- 669 Soares, E.A.A., Tatumi, S.H., & Riccomini, C. (2010) OSL age determinations of Pleistocene
670 fluvial deposits in Central Amazonia. *Anais da Academia Brasileira de Ciências*, **82**, 691–
671 699.
- 672 Taberlet, P. & Cheddadi, R. (2002) Ecology. Quaternary refugia and persistence of biodiversity.
673 *Science*, **297**, 2009–2010.
- 674 Thomaz, A.T., Christie, M.R., & Knowles, L.L. (2016) The architecture of river networks can
675 drive the evolutionary dynamics of aquatic populations. *Evolution; International Journal of*
676 *Organic Evolution*, **70**, 731–739.
- 677 Thom, G., Do Amaral, F.R., Hickerson, M.J., Aleixo, A., Araujo-Silva, L.E., Ribas, C.C.,
678 Choueri, E., & Miyaki, C.Y. (2018) Phenotypic and genetic structure support gene flow
679 generating gene tree discordances in an Amazonian Floodplain Endemic Species.
680 *Systematic Biology*, **67**, 700–718.
- 681 Thom, G., Smith, B.T., Gehara, M., Montesanti, J., Lima-Ribeiro, M.S., Piacentini, V.Q.,
682 Miyaki, C.Y., & do Amaral, F.R. (2020a) Climatic dynamics and topography control
683 genetic variation in Atlantic Forest montane birds. *Molecular Phylogenetics and Evolution*,
684 **148**, 106812.
- 685 Thom, G., Xue, A.T., Sawakuchi, A.O., Ribas, C.C., Hickerson, M.J., Aleixo, A., & Miyaki, C.
686 (2020b) Quaternary climate changes as speciation drivers in the Amazon floodplains.
687 *Science Advances*, **6**, eaax4718.
- 688 Wisz, M.S., Pottier, J., Kissling, W.D., et al. (2013) The role of biotic interactions in shaping
689 distributions and realised assemblages of species: implications for species distribution
690 modelling. *Biological Reviews of the Cambridge Philosophical Society*, **88**, 15–30.

Zimmer, K. & Isler, M.L. (2016) Typical Antbirds (Thamnophilidae). *Handbook of the Birds of the World Alive*. J. del Hoyo, A. Elliott, J. Sargatal, D.A. Christie, E. de Juana (Eds). Lynx Edicions: Barcelona. Available at <http://www.hbw.com/node/52291> [Verified 29 July 2016].

Table 1: Summary of results obtained with rangeExpansion for three species of birds occurring across the Solimoes River.

Parameters	<i>M. lugubris</i>	<i>T. cryptoleucus</i>	<i>M. assimilis</i>
longitude	-62.95	-73.22144	-75.64357
latitude	-4.021451	-9.999475	-3148
r1	0.999	0.999	0.999

r10	0.996	0.991	0.997
r100	0.964	0.918	0.975
d1	27.19	11.45	40.80
r ²	0.45	0.89	0.31
p-value	36.17	2.10e-23	11.26

717 r1, r10, and r100 - decrease in diversity at 1, 10, and 100 kilometers of distance, respectively; d1
718 – effective founder distance, which represents the deme size in km for the effective population
719 size to be reduced by 1% in the expansion front; r² and p-value - correlation coefficient and
720 correlation p-value for the most likely origin, respectively.

721

722

723

724 Figure 1: Geographic distribution of sampled individuals of *Myrmoborus lugubris* (l),
725 *Thamnophilus cryptoleucus* (c), and *Myrmotherula assimilis* (a) across the Solimões River Basin.
726 Numbers in the map represent localities as in Table S1. In red is the putative zone of unstable

727 habitats following the range expansion hypothesis.

728
729 Figure 2: Estimated effective migration surface (EEMS) and population structure and individual
730 coefficient of ancestry (bars) inferred with sNMF for the three species studied. Colors in the map
731 represent a log10 scale from the average effective migration (gray). Codes below the bars in the
732 admixture plot represent identification numbers of individuals as in Table S1. Note that for *M.*
733 *assimilis*, K=2 was the model with the second-highest cross-entropy value.

734

Figure 3: Interpolated Harpending's raggedness index estimated from the geographic spectrum of shared alleles (GSSA) histograms. Lower values represent the most likely source of a serial range expansion.

Figure 4: Average river dynamism on 1000 meters radius around randomly drawn points and occurrence records for *M. assimilis*, *M. lugubris*, and *T. cryptoleucus*. River dynamism was calculated based on the "Water Occurrence Change" layer available at <https://global-surface-water.appspot.com/>. Higher values represent higher river dynamism. The pairwise comparisons show p-values for Wilcoxon tests between species and random points.