Intact reading ability in spite of a spatially distributed visual word form ‘area’ in an individual born without the left superior temporal lobe

Authors

Jin Li¹, Evelina Fedorenko², Zeynep Saygin^{1*}

Affiliations

¹Department of Psychology, The Ohio State University, Columbus, OH, USA.

²Department of Brain and Cognitive Sciences / McGovern Institute for Brain Research, MIT, Cambridge, MA, USA

*Correspondence to: li.9361@osu.edu and saygin.3@osu.edu

Acknowledgements

We would like to thank our participant – EG – for her time and willingness to participate in brain research. We thank Greta Tuckute, Alexander Paunov, and Benjamin Lipkin for their help in coordinating the collaboration as well as their suggestions for data analysis. We also thank Idan Blank and Zachary Mineroff for help with EG’s fMRI and behavioral data collection and members of Saygin Developmental Cognitive Neuroscience Lab for helping with control data collection. We would like to acknowledge the Athinoula A. Martinos Imaging Center at the McGovern Institute for Brain Research, MIT and the support from Center for Cognitive and Behavioral Brain Imaging (CCBBI) and Ohio Supercomputer Center (OSC). E.F. was supported by NIH awards R01 DC016607, R01 DC016950, and research funds from the McGovern Institute for Brain Research and the Department of Brain and Cognitive Sciences at MIT. Z.M.S. was supported by the Alfred P. Sloan Foundation and OSU’s Arts & Sciences College and Chronic Brain Injury initiative.

Conflict of interest

The authors declare no competing financial interests.

32

33

Abstract

34

35 The visual word form area (VWFA) is an experience-dependent brain region in the left ventral
 36 temporal cortex of literate adults that responds selectively to visual words. Why does it emerge
 37 in this stereotyped location? Past research has shown that the VWFA is preferentially connected
 38 to the left-lateralized frontotemporal language network. However, it remains unclear whether the
 39 presence of a typical language network and its connections with ventral temporal cortex (VTC)
 40 are critical for the VWFA's emergence, and whether alternative functional architectures may
 41 support reading ability. We explored these questions in an individual (EG) born without the left
 42 superior temporal lobe but exhibiting normal reading ability. Using fMRI, we recorded brain
 43 activation to visual words, objects, faces, and scrambled words in EG and neurotypical controls.
 44 We did not observe word selectivity either in EG's right homotop of the VWFA (rVWFA)—the
 45 most expected location given that EG's language network is right-lateralized—or in her spared
 46 left VWFA (lVWFA), in the presence of typical face selectivity in both the right and left
 47 fusiform face area (rFFA, lFFA). Interestingly, multivariate pattern analyses revealed voxels in
 48 EG's rVWFA and lVWFA that showed 1) higher within- than between-category correlations for
 49 words (e.g., Words-Words>Words-Faces), and 2) higher within-category correlations for words
 50 than other categories (e.g., Words-Words>Faces-Faces). These results suggest that a typical left-
 51 hemisphere language network may be necessary for the emergence of focal word selectivity
 52 within ventral temporal cortex, and that orthographic processing may depend on a distributed
 53 neural code, which appears capable of supporting reading ability.

54

55 **Keywords:** lesion study; Visual Word Form Area (VWFA); brain plasticity; word selectivity;
 56 multivariate pattern analysis (MVPA); category selectivity

Introduction

In the past two decades, numerous regions in the ventral temporal cortex (VTC) have been identified and characterized that respond selectively to different high-level visual categories (e.g., faces: Kanwisher et al., 1997; places: Epstein & Kanwisher, 1998; bodies: Downing et al., 2001; for review, see Kanwisher, 2010). What are the origins of these specialized regions? How do human brains develop this functional topography? Some have hypothesized that the functional organization of the VTC may be innate and related to the evolutionary importance of certain high-level visual categories. Indeed, face perception and recognition abilities appear to be heritable (Wilmer et al., 2010; Zhu et al., 2010), and 4-6 month-old infants already show face- and place-responsive areas within expected locations in the VTC (Deen et al., 2017). A related hypothesis is that pre-existing biases for certain visual attributes (e.g., retinotopy, Hasson et al., 2002; Malach et al., 2002; rectilinearity, Nasr et al., 2014) or perceptual dimensions (e.g., real-world size and animacy, Konkle & Caramazza, 2013) may predispose a brain region to become selective for particular visual categories. However, evolutionary pressures cannot explain the existence of a brain region that specializes for orthography (Hannagan et al., 2015)—the visual word form area (VWFA). The VWFA is a small region in the left lateral VTC that shows strong selectivity for visual words and letter strings in literate individuals (e.g., L. Cohen et al., 2003; Dehaene & Cohen, 2011; Hamamé et al., 2013). Perhaps surprisingly—given its late emergence—the VWFA is located in approximately the same location across individuals and languages (Baker et al., 2007). What sets word forms apart from other visual categories and why does the VWFA land in this stereotyped location?

One compelling possibility is that the specialization of category-selective regions in the VTC is constrained by their differential connectivity to the rest of the brain (the Connectivity Hypothesis; Mahon & Caramazza, 2011; Martin, 2006). Indeed, previous work showed that category-selective responses can be predicted from both structural and functional connectivity (Osher et al., 2015; Saygin et al., 2012); further, these distinct connectivity patterns may already exist at birth and may drive future functional specialization (e.g., newborns show functional connectivity differences between lateral VTC which houses the VWFA vs. medial VTC: Barttfeld et al., 2018). Thus, written words may be processed in a stereotyped region within the left VTC due to this region's pre-existing connectivity with the left-lateralized language network (e.g., Behrmann & Plaut, 2013; Dehaene et al., 2015; Martin, 2006). This network consists of left lateral frontal and lateral temporal areas and selectively supports language comprehension and production (e.g., Fedorenko et al., 2011).

Consistent with this idea, a number of studies have reported both anatomical and functional connections between the VWFA and the language network in neurotypical adults. For example, compared to the adjacent FFA, the VWFA shows stronger anatomical connectivity to the left superior temporal, anterior temporal, and inferior frontal areas (perisylvian putative language regions) (Bouhali et al., 2014). Several candidate white matter fascicles may serve to connect the VWFA with frontal and superior temporal language cortex (Wandell et al., 2012; Yeatman & Feldman, 2013). Similarly, using resting-state functional connectivity, Stevens et al. (2017) found that the individually defined VWFA connects to the posterior left inferior frontal gyrus as well as the left planum temporale, both part of the distributed left language network.

102
 103 Moreover, a longitudinal study in children showed that the location of the VWFA could be
 104 successfully predicted by its connectivity patterns at the pre-reading stage (before any word
 105 selectivity within the VTC is observed), and that the most predictive connections of the relevant
 106 area of the VTC were with the frontal and temporal language areas (Saygin et al., 2016). Even
 107 more strikingly, this pattern of preferential connectivity appears to be already present in neonates
 108 (Li et al., 2020). These connections between the VWFA and the frontotemporal language
 109 network appear to be functionally important such that their damage or stimulation leads to
 110 reading difficulties. For example, a case study of a child with a missing arcuate fasciculus (AF,
 111 presumably connecting the VTC and other parts of the temporal cortex to parietal and frontal
 112 areas; Wandell et al., 2012) found impaired reading ability (Rauschecker et al., 2009). Similarly,
 113 a lesion in the white matter ventral to the angular gyrus resulted in alexia without agraphia,
 114 presumably disrupting the VWFA's connections with the lateral temporal language areas
 115 (Greenblatt, 1976).

116
 117 A different way to assess the importance of a language network in developing visual word
 118 selectivity is to ask whether language regions and the VWFA occupy the same hemisphere. In
 119 the majority of individuals, the language regions and the VWFA co-lateralize to the left
 120 hemisphere (LH) (e.g., Cai et al., 2010; Gerrits et al., 2019). In rare instances where neurotypical
 121 individuals show right-hemispheric (RH) language dominance, VTC activation during reading
 122 tasks also tends to be right-lateralized (e.g., Cai et al., 2008; Van der Haegen et al., 2012).
 123 Another population where the language network is right-lateralized are individuals with
 124 congenital or early left hemisphere (LH) damage (e.g., Asaridou et al., 2020). In the presence of

early LH damage, linguistic abilities tend to develop normally (e.g., Newport et al., 2017; Staudt, 2010; Staudt et al., 2001; see François et al., 2021 for a review). However, little is known about the effects of early LH damage on reading ability and on the neural architecture of visual word processing. In particular, if the left VTC is completely deafferented from the downstream LH language cortex at birth, does the VWFA emerge in the right VTC when the language network is forcibly remapped to the right hemisphere, or does it still emerge in the left VTC, due to some pre-existing bias (e.g., innate connectivity with any spared LH cortex)? Indeed, some previous studies show that language processing and reading may engage opposite hemispheres (e.g., Van der Haegen et al., 2012). Or—perhaps even more drastically—does the VWFA look atypical (e.g., smaller or less functionally selective), whatever hemisphere it emerges in? Do these potential differences affect selectivity for other high-level visual categories in the VTC? And if the VWFA manifests atypically but reading ability is normal, what does this tell us about the implementation-level flexibility with respect to orthographic processing?

Here we investigate possible functional reorganization of the visual word selectivity in the absence of a typical left-lateralized language network. We have a unique opportunity to examine fMRI responses to stimuli from different visual categories in an individual (EG) born without the left superior temporal lobe (likely due to pre/perinatal stroke) but with IVTC largely intact. EG's frontotemporal language network is completely remapped to the right hemisphere; no language-related responses, as assessed with fMRI, were observed in the remaining parts of EG's left hemisphere (Tuckute et al., 2021). EG's reading abilities (as well as other linguistic abilities) are intact. We here investigated a) whether, in the presence of a right-lateralized language network, a typical VWFA would emerge in the right VTC (in a typical location, or perhaps in other parts of

the right VTC), and b) whether any word-selectivity is observed in the (spared) left VTC. To foreshadow the results, no word selectivity was observed in EG's right or left VTC, in the presence of typical selectivity for other visual categories. We then explored the possibility that orthographic processing is supported by still selective but more distributed neural populations using multivariate analyses, and indeed observed such multivariate selectivity bilaterally, but manifesting more strongly in the right VTC.

Methods

Participants

Critical participant. The participant EG (fake initials; right-handed female with an advanced professional degree, 54 year-old at the time of testing) contacted Dr. Fedorenko's lab to participate in brain research studies. Based on her own report, the lack of the left superior temporal lobe (**Figure 1**) was discovered when she was 25 years old (in her first MRI scan in 1987) and being treated for depression. No known head traumas or injuries were reported as a child or adult. Several medical MRI scans were performed in subsequent years (1988, 1998, and 2013) and no change was observed compared to the initial scans. Importantly, EG did not report any difficulties in reading or general language abilities (see details below). She had also acquired fluency in a second language (Russian). EG was invited to participate in a series of behavioral and fMRI assessments at MIT in October 2016. Written informed consent was obtained from EG, and the study was approved by MIT's Committee on the Use of Humans as Experimental Subjects (COUHES).

To formally evaluate EG's linguistic abilities, five standardized language assessment tasks were administered: i) an electronic version of the Peabody Picture Vocabulary Test (PPVT-IV) (Dunn & Dunn, 2007); ii) an electronic version of the Test for Reception of Grammar (TROG-2) (Bishop, 2003); iii) the Western Aphasia Battery-Revised (WAB-R) (Kertesz, 2006); iv) the reading and spelling components of PALPA (Kay et al., 1992); and v) an electronic version of the verbal components of the Kaufman Brief Intelligence Test (KBIT-2) (Kaufman & Kaufman, 2004). PPVT- IV and TROG-2 target receptive vocabulary and grammar, respectively. In these tasks, the participant is shown sets of four pictures accompanied by a word (PPVT-IV, 72 trials) or sentence (TROG-2, 80 trials) and has to choose the picture that corresponds to the word/sentence by clicking on it. WAB-R (Kertesz, 2006) is a more general language assessment developed for persons with aphasia. It consists of 9 subscales, assessing 1) spontaneous speech, 2) auditory verbal comprehension, 3) repetition, 4) naming and word finding, 5) reading, 6) writing, 7) apraxia, 8) construction, visuospatial, and calculation tasks, and 9) supplementary writing and reading tasks. The verbal components of KBIT-2 include 1) the Verbal Knowledge subtest, which consists of 60 items measuring receptive vocabulary and general information about the world, and 2) the Riddles subtest consists of 48 items measuring verbal comprehension, reasoning, and vocabulary knowledge. Most relevant to the current investigation, the reading component of WAB-R includes comprehension of written sentences and reading commands; the supplementary reading tasks include reading of irregular words and nonwords. The reading component of PALPA (Kay et al., 1992) consists of 21 tasks (tasks #18 through 38), which focus on single letter recognition, visual lexical decision, out-loud reading (of words with diverse lexical properties and sentences), and homophone definition (see Supplementary Table 2 for details).

193

194 In line with her self-report, EG performed within normal range on all language assessment tasks.

195 She got 90% correct on PPVT, 99% correct on TROG-2, 97.6, 98.6, and 98.4 on the aphasia,

196 language, and cortical quotients of the WAR-B (the criterion cut-off score for diagnosis of

197 aphasia is an aphasia quotient of 93.8), and a score of 130 (98th percentile) on the KBIT-verbal.

198 EG's performance was therefore not distinguishable from the performance of neurotypical

199 controls. With respect to the reading assessments, EG made no errors on the main reading

200 component of WAB-R, no errors in the reading of irregular words, and one error in the reading

201 of nonwords (reading 'gobter' instead of 'globter'). For the PALPA tasks, she made no errors on

202 tasks that focus on single letters (tasks #18-23), no errors on the visual decision tasks (tasks #24-

203 27), no errors on the out-loud reading tasks (tasks #29-37), and no errors on the homophone

204 definition task (task #38). For the homophone decision task (task #28), EG made three errors

205 (out of the 60 trials; all were made on nonword pairs: she did not judge the following pairs as

206 sounding the same: heem-heam, byme-bime, and phex-feks). Based on this comprehensive

207 assessment, EG appears to have intact visual letter and word recognition ability.

208

209 With respect to fMRI tasks that relevant to the current study, EG completed five runs of the

210 VWFA localizer; see *The VWFA localizer task* section below).

211

212 *Neurotypical controls.* Twenty-five adults (11 female, mean age = 23.6 years old; age range 18-

213 38 years; standard deviation 5.21 years) from The Ohio State University (OSU) and the

214 surrounding community were included in the present study. As part of ongoing projects

215 exploring the relationship between brain function and connectivity, all participants completed a

battery of fMRI tasks, including, critically, the same VWFA localizer task that EG completed (see *The VWFA localizer task* section below). All participants had normal or corrected-to-normal vision, and reported no neurological, neuropsychological, or developmental diagnoses. Written informed consent was obtained from all participants and the study was approved by Institutional Review Board at OSU. (It is worth noting that although the control group participants were younger than EG, further examination revealed no significant correlation between age and word-selectivity (responses to Words minus the average of the other conditions) in our control group ($r=-0.259$, $p=0.094$; p -value was obtained by a permutation test).

Data acquisition

EG's data were collected on a 3 Tesla Siemens Trio scanner with a 32-channel head coil at the Athinoula A. Martinos Imaging Center at the McGovern Institute for Brain Research at MIT. Data of the control group participants were acquired on a 3 Tesla Siemens Prisma scanner with a 32-channel head coil at the Center for Cognitive and Behavioral Brain Imaging at OSU. To ensure that any differences between EG and our control group were not due to scanner differences, we compared word-selectivity in the current control group to a smaller group of adults recruited at MIT and scanned with the same scanner and protocols as EG; no differences were observed (Supplementary Table 1). For both EG and controls, a whole-head, high resolution T1-weighted magnetization-prepared rapid acquisition with gradient echo (MPRAGE) scan was acquired (EG: repetition time (TR) = 2530 ms, echo time (TE) = 3.48 ms; voxel resolution = 1.0 mm^3 ; the control group: TR = 1390 ms, TE = 4.62 ms, voxel resolution = 1.0 mm^3). Functional images were acquired with the same echo-planar imaging (EPI) sequence for both EG and controls: TR=2000ms, TE=30ms, 172 TRs, 100×100 base resolution, voxel

resolution=2.0 mm³, 25 slices approximately parallel to the base of the temporal lobe to cover the entire ventral temporal cortex.

The VWFA localizer task

A VWFA localizer was used to define high-level category-selective regions and to measure category-selective responses (see Saygin et al., 2016, for details). Briefly, black and white line drawings of words, scrambled words, objects, and faces, along with the fixation condition were shown in a blocked design. A grid was overlaid on top of the stimuli so that all stimulus types (not just scrambled words) had edges. Each stimulus was presented for 500ms (ISI=0.193s) and overlaid on a different single-color background, and 26 stimuli (including 2 repetitions) were presented in each block. Each run consisted of 19 blocks (4 blocks per stimulus type and 3 fixation blocks), and participants performed a one-back task. The stimuli are available for download at <http://www.zevnepsaygin.com/ZlabResources.html>. EG completed 5 runs, and participants in the control group completed 2 runs. Note that previous studies using the same task indicated that 2 runs of data are sufficient to successfully identify the VWFA in a neurotypical population (Saygin et al., 2016); here, we acquired more runs for EG to ensure that we had sufficient power and that the results obtained for EG were stable across runs (see *Definition of functional regions of interest and univariate analyses* section for details).

Preprocessing and fMRI analysis

Data were analyzed with Freesurfer v.6.0.0 (<http://surfer.nmr.mgh.harvard.edu/>), FsFast (<https://surfer.nmr.mgh.harvard.edu/fswiki/FsFast>), FSL (<https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FLIRT>) and custom MatLab code. All structural MRI data

were processed using a semiautomated processing stream with default parameters (recon-all function in Freesurfer: <https://surfer.nmr.mgh.harvard.edu/fswiki/recon-all/>), which includes the following major steps: intensity correction, skull strip, surface co-registration, spatial smoothing, white matter and subcortical segmentation, and cortical parcellation. Cortical gray matter and ventral temporal cortex masks were created based on the Desikan-Killiany (Desikan et al., 2006) parcellation in native anatomy for each subject. The success of cortical reconstruction and segmentation was visually inspected for EG.

Functional images were motion-corrected (time points where the difference in total vector motion from the previous time point exceeded 1mm were excluded), data from each run were registered to each individual's anatomical brain image using bbrregister, and resampled to $1.0 \times 1.0 \times 1.0 \text{ mm}^3$. For EG, instead of registering the functional image of each run to the anatomical brain separately, we aligned the functional images of the first four runs to the last run (which had successful functional-anatomical cross-modal registration with bbrregister) with linear affine transformation (FLIRT); then the functional-anatomical transformation for the last run was applied to all functional runs and was visually inspected (tkregisterfv) for functional to anatomical alignment.

The preprocessed functional data were then entered into a first-level analysis. Specifically, data were detrended, smoothed (3mm FWHM kernel), and the regressor for each experimental condition (Words, Scrambled Words, Objects, and Faces) was defined as a boxcar function (i.e., events on/off) convolved with a canonical hemodynamic response function (a standard gamma function ($d = 2.25$ and $t = 1.25$)). Orthogonalized motion measures from the preprocessing stage

were used as nuisance regressors for the GLM. Resulting beta estimates for each condition and contrasts of interest between categories (i.e., Words > Others other categories and Faces > Others other categories) were used in further analyses. For multivariate analyses, no spatial smoothing was applied.

Definition of functional regions of interest and univariate analyses

The subject-specific functional regions of interests (fROIs) were defined with a group-constrained subject-specific (GcSS) approach (Fedorenko et al., 2010). In this approach, individual fROIs are defined by intersecting participant-specific fMRI contrast maps for the contrast of interest (e.g., Words > Other conditions) with some spatial constraint(s) or ‘parcel(s)’, which denote the area(s) in the brain within which most individuals show responses to the relevant contrast (based on past studies with large numbers of participants). In the present study, the VWFA and the FFA parcels were derived from previous studies (FFA, Julian et al., 2012; VWFA, Saygin et al., 2016) (**Figure 1**) and were generated based on probabilistic maps of functional activation for the relevant contrasts in independent groups of participants. We registered these parcels to our participants’ own anatomy using the combined volume and surface-based (CVS) non-linear registration method (mri_cvs_register; Postelnicu et al., 2009). After mapping the functional parcels to each participant’s brain, we defined the VWFA fROI by selecting within the VWFA parcel the top 10% of most active voxels for the contrast Words > Other conditions (i.e., scrambled words, objects, and faces). Similarly, we defined the FFA by selecting within the FFA parcel the top 10% of most active voxels for the contrast Faces > Other conditions. For the control participants, we used run 1 to define the fROIs, and the run 2 to extract percent signal changes (PSCs; beta estimates divided by baseline activation) for each of

four experimental conditions. For EG, who had 5 runs of data, this procedure was performed iteratively for every 2-run/3-run split (e.g., defining fROIs with runs 1&2 and extracting PSCs from runs 3-5, defining with runs 1&3 and extracting PSCs from runs 2, 4, and 5, etc.; 10 combinations in total), and the results were averaged to derive a single estimate per condition per fROI. Note that although the parcels are relatively large, by design (in order to accommodate inter-individual variation in the precise location of the functional region), and can overlap, the resulting fROIs within an individual are small and do not overlap (Saygin et al., 2016).

Multivariate analyses: Split-half correlations

To further examine whether visual words may be represented and processed in a spatially distributed manner, we performed a multivariate pattern analysis (MVPA) to measure distinctive activation patterns for different conditions. The analyses were performed with CoSMoMVPA toolbox (Oosterhof et al., 2016) (<https://www.cosmomvpa.org/>). In line with the approach introduced in Haxby et al. (2001), we examined split-half within-category and between-category correlations. In particular, a searchlight (radius = 3 voxels) was created for each voxel within the VTC, and response patterns (i.e., beta estimates for each of the four conditions, normalized by subtracting mean responses across all conditions) were extracted from each searchlight.

Before performing the critical analysis, we asked whether the overall multivariate representation structure of the VTC is typical in EG. To do so, we constructed a representational similarity matrix (RSM) from pairwise similarities (i.e., correlations) based on the voxel-wise response patterns in a given searchlight area (defined above) to different conditions (e.g., the correlation between activation patterns across voxels to Words and Faces in a given searchlight area).

Correlations of all searchlights within the VTC were then Fisher z-transformed and averaged. This resulted in one 4x4 RSM from two runs of data for each participant in the control group; for EG, RSMs from ten run combinations were averaged to get a single RSM. We then calculated RSM similarity (correlation) between EG and controls, and tested whether this correlation was different from the correlations between any given control individual and the rest of the control group (within-controls correlations; similarity of RSM of each subject to the average RSM of the remaining control subjects).

Then, similarity of response patterns within a category (e.g., Words-Words or Faces-Faces) vs. between categories (e.g., Words-Faces) was calculated within the VWFA and FFA parcel boundaries by Pearson's correlation in a pair-wise manner, and then Fisher z-transformed. We identified voxels of interest that satisfied the following two criteria: 1) voxels whose local neighborhoods showed higher within- than between-category correlations (and which therefore represent categories distinctively); and 2) voxels whose local neighborhoods showed higher within-condition correlations for a particular category (e.g., Words-Words) than within-condition correlations for other categories (e.g., Faces-Faces). The second criterion identified voxels which represent a particular category (e.g., visual words) in a more selective fashion. We refer to such voxels as multivariate-selective voxels.

To examine hemispheric differences, we computed the number of voxels that exhibited multivariate selectivity for words or faces in the two hemispheres (within the relevant parcels). To control for the difference in the size of the search spaces (i.e., the parcels), we divided the number of the multivariate-selective voxels by the total size of the relevant parcel.

354

355 *Statistical analyses*

356 Paired t-tests were used for comparisons between conditions for EG (across ten run split
357 combinations) and within the control group. For all analyses where we compared EG's response
358 to the control group, we used a Crawford-Howell modified t-test (Crawford & Howell, 1998),
359 which is widely used in single-case studies because it accounts for the variance of the control
360 group, and the percentage of false positives remains the same for various sizes of the control
361 sample (N=5-50) (Crawford et al., 2009). This frequentist approach provided the point estimate
362 (p-value) for the proportion of the control population that will obtain a score lower than the
363 patient's score. One-tailed tests were reported for the critical (directional) hypothesis tested in
364 the current study (unless noted otherwise). In addition, we computed the Bayesian 95% credible
365 interval (a Bayesian alternative to the modified t-test; Crawford & Garthwaite, 2007) to
366 demonstrate the range of p-values based on 10,000 iterations. To assess the significance of RSM
367 correlations between EG and controls, we generated a null distribution of correlation values by
368 shuffling the matrix of EG and controls (i.e., randomizing the labels of values in the RSMs) and
369 then correlating the new shuffled matrices. This procedure was repeated 10,000 times to create
370 the null distribution of the correlation values. The p-value was calculated by counting the
371 number of correlations in the null-distribution that were higher than the correlation value based
372 on the correct category labels, and then divided by 10,000.

373

374 **Results**

375 *Is word selectivity remapped to the right hemisphere when the language network is remapped to*
376 *the right hemisphere from birth?*

We first examined univariate word selectivity in EG's rVWFA. With the VWFA localizer, we defined the individual-specific rVWFA fROIs, in EG and controls, within a spatial constraint (rVWFA parcel; see **Figure 1** and Methods) by contrasting Words versus all other categories (Words > Others). We then extracted the activation to all four conditions from independent data (see Methods for details). In neurotypical literate individuals, word selectivity is strongly left-lateralized (McCandliss et al., 2003); selective responses to visual words in the right homotop of lVWFA are less frequently observed and are less spatially consistent across individuals. Results from our control group are in line with this picture: we found no selective activation to Words as compared to other visual categories in the rVWFA (Words vs. Scrambled Words: $t(24)=-0.964$, $p=0.345$; Words vs. Objects: $t(24)=-0.780$, $p=0.443$; Words vs. Faces: $t(24)=-0.523$, $p=0.605$; **Figure 2A**).

391

392 Figure 1. EG's MRI showing the missing cortex and the parcels for the right and left VWFA and FFA.
 393 Top, T1-weighted images in coronal, sagittal, and axial views. Bottom, the VWFA (purple) and FFA
 394 (blue) parcels are projected on EG's reconstructed surface. The parcels served as spatial constraints in
 395 defining the fROIs (see Methods), but we also explored the entire VTC for category selectivity (outlined
 396 with black solid lines). By design, the parcels are relatively large (to accommodate inter-individual
 397 variability in the precise locations of these areas) and therefore can overlap, but the individual fROIs are
 398 small and do not overlap. Note that even though part of the anterior IVTC is missing in EG, the
 399 stereotypical locations for both the VWFA and FFA are spared.

401

402 Critically, in EG, whose language network is completely reorganized to the right hemisphere
 403 (Tuckute et al., 2021), we asked whether word selectivity was also remapped to the right
 404 hemisphere. Surprisingly, no word selectivity was observed in EG's rVWFA: activation to
 405 Words in her rVWFA fROI did not significantly differ from those to Scrambled Words and
 406 Faces (Words vs. Scrambled Words: $t(9)=-0.913$, $p=0.385$; Words vs. Faces: $t(9)=-0.966$, $p=$
 407 0.359) and was significantly *lower* than Objects (Words vs. Objects: $t(9)=-2.346$, $p=0.044$)
 408 (**Figure 2B**). To ensure that this result is not due to the choice of a particular threshold (i.e., top
 409 10%) used to define the rVWFA fROI, we performed the same analysis as above at a range of
 410 thresholds (5%-50%). The lack of word selectivity was stable across thresholds (**Figure 2C**). We
 411 then compared EG to the controls, and found that there was a trend whereby responses to Words
 412 (relative to baseline) were slightly higher in EG compared to the control group ($t(24)=0.503$, $p=$
 413 0.310 , modified t-test; 95% Bayesian CI [0.17, 0.45]), but this difference was not significant.

414

415 To test whether selectivity for other visual categories in the right VTC is typical in EG, we
 416 examined face selectivity in EG's rFFA, which is spatially proximal to the rVWFA. Similar to
 417 the rVWFA analysis above, we defined the rFFA fROI by contrasting Faces versus other
 418 categories (Faces > Others), and extracted the activation to the four conditions from independent

data (see Methods for details). EG's face selectivity remained intact (**Figure 2E**) and did not differ from that of the control group (**Figure 2D**; $t(24)=-0.031$, $p=0.488$, modified t-test, Bayesian 95% CI of EG's activation [0.35 0.50]). EG's rFFA fROI showed significantly higher responses to Faces than to other conditions (Faces vs. Words: $t(9)=24.813$, $p=1.346\times 10^{-9}$; Faces vs. Scrambled Words: $t(9)=16.850$, $p=4.092\times 10^{-8}$; Faces vs. Objects: $t(9)=20.312$, $p=7.921\times 10^{-9}$; **Figure 2E**), just like what we observed in the control group (Faces vs. Words: $t(24)=15.299$, $p=7.054\times 10^{-14}$; Faces vs. Scrambled Words: $t(24)=12.915$, $p=2.689\times 10^{-12}$; Faces vs. Objects: $t(24)=10.724$, $p=1.235\times 10^{-10}$; **Figure 2D**). Moreover, the face selectivity in EG's rFFA was observed across all thresholds used to define the rFFA (**Figure 2F**).

Figure 2. Responses to four conditions in the rVWFA and rFFA for EG and the control group. (A) Bar plots show mean PSCs to the four conditions estimated in independent data within individually defined rVWFA fROIs (i.e., top 10% word-selective voxels within the rVWFA parcel) for the control group. (B) Mean PSCs to the four conditions estimated in independent data within the individually defined rVWFA fROI for EG. Here and in E, the results are averaged across run combinations. (C) Parametrically decreasing the threshold for defining the rVWFA fROI from the top 5% to 50% word-selective voxels within the rVWFA parcel. Number of voxels in the rVWFA fROIs: 5%=384 voxels, 50%=3007 voxels. Here and in F, average PSCs across run combinations are shown for each threshold. (D) Mean PSCs to the four conditions estimated in independent data within individually defined rFFA fROIs for the control group. (E) Mean PSCs to the four conditions estimated in independent data within the individually defined rFFA fROI for EG. (F) Parametrically decreasing the threshold for defining the rFFA fROI from the top 5% to 50% face-selective voxels within the rFFA parcel. Number of voxels in the defined rFFA fROIs: 5%=90 voxels, 50%=699 voxels. In the bar plots, dots correspond to individual data points for each condition (controls: n=25 participants; EG: n=10 run combinations, from ten iterations). Horizontal bars reflect significant paired t-tests $p < 0.05$. Error bars in both the bar and line plots denote standard errors of the mean by participants (for the control group) and by run combinations (for EG).

To ensure that we did not miss any possible word-selective voxels by applying a predefined spatial constraint (i.e., VWFA parcel) and to account for the possibility that EG's VWFA is located in a different part of the visual cortex, we searched for word selectivity within the entire rVTC mask for EG. Specifically, different thresholds from top 1% to top 10% were used to define the most word-selective voxels (Words > Others). Even within this broad mask, no word-selective responses were observed in independent data across all thresholds; in fact, the responses were lowest to words than the other three conditions (**Figure 3A**). In contrast, robust face-selective responses were observed in independent data across all thresholds when searching for face-selective voxels (Faces > Others) (**Figure 3B**).

Finally, we supplemented our main analyses, which rely on the Words > Others contrast to define the VWFA, as is commonly done in the literature (e.g., Dehaene-Lambertz et al., 2018; Rosenke et al., 2021), with an analysis, which relies on a less stringent localizer contrast (Words vs. Scrambled Words; Glezer et al., 2009; Lerma-Usabiaga et al., 2018). Unlike the Words >

Others contrast, this contrast does not control for semantics or visual stimulus complexity. Even with this broader contrast, we found no word selectivity within EG's rVTC across thresholds (Supplementary Figure 1).

Figure 3. Mean PSCs in the rVTC and IVTC at different thresholds for EG. (A-B) Parametrically decreasing the threshold for defining word-selective (Words > Others) and face-selective (Faces > Others) voxels from the top 1% to 10% within the rVTC. Mean PSCs across run combinations (from 10 iterations) are shown for each threshold. (C-D) Parametrically decreasing the threshold for defining word-selective and face-selective voxels from the top 1% to 10% within the IVTC. Mean PSCs across run combinations are shown for each threshold. Number of selected voxels: rVTC: 1%=216 voxels, 10%=2164 voxels; IVTC: 1%=154 voxels, 10%=1536 voxels.

Is there any word selectivity in the spared left VTC?

Because we did not observe a right-lateralized VWFA and because the left VTC was largely intact, we also asked whether a canonical IVWFA may have developed in EG (perhaps due to some specific visual features of word forms that are better represented in the left hemisphere/right visual field (e.g., Hsiao & Lam, 2013; Seghier & Price, 2011; Tadros et al., 2013). In the control group, we observed robust word selectivity in the IVWFA (as expected from previous studies), with Words eliciting greater activation than each of the other conditions: Words vs. Scrambled Words: $t(24)=5.413$, $p=1.467\times 10^{-5}$; Words vs. Objects: $t(24)=4.077$, $p=4.333\times 10^{-4}$; Words vs. Faces: $t(24)=5.942$, $p=3.935\times 10^{-6}$; **Figure 4A**). In contrast, we found no word selectivity in EG's IVWFA (**Figure 4B**): activation to Words was around baseline and significantly lower than those to Scrambled Words and Objects (Words vs. Scrambled Words: $t(9)=-3.092$, $p=1.289\times 10^{-2}$; Words vs. Objects: $t(9)=-2.432$, $p=3.785\times 10^{-2}$) and did not significantly differ from that to Faces (Words vs. Faces: $t(9)=-0.699$, $p=0.503$). Moreover, EG's activation to Words was significantly lower than the control group's ($t(24)=-1.616$, $p=5.961\times 10^{-2}$, modified t-test; 95% Bayesian credible interval (CI) [0.01, 0.13]). This result was stable across thresholds that were used to define the IVWFA fROI (**Figure 4C**), and we did not observe word selectivity in EG when we searched across the entire IVTC with varying thresholds (**Figure 3C**), or when we used a less stringent contrast (Words vs. Scrambled Words; Supplementary Figure 1).

Figure 4. Responses to four conditions in the IVWFA and IFFA for EG and the controls group. (A), Bar plots show mean PSCs to four conditions estimated in independent data within individually defined IVWFA fROIs (i.e., top 10% word-selective voxels within IVWFA parcel) for the control group. (B), Mean PSCs to the four conditions estimated in independent data within individually defined IVWFA fROIs for EG. Here and in E, the results are averaged across run combinations.. (C), Parametrically decreasing the threshold for defining the IVWFA fROIs from the top 5% to 50% word-selective voxels within the IVWFA parcel. Number of voxels in the defined IVWFA fROIs: 5%=295 voxels, 50%=2402 voxels. Here and in F, average PSCs across run combinations are shown for each threshold. (D), Mean PSCs across participants to the four conditions estimated in independent data within individually defined IFFA fROIs for the control group. (E), Mean PSCs to four conditions estimated in independent data within individually defined IFFA fROIs for EG. (F), Parametrically decreasing the threshold for defining IFFA fROI from the top 5% to 50% face-selective voxels within the IFFA parcel. Number of voxels in the defined IFFA fROIs: 5%=18 voxels, 50%=185 voxels. In the bar plots, dots correspond to individual data points (controls: n=25 subjects; EG: n=10 run combinations, from ten iterations). Horizontal bars reflect significant paired t-tests $p < 0.05$. Error bars in both the bar and line plots denote standard errors of the mean by participants (for the control group) and by run combinations (for EG).

Similar to what we did for the rVTC, to test whether selectivity for other visual categories in the left VTC is typical in EG, we examined face selectivity in EG's IFFA. EG's face selectivity remained intact (**Figure 4E**) and did not differ from that of the control group (**Figure 4D**; $t(24)=-$

0.275, $p = 0.393$, modified t-test; 95% Bayesian CI, [0.27 0.50]). EG's IFFA fROI showed significantly higher responses to Faces than to other conditions (Faces vs. Words: $t(9) = 47.571$, $p = 4.015 \times 10^{-12}$; Faces vs. Scrambled Words: $t(9) = 24.681$, 1.411×10^{-9} ; Faces vs. Objects: $t(9) = 34.494$, $p = 7.141 \times 10^{-11}$; **Figure 4E**), just like what we observed in the control group (Faces vs. Words: $t(24) = 7.076$, $p = 1.467 \times 10^{-5}$; Faces vs. Scrambled Words: $t(24) = 8.102$, $p = 2.518 \times 10^{-8}$; Faces vs. Objects: $t(24) = 5.517$, $p = 1.130 \times 10^{-5}$; **Figure 4D**). As was the case for the rFFA, the face selectivity in EG's IFFA was observed across all thresholds used to define the IFFA (**Figure 4F**).

Altogether, the examination of EG's right and left VTC suggests that without the typical left-hemisphere frontotemporal language network from birth—and presumably without the necessary connections between these areas and parts of the VTC—a canonical VWFA, a word-selective area, does not develop in either hemisphere.

Distributed neural representation of visual words: multivariate pattern analysis (MVPA)

Our univariate analyses measured the average activation across the voxels most responsive to the visual category in question. Although this is the classic approach to demonstrate category selectivity, univariate analyses may be insensitive to potentially meaningful distributed representation patterns in suprathreshold and/or subthreshold voxels. Previous studies have found that categorical information can be reliably decoded by comparing within-category versus between-category correlations in the VTC (e.g., Haxby et al., 2001). Moreover, a recent study

found mature representational similarity structures via multivariate patterns in children with no univariate selectivity, suggesting that distributed representations may developmentally precede category selectivity (Cohen et al., 2019). To explore whether words may be represented in a distributed fashion in EG, we performed a series of multivariate pattern analyses (MVPA).

We first examined the representational similarity matrices (RSMs) in the entire VTC (see Methods) to investigate whether multivariate representational structure for visual categories was preserved in EG. Indeed, we found that EG's rVTC RSM was strongly and significantly correlated with that of the control group ($r=0.918$, $p=9.100 \times 10^{-3}$ (permutation test); Supplementary Figure 3). We tested whether this correlation between EG and the control group was different from the correlations between any given control individual and the rest of the control group (see Methods). Single case comparisons showed that the RSM correlation for EG vs. controls did not significantly differ from the within-controls correlations ($t(24)=0.797$, $p=0.217$, Bayesian 95% CI [0.09 0.351]). Similar results were found for the IVTC: the RSMs of EG and the control group were strongly and significantly correlated ($r=0.888$, $p=7.300 \times 10^{-3}$), and the correlation for EG vs. controls did not significantly differ from the within-controls correlations ($t(24)=0.835$, $p=0.206$, Bayesian 95% CI [0.09 0.34]).

We then asked whether EG's 'VWFA' may contain voxels that show distinct distributed activation patterns to visual words. Specifically, as described in Methods, we searched for voxels that satisfied the following criteria: the searchlight around a given voxel should show 1) distinctive response patterns to Words vs. other categories (e.g., the Words-Words correlation should be higher than the Words-Faces correlation); and 2) stronger within-category correlations

for the preferred category (e.g., the Words-Words correlation should be higher than the Faces-Faces, Objects-Objects, and Scrambled Words-Scrambled Words correlations). Previous studies have shown distributed representations within category-selective regions (e.g., the FFA) of non-preferred categories (e.g., places), and debate is ongoing over whether this information has functional relevance (e.g., Kanwisher, 2010). Our second criterion was included to identify voxels that show more stable multivariate representations for the category of interest (e.g., Words) compared to other categories. Indeed, we identified a set of voxels within the rVWFA and lVWFA parcels that showed a reliable distributed code for words in both controls and EG (Supplementary Figure 2A, 2C; Supplementary Table 3). In addition, mirroring the univariate analyses, we also identified a set of voxels that showed a reliable distributed code for faces within the rFFA and lFFA parcels in both controls and EG (Supplementary Figure 2B, 2D; Supplementary Table 3).

Critically, to test whether these distributed responses to visual words differ between EG and the controls, we subtracted the average within-category correlations for all non-word categories from the within-category correlations for Words. This difference tells us how much stronger the within-category correlation is for the preferred compared to the non-preferred categories. EG showed comparable within-category correlation differences to the controls in both the rVWFA ($t(24)=1.081$, $p=0.145$, modified t-test, 95% Bayesian CI [0.05, 0.26] and lVWFA ($t(24)=0.184$, $p=0.428$, modified t-test, 95% Bayesian CI [0.30, 0.50]); **Figure 5A**). Similar within-category correlation differences were also observed between EG and the controls in the rFFA ($t(24)=-0.802$, $p=0.215$; 95% Bayesian CI [0.09, 0.34]) and lFFA ($t(24)=0.488$, $p=0.315$; 95% Bayesian CI [0.18, 0.46]); **Figure 5B**).

Figure 5. Results of the multivariate pattern analysis (MVPA). (A) Within-category correlation differences between preferred (i.e., Words-Words) and non-preferred (i.e., average within-category correlation of non-word conditions) conditions for EG and the controls in the rVWFA and lVWFA parcels. Here and in B, correlation values were Fisher's z-transformed. (B) Within-category correlation differences between preferred (i.e., Faces-Faces) and non-preferred (i.e., average within-category correlation of non-face conditions) conditions for EG and the controls in the rFFA and lFFA parcels. (C) Proportion of voxels that show multivariate selectivity for Words in the rVWFA and lVWFA parcels for EG and the controls. (D) Proportion of voxels that show multivariate selectivity for Faces in the rFFA and lFFA parcel for EG and the controls. In the bar plots, dots correspond to individual data points (controls: n=25 subjects; EG: n=10 run combinations from ten iterations). Horizontal bars reflect significant LH-RH differences, $p < 0.05$. Error bars denote standard errors of the mean by participants (for the control group) and by run combinations (for EG).

Finally, we explored potential hemispheric asymmetries with respect to multivariate selectivity for words and faces. Left-hemisphere dominance for words and right-hemisphere dominance for faces are well established with univariate measures. Do multivariate patterns also show these asymmetries? We calculated the number of voxels that show multivariate selectivity for words and faces within the left and right VWFA and FFA parcels, respectively (controlling for the size of the search space; see Methods). For Words, mirroring the univariate results in the past literature, a numerically larger proportion of voxels showing multivariate word selectivity was found in the left compared to the right VWFA in the controls (although the difference did not reach significance: $t(24)=1.181$, $p=0.249$). Interestingly, the opposite pattern was observed in EG, with a larger proportion of voxels showing multivariate word selectivity found in the right VWFA ($t(9)= -2.469$, $p= 3.564 \times 10^{-2}$) (**Figure 5C**). For Faces, a numerically larger proportion of voxels showing multivariate face selectivity was found in the right compared to the left FFA in the controls (although, like for Words, the difference did not reach significance: $t(24)=0.367$, $p=0.717$), and EG displayed the opposite pattern: a significantly larger proportion of voxels showing multivariate face selectivity in the left FFA ($t(9)=8.983$, $p=8.672 \times 10^{-6}$) (**Figure 5D**).

Discussion

Case studies provide valuable insights for understanding the functional organization of the human brain and the patterns of reorganization that follow neurological damage (Vaidya et al., 2019). In the current study, we had a unique opportunity to collect fMRI data from an individual (EG) born without her left superior temporal lobe in order to ask whether category-selective

responses to visual words within the ventral temporal cortex would be affected. Specifically, we asked: in the absence of a typical left-hemisphere language network, does word selectivity emerge, and if so, where? Does the VWFA emerge in the right VTC given that EG's language network is reorganized to the right hemisphere (Tuckute et al., 2021)? Or do we instead, or additionally, observe word selectivity in the canonical VWFA within the spared IVTC? Surprisingly, we found no univariate word selectivity within the rVWFA, the right homotope of the canonical IVWFA, and even in an expanded search space of the entire rVTC. We also did not observe word-selective responses in the IVWFA or in any other spared parts of the IVTC. Moreover, the response magnitude to Words (relative to baseline) was significantly lower in EG's IVWFA than that of controls. Importantly, this lack of category selectivity was specific to Words: selective responses to Faces remained intact in EG's rFFA and IFFA.

Absent univariate word selectivity, we then explored multivariate representations of visual words in EG. We found that EG's VTC showed an overall similar representational structure to that of the control group, and that, like the control group, EG had a set of voxels whose local neighborhoods robustly differentiated between Words and other visual categories in their patterns of response. Critically, these voxels also showed a higher within-category correlation (across runs) for Words compared to the within-category correlations for other categories, and the degree of this 'multivariate selectivity' was similar between EG and the controls. Interestingly, however, in contrast to the typically observed left-hemispheric dominance for words and right-hemispheric dominance for faces, EG had a larger proportion of voxels that showed multivariate selectivity for words in her right than her left VWFA, and a larger

proportion of voxels that showed multivariate selectivity for faces in her left FFA than her right FFA.

Altogether, the current study suggests that in the absence of a typical left-lateralized language network—at least in cases of early left-hemispheric damage, when language processing is ‘forced’ to the right hemisphere—neural selectivity for visual words is, or at least can be, highly atypical. In our participant with extensive left-hemispheric temporal damage and a right-lateralized language network, we observed no word selectivity in the univariate analyses in either the right or the spared left VTC. The absence of such selectivity, combined with EG’s intact reading ability, suggests that successful orthographic processing may depend on a more distributed and more right-lateralized neural representation than that observed in typical literate adults.

Canonical univariate word selectivity may not emerge when a left-hemisphere language network is missing from birth

The interesting case of EG allowed us to investigate how visual word processing within the VTC can be affected by a congenital or early left-hemisphere lesion outside of visual cortex. Our results provide the first evidence of atypical visual word selectivity in the VTC when the left-lateralized language cortex is missing, and when the language network consequently develops in the right hemisphere (Tuckute et al., 2021) during the early stages of language learning and prior to learning to read. Our study also suggests that even with some remaining anatomical connections between the spared IVTC and the frontal and temporal areas (presumably via local

U-fibers or remaining long-range fibers), the IVWFA will not emerge at the stereotypical location when the left hemisphere does not support language processing.

How does a left language network contribute to the development of the VWFA? One possibility is that in a typical brain, the site of the putative VWFA is predisposed to written scripts via intrinsic co-activation with speech and/or language/semantic areas in the left temporal (and maybe frontal) cortex. The processing of speech sounds is tightly linked to reading, and impaired coding of speech sounds (e.g., phonological awareness) is often considered a precursor of dyslexia (e.g., Pennington & Bishop, 2008; Shaywitz et al., 2002). In the absence of regions that typically support speech processing (within the left superior and middle temporal gyri (STG and MTG); Raschle et al., 2012), EG's left VTC lacked early interactions with these regions. Further, a typical IVWFA communicates visual orthographic information to higher-level left temporal (and maybe frontal) regions that integrate visual and auditory information, like the amodal language regions that process lexical-semantic and combinatorial linguistic information (e.g., Fedorenko et al., 2020) and the regions that support abstract conceptual processing (e.g., Binder et al., 2009; Patterson et al., 2007)—both sets of areas that were also missing in EG's left hemisphere. Consistent with the idea that connections between the left VTC and the ipsilateral high-level language areas may be critical for the emergence of the IVWFA, a recent study found that in newborns, the putative site of the IVWFA already shows preferential functional connectivity to the areas that will later respond to high-level linguistic information, compared to adjacent regions (Li et al., 2020); this pre-existing functional coupling may further strengthen during language and reading acquisition. Because EG was missing both i) speech-responsive areas, and ii) higher-level language/conceptual areas in her left hemisphere, we cannot evaluate

the relative contributions of these two sets of areas and their connections with the VTC to the emergence of a canonical VWFA. We speculate that both may be important.

Right-hemispheric neural correlates that support visual word processing

Because EG's language network resides in the right hemisphere (Tuckute et al., 2021), we expected to find a VWFA in the right VTC. Surprisingly, no univariate word selectivity was observed in EG's VTC, and this result could not be explained by either a lack of functional specialization *anywhere* in the rVTC (intact face selectivity was observed), or our particular procedure for defining the VWFA (the results held across a range of thresholds, when using a larger search space, and across different contrasts). Interestingly, neurotypical individuals with right-dominant language activation sometimes also lack a VWFA in the right VTC (e.g., Cai et al., 2008; Van der Haegen et al., 2012); in these individuals, IVTC appears to be engaged during word recognition, presumably due to stronger frontotemporal anatomical connections in the LH than in the RH, and any language activation on the left (even if it's non-dominant) would engage the IVTC for reading (Powell et al., 2006). In the case of EG, her right-hemisphere language network may have lacked early interactions with the rVTC, and her left-hemisphere language network was altogether lacking, resulting in the atypical word selectivity (in both right and left) that we observed here.

Some have also argued that the development of word-selective cortex directly competes with face-selective cortex for neural resources, thus contributing to right-hemispheric dominance for face processing and left-hemispheric dominance for word processing in neurotypical individuals (e.g., Behrmann & Plaut, 2015; Dehaene & Cohen, 2007). Interestingly, EG showed typical face

selectivity in the right FFA, not different from controls. It is therefore possible that a focal word-selective area failed to emerge in EG's rVTC because the relevant cortical tissue had already been 'assigned' to face processing, perhaps due to some specific visual features of faces that are better represented in the right hemisphere/left visual field (e.g., holistic face processing; Li et al., 2017; Rossion et al., 2000). Interestingly, although the univariate response magnitude to faces was higher in EG's right than left FFA (similar to controls), more multivariate-selective face voxels were present in EG's IFFA, compared to her rFFA, in sharp contrast to controls. The functional significance of the latter is at present unclear, but can be explored in future work. On the other hand, EG had more multivariate-selective word voxels in her rVWFA than her IVWFA, presumably related to the fact that her language network resides in the right hemisphere. These data suggest that the hemispheric dominance of the language network drives the laterality of visual processing in the VTC (be it implemented focally, or in a distributed fashion), at least for words, but perhaps also for faces (Behrmann & Plaut, 2020; Dehaene & Cohen, 2007).

Multivariate responses to words

Despite the absence of typical univariate word selectivity, EG's VTC was similar in its overall representation similarity structure—across different visual stimulus classes—to that of the control group. Consistent with our results, Liu et al. (2019) also found typical representational structure in category-selective regions after resections within or outside VTC; in addition, a recent study found mature representational similarity structures in children (5-7 year-olds) with no univariate selectivity, suggesting that distributed representations precede category selectivity (M. A. Cohen et al., 2019). Our results provide another case where a typical multivariate representational structure is observed in the absence of univariate selectivity for words.

Moreover, our results showed that EG (like typical adults) displayed distinctive activation patterns within the general vicinity of both right and left VWFA that could distinguish visual words from other conditions, with higher within- than between-category correlations (Haxby et al., 2001) and more robust (consistent over time) representations of Words than other conditions. Supporting the idea that multivariate selectivity for words may be functionally useful, Stevens et al. (2017) found the VWFA (defined in individual participants using a standard univariate contrast) discriminated words from pseudowords, and did so more strongly than other control regions (e.g., the FFA).

Numerous questions remain about the multivariate word representations observed in EG. We will highlight two here. First, what is the relationship between univariate and multivariate representations? Univariate selectivity may partially depend on a connectivity scaffold or other genetically defined instructions to determine the location of functional specialized areas (Deen et al., 2017; Kamps et al., 2020; Li et al., 2020). In contrast, distributed representations like those found in EG may rely less on a connectivity scaffold. Do distributed representations reflect a fundamentally different neural code from that associated with focal representations? Some have argued that the answer is no, and that multivariate analyses may simply be more sensitive than univariate approaches given that they consider the heterogeneity of response across voxels within a region as well as potential subthreshold voxels (Davis et al., 2014). On the other hand, multivariate representations may be truly distinct and reflect information derived from bottom-up visual statistics to a greater extent; and thus perhaps in typical development, representational similarity structures precede univariate selectivity (M. A. Cohen et al., 2019) and emergence of univariate selectivity requires experience-dependent interactions with higher-level areas (e.g., the

language cortex in case of the VWFA). Speculating further, perhaps during EG's early stages of language development (while her language system was maturing in the right hemisphere), the right visual cortex lacked the critical early interaction with the language cortex, due to the lack of strong innate connectivity between the two in the right hemisphere (cf. the left hemisphere; e.g., Li et al., 2020; Powell et al., 2016), resulting in a more distributed representation for visual words.

And second, how are multivariate representations read out by downstream brain regions for further processing (Williams et al., 2007); specifically, whereas the behavioral relevance of univariate activity has been extensively investigated in both adults and pediatric populations (e.g., Furl et al., 2011; Grill-Spector et al., 2004; Huang et al., 2014; Song et al., 2015), the behavioral relevance of multivariate representations is still under debate. One study found that performance in a visual search task among different pairs of object categories could be predicted from underlying neural similarity structures (M. A. Cohen et al., 2016). At this point, it is unclear how multivariate representations for words contribute to EG's reading ability, and whether the right-hemisphere representations are more important—questions that will be a focus of future studies.

Limitations and future directions

Overall, our study provides unique evidence that in the absence of a typical left-hemisphere language network—at least in cases of early left-hemisphere damage—the canonical word-selective IVWFA may not develop. Some limitations are worth noting. First, our analyses were restricted to the ventral temporal cortex, which was the focus of our investigation and where we had coverage. Although we show that normal reading ability is possible in the absence of focal

selectivity for word processing in the VTC, there may be other pathways or neural structures outside of the VTC that are important for EG's reading ability? For example, Seghier et al. (2012) reported a patient who acquired dyslexia following extensive left ventral occipitotemporal cortex (LvOT) resection, but then regained reading ability; they provided evidence to suggest that the patient's reading ability was now supported via a direct connection between the occipital visual cortex and the left superior temporal sulcus (STS) without the involvement of the ventral visual stream. It is therefore possible that the STS in the language-dominant hemisphere (i.e., the right hemisphere) may support visual word processing through its connections with the occipital cortex. Future work will explore word processing in EG using whole-brain coverage. Second, EG's missing left superior temporal cortex led to right-lateralized speech and language processing. As discussed earlier, it remains unclear whether speech and higher-level language areas contribute equally to the emergence of the IVWFA, and whether temporal language regions may be more important than the frontal ones. Finally, here we reported a single case that sheds new light on the role of speech and language areas in developing a VWFA. Generalization to other, similar cases, and data from other methods like noninvasive brain stimulation will help us better understand the causal role of language cortex in developing visual word selectivity.

References

- Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A Domain-General Cognitive Core Defined in Multimodally Parcellated Human Cortex. *Cerebral Cortex*, 30(8), 4361–4380. <https://doi.org/10.1093/cercor/bhaa023>
- Baker, C., Liu, J., Wald, L., Kwong, K. K., Benner, T., & Kanwisher, N. (2007). Visual word processing and experiential origins of functional selectivity in human extrastriate cortex. *Proc Natl Acad Sci USA*. <https://www.pnas.org/content/104/21/9087>
- Barttfeld, P., Abboud, S., Lagercrantz, H., Adén, U., Padilla, N., Edwards, A. D., Cohen, L., Sigman, M., Dehaene, S., & Dehaene-Lambertz, G. (2018). A lateral-to-mesial organization of human ventral visual cortex at birth. *Brain Structure and Function*, 223(7), 3107–3119. <https://doi.org/10.1007/s00429-018-1676-3>
- Behrmann, M., & Plaut, D. C. (2013). Distributed circuits, not circumscribed centers, mediate visual recognition. *Trends in Cognitive Sciences*, 17(5), 210–219. <https://doi.org/10.1016/j.tics.2013.03.007>
- Behrmann, M., & Plaut, D. C. (2015). A vision of graded hemispheric specialization: Graded hemispheric specialization. *Annals of the New York Academy of Sciences*, 1359(1), 30–46. <https://doi.org/10.1111/nyas.12833>
- Behrmann, M., & Plaut, D. C. (2020). Hemispheric Organization for Visual Object Recognition: A Theoretical Account and Empirical Evidence. *Perception*, 49(4), 373–404. <https://doi.org/10.1177/0301006619899049>
- Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where Is the Semantic System? A Critical Review and Meta-Analysis of 120 Functional Neuroimaging Studies. *Cerebral Cortex*, 19(12), 2767–2796. <https://doi.org/10.1093/cercor/bhp055>

- 831 Bishop, D. (2003). *Test for Reception of Grammar version 2 (TROG-2)*.
832
- 833 Blank, I., Kanwisher, N., & Fedorenko, E. (2014). A functional dissociation between language
834 and multiple-demand systems revealed in patterns of BOLD signal fluctuations. *Journal*
835 *of Neurophysiology*, 112(5), 1105–1118. <https://doi.org/10.1152/jn.00884.2013>
- 836 Bouhali, F., Thiebaut de Schotten, M., Pinel, P., Poupon, C., Mangin, J.-F., Dehaene, S., &
837 Cohen, L. (2014). Anatomical Connections of the Visual Word Form Area. *Journal of*
838 *Neuroscience*, 34(46), 15402–15414. <https://doi.org/10.1523/JNEUROSCI.4918-13.2014>
- 839 Cai, Q., Lavidor, M., Brysbaert, M., Paulignan, Y., & Nazir, T. A. (2008). Cerebral
840 Lateralization of Frontal Lobe Language Processes and Lateralization of the Posterior
841 Visual Word Processing System. *Journal of Cognitive Neuroscience*, 20(4), 672–681.
842 <https://doi.org/10.1162/jocn.2008.20043>
- 843 Cai, Q., Paulignan, Y., Brysbaert, M., Ibarrola, D., & Nazir, T. A. (2010). The Left Ventral
844 Occipito-Temporal Response to Words Depends on Language Lateralization but Not on
845 Visual Familiarity. *Cerebral Cortex*, 20(5), 1153–1163.
846 <https://doi.org/10.1093/cercor/bhp175>
- 847 Cohen, L., Martinaud, O., Lemer, C., Lehericy, S., Samson, Y., Obadia, M., Slachevsky, A., &
848 Dehaene, S. (2003). Visual Word Recognition in the Left and Right Hemispheres:
849 Anatomical and Functional Correlates of Peripheral Alexias. *Cerebral Cortex*, 13(12),
850 1313–1333. <https://doi.org/10.1093/cercor/bhg079>
- 851 Cohen, M. A., Alvarez, G. A., Nakayama, K., & Konkle, T. (2016). Visual search for object
852 categories is predicted by the representational architecture of high-level visual cortex.
853 *Journal of Neurophysiology*, 117(1), 388–402. <https://doi.org/10.1152/jn.00569.2016>

854 Cohen, M. A., Dilks, D. D., Koldewyn, K., Weigelt, S., Feather, J., Kell, A. JE., Keil, B., Fischl,
855 B., Zöllei, L., Wald, L., Saxe, R., & Kanwisher, N. (2019). Representational similarity
856 precedes category selectivity in the developing ventral visual pathway. *NeuroImage*, 197,
857 565–574. <https://doi.org/10.1016/j.neuroimage.2019.05.010>

858 Crawford, J. R., & Garthwaite, P. H. (2007). Comparison of a single case to a control or
859 normative sample in neuropsychology: Development of a Bayesian approach. *Cognitive*
860 *Neuropsychology*, 24(4), 343–372. <https://doi.org/10.1080/02643290701290146>

861 Crawford, J. R., Garthwaite, P. H., & Howell, D. C. (2009). On comparing a single case with a
862 control sample: An alternative perspective. *Neuropsychologia*, 47(13), 2690–2695.
863 <https://doi.org/10.1016/j.neuropsychologia.2009.04.011>

864 Crawford, J. R., & Howell, D. C. (1998). Comparing an Individual's Test Score Against Norms
865 Derived from Small Samples. *The Clinical Neuropsychologist*, 12(4), 482–486.
866 <https://doi.org/10.1076/clin.12.4.482.7241>

867 Davis, T., LaRocque, K. F., Mumford, J. A., Norman, K. A., Wagner, A. D., & Poldrack, R. A.
868 (2014). What do differences between multi-voxel and univariate analysis mean? How
869 subject-, voxel-, and trial-level variance impact fMRI analysis. *NeuroImage*, 97, 271–283.
870 <https://doi.org/10.1016/j.neuroimage.2014.04.037>

871 Deen, B., Richardson, H., Dilks, D. D., Takahashi, A., Keil, B., Wald, L. L., Kanwisher, N., &
872 Saxe, R. (2017). Organization of high-level visual cortex in human infants. *Nature*
873 *Communications*, 8(1), 13995. <https://doi.org/10.1038/ncomms13995>

874 Dehaene, S., & Cohen, L. (2007). Cultural Recycling of Cortical Maps. *Neuron*, 56(2), 384–398.
875 <https://doi.org/10.1016/j.neuron.2007.10.004>

- Dehaene, S., & Cohen, L. (2011). The unique role of the visual word form area in reading.
Trends in Cognitive Sciences, 15(6), 254–262. <https://doi.org/10.1016/j.tics.2011.04.003>
- Dehaene, S., Cohen, L., Morais, J., & Kolinsky, R. (2015). Illiterate to literate: Behavioural and cerebral changes induced by reading acquisition. *Nature Reviews Neuroscience*, 16(4), 234–244. <https://doi.org/10.1038/nrn3924>
- Dehaene-Lambertz, G., Monzalvo, K., & Dehaene, S. (2018). The emergence of the visual word form: Longitudinal evolution of category-specific ventral visual areas during reading acquisition. *PLOS Biology*, 16(3), e2004103.
<https://doi.org/10.1371/journal.pbio.2004103>
- Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, 31(3), 968–980.
<https://doi.org/10.1016/j.neuroimage.2006.01.021>
- Downing, P. E., Jiang, Y., Shuman, M., & Kanwisher, N. (2001). A cortical area selective for visual processing of the human body. *Science (New York, N.Y.)*, 293(5539), 2470–2473.
<https://doi.org/10.1126/science.1063414>
- Duncan, J., Assem, M., & Shashidhara, S. (2020). Integrated Intelligence from Distributed Brain Activity. *Trends in Cognitive Sciences*, 24(10), 838–852.
<https://doi.org/10.1016/j.tics.2020.06.012>
- Dunn, L. M., & Dunn, D. M. (2007). *The Peabody Picture Vocabulary Test, Fourth Edition*. NCS
Pearson, Inc.

899 Epstein, R., & Kanwisher, N. (1998). A cortical representation of the local visual environment.
900 *Nature*, 392(6676), 598–601. <https://doi.org/10.1038/33402>

901 Fedorenko, E., Behr, M. K., & Kanwisher, N. (2011). Functional specificity for high-level
902 linguistic processing in the human brain. *Proceedings of the National Academy of*
903 *Sciences*, 108(39), 16428–16433. <https://doi.org/10.1073/pnas.1112937108>

904 François, C., Garcia-Alix, A., Bosch, L., & Rodriguez-Fornells, A. (2021). Signatures of brain
905 plasticity supporting language recovery after perinatal arterial ischemic stroke. *Brain and*
906 *Language*, 212, 104880. <https://doi.org/10.1016/j.bandl.2020.104880>

907 Furl, N., Garrido, L., Dolan, R. J., Driver, J., & Duchaine, B. (2011). Fusiform Gyrus Face
908 Selectivity Relates to Individual Differences in Facial Recognition Ability. *Journal of*
909 *Cognitive Neuroscience*, 23(7), 1723–1740. <https://doi.org/10.1162/jocn.2010.21545>

910 Gerrits, R., Van der Haegen, L., Brysbaert, M., & Vingerhoets, G. (2019). Laterality for
911 recognizing written words and faces in the fusiform gyrus covaries with language
912 dominance. *Cortex*, 117, 196–204. <https://doi.org/10.1016/j.cortex.2019.03.010>

913 Glezer, L. S., Jiang, X., & Riesenhuber, M. (2009). Evidence for Highly Selective Neuronal
914 Tuning to Whole Words in the “Visual Word Form Area.” *Neuron*, 62(2), 199–204.
915 <https://doi.org/10.1016/j.neuron.2009.03.017>

916 Greenblatt, S. H. (1976). Subangular alexia without agraphia or hemianopsia. *Brain and*
917 *Language*, 3(2), 229–245. [https://doi.org/10.1016/0093-934X\(76\)90019-5](https://doi.org/10.1016/0093-934X(76)90019-5)

918 Grill-Spector, K., Knouf, N., & Kanwisher, N. (2004). The fusiform face area subserves face
919 perception, not generic within-category identification. *Nature Neuroscience*, 7(5), 555–
920 562. <https://doi.org/10.1038/nn1224>

921 Hamamé, C. M., Szwed, M., Sharman, M., Vidal, J. R., Perrone-Bertolotti, M., Kahane, P.,
922 Bertrand, O., & Lachaux, J.-P. (2013). Dejerine's reading area revisited with intracranial
923 EEG: Selective responses to letter strings. *Neurology*, 80(6), 602–603.
924 <https://doi.org/10.1212/WNL.0b013e31828154d9>

925 Hannagan, T., Amedi, A., Cohen, L., Dehaene-Lambertz, G., & Dehaene, S. (2015). Origins of
926 the specialization for letters and numbers in ventral occipitotemporal cortex. *Trends in*
927 *Cognitive Sciences*, 19(7), 374–382. <https://doi.org/10.1016/j.tics.2015.05.006>

928 Hasson, U., Levy, I., Behrmann, M., Hendler, T., & Malach, R. (2002). Eccentricity Bias as an
929 Organizing Principle for Human High-Order Object Areas. *Neuron*, 34(3), 479–490.
930 [https://doi.org/10.1016/S0896-6273\(02\)00662-1](https://doi.org/10.1016/S0896-6273(02)00662-1)

931 Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001).
932 Distributed and Overlapping Representations of Faces and Objects in Ventral Temporal
933 Cortex. *Science*, 293(5539), 2425–2430. <https://doi.org/10.1126/science.1063736>

934 Hsiao, J. H., & Lam, S. M. (2013). The Modulation of Visual and Task Characteristics of a
935 Writing System on Hemispheric Lateralization in Visual Word Recognition—A
936 Computational Exploration. *Cognitive Science*, 37(5), 861–890.
937 <https://doi.org/10.1111/cogs.12033>

938 Huang, L., Song, Y., Li, J., Zhen, Z., Yang, Z., & Liu, J. (2014). Individual differences in
939 cortical face selectivity predict behavioral performance in face recognition. *Frontiers in*
940 *Human Neuroscience*, 8. <https://doi.org/10.3389/fnhum.2014.00483>

941 Julian, J. B., Fedorenko, E., Webster, J., & Kanwisher, N. (2012). An algorithmic method for
942 functionally defining regions of interest in the ventral visual pathway. *NeuroImage*, 60(4),
943 2357–2364. <https://doi.org/10.1016/j.neuroimage.2012.02.055>

944 Kamps, F. S., Hendrix, C. L., Brennan, P. A., & Dilks, D. D. (2020). Connectivity at the origins
945 of domain specificity in the cortical face and place networks. *Proceedings of the National*
946 *Academy of Sciences*, 117(11), 6163–6169. <https://doi.org/10.1073/pnas.1911359117>

947 Kanwisher, N. (2010). Functional specificity in the human brain: A window into the functional
948 architecture of the mind. *Proceedings of the National Academy of Sciences*, 107(25),
949 11163–11170. <https://doi.org/10.1073/pnas.1005062107>

950 Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The Fusiform Face Area: A Module in
951 Human Extrastriate Cortex Specialized for Face Perception. *The Journal of Neuroscience*,
952 17(11), 4302–4311. <https://doi.org/10.1523/JNEUROSCI.17-11-04302.1997>

953 Kaufman, A. S., & Kaufman, N. L. (2004). *Kaufman Brief Intelligence Test (2nd ed.)*. American
954 Guidance Service.

955 Kay, J., Coltheart, M., & Lesser, R. (1992). *Psycholinguistic assessments of language processing*
956 *in aphasia (PALPA) (1st Edition)*. Psychology Press.

957 Konkle, T., & Caramazza, A. (2013). Tripartite Organization of the Ventral Stream by Animacy
958 and Object Size. *Journal of Neuroscience*, 33(25), 10235–10242.

959 Lerma-Usabiaga, G., Carreiras, M., & Paz-Alonso, P. M. (2018). Converging evidence for
960 functional and structural segregation within the left ventral occipitotemporal cortex in
961 reading. *Proceedings of the National Academy of Sciences*, 115(42), E9981–E9990.
962 <https://doi.org/10.1073/pnas.1803003115>

963 Li, J., Huang, L., Song, Y., & Liu, J. (2017). Dissociated neural basis of two behavioral
964 hallmarks of holistic face processing: The whole-part effect and composite-face effect.
965 *Neuropsychologia*, 102, 52–60. <https://doi.org/10.1016/j.neuropsychologia.2017.05.026>

966 Li, J., Osher, D. E., Hansen, H. A., & Saygin, Z. M. (2020). Innate connectivity patterns drive
967 the development of the visual word form area. *Scientific Reports*, 10(1), 18039.
968 <https://doi.org/10.1038/s41598-020-75015-7>

969 Liu, T. T., Freud, E., Patterson, C., & Behrmann, M. (2019). Perceptual Function and Category-
970 Selective Neural Organization in Children with Resections of Visual Cortex. *Journal of*
971 *Neuroscience*, 39(32), 6299–6314. <https://doi.org/10.1523/JNEUROSCI.3160-18.2019>

972 Mahon, B. Z., & Caramazza, A. (2011). What drives the organization of object knowledge in the
973 brain? *Trends in Cognitive Sciences*, 15(3), 97–103.
974 <https://doi.org/10.1016/j.tics.2011.01.004>

975 Malach, R., Levy, I., & Hasson, U. (2002). The topography of high-order human object areas.
976 *Trends in Cognitive Sciences*, 6(4), 176–184. [https://doi.org/10.1016/S1364-](https://doi.org/10.1016/S1364-6613(02)01870-3)
977 [6613\(02\)01870-3](https://doi.org/10.1016/S1364-6613(02)01870-3)

978 Martin, A. (2006). Shades of Déjerine—Forging a Causal Link between the Visual Word Form
979 Area and Reading. *Neuron*, 50(2), 173–175. <https://doi.org/10.1016/j.neuron.2006.04.004>

980 Mbwana, J., Berl, M. M., Ritzl, E. K., Rosenberger, L., Mayo, J., Weinstein, S., Conry, J. A.,
981 Pearl, P. L., Shamim, S., Moore, E. N., Sato, S., Vezina, L. G., Theodore, W. H., &
982 Gaillard, W. D. (2009). Limitations to plasticity of language network reorganization in
983 localization related epilepsy. *Brain*, 132(2), 347–356.
984 <https://doi.org/10.1093/brain/awn329>

985 McCandliss, B. D., Cohen, L., & Dehaene, S. (2003). The visual word form area: Expertise for
986 reading in the fusiform gyrus. *Trends in Cognitive Sciences*, 7(7), 293–299.
987 [https://doi.org/10.1016/S1364-6613\(03\)00134-7](https://doi.org/10.1016/S1364-6613(03)00134-7)

988 Morais, J., Cary, L., Alegria, J., & Bertelson, P. (1979). Does awareness of speech as a sequence
989 of phones arise spontaneously? *Cognition*, 7(4), 323–331. [https://doi.org/10.1016/0010-](https://doi.org/10.1016/0010-0277(79)90020-9)
990 0277(79)90020-9

991 Nasr, S., Echavarria, C. E., & Tootell, R. B. H. (2014). Thinking Outside the Box: Rectilinear
992 Shapes Selectively Activate Scene-Selective Cortex. *Journal of Neuroscience*, 34(20),
993 6721–6735. <https://doi.org/10.1523/JNEUROSCI.4802-13.2014>

994 Newport, E. L., Landau, B., Seydell-Greenwald, A., Turkeltaub, P. E., Chambers, C. E.,
995 Dromerick, A. W., Carpenter, J., Berl, M. M., & Gaillard, W. D. (2017). Revisiting
996 Lenneberg’s Hypotheses About Early Developmental Plasticity: Language Organization
997 After Left-Hemisphere Perinatal Stroke. *Biolinguistics*, 11, 407–422.

998 Oosterhof, N. N., Connolly, A. C., & Haxby, J. V. (2016). CoSMoMVPA: Multi-Modal
999 Multivariate Pattern Analysis of Neuroimaging Data in Matlab/GNU Octave. *Frontiers in*
1000 *Neuroinformatics*, 10. <https://doi.org/10.3389/fninf.2016.00027>

1001 Osher, D. E., Saxe, R., Koldewyn, K., Gabrieli, J. D. E., Kanwisher, N., & Saygin, Z. M. (2015).
1002 Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple Visual
1003 Categories across Cortex. *Cerebral Cortex*, 26(4), 1668–1683.

1004 Patterson, K., Nestor, P. J., & Rogers, T. T. (2007). Where do you know what you know? The
1005 representation of semantic knowledge in the human brain. *Nature Reviews Neuroscience*,
1006 8(12), 976–987. <https://doi.org/10.1038/nrn2277>

1007 Pennington, B. F., & Bishop, D. V. M. (2008). Relations Among Speech, Language, and
1008 Reading Disorders. *Annual Review of Psychology*, 60(1), 283–306.
1009 <https://doi.org/10.1146/annurev.psych.60.110707.163548>

1010 Postelnicu, G., Zollei, L., & Fischl, B. (2009). Combined Volumetric and Surface Registration.
1011 *IEEE Transactions on Medical Imaging*, 28(4), 508–522.
1012 <https://doi.org/10.1109/TMI.2008.2004426>

1013 Powell, H. W. R., Parker, G. J. M., Alexander, D. C., Symms, M. R., Boulby, P. A., Wheeler-
1014 Kingshott, C. A. M., Barker, G. J., Noppeney, U., Koepp, M. J., & Duncan, J. S. (2006).
1015 Hemispheric asymmetries in language-related pathways: A combined functional MRI and
1016 tractography study. *NeuroImage*, 32(1), 388–399.
1017 <https://doi.org/10.1016/j.neuroimage.2006.03.011>

1018 Raschle, N. M., Zuk, J., & Gaab, N. (2012). Functional characteristics of developmental dyslexia
1019 in left-hemispheric posterior brain regions predate reading onset. *Proceedings of the*
1020 *National Academy of Sciences*, 109(6), 2156–2161.
1021 <https://doi.org/10.1073/pnas.1107721109>

1022 Rauschecker, A. M., Deutsch, G. K., Ben-Shachar, M., Schwartzman, A., Perry, L. M., &
1023 Dougherty, R. F. (2009). Reading impairment in a patient with missing arcuate fasciculus.
1024 *Neuropsychologia*, 47(1), 180–194.
1025 <https://doi.org/10.1016/j.neuropsychologia.2008.08.011>

1026 Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K., & Goebel, R. (2021). A
1027 Probabilistic Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cerebral*
1028 *Cortex*, 31(1), 603–619. <https://doi.org/10.1093/cercor/bhaa246>

1029 Rossion, B., Dricot, L., Devolder, A., Bodart, J.-M., Crommelinck, M., Gelder, B. de, &
1030 Zoontjes, R. (2000). Hemispheric Asymmetries for Whole-Based and Part-Based Face
1031 Processing in the Human Fusiform Gyrus. *Journal of Cognitive Neuroscience*, 12(5),
1032 793–802. <https://doi.org/10.1162/089892900562606>

- 1033 Saygin, Z. M., Osher, D. E., Koldewyn, K., Reynolds, G., Gabrieli, J. D. E., & Saxe, R. (2012).
1034 Anatomical connectivity patterns predict face selectivity in the fusiform gyrus. *Nature*
1035 *Neuroscience*, 15, 321–327.
- 1036 Saygin, Z. M., Osher, D. E., Norton, E. S., Youssoufian, D. A., Beach, S. D., Feather, J., Gaab,
1037 N., Gabrieli, J. D. E., & Kanwisher, N. (2016). Connectivity precedes function in the
1038 development of the visual word form area. *Nature Neuroscience*, 19(9), 1250–1255.
1039 <https://doi.org/10.1038/nn.4354>
- 1040 Seghier, M. L., Neufeld, N. H., Zeidman, P., Leff, A. P., Mechelli, A., Nagendran, A., Riddoch, J.
1041 M., Humphreys, G. W., & Price, C. J. (2012). Reading without the left ventral occipito-
1042 temporal cortex. *Neuropsychologia*, 50(14), 3621–3635.
1043 <https://doi.org/10.1016/j.neuropsychologia.2012.09.030>
- 1044 Seghier, M. L., & Price, C. J. (2011). Explaining Left Lateralization for Words in the Ventral
1045 Occipitotemporal Cortex. *Journal of Neuroscience*, 31(41), 14745–14753.
1046 <https://doi.org/10.1523/JNEUROSCI.2238-11.2011>
- 1047 Shaywitz, B. A., Shaywitz, S. E., Pugh, K. R., Mencl, W. E., Fulbright, R. K., Skudlarski, P.,
1048 Constable, R. T., Marchione, K. E., Fletcher, J. M., Lyon, G. R., & Gore, J. C. (2002).
1049 Disruption of posterior brain systems for reading in children with developmental dyslexia.
1050 *Biological Psychiatry*, 52(2), 101–110. [https://doi.org/10.1016/S0006-3223\(02\)01365-3](https://doi.org/10.1016/S0006-3223(02)01365-3)
- 1051 Song, Y., Zhu, Q., Li, J., Wang, X., & Liu, J. (2015). Typical and Atypical Development of
1052 Functional Connectivity in the Face Network. *Journal of Neuroscience*, 35(43), 14624–
1053 14635. <https://doi.org/10.1523/JNEUROSCI.0969-15.2015>
- 1054 Staudt, M. (2010). Reorganization after pre- and perinatal brain lesions*. *Journal of Anatomy*,
1055 217(4), 469–474. <https://doi.org/10.1111/j.1469-7580.2010.01262.x>

1056 Staudt, M., Grodd, W., Niemann, G., Wildgruber, D., Erb, M., & Krägeloh–Mann, I. (2001).
1057 Early left periventricular brain lesions induce right hemispheric organization of speech.
1058 *Neurology*, 57(1), 122–125. <https://doi.org/10.1212/WNL.57.1.122>

1059 Stevens, W. D., Kravitz, D. J., Peng, C. S., Tessler, M. H., & Martin, A. (2017). Privileged
1060 Functional Connectivity between the Visual Word Form Area and the Language System.
1061 *The Journal of Neuroscience*, 37(21), 5288–5297.
1062 <https://doi.org/10.1523/JNEUROSCI.0138-17.2017>

1063 Tadros, K., Dupuis-Roy, N., Fiset, D., Arguin, M., & Gosselin, F. (2013). Reading laterally: The
1064 cerebral hemispheric use of spatial frequencies in visual word recognition. *Journal of*
1065 *Vision*, 13(1), 4–4. <https://doi.org/10.1167/13.1.4>

1066 Tuckute, G., Paunov, A., Kean, H., Small, H., Mineroff, Z., Blank, I., & Fedorenko, E. (2021).
1067 Frontal language areas do not emerge in the absence of temporal language areas: A case
1068 study of an individual born without a left temporal lobe. *BioRxiv*, 2021.05.28.446230.
1069 <https://doi.org/10.1101/2021.05.28.446230>

1070 Vaidya, A. R., Pujara, M. S., Petrides, M., Murray, E. A., & Fellows, L. K. (2019). Lesion
1071 Studies in Contemporary Neuroscience. *Trends in Cognitive Sciences*, 23(8), 653–671.
1072 <https://doi.org/10.1016/j.tics.2019.05.009>

1073 Van der Haegen, L., Cai, Q., & Brysbaert, M. (2012). Colateralization of Broca’s area and the
1074 visual word form area in left-handers: fMRI evidence. *Brain and Language*, 122(3),
1075 171–178. <https://doi.org/10.1016/j.bandl.2011.11.004>

1076 Wandell, B. A., Rauschecker, A. M., & Yeatman, J. D. (2012). Learning to See Words. *Annual*
1077 *Review of Psychology*, 63(1), 31–53. [https://doi.org/10.1146/annurev-psych-120710-](https://doi.org/10.1146/annurev-psych-120710-100434)
1078 100434

1079 Williams, M. A., Dang, S., & Kanwisher, N. G. (2007). Only some spatial patterns of fMRI
1080 response are read out in task performance. *Nature Neuroscience*, 10(6), 685–686.
1081 <https://doi.org/10.1038/nn1900>

1082 Wilmer, J. B., Germine, L., Chabris, C. F., Chatterjee, G., Williams, M., Loken, E., Nakayama,
1083 K., & Duchaine, B. (2010). Human face recognition ability is specific and highly
1084 heritable. *Proceedings of the National Academy of Sciences*, 107(11), 5238–5241.
1085 <https://doi.org/10.1073/pnas.0913053107>

1086 Yeatman, J. D., & Feldman, H. M. (2013). Neural plasticity after pre-linguistic injury to the
1087 arcuate and superior longitudinal fasciculi. *Cortex*, 49(1), 301–311.
1088 <https://doi.org/10.1016/j.cortex.2011.08.006>

1089 Zhu, Q., Song, Y., Hu, S., Li, X., Tian, M., Zhen, Z., Dong, Q., Kanwisher, N., & Liu, J. (2010).
1090 Heritability of the Specific Cognitive Ability of Face Perception. *Current Biology*, 20(2),
1091 137–142. <https://doi.org/10.1016/j.cub.2009.11.067>
1092