E3 ligase TRIM47 positively regulates endothelial activation and pulmonary inflammation through potentiating the K63-linked ubiquitination

3

4 Yisong Qian^{1,2}, Ziwei Wang¹, Hongru Lin¹, Tianhua Lei², Zhou Zhou², Weilu Huang¹,
5 Xuehan Wu¹, Li Zuo¹, Jie Wu¹, Yu Liu¹, Ling-Fang Wang¹, Xiao-Hui Guan¹, Ke-Yu Deng¹,
6 Mingui Fu^{2,*}, Hong-Bo Xin^{1,*}

7

8 ¹ The National Engineering Research Center for Bioengineering Drugs and the Technologies,
9 Institute of Translational Medicine, Nanchang University, 1299 Xuefu Rd, Honggu District,
10 Nanchang 330031, China

11 ² Department of Biomedical Science, School of Medicine, University of Missouri Kansas
12 City, 2411 Holmes Street, Kansas City, MO 64108, USA

13

14 *Correspondence: Hong-Bo Xin, The National Engineering Research Center for
15 Bioengineering Drugs and the Technologies, Institute of Translational Medicine, Nanchang
16 University, 1299 Xuefu Rd, Honggu District, Nanchang 330031, China. Tel:
17 86-791-83827165; E-mail: xinhb@ncu.edu.cn.

18 Mingui Fu, Shock/Trauma Research Center & Department of Basic Medical Science, School
19 of Medicine, University of Missouri Kansas City, MO 64108. Tel: 816-235-2193; Fax:
20 816-235-6444; E-mail: fum@umkc.edu.

21

22

23 **Abstract**

24 Endothelial activation plays an essential role in the pathology of sepsis-induced acute lung
 25 injury, but the detailed regulatory mechanisms remain largely unknown. Here, we
 26 demonstrated that TRIM47, an ubiquitin E3 ligase of tripartite protein family, is highly
 27 expressed in vascular endothelial cells and is up-regulated during TNF α -induced endothelial
 28 activation. Knockdown of TRIM47 in endothelial cells prevents the transcription of multiple
 29 pro-inflammatory cytokines, reduces monocyte adhesion and the expression of adhesion
 30 molecules, and inhibits the secretion of IL-1 β and IL-6 into the supernatant. By contrast,
 31 overexpression of TRIM47 promotes inflammatory response and monocyte adhesion upon
 32 TNF α stimulation. TRIM47 modulates the activation of NF- κ B and MAPK signaling
 33 pathways during endothelial activation. Further experiment confirmed that TRIM47 interacts
 34 with TRAF2 and mediates K63-linked ubiquitination. In addition, TRIM47-deficient mice are
 35 more resistant to lipopolysaccharide-induced acute lung injury and death, due to attenuated
 36 pulmonary inflammation. Taken together, our studies suggest that TRIM47 promotes
 37 pulmonary inflammation and injury at least partly through potentiating the K63-linked
 38 ubiquitination of TRAF2, which in turn activates NF- κ B and MAPK signaling pathways to
 39 trigger inflammatory response in endothelial cells.

40 **Keywords:** TRIM47; endothelial cells; inflammatory response; ubiquitination; acute lung
 41 injury

42

Introduction

Acute Lung injury (ALI) comprise a uniform response of the lung to inflammatory or chemical insults and is therefore commonly generated by systemic illness including sepsis or trauma, infection with pathogens and toxic gas inhalation¹. Despite a great deal of effort has been devoted to target the immune response to infection and adjunct approaches, there are few effective therapeutic strategies due to the emergence of new pathogens such as the global pandemic of novel coronavirus pneumonia, as well as the continued rise of drug resistance².

The pulmonary endothelium is critically implicated in the pathogenesis of ALI as a main target of circulating cells and humoral mediators under injury¹. The interaction between endothelial cells and leukocytes is a key step in the development of ALI. Leukocyte adhesion to endothelial cells and migration across endothelial cells are mediated by the interaction of complementary adhesion molecules on leukocytes and endothelial cells. The increased expression or release of endothelial cell adhesion molecules is a hallmark of endothelial cell activation^{3,4}. Upon the access of leukocytes into the lung parenchyma, they can release inflammatory mediators to destroy pathogens, but the over-activated immune response have potential to result in the imbalance of the pro-inflammatory and anti-inflammatory mechanisms, triggering "cytokine storm" and subsequent tissue damage^{2,5}. Considering the critical role of endothelial response during ALI, strategies that targeting endothelial components, including cell surface receptors⁶, signaling pathways, transcriptional networks, and endothelial cell gene products, have been recently proposed to attenuate endothelial activation and improve endothelial dysfunction⁷.

Tripartite motif-containing (TRIM) proteins, a subfamily of E3 ubiquitin ligases, have been widely involved in many physiological processes including cell proliferation and differentiation, innate immunity and autophagy⁸. Multiple TRIM proteins have been found to participate in innate immunity through positive or negative regulation of cytokines, toll like receptors, pattern recognition receptors, intracellular signaling pathways and transcription factors⁹. Numerous studies have focused on the roles and regulatory mechanisms of TRIM in immune cells¹⁰⁻¹⁴, there are relatively few studies on TRIM proteins involved in the regulation of endothelial inflammation. It has been shown that TRIM28 is abundant in endothelial cells, and interfering with TRIM28 expression has anti-inflammatory and anti-angiogenic phenotypes¹⁵. However, the precise roles of TRIM family members in regulating of endothelial function remain largely unknown.

TRIM47 was originally found in brain astrocytomas and was named GOA (gene overexpressed in astrocytoma). It is prominently located in nucleus and its LXXLL motifs are thought to be closely related to nuclear receptor binding¹⁶. Emerging evidence showed that TRIM47 have functions on tumorigenesis and progression¹⁷⁻¹⁹, viral resistance processes²⁰, and cerebral ischemia-reperfusion injury²¹. A recent genome-wide association study showed that the SNPs of TRIM47 and TRIM65, located in the adjacent position of human chromosome 17, are closely related to white matter hyperintensities, which are thought to reflect ischemic damage to the small deep cerebral vessels²². These findings suggested that TRIM47 and TRIM65 may coordinately or independently participate in the regulation of cerebrovascular injury. Our previous work demonstrated that TRIM65, as an E3 ubiquitin ligase, selectively targets VCAM-1 and promote its ubiquitination and degradation, thus reducing lung inflammation and damage caused by sepsis²³, but the role of TRIM47 in

endothelial inflammation remains to be elucidated.

In the present study, we investigate the effects of TRIM47 in endothelial activation in an *in vitro* model of inflammation induced by TNF α . The global TRIM47 knockout mice was constructed to confirm its role in ALI. The E3 ubiquitin ligase activity and substrate molecules of TRIM47, as well as the signaling pathways involved were also explored.

Materials and methods

Reagents

Human recombinant TNF α and LPS (from *Escherichia coli* O111:B4) were purchased from Sigma. VCAM-1 (sc-13160), ICAM-1 (sc-1511-R), β -actin (sc-47778), Histone H2A (sc-10807) antibodies were from Santa Cruz Biotechnology. TRIM47 antibody (26885-1-AP) was purchased from Proteintech. Phospho-p65 (3033), p65 (8242), I κ B α (4812), phospho-I κ B α (2859), phospho-IKK α / β (2078), IKK α (11930), IKK β (8943), phospho-JNK (4668), JNK (9252), phospho-ERK1/2 (4370), ERK1/2 (4695), phospho-p38 (4511), p38 (8690), Flag (8146 and 2368) and α -tubulin (2125) antibodies were purchased from Cell Signaling Technology. Ubiquitin (ab7780), ubiquitin (K48, ab140601), and ubiquitin (K63, ab179434) antibodies were purchased from Abcam.

Cell culture, infection and treatment

Human Umbilical Vein Endothelial Cells (HUVEC) and THP-1 cells were purchased from Lonza Walkersville Inc. HUVECs were cultured in EGM medium according to the manufacturer instruction, and used for experiment in less than eight passages. THP-1 was cultured in RPMI-1640 medium supplemented with 10% FBS and 2-mercaptoethanol to a final concentration of 0.05 mM. RAW264.7, EA.hy926, bEnd.3, and HEK293 cells were purchased from ATCC and cultured in DMEM supplemented with 10% FBS. U251, HeLa, MDA-MB-23, 3T3-L1 and A549 cell lines were from National Collection of Authenticated Cell Cultures (Shanghai, China) and cultured in DMEM supplemented with 10% FBS. The hCMEC/D3 cell line was purchased from BeNa Culture Collection (Beijing, China) and cultured in RPMI-1640 medium supplemented with 10% FBS. HL-1 cardiac muscle cell line was obtained from Sigma Aldrich and cultured in Claycomb medium supplemented with 100 μ M norepinephrine, 4 mM l-glutamine and 10% FBS. The siRNA target sequence was selected in Human TRIM47 gene (GenBank accession NM_033452.2) as follows: siRNA1: 5'-TGAAGCTCCCAGGGACTATTT-3', and siRNA2: 5'-TACTGGGAGGTGGAGATTATC-3'. TRIM47 siRNA was constructed into the lentivirus expression vector pLV[shRNA]-EGFP:T2A:Puro-U6. A universal sequence was used as a negative control for RNA interference. Human TRIM47 gene was constructed into pLV[Exp]-EGFP:T2A:Puro-EF1A vector to obtain the expression lentiviral vector. The viral particles were produced by third generation packaging in 293T cells and Lentiviral stocks were concentrated using ultracentrifugation. HUVECs (5×10^4 /ml) were prepared and infected at a Multiplicity of Infection (MOI) of 50 with negative control, TRIM47 siRNA1, TRIM47 siRNA2 or TRIM47 overexpression lentiviruses for 16 h at 37°C in the presence of 10 mg/ml polybrene. The cultures were then washed and cultured in fresh medium for 72 h. GFP expression was detected to calculate the infection efficiency. Then, cells were treated with 10 ng/ml TNF α for indicated times, and mRNAs or proteins from those cells were

extracted and detected.

RNA isolation and QPCR

Total tissue or cellular RNA was isolated using TRIzol reagents, according to the manufacturer's instructions (Life Technologies, CA, USA). One microgramme of total RNA was reverse-transcribed using a One Step PrimeScript™ RT-PCR Kit (Takara, Liaoning, China) with a thermocycler. The mRNA levels were determined by SYBR Green dye using an ABI 7500 sequence detection system with a reaction mixture that consisted of SYBR Green 2×PCR Master Mix (Applied Biosystems, CA, USA), cDNA template, and the forward and reverse primers. Primer sequences are listed in Table S1. The PCR protocol consisted of 40 cycles of denaturation at 95 °C for 15 s followed by 60 °C for 1 min to allow extension and amplification of the target sequence. Data were analyzed using ABI 7500 sequence detection system software. The amount of mRNA was normalized to GAPDH using the $2^{-\Delta\Delta CT}$ method. The results were from three independent experiments performed in triplicate.

Protein isolation and western blot

Tissue extracts and whole-cell lysates were prepared in radioimmunoprecipitation assay buffer (Thermo Scientific) supplemented with 1 mM PMSF. Nuclear and cytoplasmic protein fractions from cells were extracted by Nuclear-Cytosol Extraction Kit (Applygen Technologies Inc, Beijing, China), according to the manufacturer's instructions. Fifty micrograms protein per sample was loaded in each lane and separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to nitrocellulose membranes (Pall Corporation, USA) in Tris-glycine buffer (48 mM Tris, 39 mM glycine, pH 9.2) containing 20% methanol. The membranes were blocked with skimmed milk for 1 h, washed in Tris buffered saline containing 0.1% Tween-20 (TBST), and incubated with primary antibodies overnight at 4 °C. After washing in TBST for three times, nitrocellulose membranes were incubated for 1 h at room temperature with the horseradish peroxidase conjugated IgG (1:5000; Santa Cruz Biotechnology, Inc, CA, USA). The bands were visualized by the SuperSignal West Pico Chemiluminescent Substrate Trial Kit (Pierce, Rockford, IL, USA). The immunodetected protein bands were then analyzed using ChemiDoc XRS system with Quantity One software (Bio-Rad, Richmond, CA, USA).

Immunocytochemistry

At the end of the treatment, cells were rinsed with phosphate-buffered saline (PBS) three times, fixed with 4% paraformaldehyde for 30 min at room temperature and permeabilised in 0.1% Triton X-100 for 10 min. An incubation in 5% bovine serum albumin (BSA) in PBS for 1 h was performed to prevent antibody non-specific binding. The cultures were incubated with primary antibodies overnight at 4 °C. After incubation with primary antibodies, cells were incubated with fluorescein isothiocyanate (FITC)-conjugated goat anti-rabbit IgG (Alexa 488; 1:1000; invitrogen) and the nuclei were stained with DAPI. Immunostained cells were examined under a fluorescence microscope (Olympus IX71, Tokyo, Japan).

Ubiquitination assay with Co-IP

HUVECs were transfected with Flag-tagged TRIM47 plasmid or empty vector using

electroporation. The protease inhibitor MG132 (10 mM; Sigma-Aldrich) was added 4 h before harvest. At 24 h posttransfection, cells were lysed in CellLytic M Cell lysis buffer with protease inhibitors, phosphatase inhibitors, NEM, and ubiquitin aldehyde. For endogenous IP, HUVECs were harvested after 15 min-TNF α (10 ng/mL) exposure. The immune complexes were collected by incubation (2 h, 4°C) with protein G-agarose (Sigma). Co-IP assays were performed by using PierceTM Protein G Agarose (Thermo Fisher) followed the manufacturer's instruction. After extensive washing, the electrophoresis loading buffer was added to the complexes and incubated for 5 minutes at 95°C. Immunoprecipitated proteins were resolved by SDS-PAGE and analyzed by immunoblotting with indicated antibodies.

Generation of TRIM47 knockout mice

To define the physiological role of TRIM47 in vivo, we have obtained the mice with heterozygous TRIM47-targeted allele by using CRISPR/Cas9 to remove all exons of TRIM47 gene (Fig. 5A). The mice with homozygous TRIM47-targeted alleles were generated by interbreeding. The mice were created in C57BL/6 background. Genotyping was done with the following primers: Trim47-F: 5'-GGTAAACACAGTCGCTAAGAGGTCAA-3', Trim47-R: 5'-TGGTCTAGGGATGCCAGGGTTCT-3' and Trim47-Wt/He-F: 5'-AGTCAGAGTGAGCAGGCAGGAGAATA-3' (Fig. 5B). Wild type and TRIM47 knockout mice were housed in the Animal Centre of Institute of Translational Medicine, Nanchang University, with a 12 h light-dark cycle, optimal temperature and humidity, filtered water, and appropriate nutrient feed. All procedures related to the care of animals were performed according to the National Institutes of Health Guide for the Care and Use of Laboratory Animals. All experimental protocols were approved by Institutional Animal Care and use Committee of Nanchang University. The lung tissue extracts were examined by western blot analysis and the results showed that TRIM47 protein was abolished in the homozygous targeted allele (Fig. 5C), which generated the global knockout mouse (designated as TRIM47^{-/-}).

LPS challenge in mice

Age-matched mice (7-9 weeks) were randomly assigned to control or experimental groups. Wild type and TRIM47^{-/-} mice underwent an intraperitoneal injection of LPS (15 mg/kg) to induce lethal endotoxic shock. The control group received injections of the equivalent volume of 0.9% NaCl solution. After injection, the mice were closely monitored for general condition and survival for 7 days.

Measurements of inflammatory cytokines

The concentrations of TNF α , IL-1 β and IL-6 in the serum, and IL-1 β and IL-6 levels in the supernatant of HUVECs were measured using the specific ELISA kits according to the manufacturer's instructions (Neobioscience Technology Co., Ltd., Shenzhen, China). Absorbance at 450 nm wave length was measured, and the protein concentration was determined by interpolation on to absorbance curves generated by recombinant protein standards using iMarkTM Microplate Absorbance Reader (Bio-Rad).

Histological analysis

The right upper lungs were removed after 24 h of LPS challenge and were fixed in 4% phosphate-buffered paraformaldehyde. The 4- μ m paraffin tissue sections were cut and stained with H&E as previously described²⁴. Photomicrographs were taken by a light microscope (Olympus BX51). Lung injury was evaluated by an independent pathologist who was blinded to the grouping, taking into account haemorrhage in the lung tissue, alveolar congestion, edema, infiltration of macrophages and neutrophils, and morphological changes in the alveolar wall.

Measurement of lung wet/dry weight ratio

To evaluate the magnitude of pulmonary edema, the wet-to-dry weight ratios at 24 h after LPS challenge were determined. The left lung tissue samples were weighed immediately after removal (wet weight) and then subjected to desiccation in an oven at 50 °C until a stable dry weight was achieved after 72 h. The ratio of the wet/dry weight was then calculated.

Statistics

Statistical analysis was performed with GraphPad Prism software (GraphPad, San Diego, CA). Data were expressed as mean $\pm$ SD. For comparison between two groups, the unpaired Student's t-test was used. For multiple comparison, one way ANOVA followed by Turkey's post hoc analysis was used. A value of $p < 0.05$ was considered significant.

Results

TRIM47 is highly expressed in vascular endothelial cells

The mRNA and protein levels of TRIM47 were detected in various tissues of mice. Results showed that TRIM47 mRNA was highly expressed in heart, lung, kidney and epididymal white adipose tissue (eWAT, Fig. 1A). TRIM47 protein levels were abundant in heart, lung, stomach and testis (Fig. 1B). The immunohistochemistry demonstrated that TRIM47 had a high expression in lung, kidney tubules, heart and eWAT, moderate expression in brain, stomach, skin and colon, and low expression in liver, testis, spleen and thymus (Fig. S1). In particular, an obvious positive staining was observed in the vascular lining of multiple tissues, as indicated by the arrows, including lung, brain, subcutaneous tissue and colon (Fig. 1C). Next, we detected TRIM47 expression in different cells. Real-time PCR and western blot results revealed that TRIM47 was specifically expressed in human umbilical vein endothelial cells (HUVEC and EA.hy926) and brain microvascular endothelial cells (bEnd.3), whereas TRIM47 showed low expression in monocyte/macrophages (Fig. 1D and 1E). In addition, we measured the mRNA levels of 56 TRIM family members in HUVECs and hCMECs, respectively. TRIM47 exhibited a moderate expression in both cells (Fig. 1F and 1G). These results suggested that TRIM47 is highly expressed in vascular endothelial cells.

TRIM47 is induced by inflammatory stimulation in endothelial cells

We further investigated the expression changes of TRIM47 following exogenous stimuli. Result showed that the mRNA and protein levels of TRIM47 were significantly up-regulated by LPS, H₂O₂ and TNF α (Fig. 2A-C). The expression of TRIM47 was increased since 2 h after TNF α stimulation and peaked at 12 h (Fig. 2C). Similarly, TRIM47 expression was also up-regulated after TNF α challenge in bEnd.3 cells upon various stimulation, including

inflammation and hypoxia (Fig. S2A-D). However, TRIM47 was not sensitive to TNF α challenge but remarkably decreased by LPS exposure in macrophages (Fig. S2E-H). The localization of TRIM47 was measured by western blot in cytosolic and nuclear fractions, respectively. The nucleus contains a prominent proportion of TRIM47, which was significantly up-regulated after TNF α exposure. The cytosolic TRIM47 was also induced after TNF α stimulation (Fig. 2D). The immunocytochemistry further confirmed that TRIM47 mainly but not solely located in the nucleus of HUVECs. One-hour stimulation of TNF α resulted in an increase in fluorescence density, which was partly resumed at 8 h of TNF α exposure (Fig. 2E). TRIM47 also responded to LPS challenge in HUVECs (Fig. S3A), and was up-regulated in hCMEC/D3 following TNF α and LPS incubation (Fig. S3B and S3C). The expression profile of TRIM47 indicated that it may play a role during inflammatory stimulation.

TRIM47 promotes TNF α -induced endothelial activation

The siRNA and overexpression vectors were introduced to explore the role of TRIM47 in endothelial activation. Knockdown of TRIM47 significantly reduced TNF α -induced mRNA expression of multiple adhesion molecules and pro-inflammatory cytokines (Fig. 3A), which were elevated in TRIM47-overexpressed HUVECs. TRIM47 siRNAs obviously decreased the protein levels of ICAM-1 and VCAM-1, and suppressed THP-1 adhesion to HUVECs, whereas overexpression of TRIM47 had opposite effects (Fig. 3C and 3D). TRIM47 siRNAs also suppressed the secretion of pro-inflammatory cytokines IL-1 β and IL-6 into the supernatant (Fig. 3E and 3F). By contrast, overexpression of TRIM47 promotes the production of pro-inflammatory cytokines (Fig. 3G and 3H). In addition, knockdown of TRIM47 remarkably inhibited cell proliferation and migration induced by TNF α in HUVECs (Fig. S4). These results suggested that TRIM47 is a positive regulator of endothelial activation.

TRIM47 modulates NF- κ B and MAPK pro-inflammatory signaling pathways

The potential signaling pathway involved in TRIM47-mediated endothelial activation was investigated, including NF- κ B and MAPK. Results showed that knockdown of TRIM47 significantly inhibited the phosphorylation of I κ B α , IKK α / β and p65 subunit, and prevented the degradation of I κ B α (Fig. 4A), whereas overexpression of TRIM47 promotes the activation of NF- κ B signaling pathway (Fig. 4B). TRIM47 siRNAs also suppressed the activation of JNK and p38 MAPK signal pathways but had no obvious effect on ERK (Fig. 4C). Overexpression of TRIM47 further activated JNK and p38 signal pathways after TNF α stimulation (Fig. 4D). These results suggested that TRIM47 mediates endothelial activation at least partly through NF- κ B and MAPK activation.

TRIM47 mediates K63-linked Ubiquitylation and interacts with TRAF2

Emerging evidence showed that TRIM proteins mediated K48- or K63-linked ubiquitination to activate NF- κ B signaling pathway in response to exogenous stimulation^{12, 25}. Therefore, the ubiquitination pattern of TRIM47 involved in endothelial activation was analyzed. We observed that TNF- α induced total ubiquitination and K63-linked ubiquitination of TRIM47 in HUVECs by Co-immunoprecipitation with the TRIM47 antibody (Fig. 5A).

HUVECs were transfected with Flag or Flag-TRIM47 vector for 48 h. Co-immunoprecipitation was performed with the Flag antibody. The exogenous assay further confirmed that TRIM47 was involved in K63-linked ubiquitination rather than K48 (Fig. 5B). Tumor necrosis factor receptor-associated factor 2 (TRAF2) is a key adaptor molecule in TNFR signaling complexes that promotes downstream signaling cascades, such as NF- κ B and MAPK activation²⁶, whereas TRAF6 is the major transducer of IL-1 receptor/TLR signaling²⁷. It has been reported that the K63-linked polyubiquitin chains could be attached to TRAF2, serving as a scaffold to recruit TAK1, TAB1, and TAB2. The active TAK1 further phosphorylates the MAPKs and IKK complex to initiate MAPK and NF- κ B cascades²⁸. We therefore examined the possible binding proteins of TRIM47 involved in this signal pathway. As shown in Fig. 5C, TRIM47 bind with TRAF2 but not TRAF6. TRIM47 did not interact with the downstream proteins including TAK1, IKK γ and I κ B α . In addition, TRIM47 did not induce the degradation of I κ B α , a classical target of K48-linked ubiquitination (Fig. S5). These results suggest that TRIM47 regulated NF- κ B and MAPK activation by enhancing the K63-linked ubiquitination of TRAF2.

TRIM47 deficiency alleviates acute lung injury and inflammatory response in LPS-challenged mice

We produced the global TRIM47 knockout mice to investigate its role in systemic inflammatory response and organ injury (Fig 6A-C). The TRIM47 KO mice had no significant changes in viscera index (Table. S2) or histology (Fig. S6) compared with the WT mice. After 24 h of LPS injection, the TRIM47 KO mice showed reduced pulmonary edema compared with the WT animals (Fig. 6D). In addition, TRIM47 deficiency improved the survival rate of mice after LPS challenge (Fig. 6E). HE staining showed that there were no significant differences in lung histology between WT and TRIM47 KO mice under normal conditions. A significant tissue damage appeared in the lungs of WT mice, including neutrophil infiltration, alveolar wall thickening, hemorrhage, alveolar edema and alveolar disruption. The histological changes were significantly alleviated in TRIM47 KO mice (Fig. 6F).

TRIM47 and various pro-inflammatory cytokines in the lung and serum were measured after 6 h and 24 h following LPS injection, respectively. The mRNA levels of pro-inflammatory cytokines were much lower in TRIM47 KO mice than that in WT mice (Fig. 6G). The knockout mice also had lower levels of TNF α , IL-1 β and IL-6 content in the serum (Fig. 6H). These results indicated that TRIM47 deficiency attenuates systemic inflammatory and acute lung injury during LPS challenge.

Discussion

Here we demonstrated that TRIM47 play crucial roles in TNF α -induced endothelial activation. As a potential E3 ubiquitin ligase, TRIM47 interacts with TRAF2 and mediates K63-linked ubiquitination, activating NF- κ B and MAPK signaling pathways. In LPS challenged mice, TRIM47 deficiency significantly reduced pulmonary inflammation and tissue damage. These results strongly suggested that TRIM47 is a positive regulator of endothelial activation.

TRIM47 was originally found to be overexpressed in astrocytoma tumor cells and

astrocytes of fetal brain, with prominent nuclear staining, but virtual absence in mature astrocytes. The mRNA levels of TRIM47 in normal tissues were low except for in kidney¹⁶. Here we showed that TRIM47 has relatively high expression in heart, lung, kidney and eWAT by detecting the mRNA and protein levels respectively. The immunohistochemistry also demonstrated its extensive distribution in these tissues. Interestingly, the vascular endothelial cells from different tissues exhibited remarkable positive staining of TRIM47. The samples from cell cultures further confirmed its prominent expression in various endothelial cells. Therefore TRIM47 may be identified as a specifically expressed protein in endothelial cells where may take its functions.

A various TRIM proteins have been proved to be implicated in innate immunity and inflammatory response. Our previous work showed that TRIM47 is down-regulated upon TNF α exposure in THP1-derived macrophages²⁹. According to the present study, TRIM47 was significantly induced by multiple stimuli, including TNF α , LPS, hypoxia and oxidative stress in endothelial cells, indicating that TRIM47 may be a sensor in response to exogenous stimulation. In agreement with the previous report¹⁶, we found that TRIM47 was prominent located in the nucleus of HUVECs and was expressed in cytoplasm upon TNF α challenge. These result suggested TRIM47-mediated endothelial activation may depend on its cytoplasm location, although its role in nucleus remain unclear.

Different from the anti-inflammatory effect of TRIM65, another constitutive expressed protein in endothelial cells²³, TRIM47 promotes inflammatory response both *in vitro* and *in vivo*. The *in vivo* results further confirmed that the pro-inflammatory phenotype primarily depend on endothelial activation rather than macrophages. Since the previous GWAS results suggested that both TRIM47 and TRIM65 contribute to the cerebral vessels injury²², we may suppose that TRIM47 and TRIM65 are critically involved in the regulation of endothelial inflammation and injury by different mechanisms.

Proteins covalently modified with K48-linked poly-ubiquitin are targeted for proteasomal degradation, whereas proteins covalently modified with K63-linked poly-ubiquitin generally become functionally activated⁹. Recent work showed that TRIM proteins positively or negatively mediate NF- κ B activation primary through K48- or K63-linked poly-ubiquitin. For example, TRIM5 interacts with TAK1-containing kinase complex and positively regulate NF- κ B activation by mediating K63 poly-ubiquitin chain synthesis¹². TRIM25 promote NF- κ B activation by enhancing the K63-linked ubiquitination of TRAF2 and bridging the interaction of TRAF2 and TAK1 or IKK β ²⁸. TRIM14 enhances NF- κ B activation in endothelial cells via directly binding to NEMO and promotes the phosphorylation of I κ B α and p65, which is dependent on its K63-linked ubiquitination³⁰. TRIM47 is structurally similar to TRIM25, with a Ring-finger domain, a B2 box and its associated coil-coiled region at the N terminus, and a PRY domain at the C terminus⁸. Functionally like TRIM25, TRIM47 also interacts with TRAF2 and promotes MAPK and NF- κ B activation through K63-linked ubiquitination. These findings provide the underlying mechanisms by which TRIM47 specifically meditates endothelial activation (Fig. 7).

To sum up, we identified a novel endothelial activation factor TRIM47, which mediates inflammatory response in endothelial cells and promote inflammation and tissue damage during ALI through endothelial TRAF2-MAPK/NF- κ B pro-inflammatory axis. Currently there remain no compounds targeting TRIM proteins at the laboratory or clinical level but it

is important to develop inhibitors of TRIM proteins for their use as a therapeutic tools in multiple diseases⁸. TRIM47 would be an attractive drug target for endothelial inflammation and ALI. Further detailed analysis of TRIM47 is needed for its use for effective therapy and to eliminate side effects.

Acknowledgement

This work was supported by the National Natural Science Foundation of China (82070080 and 81860020 to YQ, 81873659 to HBX, and 81760140 and 81970256 to KYD), Foundation for the National Institutes of Health (1R15AI138116 to M.F), and the financial support provided by China Scholarship Council (201906825031 to YQ).

CRedit authorship contribution statement

Yisong Qian, Ziwei Wang, Hongru Lin, Tianhua Lei, Zhou Zhou, Weilu Huang, Xuehan Wu, Li Zuo, Jie Wu, Yu Liu: Investigation, Methodology. Ling-Fang Wang, Xiao-Hui Guan and Ke-Yu Deng: Resources. Yisong Qian: Writing -original draft, Conceptualization, Supervision. Mingui Fu and Hong-Bo Xin: Conceptualization, Supervision, Writing -review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Maniatis NA, Kotanidou A, Catravas JD, Orfanos SE. Endothelial pathomechanisms in acute lung injury. *Vascular pharmacology*. Oct-Dec 2008;49(4-6):119-33. doi:10.1016/j.vph.2008.06.009
- Tisoncik JR, Korth MJ, Simmons CP, Farrar J, Martin TR, Katze MG. Into the eye of the cytokine storm. *Microbiology and molecular biology reviews : MMBR*. Mar 2012;76(1):16-32. doi:10.1128/MMBR.05015-11
- Orfanos SE, Mavrommati I, Korovesi I, Roussos C. Pulmonary endothelium in acute lung injury: from basic science to the critically ill. *Intensive care medicine*. Sep 2004;30(9):1702-14. doi:10.1007/s00134-004-2370-x
- Vallet B. Bench-to-bedside review: endothelial cell dysfunction in severe sepsis: a role in organ dysfunction? *Critical care*. Apr 2003;7(2):130-8. doi:10.1186/cc1864
- Ince C, Mayeux PR, Nguyen T, et al. The Endothelium in Sepsis. *Shock*. Mar 2016;45(3):259-70. doi:10.1097/SHK.0000000000000473
- Gleeson EM, O'Donnell JS, Preston RJ. The endothelial cell protein C receptor: cell surface conductor of cytoprotective coagulation factor signaling. *Cell Mol Life Sci*. Mar 2012;69(5):717-26. doi:10.1007/s00018-011-0825-0
- Aird WC. The role of the endothelium in severe sepsis and multiple organ dysfunction syndrome. *Blood*. May 15 2003;101(10):3765-77. doi:10.1182/blood-2002-06-1887
- Hatakeyama S. TRIM Family Proteins: Roles in Autophagy, Immunity, and Carcinogenesis. *Trends in biochemical sciences*. Apr 2017;42(4):297-311. doi:10.1016/j.tibs.2017.01.002

9. Versteeg GA, Benke S, Garcia-Sastre A, Rajsbaum R. InTRIMsic immunity: Positive and negative regulation of immune signaling by tripartite motif proteins. *Cytokine Growth Factor Rev.* Oct 2014;25(5):563-76. doi:10.1016/j.cytogfr.2014.08.001
10. Hoffpauir CT, Bell SL, West KO, et al. TRIM14 Is a Key Regulator of the Type I IFN Response during Mycobacterium tuberculosis Infection. *J Immunol.* Jul 1 2020;205(1):153-167. doi:10.4049/jimmunol.1901511
11. Maarifi G, Smith N, Maillet S, et al. TRIM8 is required for virus-induced IFN response in human plasmacytoid dendritic cells. *Sci Adv.* Nov 2019;5(11):eaax3511. doi:10.1126/sciadv.aax3511
12. Pertel T, Hausmann S, Morger D, et al. TRIM5 is an innate immune sensor for the retrovirus capsid lattice. *Nature.* Apr 21 2011;472(7343):361-5. doi:10.1038/nature09976
13. Yu T, Kuang H, Chen J, et al. Tripartite-motif family protein 35-28 regulates microglia development by preventing necrotic death of microglial precursors in zebrafish. *J Biol Chem.* Jun 26 2020;295(26):8846-8856. doi:10.1074/jbc.RA119.012043
14. Zhao W, Wang L, Zhang M, Yuan C, Gao C. E3 ubiquitin ligase tripartite motif 38 negatively regulates TLR-mediated immune responses by proteasomal degradation of TNF receptor-associated factor 6 in macrophages. *J Immunol.* Mar 15 2012;188(6):2567-74. doi:10.4049/jimmunol.1103255
15. Wang Y, Li J, Huang Y, et al. Tripartite motif-containing 28 bridges endothelial inflammation and angiogenic activity by retaining expression of TNFR-1 and -2 and VEGFR2 in endothelial cells. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology.* May 2017;31(5):2026-2036. doi:10.1096/fj.201600988RR
16. Vandeputte DA, Meije CB, van Dartel M, et al. GOA, a novel gene encoding a ring finger B-box coiled-coil protein, is overexpressed in astrocytoma. *Biochemical and biophysical research communications.* Aug 24 2001;286(3):574-9. doi:10.1006/bbrc.2001.5431
17. Wang Y, Liu C, Xie Z, Lu H. Knockdown of TRIM47 inhibits breast cancer tumorigenesis and progression through the inactivation of PI3K/Akt pathway. *Chemico-biological interactions.* Feb 1 2020;317:108960. doi:10.1016/j.cbi.2020.108960
18. Liang Q, Tang C, Tang M, Zhang Q, Gao Y, Ge Z. TRIM47 is up-regulated in colorectal cancer, promoting ubiquitination and degradation of SMAD4. *Journal of experimental & clinical cancer research : CR.* Apr 12 2019;38(1):159. doi:10.1186/s13046-019-1143-x
19. Han Y, Tian H, Chen P, Lin Q. TRIM47 overexpression is a poor prognostic factor and contributes to carcinogenesis in non-small cell lung carcinoma. *Oncotarget.* Apr 4 2017;8(14):22730-22740. doi:10.18632/oncotarget.15188
20. Wang Y, Kuang M, Lu Y, Lin L, Liu X. Characterization and biological function analysis of the TRIM47 gene from common carp (Cyprinus carpio). *Gene.* Sep 5 2017;627:188-193. doi:10.1016/j.gene.2017.06.017
21. Hao MQ, Xie LJ, Leng W, Xue RW. Trim47 is a critical regulator of cerebral ischemia-reperfusion injury through regulating apoptosis and inflammation. *Biochemical and biophysical research communications.* Aug 6 2019;515(4):651-657. doi:10.1016/j.bbrc.2019.05.065
22. Fornage M, Debette S, Bis JC, et al. Genome-wide association studies of cerebral white

matter lesion burden: the CHARGE consortium. *Ann Neurol.* Jun 2011;69(6):928-39.
doi:10.1002/ana.22403

23. Li Y, Huang X, Guo F, et al. TRIM65 E3 ligase targets VCAM-1 degradation to limit LPS-induced lung inflammation. *Journal of molecular cell biology.* Apr 24 2020;12(3):190-201. doi:10.1093/jmcb/mjz077

24. Liang J, Saad Y, Lei T, et al. MCP-induced protein 1 deubiquitinates TRAF proteins and negatively regulates JNK and NF-kappaB signaling. *J Exp Med.* Dec 20 2010;207(13):2959-73. doi:10.1084/jem.20092641

25. van Gent M, Sparrer KMJ, Gack MU. TRIM Proteins and Their Roles in Antiviral Host Defenses. *Annu Rev Virol.* Sep 29 2018;5(1):385-405.
doi:10.1146/annurev-virology-092917-043323

26. Park ES, Choi S, Shin B, et al. Tumor necrosis factor (TNF) receptor-associated factor (TRAF)-interacting protein (TRIP) negatively regulates the TRAF2 ubiquitin-dependent pathway by suppressing the TRAF2-sphingosine 1-phosphate (S1P) interaction. *J Biol Chem.* Apr 10 2015;290(15):9660-73. doi:10.1074/jbc.M114.609685

27. Xing Y, Yao X, Li H, et al. Cutting Edge: TRAF6 Mediates TLR/IL-1R Signaling-Induced Nontranscriptional Priming of the NLRP3 Inflammasome. *J Immunol.* Sep 1 2017;199(5):1561-1566. doi:10.4049/jimmunol.1700175

28. Liu Y, Liu K, Huang Y, et al. TRIM25 Promotes TNF-alpha-Induced NF-kappaB Activation through Potentiating the K63-Linked Ubiquitination of TRAF2. *J Immunol.* Mar 15 2020;204(6):1499-1507. doi:10.4049/jimmunol.1900482

29. Jiang MX, Hong X, Liao BB, et al. Expression profiling of TRIM protein family in THP1-derived macrophages following TLR stimulation. *Sci Rep.* Feb 17 2017;7:42781. doi:10.1038/srep42781

30. Huang X, Li Y, Li X, Fan D, Xin HB, Fu M. TRIM14 promotes endothelial activation via activating NF-kappaB signaling pathway. *Journal of molecular cell biology.* Apr 24 2020;12(3):176-189. doi:10.1093/jmcb/mjz040

Figure legends

Fig. 1 TRIM47 is highly expressed in vascular endothelial cells. (A) The mRNA levels of TRIM47 in various tissues of mice were detected by real-time PCR and visualized by agarose gel electrophoresis. (B) The protein levels of TRIM47 were detected in various tissues of mice by western blot. (C) The immunohistochemistry showed positive staining of TRIM47 in the vascular lining of multiple tissues. (D) TRIM47 mRNA expression in different cells detected by real-time PCR. (E) The protein levels of TRIM47 in different cells detected by western blot. (F) The mRNA levels of 56 TRIM family members in HUVECs and hCMEC/D3 detected by real-time PCR.

Fig. 2 TRIM47 is induced by inflammatory stimulation in endothelial cells. TRIM47 mRNA and protein levels in HUVECs after (A) LPS, (B) H₂O₂ and (C) TNF α stimulation were examined by real-time PCR and western blot, respectively. (D) The translocation of TRIM47 following TNF α treatment was measured by western blot in cytosolic and nuclear fractions. (E) The distribution of TRIM47 in HUVECs was detected by immunocytochemistry in the absence or presence of TNF α .

Fig. 3 TRIM47 promotes inflammatory response in endothelial cells. The mRNA expression of multiple adhesion molecules and pro-inflammatory cytokines was detected by real-time PCR in (A) TRIM47 siRNA- and (B) overexpression vector-treated HUVECs. (C) The protein levels of ICAM-1 and VCAM-1 were measured by western blot in TRIM47 siRNA- and overexpression vector-treated HUVECs. (D) TRIM47 siRNA or expression vector was transfected into HUVECs, and the transfected cells were incubated with TNF- α or PBS for 8 h and then co-cultured with Zombi-ER-labeled THP-1 cells for 1 h. The adhesion of THP-1 was observed under a fluorescence microscope. The contents of (E) IL-1 β and (F) IL-6 in siRNA-treated cells, and (G) IL-1 β and (H) IL-6 in overexpression vector-treated cells were measured by ELISA.

Fig. 4 TRIM47 modulates endothelial activation through NF- κ B and MAPK signaling pathway. The activation of NF- κ B was detected by western blot in (A) TRIM47 siRNA- and (B) overexpression vector-treated HUVECs. The activation of MAPK, including ERK, JNK and p38 signaling pathways was detected by western blot in (C) TRIM47 siRNA- and (D) overexpression vector-treated HUVECs.

Fig. 5 TRIM47 mediates K63-linked Ubiquitylation and interacts with TRAF2. (A) HUVECs were treated with 10 ng/ml TNF- α for 0 and 15 min. Cell lysates then were immunoprecipitated using TRIM47 antibody, followed by western blot analysis using the indicated antibodies. (B) HUVECs were transiently transfected with empty vector or TRIM47 plasmid for 48 h. Whole-cell lysates were immunoprecipitated with Flag antibody and the precipitates were immunoblotted with Ub, TRAF2, TRAF5, TAK1, IKK γ , and I κ B α antibodies.

Fig. 6 TRIM47 deficiency reduces endotoxemia-induced acute lung injury and pulmonary inflammation in mice. (A) Schematic strategy of generation of TRIM47

knockout mice. (B) Genotyping of TRIM47^{+/+}, TRIM47^{+/-}, and TRIM47^{-/-} mice. (C) Western blot analysis for TRIM47 in lungs from TRIM47^{+/+} and TRIM47^{-/-} mice. (D) Pulmonary edema was represented as lung wet-to-dry ratio. (E) Survival rate of mice challenged with 15 mg/kg LPS (i.p.). (F) The histological changes in lungs after LPS challenge were examined by HE staining. (G) The mRNA levels of TRIM47 and various pro-inflammatory cytokines in the lung were measured by real-time PCR. (F) The content of IL-1β, IL-6 and TNFα in the serum was assayed by ELISA.

Fig.7 The schematic diagram of the mechanism by which TRIM47 promotes TNFα-induced endothelial inflammation and acute lung injury.

A **B** **D** **C** **E** **F** **G**

A **B** **C** **D** **E**

A **B** **C** **D** **E** **F** **G** **H**

A **B** **C** **D**

A

B

C