Evolution of degrees of carnivory and dietary specialization across Mammalia and their effects on speciation

Matthew D. Pollard^{1,2} and Emily E. Puckett^{1,2}

¹Department of Biological Sciences, University of Memphis, Memphis, TN 38152

²Center for Biodiversity Research, University of Memphis, Memphis, TN 38152

ABSTRACT

Conflation between omnivory and dietary generalism limits ecological and evolutionary analyses of diet, including estimating contributions to speciation and diversification. Additionally, categorizing species into qualitative dietary classes leads to information loss in these analyses. Here, we constructed two continuous variables – degree of carnivory (i.e., the position along the continuum from complete herbivory to complete carnivory) and degree of dietary specialization (i.e., the number and variety of food resources utilized) – to elucidate their histories across Mammalia, and to tease out their independent contributions to mammalian speciation. We observed that degree of carnivory significantly affected speciation rate across Mammalia, whereas dietary specialization did not. We further considered phylogenetic scale in diet-dependent speciation and saw that degree of carnivory significantly affected speciation in ungulates, carnivorans, bats, eulipotyphlans, and marsupials, while the effect of dietary specialization was only significant in carnivorans. Across Mammalia, omnivores had the lowest speciation rates. Our analyses using two different categorical diet variables led to contrasting signals of diet-dependent diversification, and subsequently different conclusions regarding diet’s macroevolutionary role. We argue that treating variables such as diet as continuous instead of categorical reduces information loss and avoids the problem of contrasting macroevolutionary signals caused by differential discretization of biologically continuous traits.

Keywords:

Carnivory, diet, speciation rate, mammal, specialization

INTRODUCTION

Diet is a powerful selective force acting during evolution, with consequences for a species' physiology, morphology, behavior, persistence time, and diversification rate. Examples of physiological traits shaped by diet include detoxification capacity (Robbins *et al.* 1991; Haywood *et al.* 2005), digestive enzyme production (Janiak *et al.* 2018), essential nutrient requirements (Knopf *et al.* 1978), and taste receptor function (Zhao *et al.* 2010; Jiang *et al.* 2012; Liu *et al.* 2016). Morphological adaptations that have arisen in response to diet vary in scale from small alterations that modify bite force (Santana *et al.* 2010; Law *et al.* 2018), to the extreme modifications observed in anteaters (Naples 1999) and mysticete whales (Deméré *et al.* 2008). High trophic level and narrow dietary breadth have been implicated in increased extinction risk (e.g., Purvis *et al.* 2000; Boyles and Storm 2007), leading to reduced species durations (Balisi *et al.* 2018).

Earlier work investigated the effect of diet on macroevolutionary dynamics. Price *et al.* (2012) estimated differential diversification rates between herbivores, omnivores, and carnivores. Specifically, omnivores had the lowest diversification rates while herbivores had the highest, and transitions into omnivory were more frequent than into either other category. Similarly, herbivorous birds (Burin *et al.* 2016), insects (Wiens *et al.* 2015), and crustaceans (Poore *et al.* 2017) also show elevated diversification rates. Birds show a similar pattern of low omnivore diversification rate, but a high rate of transitions into omnivory (Burin *et al.* 2016). Omnivory has thus been posited as a macroevolutionary sink, with the suggestion that this pattern may be widespread among higher taxonomic levels of tetrapods.

The effects of ecological factors on mammalian taxonomic diversification change across phylogenetic scales, producing clade- and scale-specific diversification dynamics (Machac *et al.* 2017). At lower taxonomic levels, increasing specialization for frugivory was associated with higher diversification rates in bats (Rojas *et al.* 2012), but a diet-dependent diversification dynamic was not observed in murine rodents (Alhajeri and Steppan 2018). In ruminants, generalist herbivores showed higher diversification rates than browse-specialists (Cantalapiedra *et al.* 2014), which contradicts the prior notion of a positive relationship between dietary specialization and diversification rate in mammals. The conflicting results of the previous studies may be reconciled by the understanding that diet is capable of varying diversification relationships at different taxonomic levels.

Categorizing species into qualitative classes based on diet – especially using a coarse classification system of herbivore, carnivore, and omnivore – leads to a loss of information (Pineda-Munoz and Alroy 2014). Species with highly varied diets are clustered into potentially uninformative groupings. An additional problem with using a categorical variable is the need to choose diet categories and set boundaries between them, resulting in a lack of standardization of diet categorization in the literature. Between different studies, authors set conflicting cutoffs for clustering species. For example, one classification system separates mammals into hypercarnivores (>70% meat), mesocarnivores (50-70% meat), and hypocarnivores (>70% nonvertebrate foods) (Van Valkenburgh 1988, 2007). However, another system classified carnivores as those consuming only animal-based foods, herbivores as those consuming only plant-based foods, and omnivores as everything in between (Price *et al.* 2012). Other studies expand the number of different categories, further subdividing diet (e.g., Cantalapiedra *et al.* 2014; Burin *et al.* 2016). Comparisons between different studies are therefore made difficult by disagreement in

how species should be grouped. A quantitative, continuous classification of diet based on food consumption percentages may be preferred, as downstream results are less impacted by arbitrary cutoffs decided upon earlier in the study (Pineda-Munoz and Alroy 2014).

Omnivory and dietary generalism are distinct yet conflated in some of the literature. Omnivores necessarily show more dietary generalism than an organism specialized to consume a single food type, yet this does not preclude herbivores or carnivores from consuming a broad range of foodstuffs. For example, a strictly herbivorous species (i.e., not an omnivore) may consume several distinct plant-based food items, making it a dietary generalist. Thus, we distinguish the macroevolutionary effects of degree of dietary specialization as independent of the effects of degree of carnivory (i.e., where a species is positioned on the continuum from complete herbivory to complete carnivory).

The independent macroevolutionary contributions of the degree of carnivory and degree of dietary specialization may vary in direction and extent across phylogenetic scales. In this study, our objectives are three-fold. First, we elucidate the history of diet across Mammalia by identifying shifts in the evolution of degrees of carnivory and dietary specialization. Second, we infer the relative importance of the degree of carnivory and degree of dietary specialization to evolution in different mammalian clades. Third, we identify and characterize significant relationships between diet and speciation rate across different taxonomic scales in Mammalia.

METHODS

Preparation of dietary data and tree pruning

We utilized the EltonTraits dataset (Wilman *et al.* 2014) containing dietary data for 5,400 extant mammals. EltonTraits stores dietary preferences as the proportion of diet comprising specific food types. The number of food types was condensed to six categories, giving the proportions of invertebrates, vertebrates, fruit, seeds, nectar, and other plant material (e.g., leaves).

We cross-referenced the species names in EltonTraits with those in the consensus version of the most recent mammal phylogeny ($n = 4,098$) (Upham *et al.* 2019). Mismatched and outdated species names were resolved using the *get_gbif_taxonomy* function of the *traitdataform* R package (Schneider 2018). We removed species not included in both datasets. We pruned the phylogeny using the *drop.tip* function of the *ape* R package (Paradis and Schliep 2018), leaving 3,649 species shared between the phylogeny and dietary dataset.

Construction of “degree of carnivory” variable

We built a continuous variable to represent the degree of carnivory. We first summed the proportions of fruit, seed, nectar, and other plant material into a single value representing the proportion of plant matter. Because the proportion variables sum to 100%, we transformed them into z-scores (Fraser *et al.* 2018). To convert the proportion of invertebrates, vertebrates, and plant matter into a continuous scale, we used discriminant analysis of principal components (DAPC) using the *adegenet* R package (Jombart and Ahmed 2011). The separation between groups is maximized during DAPC compared to principal component analysis (PCA) (Jombart *et al.* 2010). At two clusters, carnivores and herbivores were discriminated, while a third cluster distinguished extreme vertivores and invertivores (Fig. S1). The resulting first discriminant function value was rescaled to a 0 – 100 scale, where 0 signifies complete herbivory and 100 complete carnivory.

Construction of “degree of dietary specialization” variable

We made a second continuous variable to represent dietary specialization. We generated three new variables from the EltonTraits dataset that represent dietary breadth. First, we extracted the percentage (from 20 – 100%) of a species’ most common food type from the original six categories. Second, we summed the number of food types (from 1 – 6) included in each species’ diet. Third, to represent trophic breadth, we gave species that consumed only plants or animals a score of 1, and those that consumed both food types were given a score of 2. We performed a PCA on these three variables in R using the MASS package (Venables and Ripley 2002), then rescaled the first principal component (PC1) to represent the degree of dietary specialization. A specialization score of 0 signifies a complete dietary generalist, whereas a 100 represents a complete dietary specialist.

Regime shift analyses

Regime shifts represent changes in phenotypic optima on a phylogenetic tree under an Ornstein-Uhlenbeck (OU) model of trait evolution. We identified shifts during the evolution of carnivory and dietary specialization across Mammalia using the ℓ 1ou method implemented in the l1ou R package (Khabbazian *et al.* 2016). The ℓ 1ou method fits a multi-optimum OU model of trait evolution using a phylogenetic lasso approach and does not require prior hypotheses regarding the location of shifts. OU models assume random change in a trait value across the tree, but with a centralizing tendency back to an optimum value (Butler and King 2004). A multi-optimum OU model assumes the presence of multiple regime shifts toward different trait optima throughout the phylogeny. The ℓ 1ou method detects the scale and location of shifts in the optimum trait value.

We used a Bayesian Information Criterion (BIC) approach to select the best-fitting configuration of evolutionary regimes. To reduce the number of false shifts detected while using the BIC, we reduced the maximum number of shifts allowed to $n/10$, where n was the number of tips in the tree. This represents a compromise between the default maximum of $n/2$ and the conservative maximum of $\sqrt{n} + 5$ recommended by alternative regime shift analysis methods (Bastide *et al.* 2018).

The ℓ 1ou method can handle large phylogenetic trees comprising thousands of taxa (Khabbazian *et al.* 2016); however, the analyses did not complete when using our pruned mammal phylogeny as input (analysis cancelled after 28 d with 24 CPUs). Therefore, we subset the tree into 13 clades of suitable size for ℓ 1ou. Clades included the orders Artiodactyla, Carnivora, Chiroptera, Eulipotyphla, and Lagomorpha, as well as the higher taxonomic groupings of Atlantogenata, Euarchonta, Marsupialia, and Monotremata. Due to its large size, we separated Rodentia into four clades – Castorimorpha, Hystricomorpha, Sciuromorpha, and Myomorpha + Anomaluromorpha. We analyzed the regime shifts on each clade for both degrees of carnivory and dietary specialization.

Phylogenetic half-lives of degrees of carnivory and dietary specialization

The phylogenetic half-life ($t_{1/2}$) of a trait represents the time it takes for a species entering a new niche or regime to evolve halfway to the new optimum trait value (Hansen 1997). The α parameter returned by each ℓ 1ou analysis represents selection strength and was used to calculate phylogenetic half-life for the degrees of carnivory and dietary specialization in each clade. Because ℓ 1ou rescales the phylogenetic tree height to one, the calculated $t_{1/2}$ values are percentages of the original clade age (Ho and Ané 2014). A short $t_{1/2}$ relative to tree height indicates that evolution towards

new optimum carnivory or dietary specialization scores was fast, with weak residual phylogenetic correlations. Conversely, if $t_{1/2}$ was large relative to tree height, the strength of the OU process was weak and may be indistinguishable from Brownian motion (Cooper *et al.* 2016). Indeed, an α value of zero indicates that the trait did not evolve under an OU process in that clade (Hansen *et al.* 2008; Uyeda and Harmon 2014).

Continuous trait-dependent speciation rates

We characterized the impact of the degree of carnivory and dietary specialization on diversification using the quantitative state speciation and extinction (QuaSSE) method in the diversitree R package (FitzJohn 2010, 2012). For both traits, we constructed various models, each fitting a different relationship between the trait of interest and speciation rate, including: a constant speciation rate independent of the trait value, a linear relationship between trait value and speciation rate, a sigmoidal relationship, and a unimodal relationship. Given known difficulties with extinction rate detection in QuaSSE (FitzJohn 2010) and broader issues with the estimation of extinction from molecular phylogenies (Rabosky 2010), we decided to fit a constant extinction rate instead of allowing the rate to vary with the trait (as in Velasco *et al.* 2016). Thus, if we observed an effect of diet on diversification, it was due to a diet-dependent speciation dynamic. We selected the best-fitting model using the Akaike information criterion (AIC). To account for the influence of phylogenetic uncertainty on our results (Pagel and Lutzoni 2002), we first performed these analyses using the pruned consensus phylogeny, then repeated the analyses on 10 trees sampled from a credible set of 10,000 DNA-only mammalian phylogenies produced by Upham *et al.* (2019). These trees were pruned in the same way as the consensus tree (see above). We fit QuaSSE models for Mammalia and all previously described lower clades, except Rodentia, which was treated as a single clade for this analysis.

We evaluated each QuaSSE model with a simulation approach (Inostroza-Michael *et al.* 2018) to address concerns with high Type 1 error (i.e., that neutral traits are significantly associated with diversification rate (Rabosky and Goldberg 2015)). We used the *fastBM* function in the phytools R package (Revell 2012) to simulate random traits on each phylogeny 100 times under a model of Brownian motion. For these simulations, we used the empirically estimated diffusion parameters from the QuaSSE models with a constant relationship between speciation rate and trait. The ancestral root values for the simulations were estimated using the *fastAnc* function in phytools. We fit the constant model and the best-fitting model for the observed data to each set of simulated trait values and calculated log-likelihood differences between the pairs of models. A *P*-value was created using the proportion of simulated values that achieved a log-likelihood difference as large as that obtained in the observed data. We considered a clade's diet-dependent speciation dynamic to be robust to phylogenetic uncertainty if the *P*-value of the best fitting QuaSSE model was ≤ 0.05 in at least eight out of the eleven phylogenies. To visualize the overall relationship between trait and rate for each robust speciation dynamic, we averaged across the best-fitting models.

Categorical trait-dependent diversification rates

Our work is not directly comparable to previous studies since we created a synthetic continuous variable instead of using categories. Thus, to compare our results, we used the multiple state speciation and extinction (MuSSE) method in the diversitree R package (Fitzjohn 2012) to characterize diet-dependent diversification using different qualitative categorizations of diet. We categorized diet in two ways; first, we used the species and groupings in Price *et al.* (2012) (we will refer to this as Price categorization). There, data was based on literature searches of direct

199 dietary composition from gut, feces, or observational data, which was then binned into plants only
200 (herbivore), plant and animal (omnivore), and animal matter only (carnivore). Second, we
201 replicated the Price categorization but used the raw EltonTraits data (and not our continuous
202 carnivory score), where species with diets composed of 100% animal matter were carnivores, 0%
203 animal matter were herbivores, and 1 – 99% animal matter were omnivores (we will refer to this
204 as Replicated categorization). We limited our MuSSE analyses to species included in both the
205 Price categorization and EltonTraits data (n = 1,323).

206 **RESULTS**

207 *Constructed dietary variables*

208 We created a synthetic variable to quantify degree of carnivory ranging from completely
209 herbivorous (0) to completely carnivorous (100; Fig. 1). Notably, herbivorous dietary guilds
210 (folivores, frugivores, granivores, and nectarivores) had a median score of 0. The median carnivory
211 scores for invertivores and vertivores were 87 and 95, respectively, with overlap between the two
212 dietary guilds. Although both groups are considered carnivorous, the constructed variable
213 attributed a higher degree of carnivory to a vertebrate-consuming species than a species whose diet
214 comprises an equivalent proportion of invertebrates. Species with carnivory scores of 100 were
215 therefore completely vertivorous, an artifact from the variable construction (Fig. S1).

216 We constructed a variable representing the degree of dietary specialization using PC1, which
217 explained 83.2% of the variance (Fig. S2). Lower dietary specialization scores indicated increased
218 dietary generalism. Species with scores of 100 had diets composed of a single food type, whereas
219 0 represented a diet comprising all six food types, with the most abundant category comprising
220 20% of the diet. Dietary specialization showed a convex curvilinear relationship with carnivory,
221 indicating that highly specialized species were also extreme herbivores or extreme carnivores.
222 Because of the higher number of herbivore dietary guilds in EltonTraits, the relationship between
223 the degrees of carnivory and dietary specialization was skewed to the right (Fig. 1A).

Figure 1. (A) The relationship between constructed variables measuring the degree of carnivory and degree of dietary specialization. To visualize how species ($n = 3,649$) were distributed across the variable space, they were grouped into dietary guilds. Species were assigned to a specific dietary guild if their diet comprised $\geq 50\%$ of that food type. If no food type comprised $\geq 50\%$ of a species' diet, they were assigned to the generalist guild. Hypercarnivores are categorized as species that consume $\geq 80\%$ animal matter in their diet, mesocarnivores consume 60 – 80% animal matter, omnivores consume 40 – 60% animal matter, mesoherbivores consume 20 – 40% animal matter, and hyperherbivores consume $\leq 20\%$ animal matter. (B) Distribution of species along the dietary specialization spectrum. (C) Distribution of species along the carnivory spectrum. (D) Distribution of dietary guilds along the carnivory score spectrum.

Regime shifts in degrees of carnivory and dietary specialization across Mammalia

We detected 255 and 227 regime shifts in degree of carnivory and dietary specialization, respectively (Fig. 2-3). The number of regime shifts in each clade varied, and although larger clades tended to have more, this did not translate to a greater proportion of shifts relative to clade size (Table S1). Variation in the optimum value of carnivory and dietary specialization differed between clades. For example, although the order Eulipotyphla had one of the highest frequencies of regime shifts, it had a low standard deviation in optimum carnivory scores among its member species. This shows that, although frequent, the size of the shifts in carnivory score within Eulipotyphla were not large. Most clades had more shifts in carnivory score than they did for dietary specialization. Despite this, the number of clades showing more variation in carnivory

score was equal to the number showing more variation in optimum dietary specialization score. Phylogenetic half-life ($t_{1/2}$) varied greatly between clades, and between carnivory and dietary specialization regimes for a given clade (Table S2). We provide detailed descriptions and interpretations of regime shift results for each clade in the Supporting Information (Figs. S11-S22).

Figure 2. Regime shifts in optimum carnivory score in Mammalia. Branch colors indicate the value of the optimum carnivory score, with blue indicating a low score, and red indicating a high score. The positions of branch color changes show the location of the regime shifts. A gray branch represents lack of data regarding regime occupancy for that branch because it was not included in any regime shift analyses. The numbered black and gray bars encircling the phylogeny indicate the individual clades on which regime shift analyses were performed. 1, Carnivora + Pholidota; 2, Chiroptera; 3, Eulipotyphla; 4, Atlantogenata; 5, Marsupialia; 6, Monotremata; 7, Castorimorpha; 8, Myomorpha + Anomaluromorpha; 9, Hystricomorpha; 10, Sciuromorpha; 11, Lagomorpha; 12, Euarchonta; 13, Artiodactyla + Perissodactyla. Images source: Wikimedia Commons.

Figure 3. Regime shifts in dietary specialization score in Mammalia. Branch colors indicate the value of the optimum specialization score, with yellow indicating a low score, and blue indicating a high score. The positions of branch color changes show the location of the regime shifts. A gray branch represents lack of data regarding regime occupancy for that branch because it was not included in any regime shift analyses. The numbered black and gray bars encircling the phylogeny indicate the individual clades on which regime shift analyses were performed. 1, Carnivora + Pholidota; 2, Chiroptera; 3, Eulipotyphla; 4, Atlantogenata; 5, Marsupialia; 6, Monotremata; 7, Castorimorpha; 8, Myomorpha + Anomaluromorpha; 9, Hystricomorpha; 10, Sciuromorpha; 11, Lagomorpha; 12, Euarchonta; 13, Artiodactyla + Perissodactyla. Images source: Wikimedia Commons.

Diet-dependent speciation rates

We tested if carnivory or dietary specialization was related to speciation rate within each clade using QuaSSE (Table 1). We found significant relationships between speciation rate and degree of carnivory for Artiodactyla + Perissodactyla, Carnivora + Pholidota, Chiroptera, Eulipotyphla, Marsupialia, and Mammalia as a whole; moreover, these relationships were robust to phylogenetic uncertainty. Specifically, most significant QuaSSE models for Mammalia fitted an inverse unimodal relationship between trait and rate (Figs. 4 and S3). Across these models, speciation rate was lowest for intermediate carnivory scores and increased towards both extremes of the carnivory spectrum. However, speciation rates were higher for extreme herbivores than extreme carnivores. Artiodactyla + Perissodactyla, Carnivora + Pholidota, Chiroptera, and Marsupialia also had inverse unimodal relationships, whereas Eulipotyphla had a sigmodal relationship where speciation rate showed a sharp increase at the hypercarnivorous end of the carnivory spectrum (Figs. S4-S8). For dietary specialization, a robust and significant association with speciation rate existed only for Carnivora + Pholidota (Fig. S9). Carnivora + Pholidota had a positive linear relationship where speciation rate was highest for complete dietary specialization and lowest for extreme dietary generalism, reaching zero lineages per Myr for specialization scores below 20.

Table 1. Diet-dependent speciation dynamics as inferred by simulation-based evaluation of best-fitting QuaSSE models using eleven mammal phylogenies. The number of statistically significant QuaSSE models (out of 11) is shown for each taxa and our two variables. Analyses robust to phylogenetic uncertainty are in bold and underlined. Clades for which the best-fitting QuaSSE models were the null were not evaluated via simulations. *P*: the proportion of Brownian motion simulations that produced log-likelihood differences as large as obtained in the observed data.

Clade	Carnivory	Dietary specialization
	No. trees/models with $P \leq 0.05$	No. trees/models with $P \leq 0.05$
Mammalia	<u>10</u>	3
Artiodactyla + Perissodactyla	<u>11</u>	0
Atlantogenata	NA	NA
Carnivora + Pholidota	<u>11</u>	<u>11</u>
Chiroptera	<u>8</u>	5
Euarchonta	1	0
Eulipotyphla	<u>2</u>	0
Lagomorpha	NA	NA
Marsupialia	<u>11</u>	6
Rodentia	6	3

Figure 4. The relationship between degree of carnivory and speciation rate in Mammalia. Across the eleven trees used in this study, ten were associated with a statistically significant best-fitting QuaSSE model. The transparent blue lines show the relationships predicted by these models. The dark blue line represents the relationship produced by averaging over each significant model.

Diet-dependent diversification using categorical variables

Given our creation of a synthetic carnivory score, we decided to compare the results using our continuous scale to a previous categorical analysis. We observed that the diversification rate varied depending on how diet was classified (Fig. 5). Using the Price categorization, herbivores had the highest diversification rates, while omnivores had the lowest (Fig. 5A), concordant with previously published results. While we were able to reproduce the pattern from the previous study using the more recent phylogeny, our estimates of diversification rate were lower for each category compared to Price *et al.* (2012). However, when using the EltonTraits categorization, carnivores had the lowest diversification rates, while omnivores and herbivores had overlapping rates (Fig. 5B).

Transition rates between dietary guilds also changed depending on classification system. Only the MuSSE analysis using the Price categorization reported highest transition rate from herbivory to omnivory (Fig. S10A). Analysis using our EltonTraits categorization reported highest transition rates from omnivory to herbivory (Fig. S10B).

Figure 5. Posterior probability densities for net diversification rate of herbivores, omnivores, and carnivores ($n = 1,323$), using two diet classification schemes: (A) The diet groupings of Price *et al.* (2012) were reused with the most recent mammalian phylogeny (Upham *et al.* 2019), (B) The classification method of Price *et al.* (2012) replicated using EltonTraits data (animal matter: 0% herbivore, 1-99% omnivore, 100% carnivore).

DISCUSSION

Mammalian diversification is shaped more by degree of carnivory than dietary specialization

Our analyses teased apart the independent contributions to mammalian diversification of degree of carnivory and degree of dietary specialization, two aspects of diet that have typically been conflated. We observed that the effect of diet on speciation varied with phylogenetic scale, and that the selective importance of degrees of carnivory and dietary specialization differed among taxonomic groups (Table S2). Notably, degree of carnivory showed a statistically significant relationship with speciation rate when assessed across all of Mammalia (Fig. 4), as well as in five lower clades (Figs. S4-S8). Across mammals, omnivorous species had the lowest rates (Figs. 4 and S3), which is a pattern previously observed in mammals and birds (Price *et al.* 2012; Burin *et al.* 2016). Our findings show that this pattern can extend to clades at lower taxonomic levels (i.e., Artiodactyla + Perissodactyla, Carnivora + Pholidota, Chiroptera, Eulipotyphla, and Marsupialia), but that it does not describe all lower groupings. Conversely, there was no significant relationship between dietary specialization and speciation across Mammalia as a whole, and only Carnivora + Pholidota showed a significant impact of dietary specialization at lower taxonomic levels (Fig. S9). Thus, we find that diet does not have a homogeneous effect on diversification at different hierarchical levels of Mammalia. This is consistent with the scale-dependent effects of other ecological factors that govern mammalian diversification (Machac *et al.* 2017).

Five clades showed no significant diversification relationship with diet. Monotremata comprised only three species in our study, so it was not included in our analyses. Diets within Lagomorpha are relatively homogeneous, with few regime shifts in degree of carnivory or dietary specialization throughout the clade (Fig. S16). Thus, lagomorph speciation was unlikely to be shaped by variation in diet. Members of Atlantogenata occupy the breadth of the carnivory spectrum, but our phylogenetic half-life analyses indicated that diet was a weaker selective force acting in this clade than in other parts of Mammalia (Table S2). Although Euarchonta and Rodentia showed no diet-dependent speciation dynamics, we observed strong selection pressure acting on diet evolution in these clades during our phylogenetic half-life analyses (Table S2). Lack of a relationship between a particular trait and speciation rate does not necessarily indicate that the trait is irrelevant to speciation, and by extension, diversification. While a trait may not be the rate-limiting control, it may still play an important role in the evolution of a clade (Rabosky 2015). Other aspects of an organism's biology may be more important in shaping speciation and diversification. For example, locomotory mode was found to impact diversification of murine rodents, with less specialized modes associated with higher diversification rates (Nations *et al.* 2020).

Previous explanations for decreased diversification of omnivores have focused on the effect of dietary specialization on diversification rate, rather than the effect of degree of carnivory. Specialists show increased competitive ability over generalists when in their optimal environment (Wilson and Yoshimura 1994; Büchi and Vuilleumier 2014). Generalists utilize several resources instead of being most effective at using a single one, so will be excluded from the optimal environments of specialist species (MacArthur and Levins 1964). Such outperformance by specialists should have macroevolutionary costs for omnivores, resulting in low speciation and/or high extinction rates (Burin *et al.* 2016). Specialized species are more prone to allopatric speciation because their narrow niches cause fragmented distributions and isolate formation as environments change (Futuyma and Moreno 1988; Allmon 1992). Specialists also suffer resource limitations more frequently than generalists, so experience stronger directional selection in a changing environment (Vrba 1980, 1987). These explanations may apply to Carnivora + Pholidota, for which dietary generalism was associated with lower rates, and specialization with higher rates. However, in most clades and in Mammalia as a whole, dietary specialization did not have a significant relationship with diversification. As dietary generalism is not directly associated with lower rates in most clades, these suggested explanations are insufficient.

Reduced omnivore diversification is not due to a purely generalism-dependent speciation dynamic. Instead, consumption of both plant and animal matter causes lowered speciation rates in mammals. Future explanations for the role of omnivory in reducing diversification rate should seek to answer how differences in degree of carnivory lead to different speciation rates. Increased herbivore diversification may be explained by primary producers comprising the largest proportion of biomass on Earth (Bar-On *et al.* 2018), suggesting that more herbivorous species can be supported globally. Ecological factors limit diversity and species richness within a clade, and net diversification rate slows down over time in a diversity-dependent manner as this limit is reached (Rabosky 2009). By extension, herbivores should experience a greater net diversification rate than omnivores as it takes them longer to reach the diversity limit imposed on them. Extreme carnivory was also associated with high speciation rates, despite typically being associated with increased extinction (Purvis *et al.* 2000; Estes *et al.* 2011). This is unsurprising given that speciation and extinction rates are often correlated, with traits that promote speciation also increasing extinction risk (Stanley 1990; Greenberg and Mooers 2017). High rates in carnivores may be attributed to

increased diversification of prey species at lower trophic levels, which drives evolution and diversification of predatory species in an upwards adaptive radiation cascade (Brodersen *et al.* 2018; Pontarp 2020).

The relationship between diet and speciation could be indirect. An unmeasured trait co-distributed with diet may be more responsible for shaping mammalian diversification (Beaulieu and O'Meara 2016). Our QuaSSE analyses cannot account for a hidden factor that affects speciation rate when the degree of carnivory or dietary specialization varies. For example, body size is associated with degree of carnivory in mammals, with herbivores having the largest masses and carnivores the smallest (Price and Hopkins 2015). Diversification rates are also higher for smaller species (Martin 2017); thus, body size may be an important factor related to increased carnivore diversification, but may not explain increased diversification across herbivores. Dispersal ability is also correlated with degree of carnivory, with carnivores having greater dispersal distances than omnivores and herbivores (Sutherland *et al.* 2000; Santini *et al.* 2013). Significant associations between dispersal ability and speciation rates have been observed in both mammals and birds (Claramunt *et al.* 2012; Upham *et al.* 2020), but the correlation between trait and rate is negative, with fastest rates in low-vagility species. Thus, the co-distribution of diet and dispersal ability could explain the high herbivore speciation rates observed in our study, but not those of carnivores. If the link between degree of carnivory and speciation is indeed indirect, the patterns we observed may manifest through diet's relationship with multiple traits, potentially with different mechanisms operating in different regions of the phylogeny. We include additional discussion of clade-specific, indirect mechanisms of diet-dependent speciation in our Supporting Information.

Limitations

The specialization score does not consider the full scope of dietary specialization because it is limited by the data used to construct it. The EltonTraits dataset differentiates between vertebrates, invertebrates, fruits, nectar, seeds, and other plant material in each species' diet (Wilman *et al.* 2014), allowing us to construct our coarse estimation of dietary specialization. However, some species display extreme dietary specializations that are inadequately captured by broad food categories. For example, the giant panda is a bamboo specialist, feeding on specific parts of certain bamboo species (Wang *et al.* 2017). Despite this extreme taxonomic specialization, the panda's diet in EltonTraits is coded similarly to species that feed on multiple, varied foodstuffs, such as grasses, lichen, shrubs, and fungi. This is because these foodstuffs all fall into the 'other plant material' category. We may expect more nuanced results for diet-dependent speciation rates if detailed dietary data was used to construct a synthetic specialization score. Thus, while this is the first effort to disentangle specialization from the carnivory-to-herbivory axis, significant refinements to the specialization score could be made, particularly for smaller phylogenies where complete dietary data may be accessible. We further acknowledge that intraspecific variation in niche specialization exists, and degree of individual specialization varies between species (Bolnick *et al.* 2003). However, each species represents a single entry in the EltonTraits dataset, meaning that population specialization cannot be considered. It is possible that intraspecific differences impact mammalian speciation, but this effect is impossible to detect in our analyses. Nevertheless, our findings indicate that dietary specialization at a broader scale does not impact speciation in most mammalian clades.

A recent study demonstrated that birth-death models cannot reliably estimate speciation and extinction rates on extant timetrees (Louca and Pennell 2020). For any diversification scenario,

there are an infinite number of equally likely alternative scenarios. Despite potentially having very different diversification dynamics, these congruent scenarios cannot be distinguished from one another. Because state-dependent speciation and extinction (SSE) models such as QuaSSE are extensions of the birth-death model, it is possible that they do not accurately estimate the relationship between rates and traits. It is important to acknowledge this unidentifiability issue when considering the findings of our study, but also to note that it does not mean our results are inaccurate. We compared a small number of alternative models representing the potential relationship between diet and speciation rate for each clade. This was due necessarily to there only being four possible relationships that QuaSSE is capable of fitting (constant, linear, sigmoidal, and unimodal/humped). By only comparing these, we avoided the problem of there being many congruent models (Morlon *et al.* 2020). Instead, we infer that one of the models from our limited selection fits the data best and is the more plausible scenario of speciation. Additionally, by considering character state evolution, SSE models consider more than just the timing of branching events in lineage-through-time plots as in the models of Louca and Pennell (2020). This may allow SSE models to overcome the unidentifiability issue (Helmstetter *et al.* 2021). In any case, we were less concerned with estimating precise rate values and more interested in characterizing differences in speciation rate as diet changes. By analyzing multiple alternative trees to account for phylogenetic uncertainty and then averaging over the best-fitting models for each tree, we also account for uncertainty in model fit. In doing so, we strengthen our confidence in the accuracy of the diet-dependent speciation dynamics inferred by our analyses.

Continuous vs categorical representations of diet

Our analysis representing diet as a continuous variable was concordant with a previous study where diet was considered a categorical variable (Price *et al.* 2012); specifically, both studies observed that diversification was lowest for omnivores (Fig. 4). However, while converting the EltonTraits data into categorical carnivory bins for validation, we observed that diet-dependent diversification rates varied due to differences in dietary classification (Fig. 5). We were unable to reproduce results from Price *et al.* (2012) using our own categorical variable. Our analysis using the Replicated categorization also found that the highest transition rates between trophic strategies were out of omnivory and into herbivory or carnivory (Fig. S10B). This is the opposite result to Price *et al.* (2012) and to our analysis using the Price categorization, where transitions into omnivory had the highest rates (Fig. S10A). These results do not support the earlier hypothesis that omnivory acts as a macroevolutionary sink across higher vertebrate taxa (Burin *et al.* 2016), instead suggesting that omnivory is a source of diversity. This demonstrates that a lack of standardization in classification can lead to vastly different conclusions regarding the macroevolutionary importance of a trait.

The species assigned to herbivore, carnivore, and omnivore differ between the Replicated and Price categorization methods (Table S3). Specifically, there are fewer herbivores and more omnivores when using EltonTraits data compared to when using the Price categorization. Different methodologies for measuring diet, such as analyses of stomach contents or direct behavioral observation, use different proxies as evidence of food item consumption and may ignore other evidence, leading to varying estimations of diet (e.g., McInnis *et al.* 1983; Shrestha and Wegge 2006; Balestrieri *et al.* 2011). Moreover, different approaches record diet across varying temporal scales, leading to incongruences in diet reconstruction for different species if a dataset uses multiple proxies to obtain its dietary data (Davis and Pineda-Munoz 2016). Use of different and varied sources of diet data by Wilman *et al.* (2014) and Price *et al.* (2012) likely compounded these

incongruences, leading to disagreement regarding each species' diet and subsequently causing discrepancies between the results of our diversification analyses when using the different datasets.

These results highlight the limitation of comparing studies that use different trait categorization methods and are applicable to other macroevolutionary studies where biologically continuous traits are discretized for analyses. Multiple complex, biologically continuous traits have been treated as qualitative variables in analyses of trait-dependent diversification, including morphology (e.g., Fernández-Mazuecos *et al.* 2013; Conway and Olsen 2019), locomotory mode (e.g., Nations *et al.* 2020), and biome or habitat usage (e.g., Schwery *et al.* 2015; Heinicke *et al.* 2017; Menéndez *et al.* 2020). However, methods exist to capture the continuous nature of these complex traits, so their macroevolutionary contributions can be assessed without the introduction of biases associated with variable discretization. The feasibility of such analyses has already been demonstrated for studies of morphology (Onstein *et al.* 2016) and habitat usage (Burbrink *et al.* 2012; Velasco *et al.* 2016). Locomotory mode can be predicted by morphology (Petrović *et al.* 2017; Verde Arregoitia *et al.* 2017), which is itself a continuous trait, meaning that locomotory mode may also be treated as biologically continuous in analyses of trait-dependent diversification. Given our findings, we recommend that future studies of trait-dependent speciation and diversification consider developing methods to capture the continuous nature of their trait of interest.

ACKNOWLEDGEMENTS

We thank the administrators of the University of Memphis High Performance Computing (HPC) facilities for support during this study.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA ACCESSIBILITY STATEMENT

LITERATURE CITED

- Alhajeri, B. H., and Steppan, S. J. 2018. Ecological and ecomorphological specialization are not associated with diversification rates in muroid rodents (Rodentia: Muroidea). *Evolutionary Biology* **45**: 268-286.
- Allmon, W. D. 1992. A causal analysis of stages in allopatric speciation. In: Futuyma, D. and Antonovics, J. eds. *Oxford Survey in Evolutionary Biology*. Vol. 8. New York: Oxford University Press, pp. 219-257.
- Balestrieri, A., Remonti, L., and Prigioni, C. 2011. Assessing carnivore diet by faecal samples and stomach contents: a case study with Alpine red foxes. *Central European Journal of Biology* **6**(2): 283-292.
- Balisi, M., Casey, C., and Van Valkenburgh, B. 2018. Dietary specialization is linked to reduced species durations in North American fossil canids. *Royal Society Open Science* **5**: 171861.
- Bar-On, Y. M., Phillips, R., and Milo, R. 2018. The biomass distribution on Earth. *PNAS* **115**: 6506-6511.
- Bastide, P., Ané, C., Robin, S., and Mariadassou, M. 2018. Inference of adaptive shifts for multivariate correlated traits. *Systematic Biology* **67**(4): 662-680.
- Beaulieu, J. M., and O'Meara, B. C. 2016. Detecting hidden diversification shifts in models of trait-dependent speciation and extinction. *Systematic Biology* **65**(4): 583-601.
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., and Forister, M. L. 2003. The ecology of individuals: incidence and implications of individual specialization. *The American Naturalist* **161**: 1-28.
- Boyles, J. G., and Storm, J. J. 2007. The perils of picky eating: dietary breadth is related to extinction risk in insectivorous bats. *PLoS ONE* **2**(7): e672.
- Brodersen, J., Post, D. M., and Seehausen, O. 2018. Upward adaptive radiation cascades: predator diversification induced by prey diversification. *Trends in Ecology & Evolution* **33**: 59-70.
- Büchi, L., and Vuilleumier, S. 2014. Coexistence of specialist and generalist species is shaped by dispersal and environmental factors. *The American Naturalist* **183**: 612-624.
- Burbrink, F. T., Ruane, S., and Pyron, R. A. 2012. When are adaptive radiations replicated in areas? Ecological opportunity and unexceptional diversification in West Indian dipsadine snakes (Colubridae: Alsophiini). *Journal of Biogeography* **39**: 465-475.
- Burin, G., Kissling, W. D., Guimarães, P. R., Şekercioğlu, Ç. H., and Quental, T. B. 2016. Omnivory in birds is a macroevolutionary sink. *Nature Communications* **7**: 11250.
- Butler, M. A., and King, A. A. 2004. Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *The American Naturalist* **164**(6): 683-695.
- Cantalapiedra, J. L., FitzJohn, R. G., Juhn, T. S., Hernández Fernández, M., DeMiguel, D., Azanza, B., Morales, J. *et al.* 2014. Dietary innovations spurred the diversification of ruminants during the Cenozoic. *Proceedings of the Royal Society B* **281**: 20132746.

526 Claramunt, S., Derryberry, E. P., Remsen, J. V., and Brumfield, R. T. 2012. High dispersal ability
527 inhibits speciation in a continental radiation of passerine birds. *Proceedings of the Royal Society*
528 *B* **279**: 1567-1574.

529 Conway, M., and Olsen, B. J. 2019. Contrasting drivers of diversification rates on islands and
530 continents across three passerine families. *Proceedings of the Royal Society B* **286**: 20191757.

531 Cooper, N., Thomas, G. H., Venditti, C., Meade, A., and Freckleton, R. P. 2016. A cautionary note
532 on the use of Ornstein Uhlenbeck models in macroevolutionary studies. *Biological Journal of the*
533 *Linnean Society* **118**: 64-77.

534 Davis, M., and Pineda-Munoz, S. 2016. The temporal scale of diet and dietary proxies. *Ecology*
535 *and Evolution* **6**(6): 1883-1897.

536 Deméré, T. A., McGowen, M. R., Berta, A., and Gatesy, J. 2008. Morphological and molecular
537 evidence for a stepwise evolutionary transition from teeth to baleen in mysticete whales.
538 *Systematic Biology* **57**: 15-37.

539 Estes, J. A., Terbough, J., Brashares, J. S., Power, M. E., Berger, J., Bond, W. J., Carpenter, S. R.
540 *et al.* 2011. Trophic downgrading of planet Earth. *Science* **333**: 301-306.

541 Fernández-Mazuecos, M., Blanco-Pastor, J. L., Gómez, J. M., and Vargas, P. 2013. Corolla
542 morphology influences diversification rates in bifid toadflaxes (*Linaria* sect. *Versicolores*). *Annals*
543 *of Botany* **112**(9): 1705-1722.

544 FitzJohn, R. G. 2010. Quantitative traits and diversification. *Systematic Biology* **59**: 619-633.

545 FitzJohn, R. G. 2012. Diversitree: comparative phylogenetic analyses of diversification in R.
546 *Methods in Ecology and Evolution* **3**: 1084-1092.

547 Fraser, D., Haupt, R. J., and Barr, W. A. 2018. Phylogenetic signal in tooth wear dietary niche
548 proxies. *Ecology and Evolution* **8**: 5355-5368.

549 Futuyma, D. J., and Moreno, G. 1988. Evolution of ecological specialization. *Annual Review of*
550 *Ecology and Systematics* **19**: 207-233.

551 Greenberg, D. A., and Mooers, A. Ø. 2017. Linking speciation to extinction: diversification raises
552 contemporary extinction risk in amphibians. *Evolution Letters* **1**: 40-48.

553 Hansen, T. F. 1997. Stabilizing selection and the comparative analysis of adaptation. *Evolution*
554 **51**(5): 1341-1351.

555 Hansen, T. F., Pienaar, J., and Orzack, S. H. 2008. A comparative method for studying adaptation
556 to a randomly evolving environment. *Evolution* **62**: 1965-1977.

557 Haywood, S., Simpson, D. M., Ross, G., and Beynon, R. J. 2005. The greater susceptibility of
558 North Ronaldsay sheep compared with Cambridge sheep to copper-induced oxidative stress,
559 mitochondrial damage and hepatic stellate cell activation. *Journal of Comparative Pathology* **133**:
560 114-127.

561 Heinicke, M. P., Jackman, T. R., and Bauer, A. M. 2017. The measure of success: geographic
562 isolation promotes diversification in *Pachydactylus* geckos. *BMC Evolutionary Biology* **17**: 9.

Helmstetter, A. J., Glemin, S., Käfer, J., Zenil-Ferguson, R., Sauquet, H., de Boer, H., Dagallier, L. M. J. *et al.* 2021. Pulled diversification rates, lineages-through-time plots and modern macroevolutionary modelling. *bioRxiv*: 2021.2001.2004.424672.

Ho, L. S. T., and Ané, C. 2014. Intrinsic inference difficulties for trait evolution with Ornstein-Uhlenbeck models. *Methods in Ecology and Evolution* **5**(11): 1133-1146.

Inostroza-Michael, O., Hernández, C. E., Rodríguez-Serrano, E., Avaria-Llautureo, J., and Rivadeneira, M. M. 2018. Interspecific geographic range size-body size relationship and the diversification dynamics of Neotropical furnariid birds. *Evolution* **72**(5): 1124-1133.

Janiak, M. C., Chaney, M. E., and Tosi, A. J. 2018. Evolution of acidic mammalian chitinase genes (*CHIA*) is related to body mass and insectivory in primates. *Molecular Biology and Evolution* **35**: 607-622.

Jiang, P., Josue, J., Li, X., Glaser, D., Li, W., Brand, J. G., Margolskee, R. F. *et al.* 2012. Major taste loss in carnivorous mammals. *PNAS* **109**: 4956-4961.

Jombart, T., and Ahmed, I. 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics* **27**(21): 3070-3071.

Jombart, T., Devillard, S., and Balloux, F. 2010. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genetics* **11**: 94.

Khabbazian, M., Kriebel, R., Rohe, K., and Ané, C. 2016. Fast and accurate detection of evolutionary shifts in Ornstein-Uhlenbeck models. *Methods in Ecology and Evolution* **7**: 811-824.

Knopf, K., Sturman, J. A., Armstrong, M., and Hayes, K. C. 1978. Taurine: an essential nutrient for the cat. *The Journal of Nutrition* **108**: 773-778.

Law, C. J., Duran, E., Hung, N., Richards, E., Santillan, I., and Mehta, R. S. 2018. Effects of diet on cranial morphology and biting ability in musteloid mammals. *Journal of Evolutionary Biology* **31**: 1918-1931.

Liu, Z., Liu, G., Hailer, F., Orozco-terWengel, P., Tan, X., Tian, J., Yan, Z. *et al.* 2016. Dietary specialization drives multiple independent losses and gains in the bitter taste gene repertoire of Laurasiatherian mammals. *Frontiers in Zoology* **13**: 28.

Louca, S., and Pennell, M. W. 2020. Extant timetrees are consistent with a myriad of diversification histories. *Nature* **580**: 502-505.

MacArthur, R., and Levins, R. 1964. Competition, habitat selection, and character displacement in a patchy environment. *PNAS* **51**: 1207-1210.

Machac, A., Graham, C. H., and Storch, D. 2017. Ecological controls of mammalian diversification vary with phylogenetic scale. *Global Ecology and Biogeography* **27**: 32-46.

Martin, R. A. 2017. Body size in (mostly) mammals: mass, speciation rates and the translation of gamma to alpha diversity on evolutionary timescales. *Historical Biology* **29**: 576-593.

McInnis, M. L., Vavra, M., and Krueger, W. C. 1983. A comparison of four methods used to determine the diets of large herbivores. *Journal of Range Management* **36**(3): 302-306.

Menéndez, I., Gómez Cano, A. R., Cantalapiedra, J. L., Peláez-Campomanes, P., Álvarez-Sierra, M. Á., and Hernández Fernández, M. 2020. A multi-layered approach to the diversification of squirrels. *Mammal Review* **51**: 66-81.

Morlon, H., Hartig, F., and Robin, S. 2020. Prior hypotheses or regularization allow inference of diversification histories from extant timetrees. *bioRxiv*: 2020.2007.2003.185074.

Naples, V. L. 1999. Morphology, evolution and function of feeding in the giant anteater (*Myrmecophaga tridactyla*). *Journal of Zoology* **249**: 19-41.

Nations, J. A., Mount, G. G., Morere, S. M., Achmadi, A. S., Rowe, K. C., and Esselstyn, J. A. 2020. Locomotory mode transitions alter phenotype evolution and lineage diversification in an ecologically rich clade of mammals. *Evolution* **75**: 376-393.

Onstein, R. E., Jordan, G. J., Sauquet, H., Weston, P. H., Bouchenak-Khelladi, Y., Carpenter, R. J., and Linder, H. P. 2016. Evolutionary radiations of Proteaceae are triggered by the interaction between traits and climates in open habitats. *Global Ecology & Biogeography* **25**: 1239-1251.

Pagel, M., and Lutzoni, F. 2002. Accounting for phylogenetic uncertainty in comparative studies of evolution and adaptation. In: Lässig, M. and Valleriani, A. eds. *Biological Evolution and Statistical Physics*. Berlin: Springer, pp. 148–161.

Paradis, E., and Schliep, K. 2018. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**: 526-528.

Petrović, T. G., Vukov, T. D., and Kolarov, N. T. 2017. Morphometric ratio analyses: locomotor mode in anurans. *Comptes Rendus Biologies* **340**(4): 250-257.

Pineda-Munoz, S., and Alroy, J. 2014. Dietary characterization of terrestrial mammals. *Proceedings of the Royal Society B* **281**: 20141173.

Pontarp, M. 2020. Ecological opportunity and upward prey-predator radiation cascades. *Scientific Reports* **10**: 10484.

Poore, A. G. B., Ahyong, S. T., Lowry, J. K., and Sotka, E. E. 2017. Plant feeding promotes diversification in Crustacea. *PNAS* **114**: 8829-8834.

Price, S. A., and Hopkins, S. S. B. 2015. The macroevolutionary relationship between diet and body mass across mammals. *Biological Journal of the Linnean Society* **115**: 173-184.

Price, S. A., Hopkins, S. S. B., Smith, K. K., and Roth, V. L. 2012. Tempo of trophic evolution and its impact on mammalian diversification. *PNAS* **109**: 7008-7012.

Purvis, A., Gittleman, J. L., Cowlishaw, G., and Mace, G. M. 2000. Predicting extinction risk in declining species. *Proceedings of the Royal Society B* **267**: 1947-1952.

Rabosky, D. L. 2009. Ecological limits and diversification rate: alternative paradigms to explain the variation in species richness among clades and regions. *Ecology Letters* **12**: 735-743.

Rabosky, D. L. 2010. Extinction rates should not be estimated from molecular phylogenies. *Evolution* **64**(6): 1816-1824.

- Rabosky, D. L. 2015. Reproductive isolation and the causes of speciation rate variation in nature. *Biological Journal of the Linnean Society* **118**: 13-25.
- Rabosky, D. L., and Goldberg, E. E. 2015. Model inadequacy and mistaken inferences of trait-dependent speciation. *Systematic Biology* **64**(2): 340-355.
- Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**(2): 317-223.
- Robbins, C. T., Hagerman, A. E., Austin, P. J., McArthur, C., and Hanley, T. A. 1991. Variation in mammalian physiological responses to a condensed tannin and its ecological implications. *Journal of Mammalogy* **72**: 480-486.
- Rojas, D., Vale, Á., Ferrero, V., and Navarro, L. 2012. The role of frugivory in the diversification of bats in the Neotropics. *Journal of Biogeography* **39**: 1948-1960.
- Santana, S. E., Dumont, E. R., and Davis, J. L. 2010. Mechanics of bite force production and its relationship to diet in bats. *Functional Ecology* **24**: 776-784.
- Santini, L., Di Marco, M., Visconti, P., Baisero, D., Boitani, D., and Rondinini, C. 2013. Ecological correlates of dispersal distance in terrestrial mammals. *Hystrix* **24**(2): 181-186.
- Schneider, F. D. 2018. Package 'traitdataform'. <https://ecologicaltraitdata.github.io/traitdataform/>.
- Schwery, O., Onstein, R. E., Bouchenak-Khelladi, Y., Xing, Y., Carter, R. J., and Linder, H. P. 2015. As old as the mountains: the radiations of Ericaceae. *New Phytologist* **207**: 355-367.
- Shrestha, R., and Wegge, P. 2006. Determining the composition of herbivore diets in the Trans-Himalayan rangelands: a comparison of field methods. *Rangeland Ecology & Management* **59**: 512-518.
- Stanley, S. M. 1990. The general correlation between rate of speciation and rate of extinction: fortuitous causal linkages. In: Ross, R.M. and Allmon, W.D. eds. *Causes of Evolution: A Paleontological Perspective*. Chicago, USA: Chicago University Press, pp. 103-172.
- Sutherland, G. D., Harestad, A. S., Price, K., and Lertzman, K. P. 2000. Scaling of natal dispersal distances in terrestrial birds and mammals. *Conservation Ecology* **4**(1): 16.
- Upham, N. S., Esselstyn, J. A., and Jetz, W. 2019. Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PloS Biology* **17**(12): e3000494.
- Upham, N. S., Esselstyn, J. A., and Jetz, W. 2020. Ecological causes of uneven speciation and species richness in mammals. *bioRxiv*: 504803.
- Uyeda, J. C., and Harmon, L. J. 2014. A novel bayesian method for inferring and interpreting the dynamics of adaptive landscapes from phylogenetic comparative data. *Systematic Biology* **63**(6): 902-918.
- Van Valkenburgh, B. 1988. Trophic diversity in past and present guilds of large predatory mammals. *Paleobiology* **14**: 155-173.

- 672 Van Valkenburgh, B. 2007. Déjà vu: the evolution of feeding morphologies in the Carnivora.
673 *Integrative and Comparative Biology* **47**: 147-163.
- 674 Velasco, J. A., Martínez-Meyer, E., Flores-Villela, O., García, A., Algar, A. C., Köhler, G., and
675 Daza, J. M. 2016. Climatic niche attributes and diversification in *Anolis* lizards. *Journal of*
676 *Biogeography* **43**: 134-144.
- 677 Venables, W. N., and Ripley, B. D. 2002. *Modern Applied Statistics with S*. 4th ed. New York,
678 USA: Springer.
- 679 Verde Arregoitia, L. D., Fisher, D. O., and Schweizer, M. 2017. Morphology captures diet and
680 locomotor types in rodents. *Royal Society Open Science* **4**: 160957.
- 681 Vrba, E. S. 1980. Evolution, species and fossils: How does life evolve? *South African Journal of*
682 *Science* **76**: 61-84.
- 683 Vrba, E. S. 1987. Ecology in relation to speciation rates: some case histories of Miocene-Recent
684 mammal clades. *Evolutionary Ecology* **1**: 283-300.
- 685 Wang, H., Zhong, H., Hou, R., Ayala, J., Liu, G., Yuan, S., Yan, Z. *et al.* 2017. A diet diverse in
686 bamboo parts is important for giant panda (*Ailuropoda melanoleuca*) metabolism and health.
687 *Scientific Reports* **7**: 3377.
- 688 Wiens, J. J., Lapoint, R. T., and Whiteman, N. K. 2015. Herbivory increases diversification across
689 insect clades. *Nature Communications* **6**: 8370.
- 690 Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., and Jetz, W. 2014.
691 EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology* **95**:
692 2027.
- 693 Wilson, D. S., and Yoshimura, J. 1994. On the coexistence of specialists and generalists. *The*
694 *American Naturalist* **144**: 692-707.
- 695 Zhao, H., Yang, J.-R., Xu, H., and Zhang, J. 2010. Pseudogenization of the umami taste receptor
696 gene *Tas1r1* in the giant panda coincided with its dietary switch to bamboo. *Molecular Biology*
697 *and Evolution* **27**: 2669-2673.