White Matter Signals Reflect Information Transmission Between Brain Regions During Seizures

Andrew Y. Revell^{*,1,2}, Alexander B. Silva^{2,3}, Dhanya Mahesh^{1,2}, Lena Armstrong², T. Campbell Arnold^{2,3}, John M. Bernabei^{2,3}, Joel M. Stein^{2,5}, Sandhitsu R. Das^{2,4}, Russell T. Shinohara^{6,7}, Dani S. Bassett^{1,2,3,4,9,10,11,12}, Brian Litt^{2,3,4}, and Kathryn A. Davis^{1,2,4}

¹Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

²Center for Neuroengineering and Therapeutics, University of Pennsylvania, Philadelphia, PA 19104 USA

³Department of Bioengineering, School of Engineering and Applied Science, University of Pennsylvania, Philadelphia, PA 19104 USA

⁴Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

⁵Department of Radiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

⁶Department of Biostatistics, Epidemiology, and Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

⁷Penn Statistics in Imaging and Visualization Endeavor, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

⁸Center for Biomedical Image Computing and Analytics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

⁹Department of Electrical and Systems Engineering, School of Engineering and Applied Science, University of Pennsylvania, Philadelphia, PA 19104 USA

¹⁰Department of Physics and Astronomy, College of Arts and Sciences, University of Pennsylvania, Philadelphia, PA 19104 USA

¹¹Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA

¹²Santa Fe Institute, Santa Fe, NM 87501 USA

*Corresponding author: andrew.revell@pennmedicine.upenn.edu

White matter supports critical brain functions such as learning and memory, modulates the distribution of action potentials, and acts as a relay of neural communication between different brain regions. Interestingly, neuronal cell bodies exist in deeper white matter tissue, neurotransmitter vesicles are released directly in white matter, and white matter blood-oxygenation level dependent (BOLD) signals are detectable across a range of different tasks—all appearing to reflect intrinsic electrical signals in white matter. Yet, such signals within white matter have largely been ignored. Here, we elucidate the properties of white matter signals using intracranial EEG. We show that such signals capture the communication between brain regions and differentiate pathophysiologies of epilepsy. In direct contradiction to past assumptions about white matter *in*-activity, we show that white matter recordings can elucidate brain function and pathophysiology *not* apparent in gray matter. Broadly, white matter functional recordings acquired through implantable devices may provide a wealth of currently untapped knowledge about the neurobiology of disease.

White Matter | Epilepsy | Networks

Introduction

Historically, neuroscientists have “exhibited little interest in white matter”¹ because it was thought to be an inert, passive tissue, despite comprising approximately 50% of the brain’s volume^{2,3}. Instead, investigators have focused on gray matter functional activity because neural computation primarily takes place in gray matter, and gray matter signals can be orders of magnitude more powerful than white matter^{4–6}. Only recently has white matter been appreciated to support many critical brain functions⁷, modulate the distribution of action potentials⁸, and act as a relay of neural communication between different brain regions⁹. A limited number of research groups have shown that neuronal cell bodies exist in deeper white matter tissue (>2–3mm from the cortical layers)¹⁰; neurotransmitter vesicles are released directly in white matter, affecting neural communication¹¹; and white matter blood-oxygenation level dependent (BOLD) signals are detectable across a range of different tasks¹². All of these findings are evidence for intrinsic neural white matter activity. White matter is a dynamic, active tissue and its recordings may reveal important clues to brain function and pathophysiology¹³.

Stereoelectroencephalography (SEEG) is an invasive procedure that captures local field potentials (LFPs)^{15–17}, and provides vital neural information in medically refractory epilepsy patients to aid in localization of the seizure onset zone (SOZ) for its eventual resection, ablation, or modulation^{18,19}. The electrodes are stereotactically inserted into the skull (Fig. 1a).

This implantation modality is now favored in clinical practice because it decreases morbidity and is better tolerated by patients²⁰. Yet, SEEG differs from more traditional implantation modalities, such as electrocorticography (ECoG), creating potential problems for clinical translation of the literature based on ECoG²¹. For example, SEEG provides coverage to deeper brain structures and captures neural activity in many tissue types, including white matter (Fig. 1b).

White matter recordings are ignored both in the clinic and in epilepsy research because no validated studies demonstrate appropriate interpretation and analysis of white matter contacts, despite the fact that approximately half of all SEEG contacts localize to white matter. Clinicians in many epilepsy centers typically exclude white matter contacts when creating intracranial EEG (iEEG) montages to interpret, diagnose, and localize the SOZ. Research studies^{21–25} utilizing SEEG will also typically exclude white matter contacts; their signals may prove problematic when creating computational models sensitive to EEG features if white matter signals are functionally unique. Indeed, analyses on *interictal* recordings of select EEG features have shown differences in white matter (Fig. 1c)^{2,4–6}. Any differences in amplitude, frequency, or timing of activity can be problematic for localizing the SOZ, which is a vital step in achieving eventual seizure freedom in medically refractory epilepsy patients¹⁴.

How should clinicians interpret white matter signals in implanted medically refractory epilepsy patients? Are the recordings merely redundant, noisy signals picked up from nearby

Fig. 1. White matter recordings may elucidate brain function and pathophysiology. | **a**, A CT scan of an epilepsy patient who underwent bilateral stereoelectroencephalography (SEEG) implantation. **b**, SEEG records from both gray matter (GM) and white matter (WM). Recordings can appear visually different along the depth of an SEEG electrode, which can be problematic for localizing the seizure onset zone¹⁴. A power analysis (inset) shows that WM recordings have lower power than GM recordings (interictal, preictal, ictal, postictal). **c**, Power in WM decreases farther from GM tissue (top). WM contacts ≥ 6 mm from GM tissue have lower power during seizures compared to baseline (i.e. interictal power, $p < 5 \times 10^{-5}$) than contacts in GM tissue (bottom). However, power remains elevated *postictally* in WM, while power in GM tissue returns to baseline ($p < 0.05$). **d**, Here, we investigate WM recordings in three parts: (1) assess differences between the signals and networks of GM and WM, (2) understand what drives WM signals, and (3) differentiate good and poor outcome epilepsy patients with WM recordings. SNR, signal-to-noise ratio; ii, interictal; pi, preictal; ic, ictal; po, postictal.

gray matter tissue, as is often thought to be the case? Or do informative white matter signals exist and offer unique insight into neurobiology and epilepsy pathophysiology? Here, we investigate white matter recordings through a multi-pronged approach (Fig. 1d): (1) characterizing white matter and gray matter signal and network differences across distinct brain states, (2) testing the hypothesis⁵ that white matter signals capture the communication between gray matter regions using multimodal structural and functional data, and (3) demonstrating that white matter activity differs across outcomes of epilepsy patients who underwent surgery to remove the SOZ. We provide ample evidence highlighting the importance of white matter signals, which may harbor key information regarding neurobiology, disease, and diagnoses.

Results

The power of white matter iEEG signals is lower during seizures, but remains elevated postictally compared to baseline. A power analysis (Fig. 1b, inset) shows both gray mat-

ter (GM) and white matter (WM) recordings (≥ 2 mm from GM) increase in broadband power from preictal to ictal states ($p < 5 \times 10^{-5}$). GM recordings also have higher power than WM recordings in all peri-ictal states (interictal, preictal, ictal, postictal, $p < 5 \times 10^{-5}$). This observation confirms the intuition that WM recordings have different signal properties. We show that power across all frequency bands decreases in WM tissue as a function of distance from GM tissue (Fig. 1c). When comparing power to baseline (i.e. interictal) power, WM contacts ≥ 6 mm from GM tissue have lower power during seizures ($p < 5 \times 10^{-5}$) than contacts in GM tissue. However, power remains elevated postictally in these WM recordings, while power in GM tissue returns to baseline ($p < 0.05$). This elevation in WM power postictally is evidence that WM recordings are functionally distinct from GM recordings in epilepsy patients. This may also indicate that inter-regional communication, detected by contacts inserted in the white matter communication pathways themselves, is abnormal postictally, even when gray matter activity might return to baseline.

Fig. 2. White matter (WM) correlations increase more than gray matter (GM) during seizures. | a, SEEG with labeled electrodes. b, The high- γ -band cross correlation matrix ordered by electrode labels. Same color as in panel a. Bottom color bar indicates which contacts are GM versus WM. Distribution of FC (functional connectivity) values is plotted. c, Re-ordered adjacency matrix by WM distance. FC distribution is separated by GM or WM values showing WM contributes to the tail. d, Empirical cumulative density function (ECDF) is plotted. ΔT is the difference in the median WM and GM FC values. e, Pairwise Euclidean distance between contacts (left). Contacts closer together have higher FC (right); therefore we use distance between contacts as a covariate in understanding the relationship between WM distance from GM and the increase in FC seen in panel d. f, WM contacts are defined using two definitions (Fig. S2). WM FC increases as a function of the respective WM definition after controlling for contact distance. Each WM definition is significantly related to FC. g, The ECDF plots of WM and GM sub-networks during four states. The distribution of bootstrapped GM and WM FC values are inset. h, ΔT is plotted for each seizure and corresponding peri-ictal state (interictal, preictal, postictal). Error bars are 95% confidence intervals. Mean ΔT for each state of all bootstraps ($N = 10,000$) are plotted. i, Summary of findings: WM correlations are higher than GM, and the increase in those correlations during seizures is greater in WM than in GM.

White matter signals are more correlated with each other than gray matter signals. Next, we compared GM and WM functional connectivity (Fig. 2). We defined functional connectivity (FC) using high- γ -band cross correlation (100-128Hz) in a bipolar montage (see Methods on frequency definitions and montaging). This band along with bipolar montaging represents relatively local activity dynamics that are largely unaffected by volume conduction²⁸. The band is particularly relevant to our study because a well-developed literature implicates high-frequency oscillations and γ -band activity as drivers of epileptic activity^{29,30}.

We localized SEEG contacts (Fig. 2a) to GM or WM (see the tissue localization subsection of the Methods), and plotted the distribution of all FC values (Fig. 2b) during the ictal state. We separated out tissue sub-networks (GM-only

and WM-only) and observed that WM contributes to the tail of the distribution (Fig. 2c). Given that WM has lower power than GM and may record redundant noisy signals⁴, we initially hypothesized that WM FC would be *lower* than GM FC. We tested the null hypothesis that the two sub-network distributions of FC values were identical during the ictal period. Instead, we found that WM contacts have *higher* FC values ($p < 5 \times 10^{-14}$, Fig. 2d) during the ictal state.

Controlling for distances between contacts. Although contacts in medial GM structures (such as the cingulate gyrus or limbic structures) are frequently targeted for bilateral implantation (thus these GM contacts are physically close), it is reasonable to suspect that WM contacts are closer together than GM contacts and may explain the increased FC values.

Fig. 3. White matter (WM) tissue sub-networks have different network properties than gray matter (GM). | **a**, Network visualization of an example patient during the ictal state. WM contacts are ordered clockwise by increasing WM distance from GM contacts. Another example patient during all four states is shown (bottom), illustrating the increase in functional connectivity during the ictal state. **b**, Network measures of density, characteristic path length, and transitivity during each state are plotted. Density and transitivity are higher in WM than GM tissue sub-networks ($p < 0.05$); however, the characteristic path length is not different. WM networks are more dense and clustered. Density and transitivity is higher in WM than in GM sub-networks for the interictal and preictal state. (trending, $p < 0.1$ indicated by an x).

Indeed, contacts closer together have higher FC than contacts further away (Fig. 2e, Spearman Rank Correlation $r = 0.31$, $p < 5 \times 10^{-10}$), even in the high- γ -band after bipolar montaging. Thus, we used contact distance as a covariate. We fit an ordinary least squares regression model to predict FC as a function of the coefficients' contact distance and either (1) distance from GM or (2) WM depth (see WM definitions in methods and Fig. S1 and S2). Briefly, we used two alternative definitions of WM because the limited number of studies analyzing interictal WM SEEG signals also used multiple definitions, including distance to GM and WM depth (depth accounts for the surrounding tissue composition. A higher depth means the contact is deeply embedded within WM and has little surrounding GM tissue)^{5,6}. Alternative definitions have shown similar results when comparing GM and WM signals in previous studies⁶, however, we still provide complimentary analyses using both definitions here. We found that both distance from GM ($p < 5 \times 10^{-10}$) and WM depth ($p < 0.05$) are still significantly related to FC (Fig. 2f). FC increases more in WM even after accounting for the distance between the WM contacts.

White matter correlations increase more than gray matter signals during seizures. Next we show the changes in FC between the four peri-ictal states (Fig. 2g-h). We show through bootstrapping (see the bootstrapping subsection of the Methods) that WM FC is slightly lower than GM FC during the interictal and preictal states, but increases dramatically more than GM during the ictal state ($p < 0.05$, Fig. 2g ECDF plots and inset histograms). The distributions of FC values in WM become similar to those in GM during the postictal state ($p > 0.05$). We test the null hypothesis that the difference in GM and WM median FC values during the ictal state is not zero ($\Delta T_{ic} \neq 0$, Fig. 2h). We found evidence that the median WM FC is higher than the median GM FC ($\Delta T_{ic} > 0$; $p < 0.005$). Although both GM and WM FC increase from the preictal to the ictal states, we also show that the increase is larger for WM ($p < 0.05$). In summary (Fig. 2i), we show that WM FC is higher than GM FC, and that the increase in WM FC from the preictal to the ictal states is larger than in GM FC. We repeat our key results in broadband signals and show that the observed differences are even larger (Fig. S3). The values of ΔT for all bands and for cross correlation, coherence, and Pearson correlations are shown in Fig. S4 and

Fig. 4. White matter (WM) recordings capture the relay of communication between gray matter (GM) regions | **a**, WM tracts are visualized between the left and right hippocampi for an example patient. **b**, Histogram distributions and ECDFs are shown for the different tissue sub-networks similar to Fig. 2c. **c**, Structure-function correlation (SFC) is computed for each sub-network. r = Spearman Rank correlation. **d**, SFC for each state and sub-network is shown (top). Preictal and ictal SFCs are compared. Bootstrapped means ($N = 10,000$) of SFC for each state and each sub-network is shown (bottom). **e**, The distribution of bootstrapped means ($N = 10,000$) of ΔSFC ($SFC_{ictal} - SFC_{preictal}$) is plotted showing ΔSFC is highest for GM-WM connections and lowest for WM-only sub-networks. **f**, Summary of findings: (1) GM functional activity changes during seizures are correlated to other GM regions (and are documented through other studies^{26,27}). (2) WM functional activity changes during seizures are correlated to the structural connectivity from GM regions. (3) WM functional activity changes during seizures have the lowest correlation to the structural connectivity from other WM regions (see confidence intervals in the main text). This observation supports the notion that activity in WM is driven by activity in GM and may be explained by the structural connections between regions. **SFC**, Structure-Function Correlation.

support that these findings are not limited to specific bands or FC definitions.

White matter sub-networks have different network properties than gray-matter sub-networks. As part of our multi-pronged approach, we compared GM and WM signals with a network analysis of the tissue sub-networks (Fig. 3). We measured density, characteristic path length, and transitivity of the full network which includes all contacts, the GM-only sub-network, and the WM-only sub-network. We show that density and transitivity are both higher in WM-only sub-networks than in GM-only sub-networks during the ictal state ($p < 0.05$). Networks composed of only WM contacts are more clustered and have different network properties than networks composed of only GM contacts. This finding is crucial for informing research applying network models in SEEG. It remains to be seen whether the utility for network models excluding or including WM recordings affects clinical translation. However, we show here that functional network properties derived from data of both GM and WM tissue are affected.

White matter recordings capture the communication between gray matter regions. The previous analyses above illustrate the altered signaling and seizure dynamics of GM and WM

recordings: WM signals have increased FC during seizures. We next sought to explain what drives this activity. Does this increased WM FC result from the activity of other GM tissue (for example, from the SOZ), or is the increase in FC intrinsic to WM tissue itself during the ictal state? We applied a structure-function analysis by assessing the statistical correlation between structural and functional connections to determine if the increased FC in WM is explained by the structural connections from GM. If so, we and others hypothesize that WM signals reflect the communication between GM regions^{5,6}. In patients with high angular diffusion imaging (HARDI, $N = 16$), we measured the structural connectivity between contacts (Fig. 4a). We also show the distribution of functional connectivity values in Fig. 4b. We calculated the structure-function correlation (SFC) as a correlation coefficient between the upper triangle of the structural matrix and the upper triangle of the functional matrix, separately for the four tissue networks (Fig. 4c). We show that the SFC increases for the full, GM-only, and GM-WM networks ($p < 0.05$), but not for the WM-only sub-network (Fig. 4d, top). We bootstrapped our data 10,000 times (Fig. 4d, bottom) and calculated the mean SFC for each simulation. We also calculated the change in SFC (ΔSFC) from preictal

Fig. 5. Good outcome epilepsy patients have lower white matter (WM) functional connectivity than poor outcome patients. | a, Diagram summarizing results for the top row: WM and GM FC increase during seizures for both good and poor outcome patients. WM FC increases more than GM FC (see Fig. 2i), and this increase is even larger for poor outcome patients. b, Empirical cumulative density functions (ECDF) are shown for the sub-networks in poor and good outcome patients. FC is higher during the ictal state for poor outcome patients in the full, GM-WM, and WM-only tissue sub-networks. However, FC is not significantly different for the GM-only tissue sub-network. c, Differences in GM and WM FC values (ΔT) during the ictal period are larger for poor outcome patients than for good outcome patients, consistent with Fig. 2h. Error bars represent 95% confidence intervals. d, Re-sampling hypothesis testing (top) confirming the analysis of Fig. 5c (see the re-sampling hypothesis testing subsection of the Methods). Dashed vertical line: T . The distribution of bootstrapped means of the ictal ΔT is also plotted comparing good and poor outcome patients (bottom). e, Diagram summarizing results for the bottom row: ablated regions in good outcome patients have lower changes in FC to distant WM contacts than poor outcome patients during seizures. f, ECDF of the FC values between the contacts in the ablation zone and all WM contacts (left). The median FC values for each re-sampled patient is compared (right). g, Same as in panel d but now comparing the changes in FC between ablated and WM contacts. h, The FC values between the contacts in the ablation zone and WM contacts as a function of distance. We show that the FC of ablated contacts to distant WM contacts increases more during seizures for poor than good outcome patients. T^* , test statistic of permuted samples; T , test statistic of original data.

to ictal states ($\Delta SFC = SFC_{ic} - SFC_{pi}$) and plot the distributions of ΔSFC over 10,000 bootstrap simulations (Fig. 4e). The SFC increases the most from preictal to ictal states for GM-WM sub-networks and the least for WM-only sub-networks (ΔSFC means with 95% confidence intervals: Full, 0.0537 [0.0529-0.0544]; GM-only, 0.0525 [0.0516-0.0533]; GM-WM, 0.0771 [0.0758-0.0783]; WM-only, 0.0312 [0.0299-0.0324]). These results support the hypothesis that the change in WM at seizure onset is driven by the structural connections from GM regions (e.g., the SOZ or other regions involved in seizure activity, Fig. 4f).

Surgical epilepsy patients with poor outcomes have higher white matter connectivity than patients with good outcomes. A total of 17 patients had outcome scores ≥ 2 years after epilepsy surgery to remove the SOZ (good outcome $N = 10$; poor outcome $N = 7$). We show that poor outcome patients have a stronger increase in WM FC during the ictal state (see overview in Fig. 5a). During the ictal state, the full, GM-WM, and WM-only sub-networks have higher connectivity ($p < 0.05$) in poor outcome patients than in good outcome patients (Fig. 5b). This increase is not apparent in the GM-only sub-network, which is the sole network available for patients that only have

GM implanted or for analyses that only consider GM contacts ($p > 0.05$). We repeated the analysis in Fig. 2h, separating good and poor outcome patients (Fig. 5c). We show that the differences between GM and WM sub-networks during the ictal period become more pronounced for poor outcome patients than for good outcome patients using re-sampling hypothesis testing ($p < 0.05$, Fig. 5d, top) and bootstrapping 10,000 times the mean ΔT_{ictal} of good and poor outcome patients (Fig. 5d, bottom).

Ablated regions in good outcome patients have lower connectivity to white matter signals than poor outcome patients. In both good and poor outcome patients, the ablation area was targeted because its removal would give the best chance of seizure freedom despite the surgery's inherent risk. Surgery was performed after a comprehensive clinical assessment of data acquired from multiple data and imaging modalities including many MRI sequences, scalp EEG, iEEG, patient and family histories, and neuropsychological evaluation. We showed in Fig. 5 (top row) that differences in FC of good and poor outcomes are mainly driven by connections to WM, but not apparent in GM-only regions. We wanted to test if the FC between ablated regions and WM contacts could differentiate

between good and poor outcome patients (Fig. 5e). We show that FC between the ablated regions and WM contacts is higher for poor outcome patients than good outcome patients ($p < 0.05$, Fig. 5f-g). This finding may indicate that the suspected SOZ in poor outcome patients (which would eventually be ablated) is not a good target given that its activity has a high correlation with the activity of WM contacts.

White matter connectivity changes during seizures are elevated far from the ablation zone in poor outcome patients.

We next sought to determine whether the correlations remain elevated in WM contacts distant from the ablation site in poor outcome patients but not in good outcome patients (Fig. 5h). After bootstrap re-sampling of patients and seizures, we show that WM FC to the contacts contained in the *eventual* ablation zone remains elevated in poor outcome patients. This increase is most apparent for WM contacts 40–60 mm (approximately a half hemisphere away) and 80–100 mm (approximately on the contralateral hemisphere) from the ablation zone. Future analyses should further contextualize the location of these WM contacts to canonical WM pathways.

Discussion

We demonstrated that white matter intracranial EEG (iEEG) recordings reveal underlying seizure dynamics that are not captured in gray matter recordings alone. We compared gray matter and white matter recordings from stereoelectroencephalography (SEEG) implantations of 29 patients and found differences in signal and network properties of white matter contacts (Figs. 1-3). We showed that white matter contacts capture neural communication along the structural pathways connecting brain regions in 16 of the SEEG patients who acquired high angular resolution diffusion imaging (HARDI, Fig. 4). These white matter recordings also have different seizure dynamics in good and poor outcome epilepsy patients at distant sites from the ablated regions (Fig. 5). Our study provides evidence that the analysis of white matter functional signals may prove worthwhile when researching and developing new diagnostic and treatment strategies of medically refractory epilepsy patients. More broadly, white matter functional signals may also reveal important clues to brain function and pathophysiology of other neurological disorders. To facilitate reproducibility and further investigation of these SEEG patients, we have made the iEEG data, metadata, and de-identified processed structural tractography neuroimaging data compliant with Brain Imaging Data Structure (BIDS) and publicly available (see Methods and 'Data availability and reproducibility').

Hidden biology revealed in white matter. Both in the clinic and in epilepsy research, implantations in white matter are ignored. Clinicians in many epilepsy centers typically exclude white matter contacts when creating iEEG montages to interpret, diagnose, and localize the seizure onset zone (SOZ). Research studies^{21–25} utilizing SEEG will typically exclude white matter contacts. The exclusion of white matter contacts is perhaps not unexpected given the fact that no validated studies exist that demonstrate a rigorous approach to incorporating white matter contacts.

Across neuroscience, white matter functional activity is also largely ignored¹³. Historically, white matter was thought to

be an inert, passive tissue and, thus, neuroscientists “exhibited little interest in white matter”¹. Only a limited number of studies over the last 20 years from a small cohort of research groups explore the nature of white matter activity through functional MRI (fMRI)^{12,31–39}. The focus, instead, has been on the analysis of gray matter functional activity as it relates to neurobiology and disease, because the signals can be orders of magnitude larger (Fig. 1b-c)^{4–6}, and gray matter is predominantly thought to be where neural computation takes place. Excitingly, some research groups have shown the existence of neuronal cell bodies in deeper white matter tissue¹⁰, the release of neurotransmitter vesicles in white matter¹¹, and the existence of detectable white matter blood-oxygenation level dependent (BOLD) signals across a range of different tasks¹². These facts appear to reflect intrinsic neural white matter activity. These findings raise several questions. What are the functions of white matter neurons in epilepsy? What are the consequences of altered neural communication resulting from the release of neurotransmitters along the axons of white matter? Can white matter activity recorded by fMRI, iEEG, and other technologies reveal clues to epilepsy pathophysiology and, more generally, neurophysiology?

Here in our study, we have provided examples of white matter’s utility for neuroscience inquiry through a multi-pronged approach (from signals to networks) across many brain states (from interictal to postictal) and with multimodal data (from structure to function) in different outcomes of epilepsy patients (from good to poor). We provide ample evidence highlighting the importance of white matter signals, which could be further investigated in future studies to reveal novel insights into neurobiology, disease, and diagnoses.

The Distributed Epileptic Network Hypothesis. We provided evidence that white matter recordings reveal differences in epilepsy pathophysiology of good and poor outcome epilepsy patients not apparent in gray matter recordings (Fig. 5). Poor outcome patients have higher FC between the *suspected* (and eventually ablated) SOZ and more distant white matter recordings than good outcome patients (Fig. 5h). We attribute these differences to the potential non-focal seizure pathophysiology of poor outcome patients. The areas ablated in poor outcome patients were chosen because of supporting clinical evidence that the primary target for ablation was a well-localized, focal SOZ; otherwise, patients would not have been ideal candidates for surgery. Yet, patients continue having seizures after ablation. The inefficacy of the surgery could be due to two factors: (1) the SOZ is still a focal region of the pathology, but was missed in the ablation, or (2) the SOZ appears focal to clinicians, but the underlying pathophysiology in some patients is more distributed across brain regions and interconnecting white matter⁴⁰. This latter factor can occur when a patient does not have a focal pathophysiology, but one that stems from an “epileptogenic”⁴¹ or “epileptic”²¹ network.

The traditional model of the epileptogenic focus is too simplistic to capture the spatiotemporal organization of seizures⁴². Thus, many investigators have begun to apply complex system approaches and network models to understand the underlying neurophysiological mechanisms in epilepsy^{26,43–49}. Here, we provided evidence for the epileptic network captured in the white matter of poor outcome patients. We build on the concept of the epileptic network to explain *why* some

patients have poor outcomes. We hypothesize that, in these patients, the true SOZ is not a single focal anatomical location, but rather a *distributed epileptic network* where many brain regions interact, causing patient's neural dynamics to bifurcate⁵⁰ or spontaneously devolve into seizure states. Here, we take the opportunity to name the hypothesis we proposed in the previous sentence the “**Distributed Epileptic Network Hypothesis**” to provide concrete language for discussing the pathophysiology of some epilepsy patients, in hopes of spurring new areas of research and therapeutic treatment options for these patients. In these poor outcome patients, important nodes of the epileptic network were eliminated during surgery; thus, they may be considered Engel III with “marked seizure reduction” and reduced disease burden. However, seizures eventually reoccur because the epileptic network was not entirely hindered. Instead, only select important nodes were eliminated. The network then reacts or it reforms to produce seizures, albeit not as strong or as frequent in Engel II/III. Here in our paper, we show evidence for a distributed epileptic network in poor outcome patients that is captured in white matter recordings.

Limitations. Our study is not without limitations. They include: (1) The choice of montaging and frequency bands. We attempted to extract the intrinsic white matter signals or other signals coming from distant brain areas, and not from volume conduction of nearby gray matter. Regardless, volume conduction from other confounding extrinsic sources cannot be eliminated completely and may increase FC between certain brain regions, thereby affecting our findings. (2) A small sample size of poor outcome patients and definitions of outcomes. We aimed to mitigate the biases of patients, different implantation schemes, and different tissue sampling techniques through bootstrapping, another approach using re-sampling hypothesis testing, and more stringent exclusion criteria of outcomes ≤ 2 years. (3) Seizure temporal dynamics. We computed FC over entire segments of interictal, preictal, ictal, and postictal states. Seizure dynamics are not the same between early and late stages of seizure evolution. FC differences between white matter/gray matter and good/poor outcome patients could lead to a different interpretation when involving more complex time-dynamical analyses. (4) White matter location. We did not include an analysis of the anatomical structures of white matter contacts or test if specific structures are mainly driving white matter FC (possibly supported by Fig. 5h). For example, we did not analyze the corpus callosum or corona radiata fibers that are typically captured in SEEG implantations along the trajectory to deeper gray matter targets. Note that the limited research of WM activity in fMRI has focused on the corpus callosum³¹. (5) Sampling technology. Other implantable devices such as microelectrode-arrays (MEAs) or Utah arrays⁵¹, which can record at higher densities, altered to target white matter may provide more robust findings about white matter signals. In contrast, SEEG records LFPs at the meso-scale.

Future Directions. White matter functional recordings acquired through many technologies (including iEEG, fMRI, MEG, PET, ASL, and others) may provide a wealth of *currently untapped* knowledge about neurobiology and disease. Our work has ramifications across the field of neuroscience and

highlights the possibility of new areas to explore: stimulation of white matter to modulate brain function; biomarkers for monitoring white matter functional signals in the progression, and evolution of, other neurological and psychiatric diseases such as multiple sclerosis, dementia, and schizophrenia; the utility of targeting white matter structures directly, rather than coincidentally acquiring signals in targeting gray matter structures; direct ablation or surgical intervention of white matter areas to improve outcomes, rather than focusing on gray matter structures only; drug therapies that focus on the modulation of abnormal white matter activity, rather than using gray matter activity benchmarks. More research is needed to assess the viability of all these options. The focus of treatment strategies can be on white matter, too.

References

- [1] R.DouglasFields. White matter. *SCIENTIFIC AMERICAN* 54–61 (2008).
- [2] Li, G. *et al.* Detection of human white matter activation and evaluation of its function in movement decoding using stereo-electroencephalography (seeg). *Journal of neural engineering* **18** (2021).
- [3] Fields, R. D. White matter in learning, cognition and psychiatric disorders. *Trends in neurosciences* **31**, 361–370 (2008).
- [4] Greene, P., Li, A., González-Martínez, J. & Sarma, S. V. Classification of stereo-EEG contacts in white matter vs. gray matter using recorded activity. *Frontiers in Neurology* **11** (2021).
- [5] Mercier, M. R. *et al.* Evaluation of cortical local field potential diffusion in stereotactic electroencephalography recordings: A glimpse on white matter signal. *NeuroImage* **147**, 219–232 (2017).
- [6] Young, J. J. *et al.* Quantitative signal characteristics of electrocorticography and stereoelectroencephalography: The effect of contact depth. *Journal of clinical neurophysiology : official publication of the American Electroencephalographic Society* **36**, 195–203 (2019).
- [7] Bengtsson, S. L. *et al.* Extensive piano practicing has regionally specific effects on white matter development. *Nature neuroscience* **8**, 1148–1150 (2005).
- [8] Sigaard, R. K., Kjaer, M. & Pakkenberg, B. Development of the cell population in the brain white matter of young children. *Cerebral cortex (New York, N.Y. : 1991)* **26**, 89–95 (2016).
- [9] Mollink, J. *et al.* The spatial correspondence and genetic influence of interhemispheric connectivity with white matter microstructure. *Nature neuroscience* **22**, 809–819 (2019).
- [10] García-Marín, V., Blázquez-Llorca, L., Rodríguez, J. R., González-Soriano, J. & DeFelipe, J. Differential distribution of neurons in the gyral white matter of the human cerebral cortex. *The Journal of comparative neurology* **518**, 4740–4759 (2010).
- [11] Kukley, M., Capetillo-Zarate, E. & Dietrich, D. Vesicular glutamate release from axons in white matter. *Nature Neuroscience* **10**, 311–320 (2007).
- [12] Li, M., Newton, A. T., Anderson, A. W., Ding, Z. & Gore, J. C. Characterization of the hemodynamic response function in white matter tracts for event-related fMRI. *Nature communications* **10**, 1140 (2019).
- [13] Grajauskas, L. A., Frizzell, T., Song, X. & D'Arcy, R. C. N. White matter fMRI activation cannot be treated as a nuisance regressor: Overcoming a historical blind spot. *Frontiers in Neuroscience* **13** (2019).
- [14] Chauvel, P., Rheims, S., McGonigal, A. & Kahane, P. French guidelines on stereoelectroencephalography (seeg): Editorial comment. *Neurophysiologie clinique = Clinical neurophysiology* **48**, 1–3 (2018).
- [15] Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and currents—eeg, ecog, lfp and spikes. *Nature reviews. Neuroscience* **13**, 407–420 (2012).
- [16] Herreras, O. Local field potentials: Myths and misunderstandings. *Frontiers in neural circuits* **10**, 101 (2016).
- [17] Kajikawa, Y. & Schroeder, C. How local is the local field potential. *Neuron* **72**, 847–858 (2011).
- [18] Cardinale, F. *et al.* Stereoelectroencephalography: retrospective analysis of 742 procedures in a single centre. *Brain : a journal of neurology* (2019).
- [19] Panzica, F., Varotto, G., Rotondi, F., Spreafico, R. & Franceschetti, S. Identification of the epileptogenic zone from stereo-EEG signals: A connectivity-graph theory approach. *Frontiers in Neurology* **4**, 6–11 (2013).
- [20] Mullin, J. P. *et al.* Is seeg safe? a systematic review and meta-analysis of stereoelectroencephalography-related complications. *Epilepsia* **57**, 386–401 (2016).
- [21] Bernabei, J. M. *et al.* Electrocorticography and stereo EEG provide distinct measures of brain connectivity: implications for network models. *Brain Communications* **3** (2021).
- [22] Li, G. *et al.* Optimal referencing for stereo-electroencephalographic (seeg) recordings. *NeuroImage* **183**, 327–335 (2018).
- [23] Ter Wal, M. *et al.* Characterization of network structure in stereoEEG data using consensus-based partial coherence. *NeuroImage* **179**, 385–402 (2018).
- [24] Li, A. *et al.* Neural fragility as an EEG marker of the seizure onset zone. *Nature neuroscience* (2021).
- [25] Frauscher, B. *et al.* Atlas of the normal intracranial electroencephalogram: neurophysiological awake activity in different cortical areas. *Brain : a journal of neurology* **141**, 1130–1144 (2018).
- [26] Shah, P. *et al.* Characterizing the role of the structural connectome in seizure dynamics. *Brain : a journal of neurology* (2019).

- [27] Ashourvan, A. *et al.* Pairwise maximum entropy model explains the role of white matter structure in shaping emergent co-activation states. *Commun Biol* **4** (2021).
- [28] Khambhati, A. N., Davis, K. A., Lucas, T. H., Litt, B. & Bassett, D. S. Virtual cortical resection reveals push-pull network control preceding seizure evolution. *Neuron* **91**, 1170–1182 (2016).
- [29] Warren, C. P. *et al.* Synchrony in normal and focal epileptic brain: the seizure onset zone is functionally disconnected. *Journal of neurophysiology* **104**, 3530–3539 (2010).
- [30] Worrell, G. A. *et al.* High-frequency oscillations in human temporal lobe: simultaneous micro- and clinical macroelectrode recordings. *Brain : a journal of neurology* **131**, 928–937 (2008).
- [31] Gawryluk, J. R., Mazerolle, E. L. & D'Arcy, R. C. N. Does functional mri detect activation in white matter? a review of emerging evidence, issues, and future directions. *Frontiers in neuroscience* **8**, 239 (2014).
- [32] Gore, J. C. *et al.* Functional mri and resting state connectivity in white matter - a mini-review. *Magnetic resonance imaging* **63**, 1–11 (2019).
- [33] Tettamanti, M. *et al.* Interhemispheric transmission of visuomotor information in humans: fmri evidence. *Journal of neurophysiology* **88**, 1051–1058 (2002).
- [34] Mazerolle, E. L. *et al.* Confirming white matter fmri activation in the corpus callosum: Colocalization with dti tractography. *NeuroImage* **50**, 616–621 (2010).
- [35] Schlösser, R. G. M. *et al.* White matter abnormalities and brain activation in schizophrenia: a combined dti and fmri study. *Schizophrenia research* **89**, 1–11 (2007).
- [36] Özbay, P. S. *et al.* Contribution of systemic vascular effects to fmri activity in white matter. *NeuroImage* **176**, 541–549 (2018).
- [37] Gawryluk, J. R., Mazerolle, E. L., Beyea, S. D. & D'Arcy, R. C. N. Functional mri activation in white matter during the symbol digit modalities test. *Frontiers in human neuroscience* **8**, 589 (2014).
- [38] Fabri, M., Polonara, G., Mascioli, G., Salvolini, U. & Manzoni, T. Topographical organization of human corpus callosum: an fmri mapping study. *Brain research* **1370**, 99–111 (2011).
- [39] Aramaki, Y., Honda, M., Okada, T. & Sadato, N. Neural correlates of the spontaneous phase transition during bimanual coordination. *Cerebral cortex (New York, N.Y. : 1991)* **16**, 1338–1348 (2006).
- [40] Bernhardt, B. C., Bonilha, L. & Gross, D. W. Network analysis for a network disorder: The emerging role of graph theory in the study of epilepsy. *Epilepsy & behavior : E&B* **50**, 162–170 (2015).
- [41] Bartolomei, F. *et al.* Defining epileptogenic networks: Contribution of seed and signal analysis. *Epilepsia* **58**, 1131–1147 (2017).
- [42] Davis, K., Jirsa, V. & Schevon, C. Wheels within wheels: Theory and practice of epileptic networks. *Epilepsy Curr* 15357597211015663 (2021).
- [43] Kramer, M. A. *et al.* Coalescence and fragmentation of cortical networks during focal seizures. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **30**, 10076–10085 (2010).
- [44] Rosch, R. E., Hunter, P. R., Baldeweg, T., Friston, K. J. & Meyer, M. P. Calcium imaging and dynamic causal modelling reveal brain-wide changes in effective connectivity and synaptic dynamics during epileptic seizures. *PLoS computational biology* **14**, e1006375 (2018).
- [45] Sinha, N. *et al.* Predicting neurosurgical outcomes in focal epilepsy patients using computational modelling. *Brain : a journal of neurology* **140**, 319–332 (2017).
- [46] Scheid, B. *et al.* Time-evolving controllability of effective connectivity networks during seizure progression. *Proc Natl Acad Sci U S A* **118** (2021).
- [47] Girardi-Schappo, M. *et al.* Altered communication dynamics reflect cognitive deficits in temporal lobe epilepsy. *Epilepsia* (2021).
- [48] An, S., Bartolomei, F., Guye, M. & Jirsa, V. Optimization of surgical intervention outside the epileptogenic zone in the virtual epileptic patient (vep). *PLoS computational biology* **15**, e1007051 (2019).
- [49] Chang, W.-C. *et al.* Loss of neuronal network resilience precedes seizures and determines the ictogenic nature of interictal synaptic perturbations. *Nature neuroscience* **21**, 1742–1752 (2018).
- [50] Saggio, M. *et al.* A taxonomy of seizure dynamotypes. *Elife* **9** (2020).
- [51] Schevon, C. A. *et al.* Evidence of an inhibitory restraint of seizure activity in humans. *Nature Communications* **3** (2012).
- [52] A multimodal platform for cloud-based collaborative research (IEEE, 2013).
- [53] Kini, L. G., Davis, K. A. & Wagenaar, J. B. Data integration: Combined imaging and electrophysiology data in the cloud. *NeuroImage* **124**, 1175–1181 (2016).
- [54] Revell, A. Y. *et al.* A framework for brain atlases: Lessons from seizure dynamics (2021).
- [55] Litt, B. *et al.* Epileptic seizures may begin hours in advance of clinical onset. *Neuron* **30**, 51–64 (2001).
- [56] Azarion, A. A. *et al.* An open-source automated platform for three-dimensional visualization of subdural electrodes using ct-mri coregistration. *Epilepsia* **55**, 2028–2037 (2014).
- [57] Makin, J. G., Moses, D. A. & Chang, E. F. Machine translation of cortical activity to text with an encoder-decoder framework. *Nature neuroscience* **23**, 575–582 (2020).
- [58] SciPy. C. *et al.* Scipy 1.0: fundamental algorithms for scientific computing in python. *Nature Methods Nat Methods* **17**, 261–272 (2020).
- [59] Newson, J. J. & Thiagarajan, T. C. Eeg frequency bands in psychiatric disorders: A review of resting state studies. *Frontiers in Human Neuroscience* **12**, 521 (2019).
- [60] Cieslak, M. *et al.* Qsiprep: An integrative platform for preprocessing and reconstructing diffusion mri. *Biorxiv* (2020).
- [61] Fang-Cheng, Y., Wedeen, V. J. & Tseng, W.-Y. I. Generalized q-sampling imaging. *IEEE Transactions on Medical Imaging* **29**, 1626–1635 (2010).
- [62] Otsu, N. A threshold selection method from gray-level histograms. *IEEE Transactions on Systems, Man, and Cybernetics* **9**, 62–66 (1979).
- [63] Zalesky, A. *et al.* Whole-brain anatomical networks: does the choice of nodes matter? *Neuroimage* **50**, 970–983 (2010).
- [64] Tzourio-Mazoyer, N. *et al.* Automated anatomical labeling of activations in spm using a macroscopic anatomical parcellation of the mni mri single-subject brain. *NeuroImage* **15**, 273–289 (2002).
- [65] Rolls, E. T., Joliot, M. & Tzourio-Mazoyer, N. Implementation of a new parcellation of the orbitofrontal cortex in the automated anatomical labeling atlas. *NeuroImage* **122**, 1–5 (2015).
- [66] Rolls, E. T., Huang, C.-C., Lin, C.-P., Feng, J. & Joliot, M. Automated anatomical labelling atlas 3. *NeuroImage* **206**, 116189 (2020).
- [67] Wirsich, J. *et al.* Whole-brain analytic measures of network communication reveal increased structure-function correlation in right temporal lobe epilepsy. *NeuroImage. Clinical* **11**, 707–718 (2016).
- [68] Bonilha, L., Gleichgerrcht, E., Nesland, T., Rorden, C. & Fridriksson, J. Gray matter axonal connectivity maps. *Frontiers in Psychiatry* **6** (2015).
- [69] Park, B., Eo, J. & Park, H.-J. Structural brain connectivity constrains within-a-day variability of direct functional connectivity. *Frontiers in Human Neuroscience* **11**, 408 (2017).
- [70] Taylor, P. N. *et al.* The impact of epilepsy surgery on the structural connectome and its relation to outcome. *NeuroImage. Clinical* **18**, 202–214 (2018).
- [71] Shah, P. *et al.* Structural and functional asymmetry of medial temporal subregions in unilateral temporal lobe epilepsy: A 7t mri study. *Human brain mapping* **40**, 2390–2398 (2019).
- [72] Engel, J. Update on surgical treatment of the epilepsies. summary of the second international palm desert conference on the surgical treatment of the epilepsies (1992). *Neurology* **43**, 1612–1617 (1993).
- [73] Kini, L. G. *et al.* Virtual resection predicts surgical outcome for drug-resistant epilepsy. *Brain : a journal of neurology* **142**, 3892–3905 (2019).
- [74] Steinbart, D., Steinbrenner, M., Oltmanns, F. & Holtkamp, M. Prediction of seizure freedom after epilepsy surgery - critical reappraisal of significance of intracranial eeg parameters. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology* **131**, 2682–2690 (2020).
- [75] Alhilani, M. *et al.* Ictal and interictal source imaging on intracranial eeg predicts epilepsy surgery outcome in children with focal cortical dysplasia. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology* **131**, 734–743 (2020).
- [76] Knowlton, R. C. *et al.* Functional imaging: II. prediction of epilepsy surgery outcome. *Annals of neurology* **64**, 35–41 (2008).
- [77] Tomlinson, S. B., Porter, B. E. & Marsh, E. D. Interictal network synchrony and local heterogeneity predict epilepsy surgery outcome among pediatric patients. *Epilepsia* **58**, 402–411 (2017).
- [78] Kuroda, N. *et al.* Objective interictal electrophysiology biomarkers optimize prediction of epilepsy surgery outcome. *Brain communications* **3**, icab042 (2021).
- [79] James, G., Witten, D., Hastie, T. & Tibshirani, R. *An Introduction to Statistical Learning* (Springer, 2021).

Materials and Methods

Human Dataset. Twenty-nine drug-resistant epilepsy patients underwent stereoelectroencephalography (SEEG) implantation for clinical purposes at the Hospital of the University of Pennsylvania. Of these patients, 88 seizures were recorded, with distributions of seizures per patient and seizure lengths shown in Fig. S5. Sixteen of the SEEG patients had MRI data collected for SFC analyses. Inclusion criteria consisted of all individuals who agreed to participate in our research scanning protocol, and allowed their de-identified intracranial EEG (iEEG) data to be publicly available for research purposes on the International Epilepsy Electrophysiology Portal (<https://www.ieeg.org>)^{52,53}. Seizure evaluation was determined via comprehensive clinical assessment, which included multimodal imaging, scalp and intracranial video-EEG monitoring, and neuropsychological testing. This study was approved by the Institutional Review Board of the University of Pennsylvania, and all subjects provided written informed consent prior to participating. See Table S1 for subject demographics. Sample sizes were determined by attempting to gather as many patients as possible with validated outcome scores ≥ 2 years, annotated seizure markings, and high angular resolution diffusion imaging (HARDI). No statistical methods were used to predetermine sample sizes, but our sample sizes are larger than those reported in previous publications using both iEEG and diffusion imaging data^{26,27,54}.

Intracranial EEG Acquisition. Stereotactic depth electrodes were implanted in patients based on clinical necessity. Continuous SEEG signals were obtained for the duration of each patient's stay in the epilepsy monitoring unit. Intracranial data was recorded at 256, 512, or 1024 Hz for each patient. Seizure onset times were defined by the unequivocal electrographic onset (UEO)⁵⁵. All annotations were verified by neurologists and consistent with detailed clinical documentation.

Electrode Localization. See Fig. S1 for an overview of electrode localization. In-house software⁵⁶ was used to assist in localizing electrodes after registration of pre-implant and post-implant images (T1w and CT images). All electrode coordinates and labels were

saved and matched with the electrode names on IEEG.org. All electrode localizations were verified by a board-certified neuroradiologist (J.S.).

Tissue segmentation. See Fig. S1 for an overview of tissue segmentation. Freesurfer recon-all tissue segmentation output was used. The pre-implant T1w image specified in the imaging protocol below was used for tissue segmentation. If no T1w image from this protocol was acquired, the T1w image from the pre-implant clinical scan was used. The tissue segmentation (the "aseg" file output from Freesurfer) file which includes cortical GM, subcortical GM, WM, and other anatomical labels was converted to 4 classes: GM, WM, CSF, and outside. All cortical and subcortical GM structures were combined to a single GM class. WM structures were all combined to a single WM class. Voxels contained within the Freesurfer brain extraction file, but outside GM or WM structures, were labeled as CSF. All other voxels not in WM, GM, or inside the brain extraction file were labeled as outside the cranium.

Contact tissue localization. The post-implant T1w image used for electrode localization underwent linear registration to the pre-implant T1w image used for tissue segmentation above. Linear registration was performed using *fniirt* in the FSL neuroimaging software. Contacts were classified to the tissue classes defined in the section below. Only contacts where the center coordinate fell into GM or WM labeled voxels were used. CSF labeled contacts, which generally included those between the skull and brain, and all other contacts were excluded from analysis. Note that for tissue localization purposes, implanted contacts were rarely localized to non-mesial temporal subcortical structures (i.e. the thalamus, putamen, etc.). These structures are rarely targeted in epilepsy patients. Thus accurate tissue segmentation of these subcortical structures were not vital for our study.

GM and WM definitions by distance or depth. Contacts localized to WM voxels may not necessarily record exclusively from WM, especially when neighboring voxels are GM cortical structures. This was addressed in Mercier et al. 2017 using a measure called proximal tissue density (PTD)⁵. This "fuzzy" approach is used to (1) mitigate volume conduction (2) mitigate the degree of uncertainty in the location of the contact (can be due to small errors in registration). Here, we follow a similar fuzzy approach to define GM and WM contacts. See Fig. S2 for an overview.

Definitions:

1. **Distance:** Distance to the nearest GM voxel. Typical range 0-10mm. Contacts that were 0mm away (i.e. the contact centroid coordinate was in a GM voxel) were labeled as GM. Analyses that relied on a definition of distance used the following criteria:
 - (a) GM: = 0 mm
 - (b) WM: ≥ 2 mm
 - (c) contacts between 0 and 2mm were excluded from analysis comparing GM and WM. This follows the Mercier et al. 2017⁵ approach where PTD the extreme values of -1 and 1 were used.
2. **Depth:** Percent of surrounding voxels that are WM. Typical range: 0-1 (0-100%). A higher WM contact depth equates to a contact deeply embedded into WM tissue with little surrounding GM tissue. This convention is similar to the proximal tissue density (PTD) used in Mercier et al. 2017⁵. A sphere with a diameter of 1cm around the centroid coordinate of the contact was used to calculate the percent WM contained within that sphere. Analyses that relied on a definition of depth used the following criteria:
 - (a) GM: ≤ 0.5 (or 50%).
 - (b) WM: ≥ 0.9 (or 90%)
 - (c) contacts between 0.5 and 0.9 were excluded from analysis comparing GM and WM

The above definitions, along with bipolar montaging (below) help ensure the recorded signals are a good representation of activity of that tissue and contact location. We chose a definition of WM

depth for figure analyses in the main text because it is similar to PTD, unless specified WM distance was used (i.e. power analysis vs distance and SNR in Fig. 1c, and controlling for WM distance in Fig. 2f). Please see the main text for a discussion on why two definitions were used.

Briefly, alternative definitions provide different perspectives when each respective tissue definition may have its own limitations (Fig. S2). For example, definition 1 (distance) may incorrectly say a WM contact is close to GM because a single GM voxel was mis-labeled due to imperfections of the Freesurfer tissue segmentation or due to small errors in registration. Regardless, previous studies⁶ and our results in "Controlling for distances between contacts" show similar findings using alternate definitions. For subsequent analyses, we used definition 2 (depth) because it is a similar and complimentary definition to PTD used in Mercier et al. 2017⁵, yet still provides a different perspective in this nascent field for understanding WM iEEG activity where a standard WM iEEG definition is not set. The boundaries for classifying GM and WM for each definition (e.g. ≥ 2 mm) was slightly arbitrary and set before conducting the study: two millimeters for definition 1 was chosen because that is approximately where "deep" white matter is defined in the histological field¹⁰. For definition 2, 50% for GM was chosen because that seemed appropriate where the minority of surrounding voxels are WM. Similarly, 90% for WM was chosen because that seemed appropriate where extreme values of PTD were used in prior studies. Future studies should explore these exact parameters, however, it is out of the scope of this study given that our work and other groups have already shown that the nuances of tissue definitions are not necessarily driving the biological findings.

Contact localization to ablation zone. Contacts contained within the eventual ablation zone were manually identified. Post-surgical T1w images were acquired and manually segmented using ITK-SNAP for the areas ablated. The post-surgical T1w image was linearly registered to pre-implant image. Contact coordinates were found in pre-implant space (electrode localization pipeline above). Contacts that fell into the ablation segmentation mask were labeled as "ablated contacts."

Montaging. While epileptologist routinely use bipolar montage in EEG interpretation, the research community does not. A few studies^{5,6} noted differences in EEG features like power, phase-amplitude coupling, and functional connectivity measurements between the different montages. Makin et al. 2020⁵⁷ emphasize that if the electrodes are common average referenced, it can "smear the effects of some electrodes across the grid". Bipolar referencing was noted to improve overall performance using their deep learning approach. Here we use bipolar referencing to mitigate the effects of volume conduction, not in line with previous studies using common average referencing to create functional connectivity matrices to study epilepsy. To perform bipolar referencing, we started with the contact at the most distal end of the electrode (e.g. LA01) and subtracted the adjacent contact on that electrode (e.g. LA01 = LA01 - LA02). We excluded the last contact on the electrode (there is nothing to subtract), or any bipolar pairs that included a contact excluded through any methods mentioned throughout this methods section (e.g. an artifact contact or a contact not in GM or WM). A critical note is that bipolar montaging, along with the GM and WM contact definitions (above) help ensure the recorded signals are a good representation of activity of that tissue and contact location.

Power Spectral Density Calculation. Following removal of artifact-ridden electrodes, SEEG signals were bipolar referenced. Signals were notch-filtered at 60 Hz to remove power line noise and low-pass and high-pass filtered at 127 Hz and 1Hz to account for noise and drift. Power spectral density (PSD) was calculated using the Scipy Python package version 1.5⁵⁸, and the function `scipy.signal.welch` (default parameters, with FFT epoch length equal to 1s). For the power vs distance analysis, power was taken over the entire state (interictal, preictal, ictal, postictal). For the SNR analysis, power was taken over 1s intervals. Power was log10 transformed on a V^2/Hz scale.

Signal-to-Noise Ratio Calculation. Here we define signal-to-noise ratio (SNR) as the ratio of the ictal power ('signal') to interictal power ('noise'). This definition follows the framework in which epileptologists compare background interictal or preictal activity with ictal activity using EEG features such as amplitude, frequency, and timing. Comparison of any epileptiform signals to this background noise aids in seizure diagnosis and localization.

Over each state, we normalized all times to a length of 100. For example, all interictal, preictal, and postictal segments were 180 seconds (see functional connectivity network generation below). The SNR vectors (180 one-second intervals long) were normalized to a length of 100 for each patient. For ictal segments of variable length of each patient, vectors were normalized to a length of 100. Thus 50% on the x-axis of Fig. 1c represents half-way through each segment (i.e. 90s into the interictal segment, or 50s into a 100s long seizure). Patients' SNR was averaged and plotted. For the GM vs WM SNR comparison, the boxplot of all patients' SNR was plotted at 50% into each state. We then performed a Mann-Whitney U rank test to compare the SNR of GM (distance from GM = 0) and WM ≥ 6 mm from GM at the 50% time interval.

Functional Connectivity Network Generation. Functional connectivity networks were generated from four states: interictal (ii), preictal (pi), ictal (ic), and postictal (po). (1) The interictal state consisted of the time approximately 6 hours before the ictal state. (2) The preictal state consisted of the time immediately before the ictal state. (3) The ictal state consisted of the time between the seizure unequivocal onset and seizure termination. (4) The postictal state consisted of the time immediately after the ictal state. Interictal, preictal, and postictal periods were 180 seconds in duration. To generate functional networks defined by cross correlations between SEEG signals, we carried out a method described in detail previously^{26,28}. SEEG signals were preprocessed using methods outlined in the PSD section above (i.e. artifact removal, bipolar referencing, and filtering). Signals were then pre-whitened using a first-order autoregressive model to account for slow dynamics. Functional networks were then generated by applying a normalized cross correlation function ρ between the signals of each pair of electrodes within each of the four states, using the formula:

$$\rho_{xy} = \max_{\tau} \left[\frac{1}{T} \sum_{t=1}^T \frac{[x_k(t) - \bar{x}_k] * [y_k(t + \tau) - \bar{y}_k]}{\sigma_{x_k} \sigma_{y_k}} \right] \quad [1]$$

where x and y are signals from two electrodes, k is the state, t is one of the T samples during the time window, and τ is the time lag between signals, with a maximum lag of 0.5 s. The absolute value of the cross correlation values were taken. For more details, please see prior publications^{26,28}.

Frequency band definition. Frequency bands were defined using typical ranges reviewed in Newson and Thiagarajan 2019⁵⁹, rounded to the nearest integer:

1. Broadband: 1-128Hz
2. δ (delta) band: 1-4Hz
3. θ (theta) band: 4-8Hz
4. α (alpha) band: 8-13Hz
5. β (beta) band: 13-30Hz
6. γ (gamma) band: ≥ 30 Hz
 - (a) low- γ 30-40Hz
 - (b) mid- γ 70-100Hz
 - (c) high- γ 100-128Hz

The frequency band with the widest range reported in Newson and Thiagarajan 2019⁵⁹ after reviewing 184 studies was the γ -band. A typical γ -band was 30-40 Hz. We subdivided this band further^{26,28}. Low- γ equates to the typical γ -band. Mid- γ skips the 60Hz power line noise (or any potential imperfections in its elimination) and starts at a lower bound of 70Hz. High- γ starts at 100Hz and the upper limit was chosen as 128Hz because that was half the sampling frequency of the patient with the lowest sampling frequency. We chose to analyze the high- γ frequency band because of minimal

volume conduction, and well-developed literature studying high- γ in epilepsy (however, it is less consistently defined across the literature, but typically >100 Hz). Unfortunately, we could not analyze higher frequency bands due to non-uniform sampling frequencies across all patients.

GM and WM functional connectivity (FC) differences. To understand GM and WM FC differences, we subdivided the full network (all contacts included) into tissue sub-networks. Four networks in total were studied: (1) the full network (2) GM-only (3) GM-WM, which included only the connections from GM to WM contacts, and (4) WM-only. We analyzed the distribution of FC values by taking the top triangle half of the symmetrical, undirected adjacency matrix, except for the GM-WM tissue sub-network where all FC values were taken (Fig. 2b, 2c, and 4b histograms). We plotted the empirical cumulative density functions (ECDF) of the sub-network FC values (Fig. 2d and 4b). These plots (Fig. 2b-d) show a representative example patient and seizure. Subsequent plots (Fig. 2e, right to 2h) show at the subject-level.

We wanted to answer the biological question if WM connections are higher (or lower, as hypothesized initially) during the ictal state. Continuing our illustration of the example patient plotted in Fig. 2d, we performed a two-sided Kolmogorov-Smirnov test to test the null hypothesis that the two sub-network distributions of FC values are identical during the ictal period. We found evidence against the null hypothesis.

Across all subjects, we took two approaches to visualizing the differences in the distributions of WM and GM FC values. (1) We pooled all FC values within each tissue sub-network across all patients (i.e. all the WM FC values from patient 1 would be pooled with all the WM FC values from patient 2, and so forth for all patients; $N = 29$). Then (similar to Fig. 2d), we plotted the pooled FC distributions across all patients and the two tissue sub-networks for each state (Fig. 2d, ECDF plots). We see in the ictal state, the WM (blue dotted line) is pulled to the right. (2) In the second visualization approach, we bootstrapped patients and seizures instead of pooling FC values. For each re-sampled patient, we found the median FC value for each tissue sub-network and state. We then took the mean across all patients for each tissue sub-network and state. We plotted the means for 10,000 bootstrap simulations for each state in the inset histograms in Fig. 2d. We see that WM FC is lower during the interictal and preictal states, however, WM FC becomes higher during the ictal state.

To directly compare the GM-only and WM-only sub-networks across each patient, we defined ΔT as the difference in the median FC value for each patient. Here we had two null hypotheses: (1) ΔT during the ictal state (ΔT_{ic}) equals zero, meaning that there is not a difference in the tissue sub-network FC values during the ictal state and (2) $\Delta T_{ic} - \Delta T_{pi}$ equals zero. In other words, rejection of our second null hypothesis would support our alternative hypothesis that the differences in the tissue sub-network FC values become larger at the transition to a seizure state.

We show ΔT for all patients and states (Fig. 2h, left). To test our two null hypotheses, we performed a Mann-Whitney U rank test for the first hypothesis and a Wilcoxon signed-rank test for the second hypothesis (paired test between preictal and ictal). The means of 1,000 bootstrap simulations of ΔT for each state were plotted in Fig. 2h, right.

Controlling for distances between contacts. Although mid-line GM structures (such as the cingulate gyrus) or more medial structures (such as the limbic system) are targets for implantation, it is reasonable to suspect WM contacts are closer together and may explain the increased FC values. Thus, a covariate in our analyses is distance between contacts. We measured euclidean distance between contacts from the coordinates produced in the electrode localization pipeline above. Although we mitigated the influence of passive volume conduction through bipolar montaging explained above, adjacent contacts may still measure related signals even if montaging methods completely eliminated volume conduction because adjacent neural tissue may be performing similar, independent computation. Contacts closer together have higher FC than contacts further away (Fig. 2e) after fitting an ordinary least squares linear regression line (contact distance is first $\log_{10}$ transformed).

Here we wanted to test if WM sub-networks still have increasing FC farther from GM tissue (we quantified "farness" using two definitions, distance and depth defined above). We included the distance between contacts as a covariate in both ordinary least squares models using either the WM distance or WM depth as the other covariate (in Python package statsmodel, we used OLS.from_formula: FC value ~ WM definition + contact distance). P-values reported in Fig. 2f show the probability that the covariate of WM distance or depth have no effect on FC values.

Network Analysis. We took a network analysis approach in studying GM and WM differences. We compared the Full, GM-only, and WM-only networks. We could not feasibly compare the GM-WM networks taking a network analysis approach because these networks are not symmetric (they are not an NxN matrix where N represents the number of nodes in the network). All network measures were calculated using the Brain Connectivity Toolbox for Python, including density, characteristic path length, and transitivity. These network measures were chosen to understand how basic network properties differ between the sub-networks. We also wanted to answer if WM networks are more clustered than GM networks based on the results in Fig. 2 through a measure of transitivity. We performed network analysis on an initial trial of 8 patients (marked in Table 1) and results are shown in Fig. 3. We thresholded the broadband cross correlation functional connectivity matrices at 0.2 and binarized the matrices. After computing the network measures, we compared GM-only and WM-only tissue sub-networks for each state using the Wilcoxon signed-rank test.

Imaging protocol for structure-function analysis. Prior to electrode implantation, MRI data were collected on a 3T Siemens Magnetom Trio scanner using a 32-channel phased-array head coil. High-resolution anatomical images were acquired using a magnetization prepared rapid gradient echo (MPRAGE) T1-weighted sequence (repetition time = 1810 ms, echo time = 3.51m, flip angle = 9, field of view = 240mm, resolution = 0.94x0.94x1.0 mm3). High Angular Resolution Diffusion Imaging (HARDI) was acquired with a single-shot EPI multi-shell diffusion-weighted imaging (DWI) sequence (116 diffusion sampling directions, b-values of 0, 300, 700, and 2000s/mm2, resolution = 2.5x2.5x2.5 mm3, field of view = 240mm). Following electrode implantation, spiral CT images (Siemens) were obtained clinically for the purposes of electrode localization. Both bone and tissue windows were obtained (120kV, 300mA, axial slice thickness = 1.0mm).

Diffusion Weighted Imaging (DWI) Preprocessing. HARDI images were subject to preprocessing pipeline QSIprep to ensure reproducibility and implementation of the best practices for processing of diffusion images⁶⁰. Briefly, QSIprep performs advanced reconstruction and tractography methods in curated workflows using tools from leading software packages, including FSL, ANTs, and DSI Studio with input data specified in the Brain Imaging Data Structure (BIDS) layout.

Structural Network Generation. DSI-Studio (<http://dsi-studio.labsolver.org>, version: March 2021) was used to reconstruct the orientation density functions within each voxel using generalized q-sample imaging with a diffusion sampling length ratio of 1.25⁶¹. Deterministic whole-brain fiber tracking was performed using an angular threshold of 35 degrees, step size of 1mm, and quantitative anisotropy threshold based on Otsu's threshold⁶². Tracks with length shorter than 30mm or longer than 800mm were discarded, and a total of 1,000,000 tracts were generated per brain. Deterministic tractography was chosen based upon prior work indicating that deterministic tractography generates fewer false positive connections than probabilistic approaches, and that network-based estimations are substantially less accurate when false positives are introduced into the network compared with false negatives⁶³. To calculate structural connectivity between electrode contacts, ROIs of 7mm radius surrounding the contacts were created (Fig. 4a). The ROI volume was selected to equate to the volume of an average ROI in another commonly used neuroimaging atlas, the Automated Anatomical Labeling (AAL) atlas⁶⁴⁻⁶⁶. This soft-boundary⁵⁴, contact-specific atlas was used to because we were

studying individual signals from each contact and the tissue they recorded from. Structural networks were generated by computing the number of streamlines passing through each pair of ROIs. Streamline counts were log-transformed and normalized to the maximum streamline count, as is common in prior studies⁶⁷⁻⁷⁰.

Structure-Function Correlation. To quantify the relationship between structure and function, we computed the Spearman Rank correlation coefficient between the edges of the structural connectivity networks and the edges of the functional connectivity network (Fig. 4c). Broadband cross correlation was used because it has shown the most robust SFC in epilepsy patients⁷¹. We divided each structural and functional network in tissue sub-networks: (1) A full network representing equal treatment of GM and WM contacts (2) a GM-only network. (3) and GM-WM network (4) a WM-only network. We correlated each structural and functional sub-network. We re-sampled our data set (see bootstrapping methods below) and compared the the SFC of each sub-network across the four states. We performed the Wilcoxon signed-rank test between the ictal and preictal states for each sub-network to test the following alternative hypothesis: $\Delta SFC_{tissue} \neq 0$, where ΔSFC_{tissue} represents the difference between the ictal and preictal SFC ($SFC_{ic} - SFC_{pi}$) for the given tissue sub-network. We hypothesized that ΔSFC is higher in any tissue sub-network containing GM because GM is the neural tissue that drives the activity of the brain. Thus WM ΔSFC would be lower. We bootstrapped this experiment 1000 times and plot the distribution of the mean ΔSFC_{tissue} across the re-sampled patients and seizures. We report confidence intervals.

Outcome Classification. We determined the outcomes of epilepsy surgery using the Engel classification scoring system⁷². We included patients we were ≥ 2 years from surgery to account time where seizure occurrences may relapse. Clinical studies monitoring outcomes are ongoing, however, ≥ 2 years was selected because we felt this was ample time to increase the accuracy of the ground-truth effectiveness of the epilepsy surgery. We separated good (Engel I) and poor (Engel II-IV) outcome patients typically done in the literature^{45,70,73-78}. Note that we dichotomized good and poor outcomes because we assume that in good outcome patients, the ablated regions contained a well-localized seizure generating area that was correctly targeted. We wanted to understand how this targeted area is related to WM activity. Note that a poor outcome patient does not necessarily mean a "poor" outcome for the patient. Engel III (a poor outcome class in this study) means marked seizure improvement, which greatly enhances quality of life in some patients and would be considered a "good" outcome to them. Here we dichotomize Engel I and II-IV, not from a patient point of view, but rather from a neurosurgical point of view in accurately localizing a circumscribed SOZ. In the case of Engel II-IV, patients continue having seizures, and thus the true SOZ was missed (see the discussion in the main text "The Distributed Epileptic Network Hypothesis"). In this study, poor outcome patients were not recommended for palliative care, meaning their overall clinical findings (including their SEEG recordings) indicated a reasonable suspicion for the target area. Seizure freedom may last for a short time (≤ 2 years). After a unspecified time, brain regions may adapt and seizures may continue. Hence, this is one reason why we included patients with outcomes of follow-up longer than previous studies.

Functional connectivity of tissue sub-networks in good and poor outcomes. We compared the tissue sub-networks in good and poor outcome patients using methods similar to Fig. 2. We compared the distribution of FC values in (1) the full (2) GM-only (3) GM-WM and (4) WM-only networks between good and poor outcomes during the preictal and ictal states. We tested the null hypothesis that the distribution of FC values in good and poor outcome patients with respect to each sub-network and state are similar. We performed a two-sided Kolmogorov-Smirnov test and outputs are shown in Fig. 5b. We also compared the change in FC (ΔFC) from preictal to ictal states ($\Delta FC = FC_{ictal} - FC_{preictal}$) from the contacts contained within the ablation zone to other WM contacts (Fig. 5f). We pooled all FC values from the bootstrapped patients, seizures, and connectivity matrices (see bootstrapping methods below) and plotted the distributions (Fig. 5f, left) similar to Fig. 2g. To

compare ΔFC on a patient level, we took the median FC value for each patient, and plotted the results in Fig. 5f, right. To test the null hypothesis that good outcome patients have similar FC values from the ablation zone to all WM contacts, we performed the Mann-Whitney U test. To provide more robust statistical methods considering our small sample size of poor outcome patients, re-sampling hypothesis testing was also performed and is explained in detail below.

Hypothesis testing comparing good and poor outcome patients. We wanted to know if WM activity is different between good and poor outcome epilepsy patients.

We repeated the analysis in Fig. 2h separating good and poor outcome patients with the null and alternative hypotheses defined below:

1. $H_0: Poor_{\Delta T_{ic}} - Good_{\Delta T_{ic}} = 0$. The differences in GM functional connectivity and WM functional connectivity (ΔT , differences in *Tissue*) is similar for good and poor outcome patients during the ictal period.
2. $H_1: Poor_{\Delta T_{ic}} - Good_{\Delta T_{ic}} \neq 0$. The differences in GM functional connectivity and WM functional connectivity (ΔT , differences in *Tissue*) is NOT similar for good and poor outcome patients during the ictal period.

After the results from Fig. 5c, we wanted to test the hypothesis that ablated regions in good outcome patients do not increase in correlation to WM regions as much as poor outcome patients during seizures. This hypothesis helps answer the biological question if WM electrodes reveal a distributed, less-localizing nature in the pathophysiology of their epilepsy. Here we formally define the null and alternative hypotheses:

1. $H_0: Poor_{\Delta FC_{ablated\ to\ WM}} - Good_{\Delta FC_{ablated\ to\ WM}} = 0$. The change in functional connectivity from preictal to ictal states (ΔFC) from the ablated GM regions to WM regions is similar for good and poor outcome patients.
2. $H_1: Poor_{\Delta FC_{ablated\ to\ WM}} - Good_{\Delta FC_{ablated\ to\ WM}} \geq 0$. The change in functional connectivity from preictal to ictal states (ΔFC) from the ablated GM regions to WM regions is greater for poor outcome patients than good outcome patients. Using knowledge and expectations based on prior results, we hypothesize poor outcome patients are greater, therefore, we do a one sided test (see re-sampling hypothesis testing below). We omit the absolute value in eq. 3.

An important note: given the sampling of our data (unequal seizure number across patients, non-uniform implantation schemes, unequal tissue sampling, only 17 patients with ≥ 2 years of outcomes reported, etc), we performed both bootstrapping methods and re-sampling hypothesis testing for a robust analysis (see relevant sections below).

Bootstrapping. The bootstrap is a widely applied and powerful statistical tool⁷⁹. It is useful in our data set where a measure of variability is otherwise difficult to obtain. In our data set, we have a variable amount of seizures per patient. Each patient has a variable implantation strategy with a variable number of recording contacts. Therefore, each patient will also have a different sampling of tissue types with resulting different network sizes. To mitigate these biases, or over-/under-sampling, of any one patient or seizure, we re-sampled with replacement at each level of patient and seizure. For example, we would sample with replacement N patients from our cohort equal to 1x the number of total patients. Then, we would sample with replacement M seizures from each patient equal to the number of seizures that patient has. In this case, because patients are already bootstrapped, seizures can be considered independent. We would bootstrap 10,000 times to ensure patients have equal sampling of seizures.

Re-sampling hypothesis testing. We employed re-sampling hypothesis testing to understand the differences in WM seizure activity in good and poor outcome patients in Fig. 5. Due to a small sample size of our patients with validated outcome scores ≥ 2 years, we follow the permutation approach to the p-value outlined in James et

al., 2021.⁷⁹. As illustrated in their text, this approach is preferable and is considered reliable; we follow Algorithm 13.3 of the text exactly and written here for convenience:

1. Compute the test statistic, T , on the original data $x_1, \dots, x_{n_X}$, and $y_1, \dots, y_{n_Y}$.
2. For $b = 1, \dots, B$:
 - (a) Permute the $n_X + n_Y$ observations at random. Call the first n_X permuted observations $x^*_1, \dots, x^*_{n_X}$, and call the remaining observations $y^*_1, \dots, y^*_{n_Y}$.
 - (b) Compute T on the permuted data $x^*_1, \dots, x^*_{n_X}$ and $y^*_1, \dots, y^*_{n_Y}$, and call the result T^{*b} .
3. The p-value is given by eq. 3 below.

We computed the test statistic as the two-sample t -statistic Following James et al., 2021.⁷⁹:

$$T = \frac{\hat{\mu}_X - \hat{\mu}_Y}{s \sqrt{\frac{1}{n_X} + \frac{1}{n_Y}}} \quad [2]$$

where $\hat{\mu}_X = \frac{1}{n_X} \sum_{i=1}^{n_X} x_i$, $\hat{\mu}_Y = \frac{1}{n_Y} \sum_{i=1}^{n_Y} y_i$, $s = \sqrt{\frac{(n_X-1)s_X^2 + (n_Y-1)s_Y^2}{n_X + n_Y - 2}}$ and s_X^2 and s_Y^2 are unbiased estimators of the variances in poor and good outcome patients, respectively. We randomly permute the $n_X + n_Y$ observations B times, and each time we compute eq. 2. We let $T^{*1}, \dots, T^{*B}$ denote the values of eq. 2 on the permuted data. These can be viewed as an approximation of the null distribution of T under H_0 . To compute the p-value, we do

$$p - value = \frac{\sum_{b=1}^B 1_{|T^{*b}| \geq |\bar{T}|}}{B} \quad [3]$$

This computes the fraction of permuted datasets for which the value of the test statistic is at least as extreme as the value observed on the original data's test statistic, T . For our re-sampling hypothesis testing approach, we treated each seizure as an independent sample. We also averaged over patients-level and show significant ($p < 0.05$) or trending ($p < 0.1$) results in Fig. S6. Note that we approached the analysis comparing GM and WM throughout the manuscript using multiple methods to ensure we lower our Type II error rates testing the overall question if WM is different from GM and provides unique information about epilepsy.

Functional connectivity changes in WM as a function of distance from ablation zone between good and poor outcome patients. We measured the change in functional connectivity (ΔFC) from preictal to ictal states between all electrode contacts. We determined which contacts recorded neural activity in the suspected seizure onset zone and subsequently ablated. We then determined the FC between the ablated contacts and WM contacts. We measured the distance between those contacts and took the median FC of all the contacts less than or equal to the distance specified on the x-axis of Fig. 2h. Ninety-five percent confidence intervals are plotted. The results shown is the outcome of one bootstrap simulation (see bootstrapping methods above).

A note on the ablation zone: In good outcome patients, we assume that the ablation zone contained the "true" seizure onset zone, or alternatively, contained a seizure area that was well localized to that region. In poor outcome patients, the ablated regions were targeted because it followed the above assumptions in good outcome patients, but the outcomes would not be known yet. It was the suspected SOZ. However, our results show that in poor outcome patients, this suspected SOZ becomes more correlated to distant WM regions, pointing to a more distributed, non-localizing nature of their seizure pathophysiology even in the face of all clinical evidence pointing to otherwise (or else they would not have been recommended epilepsy surgery, but rather alternative treatment options such as implantable neurostimulating devices).

Data availability and Reproducibility. All code files used in this manuscript are available at <https://github.com/andyrevell/revellLab>. All de-identified raw and processed data (except for patient MRI imaging) are available for download on Box. Link provided on GitHub. The GitHub repository used to analyze the data is also contained within Box. Raw imaging data is available upon reasonable request from Principal Investigator K.A.D.; tractography files generated from the imaging data are readily available on Box. iEEG snippets used specifically in this manuscript are contained within the Box data folder, while full iEEG recordings are publicly available at <https://www.ieeg.org>. The Python environment for the exact packages and versions used in this study is contained in the environment directory within the GitHub. The QSIPrep docker container was used for DWI preprocessing.

ACKNOWLEDGMENTS

We thank Adam Gibson, Carolyn Wilkinson, Jacqueline Boccanfuso, Magda Wernovsky, Ryan Archer, Kelly Oechsel, members of Andrew's Thesis Committee, Braden Kelner, and all other members and staff of the Center for Neuroengineering and Therapeutics for their continued help and support in this work.

Funding

This work was supported by National Institutes of Health grants 5-T32-NS-091006-07, 1R01NS116504, 1R01NS099348, 1R01NS085211, and 1R01MH112847. We also acknowledge support by the Thornton Foundation, the Mirowski Family Foundation, the ISI Foundation, the John D. and Catherine T. MacArthur Foundation, the Sloan Foundation, the Pennsylvania Tobacco Fund, and the Paul Allen Foundation.

Competing Interests

The authors declare no competing interests.

Supplementary Materials

Please see supplemental materials below.

- Figures
 - [Fig. S1](#): Overview of Localization and Segmentation
 - [Fig. S2](#): White Matter Definitions: Distance and Depth
 - [Fig. S3](#): GM and WM distribution of functional connectivity value: Select frequency bands and other functional connectivity measurements
 - [Fig. S4](#): Differences in GM and WM (ΔT) for all frequency bands and functional connectivity measurements
 - [Fig. S5](#): Distribution of seizure occurrences and lengths
 - [Fig. S6](#): Repeat analysis of Fig. 5 on a per-patient level.
- Tables
 - [Table. S1](#): Patient table
- Other materials
 - [Glossary](#)

Glossary.

1. **Ablated contacts:** Contacts which fell in the (eventually) ablated region
2. **GM:** Gray Matter. See tissue definitions in the methods section for details on how contacts were classified as either GM or WM.
3. **WM:** White Matter. See tissue definitions in the methods section for details on how contacts were classified as either GM or WM.
4. **GM contact.** A contact defined as GM, using tissue definitions detailed in the methods section.
5. **WM contact.** A contact defined as WM, using tissue definitions detailed in the methods section.
6. Δ **SFC:** The change in SFC between ictal and preictal stats ($SFC_{ictal} - SFC_{preictal}$). This indicates whether or not the change in functional connectivity is congruent with the underlying structural connectivity.
7. Δ **FC.** The change in functional connectivity between ictal and preictal states. $FC_{ictal} - FC_{preictal}$. This measures if two contacts are increasing or decreasing in the functional connectivity values when a seizure evolves.
8. **ii, pi, ic, po.** Interictal, preictal, ictal, postictal states.
9. **T1w.** T1-weighted MRI image.
10. **Contact.** A single sensor on an electrode that records LFP. Not to be confused with the electrode itself (an entire "stick" of an SEEG)
11. **Electrode.** Not to be confused with contact.
12. **SEEG:** Stereoelectroencephalography.
13. **SNR:** Signal-to-noise ratio. See methods section.
14. **ECoG:** Electrocorticography.
15. **Functional connectivity (FC).** The statistical relationship between two signals (two contacts in this study). Many FC measurements exists, including but not limited to Pearson correlation, Spearman Rank correlation, Cross correlation, and Coherence. We primarily use Cross correlation in this manuscript.
16. **Structural connectivity (SC).** The physical relationship between two brain regions. We use streamline counts in this manuscript from High Angular Resolution Diffusion Imaging.
17. **Bootstrapping.** A robust statistical tool. See methods section.
18. **Re-sampling hypothesis testing.** A robust statistical analysis technique for testing hypothesis using permutation approaches. See methods section.

Fig. S1. Overview of Localization and Segmentation. | a, Implant T1w MRI. b, Implant CT. c, Implant CT SEEG contacts are segmented and localized in native coordinates. d, Pre-implant 3T T1w MPRAGE. Usually higher quality imaging (1mm isotropic) was used for tissue segmentation. If patient did not undergo high resolution imaging, then a clinical T1w pre-implant image is used. e, Segmentation output by Freesurfer showing tissue parcellations. f, Conversion of tissue parcellation into labels of GM, WM, CSF, or outside of the brain. g, Combination of entire pipeline to localize electrode contacts to tissue. Images (a) and (b) are linear registered using FSL's flirt command so that contact coordinates localized in native CT space can be converted to image (a)'s space. Images (a) and (d) are linear registered so that contact coordinates in image (a)'s space can be converted to image (d)'s space. Once contacts are localized to image (d)'s space and image (d) has been segmented, coordinates and segmentation are combined to produce image (g). WM classification is determined using two definitions shown in Fig. S2.

White Matter Definitions: Distance and Depth

Fig. S2. White Matter Definitions: Distance and Depth. | a, Contacts are localized to tissue. See Fig. S1. b, Binary classification of WM and GM contacts can be found by measuring the distance of the contact's center to the nearest GM voxel. c, Contacts can be classified by WM depth. This is the percent of surrounding voxels (within a 1cm diameter sphere) that are WM. This is closely related to the PTD measurement in Mercier et al. 2017⁵ (see Methods). A benefit of WM depth is that it accounts for cortical folding and small errors in tissue segmentation. WM contacts near the cortical surface, but a few voxels from GM will have low WM depth because they are still surrounded by brain sulci, and thus many GM voxels. Furthermore, a single voxel mis-segmented as GM can affect WM distance. Therefore, we primarily used WM depth.

Fig. S3. GM and WM distribution of functional connectivity value: Select frequency bands and other functional connectivity measurements. | We repeat analysis from Fig. 2g using select frequency bands and functional connectivity measurements. For all frequency bands and functional connectivity measurements, repeating Fig. 2h, see Fig. S4. We focus on high- γ -band and cross correlation in the main text as detailed previously. Our central hypotheses related to seizure activity, therefore, we tested our findings using high- γ . However, we note more striking differences in other frequency bands and functional connectivity measurements. We provide information here and in Fig. S4 to aid research using this manuscript for a basis.

Fig. S4. Differences in GM and WM (ΔT) for all frequency bands and functional connectivity measurements. | We repeat analysis from Fig. 2h showing all frequency bands and functional connectivity measurements. Top row: Pearson correlation. Middle row: Coherence. Bottom row: Cross Correlation. Left-to-right: Broadband, delta, theta, alpha, beta, low-gamma, mid-gamma, high-gamma. See methods on frequency band definitions. We note results may be biased based on the choice of frequency bands, therefore, we provide a more comprehensive analysis here. We selected high- γ -band and cross correlation prior to conduction of the study based on prior knowledge about epilepsy and functional connectivity measurements from the literature. Error bars are 95% confidence intervals. Asterisk represent significant findings between the statistical comparison of ictal and preictal ΔT . Multiple comparison correction is not performed because we do not use these results subsequently in the manuscript, but we provide the analysis here for reference in other studies.

Fig. S5. Distribution of seizure occurrences and lengths. | **a**, The distribution of seizures for each patient. Most patients ($N=13$) had one seizure annotated. Two patients had 11 seizures annotated. **b**, The distribution of seizure lengths. Bins of 60 seconds are shown on the left, and bins of 10 seconds are shown on the right. One seizure was approximately 16 minutes long while most seizures were between 1 and 2 minutes long.

Fig. S6. Repeat analysis of Fig. 5 on a per-patient level. | **a**, Repeat of Fig. 5c. **b**, Repeat of Fig. 5d, top. **c**, Repeat of Fig. 5f, right. **d**, Repeat of Fig. 5g, top. An important point to stress is that the Distributed Epileptic Hypothesis described in the main text may apply too only *some* poor outcome patients. The reasoning is that 1) the SOZ is still a focal region, but missed during ablation surgery, or (2) the SOZ appears focal to clinicians, but the underlying pathophysiology in some patients points to a more distributed interaction of brain regions and interconnecting WM pathways⁴⁰. The heterogeneity of poor outcome patients itself proves problematic when poor outcome patients are less common than good outcome patients across epilepsy centers. We show low variation in WM measurements in good outcome patients, but high variability in poor outcome patients. This indicates that different pathophysiological processes are going on in poor outcome patients, however, similar pathophysiological processes are going on in good outcome patients. WM recordings gives better clues to the underlying pathophysiology of epilepsy and thus more research needs to be done. We provide a seizure-level analysis in the main text rather than on a patient-level shown here to understand the variation amongst all seizures. A study with patients from a combination of many epilepsy centers will need to be conducted to improve our Type II error rate. Here we show significance ($p<0.05$) and trends ($p<0.1$) in our findings with significance defined as an arbitrary cutoff of $p<0.05$. One-sided independent t-test are performed. We caution to base future work and hypotheses on a single plot in this manuscript, however, we provide ample evidence from multiple analysis approaches to robustly claim that GM and WM have differences related to epilepsy pathophysiology, and that WM recordings should not be ignored.

Subject	Sex	Intervention	Engel	HARDI
RID0194	M	RNS	–	YES
RID0238	F	Laser	IA	–
RID0267	F	Laser	IA	–
RID0274	M	Laser	IIA	–
RID0278	M	Laser	IVC	YES
RID0279	F	Laser	IA	–
RID0294	M	Laser	IC	–
RID0307	M	Laser	IA	–
RID0309	M	Laser	IA	YES
RID0320	F	Laser	IB	YES
RID0322	M	Laser	IIIA	–
RID0365	M	Laser	IB	YES
RID0371	M	Laser	IIA	–
RID0382	M	Laser	IIIA	–
RID0405	M	Resection	IIIA	–
RID0420	F	DBS	–	YES
RID0424	F	Resection	IA	–
RID0440	F	Laser	IB	YES
RID0442	M	Laser	IIIA	–
RID0502	F	RNS	–	YES
RID0508	M	NR	–	YES
RID0536	F	NR	–	YES
RID0572	F	NR	–	YES
RID0583	F	RNS	–	YES
RID0595	F	NR	–	YES
RID0596	M	Laser	–	YES
RID0648	M	Laser	–	YES
RID0650	M	NR	–	–
RID0652	M	NR	–	YES

Table S1. Patient table. | Patients who had no surgery may not have had an intervention at time of writing. Only two patients with Engel outcome scores underwent resection rather than ablation, however, in the main text we generalize to writing about ablation only despite a very small minority of patients underwent resection. **M**, Male; **F**, Female; **RID**, Research ID number; **DBS**, Deep brain stimulation; **L**, left; **R**, Right; **St. Dev**, Standard deviation. **NR**, Not reported or still on-going decision for surgery.