Structural analysis and construction of thermostable antifungal chitinase

Dan Kozome^{1, 4}, Keiko Uechi¹, Toki Taira¹, Harumi Fukada², Tomomi Kubota³, Kazuhiko Ishikawa^{3*}

¹ Department of Bioscience and Biotechnology, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa, 903-0213, Japan

² Graduate School of Science, Osaka Prefecture University, 1-1 Gakuencho, Sakai, Osaka 599-8531, Japan

³ Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Central-6, 1-1 Higashi, Tsukuba, Ibaraki, 305-8566, Japan

⁴ Okinawa Institute of Science and Technology Graduate University, Okinawa, 904-0412, Japan (current affiliation)

Corresponding author: Kazuhiko Ishikawa

Email: kazu-ishikawa@aist.go.jp

ABSTRACT

Chitin is a biopolymer of *N*-acetyl-D-glucosamine with β -1,4-bond and is the main component of arthropod exoskeletons and the cell walls of many fungi. Chitinase (EC 3.2.1.14) is an enzyme that hydrolyzes the β -1,4-bond in chitin and degrades chitin into oligomers. It has been found in a wide range of organisms. Chitinase from *Gazymaru* (*Ficus microcarpa*) latex exhibits antifungal activity by degrading chitin in the cell wall of fungi and is expected to be used in medical and agricultural fields. However, the enzyme's thermostability is an important factor; chitinase is not thermostable enough to maintain its activity under the actual applicable conditions. We solved the crystal structure of chitinase to explore the target sites to improve its thermostability. Based on the crystal structure and sequence alignment among other chitinases, we rationally introduced proline residues, a disulfide bond, and salt bridge in the chitinase using protein engineering methods. As a result, we successfully constructed the thermostable mutant chitinases rationally with high antifungal and specific activities. The results provide a useful strategy to enhance the thermostability of this enzyme family.

Keywords: protein engineering; proline; thermostable; disulfide bond; salt bride; chitin; antifungal activity; crystal structure

IMPORTANCE

We solved the crystal structure of the chitinase from *Gazymaru* (*Ficus microcarpa*) latex exhibiting antifungal activity. Furthermore, we demonstrated that the thermostable mutant enzyme with a melting temperature (T_m) 6.9 °C higher than wild type (WT) and a half-life at 60°C that is 15 times longer than WT was constructed through 10 amino acid substitutions, including five proline residues substitutions, making disulfide bonding, and building a salt bridge network in the enzyme. These mutations do not affect its high antifungal activity and chitinase activity, and the principle for the construction of the thermostable chitinase was well explained by its crystal structure. Our results provide a useful strategy to enhance the thermostability of this enzyme family and to use the thermostable mutant as a seed for antifungal agents for practical use.

INTRODUCTION

It has been reported that nearly 80% of plant pathogens are fungi (1), and their damage to crops has become a significant problem in the agricultural industry. Up to now, chemical fungicides have been extensively adopted in combating current plant diseases. However, since fungi are eukaryotes like mammals, many chemical fungicides are highly toxic to human and fungi, and their utilization has been associated with risk. Therefore, the development of fungus-specific antifungal agents is in high demand. In a pathogenic attack, plants produce pathogenesis-related (PR) proteins as a part of systemic acquired resistance (2). Plant chitinase is considered to be one of these PR proteins. Chitinase, an enzyme that degrades chitin into oligomers, has been found to exist in a wide range of organisms. Since chitin is absent in animal cells compared to fungal cells, chitinases exhibiting the antifungal activity through the enzymatic hydrolysis of fungal cell walls (3) are expected to be used as a fungus-specific antifungal agent. According to the CAZy database (<http://www.cazy.org/>), chitinases (EC 3.2.1.14) are divided into two families, glycoside hydrolase (GH) family 18 (GH18) and 19 (GH19) based on amino acid sequences of their catalytic regions (4). Furthermore, according to an independent classification system for plant chitinases, they are grouped into at least five classes (classes I, II, III, IV, and V) based on their domain organization and loop deletions (5).

There are many reports of antifungal activity in plant GH19 chitinases; however, reports of antifungal activity in plant GH18 chitinases are minimal. Taira et al. have screened various tropical plants that produce latex for chitinase activity (6). Among them, latex from gazyumaru (*Ficus microcarpa*), a woody flowering plant distributed in subtropical/tropical regions in Asia, showed the highest catalytic activity of all samples assayed. Moreover, gazyumaru latex exhibits strong antifungal activity. Three types of chitinase were purified from gazyumaru latex; Gazyumaru latex chitinase-A, B, and -C (GlxChiA, B, and C) belong to class III, class I, and class III chitinase, respectively. Among them, GlxChiB, a basis class I chitinase (32 kDa, pI 9.3), exhibited the highest antifungal activity (6). GlxChiB consists of two domains, carbohydrate-binding module family 18 (CBM18) as a chitin-binding domain (ChBD) and GH19 domain as a catalytic

domain (7). The optimum temperature for GlxChiB was 50 – 60 °C, but this enzyme was unstable above 60 °C (6).

For industrial use of the enzyme, its high activity and thermostability are important factors. The thermostability of many enzymes has been improved by protein engineering methods for their industrial use. Random mutagenesis is a typical tool for increasing thermostability (8-10). Recently computational design can be applied to the protein design (11, 12). Compared with such trial-and-error methods, rational protein design using the protein model or structural data is developing as a powerful and efficient tool for constructing thermostable enzymes (13-15). In this study, we solved the crystal structure of the catalytic domain of both WT and a thermostable mutant of GlxChiB, which was rationally designed by the protein engineering method not to affect the enzyme activity. Furthermore, based on the crystal structures of the WT and the thermostabilized enzyme, the mechanism of each mutation and new insight on the thermostabilizing effect are discussed.

RESULTS

Structural aspect and crystal structure of the chitinase

GlxChiB was found to belong to the GH19 family containing the chitin-binding domain (ChBD) at the N-terminus and exhibited high catalytic activity on chitin degradation and antifungal activities (7). Multiple sequence alignments of GH19 chitinases from the plant are shown in Figure 1. Only the truncated catalytic domain of WT GlxChiB (CD-WT; Amino acid numbers 45-289) was the object for the crystal structure determination in this study due to the low molecular weight ChBD (amino acid numbers 1-39) containing four disulfide bonds appearing to be a thermostable protein as observed in hevein, a small disulfide-rich protein from *Hevea brasiliensis* (16). The recombinant catalytic domain of the enzyme was prepared and purified as described previously (7). Crystals approximately 200 × 200 × 200 μm in size were obtained after more than one week at 298 K (see Materials and methods for the crystallization conditions). The diffraction data set was collected at a wavelength of 1.000 Å at the KEK PF BL5A. Data collection and refinement statistics are shown in Table 1. The diffraction data were collected up to 1.6 Å resolution. The crystals belonged to orthorhombic of space group P2(1)2(1)2(1), with unit cell $a =$

90.900 Å, $b = 106.583$ Å, $c = 107.656$ Å. The initial phase of the structure was determined by molecular replacement with Phaser in the CCP4 package using the crystal structure of chitinase (GH19) from *Vigna unguiculata* (PDB: 4TX7) (17) as the search model. In the asymmetric unit, four molecules of CD-WT were identified; the Matthews coefficient (VM) (18) was calculated as $2.32 \text{ Å}^3 \text{ Da}^{-1}$, with a solvent content of 47% (v/v). After refinement using Refmac5, R factors of the model were estimated as Rwork 22.7% and Rfree 25.3%. The value of Rfree was obtained from a test set consisting of 5% of all reflections. Ramachandran plot (19) for the model showed that 98% of the residues were in the most favored regions, with 2% of the residues in additional allowed regions. Trp121 is in Ramachandran outliers regions, with unambiguous electron density. The coordinates and structure factors for CD-WT were deposited in the Protein Data Bank under the accession code 7V91. The overall structures of four molecules of GlxChiB were built from 45 to 288 residues (Figure 2A left), and the root means square deviation (RMSD) of the C α atoms was less than 0.457 Å over 300 C α atoms among them. The RMSD value indicated that the overall structures of the four molecules of GlxChiB were almost identical.

Design of thermostable mutants of GlxChiB

Sequence alignment among the chitinases; class-I chitinase from *Oryza sativa* (PDB ID: 2DKV) (20), catalytic domain of class I chitinase from *Vigna unguiculata* (PDB ID:4TX7) (17), class II chitinase from *Secale cereale* (rye) seed (PDB ID:4DWX) (21), and class II chitinase from *Carica papaya* (PDB ID:3CQL) (22) are shown in Figure 1. In the catalytic domains of chitinase sequences, catalytic residues, disulfide bonds, and overall structures are well conserved. Proline is the only amino acid with a secondary amine, in that the side chain is directly connected to nitrogen of the main chain, preventing the rotation of phi angles of the peptide bond. Therefore, proline residues in proteins make the main chain at loop regions rigid, and several of them should be critical to the stability, including the thermostability of the protein. Furthermore, it has been proposed that the thermostability of a protein can be increased by introducing a proline that decreases the configurational entropy of unfolding (23). In the catalytic domains of GlxChiB sequence, some residues where proline residue is conserved in some other chitinases are replaced with other amino

acids (Figure 1). Therefore, the introduction of proline residues in CD-WT appears tolerant and responsible for increasing its thermostability. We first took notice of the proline residues for the construction of thermostable mutants. The five positions (Ala117, Ala151, Gln171, Ala233, and Ala254) in which proline residue are not conserved in CD-WT among these enzymes (Figure 1 and 3) are remarkable points for the mutations. Referring to the crystal structure of CD-WT, Ala117 is located at the second site of a beta-turn I (Figure 4A), Ala151 is located at the second site of a beta-turn II (Figure 4B), Gln171 is located in the middle of the helix structure (Figure 4C), and Ala233 and Ala254 are located at the loop region preceding conserved helix structure (Figure 4D and E). The following five substitutive mutations were employed in CD-WT: A117P, A151P, Q171P, A233P, and A254P.

A disulfide bridge linking the two cysteines and their respective main peptide chains can restrict the motion of the unfolded, random coil of protein or stabilize the folded state of the protein. One disulfide bridge can contribute 2.3–5.2 kcal/mol to the thermodynamic stability of proteins (24). Considerable evidence has demonstrated the thermostability effects of engineered disulfide bridges in protein. The crystal structural analysis clarified that CD-WT contains three disulfide bonds and free Cys residues. Using the program SSBOND (25) referring to the crystal structure of CD-WT, we cannot find the suitable position in which the additional disulfide bond can be introduced in CD-WT. We have no structural information of the linker region (amino acid numbers 40-44) between ChBD and CD-WT. However, from the crystal structure, it is estimated that the distance between C α s of N-terminal Asp45 and Lys83 is 6-8 Å in all CD-WT molecules, and then the distance between C α s of Gly44 (in the linker region) preceding Asp45 and Lys83 is considered to be similar. Therefore, the disulfide bond between Cys44 and Cys83 could form (Figure 3) and the mutant employed in G44C/K83C. In addition, it has been reported that salt bridges play an important role in the thermostability of many proteins. The ionizable side chains frequently form ion pairs in many protein structures. Since electrostatic attraction between opposite charges is strong *per se*, salt bridges can be regarded as an important factor stabilizing the protein structure. In addition, many salt bridges were observed at the surface of thermophilic enzymes (26-28). The loop region

250-255 located at the surface loop region between two helices appeared flexible. We targeted the loop region 250-255 located at the surface loop region between two helices in GlxChiB (Figure 3). Significant salt bridges were observed at the corresponding region in 2DKV, 4TX7, 4DWX, and 3CQL but not in GlxChiB according to the multiple sequence alignment (Figure 1). Therefore, we tried to modify the three amino acids Q250/K253/Q255 to introduce salt bridges at the region in CD-WT by referring to these structures. Two mutants, Q250K/K253D/Q255R and Q250R/K253D/Q255R were prepared.

Comparison of the thermostability of the mutants

Recombinant GlxChiB and their mutants described above were prepared and purified by the same method previously reported (7). The enzymatic activity assays were performed as described in the material method. Their specific activities at 37 °C were not meaningfully affected by these mutations (Table 2). WT GlxChiB exhibited the maximum activity at the temperatures around 55 °C under the conditions employed at pH 7.0 for 15 minutes (Figure S1). The thermostability was examined by the residual activities of enzymes after the incubation at 60 °C. All proline substitution mutants showed a longer half-life than that of WT (Table 2). Furthermore, as for the melting temperature (T_m) of the mutants measured by differential scanning calorimetry (DSC), all proline substitution mutants exhibited higher T_m values than that of WT (Table 2). For G44C/K83C, T_m did not change significantly; however, its half-life at 60 °C was elongated (Table 2). For the two mutants with a salt bridge network introduced (Q250K/K253D/Q255R and Q250R/K253D/Q255R), the specific activities at 37 °C were slightly increased, and the half-life at 60 °C was elongated (Table 2).

The integrated Pro mutant and the crystal structure of the mutant

For the thermostable proline substitute mutants (A117P, A151P, Q171P, A233P, and A254P), no decrease of the enzymatic or antifungal activity was observed (Table 2 and Figure 5). Table 2 shows that these substitutions did not affect both activities. All proline substitutions elongated half-life at 60 °C 1.4-2.4 times longer than WT, and the individual

effects on thermostability were considered cumulative. To examine the structural effects of proline substitutions on the enzyme's thermostability, we prepared the integrated mutant, mt5 (A117P/A151P/Q171P/A233P/A254P) and solved the crystal structure of the catalytic domain of mt5 (CD-mt5). CD-mt5 was prepared and purified as the same as for CD-WT. The crystals of CD-mt5 were grown under different conditions from those for CD-WT. Data collection and refinement statistics are shown in Table 1. The diffraction data was collected to a resolution of 1.80 Å. The crystals belonged to space group C2, with unit cell $a = 140.58 \text{ Å}$, $b = 38.89 \text{ Å}$, $c = 177.74 \text{ Å}$, $\beta = 96.03^\circ$; note this the crystal system is completely different from that of CD-WT. The coordinates and structure factors for CD-mt5 were deposited in the Protein Data Bank under the accession code 7V92. The overall structures of four molecules in an asymmetric unit were built from 45 to 288 (Figure 2A, right), and the RMSD of the C α atoms were less than 0.455 Å over 300C α atoms among them. The RMSD value indicated that the overall structures of the four molecules of CD-mt5 were almost identical. Furthermore, the RMSD of the C α atoms were less than 0.530 Å on average over 300C α atoms between CD-WT and CD-mt5 in chain A. This result shows that the overall structures and the active site between WT and mt5 were almost identical. Figure 2B and Figure 4 show the structural differences between CD-WT and CD-mt5 for the five mutation sites (A117P, A151P, Q171P, A233P, and A254P) and the active site residues, Glu111 and Glu133. The structures of the main chain and side chains around the mutation points were not influenced by these mutations. This result shows that all positions for the introduction of proline residue into WT are ideal for the construction of thermostable enzymes.

It was proved that the additional mutations for G44C/K83C and Q250K/K253D/Q255R into mt5 were also ideal for the construction of the thermostable enzymes by crystal structural analysis. Therefore, all positive mutations (mt5, G44C/K83C, Q250K/K253D/Q255R) were integrated and inspected. Table 2 shows that these mutations do not influence the specific activity and the individual effect for thermostability caused by mutations was additive. Half-life at 60 °C was elongated by accumulating mutations: half-life at 60 °C of mt5, mt5/G44C/K83C (mt5ss), and mt5ss/Q250K/K253D/Q255R were about 7, 11 and 15 times longer than that of WT, respectively (Table 2 and Figure 6A).

mt5ss/Q250K/K253D/Q255R was proved to be the best enzyme for this study (Table 2). For the 250th position, however, histidine residue was observed in other GH19 chitinases (2DKV and 4DWX) (Figure 1). In addition, the alternative mutant mt5ss/Q250H/K253D/Q255R was examined (Table 2). There was no significant difference in half-life at 60 °C of mt5ss/Q250H/K253D/Q255R and mt5ss/Q250K/K254D/Q255R, but half-life at 65 °C of mt5ss/Q250H/K253D/Q255R was about two times longer than that of mt5ss/Q250K/K254D/Q255R (Table 2 and Figure 6B). mt5ss/Q250H/K253D/Q255R showed the highest T_m (71.1 °C: 6.9 °C higher than that of WT) among the mutants and elongated its half-life at 60 °C 15 times and at 65 °C 90 times more than that of WT, respectively (Table 2).

Comparison of the antifungal activity of the mutants

The antifungal activities of GlxChiB and its mutants were determined by using the hyphal re-extension inhibition assay with *Trichoderma viride* as the test fungus (Figure 5). We define IC_{50} as the concentration where 50% of the hyphal re-extension areas are inhibited. As shown in Table 2 and Figure 5, the IC_{50} of WT is $1.56 \pm 0.562 \mu M$, and all mutants exhibit the same level of WT for their antifungal activities. It shows that all thermostabilized mutations do not affect their antifungal activities significantly.

DISCUSSION

Obtaining thermostable mutants

It is ideal for efficient antifungal activity to degrade chitin for actual use, which is realized by having the enzyme perform under various conditions. Thus, enzymes must be resistant to high temperature, organic solvents, not-neutral pH, and other chemicals. Thermostabilized chitinases are expected to exhibit excellent performance under such conditions. GlxChiB obtained from the gazyumar latex has the highest antifungal activity among the other chitinases isolated from the latex and is expected to potentially be applied practically. While the optimum temperature for GlxChiB is 50 – 60 °C, this enzyme is unstable above 60 °C. Industrial use of this enzyme requires improving their thermostability without decreasing their enzymatic performance. However, there is often a

trade-off between thermostability and enzyme activity (29-31). In this study, we successfully obtained several thermostable mutants without decreasing their activities by rational design based on sequence and structural comparison among homologous enzymes. The thermostable GlxChiB created in this study is promising and has a high potential for its application. To date, two correlations about the primary sequences of an enzyme and its thermostability have been proved. One-point mutation can improve the thermostability of an enzyme, but this effect is in general minuscule (23, 32). The second is that such effects caused by individual structural changes from one-point mutations will be cumulatively counted and added (13-15, 33). These facts suggest that an enzyme is heat adaptable (34). Several studies (35-37) provided this trend about heat adaptation; however, a fundamental principle for the design of heat adaptable mutant enzymes has not been discovered (38).

Structural interpretation of the thermostability by Proline substitution

Proline is a unique amino acid residue in that the side chain is covalently bound to the preceding peptide bonded nitrogen and the five-membered ring imposes rigid constraints on the N-C α rotation (39). Thus, it is proposed that the substitution of proline for appropriate amino acids can increase the stability of a protein by decreasing the conformational entropy of unfolding (23, 37, 40), which is known as the “proline theory.” From the sequence homology among the four chitinases (Figure 1), we noticed the five residues where proline residue is conserved were in some other chitinases but not in GlxChiB, and we designed five mutants (A117P, A151P, Q171P, A233P, and A254P). All five mutants and integrated mutant, mt5, exhibited a positive effect on the thermostability (Table 2 and Figure 6). From the structural analysis (Figure 4), it was clarified that Ala117 is located at the second site of beta-turn I, Ala151 is located at the second site of beta-turn II, Gln171 is in the conserved helix structure, Ala233 and Ala254 are located at the loop region preceding conserved helix structure in CD of WT-GlxChiB, and Ala254 is located at N-terminal caps of α -helices. Proline residues contributing to thermal stabilization favor second sites of β -turns, loop regions, and N-terminal caps of α -helices (23, 36). Therefore, the construction of the thermostable mutants (A117P, A151P, A233P, and A254P) by the substitution of proline is well explained as the proline theory (36, 41).

Q171P is in a different circumstance; the residue is located in the middle of the alpha-helices structure, and proline residue is well known as the alpha-helices breaker (28). However, the Gly170 preceding the Pro171 appears to release the distorted structure by Pro171 (Figure 4C). In addition, Tyr147 and Trp200 exhibit the hydrophobic interaction toward Pro171 (Figure 7). The hydrophobic interaction appears to contribute to the thermostability of Q171P. For position 147, aromatic residues (Tyr, Phe, and Trp) are well conserved in the other chitinase, and Trp200 is also conserved in the enzyme (Figure 1). It is noteworthy that this hydrophobic interaction strengthens the connection between two helices and a loop region, contributing to the thermostability.

Structural interpretation of the thermostability by SS bond

A disulfide bridge linking the two cysteines stabilizes the folded state of the protein. From the crystal structural data of CD-WT and the sequence homology among the enzymes, we cannot find the suitable position in which the additional disulfide bonding can be introduced in the catalytic domain. However, it was estimated that the disulfide bonding between Gly44 (in the linker region) and Lys83 could form. In the DSC analysis, the melting temperature (T_m) of the mutant G44C/K83C was not higher than that of WT. However, the half-life at 60 °C was elongated without depressing their activities (Table 2 and Figure 6). By introducing the disulfide bonding, the thermostability of the enzyme is not improved; however, irreversible inactivation appears to be suppressed.

Structural interpretation of the thermostability by Salt bridge

It is well known that salt bridges contribute to the stability of proteins (42, 43). Ideally, the introduction of salt bridges can increase the thermostability of the enzyme. However, the fundamental problem is that the formation of salt bridges depends on the ionization properties of the participating groups, which are significantly influenced by the environmental changes around the proteins (44). Furthermore, salt bridges experience thermal fluctuations, continuously break and reform, and the flexibility of the protein governs their lifespan in solution. Nevertheless, proteins from thermophiles and hyperthermophiles exhibit more frequently networked salt bridges than proteins from the

mesophilic counterparts (43, 45, 46). Increasing the thermostability of proteins by optimizing charge-charge interactions is a good example of an evolutionary solution utilizing physical factors. For the two mutants with a salt bridge network introduced (Q250K/K253D/Q255R and Q250R/K253D/Q255R), specific activities at 37 °C were slightly increased. In addition, the half-life time at 60 °C was increased (Table2 and Figure 6). Figure 7B shows a model structure of the loop region (250Q-255Q) between two helices in the catalytic domain of mt5ss/Q250K/K253D/Q255R based on the crystal structure of mt5. It is speculated that the salt bridge networks among Lys250, Asp253, and Arg255 are constructed by inferring the salt bridge network in other GH19 structures (Figure S2). It should be noted that this region involves the proline substitution site (A254P). Thus, the performance of mt5ss/Q250K/K253D/Q255R is achieved by multiple effects of the hydrophobic interaction between Trp252 and Pro254, the stabilized peptide chain by Pro254, and the salt bridge network among Lys250, Asp253, and Arg255. In addition, Asp253 in mt5ss/Q250K/K253D/Q255R appears to form ASX turn preceding α -helices (Figure 1 and 7B) (47). The α -helix has an overall dipole moment due to the aggregate effect of the individual microdipoles. Therefore, α -helices often occur with the N-terminal end bound by a negatively charged group (48). It is speculated that Asp253 (negative charge) in mt5ss/Q250K/K253D/Q255R also increases the stability of the α -helices in the enzyme. We cannot explain why the thermostability of mt5ss/Q250H/K253D/Q255R is better than mt5ss/Q250K/K253D/Q255R. However, this 250th histidine residue is conserved in some chitinases (Figure 1), and an additional mechanism of the thermostability seems to be involved. The detailed study for the position is in progress.

Thermostable mutation effects on the catalytic and antifungal activity

The trade-off between the stability and function of the enzyme is widely recognized from the observation that the substitution of catalytic residues can dramatically improve its stability at the expense of its activity (29-31). Improvement of thermostability without reducing enzyme activity is highly demanded in industrial applications. We targeted the mutation sites that are apart from the catalytic residues (Figure 3). As a result, all mutations applied to GlxChiB in this study did not reduce its catalytic activity and antifungal activity

(Table 2). Furthermore, some mutants exhibit higher antifungal activity than WT. It has been reported that decreasing the conformational entropy of unfolding contributes to a resistance to protease (49-51). Therefore, some thermostable mutations appear to affect the activity itself but improve the resistance to protease secreted from the fungus, resulting in the improvement of the antifungal activity. In summary, we have succeeded in making the enzyme thermostable without decreasing its high catalytic activity and antifungal activity by using protein engineering methods. This study provides a successful strategy to improve the thermostability of GH19 chitinase and identifies the thermostable mutants of GlxChiB as a good seed for industrial applications.

MATERIALS AND METHODS

Construction of mutant protein genes

The genes encoding GlxChiB and its mutants were cloned into pET22b (Novagen) at the NdeI and BamHI restriction sites. All mutant genes were constructed by polymerase chain reactions using a QuickChange Site-Directed Mutagenesis kit (Stratagene) and primers (Table S1). Confirmation of the plasmid DNA sequences was outsourced to Fasmac Japan (Kanagawa, Japan).

Expression and purification of the recombinant protein

Plasmids encoding GlxChiB and its mutants were transformed into *E.coli* SHuffle T7 (DE3) cells (New England Biolabs). Cells harboring the plasmids were cultured at 37 °C in LB medium containing 100 mg/L ampicillin-Na for about 3.5 hours to reach OD600 of 0.6-0.8 and then were induced to express recombinant protein by adding IPTG to a final concentration of 0.1 mM. The culture was incubated for an additional 60 hours at 18 °C, and then the cells were harvested and disrupted by sonication in 20 mM Tris-HCl buffer, pH 8.0. The sonicated extract was separated into soluble and insoluble fractions by centrifugation at 12,000×g for 15 minutes at 4 °C. The soluble fraction was dialyzed against 10 mM sodium phosphate buffer, pH 7.0, and apply to a RESOURCE S column (6 mL, GE Healthcare) previously equilibrated with the same buffer. The elution was done with a linear gradient of NaCl from 0 to 0.3 M in the same buffer. The fractions containing

recombinant protein were collected and dialyzed against 10 mM sodium phosphate buffer, pH 7.0. The purity of the recombinant protein was analyzed by SDS-PAGE by the Laemmli method (52) using 12.5% polyacrylamide gels.

Protein assay

All protein concentrations were determined with the bicinchoninic acid (BCA) method (53). The protein concentration was determined with the Pierce BCA Protein Assay Kit (Thermo Scientific) using bovine serum albumin as the protein standard.

Chitinase activity assay

Chitinase activity was assayed colorimetrically with glycol chitin as a substrate. Glycol chitin was prepared by the method described by Yamada & Imoto (54). Ten microliters of the sample solution were added to 250 μ L of 0.2% (w/v) glycol chitin solution in 0.1 M sodium phosphate buffer, pH 7.0. After incubation of the reaction mixture at 37 °C for 15 minutes, the reducing power of the mixture was measured with ferric ferrocyanide reagent by the method of Imoto & Yagishita (55). One unit of activity was defined as the enzyme activity that produced 1 μ mol of GlcNAc per minute at 37 °C. The thermal stabilities of the enzymes were assessed by measuring the residual activities after incubation in 10 mM sodium phosphate buffer, pH 7.0 at 60 °C and 65 °C for the appropriate length of time. The residual activities were measured under the standard condition.

Quantitative antifungal activity assay (Taira et al., 2002) (56)

Hyphal re-extension inhibition assay was done by using *Trichoderma viride*. Agar disks (4 mm in diameter) containing the fungal hyphae, which were derived from the resting part of the fungus previously cultured on potato dextrose broth containing 1.5% (w/v) agar (PDA), were put on another PDA plate with the hyphae attached side down. Five microliters of sterile water or sample solution were overlaid on the agar disks, and then the plate was incubated at 25 °C. for 12 hours. After incubation, images of the plates were scanned using an image scanner. The areas of the re-extended hyphae were calculated as numbers of pixels by GNU Image Manipulation Program (GIMP, ver. 2.0). The protein concentration

required for inhibiting the growth of the fungus by 50% was defined as IC₅₀ and determined by constructing dose-response curves (percentage of growth inhibition versus protein concentration).

Differential scanning calorimetry (DSC)

The thermal stability of the enzymes was examined using differential scanning calorimetry (DSC). GlxChiB or its mutants in 10 mM sodium phosphate buffer (pH 7.0) were used at a final concentration of 1.0 mg/mL. A Nano DSC instrument (TA Instruments) was used at a scanning speed of 60 °C/h. Control runs in the absence of protein were carried out before and after each sample run. DSC scans in the presence of protein were performed two or three times for each protein examined.

Crystallization

The recombinant catalytic domain of the enzyme was prepared and purified as described previously (7). The purified proteins were dialyzed against 5 mM Tris-HCl buffer (pH 8.0) and concentrated to 10 mg/mL. Initial crystallization screening of the mutant protein was performed using various crystallization screening kits commercially available. The protein solution drop (0.15 µL) was mixed with 0.15 µL of a reservoir solution and then equilibrated with 50 µL of the reservoir solution. The crystallization was carried out according to the hanging-drop vapor diffusion method at 293 K in 96-well plates. After a week, well-formed crystals of CD-WT were obtained from the optimized condition (3% (w/v) gamma-polyglutamic acid low molecule, 25%(w/v) 2-Methy-2, 4-pentanediol, 0.1 M HEPES (pH 7.5), and 0.5 M Ammonium sulfate). Well-formed crystals of CD-mt5 were obtained from the different condition than CD-WT (18% (w/v) Polyethylene glycol 20K, 0.1 M sodium citrate (pH 5.0), and 3% (v/v) glycerol).

X-ray data collection

The crystal of CD-WT and CD-mt5 were frozen in liquid nitrogen. The diffraction data sets were collected at a wavelength of 1.000 Å at BL5A and NE3A beamline of the Photon

Factory in KEK, Japan, respectively. Data was processed by the program HKL2000 (57) for CD-WT and XDS (58) for CD-mt5.

Structure solution and refinement

General data handling was carried out with the CCP4 package (59). The initial model was solved by molecular replacement using Phaser (60) with the crystal structure of chitinase (GH19) from *Vigna unguiculata* (PDB: 4TX7) (17) as the search model for CD-WT. The model building was carried out with *Coot* (61) and refinement using REFMAC5 (62). Structural figures are described and rendered by the PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC.

Acknowledgments

This work was performed as a part of Projects of Okinawa innovation system building business of science and technology, supported by the Okinawa Science and Technology Promotion Center.

Author contributions

DK, HF and TK performed the experiments. DK and KI wrote the manuscript. KI, TT, TK, and KU thoroughly revised the manuscript. KI and TT designed and supervised the project. All authors read and approved the manuscript.

References

1. Oerke EC. 2005. Crop losses to pests. *The Journal of Agricultural Science* 144:31-43.
2. van Loon LC, Pierpoint WS, Boller T, Conejero V. 1994. Recommendations for naming plant pathogenesis-related proteins. *Plant Molecular Biology Reporter* 12:245-264.
3. Schlumbaum A, Mauch F, Vögeli U, Boller T. 1986. Plant chitinases are potent inhibitors of fungal growth. *Nature* 324:365-367.

4. Henrissat B, Bairoch A. 1993. New families in the classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem J* 293 (Pt 3):781-8.
5. Taira T. 2010. Structures and antifungal activity of plant chitinases. *J Appl Glycosci* 57:167-176.
6. Taira T, Ohdomari A, Nakama N, Shimoji M, Ishihara M. 2005. Characterization and Antifungal Activity of Gazyumaru (*Ficus microcarpa*) Latex Chitinases: Both the Chitin-Binding and the Antifungal Activities of Class I Chitinase Are Reinforced with Increasing Ionic Strength. *Bioscience, Biotechnology, and Biochemistry* 69:811-818.
7. Takashima T, Henna H, Kozome D, Kitajima S, Uechi K, Taira T. 2021. cDNA cloning, expression, and antifungal activity of chitinase from *Ficus microcarpa* latex: difference in antifungal action of chitinase with and without chitin-binding domain. *Planta* 253:120.
8. Choi JG, Ju YH, Yeom SJ, Oh DK. 2011. Improvement in the thermostability of D-psicose 3-epimerase from *Agrobacterium tumefaciens* by random and site-directed mutagenesis. *Appl Environ Microbiol* 77:7316-20.
9. Kohno M, Enatsu M, Funatsu J, Yoshiizumi M, Kugimiya W. 2001. Improvement of the optimum temperature of lipase activity for *Rhizopus niveus* by random mutagenesis and its structural interpretation. *J Biotechnol* 87:203-10.
10. Zhang W, Jia M, Yu S, Zhang T, Zhou L, Jiang B, Mu W. 2016. Improving the Thermostability and Catalytic Efficiency of the d-Psicose 3-Epimerase from *Clostridium bolteae* ATCC BAA-613 Using Site-Directed Mutagenesis. *J Agric Food Chem* 64:3386-93.
11. Cui Y, Chen Y, Liu X, Dong S, Tian Ye, Qiao Y, Mitra R, Han J, Li C, Han X, Liu W, Chen Q, Wei W, Wang X, Du W, Tang S, Xiang H, Liu H, Liang Y, Houk KN, Wu B. 2021. Computational Redesign of a PETase for Plastic Biodegradation under Ambient Condition by the GRAPE Strategy. *ACS*

Catalysis 11:1340-1350.

12. Musil M, Stourac J, Bendl J, Brezovsky J, Prokop Z, Zendulka J, Martinek T, Bednar D, Damborsky J. 2017. FireProt: web server for automated design of thermostable proteins. *Nucleic Acids Res* 45:W393-w399.
13. Watanabe M, Fukada H, Ishikawa K. 2016. Construction of Thermophilic Xylanase and Its Structural Analysis. *Biochemistry* 55:4399-409.
14. Nakabayashi M, Kamachi S, Malle D, Yanamoto T, Kishishita S, Fujii T, Inoue H, Ishikawa K. 2019. Construction of thermostable cellobiohydrolase I from the fungus *Talaromyces cellulolyticus* by protein engineering. *Protein Eng Des Sel* 32:33-40.
15. Ishikawa K, Kataoka M, Yanamoto T, Nakabayashi M, Watanabe M, Ishihara S, Yamaguchi S. 2015. Crystal structure of β -galactosidase from *Bacillus circulans* ATCC 31382 (BgaD) and the construction of the thermophilic mutants. *Febs j* 282:2540-52.
16. Hernández-Arana A, Rojo-Domínguez A, Soriano-García M, Rodríguez-Romero A. 1995. The thermal unfolding of hevein, a small disulfide-rich protein. *Eur J Biochem* 228:649-52.
17. Landim PGC, Correia TO, Silva FDA, Nepomuceno DR, Costa HPS, Pereira HM, Lobo MDP, Moreno FBMB, Brandão-Neto J, Medeiros SC, Vasconcelos IM, Oliveira JTA, Sousa BL, Barroso-Neto IL, Freire VN, Carvalho CPS, Monteiro-Moreira ACO, Grangeiro TB. 2017. Production in *Pichia pastoris*, antifungal activity and crystal structure of a class I chitinase from cowpea (*Vigna unguiculata*): Insights into sugar binding mode and hydrolytic action. *Biochimie* 135:89-103.
18. Matthews BW. 1968. Solvent content of protein crystals. *Journal of Molecular Biology* 33:491-497.
19. Ramachandran GN, Sasisekharan V. 1968. Conformation of polypeptides and proteins. *Adv Protein Chem* 23:283-438.
20. Kezuka Y, Kojima M, Mizuno R, Suzuki K, Watanabe T, Nonaka T. 2010. Structure of full-length class I chitinase from rice revealed by X-ray

- crystallography and small-angle X-ray scattering. *Proteins* 78:2295-305.
21. Ohnuma T, Numata T, Osawa T, Inanaga H, Okazaki Y, Shinya S, Kondo K, Fukuda T, Fukamizo T. 2012. Crystal structure and chitin oligosaccharide-binding mode of a 'loopful' family GH19 chitinase from rye, *Secale cereale*, seeds. *Febs j* 279:3639-3651.
22. Huet J, Rucktooa P, Clantin B, Azarkan M, Looze Y, Villeret V, Wintjens R. 2008. X-ray structure of papaya chitinase reveals the substrate binding mode of glycosyl hydrolase family 19 chitinases. *Biochemistry* 47:8283-91.
23. Matthews BW, Nicholson H, Becktel WJ. 1987. Enhanced protein thermostability from site-directed mutations that decrease the entropy of unfolding. *Proceedings of the National Academy of Sciences* 84:6663-6667.
24. Tidor B, Karplus M. 1993. The contribution of cross-links to protein stability: A normal mode analysis of the configurational entropy of the native state. *Proteins: Structure, Function, and Bioinformatics* 15:71-79.
25. Hazes B, Dijkstra BW. 1988. Model building of disulfide bonds in proteins with known three-dimensional structure. *Protein Engineering, Design and Selection* 2:119-125.
26. Arnott MA, Michael RA, Thompson CR, Hough DW, Danson MJ. 2000. Thermostability and thermoactivity of citrate synthases from the thermophilic and hyperthermophilic archaea, *Thermoplasma acidophilum* and *Pyrococcus furiosus*. *J Mol Biol* 304:657-68.
27. Christodoulou E, Rypniewski WR, Vorgias CR. 2003. High-resolution X-ray structure of the DNA-binding protein HU from the hyper-thermophilic *Thermotoga maritima* and the determinants of its thermostability. *Extremophiles* 7:111-22.
28. Tanaka T, Sawano M, Ogasahara K, Sakaguchi Y, Bagautdinov B, Katoh E, Kuroishi C, Shinkai A, Yokoyama S, Yutani K. 2006. Hyper-thermostability of CutA1 protein, with a denaturation temperature of nearly 150 degrees C. *FEBS Lett* 580:4224-30.
29. Beadle BM, Shoichet BK. 2002. Structural Bases of Stability-function

Tradeoffs in Enzymes. *Journal of Molecular Biology* 321:285-296.

30. Meiering EM, Serrano L, Fersht AR. 1992. Effect of active site residues in barnase on activity and stability. *Journal of Molecular Biology* 225:585-589.
31. Shoichet BK, Baase WA, Kuroki R, Matthews BW. 1995. A relationship between protein stability and protein function. *Proceedings of the National Academy of Sciences* 92:452-456.
32. Alber T. 1989. Mutational effects on protein stability. *Annual review of biochemistry* 58:765-798.
33. Wells JA. 1990. Additivity of mutational effects in proteins. *Biochemistry* 29:8509-8517.
34. Argos P, Rossman MG, Grau UM, Zuber H, Frank G, Tratschin JD. 1979. Thermal stability and protein structure. *Biochemistry* 18:5698-703.
35. Matthews BW. 1987. Genetic and structural analysis of the protein stability problem. *Biochemistry* 26:6885-6888.
36. Suzuki Y, Oishi K, Nakano H, Nagayama T. 1987. A strong correlation between the increase in number of proline residues and the rise in thermostability of five *Bacillus* oligo-1,6-glucosidases. *Applied Microbiology and Biotechnology* 26:546-551.
37. Watanabe K, Masuda T, Ohashi H, Mihara H, Suzuki Y. 1994. Multiple proline substitutions cumulatively thermostabilize *Bacillus cereus* ATCC7064 oligo-1,6-glucosidase. Irrefragable proof supporting the proline rule. *Eur J Biochem* 226:277-83.
38. Jaenicke R, Závodszky P. 1990. Proteins under extreme physical conditions. *FEBS Letters* 268:344-349.
39. MacArthur MW, Thornton JM. 1991. Influence of proline residues on protein conformation. *J Mol Biol* 218:397-412.
40. Schimmel PR, Flory PJ. 1968. Conformational energies and configurational statistics of copolypeptides containing l-proline. *Journal of Molecular Biology* 34:105-120.

- 584 41. Chou PY, Fasman GD. 1974. Prediction of protein conformation.
585 Biochemistry 13:222-245.
- 586 42. Kumar S, Nussinov R. 1999. Salt bridge stability in monomeric proteins.
587 Journal of molecular biology 293:1241-1255.
- 588 43. Xiao L, Honig B. 1999. Electrostatic contributions to the stability of
589 hyperthermophilic proteins. Journal of molecular biology 289:1435-1444.
- 590 44. Kumar S, Nussinov R. 2002. Relationship between ion pair geometries
591 and electrostatic strengths in proteins. Biophysical journal 83:1595-1612.
- 592 45. Dominy BN, Minoux H, Brooks III CL. 2004. An electrostatic basis for the
593 stability of thermophilic proteins. Proteins: Structure, Function, and
594 Bioinformatics 57:128-141.
- 595 46. Zhou H-X, Dong F. 2003. Electrostatic contributions to the stability of a
596 thermophilic cold shock protein. Biophysical journal 84:2216-2222.
- 597 47. Richardson JS. 1981. The anatomy and taxonomy of protein structure.
598 Adv Protein Chem 34:167-339.
- 599 48. Motoshima H, Mine S, Masumoto K, Abe Y, Iwashita H, Hashimoto Y,
600 Chijiiwa Y, Ueda T, Imoto T. 1997. Analysis of the stabilization of hen
601 lysozyme by helix macrodipole and charged side chain interaction. J
602 Biochem 121:1076-81.
- 603 49. Fontana A, Polverino de Laureto P, De Filippis V, Scaramella E, Zamboni
604 M. 1997. Probing the partly folded states of proteins by limited proteolysis.
605 Fold Des 2:R17-26.
- 606 50. Whitlow M, Bell BA, Feng SL, Filpula D, Hardman KD, Hubert SL,
607 Rollence ML, Wood JF, Schott ME, Milenic DE, et al. 1993. An improved
608 linker for single-chain Fv with reduced aggregation and enhanced
609 proteolytic stability. Protein Eng 6:989-95.
- 610 51. Parsell DA, Sauer RT. 1989. The structural stability of a protein is an
611 important determinant of its proteolytic susceptibility in *Escherichia coli*.
612 J Biol Chem 264:7590-5.
- 613 52. Laemmli UK. 1970. Cleavage of Structural Proteins during the Assembly

of the Head of Bacteriophage T4. *Nature* 227:680-685.

53. Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, Fujimoto EK, Goeke NM, Olson BJ, Klenk DC. 1985. Measurement of protein using bicinchoninic acid. *Anal Biochem* 150:76-85.
54. Yamada H, Imoto T. 1981. A convenient synthesis of glycolchitin, a substrate of lysozyme. *Carbohydr Res* 92:160-2.
55. Imoto T, Yagishita K. 1971. A Simple Activity Measurement of Lysozyme. *Agricultural and Biological Chemistry* 35:1154-1156.
56. Taira T, Ohnuma T, Yamagami T, Aso Y, Ishiguro M, Ishihara M. 2002. Antifungal Activity of Rye (*Secale cereale*) Seed Chitinases: the Different Binding Manner of Class I and Class II Chitinases to the Fungal Cell Walls. *Bioscience, Biotechnology, and Biochemistry* 66:970-977.
57. Otwinowski Z, Minor W. 1997. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol* 276:307-326.
58. Kabsch W. 2010. XDS. *Acta crystallographica Section D, Biological crystallography* 66:125-132.
59. Potterton E, Briggs P, Turkenburg M, Dodson E. 2003. A graphical user interface to the CCP4 program suite. *Acta Crystallogr D Biol Crystallogr* 59:1131-7.
60. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. 2007. Phaser crystallographic software. *J Appl Crystallogr* 40:658-674.
61. Emsley P, Cowtan K. 2004. Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* 60:2126-32.
62. Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, Winn MD, Long F, Vagin AA. 2011. REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr* 67:355-67.

Tables

Table 1. Data Collection and Refinement Statistics for CD-WT and CD-mt5

	CD-WT	CD-mt5
PDB Entry	7V91	7V92
Data collection		
Wavelength (Å)	1.0000	1.0000
Resolution (Å)	50.00 – 1.60 (1.63-1.60)	47.58-1.8 (1.70-1.60)
Space group	$P2_12_12_1$	$C2$
Unit cell dimensions (Å/°)	$a = 90.9, b = 106.58$ $c = 107.66$	$a = 140.58, b = 38.89$ $c = 177.74, \beta = 96.03$
No. of total reflections	1,000,279	799,385
No. of unique reflections	137,756	235,922
Rmerge*(%)	6.1 (38.1)	5.0 (39.9)
Completeness (%)	99.8 (99.7)	96.6 (88.5)
Redundancy	7.3 (3.8)	3.4 (3.16)
Average I/σ (I)	46.0 (5.6)	12.0 (2.3)
molecules/ASU	4	4
Matthews coefficient (Å ³ /Da)	2.5	2.32
Solvent (%)	51	47
Refinement		
Rwork**/Rfree*** (%)	22.7/25.3	21.3/24.3
RMS deviations from ideal values		
Bond angle (°)	1.400	1.365
Bond length (Å)	0.006	0.006
Ramachandran plot		
Favored (%)	98	97
Allowed (%)	2	2
Outliers (%)	0 (4 residues)	0 (2 residues)

The numbers in parentheses are for the outer shells.

* $R_{\text{merge}} = \sum_{hkl} \sum_j |I_{j(hkl)} - \langle I_{(hkl)} \rangle| / \sum_{hkl} \sum_j I_{j(hkl)}$, where $I_{j(hkl)}$ is the j th measurement of reflection hkl , including symmetry-related observations of a unique reflection, and $\langle I_{(hkl)} \rangle$ is the average.

** $R_{\text{work}} = \sum |F_o - F_c| / \sum F_o$ calculated for the working set.

*** $R_{\text{free}} = \sum |F_o - F_c| / \sum F_o$ calculated for the test set (approximately 5% of reflections that were excluded from the refinement).

Table 2. Summary of the enzymatic activities and thermodynamic parameters of the GlxChiB and its mutants.

			DSC*	Specific activity**	Antifungal activity	Half-life (min)	
			T _m (°C)	(U/mol)	(μM)	60°C	65°C
WT			64.2	5.40×10^9	1.56 ± 0.562	6.9 ± 0.51	0.7 ± 0.08
G44C/K83C	Disulfide bonding		64.2 (+ 0)	6.06×10^9	1.58 ± 0.200	14.5 ± 0.54	-
A117P	Pro substitution		65.7 (+ 1.5)	4.40×10^9	1.54 ± 0.310	13.8 ± 0.32	-
A151P	Pro substitution		64.8 (+ 0.6)	4.64×10^9	1.36 ± 0.064	9.9 ± 0.54	-
Q171P	Pro substitution		65.3 (+ 1.1)	5.40×10^9	1.32 ± 0.018	15.1 ± 0.59	-
A233P	Pro substitution		66.8 (+ 2.6)	5.56×10^9	1.40 ± 0.110	17.0 ± 0.46	-
A254P	Pro substitution		64.5 (+ 0.3)	4.59×10^9	1.49 ± 0.027	14.7 ± 0.89	-
Q250K/K253D/Q255R	Salt bridge		65.6 (+ 1.4)	7.38×10^9	1.21 ± 0.024	13.7 ± 1.20	-
Q250R/K253D/Q255R	Salt bridge		65.8 (+ 1.6)	7.50×10^9	1.06 ± 0.094	10.8 ± 0.19	-
mt5	All five Pro mutation		67.2 (+ 3.0)	4.74×10^9	0.69 ± 0.130	47.4 ± 1.39	3.3 ± 0.18
mt5ss	mt5/G44C/K83C		67 (+ 2.8)	5.88×10^9	0.69 ± 0.050	73.8 ± 2.49	3.9 ± 0.32
mt5ss/KDR	mt5ss/Q250K/K253D/Q255R		69.1 (+ 4.9)	4.31×10^9	1.58 ± 0.200	104.7 ± 7.21	33.2 ± 2.85
mt5ss/HDR	mt5ss/Q250H/K253D/Q255R		71.1 (+ 6.9)	5.13×10^9	1.94 ± 0.180	103.9 ± 3.34	63.3 ± 0.97

*Melting temperature (T_m) was measured using DSC followed by the procedures described as in the materials and methods.

**Specific activity was measured in the standard condition using 0.2% (w/v) glycol chitin as substrate; 37 °C for 15 minutes in 0.2 M sodium phosphate buffer (pH 7.0). One unit of activity was defined as the enzyme activity that produced 1 μmol of GlcNAc per minute at 37 °C.

Figures

Figure 1. A multiple sequence alignment of GH19 chitinases from plant.

Strictly conserved and similar residues are shown in white with a black and gray background, respectively. Two glutamic acid residues, which are conserved among GH19 chitinases and constructed catalytic sites are indicated with a blue background. α -helices and loops are represented as blue squares and black lines, respectively. All five proline

670 mutation sites, two cysteine mutation sites, and mutation sites for salt bridges are indicated
 671 with red boxes, purple boxes, and green boxes, respectively. All target sites are indicated
 672 with *. The abbreviations of proteins are used for this alignment as PDB ID as follows:
 673 2DKV, class-I chitinase from *Oryza sativa*; 4TX7, catalytic domain of class I chitinase
 674 from *Vigna unguiculata*; 4DWX, class II chitinase from *Secale cereal* (rye) seed; 3CQL,
 675 class II chitinase from *Carica papaya* (Papaya).

676

Figure 2. Crystal structure of GlxChiB CD-WT and CD-mt5.

(A) Superimposed models of four molecules of CD-WT (left) and CD-mt5 (right). Chains A-D of CD-WT and CD-mt5 are shown as cartoon loop models and colored gray, red, blue, black, magenta, cyan, orange, and green, respectively. (B) Structural alignment of CD-WT (gray) and CD-mt5 (magenta). α of them are shown as ribbon models. Two catalytic

glutamate residues (E111 and E133) and proline substitution sites (A117P, A151P, A171P, A233P, and A254P) are shown in stick models with atomic elements colors (O atoms, red; N atoms, blue; C atoms, gray in CD-WT and magenta in CD-mt5, respectively).

Figure 3. Positions of five proline substitutions and of introducing a disulfide bond and salt bridges.

All five proline mutations (red sphere), two cysteine mutations (purple sphere), and three mutations for salt bridges (green sphere) are located apart from the active sites (blue sphere). Positions are labeled with the corresponding residue in WT-GlxChiB.

Figure 4. Location of mutation sites from information of crystal structure.

Main chains of target regions and side chains of mutation sites are shown as a stick model with atomic element colors (O atoms, red; N atoms, blue; C atoms, gray in CD-WT and magenta in CD-mt5, respectively). A, Ala/Pro117 is located at second site of a beta-turn I. B, Ala/Pro151 is located at second site of a beta-turn II. C, Gln/Pro171 is located in conserved helix structure. D, Ala/Pro233 is located at the loop region preceding a

conserved helix structure. E, Ala /Pro254 is located at the loop region, preceding a conserved helix structure.

Figure 5. Antifungal activities of thermostable mutants.

Quantitative antifungal activity assays were performed as described in the methods. Five microliters of the sample were overlaid onto an agar disk containing the mycelium of *T. viride* on a PDA plate, and then, the plate was incubated at 25 °C for 12 hours. After incubation, the re-growth area of mycelia was measured. Open and closed circles indicate wild-type and mutants, respectively. (A-E) Proline substitution: A, A117P; B, A151P; C, Q171P; D, A233P; E, A254P. (F-H) Introducing disulfide bonding and salt bridges: F, G44C/K83C; G, Q250K/K253D/Q255R; H, Q250R/K253D/Q255R. (I-L) Integrated mutants: I, mt5; J, mt5ss; K, mt5ssKDR; L, mt5ssHDR. All assays were triplicate. Error bars represent $\pm$ SD (n = 3).

Figure 6. Thermostability of GlxChiB WT and mutants.

Enzyme samples in a 10 mM sodium phosphate buffer (pH 7.0) were incubated at 60 °C (A) and 65 °C (B) for each period. After the treatment, the residual activities (A_r) were

measured at 37 °C for 15 minutes in a 0.2 M sodium phosphate buffer (pH 7.0), with the initial activity without heat treatment taken as 100% (A_0). Open circle, closed circle, open square, closed square, open triangle, and closed triangle indicate WT, G44C/K83C, mt5, mt5/G44C/K83C, mt5/G44C/K83C/KDR, and mt5/G44C/K83C/HDR, respectively. Red dashed lines indicate where the value of $A_t/A_0 = 1/2$: the residual activity is 50% of initial activity. All assays were triplicate. Error bars represent $\pm$ SD ($n = 3$).

Figure 7. New interactions in thermostable mutants and the estimated salt bridges.

A, The hydrophobic interactions toward Pro171. The surface of Pro residue and two aromatic residues, Y147 and W156 were shown in gray. B, Structure of loop region from Q250 to Q255 in mt5. The salt bridge network among K250, D253, and R255 was estimated using a PyMOL mutagenesis option based on the structure of mt5 (the orientation of the side chains is adjusted to avoid steric hindrance). Side chains of original residues,

Q250, K253, and Q255 are shown as stick model colored transparent. The Estimated hydrogen bonding among K250, D253, and R255 are indicated as yellow dashed lines with labeled number as the distance (Å).

Supplemental information

Table S1. Site-directed mutagenesis primers.

Primer name		Template	Mutation	Sequence (5' to 3')
A117P	Fwd	WT	A117P	AAACTACCGGTGGATGGCCAAGTGCTCCTGATGGT
	Rev	WT	A117P	ACCATCAGGAGCACTTGGCCATCCACCGTAGTTT
A151P	Fwd	WT	A151P	TTATCCTTGTGCCCCTGGAAAGCAATATTATGGCCG
	Rev	WT	A151P	CGGCCATAATATTGCTTCCAGGGGCACAAGGATAA
Q171P	Fwd	WT	Q171P	GAACTACAACACGCGCCGTGTGAAAAGCCATAG
	Rev	WT	Q171P	CTATGGCTTTCCACACGCGCCGTAGTTGTAGTTC
A233P	Fwd	WT	A233P	GCCGGTCGAGTACCAGGATATGGTGC
	Rev	WT	A233P	GCACCATATCCTGGTACTCGACCGGC
A254P	Fwd	WT	A254P	CCAAGGTTGGAAGCCACAAGTGGAGGACC
	Rev	WT	A254P	GGTCCTCCACTTGTGGCTTCCAACCTTGG
G44C	Fwd	WT, mt5	G44C	AGCGCTTGCACATCAGCAAACTCATC
	Rev	WT, mt5	G44C	GATGTCGCAAGCGCTACTGCCACATTG
K83C	Fwd	WT, mt5	K83C	GCCGCCTGCGCCTTCCCTGGCTTTGGC
	Rev	WT, mt5	K83C	GAAGGCGCAGGCGCTGCGATGAAAGC
Q250K/K253D	Fwd	WT	Q250K/K253D	CAAAGGTTGGGACGCTCAGGTGGAGGACCG
	Rev	WT	Q250K/K253D	CGTCCCAACCTTTGCCACATTCAAGACCTC
Q250K/K253D/Q255R	Fwd	Q250K/K253D	Q255R	GACGCTCGGGTGGAGGACCGGATTGG
	Rev	Q250K/K253D	Q255R	CTCCACCCGAGCGTCCCAACCTTTGC
Q250R/K253D/Q255R	Fwd	Q250K/K253D/Q255R	K250R	CAAAGGTTGGGACGCTCGGGTGGAGGACCG
	Rev	Q250K/K253D/Q255R	K250R	CGTCCCAACCCGCGCCACATTCAAGACCTC
mt5ss/K253D/Q255R	Fwd	mt5ss	K253D/Q255R	GTTGGGATCCACGGGTGGAGGACCGGATTG
	Rev	mt5ss	K253D/Q255R	CCGTGGATCCCAACCTTGGCCACATTCAAG
mt5ss/Q250K/K253D/Q255R	Fwd	mt5ss/K253D/Q255R	Q250K	AAGGGTTGGGATCCACGGGTGGAGGACCGG
	Rev	mt5ss/K253D/Q255R	Q250K	ACCCGTGGATCCCAACCTTGGCCACATTC
mt5ss/Q250H/K253D/Q255R	Fwd	mt5ss/K253D/Q255R	Q250H	CATGGTTGGGATCCACGGGTGGAGGACCGG
	Rev	mt5ss/K253D/Q255R	Q250H	ACCCGTGGATCCCAACCATGGCCACATTC

Figure S1. Optimum temperature of WT and integrated mutants.

Specific activities were determined using 0.2% (w/v) glycol chitin as substrate in 0.2 M sodium phosphate buffer (pH 7.0) for 15 minutes. Open circles indicate WT. Open and closed squares indicate mt5ssKDR and mt5ssHDR, respectively.

A

GlxChiB mt5/KDR	250	K	G	W	D	P	R	V	E	D	R	I	G	F	Y	K	264
4DWX	206	H	G	Q	D	S	R	V	A	D	R	I	G	F	Y	K	220
4TX7	208	R	G	Q	D	S	R	V	E	D	R	I	G	F	F	K	222

750 **Figure S2. comparison of salt bridge network in other GH19 chitinases.**
751 A, Multiple sequence alignment among GlxChiB mt5/KDR, 4DWX, and 4TX7.
752 B, Structural comparison of the salt bridge network among these three GH19 chitinases.