Transcriptomics reveals the putative mycoparasitic strategy of the mushroom

Entoloma abortivum on species of the mushroom *Armillaria*

Running title: Mycoparasitism of *Armillaria* species by *Entoloma abortivum*

Rachel A. Koch¹ and Joshua R. Herr^{1,2,#}

¹Department of Plant Pathology, University of Nebraska, Lincoln, NE 68503, USA;

²Center for Plant Science Innovation, University of Nebraska, Lincoln, NE 68588, USA;

#Correspondence: Joshua R. Herr (jherr@unl.edu)

Abstract word count (including abstract and importance): 363

Text word count: 4938

ABSTRACT

During mycoparasitism, a fungus—the host—is parasitized by another fungus—the mycoparasite. The genetic underpinnings of these relationships have been best characterized in ascomycete fungi. However, within basidiomycete fungi, there are rare instances of mushroom-forming species parasitizing the reproductive structures, or sporocarps, of other mushroom-forming species, which have been rarely investigated on a genetic level. One of the most enigmatic of these occurs between *Entoloma abortivum* and species of *Armillaria*, where hyphae of *E. abortivum* are hypothesized to disrupt the development of *Armillaria* sporocarps, resulting in the formation of carpophoroids. However, it remains unknown whether carpophoroids are the direct result of a mycoparasitic relationship. To address the nature of this unique interaction, we analyzed gene expression of field-collected *Armillaria* and *E. abortivum* sporocarps and carpophoroids. Transcripts in the carpophoroids are primarily from *E. abortivum*, supporting the hypothesis that this species is parasitizing *Armillaria*. Most notably, we identified differentially upregulated *E. abortivum* β -trefoil-type lectins in the carpophoroid, which we hypothesize bind to *Armillaria* cell wall galactomannoproteins, thereby mediating recognition between the mycoparasite and the host. The most differentially upregulated *E. abortivum* transcripts in the carpophoroid code for oxalate decarboxylases—enzymes that degrade oxalic acid. Oxalic acid is a virulence factor in many plant pathogens, including *Armillaria* species, however, *E. abortivum* has evolved a sophisticated strategy to overcome this defense mechanism. The number of gene models and genes that code for carbohydrate-active enzymes in the *E. abortivum*

transcriptome were reduced compared to other closely related species, perhaps as a result of the specialized nature of this interaction.

IMPORTANCE

By studying fungi that parasitize other fungi, we can understand the basic biology of these unique interactions. Studies focused on the genetic mechanisms regulating mycoparasitism between host and parasite have thus far concentrated on a single fungal lineage within the Ascomycota. The work presented here expands our understanding of mycoparasitic relationships to the Basidiomycota and represents the first transcriptomic study to our knowledge that examines fungal-fungal relationships in their natural setting. The results presented here suggest that even distantly related mycoparasites utilize similar mechanisms to parasitize their host. Given that species of the mushroom-forming pathogen *Armillaria* cause plant root-rot diseases in many agroecosystems, an enhanced understanding of this interaction may contribute to better control of these diseases through biocontrol applications.

INTRODUCTION

Fungal mycoparasitism is a nutritional strategy where a living fungus – the host – is parasitized by and acts as a nutrient source for another fungus – the mycoparasite. Certain species of fungi in the Hypocreales (Ascomycota) are among the best-studied mycoparasites. Perhaps the most well-known of these are species of *Trichoderma* and *Clonostachys rosea*, which have biocontrol activity against plant pathogenic species of *Botrytis*, *Fusarium*, *Pythium*, and *Rhizoctonia* (1, 2). Other fungal mycoparasites in the Hypocreales include *Tolypocladium* species, many of which are parasites on the reproductive structures, or sporocarps, of species in the genus *Elaphomyces* (Eurotiales, Ascomycota) (3), *Escovopsis weberi*, which is a specialized necrotrophic parasite of fungal gardens of attine ants (4, 5) and *Hypomyces lactiflorum*, which parasitizes the mushrooms of *Russula* species and transforms them into the iconic “lobster mushroom” (6). Within the Basidiomycota, a well-studied genus of mycoparasites include *Tremella* species, which are parasites of corticiaceous basidiomycetes (7) and lichen-forming fungi (8).

Less studied examples of mycoparasitism involve mushroom-forming fungi that parasitize other mushroom-forming fungi. Fewer than 20 reported mushroom species may engage in this type of interaction, making it an incredibly rare phenomenon given the total number of mushroom-forming fungi (9, 10). Examples of this interaction whereby the parasite does not appear to impact the fitness of the host include *Volvariella surrecta*, which fruits from the pileus of its host, *Clitocybe nebularis* (11), and *Asterophora* species, which colonize their *Russula* or *Lactarius* host after it dies (10). More commonly, though, mushroom mycoparasites deform host sporocarps and likely

prevent the dispersal of their spores. *Pseudoboletus parasiticus* fruits from the sporocarps of *Scleroderma* species, which, after infection, are no longer able to mature and disperse their spores (12). *Psathyrella epimyces* causes the deformation of sporocarp tissue of its host, *Coprinus comatus* (13). Additionally, of the ten mushroom species in the genus *Squamanita*, all are known to be parasites of sporocarps of species in the genera *Cystoderma*, *Galerina*, *Hebeloma*, and *Inocybe* (10).

One of the most frequently encountered putative mycoparasitic interactions between two mushrooms involves species of *Armillaria* and *Entoloma abortivum* (Fig. 1). *Entoloma abortivum* is often encountered fruiting in soil, humus, or decaying logs in deciduous woods (14), while *Armillaria* species are facultative necrotrophs that can cause root rot in forest and agronomic systems worldwide (15, 16). *Entoloma abortivum* was originally described to occur as two morphologies: the mushroom form, which has a pinkish-gray stipe and pileus, and pink gills, and then the carpophoroid form, which is generally white and subglobose, and does not develop well-formed gills (Fig. 1). Initially, the carpophoroid form was assumed to be *E. abortivum* sporocarps that were malformed due to parasitism by *Armillaria* species (14). However, macro- and microscopic studies of carpophoroid collections determined that they are actually *Armillaria* sporocarps permeated with *E. abortivum* hyphae (17). Laboratory inoculation experiments showed that *E. abortivum* interacts with *Armillaria* sporocarps to disrupt their morphological development (17).

Whether carpophoroids are the result of a mycoparasitic relationship, where *E. abortivum* serves as the mycoparasite and *Armillaria* species serve as the host, remains unknown. To address this, we initiated a transcriptome study after we encountered all

three components of this system fruiting in close proximity—carpophoroids and individual sporocarps of *E. abortivum* and *Armillaria*. For a mycoparasite to be successful, there are several crucial steps to the utilization of a fungal host for nutrition. These steps include: 1) sensing the host; 2) recognition and attachment to host hypha; 3) initiating defense responses; and 4) the eventual demise of the host (18). Previous genomic and transcriptomic studies elucidated the genetic machinery that model mycoparasites utilize during each of these steps (18–20). In this work, we used transcriptomic and meta-transcriptomic techniques to analyze the genomic toolbox of *E. abortivum* and *Armillaria* during the carpophoroid stage. We show that the gene expression profiles of *E. abortivum* resemble those of known mycoparasitic species, as well as predict certain genes in both species that facilitate this interaction. Additionally, we used transcriptomic information to determine the species of *Armillaria* involved in this association.

RESULTS

Transcriptome assemblies of *Entoloma abortivum* and *Armillaria*. To benchmark gene diversity and baseline expression levels of the field-collected mushroom species in our study, we sequenced the sporocarp transcriptomes of *E. abortivum* and the *Armillaria* species found in close proximity to the carpophoroids. To date, there are several transcriptomic studies of *Armillaria* species (21, 22), but none for any *Entoloma* species. The assembled transcriptome of *E. abortivum* was just under 120 million base pairs. There were a total of 43,599 contigs and an N50 value of 3,527; 94.5% of benchmark universal single-copy orthologs (BUSCOs) from the Agaricales were present

in the *E. abortivum* transcriptome. Nine internal transcribed spacer (ITS) sequences were extracted from the transcriptome, accounting for 5.96×10^{-4} % to 1.82×10^{-3} % of the mapped reads; all ITS sequences were from *E. abortivum*. Within the contigs, a total of 9,728 unique gene models were recovered in the transcriptome assembly (Fig. 2) with 603 transcripts differentially upregulated in the carpophoroid tissue when compared to the sporocarp tissue and 403 transcripts differentially upregulated in the sporocarp tissue when compared to carpophoroid tissue (Fig. 3). The transcriptome contained 195 genes that code for carbohydrate-active enzymes (CAZymes). The transcriptome lacks any genes that code for cellobiohydrolases (glycoside hydrolase (GH) 6 and GH7), xylanases (GH10, GH11, GH30) and auxiliary proteins like polysaccharide monooxygenases (GH61) but does contain nine chitinases (GH18) (Fig. 2). Transcripts detected in the *E. abortivum* transcriptome that were previously determined to play important roles in mycoparasitic interactions (18–20) include: ten putative secondary metabolite gene clusters, one G-coupled protein receptor (GCPR), 38 ATP-binding cassette (ABC) transporters and 113 genes from the major facilitator superfamily (MFS) (Fig. 3). The average gene expression (in normalized units of trimmed mean of m-values (TMM)) of the ten most abundant *E. abortivum* transcripts in the sporocarp ranged from 4,333 to 17,890. Information about the ten most abundant transcripts in the sporocarps is available in Table 1.

The assembled transcriptome of *Armillaria* was just over 138 million base pairs. A total of 63,905 contigs, an N50 value of 2,845, and 97.8% of the BUSCOs representative of the Agaricales were present in the transcriptome. Eleven ITS sequences were extracted from the transcriptome, accounting for 2.77×10^{-5} % to

6.26×10⁻⁴ % of the mapped reads; six ITS sequences were from *Armillaria*, while five were from an uncharacterized species of the yeast *Kodamaea*. A total of 38,215 unique gene models were recovered, and, after bioinformatic filtering against public *Armillaria* genomes, 29,936 of these representing orthologs of *Armillaria* species (Fig. 2). In total, 2,619 transcripts were differentially upregulated in the carpophoroid tissue when compared to the sporocarp tissue, whereas 9,820 transcripts were differentially upregulated in the sporocarp tissue when compared to the carpophoroid tissue (Fig. 3). The transcriptome contained 580 genes that code for CAZymes, with 34 of those coding for chitinases (Fig. 2). Genes detected in the *Armillaria* transcriptome that might be important in mycoparasitic interactions include: 12 putative secondary metabolite gene clusters, five GCPRs, 59 ABC transporters and 144 MFS transcripts (Fig. 3). The average TMM of the ten most highly upregulated *Armillaria* transcripts in the sporocarp ranged from 8,438 to 36,477. The most differentially upregulated transcript was annotated as a cell wall galactomannoprotein (Fig. 4 and 5) (Table 2), while, notably, the fourteenth most differentially upregulated transcript in the sporocarp coded for isocitrate lyase (Table 2, Fig. 5).

Metatranscriptomic analysis of combined fungal hyphae in carpophoroid tissue.

We sequenced the metatranscriptome of the mixed tissue in the carpophoroids that are typically found when *E. abortivum* and species of *Armillaria* are found in proximity. In the carpophoroid tissue, significantly more transcriptomic reads from *E. abortivum* were identified than from *Armillaria* sp. ($t_{(8)} = 16.6$, $p = 1.77 \times 10^{-7}$, $n = 9$ replicates per species) (Fig. 6). The average number of *E. abortivum* mapped reads in the carpophoroid tissue

was 2,613,988, while the average number of *Armillaria* mapped reads was 74,880 (Fig. 6).

The average TMM of the top ten most abundant *E. abortivum* transcripts in the carpophoroid ranged from 9,075 to 68,720 (Table 1), while the average TMM of the top ten most differentially upregulated *Armillaria* transcripts in the carpophoroid ranged from 300 to 9,049 (Table 2). The first and third most highly upregulated *E. abortivum* transcripts in the carpophoroid tissue code for two oxalate decarboxylases, both of which were differentially upregulated in the carpophoroid tissue compared to the sporocarp (Figs. 3, 4, 5 and Table 1). There were other differentially upregulated *E. abortivum* transcripts in the carpophoroid that were not as highly upregulated that may also play a role in mycoparasitism. These include three β -trefoil-type lectins, three ABC transporters, two chitinases, and 14 MFS transcripts (Fig. 3, Table 1). We were unable to detect any transcripts involved in secondary metabolite gene clusters that were differentially upregulated in the carpophoroids (Fig. 3). Two *Armillaria* transcripts coding for putative senescence-associated proteins, as well as a heat shock protein associated with cell death, were differentially upregulated in the carpophoroid compared to the sporocarp (Table 2).

Phylogenetic placement of *Armillaria* reads. Phylogenomic analysis of 100 *Armillaria* BUSCOs generated in this study, in conjunction with previously published *Armillaria* genomes, shows a strongly supported sister relationship with an *A. mellea* specimen from France (100% BS) (Fig. 7A). Phylogenetic analysis of all ITS sequences

characterized as *A. mellea* from GenBank shows that this specimen is conspecific with specimens from eastern North America (Fig. 7B).

DISCUSSION

The formation of carpophoroids associated with species of *E. abortivum* have traditionally been thought to be the result of an *Armillaria* species attacking and parasitizing *Entoloma* sporocarps (14), hypothesized on the basis that *Armillaria* are wide-spread generalist forest pathogens that exhibit a broad range of host plants (15). However, subsequent studies suggested the opposite: the production of carpophoroids are the result of *E. abortivum* disrupting the development of *Armillaria* sporocarps (17). Here, we employed RNA sequencing and differential gene expression analysis on field-collected fungal tissue of each of the three components of this association to better understand the mechanistic basis of this interaction. We determined that *E. abortivum* reads in the metatranscriptome of the carpophoroid tissue—which can be interpreted as a measure of living tissue—are almost 35 times more abundant than *A. mellea* reads (Fig. 6). This finding suggests that carpophoroids are structures that result from *E. abortivum* parasitizing, and eventually killing, its *Armillaria* host under natural conditions.

Fungal-fungal necrotrophic mycoparasitic interactions are multistage processes that are best studied in model species, such as those in the genera *Trichoderma*, *Coniothyrium*, *Clonostachys* and *Tolypocladium* (20). Genomic and transcriptomic studies of necrotrophic mycoparasites show a convergence of significant genetic mechanisms at each stage (20). The *E. abortivum* genes that are differentially upregulated in the carpophoroid tissue are largely consistent with other examples of

necrotrophic mycoparasites in the Ascomycota. In the carpophoroid tissues we analyzed, *E. abortivum* appears to employ much of its energy on recognition and defense responses (Fig. 5). Inversely, the *Armillaria* sporocarps we analyzed illuminate possible mechanisms by which these two species recognize one another and how *Armillaria* responds to parasitism (Fig. 5).

The genetics of the Entoloma-Armillaria mycoparasitic interaction. A crucial step in a successful mycoparasite's life history is the ability to sense its host. Genes involved in the recognition of the fungal prey include those that code for GPCRs (18, 20). However, we did not find any of these genes that were differentially upregulated by *E. abortivum* in the carpophoroid tissue (Fig. 3). Given the significantly fewer number of *Armillaria* reads in the carpophoroid tissue compared to *E. abortivum*, we presume that these carpophoroids are relatively advanced in age, and expression of the genes used for sensing the presence of the host are no longer necessary.

We also identified three *E. abortivum* transcripts that code for β -trefoil-type lectins—proteins that bind to galactose units of sugar chains (23)—that were differentially upregulated in the carpophoroid tissue (Figs. 3 and 5). In well-studied mycoparasitic interactions, the recognition, attachment, and coiling around a fungal substrate is mediated by lectins expressed by at least one of the fungal partners (24–26). More specifically, basidiomycete β -trefoil-type lectins have previously been shown to play a role in the recognition of non-self glycans (27). Interestingly, the most abundant and differentially upregulated transcript produced in the *Armillaria* sporocarps, the substrate to which *E. abortivum* hyphae attach, codes for a cell wall

galactomannoprotein (Figs. 4 and 5). These proteins belong to a group of glycans which consist of a mannose backbone with galactose side chains and are known to make up a major part of the cell wall of some fungal species (28). This particular galactomannoprotein appears to be specific to *Armillaria* species. Watling (14) commented on the highly specific nature of this interaction and that it has only been documented occurring between *E. abortivum* and *Armillaria* species. One possible mediator of the specificity of this interaction could be the galactose sugars on the mannose protein (only known thus far from *Armillaria* species) that are the means by which *E. abortivum* β -trefoil-type lectins recognize and attach to its *Armillaria* host. However, more genome sequencing of other Agaricales species is needed to determine whether this protein is truly specific to species in the genus *Armillaria*.

During mycoparasitic interactions, the fungal host responds by mounting its own defense, and a successful mycoparasite must be able to cope with this counterattack (18, 20). Oxalic acid is a virulence factor employed by some plant pathogens, including species of *Armillaria*, to compromise the defense responses of the host plant by creating an acidic environment (29, 30). One differentially upregulated *Armillaria* transcript in the sporocarps codes for isocitrate lyase (Fig. 5), which is involved in OA biosynthesis in other fungal pathogens (31, 32), and we suspect that *Armillaria* sporocarps may utilize OA to defend against the initial infection by *E. abortivum*. However, two of the most abundant and differentially upregulated *E. abortivum* transcripts in the carpophoroid tissue code for oxalate decarboxylases—enzymes responsible for the degradation of OA (Figs. 3, 4, 5D, Table 1). In at least one known well-studied mycoparasitic interaction, OA is secreted by the fungal host, *Sclerotinia*

sclerotiorum, in reaction to penetration by its mycoparasite, *Coniothyrium minitans*. The acidic environment created by the secreted OA inhibits conidial germination and suppresses mycelial growth of *C. minitans* (33). However, *C. minitans* nullifies the growth-suppressing effects of OA or OA-mediated low pH by degrading the OA (34, 35), an enzymatic process largely mediated by oxalate decarboxylase. Because of this, oxalate decarboxylase plays an imperative role in mycoparasitism as OA degradation is vital for infection of the fungal host (36). Given the abundance of these genes produced by *E. abortivum* in the carpophoroid, we suspect a similar scenario in this mycoparasitic interaction. Subsequent downregulation of isocitrate lyase by *Armillaria* in the carpophoroids suggests that its genetic defense responses to infection were likely over at the time of our sampling.

Other ways that mycoparasites cope with the counterattack launched by their host include actively excreting host-secreted toxins. Here, we hypothesize that active extrusion of toxins secreted by the host occurs in the *E. abortivum* carpophoroid tissue via membrane transporters in the ABC superfamily (37–39). Three ABC transporters were differentially upregulated in the carpophoroid tissue (Fig. 3). Another group of genes that were differentially upregulated by *E. abortivum* in the carpophoroid belong to the major facilitator superfamily (MFS) transporters (Fig. 3). In *C. rosea*, there was selection for genes in this family that were related to drug resistance, and the transport of secondary metabolites, small organic compounds and carbohydrates (40). Their importance to mycoparasitism in *C. rosea* is predicted to invoke efflux-mediated protection against exogenous or endogenous secondary metabolites and nutrient uptake (40). MFS transporters have also been shown to be induced in other

mycoparasitic species (19, 41), but their exact biological roles have not been investigated.

In mycoparasitism, the final death of the host often results from the synergistic actions of cell wall-hydrolytic enzymes and antifungal secondary metabolites (18, 20). No secondary metabolite gene clusters identified in the *E. abortivum* transcriptome were differentially upregulated in the carpophoroid tissue (Fig. 3). In some mycoparasitic relationships, the secretion of secondary metabolites occurs early in the interaction, including in *Escovopsis weberi*, which secretes toxic compounds that kill the leafcutter ant garden before contact (5). In culture experiments between *Armillaria* isolates and *E. abortivum*, the growth of *Armillaria* was severely inhibited by the presence of *E. abortivum* (42). This suggests that *E. abortivum* may potentially secrete a toxic compound early in the interaction that inhibits the growth of *Armillaria*. Given that significantly more of the living tissue in the carpophoroids belonged to *E. abortivum* (Fig. 6), it is possible that much of the *Armillaria* tissue was killed preceding the full development of the carpophoroid.

Additionally, we hypothesize that the upregulated β -trefoil-type lectin in *E. abortivum* that may be important in hyphal recognition may also be cytotoxic towards *Armillaria*. This type of lectin has sequence homology, as well as putative structural similarity, to the B-subunit of ricin, a toxic protein from the castor bean *Ricinus communis* (43). An array of β -trefoil-type lectins have been characterized from the sporocarps of the mushroom-forming species *Clitocybe nebularis* (44), *Coprinus cinerea* (27), *Macrolepiota procera* (45) and *Boletus edulis* (46). Besides being important for non-self recognition, these same mushroom lectins also exhibit entomotoxic activity (47)

as well as nematotoxic activity (27, 45, 48). Taken together, it is possible that the *E. abortivum* β -trefoil-type lectins may also function as toxins towards *Armillaria*. While the *E. abortivum* transcripts coding for these lectins are not in the highest abundance in the carpophoroid tissue (Fig. 5), this could be because most of the *Armillaria* sporocarp tissue is already dead and the potential lethal effects produced by them are no longer necessary.

Chitin is an essential polymer in fungal cell walls (49) and is an important target during mycoparasitic attack (20). Indicative of the importance of chitinases in mycoparasitic interactions, members of the genus *Trichoderma*, as well as *Tolypocladium ophioglossoides* and *Escovopsis weberi*, have an increased number of genes coding for them (19, 50–53). Nine fungal chitinases were detected in the transcriptome of *E. abortivum*, which is fewer than the 13, 19, 20 and 29 detected in the closely related species, *Tricholoma matsutake* (54), *Clitocybe gibba* (55), *Lyophyllum atratum* (56) and *Lepista nuda* (55), respectively (Fig. 2). Only two of those *E. abortivum* chitinases were differentially upregulated in the carpophoroid tissue (Fig. 3) and were not abundant in comparison to other genes, suggesting minimal significance at this stage in carpophoroid development. One possibility for this difference in abundance could be the result of the putatively acidic pH in the carpophoroid that we infer based on the high gene expression of oxalate decarboxylases. In *C. minitans*, chitinase activity is positively correlated with ambient pH ranging from three to eight (57), so it is possible that chitinase activity in *E. abortivum* will increase after a neutral pH is restored. Another possibility for the low abundance is that most of the *Armillaria* host tissue was already broken down, reducing the need for chitinase activity.

Some of the putatively mycoparasitism-related genes outlined above were also differentially upregulated by *Armillaria* in the carpophoroid tissue. These include genes that code for MFS, ABC transporters, chitinases, and secondary metabolite gene clusters (Fig. 3). This suggests that *Armillaria* may be using many of the same genetic mechanisms to defend itself against parasitism by *E. abortivum*. Additionally, the degree of expression changes—both in the number of differentially upregulated transcripts and the log-fold change (logFC)—between the sporocarp and carpophoroid is much greater in *Armillaria* compared to *E. abortivum* (Fig. 3, 4), which could reflect an increase in the level of defense from *Armillaria*. However, this defense is apparently not enough to overcome the parasitic adaptations of *E. abortivum*.

Gene and CAZyme content of *E. abortivum*. The number of predicted gene models in the transcriptome of *E. abortivum* was 9,728, which is markedly fewer than the number of gene models in the genomes of its closest sequenced relatives (Fig. 2B). Additionally, relative to other closely related mushroom species, *E. abortivum* also exhibits a strong reduction in several gene families encoding CAZymes (Fig. 2C), and contains no cellobiohydrolases, xylanases, or polysaccharide monooxygenases. This finding is consistent with what has been observed in obligate mycoparasites and animal pathogens, which also have reduced CAZyme repertoires compared to fungi that utilize other nutritional strategies, hypothesized to be the result of their highly specialized interactions with their hosts (50, 58, 59). Therefore, it is possible that *E. abortivum* only retained the CAZymes and accessory genes necessary to interact with *Armillaria* species. While a genome sequence of *E. abortivum* will be necessary to confirm this

reduction, the BUSCO analysis verified that the *E. abortivum* transcriptome contains nearly 95% of the core set of eukaryotic genes, which suggests that our transcriptome is relatively complete, and it is unlikely that entire gene families are underrepresented.

Another possibility for the reduction in CAZymes could be explained by a broader nutritional strategy employed by *Entoloma* species, some of which form ectomycorrhizal-like structures on host plant species (60–62). Ectomycorrhizal species have a marked reduction in CAZymes in comparison to their saprotrophic ancestors (63), which we also observe with *Tricholoma matsutake* (Fig. 2C). However, microscopic analyses of *Entoloma* ectomycorrhizae-like structures suggest that some species destroy root meristems and young root cells, suggestive of a more parasitic relationship (60, 61).

One explanation is that *Entoloma* species, in general, are parasites of true ectomycorrhizae. More broadly, other species within the Entolomataceae are suspected mycoparasites, as they have been reported growing in close association with the sporocarps of other species (reviewed in (17)), including *Entoloma parasiticum* (= *Claudopus parasiticus*) and *Clitopilus fasciculatus* (64, 65). This explanation would also add credence to the evidence that Entolomataceae species are difficult to culture and are slow growing (62). Additional research utilizing genomes and laboratory studies to understand the nutritional strategy employed by this lineage will inform us as to whether fungal parasitism in this group is more common than it is currently understood to be. Lastly, it should be noted that *Entoloma* species, and *E. abortivum* in particular, form sclerotia in culture and presumably in soil (14, 17). These resting structures are perhaps a dormancy mechanism in the soil to survive seasonality when host tissue is unavailable. Because the timing and presence of *E. abortivum* fruitings have been hard

to predict, we were not able to observe or acquire a transcriptome for the sclerotia in the field.

Gene content and identity of *Armillaria* species in this interaction. The number of predicted gene models in the transcriptome of this *Armillaria* species was 38,215 (Fig. 2B). This is substantially greater than all other sequenced *Armillaria* species, which range from 14,473 to 25,704 gene models (22, 66). However, when we looked for gene models in our transcriptome that belong to known *Armillaria* species, this reduced the total number of gene models to 29,936 (Fig. 2B). The excess gene models in the *Armillaria* transcriptome, when compared to reference genomes, likely represent duplicated gene models with splice variants, a common artifact of transcriptome sequencing (67). Additionally, nearly one quarter of the gene models in the *Armillaria* sporocarps were from organisms other than *Armillaria*, including the yeast *Kodamaea*, highlighting the fact that field-collected sporocarps are not composed of tissue from a single organism. However, none of these contaminating organisms had genes that were both abundant and differentially upregulated that we predicted to play a role in this interaction.

Phylogenomic analysis of the *Armillaria* transcripts generated in this study suggest that the specific *Armillaria* species parasitized in this relationship is sister to an *A. mellea* specimen collected from western Europe (Fig. 7A). An ITS-based phylogenetic analysis shows the *Armillaria* specimen collected in this study is conspecific with other *A. mellea* collections from eastern North America (Fig. 7B). Before now, observations of *A. mellea sensu stricto*. fruiting in proximity to

carpophoroids (17) hinted it may be a host to *E. abortivum*. Herein, though, we show for the first time using genomic data that *A. mellea* s. s. can definitively serve as a host for *E. abortivum*. However, our hypothesis is that this interaction does not appear specific to just *A. mellea* s. s., as *A. gallica*, *A. ostoyae*, *A. jezoensis*, *A. sp. Nag. E* and *Desarmillaria tabescens* have been previously confirmed as hosts in this interaction as well (17, 42, 68). However, a broad geographic sampling of carpophoroids using molecular markers or genomic information could address this question more thoroughly. Interestingly, *Armillaria* species parasitized by *E. abortivum* appear to be only those present in Eastern North America and Eastern Asia (17, 42, 68).

Conclusions. Data from this study support the hypothesis that *E. abortivum* is a mycoparasite of *Armillaria* sporocarps. Three β -trefoil-type lectins are differentially upregulated by *E. abortivum* in the carpophoroid tissue, and we propose that these lectins mediate recognition with *Armillaria* sporocarps through binding to an *Armillaria*-specific galactomannoprotein. We hypothesize that by using oxalate decarboxylase, *E. abortivum* is likely defending against the secretion of OA by *Armillaria*. These strategies employed by *E. abortivum* for recognition and defense are similar to mechanisms utilized by other mycoparasites, suggesting that even distantly related mycoparasites utilize similar genetic mechanisms to mediate mycoparasitic interactions. One weakness of this study is that we were limited to the carpophoroid life stage that was available at the point of collection, which led us to speculate about what is occurring during other stages of mycoparasitism (i.e. sensing the host, initiating the interaction and killing and consuming the host). Therefore, future studies using culture methods,

isotopic analysis and metatranscriptomics of naturally collected carpophoroids at different life stages (i.e. younger and older carpophoroid specimens) will be necessary to completely tease apart the putative mycoparasitic strategies employed by *E. abortivum* and the defense responses by the *Armillaria* host. Finally, given that *Armillaria* species are pathogens in both natural and agronomic systems, a better understanding of this interaction may lead to the development of biocontrol methods for the control of *Armillaria* root rot.

MATERIALS AND METHODS

Sample collection, preparation, and sequencing. Sporocarps of *Armillaria* sp., *E. abortivum*, and the mixed-tissue carpophoroids were observed fruiting in proximity to one another on 18 September 2015 within the Baker Woodlot and Rajendra Neotropical Migrant Bird Sanctuary, Michigan State University, East Lansing, MI 48823 (42°42'56.4" N, 84°28'34.4" W) (collection accession: JRH 2446). Entire sporocarps were collected and immediately flash frozen in liquid nitrogen and subsequently stored at -80°C. At the time of processing, three biological replicates of each of the three tissue types (*Armillaria* sp. sporocarp, *E. abortivum* sporocarp, and carpophoroid), were individually ground in liquid N₂. Total RNA was then extracted from the ground tissue using the Qiagen RNeasy kit (Qiagen Inc., Hilden, Germany) according to manufacturer's protocol. RNA concentration and quality for each of the samples were assessed on a DeNovix DS-11 FX Spectrophotometer (DeNovix Inc., Wilmington, DE 19810, USA) and then shipped directly to the University of Minnesota's Genomics Center (<https://genomics.umn.edu>). Three technical replicates were sequenced for each

biological replicate. Transcriptomic and metatranscriptomic libraries were constructed with the TruSeq Standard Total RNA Library Preparation Kit with Ribo-Zero ribosomal reduction following the protocol developed by Schuierer et al. (69). Nucleotide sequencing was performed on the Illumina HiSeq 2500 System (Illumina Inc., San Diego, USA) and paired-end RNA sequence reads of 51 bp were generated for further analysis.

De novo transcriptome assembly, transcript abundance estimation and gene expression analysis. The quality of the raw reads was assessed using FastQC version 0.11.9 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc>). The range of the number of reads for each condition are as follows: *E. abortivum* sporocarps ranged from 10,584,302 to 14,473,328; *Armillaria* sporocarps ranged from 11,712,320 to 12,431,979; and the carpophoroids (containing reads from both organisms) ranged from 9,146,682 to 12,852,086. Sequencing adaptors were trimmed and PhiX contaminants were filtered for each sample using bbdut (<https://jgi.doe.gov/data-and-tools/bbtools/bb-tools-user-guide/bbdut-guide/>). Prior to transcriptome assembly, *k-mer* hash sizes were estimated with *khmer* (70). De novo assemblies were constructed independently for both *Armillaria* sp. and *E. arbotivum* with Trinity version 2.11.0 (71) using the trimmed reads generated from the respective sporocarp reads. Assembly statistics for both transcriptomes were generated with QUAST version 5 (72) and transcriptome completeness was assessed by determining the percentage of sequenced BUSCOs in each (73).

The results of the *de novo* transcriptome assemblies were used as references to perform sample-specific expression analysis. The trimmed sporocarp reads from each

of the nine replicates were mapped against their respective reference transcriptomes using Bowtie 2 (74) followed by calculation of abundance estimates using RSEM (75). The trimmed carpophoroid reads were also subsequently mapped, following the same protocol as described above, to both the *Armillaria* sp. and *E. abortivum* transcriptomes. Because of the close phylogenetic relatedness between these two species, and to filter out poorly aligned reads, we only retained mapped reads for all samples that had a MAPQ value of 30 and above, which is equivalent to reads that have a 99.9% chance of hitting the correct match. The R package edgeR (76) and “trimmed mean of M-value” (TMM) normalization (77) were used to determine differentially upregulated transcripts between: 1) *Armillaria* sporocarps and carpophoroids, and 2) *E. abortivum* sporocarps and carpophoroids. Transcripts were considered differentially upregulated if they had a logFC of two or greater and an FDR-adjusted *p*-value, or *q*-value, of < 0.05. All statistical analyses for the packages listed above were conducted using R version 4.0.3 (<http://www.r-project.org/>).

We used SAMtools (78) to determine the number of reads from the carpophoroids that mapped to our reference transcriptomes of *E. abortivum* and our particular *Armillaria* species. To understand whether the number of reads that mapped to the carpophoroids differed significantly between each fungal species, we performed an *F*-test of equality of variances, and then a two-tailed *t*-test assuming unequal variance with a *p*-value < 0.05 denoting significance.

Sporocarp Transcriptome and Carpophoroid Metatranscriptome Annotation. We annotated the *Armillaria* sp. and *E. abortivum* transcriptomes using Trinotate version 3.2

(79). Briefly, the transcripts were translated to coding protein sequences using TransDecoder version 5.5.0 (<http://transdecoder.github.io>) following identification of the longest open reading frames. To identify the most likely homologous sequence data, we used *blastx* on the transcripts and *blastp* on the predicted protein sequences (80). Using the predicted protein sequences, we also ran a HMMER (81) search against the PFAM database (82) to identify conserved domains that might be suggestive of function. We also compared these results to currently curated annotation databases such as Gene Ontology (GO) (83) and Kyoto Encyclopedia of Genes and Genomes (KEGG) (84–86). We used dbCAN2 (87) to annotate the CAZymes present in both species and compared their CAZy content to other closely related Agaricales species (22, 54–56), along with antiSMASH version 5.0 (88) to identify transcripts that belong to secondary metabolite gene clusters for both *Armillaria* and *E. abortivum*.

Finally, because this work was based off *de novo* transcriptome assemblies, there was the possibility that transcripts from contaminating organisms, including insects, yeasts, and other fungi, were also present and may be confounding the results. We approached this issue multiple ways. First, we extracted all internal transcribed spacer (ITS) regions from both transcriptomes, including the regions 18S, ITS1, 5.8S, ITS2, and 28S, using ITSx (89), and identified the origin of those sequences. To verify the number of mapped reads that represent rRNA, we also calculated the proportion of reads that mapped to the transcripts that contained the ITS sequences to the total number of mapped reads. Second, because high-quality genome assemblies of *Armillaria* exist (22, 66), we used Exonerate (90) to determine how many of the gene models in our *Armillaria* transcriptome are orthologs of *A. gallica*—the *Armillaria* species

with the largest number of known gene models. Finally, all genes that we report on as important to this interaction were manually verified to be from the target organism—*Armillaria* or *E. abortivum*—using the NCBI *blastx* tool (80) against the non-redundant protein sequences database and confirming a hit from the Physalacriaceae or Entolomataceae/Lyophyllaceae, respectively.

Phylogenetic analysis of *Armillaria* transcripts. In order to identify the specific species of *Armillaria* associated in this relationship, we identified BUSCOs (73) from the transcriptome of our *Armillaria* sporocarps along with other *Armillaria* and Physalacriaceae species with previously sequenced genomes (22, 66, 91, 92). We randomly selected 100 BUSCOs to reconstruct a phylogenomic tree from the six *Armillaria* specimens (22, 66), *Guyanagaster necrorhizus* (92) and *Cylindrobasidium torrendii* (91), which served as the outgroup. Protein-coding sequences were aligned using MAFFT version 7 (93) and non-informative sites and non-aligning regions were trimmed with Gblocks (94). The 100 BUSCOs were concatenated into a supermatrix with 64,436 sites. This supermatrix was used to infer a species tree and branch support using RAXML-NG (95), using a partitioned WAG+G model, where each data partition represented an individual BUSCO.

To expand on the phylogenomic analysis above, we used the representative *Armillaria* ITS sequence obtained from ITSx, and given the close relationship of our *Armillaria* species to *A. mellea*, we pulled all *A. mellea* ITS sequences from GenBank that included associated location metadata (96–110) (Table S1). These sequences were aligned using MAFFT version 7 (93), with refinements to the alignment performed

manually. RAxML-NG (95) was used to reconstruct this phylogeny. Taxa used to root this phylogeny included *A. gemina*, *A. sinapina*, *A. puiggarii* and *A. luteobubalina*—all members of the sister lineage to *A. mellea* *fide* (16).

Data and code availability. The raw reads and assembled transcriptomes generated during this study have been deposited in NCBI’s Gene Expression Omnibus and are accessible through GEO Series accession number GSE183699 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE183699>) or under the NCBI BioProject PRJNA761704. All other associated data, including analysis code and information on voucher collections, are available at https://github.com/HerrLab/Koch_Arma-Ento_2021.

Acknowledgements

We would like to give special thanks to Ben Lemmond and Eva Skific for providing photos of this system and to Greg Bonito at Michigan State University for generously storing sporocarp and carpophoroid tissue until we were able to transport the tissue to our laboratory. We also want to thank three anonymous reviewers who provided feedback and comments on the initial submission of this manuscript. This work was completed using the Holland Computing Center of the University of Nebraska, which receives support from the Nebraska Research Initiative. This research was directly supported by start-up funding from the University of Nebraska Agricultural Research Division and the University of Nebraska Office of Research and Economic Development. Additionally, JRH acknowledges funding from the US National Air &

Space Administration (Grant #80NSSC17K0737), the US National Science Foundation (EPSCoR Grant #1557417), and US National Institute of Justice (Grant #2017-IJ-CX-0025), all of which indirectly supported this research through the support of research in his laboratory. Funding agencies had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Conflict of Interest Statement

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Author Contributions

JRH initiated the work and sampled field collections; RAK extracted RNA from all the tissue samples and performed the laboratory work; RAK and JRH processed the experimental data, analyzed the data, designed the figures, and drafted the manuscript.

REFERENCES

1. Zeilinger S, Omann M. 2007. *Trichoderma* biocontrol: signal transduction pathways involved in host sensing and mycoparasitism. *Gene Regul Syst Bio* **1**:227–234.
2. Sun Z-B, Li S-D, Ren Q, Xu J-L, Lu X, Sun M-H. 2020. Biology and applications of *Clonostachys rosea*. *J Appl Microbiol* **129**:486–495.
3. Quandt CA, Kepler RM, Gams W, Araújo JPM, Ban S, Evans HC, Hughes D, Humber R, Hywel-Jones N, Li Z, Luangsa-Ard JJ, Rehner SA, Sanjuan T, Sato

- H, Shrestha B, Sung G-H, Yao Y-J, Zare R, Spatafora JW. 2014. Phylogenetic-based nomenclatural proposals for Ophiocordycipitaceae (Hypocreales) with new combinations in *Tolypocladium*. *IMA Fungus* **5**:121–134.
4. Currie CR, Mueller UG, Malloch D. 1999. The agricultural pathology of ant fungus gardens. *Proc Natl Acad Sci U S A* **96**:7998–8002.
5. Reynolds HT, Currie CR. 2004. Pathogenicity of *Escovopsis weberi*: the parasite of the attine ant-microbe symbiosis directly consumes the ant-cultivated fungus. *Mycologia* **96**:955–959.
6. Rogerson CT, Samuels GJ. 1994. Agaricolous species of *Hypomyces*. *Mycologia* **86**:839–866.
7. Zugmaier W, Bauer R, Oberwinkler F. 1994. Mycoparasitism of some *Tremella* species. *Mycologia* **86**:49–56.
8. Zamora JC, Millanes AM, Rico VJ, Pérez-Ortega S. 2015. Understanding lichenicolous heterobasidiomycetes: new taxa and reproductive innovations in *Tremella* s.l. *Mycologia* **108**:381–396.
9. Redhead SA, Ammirati JF, Walker GR, Norvell LL, Puccio MB. 1994. *Squamanita contortipes*, the Rosetta Stone of a mycoparasitic agaric genus. *Can J Bot* **72**:1814–1828.
10. Griffith GW, Piotr Gajda K, Detheridge AP, Douglas B, Bingham J, Turner A, Bowmaker V, Evans DA, McAdoo WG, Dentinger BTM. 2019. Strangler unmasked: Parasitism of *Cystoderma amianthinum* by *Squamanita paradoxa* and *S. pearsonii*. *Fungal Ecol* **39**:131–141.

11. Weber R, Webster J. 1996. *Volvariella surrecta*: An uncommon mycoparasite. *Mycologist* **10**:160.
12. Both EE. 2006. Personal encounters with the parasitic bolete. *Field Mycol* **7**:1–7.
13. Buller A. 1924. *Researches on Fungi*. Longmans, London, UK.
14. Watling R. 1974. Dimorphism in *Entoloma abortivum*. *Bull Soc Linn Lyon, Num Spec* **43**:449–470.
15. Baumgartner K, Coetzee MPA, Hoffmeister D. 2011. Secrets of the subterranean pathosystem of *Armillaria*. *Mol Plant Pathol* **12**:515–534.
16. Koch RA, Wilson AW, Séné O, Henkel TW, Aime MC. 2018. Resolved phylogeny and biogeography of the root pathogen *Armillaria* and its gasteroid relative, *Guyanagaster*. *BMC Evol Biol* **17**:33.
17. Lindner Czederpiltz DL, Volk TJ, Burdsall HH, Jr. 2001. Field observations and inoculation experiments to determine the nature of the carpophoroids associated with *Entoloma abortivum* and *Armillaria*. *Mycologia* **93**:841–851.
18. Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger S, Grigoriev IV, Kubicek CP. 2011. *Trichoderma*: the genomics of opportunistic success. *Nat Rev Microbiol* **9**:749–759.
19. Quandt CA, Di Y, Elser J, Jaiswal P, Spatafora JW. 2016. Differential expression of genes involved in host recognition, attachment, and degradation in the mycoparasite *Tolypocladium ophioglossoides*. *G3* **6**:731–741.
20. Karlsson M, Atanasova L, Jensen DF, Zeilinger S. 2017. Necrotrophic mycoparasites and their genomes. *Microbiol Spectr* **5**:FUNK-0016-2016.

21. Ross-Davis AL, Stewart JE, Hanna JW, Kim MS, Knaus BJ, Cronn R, Rai H, Richardson BA, McDonald GI, Klopfenstein NB. 2013. Transcriptome of an *Armillaria* root disease pathogen reveals candidate genes involved in host substrate utilization at the host–pathogen interface. *For Pathol* **43**:468–477.
22. Sipos G, Prasanna AN, Walter MC, O'Connor E, Bálint B, Krizsán K, Kiss B, Hess J, Varga T, Slot J, Riley R, Bóka B, Rigling D, Barry K, Lee J, Mihaltcheva S, LaButti K, Lipzen A, Waldron R, Moloney NM, Sperisen C, Kredics L, Vágvölgyi C, Patrignani A, Fitzpatrick D, Nagy I, Doyle S, Anderson JB, Grigoriev IV, Güldener U, Münsterkötter M, Nagy L. 2017. Genome expansion and lineage-specific genetic innovations in the forest pathogenic fungi *Armillaria*. *Nat Ecol Evol* **1**:1931–1941.
23. Dodd RB, Drickamer K. 2001. Lectin-like proteins in model organisms: implications for evolution of carbohydrate-binding activity. *Glycobiology* **11**:71R–79R.
24. Inbar J, Chet I. 1992. Biomimics of fungal cell-cell recognition by use of lectin-coated nylon fibers. *J Bacteriol* **174**:1055–1059.
25. Inbar J, Chet I. 1996. The role of lectins in recognition and adhesion of the mycoparasitic fungus *Trichoderma* spp. to its host. *Adv Exp Med Biol* **408**:229–231.
26. Neethling D, Nevalainen H. 1996. Mycoparasitic species of *Trichoderma* produce lectins. *Can J Microbiol* **42**:141–146.
27. Schubert M, Bleuler-Martinez S, Butschi A, Wälti MA, Egloff P, Stutz K, Yan S, Wilson IBH, Hengartner MO, Aebi M, Allain FHT, Künzler M. 2012. Plasticity of

- the β -trefoil protein fold in the recognition and control of invertebrate predators and parasites by a fungal defense system. *PLoS Pathog* **8**:e1002706.
28. Latgé JP, Kobayashi H, Debeaupuis JP, Diaquin M, Sarfati J, Wieruszeski JM, Parra E, Bouchara JP, Fournet B. 1994. Chemical and immunological characterization of the extracellular galactomannan of *Aspergillus fumigatus*. *Infect Immun* **62**:5424–5433.
29. Cessna SG, Sears VE, Dickman MB, Low PS. 2000. Oxalic acid, a pathogenicity factor for *Sclerotinia sclerotiorum*, suppresses the oxidative burst of the host plant. *Plant Cell* **12**:2191–2200.
30. Dumas MT, Boyonoski N, Jeng RS. 1989. Oxalic acid production by species of *Armillaria* in the boreal mixedwood forest. In: Proc. 7th International Conference on Root and Butt Rots. 9–16 Aug. 1988, Vernon and Victoria, British Columbia. Forestry Canada, Pacific Forestry Centre, Victoria, B.C. 680 p. 593–601.
31. Munir E, Jun Yoon J, Tokimatsu T, Hattori T, Shimada M. 2001. A physiological role for oxalic acid biosynthesis in the wood-rotting basidiomycete *Fomitopsis palustris*. *Proc Natl Acad Sci U S A* **98**:11126–11130.
32. Wang Y, Wang Y. 2020. Oxalic acid metabolism contributes to full virulence and pycnidial development in the poplar canker fungus *Cytospora chrysosperma*. *Phytopathology* **110**:1319–1325.
33. Wei SJ, Li GQ, Jiang DH, Wang DB. 2004. Effect of oxalic acid on spore germination and mycelial growth of the mycoparasite *Coniothyrium minitans*. *Acta Phytopathol Sin* **34**:199–203.

34. Ren L, Li GQ, Han YC, Jiang DH, Huang HC. 2007. Degradation of oxalic acid by *Coniothyrium minitans* and its effects on production and activity of β -1,3-glucanases of this mycoparasite. *Biol Control* **43**:1–11.
35. Ren L, Li GQ, Jiang DH. 2010. Characterization of some culture factors affecting oxalate degradation by the mycoparasite *Coniothyrium minitans*. *J Appl Microbiol* **108**:173–180.
36. Zeng L-M, Zhang J, Han Y-C, Yang L, Wu M-d, Jiang D-H, Chen W, Li G-Q. 2014. Degradation of oxalic acid by the mycoparasite *Coniothyrium minitans* plays an important role in interacting with *Sclerotinia sclerotiorum*. *Environ Microbiol* **16**:2591–2610.
37. Dubey M, Jensen DF, Karlsson M. 2015. The ABC transporter ABCG29 is involved in H₂O₂ tolerance and biocontrol traits in the fungus *Clonostachys rosea*. *Mol Genet Genom* **291**:677–686.
38. Ruocco M, Lanzuise S, Vinale F, Marra R, Turrà, Lois Woo S, Lorito M. 2009. Identification of a new biocontrol gene in *Trichoderma atroviride*: the role of an ABC transporter membrane pump in the interaction with different plant-pathogenic fungi. *Mol Plant Microbe In* **22**:291–301.
39. Zhao H, Zhou T, Xie J, Cheng J, Chen T, Jiang D, Fu Y. 2020. Mycoparasitism illuminated by genome and transcriptome sequencing of *Coniothyrium minitans*, an important biocontrol fungus of the plant pathogen *Sclerotinia sclerotiorum*. *Microb Genom* **6**:e000345.
40. Nygren K, Dubey M, Zapparata A, Iqbal M, Tzelepis GD, Brandström Durling M, Funck Jensen D, Karlsson M. 2018. The mycoparasitic fungus *Clonostachys*

- 711 *rosea* responds with both common and specific gene expression during

712 interspecific interactions with fungal prey. *Evol Appl* **11**:931–949.
- 713 41. Atanasova L, Le Crom S, Gruber S, Couplier F, Seidl-Seiboth V, Kubick CP,

714 Druzhinina IS. 2013. Comparative transcriptomics reveals different strategies of

715 *Trichoderma* mycoparasitism. *BMC Genom* **14**:121.
- 716 42. Cha JY, Igarashi T. 1996. Biological species of *Armillaria* and their mycoparasitic

717 associations with *Rhodophyllus abortivus* in Hokkaido. *Mycoscience* **37**:25–30.
- 718 43. Tregear JW, Roberts LM. 1992. The lectin gene family of *Ricinus communis*:

719 Cloning of a functional ricin gene and three lectin pseudogenes. *Plant Mol Biol*

720 **18**:515–525.
- 721 44. Pohleven J, Obermajer N, Sabotič J, Anžlovar S, Sepčič K, Kos J, Kralj B,

722 Štrukelj B, Brzin J. 2009. Purification, characterization and cloning of a ricin B-

723 like lectin from mushroom *Clitocybe nebularis* with antiproliferative activity

724 against human leukemic T cells. *Biochim Biophys Acta Gen Subj* **1790**:173–181.
- 725 45. Žurga S, Pohleven J, Renko M, Bleuler-Martínez S, Sosnowski P, Turk D,

726 Künzler M, Kos J, Sabotič J. 2014. A novel beta-trefoil lectin from the parasol

727 mushroom (*Macrolepiota procera*) is nematotoxic. *FEBS J* **281**:3489–3506.
- 728 46. Bovi M, Cenci L, Perduca M, Capaldi S, Carrizo ME, Civiero L, Chiarelli LR,

729 Galliano M, Monaco HL. 2013. BEL beta-trefoil: a novel lectin with antineoplastic

730 properties in king bolete (*Boletus edulis*) mushrooms. *Glycobiology* **23**:578–592.
- 731 47. Sabotič J, Ohm RA, Künzler M. 2016. Entomotoxic and nematotoxic lectins and

732 protease inhibitors from fungal fruiting bodies. *Appl Microbiol Biotechnol* **100**:91–

733 111.

48. Pohleven J, Renko M, Magister S, Smith DF, Künzler M, Strukelj B, Turk D, Kos J, Sabotič J. 2012. Bivalent carbohydrate binding is required for biological activity of *Clitocybe nebularis* lectin (CNL), the N, N'-diacetyllactosediamine (GalNAc β 1-4GlcNAc, LacdiNAc)-specific lectin from basidiomycete *C. nebularis*. *J Biol Chem* **287**:10602–10612.
49. Latgé JP. 2007. The cell wall: a carbohydrate armour for the fungal cell. *Mol Microbiol* **66**:279–290.
50. de Man TJ, Stajich JE, Kubicek CP, Teiling C, Chenthamara K, Atanasova L, Druzhinina IS, Levenkova N, Birnbaum SS, Barribeau SM, Bozick BA, Suen G, Currie CR, Gerardo NM. 2016. Small genome of the fungus *Escovopsis weberi*, a specialized disease agent of ant agriculture. *Proc Natl Acad Sci U S A* **113**:3567–3572.
51. Kubicek CP, Herrera-Estrella A, Seidl-Seiboth V, Martinez DA, Druzhinina IS, Thon M, Zeilinger S, Casas-Flores S, Horwitz BA, Mukherjee PK, Mukherjee M, Kredics L, Alcaraz LD, Aerts A, Antal Z, Atanasova L, Cervantes-Badillo MG, Challacombe J, Chertkov O, McCluskey K, Couplier F, Deshpande N, von Döhren H, Ebbole DJ, Esquivel-Naranjo EU, Fekete E, Flipphi M, Glaser F, Gómez-Rodríguez EY, Gruber S, Han C, Henrissat B, Hermosa R, Hernández-Oñate M, Karaffa L, Kostı I, Le Crom S, Lindquist E, Lucas S, Lübeck M, Lübeck PS, Margeot A, Metz B, Misra M, Nevalainen H, Omann M, Packer N, Perrone G, Uresti-Rivera EE, Salamov A, Schmoll M, Seiboth B, Shapiro H, Sukno S, Tamayo-Ramos JA, Tisch D, Wiest A, Wilkinson HH, Zhang M, Coutinho PM, Kenerley CM, Monte E, Baker SE, Grigoriev IV. 2011. Comparative genome

sequence analysis underscores mycoparasitism as the ancestral life style of

Trichoderma. *Genome Biol* **12**:R40.

52. Ihrmark K, Asmail N, Ubhayasekera W, Melin P, Stenlid J, Karlsson M. 2010.

Comparative molecular evolution of *Trichoderma* chitinases in response to

mycoparasitic interactions. *Evol Bioinform* **6**:1–26.

53. Xie BB, Qin QL, Shi M, Chen LL, Shu YL, Luo Y, Wang XW, Rong JC, Gong ZT,

Li D, Sun CY, Liu GM, Dong XW, Pang XH, Huang F, Liu W, Chen XL, Zhou BC,

Zhang YZ, Song ZY. 2014. Comparative genomics provides insights into

evolution of *Trichoderma* nutrition style. *Genome Biol Evol* **6**:379–390.

54. Miyauchi S, Kiss E, Kuo A, Drula E, Kohler A, Sánchez-García M, Morin E,

Andreopoulos B, Barry KW, Bonito G, Buée M, Carver A, Chen C, Cichocki N,

Clum A, Culley D, Crous PW, Fauchery L, Girlanda M, Hayes RD, Kéri Z, LaButti

K, Lipzen A, Lombard V, Magnuson J, Maillard F, Murat C, Nolan M, Ohm RA,

Pangilinan J, Pereira MF, Perotto S, Peter M, Pfister S, Riley R, Sitrit Y, Stielow

JB, Szöllősi G, Žifčáková L, Štursová M, Spatafora JW, Tedersoo L, Vaario LM,

Yamada A, Yan M, Wang P, Xu J, Bruns T, Baldrian P, Vilgalys R, Dunand C,

Henrissat B, Grigoriev IV, Hibbett D, Nagy LG, Martin FM. 2020. Large-scale

genome sequencing of mycorrhizal fungi provides insights into the early evolution

of symbiotic traits. *Nat Commun* **11**:5125.

55. Ruiz-Dueñas FJ, Barrasa JM, Sánchez-García M, Camarero S, Miyauchi S,

Serrano A, Linde D, Babiker R, Drula E, Ayuso-Fernández I, Pacheco R, Padilla

G, Ferreira P, Barriuso J, Kellner H, Castanera R, Alfaro M, Ramirez L, Pisabarro

AG, Riley R, Kuo A, Andreopoulos W, LaButti K, Pangilinan J, Tritt A, Lipzen A,

- He G, Yan M, Ng V, Grigoriev IV, Cullen D, Martin F, Rosso MN, Henrissat B, Hibbett D, Martínez AT. 2020. Genomic analysis enlightens Agaricales lifestyle evolution and increasing peroxidase diversity. *Mol Biol Evol* doi:10.1093/molbev/msaa301.
56. Steindorff AS, Carver A, Calhoun S, Stillman K, Liu H, Lipzen A, He G, Yan M, Pangilinan J, LaButti K, Ng V, Bruns TD, Grigoriev IV. 2020. Comparative genomics of pyrophilous fungi reveals a link between fire events and developmental genes. *Environ Microbiol* **23**:99–109.
57. Han Y-C, Li G-Q, Yang L, Jiang D-H. 2011. Molecular cloning, characterization and expression analysis of a *pacC* homolog in the mycoparasite *Coniothyrium minitans*. *World J Microbiol Biotechnol* **27**:381–391.
58. Ahrendt SR, Quandt CA, Ciobanu D, Clum A, Salamov A, Andreopoulos B, Cheng J-F, Woyke T, Pelin A, Henrissat B, Reynolds NK, Benny GL, Smith ME, James TY, Grigoriev IV. 2018. Leveraging single-cell genomics to expand the fungal tree of life. *Nat Microbiol* **3**:1417–1428.
59. Palmer JM, Drees KP, Foster JT, Lindner DL. 2018. Extreme sensitivity to ultraviolet light in the fungal pathogen causing white-nose syndrome of bats. *Nat Commun* **9**:35.
60. Agerer R, Waller K. 1993. Mycorrhizae of *Entoloma saepium*: parasitism or symbiosis? *Mycorrhiza* **3**:145–154.
61. Kobayashi H, Hatano K. 2001. A morphological study of the mycorrhiza of *Entoloma clypeatum* f. *hybridum* on *Rosa multiflora*. *Mycoscience* **42**:83–90.

62. Shishikura M, Takemura Y, Sotome K, Maekawa N, Nakagiri A, Endo N. 2021. Four mycelial strains of *Entoloma clypeatum* species complex form ectomycorrhiza-like roots with *Pyrus betulifolia* seedlings in vitro, and one develops fruiting bodies 2 months after inoculation. *Mycorrhiza* **31**:31–42.
63. Kohler A, Kuo A, Nagy LG, Morin E, Barry KW, Buscot F, Canbäck B, Choi C, Cichocki N, Clum A, Colpaert J, Copeland A, Costa MD, Doré J, Floudas D, Gay G, Girlanda M, Henrissat B, Herrmann S, Hess J, Högberg N, Johansson T, Khouja H-R, LaButti K, Lahrmann U, Levasseur A, Lindquist EA, Lipzen A, Marmeisse R, Martino E, Murat C, Ngan CY, Nehls U, Plett JM, Pringle A, Ohm RA, Perotto S, Peter M, Riley R, Rineau F, Ruytinx J, Salamov A, Shah F, Sun H, Tarkka M, Tritt A, Veneault-Fourrey C, Zuccaro A, Mycorrhizal Genomics Initiative Consortium, Tunlid A, Grigoriev IV, Hibbett DS, Martin F. 2015. Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal mutualists. *Nat Genet* **47**:410–415.
64. Noordeloos ME. 1987. *Entoloma* (Agaricales) in Europe. *Beih Nova Hedwigia* **91**:1–419.
65. Noordeloos ME. 1993. Studies in *Clitopilus* (Basidiomycetes, Agaricales) in Europe. *Persoonia* **15**:241–248.
66. Collins C, Keane TM, Turner DJ, O’Keeffe G, Fitzpatrick DA, Doyle S. 2013. Genomic and proteomic dissection of the ubiquitous plant pathogen, *Armillaria mellea*: Toward a new infection model system. *J Proteome Res* **12**:2552–2570.

67. Strickler SR, Bombarely A, Mueller LA. 2012. Designing a transcriptome next-generation sequencing project for a nonmodel plant species. *Am J Bot* **99**:257–266.
68. Fukuda M, Nakashima E, Hayashi K, Nagasawa E. 2003. Identification of the biological species of *Armillaria* associated with *Wynnea* and *Entoloma abortivum* using PCR-RFLP analysis of the intergenic region (IGR) of ribosomal DNA. *Mycol Res* **107**:1435–1441.
69. Schuierer S, Carbone W, Knehr J, Petitjean V, Fernandez A, Sultan M, Roma G. 2017. A comprehensive assessment of RNA-seq protocols for degraded and low-quantity samples. *BMC Genom* **18**:1–13.
70. Crusoe MR, Alameldin HF, Awad S, Boucher E, Caldwell A, Cartwright R, Charbonneau A, Constantinides B, Edverson G, Fay S, Fenton J, Fenzl T, Fish J, Garcia-Gutierrez L, Garland P, Gluck J, González I, Guermond S, Guo J, Gupta A, Herr JR, Howe A, Hyer A, Härpfer A, Irber L, Kidd R, Lin D, Lippi J, Mansour T, McA’Nulty P, McDonald E, Mizzi J, Murray KD, Nahum JR, Nanlohy K, Nederbragt AJ, Ortiz-Zuazaga H, Ory J, Pell J, Pepe-Rannek C, Russ ZN, Schwarz E, Scott C, Seaman J, Sievert S, Simpson J, Skennerton CT, Spencer J, Srinivasan R, Standage D, Stapleton JA, Steinman SR, Stein J, Taylor B, Trimble W, Wiencko HL, Wright M, Wyss B, Zhang Q, Zyme e, Brown CT. 2015. The khmer software package: enabling efficient nucleotide sequence analysis. *F1000Res* **4**:900.
71. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A,

Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq data. *Nat Biotechnol* **29**:644–652.

72. Mikheenko A, Pribelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics* **34**:i142–i150.

73. Seppey M, Manni M, Zdobnov EM. 2019. BUSCO: Assessing genome assembly and annotation completeness. In: Kollmar M (eds) Gene Prediction. Methods in Molecular Biology, vol 1962. Humana, New York, NW.

74. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:357–359.

75. Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform* **12**:323.

76. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139–140.

77. Robinson MD, Oshlack A. 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol* **11**:R25.

78. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:2078–2079.

79. Bryant DM, Johnson K, DiTommaso T, Tickle T, Brian Couger M, Payzin-Dogru D, Lee TJ, Leigh ND, Kuo T-H, Davis FG, Bateman J, Bryant S, Guzikowski AR, Tsai SL, Coyne S, Ye WW, Freeman RM, Jr, Peshkin L, Tabin CJ, Regev A,

Haas BJ, Whited JL. 2017. A tissue-mapped azolotl de novo transcriptome enables identification of limb regeneration factors. *Cell Rep* **18**:P762–P776.

80. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and application. *BMC Bioinform* **10**:421.

81. Wheeler TJ, Eddy SR. 2013. nhmmer: DNA homology search with profile HMMs. *Bioinformatics* **29**:3487–3489.

82. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M. 2014. Pfam: the protein families database. *Nucleic Acids Res* **42**:D222–D230.

83. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene Ontology: tool for the unification of biology. *Nat Genet* **25**:25–29.

84. Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* **28**:27–30.

85. Kanehisa M. 2019. Toward understanding the origin and evolution of cellular organisms. *Protein Sci* **28**:1947–1951.

86. Kanehisa M, Furumichi M, Sato Y, Ishiguro-Watanabe M, Tanabe M. 2021. KEGG: integrating viruses and cellular organisms. *Nucleic Acids Res* **49**:D545–D551.

87. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* **46**:W95–W101.

88. Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Yup Lee S, Medema MH, Weber T. 2019. antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res* **47**:W81–W87.
89. Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De Wit P, Sánchez-García M, Ebersberger I, de Sousa F, Amend A, Jumpponen A, Unterseher M, Kristiansson E, Abarenkov K, Bertrand YJK, Sanli K, Eriksson KM, Vik U, Veldre V, Nilsson RH. 2013. Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods Ecol Evol* **4**:914–919.
90. Slater GSC, Birney E. 2005. Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics* **6**:31.
91. Floudas D, Held BW, Riley R, Nagy LG, Koehler G, Ransdell AS, Younus H, Chow J, Chiniquy J, Lipzen A, Tritt A, Sun H, Haridas S, LaButtie K, Ohm RA, Kües U, Blanchette RA, Grigoriev IV, Minto RE, Hibbett DS. 2015. Evolution of novel wood decay mechanisms in Agaricales revealed by the genome sequences of *Fistulina hepatica* and *Cylindrobasidium torrendii*. *Fungal Genet Biol* **76**:78–92.
92. Koch RA, Yoon GM, Arayal UK, Lail K, Amirebrahimi M, LaButti K, Lipzen A, Riley R, Barry K, Henrissat B, Grigoriev IV, Herr JR, Aime MC. 2021. Symbiotic nitrogen fixation in the reproductive structures of a basidiomycete fungus. *Curr Biol* **31**:3905–3914.
93. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**:772–780.

94. Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* **17**:540–552.
95. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAXML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**:4453–4455.
96. Guo T, Wang HC, Xue WQ, Zhao J, Yang ZL. 2016. Phylogenetic analyses of *Armillaria* reveal at least 15 phylogenetic lineages in China, seven of which are associated with cultivated *Gastrodia elata*. *PLoS One* **11**:e0154794.
97. Coetzee MPA, Wingfield BD, Harrington TC, Dalevi D, Coutinho TA, Wingfield MJ. 2000. Geographical diversity of *Armillaria mellea* s. s. based on phylogenetic analysis. *Mycologia* **92**:105–113.
98. Coetzee MPA, Wingfield BD, Bloomer P, Wingfield MJ. 2005. Phylogenetic analyses of DNA sequences reveal species partitions amongst isolates of *Armillaria* from Africa. *Mycol Res* **109**:1223–1234.
99. Schneider-Maunoury L, Deveau A, Moreno M, Todesco F, Belmondo S, Murat C, Courty P-E, Jąkowski M, Selosse M-A. 2020. Two ectomycorrhizal truffles, *Tuber melanosporum* and *T. aestivum*, endophytically colonise roots of non-ectomycorrhizal plants in natural environments. *New Phytol* **225**:2542–2556.
100. Coetzee MPA, Bloomer P, Wingfield MJ, Wingfield BD. 2011. Paleogene radiation of a plant pathogenic mushroom. *PLoS One* **6**:e28545.
101. Osmundson TW, Robert VA, Schoch CL, Baker LJ, Smith A, Robich G, Mizzan L, Garbelotto MM. 2013. Filling gaps in biodiversity knowledge for

- 937 macrofungi: contributions and assessment of an herbarium collection DNA
- 938 barcode sequencing project. *PLoS One* **8**:e62419.
- 939 102. Kasper-Pakosz R, Pietras M, Łuczaj Ł. 2016. Wild and native plants and
- 940 mushrooms sold in the open-air markets of south-eastern Poland. *J Ethnobiol*
- 941 *Ethnomed* **12**:45.
- 942 103. Hughes KW, Petersen RH, Lodge DJ, Bergemann SE, Baumgartner K,
- 943 Tulloss RE, Lickey E, Cifuentes J. 2013. Evolutionary consequences of putative
- 944 intra- and interspecific hybridization in agaric fungi. *Mycologia* **105**:1577–1594.
- 945 104. Potyralska A, Schmidt O, Moreth U, Łakomy P, Siwecki R. 2002. rDNA-
- 946 ITS sequence of *Armillaria* species and a specific primer for *A. mellea*. *For Genet*
- 947 **9**:119–124.
- 948 105. Ortega A, Moreno G, Platas G, Pelaez F. 2008. A new *Armillaria mellea*
- 949 variant from tops of coastal sand dunes in southern Spain. *Mycotaxon* **104**:349–
- 950 364.
- 951 106. Vu D, Groenewald M, de Vries M, Gehrman T, Stielow B, Eberhardt U,
- 952 Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW,
- 953 Robert V, Verkley GJM. 2019. Large-scale generation and analysis of
- 954 filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals
- 955 thresholds for fungal species and higher taxon delimitation. *Stud Mycol* **92**:135–
- 956 154.
- 957 107. Binder M, Hibbett DS, Wang Z, Farnham WF. 2006. Evolutionary
- 958 relationships of *Mycaureola dilseae* (Agaricales), a basidiomycete pathogen of a
- 959 subtidal rhodophyte. *Am J Bot* **93**:547–556.

108. Haelewaters D, Dirks AC, Kappler LA, Mitchell JK, Quijada L, Vandegrift R, Buyck B, Pfister DH. 2018. A preliminary checklist of fungi at the Boston Harbor Islands. *Northeast Nat* **25**:45–76.
109. Kim M-S, Klopfenstein NB, Hanna JW, McDonald GI. 2006. Characterization of North American *Armillaria* species: genetic relationships determined by ribosomal DNA sequences and AFLP markers. *For Path* **36**:145–164.
110. Schnabel G, Ash JS, Bryson PK. 2005. Identification and characterization of *Armillaria tabescens* from the southeastern United States. *Mycol Res* **109**:1208–1222.
111. Dentinger BTM, Gaya E, O'Brien H, Suz LM, Lachlan R, Díaz-Valderrama JR, Koch RA, Aime MC. 2016. Tales from the crypt: genome mining from fungarium specimens improves resolution of the mushroom tree of life. *Biol J Linn Soc* **117**:11–32.

TABLE 1 Ten most abundant *E. abortivum* transcripts in sporocarps and carpophoroids

Gene	Annotation	Cond.	C (TMM)	S (TMM)	logFC	q-value
ENT_DN2762_c0_g3	Oxalate decarboxylase	C	68,724	238	-8.5	4.5×10^{-317}
ENT_DN1212_c0_g1	Hypothetical protein	C/S	55,939	12,590	-2.5	4.3×10^{-49}
ENT_DN3063_c0_g2	Oxalate decarboxylase	C	48,157	135	-8.8	0
ENT_DN1952_c0_g2	Acid phosphatase	C	19,999	677	-5.2	1.3×10^{-169}
ENT_DN4045_c1_g1	Hypothetical protein	C	19,965	3,374	-2.9	3.4×10^{-63}
ENT_DN5742_c0_g1	Hypothetical protein	C	11,229	23	-9.3	6.9×10^{-220}
ENT_DN4332_c0_g1	Hypothetical protein	C	11,038	607	-4.5	4.7×10^{-77}
ENT_DN1952_c0_g1	Acid phosphatase	C	10,766	673	-4.4	1.9×10^{-122}
ENT_DN2936_c0_g2	Hypothetical protein	C	9,212	3,686	NS	
ENT_DN3086_c0_g1	Hypothetical protein	C	9,076	1,563	-2.9	2.2×10^{-62}
ENT_DN1375_c0_g1	Trehalase	S	1,934	17,890	2.9	6.5×10^{-37}
ENT_DN5247_c0_g3	Hypothetical protein	S	2,718	7,204	NS	
ENT_DN761_c0_g2	Hypothetical protein	S	7,405	6,918	NS	
ENT_DN521_c0_g2	Hypothetical protein	S	6,045	6,424	NS	
ENT_DN852_c0_g1	Hypothetical protein	S	123	5,429	5.1	2.0×10^{-142}
ENT_DN3621_c0_g2	RNA polymerase	S	1,977	5,138	NS	
ENT_DN6707_c0_g1	Hypothetical protein	S	1,654	5,027	NS	
ENT_DN2379_c0_g1	TPR-like protein	S	4,862	5,017	NS	
ENT_DN466_c0_g1	Auxin efflux carrier	S	1,205	4,333	NS	

Cond.: condition in which each gene was most abundant, referring to either the carpophoroids (C) or sporocarps (S). NS: not significantly differentially upregulated. Negative logFC values are significantly differentially upregulated in the carpophoroid, whereas positive values are significantly differentially upregulated in the sporocarp. Transcripts discussed in the text include two oxalate decarboxylases (ENT_DN2762_c0_g3, ENT_DN3063_c0_g2), three β -trefoil-type lectins (ENT_DN4359_c0_g1, ENT_DN1877_c0_g1, ENT_DN4255_c0_g1), three ABC transporters (ENT_DN1537_c0_g1, ENT_DN189_c0_g1, ENT_DN3860_c0_g1), two chitinases (ENT_DN2096_c0_g1, ENT_DN4507_c0_g3) and 14 MFS transcripts (ENT_DN409_c0_g1, ENT_DN1954_c0_g1, ENT_DN1998_c0_g1,

1003 ENT_DN2070_c0_g1, ENT_DN2744_c0_g1, ENT_DN3292_c0_g1, ENT_DN3474_c0_g1, ENT_DN3861_c0_g1, ENT_DN3943_c0_g1,
1004 ENT_DN3943_c0_g2, ENT_DN3981_c0_g1, ENT_DN4751_c0_g2, ENT_DN6588_c0_g1, ENT_DN6695_c0_g1).
1005

TABLE 2 Ten most abundant *Armillaria* transcripts in sporocarps and carpophoroids

Gene	Annotation	Cond.	C (TMM)	S (TMM)	logFC	q-value
ARM_DN1755_c2_g1	Cell wall galactomannoprotein	S	0	36,479	13.5	4.0×10^{-182}
ARM_DN3840_c0_g1	Serine carboxypeptidase	S	12	29,472	6.7	7.9×10^{-100}
ARM_DN22943_c1_g1	Rab geranylgeranyltransferase	S	21	18,649	5.1	3.8×10^{-89}
ARM_DN1737_c0_g1	LysM-domain-containing protein	S	0	14,557	15.9	3.0×10^{-159}
ARM_DN1699_c3_g1	Hypothetical protein	S	9	12,848	5.8	1.6×10^{-94}
ARM_DN1314_c0_g1	Chondroitin AC/alginate lyase	S	102	11,436	2.2	3.7×10^{-25}
ARM_DN20980_c0_g1	Glycopeptide	S	0	10,449	15.4	3.4×10^{-139}
ARM_DN8135_c0_g1	Rasp f 7 allergen	S	0	9,014	11.5	1.8×10^{-58}
ARM_DN4971_c0_g2	Aldehyde dehydrogenase	S	103	8,688	NS	
ARM_DN1205_c0_g1	Hypothetical protein	S/C	9,049	8,348	-4.8	2.1×10^{-74}
ARM_DN5170_c0_g1	Hypothetical protein	C	2,086	377	-7.1	6.9×10^{-150}
ARM_DN23207_c0_g2	Senescence-associated	C	1,711	2,482	-4.1	6.6×10^{-61}
ARM_DN996_c0_g1	Senescence-associated	C	1,423	1,090	-5.0	3.0×10^{-73}
ARM_DN409_c1_g1	Elongation factor 1-alpha	C	769	1,154	-4.1	4.0×10^{-155}
ARM_DN1893_c0_g1	Hypothetical protein	C	722	58	-8.3	5.3×10^{-109}
ARM_DN693_c0_g1	Hypothetical protein	C	520	840	-3.9	1.1×10^{-48}
ARM_DN1222_c0_g1	CYS3-cystathionine gamma-lyase	C	485	717	-4.1	5.5×10^{-27}
ARM_DN1146_c0_g4	Heat shock protein 70	C	431	2,361	-2.2	1.7×10^{-54}
ARM_DN3800_c0_g1	ATP synthase F1	C	300	4,686	NS	

Cond.: condition in which each gene was most abundant, referring to either the carpophoroids (C) or sporocarps (S). NS: not significantly differentially upregulated. Negative logFC values are significantly differentially upregulated in the carpophoroid, whereas positive values are significantly differentially upregulated in the sporocarp. Transcripts discussed in the text were an isocitrate lyase (ARM_DN1_c7_g1), two putative senescence-associated proteins (ARM_DN23207_c0_g2, ARM_DN996_c0_g1), and a heat shock protein (ARM_DN1146_c0_g4).

FIG 1 The components of this fungal interaction in nature. (A) All three components—*Armillaria* and *E. abortivum* sporocarps, along with carpophoroids—fruiting synchronously. (B) A close-up photo of all of the components, with an *Armillaria* sporocarp indicated by the yellow arrow, an *E. abortivum* sporocarp indicated by the light pink arrow and a carpophoroid indicated by the magenta arrow. Scale bars in A are equal to 10 cm and in B are equal to 1 cm. Photos taken by Ben Lemmond.

FIG 2 Number of total gene models and gene copy numbers of CAZymes. (A) A representative phylogeny based off phylogenetic evidence from (22) and (111). (B) Filled in, colored dots represent the number of fungal gene models in the transcriptomes of *Armillaria* (yellow) and *E. abortivum* (pink) generated in this study, along with those of closely related species from previously generated genomic data (filled in black dots). The dashed circle associated with the *Armillaria carpophoroid* represents the total number of identified gene models (inclusive of additional non-*Armillaria* genes), showing that nearly one quarter of the total gene models in the *Armillaria* transcriptome were from organisms other than the target. (C) Heat map showing gene copy numbers of plant cell wall degrading enzymes detected in the transcriptomes of *E. abortivum* and *Armillaria*, along with other closely related species. Abbreviations: AA: Auxiliary Activities; CBM: Carbohydrate-Binding Modules; CE: Carbohydrate Esterase; GH: Glycoside Hydrolase; PL: Polysaccharide Lyase.

FIG 3 Differentially upregulated transcripts in the sporocarp and carpophoroid. (A) Venn diagram showing the number of differentially upregulated transcripts between the sporocarp and carpophoroid in *Armillaria* and *E. abortivum*; photo of *E. abortivum* taken by Eva Skific. (B) Bar graph showing the number of copies of genes important in mycoparasitic interactions. The white portion of the bar shows the number of these genes detected in the transcriptome but not differentially upregulated, while the darker colors (brown and purple, respectively) show the number of genes that are differentially upregulated by each species in the carpophoroid and the lighter colors (yellow and coral, respectively) show the number of these genes that are differentially upregulated by each species in their respective sporocarp.

FIG 4 Volcano plots. Each dot represents a transcript plotted according to its log₂ fold change and the -log₁₀ of its *p*-value. Transcripts with the highest statistical significance and the largest fold change will be represented by dots towards the top of the plot that are far to either the left- (carpophoroid) or right-hand (sporocarp) sides. All transcripts with a *q*-value < 0.05 are shown. Black dots represent a gene with a non-significant logFC (-2 < logFC < 2). (A) *Armillaria*. (B) *E. abortivum*. Dots with a black circle around them are annotated according to the abbreviations; OD: oxalate decarboxylase; CWG: cell wall galactomannoprotein.

FIG 5 Genes in both *Armillaria* and *E. abortivum* that are putatively important in the recognition and defense responses during this mycoparasitic interaction. (A) An *Armillaria* cell wall galactomannan that is differentially upregulated in the sporocarp; (B) three *E. abortivum* β-trefoil-type lectins that are differentially upregulated in the carpophoroid; (C) an *Armillaria* transcript that codes for isocitrate lyase that is differentially upregulated in its sporocarps; and (D) two *E. abortivum* transcripts that code for oxalate decarboxylases that are differentially upregulated in the carpophoroids.

1083 All results are shown as means $\pm$ standard error of the mean. TMM: normalized units of
1084 trimmed mean of m-values.

1085

1086

1087

1088

FIG 6 Boxplots of number of carpophoroid reads that mapped to *Armillaria* and *E. abortivum* when MAPQ = 30. Individual data points are indicated for each species with an open circle. The continuous line within each box represents the mean number of mapped reads. Species labeled with different letters (a to b) have a statistically significant ($p < 0.05$) different number of mapped reads in the carpophoroid.

FIG 7 Phylogenetic placement of the *Armillaria* species used in this analysis. (A) Maximum likelihood phylogeny of *Armillaria* species and *Guyanagaster necrorhizus* generated from the analysis of 100 random BUSCOs. The outgroup taxon is *Cylindrobasidium torrendii*. Each node is fully supported with 100% bootstrap supported. (B) Maximum likelihood phylogeny of *Armillaria mellea* phylogenetic tree generated from the analysis of the ITS region. Branches with 70% or more bootstrap support are thickened. Outgroup taxa include *A. gemina*, *A. luteobubalina*, *A. puiggarii* and *A. sinapina*. The *Armillaria* specimen analyzed during this study is in bold and yellow in both phylogenies.