APRILE: Exploring the Molecular Mechanisms of Drug Side Effects with Explainable Graph Neural Networks

Hao Xu¹, Shengqi Sang^{2,3}, Herbert Yao¹, Alexandra I. Herghelegiu⁴, Haiping Lu⁴, James T. Yurkovich⁵, and Laurence Yang^{1,*}

¹Department of Chemical Engineering, Queen's University, Ontario, Canada

²Department of Physics and Astronomy, University of Waterloo, Ontario, Canada

³Perimeter Institute for Theoretical Physics, Ontario, Canada

⁴Department of Computer Science, The University of Sheffield, UK

⁵Department of Bioengineering, University of California San Diego, La Jolla, CA

* Correspondence: laurence.yang@queensu.ca

Abstract

The majority of people over the age of 65 take two or more medications. While many individual drug side effects are known, polypharmacy side effects due to novel drug combinations poses great risk. Here, we present APRILE: an explainable artificial intelligence (XAI) framework that uses graph neural networks to explore the molecular mechanisms underlying polypharmacy side effects. Given a list of side effects and the pairs of drugs causing them, APRILE identifies a set of proteins (drug targets or non-targets) and associated Gene Ontology (GO) terms as mechanistic ‘explanations’ of associated side effects. Using APRILE, we generate such explanations for 843,318 (learned) and 93,966 (novel) side effect–drug pair events, spanning 861 side effects (472 diseases, 485 symptoms and 9 mental disorders) and 20 disease categories. We show that our two new metrics—pharmacogenomic information utilization and protein-protein interaction information utilization—provide quantitative estimates of mechanism complexity. Explanations were significantly consistent with state of the art disease-gene associations for 232/239 (97%) side effects. Further, APRILE generated new insights into molecular mechanisms of four diverse categories of adverse drug reactions: infection, metabolic diseases, gastrointestinal diseases, and mental disorders, including paradoxical side effects. We demonstrate the viability of discovering polypharmacy side effect mechanisms by training an XAI framework on massive biomedical data. Consequently, it facilitates wider and more reliable use of AI in healthcare.

Introduction

Polypharmacy poses a growing global health challenge. Polypharmacy is prevalent in the general population, as half of those taking any prescription medication are exposed to two or more drugs¹. For people 65 and older, this percentage increases to 75%². With 22% of the global population expected to be over 65 years old by 2050², polypharmacy is becoming increasingly pervasive.

Polypharmacy increases the risk of adverse drug reactions (ADRs). In Europe, adverse drug events are responsible for 8.6 million hospital admissions per year². In the UK, up to 11% of hospital admissions were due to adverse drug events—critically, over 70% of these admissions were in elderly patients, who are exposed to polypharmacy³. Thus, polypharmacy management is a core element of the global challenge for medication safety that was launched by the World Health Organization (WHO) in 2017².

Polypharmacy management is complex: over 200,000 approved drugs gives rise to over 200 million possible drug pairs, most of which will not be experimentally tested. Thus, Artificial Intelligence (AI)-assisted methods have been developed to predict potential polypharmacy risk^{4,5}. Specifically, such “predictors” predict the ADRs for all possible drug combinations.

Pharmacogenomic information plays an essential role for predicting and explaining the drug interactions and ADRs, and there are massive experimental data available^{6,7}. This information includes drug targets, functional associations or physical connections between or within drugs and proteins, all of which can be naturally depicted as a knowledge graph (KG). By extracting and encoding such information, graph neural network based deep learning methods are able to provide more reasonable ADR predictions based on the underlying protein interactions^{4,5}. However, the prediction accuracy of these approaches is not high enough for the model to be trusted completely and anomalous decisions can be made, causing undesired side effect in practical applications. Thus, human-interpretable explanation for making such predictions are still needed to provide data-driven solutions that help medical practitioners make better diagnoses to improve medical care⁸.

Explainable AI (XAI) seeks to enhance the model training, representation learning and decision-making processes with human-understandable explanations⁹. Recently, increasing numbers of XAI methods have been designed to improve the reliability of model predictions^{10,11}. The “explainer” methods have developed as a sub-domain of XAI that explain the predictions made by particular classes of machine/deep learning models, including even those that are non-interpretable^{12,13}, such as an explainer for all graph

neural networks¹⁴. We hypothesize that the explanations for ADR predictions based on the pharmacogenic KG can reveal new mechanistic associations between ADRs and drug targets/non-targets. Such associations make it possible to gain further pharmaceutical insights and provide reliable treatment options.

In this paper, we demonstrate that XAI can effectively reveal the mechanisms underlying polypharmacy side effects. We present APRILE (Adverse Polypharmacy Reaction Intelligent Learner and Explainer), an explainable framework in the format of predictor-explainer, that leverages pharmacogenomic information for discovering the molecular mechanisms of interactions between drugs. The workflow of APRILE is as follows: first it trains a machine learning model, APRILE-Pred, that is capable of predicting drug combination's side effects from pharmacogenomic data. Then, given a side effect prediction or a set of side effect predictions, APRILE trains another machine learning model, APRILE-Exp. This model produces a tentative molecular mechanism level explanation to those predictions—it queries APRILE-Pred to identify the most important pharmacogenomic information required to make the predictions.

Results

Here, we present APRILE, an XAI platform that provides novel drug-drug-ADR triplet predictions and mechanistic explanations for their underlying mechanisms of interaction. In the following sections, we justify the need for APRILE, detail the framework itself and provide validation for the underlying model. We then show that APRILE generates new insights into molecular mechanisms of four diverse categories of adverse drug reactions: infection, metabolic diseases, gastrointestinal diseases, and mental disorders, including paradoxical side effects of APRILE using existing data for drug interactions and mechanisms.

XAI is required to explain drug interactions

We first asked whether XAI is needed to explain why drug pairs cause a side effect based on the KG used to train APRILE. Our KG comprised 4.6 million drug-drug-ADR triplets, 18,596 drug-protein interactions and 1,431,224 protein-protein interactions (Methods). If most of the ADRs were caused simply by two drugs targeting the same protein, then AI may not be necessary as the side effect mechanism would be trivial. However, only 2.6% of drug-drug-ADR triplets had at least one shared protein target. Therefore, XAI, such as APRILE, is needed to explain the vast majority of these side effects.

APRILE: an explainable framework for discovering adverse polypharmacy reactions

The APRILE framework is shown in Figure 1 and described in the Methods section. It is composed of two modules, APRILE-Pred and APRILE-Exp, for predicting and explaining adverse drug reactions (ADRs), respectively.

APRILE-Pred is an instance of graph neural network (GNN), which is a popular machine learning model for extracting structural information and node features from graph-like data. GNN has proven to be successful at predicting polypharmacy ADR tasks^{4,5}. APRILE-Pred takes a pharmacogenomic KG as input, and predicts the polypharmacy side effects driven by all possible drug pairs. The training of APRILE-Pred is the process of tuning its parameters and learning node embeddings by capturing the information in the pharmacogenomic KG to predict ADRs.

APRILE-Exp is a variant of the GNNExplainer¹⁴ model, which is a general framework to explain GNNs. To our knowledge, our work is the first implementation of the GNN-explainer to the problem of explaining ADR predictions. However, we find that applying GNN-explainer directly results in explanations that are not biologically informative. Most of the biologically important interactions (protein-protein and protein-drug interactions) are not used for the explanation—instead, only drugs are used to explain the ADR. This is because the drug embeddings learned by the GNN dominate the model explanations. Therefore, we develop a new training strategy for GNNs to enhance explainability, which we call ‘lazy training’ (Methods). This strategy involves tuning parameterization to isolate relational information on the KG. We thus ensure that target and non-target proteins can be considered when explaining ADRs.

Moreover, APRILE-EXP suggests a tailored loss function to provide meaningful and flexible explanations, taking the multiple mechanisms by which polypharmacy can cause ADRs into account. It takes a well-trained APRILE-Pred and a set of drug-drug-ADR triplets as input. Its training amounts to looking for a subgraph of the original pharmacogenomic knowledge graph that simultaneously minimizes (i) the number of drug targets and non-targets in the subgraph, and (ii) the error of APRILE-Pred’s prediction(s), which are inputs (Equation (7) in Methods). The competition between minimizing target versus non-target proteins in objective (i) is controlled by the coefficients (α and β) of corresponding terms in the loss function, which can be manually tuned. Finally the optimal subgraph is outputted as a ‘molecular mechanism-level explanation’ of the side effects of interest.

To analyze APRILE’s ADR predictions and explanations, we design two metrics to describe the complex-

ity of ADR mechanisms: pharmacogenomic information utilization (PIU) and protein-protein interaction information utilization (PPIU) scores (Methods). These metrics quantify how important the information on drug targets or non-target proteins is respectively for APRILE-Pred's predictions, and guide the coefficients tuning in the APRILE-Exp's loss function. APRILE thus is able to suggest the identification of a given ADR's mechanism into the following two different classes of mechanisms: mechanisms involving 1) only the drug targets, and 2) target and non-target proteins.

Lazy training improves the explainability of APRILE

We found that lazy training improves the explainability of APRILE-Pred stably (Figure 2-A,C) without significantly reducing model prediction accuracy. First, we assessed how sensitive the model is to the initialization of these fixed matrices and the size of samples for training in detail (Methods). The results show that the standard deviation of the AUROC score on the corresponding testing set for different splitting rates is 0.038 ± 0.0004 (Figure 2-A, Table S1). Next, the ablation study on lazy training (Methods) also showed that there is a trade-off between model prediction accuracy and the PPIU score for each side effect (Figure 2-C). Sacrificing a little accuracy can significantly improve the potential interpretation of the model.

APRILE-Pred model tends to better predict rare side effects than common side effects (the marker size indicates the number of positive instances of the corresponding side effects in Figure 2-C). This observation is as expected because APRILE-Pred emphasizes learning from pharmacogenomic information and weakens learning from drug features and direct drug interactions.

Validation of APRILE

We built APRILE on a pharmacogenomic knowledge graph involving ~ 1.5 million relations among 284 drugs and ~ 19 thousand drug targets and non-targets. Then, it is trained and tested on ~ 4.6 million drug-drug interactions (DDIs) each associated with an ADR. In order to ensure that APRILE's performance on the seen and unseen DDIs is balanced, APRILE-Pred's model selection is based on cross validation. The selected APRILE-Pred model is trained with 90% positive samples, tested with the remaining 10%, and has the most balanced performance on the training and testing set (micro-averaged AUROC=0.85) (Figure 2-B).

APRILE learns meaningful side effect embeddings

To examine what APRILE-Pred learned, we interrogated the prediction model from side effect to category level and examined its learned embeddings. Based on the Medical Subject Headings (MeSH) database¹⁵, we classify the side effects into three main categories: *disease* (C), *mental disorder* (F) and *symptom* (S) (Supplementary Data). We found that a two-dimensional UMAP¹⁶ projection of side effect embeddings learned by the selected trained APRILE-Pred contained meaningful clusters for 20 disease subcategories and the other two categories (Figure 2-D). Clusters F (mental disorder), C07 (stomatognathic disease) and C13 (female urogenital diseases and pregnancy complications) were located at the extremities, indicating that their embeddings differed the most from other MeSH categories and from each other. Clusters close to each other are interpreted as having similar embeddings: e.g., C16 (congenital, hereditary, and neonatal disease and abnormalities) and C05 (musculoskeletal diseases). However, comparing the Euclidean distance and the Jaccard similarity of the side effects of these 22 side effect categories shows that the overlapping side effects among different categories does not dominate the clustering of these categories (Figure S2). Rather, the embeddings are based on extracting information from integrating the pharmacogenomic knowledge graph with drug adverse events.

APRILE generates valid explanations

When discovering the toxicity mechanisms of combination therapy, APRILE-Exp explains why APRILE-Pred decides that a pair of drugs caused a side effect. We perform a literature-based evaluation of the explanations' reliability. Literature evidence was found for additional explanations for ADRs in the training set (Table S2). Out of 843,318 predictions (Supplementary Data), we manually searched literature that supported explanations for the top five predictions ranked by prediction score (Methods). We found evidence for 2/3 genes for scotoma, one gene (*CD7*) for melanocytic naevus and none for elevated prostate specific antigen. Indirect evidence was found for bacterial pneumonia and post traumatic stress disorder.

When studying the general mechanism of a side effect, APRILE-Exp takes the set of all predictions on this side effect as input and gives the reasons why APRILE-Pred makes these predictions. For the example of pleural pain, the input to APRILE-Exp is $\{(\text{drug}_1, \text{drug}_2, \text{pleural pain}) \in \mathbf{A}^R\}$ (Methods), which includes all drug pairs causing pleural pain. We do this for every side effect (861 in total) and get a set of important genes, gene-gene interactions and enriched GO items (Figure 3-A, Supplementary Data) We found that 6 GO items: 'GO:0007186', 'GO:0007268', 'GO:0007204', 'GO:0007165', 'GO:0006954', 'GO:0007200' are associated with all side effects. Meanwhile, 250 GO items are associated with only

one side effect, while there are 159 side effects with unique GO items.

We then investigated the significance of the explanations by comparing them against disease-gene associations from Comparative Toxicogenomics Database (CTD)¹⁷, for 239 side effects that were found in CTD. We computed the precision and Jaccard index based on comparing the sets of genes in the APRILE-Exp explanation versus the disease-associated genes in CTD.

We then compared APRILE-Exp accuracy with 10,000 random explanations (i.e., gene sets) (Methods). We compared two types of explanations: all genes in a APRILE-Exp subgraph, or the subset of these genes that are in enriched GO terms. Additionally, two types of random models were tested: (A) one that preserves the number of drug target vs. non-target genes, and (B) one that samples from all genes in the KG. Thus, these random explanations capture protein interactions better than the naive GNN-explainer implementation.

Overall, APRILE-Exp explanations for 97% (232/239) of side effects were better than 10,000 random explanations ($P < 0.05$) (Supplementary Data). On average, the precision of APRILE-Exp explanations was 33% to 55% higher than random models, when comparing GO-enriched explanations versus drug target-agnostic random models (random model B).

For the 5 to 7 side effects where $P > 0.05$, the number of disease-associated genes in CTD was typically small with most < 60 , one case having 629 genes, and one case having 10,620 genes. For cases with very few CTD genes, we propose that APRILE-Exp explanations may contain new hypotheses for side effect-gene associations. In the last case (proctitis), the random models had a mean precision of 0.5. Thus, by randomly sampling genes from the KG, half are expected to be in the disease-associated gene list. This case suggests that proctitis may be highly complex, involving a multitude of genes, which makes a mechanistic explanation particularly challenging.

ADR classification using identified biological processes

Very few systematic methods are available for performing a global comparison of molecular mechanisms for a vast number of side effects. To address this challenge using APRILE, we constructed an ADR feature matrix based on the 1,787 biological process GO terms in APRILE's explanations for 861 side effects (Figure 3-A). We applied affinity propagation¹⁸ to cluster disease and mental disorder ADRs, based on the negative squared Euclidean distances. We then tuned the input preferences to minimize the number of clusters until all clusters contain more than one side effect. Finally, we obtained 25 side effect clusters

in total (Figure 3-B, Table S3).

We hypothesized that side effects clustered by biological process may also be related by MeSH disease categories. Enrichment analyses of the 25 clusters for 22 MeSH categories (see Methods) was used to rank order clusters by odds ratio, subject to an unadjusted enrichment $P < 0.05$ (Figure S1). The top 5 enriched clusters and their MeSH categories were panic disorder-mental disorders, embolism pulmonary-hemic and lymphatic diseases, cough-respiratory tract diseases, palpitation-cardiovascular disease, and incisional hernia-eye diseases. While all five clusters had unadjusted $P < 0.05$ and odds ratio > 4 , their FDR-adjusted P values all exceeded 0.05 (Supplementary Data). This result showed that side effects sharing similar mechanisms (i.e., biological processes) are not necessarily similar according to disease categorizations, in this case MeSH. We thus performed GO enrichment (see Methods) of each cluster and found 1 to 124 enriched GO biological processes (FDR-adjusted $P < 0.05$; median odds ratio > 4.34 (Supplementary Data). We further identified up to 64 GOs that were uniquely enriched for each cluster. Thus, side effects that are conventionally considered to be distinct disease categories may share many underlying biological processes. This view of side effect clusters may help in developing interventions against multiple, seemingly disparate side effects.

The bursitis cluster includes 20 side effects including breast cancer, sleep apnea, renal cyst, and gastroduodenal ulcers. These side effects span 10 diverse disease categories. Only one GO term was uniquely enriched: aminergic neurotransmitter loading into synaptic vesicle. In the context of breast cancer, drugs that dysregulate aminergic neurotransmitters affect immune function and cancer progression¹⁹. Clustering of the side effects sleep apnea and muscle disorder was also supported by literature: aminergic neurotransmitter factors or imbalances can cause sleep apnea²⁰, or masticatory muscle disorder²¹. Thus, non-intuitive, mechanistic similarities were captured by APRILE-Exp.

The panic disorder cluster includes 54 side effects spanning 19 disease categories, including mental disorders, nervous system diseases, and cardiovascular diseases (Supplementary Data). It is enriched for 11 unique GO terms, including GOs related to regulation of signal transduction, regulation of ion transmembrane transporter activity, tryptophan catabolism, and fatty acid metabolism and nucleoside salvage (Supplementary Data). Consistent with this clustering, tryptophan metabolism is disturbed in cardiovascular disease patients²² and also in panic disorder²³. In both cases, cytokine-based signal transduction alterations cause disrupted tryptophan metabolism.

Using PPIU scores to interpret APRILE results

We defined the protein-protein interaction information utilization (PPIU) metric to help interpret APRILE-Exp results (Methods). The PPIU score is a measure of the contribution of known protein-protein interactions to an APRILE prediction. A low score indicates that the target proteins are sufficient for explaining a side effect, while a high score indicates that additional proteins in the network, including those connected to the direct drug target, are needed to fully explain the side effect. Thus, we explored the interpretation of PPIU scores and how they can be used to better understand APRILE results.

Selecting sepsis as an example side effect, we examined the high-probability (> 0.95) APRILE predictions for which alternative explanations had low or high PPIU scores. Of the 66 predicted explanations with low PPIU (< 0.1), 64 (97%) were direct drug targets (Supplementary Data). Alterations in one such direct target, transferrin, has been reported in sepsis patients²⁴, suggesting that a perturbation to transferrin through some drug mechanism may increase the chances of experiencing sepsis as a side effect. Predicted explanations with high PPIU (> 0.9) included six target genes and one common non-target gene. All seven genes were related to choline transport and anti-inflammatory mechanisms, a neuroimmune pathway that may play a role in the development of sepsis.

APRILE predicts drug combinations with altered individual drug effects

We next explored predicted drug-drug-ADR triplets that were not in the training set, meaning they may represent yet undocumented ADRs. One such novel, high-confidence (prediction score > 0.99) example is that the combination of omeprazole and gabapentin were predicted to result in pleural pain. Omeprazole is an H⁺/K⁺ ATPase-binding proton-pump inhibitor typically used to treat gastric acid-related disorders, and gabapentin is a voltage-gated calcium channel (VGCC) blocker typically prescribed as an anticonvulsant. For this pair, APRILE predicted significant enrichment for GO terms associated with VGCCs (cellular component ‘VGCC complex’ and molecular function ‘VGCC activity’).

The inhibition of H⁺/K⁺ ATPase activity—such as by drugs like omeprazole—disrupts the sodium-calcium balance, leading to an accumulation of intracellular calcium²⁵. While VGCC blockers such as gabapentin are typically used to reduce pain by inhibiting the transmission of pain through calcium gradients, VGCC blockers have also been used to reduce intracellular calcium accumulation²⁶. Furthermore, calcium channel blockers have been thought to cause pleural effusion, which can cause pleural pain²⁷. Thus, it is possible that the combination of these two drugs results in not only differing mechanisms of the individual

drugs, but also pleural pain. While such a prediction warrants further investigation, it nevertheless suggests that APRILE has the potential to identify common drugs that result in unintended side effects when used in combination.

Predicting explanations for paradoxical side effects

Some drugs have side effects reported in some conditions but not others, or, in more severe cases, one side effect in some cases and the opposite side effect in other cases. Cases where the drug effect is opposite to the expected outcome is referred to as ‘paradoxical adverse reactions’²⁸. Such paradoxical side effects offer obvious challenges for patient care. Here, we explore the capacity of APRILE to predict mechanistic explanations for observed paradoxical side effects through two case studies.

Nicotine activates *CHRNA7*, the human alpha 7 nicotinic acetylcholine receptor. It has been shown to both cause and reduce anxiety, while also having been observed to both desensitize and activate nicotinic acetylcholine receptors (nAChRs)²⁹. Ondansetron is an antiemetic which is predicted to bind to *CHRFAM7A*³⁰, a chimeric gene partially duplicated from *CHRNA7*. APRILE correctly predicted that the combination of nicotine and ondansetron would result in anxiety. The predicted explanation included two genes that are not direct targets of either drug: *Lynx-1* and *SLURP-1*, which are both endogenous allosteric modulators of nAChRs. *SLURP-1* inhibits alpha 7 nAChR activity (ion influx through the receptor channel)³¹, while *Lynx1* inhibits nAChRs through direct binding and reducing their sensitivity to acetylcholine.

Because both drugs activate *CHRNA7*, we hypothesize that polypharmacy of nicotine and ondansetron induces nAChR hypersensitivity, resulting in increased anxiety. This hypothesis has been corroborated by animal studies, wherein hypersensitive nAChR has been shown to increase anxiety in mice³². A simple description of the mechanism is that multiple drug-protein and protein-protein activities combine in a new context, resulting in the polypharmacy side effect. Here, we propose that hypersensitivity to polypharmacy arises as the respective drug target proteins also inhibit a common protein that is not directly targeted by either drug (Figure 4b). We explored other predicted cases of paradoxical side effects, such as how the use of a combination of anti-epileptic drugs may present side effects similar to that of an excessive dosage of a single drug (Figure S3)

Suggesting mechanisms for common polypharmacy side effects

Finally, we performed a global analysis to investigate polypharmacy side effects that commonly result from pairs of drugs. Specifically, we examined three common peptic ulcers—duodenal, gastric, and esophageal—that are documented side effects for 69 drug pairs. APRILE was used to predict mechanistic explanations these three side effects have in common, resulting in seven GO terms (Table S4). *Helicobacter pylori* infection is a common cause of ulcers³³ and is thought to cause ulcers in nearly 20% of human hosts³³. This process led us to hypothesize two key mechanisms related to *H. pylori* infection in the gut microbiome that duodenal, gastric, and esophageal ulcers share.

First, APRILE predicted heme binding (GO:0020037) and lowered iron binding (GO:0005506) to be common mechanisms in these peptic ulcers. *H. pylori* virulence has previously been linked to both heme-binding proteins³⁴ and reduced iron-binding capacity³⁵. Second, APRILE predicted that several metabolic processes—arachidonic acid metabolism, geranyl diphosphate biosynthesis, and phospholipase A2 activity (GO:0004623)—contribute to the cause of these ulcers. These processes are known to play a role in breaking or maintaining the mucosal barrier, which helps defend against infection³⁶. Thus, while the virulence of the pathogen may contribute, APRILE identified important host factors that increase susceptibility to infection—ultimately increasing the risk for peptic ulcers.

Discussion

The ability to predict and mechanistically understand drug side effects resulting from polypharmacy poses a significant challenge to modern medicine. Here, we employ a novel explainable artificial intelligence (XAI) platform, APRILE, to provide meaningful, accurate, mechanistic explanations for polypharmacy side effects. APRILE computes quantitative metrics that indicate whether proteins beside drug targets are likely to be involved in an ADR mechanism. We validated APRILE predictions, and offered several mechanistic hypotheses for novel drug-drug-ADR triplets. Taken together, these results have several important implications.

First, our data-driven XAI approach has the potential to identify unreported drug pairs that pose risk to patients and further predict mechanistic explanations to support clinical decision making. There are over 20,000 approved prescription drugs giving rise to over 200 million possible drug pairs. Clinical decision support softwares such as that by uMETHOD Health³⁷ create personalized treatment plans that provide physicians with alerts and recommendations on drug interactions. A desired feature of such software is

314 proposing mechanisms that underlie the drug interactions³⁸, while also accounting for a patient-specific
315 conditions such as other chronic illnesses (see Supplementary Note 1). Thus, APRILE may be integrated
316 into clinical decision support software to enhance their effectiveness and adoption by clinicians.

317 Second, lazy training encourages APRILE’s explainability, and APRILE-Exp’s loss design provides mean-
318 ingful and flexible explanations. Lazy training enables GCNs to maintain the independence of relational
319 information, such that it encourages APRILE-Exp to pay more attention to the non-target proteins that
320 are important for APRILE-Pred to make predictions. We also present the strategy for designing alterna-
321 tive loss functions for the explainer. Users of APRILE can tailor explanations involving drug targets and
322 non-targets by exploring the loss function design space. Therefore, these ideas can help adapt the GNN
323 and GNN-based explainer¹⁴ to different application scenarios.

324 Third, an XAI platform that predicts polypharmacy ADRs can help in designing clinical trials by acting
325 as an *in silico* screening step. Robust clinical trials requires selecting a study population while balanc-
326 ing the trial’s capacity to validate the intervention and its generalization to the general population that
327 may deviate from ideal conditions³⁹. In particular, comorbidities are common for patients in real world
328 settings, and “pragmatic” clinical trials are increasingly used at or after phase 3 trials^{40–42}. Thus, while
329 designing pragmatic clinical trials, APRILE can predict drug-drug interactions that can arise for partici-
330 pants on existing medications and also provide explanatory insights as to why they occur. Overall, this
331 XAI platform for model-driven discovery of novel drug interaction mechanisms could accelerate existing
332 drug-discovery pipelines. Especially for drugs intended to be used in combination therapies (e.g., cancer
333 cocktails) or targeting populations having high polypharmacy (e.g., aging populations).

334 Using APRILE, we have reported both predicted polypharmacy drug-drug-ADR triplets and predicted
335 mechanistic explanations underlying them. While some of the predictions reported here were corrobo-
336 rated by literature evidence, many require follow-up experiments by the biomedical community before
337 being able to evaluate their clinical utility. In fact, APRILE has the capacity to generate model-driven hy-
338 potheses for mechanisms underlying 34 million drug-drug-ADR combinations using existing data—this
339 hypothesis space will grow exponentially as the knowledge graph is expanded with new data. Validating
340 these hypotheses may uncover new drug interaction mechanisms that have translational importance and
341 clinical value.

342 Here, we present a first step toward AI-guided polypharmacy side effect explanation. Using massive data

that already exists, APRILE has generated numerous model-driven hypotheses for ADR mechanisms. APRILE’s high explainability makes it useful when designing safe polypharmacy interventions, whose side effects are too complex to rely on human analysis alone. XAI such as APRILE will play an increasingly critical role for polypharmacy management in an aging population for which polypharmacy is currently the norm. Further, we believe APRILE represents an important foundation for future AI-based platforms that will aid medical practitioners in clinical diagnoses and decision making.

Methods

Pharmacogenomic knowledge graph construction

We use two subdatasets for polypharmacy-associated ADR modelling from the BioSNAP-Decagon dataset⁷ to prepare the pharmacogenomic knowledge graph : **PP (protein-protein)**: relations between proteins depend on whether they have functional association or physical connections. **GhG (drug-protein)**: relations between drugs and their targeted proteins. All drugs were referenced by CID, which is linked to the chemical structure; this annotation ensures uniqueness within the database.

In summary, the pharmacogenomic knowledge graph contains 19,365 vertices (284 drugs (D) and 19,081 proteins (P)), 1,449,820 edges. Among the edges, there are 1,431,224 protein interaction edges (P-P) and 18,596 drug protein targeting edges (P-D). We use the following data structures to represent the graph we constructed.

- $\mathbf{A}^P \in \mathbb{R}^{|P| \times |P|}$: symmetric adjacency matrix for the undirected subgraph within proteins.
- $\mathbf{A}^t \in \mathbb{R}^{|P| \times |D|}$: adjacency matrix for the directed bipartite subgraph between the proteins and drugs, whose edges are directed from proteins to drugs.

APRILE-Pred: explainable adverse polypharmacy reaction prediction with GNNs

APRILE-Pred is a Graph Neural Network (GNN) model that takes protein-protein and protein-drug interaction graphs as input and predicts the side-effects caused by drug combinations. Specifically, we first learn the protein representation in the protein interaction subgraph, and then generate the drug representation through the protein-drug interaction subgraph with drug attributes (See Algorithm 1). In addition, The input drug/protein node features/attributes $\{\mathbf{v}_{d/p} | d \in D, p \in P\}$ are set up to one-hot vectors, which are binary vectors representing the i th drug/protein if its i th element is 1. Protein and drugs are indexed respectively.

Learning Drug Representation.

Firstly, we use a 2-layer graph convolutional network (GCN)⁴³ to capture protein attributes and relations on the protein interaction subgraph $\mathbf{A}^P$. The input is the protein node features $\mathbf{h}_p^0 = \mathbf{v}_p, \forall p \in P$. For each protein node $p \in P$, the relation between two hidden layers is given by

$$\mathbf{h}_p^{k+1} = \text{ReLU} \left(\sum_{p' \in \mathcal{N}_p} \frac{1}{C_{p,p'}} \mathbf{W}_p^k \mathbf{h}_{p'}^k \right), \quad (1)$$

where $\mathcal{N}_p$ denotes the union of protein node p and its neighbors on the subgraph $\mathbf{A}^P$, $C_{p,p'} = |\mathcal{N}_p| |\mathcal{N}_{p'}|$ is coefficient calculated by the product of the degree of p and $p' \in \mathcal{N}_p$, and $\mathbf{W}_p$ are parameters. The final learned protein embeddings/representation $\mathbf{z}_p = \mathbf{h}_p^1 \oplus \mathbf{h}_p^2, \forall p \in P$ are the concatenation of the outputs of all the GCN layers.

Then, we use a graph-to-graph information propagation module⁵ to transform learned protein embeddings and drug node features into feature representation of drugs via the bipartite protein-drug integration subgraph $\mathbf{A}^I$. The protein's contribution to the drug d 's representation for all $d \in D$ is:

$$\mathbf{c}_d = \text{ReLU} \left(\frac{1}{C_d} \sum_{p' \in \mathcal{N}_d} \mathbf{W}_T \mathbf{z}_{p'} \right), \quad (2)$$

where $\mathcal{N}_d = \{p' | p' \in P, \mathbf{A}_{p',d} = 1\}$ is the set of drug d 's protein neighbours, $C_d = |\mathcal{N}_d|$, and $\mathbf{W}_T$ is a trainable parameter matrix. Besides, A linear transformation followed by an activation function is used to map the original drug node features $\mathbf{v}_d$ into the same space that protein's contributions $\mathbf{c}_d$ live in for each drug $d \in D$:

$$\mathbf{f}_d = \text{ReLU}(\mathbf{W}_F \mathbf{v}_d). \quad (3)$$

The final representation of drug $d \in D$ is: $\mathbf{z}_d = \mathbf{z}_p \oplus \mathbf{c}_d \oplus \mathbf{f}_d$, where $\oplus$ denotes concatenation.

Polypharmacy Side Effect Prediction.

We predict if drug combinations can cause any side effect. Here, DistMult factorization⁴⁴ is used as the scoring function, because it is a well-known model that models rich relationships well on standard benchmarks alone or as a decoder. The probability that drug pairs $(d, d'), \forall d, d' \in D$ cause side effects $r, \forall r \in R$ is obtained by acting the sigmoid function on the score tensor $\mathbf{G} = \{g_{d,d'}^r\} \in \mathbb{R}^{|D| \times |D| \times |R|}$ computed by the scoring function:

$$p_{d,d'}^r = \sigma(g_{d,d'}^r) = \sigma(\mathbf{z}_d^\top \mathbf{M}_r \mathbf{z}_{d'}), \quad (4)$$

Algorithm 1: APRILE-Pred’s Forward Propagation

Input: $\mathbf{V}_p, \mathbf{V}_d, \mathbf{A}^p, \mathbf{A}^t$

Parameter: $\mathbf{W}_P, \mathbf{W}_T, \mathbf{W}_F$

Output: $\mathbf{P}$

$\mathbf{H}^0 \leftarrow \mathbf{V}_p$

for $l \in L_p$ **do**

$\mathbf{H}^{l+1} \leftarrow \text{ReLU}(\text{GCN}_{pp}^l(\mathbf{H}^l, \mathbf{A}^p; \mathbf{W}_P^l))$

end

$\mathbf{Z}_p \leftarrow [\mathbf{H}^0, \dots, \mathbf{H}^L]$

$\mathbf{C} \leftarrow \text{ReLU}(\text{GCN}_{pd}(\mathbf{H}, \mathbf{A}^t; \mathbf{W}_T))$

$\mathbf{F} \leftarrow \text{ReLU}(\mathbf{V}_d \mathbf{W}_F^T)$

$\mathbf{Z}_d \leftarrow [\mathbf{Z}_p, \mathbf{C}, \mathbf{F}]$

$\mathbf{P} \leftarrow \text{Sigmoid}(\mathbf{Z}_d^T \mathbf{M} \mathbf{Z}_d)$

Algorithm 2: APRILE-Exp’s Forward Propagation

Input: $S = \{(d_n, d'_n, r_n)\}_n, \mathbf{V}_p, \mathbf{V}_d, \mathbf{W}_P, \mathbf{W}_T, \mathbf{W}_F$

$\mathbf{W}_F$

Parameter: $\hat{\mathbf{A}}^p, \hat{\mathbf{A}}^t$

Output: $\mathbf{P}_S$

$\mathbf{H}^0 \leftarrow \mathbf{V}_p$

for $l \in L_p$ **do**

$\mathbf{H}^{l+1} \leftarrow \text{ReLU}(\text{GCN}_{pp}^l(\mathbf{H}^l, \hat{\mathbf{A}}^p; \mathbf{W}_P^l))$

end

$\mathbf{Z}_p \leftarrow [\mathbf{H}^0, \dots, \mathbf{H}^L]$

$\mathbf{C} \leftarrow \text{ReLU}(\text{GCN}_{pd}(\mathbf{H}, \hat{\mathbf{A}}^t; \mathbf{W}_T))$

$\mathbf{F} \leftarrow \text{ReLU}(\mathbf{V}_d \mathbf{W}_F^T)$

$\mathbf{Z}_d \leftarrow [\mathbf{Z}_p, \mathbf{C}, \mathbf{F}]$

$\mathbf{P} \leftarrow \text{Sigmoid}(\mathbf{Z}_d^T \mathbf{M} \mathbf{Z}_d)$

$\mathbf{P}_S = \{P_{d_n, d'_n, r_n} | (d_n, d'_n, r_n) \in S\}$

where $\mathbf{M}_r$ is a trainable diagonal matrix associated with the side effect $r \in R$. We call it a single-instance prediction or a single prediction. In addition to calculating the probability that the drug pair (d, d') causes the side effect r , the proportion of variant protein related information which leads to the model to make such a prediction is also evaluated by pharmacogenomic information utilization (PIU) and protein-protein interaction information utilization (PPIU) scores:

$$\text{PIU}_{d, d'}^r = 1 - \frac{p_{d, d'}^r(\mathbf{z}_d^{\text{PIU}})}{p_{d, d'}^r(\mathbf{z}_d)}, \quad (5)$$

$$\text{PPIU}_{d, d'}^r = 1 - \frac{p_{d, d'}^r(\mathbf{z}_d^{\text{PPIU}})}{p_{d, d'}^r(\mathbf{z}_d)}, \quad (6)$$

where $\mathbf{z}_d^{\text{PIU}} = \mathbf{z}_p \oplus (\mathbf{c}_d \cdot \mathbf{0}) \oplus \mathbf{f}_d$ and $\mathbf{z}_d^{\text{PPIU}} = (\mathbf{z}_p \cdot \mathbf{0}) \oplus \mathbf{c}_d \oplus \mathbf{f}_d$.

APRILE-Pred Training Details

APRILE-Pred’s forward propagation is shown in Algorithm 1. We use the Adam optimizer with a training rate of 0.01 to minimize the cross-entropy loss. We applied full-batched end-to-end training for 80 epochs

on a single Tesla V100 (32 GB RAM) graphics processing unit (GPU).

Dataset. We use the database of polypharmacy side effects compiled by the BioSNAP-Decagon dataset^{4,7}. This database consists of 964 polypharmacy side effects from 4,651,131 drug-drug interactions, which is a subset of the TWOSIDES database⁴⁵. These polypharmacy side effects already exclude side effects attributed to an individual drug. In this work, we focus on the drugs that have at least one target protein, and the side effects that each occurred in at least 250 drug combinations. Therefore, there are 4,625,608 drug interaction edges (D-D) labeled by 861 side effects. We used as the positive polypharmacy adverse events to train and test the model.

Dataset split. The training and testing sets were stratified split based on side effects. To train and test the model, we need both positive and negative ADR knowledge instances in the form of (drug₁, drug₂, side effect). Positive instances are the observed polypharmacy side effect information, while negative ones are sampled with categorized negative sampling strategy⁴⁶.

Evaluation metrics. When comparing overall performance among trained models, we use the {Micro, Macro}-averaged receiver operator characteristic area under the curve (AUROC) and the area under the precision-recall curve (AUPRC) score for all side effects. The PIU and PPIU score of each side effect are computed based on the Macro-AUROC with the same $\mathbf{z}_d^{PIU}$ and $\mathbf{z}_d^{PPIU}$ for those of each prediction.

Lazy training. GNNs recursively incorporate information through the information propagation paths in the graph, leading to the node intermediary embeddings capturing both graph structure and node features. However, these rich embeddings learned by APRILE-Pred prevent APRILE-Exp from finding important information sources on drug targets and non-target proteins, as the pharmacogenomic information and ADRs have been learned into drug embeddings already during the back-propagation. This makes APRILE-Pred's ADR predictions difficult to be explained by APRILE-Exp. Therefore, to build a more interpretable model and reduce explanation redundancy for the side effects that have molecular origins, information on drug targets, protein relations and ADRs should be prevented from being learned in the trainable embedding matrices. Accordingly, we weaken the contribution of protein and drug attributes to APRILE prediction, and control the trainability of the model parameters. This strategy is referred to as 'lazy training'. Specifically, after initializing all the model parameters by Xavier initialization⁴⁷, we fix the parameters matrices $\mathbf{W}_p^1$ (in Equation 1) and $\mathbf{W}_F$ (in Equation 3), which are supposed to be trainable parameters, to their initial values. The fixed parameters are referred to as lazy learners.

APRILE-Exp: optimal explanations generation with GNNs explainer

We develop APRILE-Exp to explore the molecular mechanisms of side effects. The APRILE-Exp explains the existence of known knowledge by understanding APRILE-Pred model prediction. It is primarily motivated as an adaption of previous work on GNNExplainer¹⁴ for large-scale and tri-graph-like data. Given a trained APRILE-Pred model and a set of predictions we want to explain $S = \{(d_n, d'_n, r_n)\}_n$, the APRILE-Exp model generates a subgraph of the original graph (the dataset) that are most influential for the APRILE-Pred model's prediction. Therefore, the APRILE-Exp model transforms the problem of explaining why drug pairs cause side effects to finding out which protein interactions and drug interactions are the most important to predict side effects, which in turn helps us to understand the molecular mechanism of side effects.

We assigned an importance score to each edge in the P-P and P-D subgraphs by replacing the adjacency matrices $\mathbf{A}^P$ and $\mathbf{A}^T$ to $\hat{\mathbf{A}}^P$ and $\hat{\mathbf{A}}^T$, where $\hat{\mathbf{A}}^P$ and $\hat{\mathbf{A}}^T$ have the same non-zero elements as $\mathbf{A}^P$ and $\mathbf{A}^T$ but with tunable values $\hat{A}_{ij}^P = \hat{A}_{ji}^P \in [0, 1]$ and $\hat{A}_{ij}^T \in [0, 1]$. As shown in Algorithm 2, the APRILE-Exp model calculates scores of drug pairs for each side effect with the same pipeline as the prediction model but with trainable $\mathbf{X}\hat{\mathbf{A}}^P$ and $\hat{\mathbf{A}}^T$, fixing the trained parameters $\mathbf{W}_P$, $\mathbf{W}_T$, and $\mathbf{W}_F$. We minimize the following loss during training:

$$\begin{aligned} Loss = & \sum_n \log(1 - P(d_n, d'_n, r_n)) \\ & + \alpha (\sum_{i,j} \hat{A}_{i,j}^P + \sum_{i,j} \hat{A}_{i,j}^T) \\ & + \beta (\sum_{i,j} \hat{A}_{i,j}^P (1 - \hat{A}_{i,j}^P) + \sum_{i,j} \hat{A}_{i,j}^T (1 - \hat{A}_{i,j}^T)), \end{aligned} \quad (7)$$

where α and β are hyper-parameters. For the loss function, the first term encourages P to take a value close to 1, which can be replaced with any functions that decrease sharply near 1; the second term encourages the size of subgraph given by $\hat{A}_{i,j}$ to be as small as possible, and it can be replaced by any monotonically increase function; and the last term $= \hat{A}_{i,j}$ to be either 0 or 1, which can be replaced with any function that is 0 at points $x = 0$ and $x = 1$.

Gene Ontology Enrichment

To investigate the functional roles of the genes that are most influential for the interested predictions, we perform Gene Ontology (GO) enrichment analysis by using the GOATOOLS package⁴⁸. The Fisher's exact test is used to compute the p-values for each of: biological process, molecular function and cellular

component, which evaluates significant enrichment of certain GO terms. We choose the non-negative Benjamini-Hochberg multiple test correction method to correct the test, which controls the false discovery rate (FDR), and use a FDR set the significance cut-off to 0.05⁴⁹. We applied this methodology also to analyze enrichment of MeSH categories in clusters of side effects.

Experiments

Lazy training sensitive study

We assessed how sensitive the model is to the initialization of these fixed matrices and the size of samples for training in detail. To this end, we trained the APRILE-Pred models with five different parameter initialization settings and six different dataset splitting rates (the proportion of positive training samples to the whole positive samples for each side effect): 0.1, 0.2, 0.4, 0.6, 0.8 and 0.9.

Lazy training ablation study

We trained three APRILE-Pred model 90% positive samples: with lazy training, with lazy training only on W_F (in Equation 3), and without lazy training. Model performance is tested with the remaining 10% samples and the same numbers of random negative samples.

Evaluation on explaining single predictions

We perform a literature-based evaluation of explanation for both learned ADR knowledge and novel predictions based on the selected trained APRILE-Pred model. To this aim, we ask APRILE-Pred to make a prediction for every drug pair and every side effect in the knowledge graph. We then divided these predictions into two groups – learned knowledge and new prediction – according to whether predictions are in the training set or not, and constructed the corresponding ranked lists based on predicted probability for each prediction group. Then, we use APRILE-Exp to obtain a subset of drug targets, protein interactions and significant GO items for each of the twenty highest ranked predictions in both rank lists, and afterwards search biomedical literature to see if we can find supporting evidence.

Permutation test

To assess the significance of APRILE-Exp explanations, we compared them against 10,000 random explanations (i.e., gene sets). For APRILE-Exp and random explanations, we computed the precision and Jaccard index using the CTD¹⁷ disease-associated genes as validation. We compared two types of explanations: all genes in a APRILE-Exp subgraph, or the subset of these genes that are in enriched GO terms. Additionally, two types of random models were tested: (A) one that preserves the number of drug target

vs. non-target genes, and (B) one that samples from all genes in the KG. Permutation test p-values were determined as the number of random explanations whose precision (or Jaccard index) was equal to or greater than that of the APRILE-Exp explanation.

Acknowledgements

This work was supported by Queen's University, and the Natural Sciences and Engineering Research Council of Canada (NSERC) [RGPIN-2020-06325]. This research was enabled in part by support provided by Compute Ontario (www.computeontario.ca) and Compute Canada (www.computecanada.ca). Computations were performed on resources and with support provided by the Centre for Advanced Computing (CAC) at Queen's University in Kingston, Ontario. The CAC is funded by: the Canada Foundation for Innovation, the Government of Ontario, and Queen's University.

Author contributions

LY and HX conceived the study. HX and SS developed the framework. HX, LY, HY and AIH carried out the simulations and biologically interpreted results. All authors wrote and edited the manuscript.

Competing interests

The authors declare that they have no competing financial interests.

Data availability

The data for constructing pharmacogenomic knowledge graph and drug-drug interactions is available in the BioSNAP⁷ database (<http://snap.stanford.edu/biodata/>). The medical subject headings for drugs are available at <https://meshb.nlm.nih.gov>. The CTD¹⁷ disease-gene associations are available in <http://ctdbase.org/downloads/>.

Code availability

Code to train, test and select an APRILE-Pred model is available at <https://github.com/QCSB/APRILE-Pred>. Code to build and use APRILE-Exp is available at <https://github.com/QCSB/APRILE-Exp>. The published APRILE python package is available at <https://github.com/QCSB/APRILE>

Figure legends

Figure 1 - Overview of the APRILE framework: a workflow to predict and explain ADRs. **A-** the pharmacogenomic knowledge graph which contains drug-gene and gene-gene relations. **B-** protein and drug representation modules apply graph convolutional networks to extract pharmacogenomic information from **A**. **C-**, APRILE prediction module uses drug representations to predict the probability of that all drug pairs cause different side effects. The modules in **B** and **C** constitute the predictor of APRILE, APRILE-Pred, and are trained end-to-end with polypharmacy adverse event data. **D-** APRILE-Exp gives a small subgraph of **A** as an explanation for given prediction(s) e.g. polypharmacy adverse event(s), which are the most important drug targets and protein associations in making the given prediction(s), accompanied by functional roles of these proteins. PPIU and PIU score are also provided to measure the utilization of information on drug targets and protein interaction by APRILE-Pred. **E-** lazy training fixes the parameters $\mathbf{f}_p$ and $\mathbf{f}_d$ when they were initialized to prevent the relational information from being learned into the embeddings of protein, drug and side effect. **F-** gene ontology (GO) enrichment analysis is used to investigate the functional roles of the genes in APRILE-Exp's explanations.

Figure 2 - Evaluation metrics and visualization for APRILE-Pred. **A-** Sensitive evaluation for lazy training to different dataset split rates and initial parameter values. **B-** performance comparison of the selected trained APRILE-Pred model on the training and testing sets. **C-** ablation study on lazy training, which compares the distribution of AUROC for each side effect when applying no, partial and full lazy training. Marker size indicates the number of drug pairs that cause the corresponding side effect. **D-** UMAP of the category-average side effect embeddings learned by APRILE-Pred. Marker sizes are the log-scaled numbers of side effects in each categories.

Figure 3 - Side effect clustering based on GO items identified by APRILE-Exp. **A-** Association matrix for 861 side effects and 1,787 biological process GO items. C, S, F are side effect categories are the same as those in Fig 2d. **B-** side effect clustering based on the APRILE-Exp's explanation. The name of the example for each cluster (25 in total) are represented in large fonts. Each small cell is a side effect, and its colour indicates the number of GO items associated with it. (See Table S3 for details). **C-** clustering side effects in the infection disease category based on associated GO items. The side effects in the same cluster use the same marker symbol.

Figure 4 - Mechanisms of ADRs in polypharmacy. **A-** polypharmacy alters drug effects resulting in

molecular imbalance. **B-** drug effects change with context, such as drug targets inhibiting a common non-target protein. **C-** paradoxical ADRs may be caused by (i) inverse response within feedback regulatory processes, or by increased (ii) vulnerability to disruption of a biological process due to the action of a second drug.

Supplementary information

The supplementary information includes four figures and three tables. The supplementary data is available at <https://github.com/QCSB/APRILE-Exp/tree/master/supplementary-data>.

References

1. Quinn, K. J. & Shah, N. H. A dataset quantifying polypharmacy in the United States. *Scientific data* **4**, 1–11 (2017).
2. Mair, A. *et al.* Polypharmacy management by 2030: a patient safety challenge. (2017).
3. Kongkaew, C. *et al.* Risk factors for hospital admissions associated with adverse drug events. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* **33**, 827–837 (2013).
4. Zitnik, M., Agrawal, M. & Leskovec, J. Modeling polypharmacy side effects with graph convolutional networks. *Bioinformatics* **34**, i457–i466 (2018).
5. Xu, H., Sang, S. & Lu, H. Tri-graph Information Propagation for Polypharmacy Side Effect Prediction. *NeurIPS Workshop on Graph Representation Learning* (2020).
6. Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic acids research* **46**, D1074–D1082 (2018).
7. Marinka Zitnik Rok Sasic, S. M. & Leskovec, J. *BioSNAP Datasets: Stanford Biomedical Network Dataset Collection* <http://snap.stanford.edu/biodata>. Aug. 2018.
8. Stiglic, G. *et al.* Interpretability of machine learning-based prediction models in healthcare. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery* **10**, e1379 (2020).
9. Samek, W., Montavon, G., Lapuschkin, S., Anders, C. J. & Müller, K.-R. Explaining deep neural networks and beyond: A review of methods and applications. *Proceedings of the IEEE* **109**, 247–278 (2021).

- 543 10. Hastie, T., Tibshirani, R. & Wainwright, M. *Statistical learning with sparsity: the lasso and gener-*
544 *alizations* (Chapman and Hall/CRC, 2019).
- 545 11. Roscher, R., Bohn, B., Duarte, M. F. & Garcke, J. Explainable machine learning for scientific in-
546 sights and discoveries. *IEEE Access* **8**, 42200–42216 (2020).
- 547 12. Hinton, G., Vinyals, O. & Dean, J. Distilling the knowledge in a neural network. *NIPS Deep Learn-*
548 *ing and Representation Learning Workshop* (2015).
- 549 13. Du, M., Liu, N. & Hu, X. Techniques for interpretable machine learning. *Communications of the*
550 *ACM* **63**, 68–77 (2019).
- 551 14. Ying, Z., Bourgeois, D., You, J., Zitnik, M. & Leskovec, J. *Gnnexplainer: Generating explanations*
552 *for graph neural networks* in *Advances in neural information processing systems* (2019), 9244–
553 9255.
- 554 15. Lipscomb, C. E. Medical subject headings (MeSH). *Bulletin of the Medical Library Association* **88**,
555 265 (2000).
- 556 16. McInnes, L., Healy, J., Saul, N. & Grossberger, L. UMAP: Uniform Manifold Approximation and
557 Projection. *The Journal of Open Source Software* **3**, 861 (2018).
- 558 17. Davis, A. P. *et al.* Comparative Toxicogenomics Database (CTD): update 2021. *Nucleic acids re-*
559 *search* **49**, D1138–D1143 (2021).
- 560 18. Frey, B. J. & Dueck, D. Clustering by passing messages between data points. *Science* **315**, 972–977
561 (2007).
- 562 19. Fleischer, L. M., Somaiya, R. D. & Miller, G. M. Review and meta-analyses of TAAR1 expression
563 in the immune system and cancers. *Frontiers in pharmacology* **9**, 683 (2018).
- 564 20. Dempsey, J. A., Veasey, S. C., Morgan, B. J. & O'Donnell, C. P. Pathophysiology of sleep apnea.
565 *Physiological reviews* (2010).
- 566 21. Alóe, F. Sleep bruxism neurobiology. *Sleep science* **2**, 40–48 (2009).
- 567 22. Mangge, H. *et al.* Disturbed tryptophan metabolism in cardiovascular disease. *Current medicinal*
568 *chemistry* **21**, 1931–1937 (2014).
- 569 23. Quagliato, L. A. & Nardi, A. E. Cytokine alterations in panic disorder: A systematic review. *Journal*
570 *of affective disorders* **228**, 91–96 (2018).

24. Piagnerelli, M. *et al.* Rapid alterations in transferrin sialylation during sepsis. *Shock* **24**, 48–52 (2005).
25. Lees, G. Inhibition of sodium-potassium-ATPase: a potentially ubiquitous mechanism contributing to central nervous system neuropathology. *Brain Research Reviews* **16**, 283–300 (1991).
26. Gurkoff, G. G., Shahlaie, K., Lyeth, B. G. & Berman, R. F. in *New Therapeutics for Traumatic Brain Injury* 179–197 (Elsevier, 2017).
27. Kim, D. K. *et al.* A Case of Pleural Effusion and Pulmonary Edema Caused by Calcium Channel Blockers in a Patient of Systemic Hypertension. *Soonchunhyang Medical Science* **21**, 237–241 (2015).
28. Smith, S. W., Hauben, M. & Aronson, J. K. Paradoxical and bidirectional drug effects. *Drug safety* **35**, 173–189 (2012).
29. Picciotto, M. R., Brunzell, D. H. & Caldarone, B. J. Effect of nicotine and nicotinic receptors on anxiety and depression. *Neuroreport* **13**, 1097–1106 (2002).
30. Szklarczyk, D. *et al.* STITCH 5: augmenting protein–chemical interaction networks with tissue and affinity data. *Nucleic acids research* **44**, D380–D384 (2016).
31. Shulepko, M. A. *et al.* Human secreted protein SLURP-1 abolishes nicotine-induced proliferation, PTEN down-regulation and $\alpha 7$ -nAChR expression up-regulation in lung cancer cells. *International immunopharmacology* **82**, 106303 (2020).
32. Labarca, C. *et al.* Point mutant mice with hypersensitive $\alpha 4$ nicotinic receptors show dopaminergic deficits and increased anxiety. *Proceedings of the National Academy of Sciences* **98**, 2786–2791 (2001).
33. Deen, N. S., Huang, S. J., Gong, L., Kwok, T. & Devenish, R. J. The impact of autophagic processes on the intracellular fate of *Helicobacter pylori*: more tricks from an enigmatic pathogen? *Autophagy* **9**, 639–652 (2013).
34. Worst, D., Otto, B. & De Graaff, J. Iron-repressible outer membrane proteins of *Helicobacter pylori* involved in heme uptake. *Infection and immunity* **63**, 4161–4165 (1995).
35. Qujeq, D., Sadogh, M. & Savadkahi, S. Association between *helicobacter pylori* infection and serum iron profile. *Caspian journal of internal medicine* **2**, 266 (2011).

36. Berstad, A., Berstad, K. & Berstad, A. pH-Activated Phospholipase A 2: an Important Mucosal Barrier Breaker in Peptic Ulcer Disease. *Scandinavian journal of gastroenterology* **37**, 738–742 (2002).
37. Keine, D., Zelek, M., Walker, J. Q. & Sabbagh, M. N. Polypharmacy in an elderly population: enhancing medication management through the use of clinical decision support software platforms. *Neurology and therapy* **8**, 79–94 (2019).
38. Turgeon, J. & Michaud, V. Clinical decision support systems: Great promises for better management of patients' drug therapy. *Expert opinion on drug metabolism & toxicology* **12**, 993–995 (2016).
39. Umscheid, C. A., Margolis, D. J. & Grossman, C. E. Key concepts of clinical trials: a narrative review. *Postgraduate medicine* **123**, 194–204 (2011).
40. Bruhn, H. *et al.* Providing trial results to participants in phase III pragmatic effectiveness RCTs: a scoping review. *Trials* **22**, 1–32 (2021).
41. Marrie, R. A. *et al.* The challenge of comorbidity in clinical trials for multiple sclerosis. *Neurology* **86**, 1437–1445 (2016).
42. Ford, I. & Norrie, J. Pragmatic trials. *New England journal of medicine* **375**, 454–463 (2016).
43. Kipf, T. N. & Welling, M. Semi-supervised classification with graph convolutional networks. *International Conference on Learning Representations* (2017).
44. Yang, B., Yih, W.-t., He, X., Gao, J. & Deng, L. Embedding entities and relations for learning and inference in knowledge bases. *arXiv preprint arXiv:1412.6575* (2014).
45. Tatonetti, N. P., Patrick, P. Y., Daneshjou, R. & Altman, R. B. Data-driven prediction of drug effects and interactions. *Science translational medicine* **4**, 125ra31–125ra31 (2012).
46. Xu, H., Sang, S., Bai, P., Yang, L. & Lu, H. GripNet: Graph Information Propagation on Supergraph for Heterogeneous Graphs. *arXiv preprint arXiv:2010.15914* (2020).
47. Glorot, X. & Bengio, Y. *Understanding the difficulty of training deep feedforward neural networks* in *Proceedings of the thirteenth international conference on artificial intelligence and statistics* (2010), 249–256.
48. Klopfenstein, D. *et al.* GOATOOLS: A Python library for Gene Ontology analyses. *Scientific reports* **8**, 1–17 (2018).

- 627 49. Noble, W. S. How does multiple testing correction work? *Nature biotechnology* **27**, 1135–1137
628 (2009).

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Information

Supplementary Note

Predicted explanations offer novel mechanistic hypotheses

A patient's existing conditions may influence the mechanism of adverse reactions. We investigated one such example in which the combination of glipizide and lansoprazole may result in hypophosphatemia. APRILE predicted the likelihood of this side effect at 97% .

Glipizide is a second-generation oral hypoglycemic agent used to treat type 2 diabetes that targets the *CYP* gene family¹. Lansoprazole is a proton pump inhibitor commonly used to reduce gastric acid and treat gastric ulcers² and is metabolized by the liver using the P450 pathway; its targets include *PHOSPHOI*, a gene involved in the production of inorganic phosphate from organic compounds. Lansoprazole inhibits *PHOSPHOI*, disturbing this phosphorous cycle, which causes a drop in serum phosphate level. Meanwhile, Glipizide inhibits *CYP3A4*—a member of the P450 pathway—causing a delayed excretion and increased exposure to lansoprazole³, greatly reducing the amount of serum phosphate and ultimately leading to hypophosphatemia. Glipizide also inhibits *ABCC8* from promoting insulin production by the liver¹, thus having the same effect as direct insulin therapies. Insulin therapies may lead to a redistribution of phosphate due to insulin-stimulated cells actively acquiring phosphate from the blood⁴. This mechanism is not sufficient to cause hypophosphatemia but may worsen an existing condition⁵. Glipizide is used to treat type 2 diabetes, and these patients have more renal phosphate excretion than average due to osmotic diuresis⁶. This condition is a potential contributing factor to an existing hypophosphatemia adverse effect. Thus, APRILE was able to offer a comprehensive explanation for hypophosphatemia arising from the combination of glipizide and lansoprazole (Figure S4).

Supplementary Figure

Figure S1. Significant relations ($p < 0.05$) among side effect clusters and MeSH disease categories.

Figure S2. Jaccard distance between side effects vs. Euclidean distance between embeddings learned by APRILE-Pred for each pair of categories (22 in total) in Fig. 2d. The markers are shown if their Jaccard distance > 0 . The labels of markers are shown if their Jaccard distance > 0.006 .

Figure S3. APRILE explanation for panic disorder caused by drug pairs (10 possible drugs). Parameters used: regulation weight of 2.0, and probability threshold of 0.99. Drug nodes are blue and protein nodes are yellow.

Figure S4. Hypothesized mechanism for hypophosphatemia caused by glipizide-lansoprazole interaction.

Supplementary Table

Table S1. Macro-averaged AUROC for applying lazy training with different split rates and random states for parameter initialization

Split rate	Random-1	Random-2	Random-3	Random-4	Random-5
0.1	0.800364	0.800671	0.803084	0.797250	0.807832
0.2	0.844435	0.841982	0.846250	0.839958	0.850821
0.4	0.865966	0.861887	0.867489	0.864221	0.871715
0.6	0.870628	0.867482	0.869772	0.869908	0.875696
0.8	0.870904	0.867210	0.872412	0.870314	0.877834
0.9	0.872218	0.868020	0.873475	0.871330	0.878638

Table S2. Literature evidence for APRILE-Exp explanations for training data.

Side effect	Drug target?	Gene	Supported	Literature evidence
scotoma	target	PDE6B	yes	PDE6B mutations cause Retinitis Pigmentosa (RP) ⁷ . Scotoma is a symptom for RP patients. The polypharmacy may affect PDE6B similar to mutations.
scotoma	target	PDE6C	yes	PDE6C mutations cause achromatopsia ⁸ . Scotoma is a symptom of achromatopsia.
elevated prostate specific antigen	target	PDE11A	no	—
melanocytic naevus	non-target	CD7	yes	Patient with melanocytic nevus exhibited low CD7 expression levels ^{9,10}
bacterial pneumonia	target	CA4	indirect	CA4 is expressed in the lung and kidney and its inhibition alters CO ₂ equilibrium in tissues ¹¹ . In lambs with pneumonia, an insignificant decrease of carbonic anhydrase activity was observed ¹² .
post traumatic stress disorder (PTSD)	target	PDE6B	indirect	No direct evidence but PDE6B was differentially expressed in hippocampi of alcoholics, potentially affecting cortisol pathway dysregulation, which is observed in PTSD ¹³ .

Table S3. Side effect clustering based on the identified biological processes by APRILE.

GO	GO description
infection up- per respiratory	atelectasis, sleep disorder, candida infection, cholelithiasis, migraine, road traffic accident, osteoarthritis, arthritis, aptyalism, balance disorder, mod, sinusitis, intervertebral disc herniation, bruxism, arthropathy, gastric inflammation, dysuria, phlebothrombosis, infection upper respiratory, hive
aching joints	back ache, hyperglycaemia, hypoventilation, head ache, pain, dizziness, chest pain, high blood pressure, pleural pain, aching joints, edema extremities, deglutition disorder, infection urinary tract, anxiety, weight gain
dyspnea exer- tional	lung edema, pulmonary hypertension, bladder inflammation, allergic dermatitis, epistaxis, haemorrhage rectum, muscle weakness, dyspnea exertional, osteomyelitis, neuropathy peripheral, cardiomyopathy, aseptic necrosis bone, blood in urine, pancreatitis, arrhythmia
palpitation	agitated, allergies, bradycardia, arterial pressure nos increased, chronic obstructive airway disease, aching muscles, heart attack, acid reflux, palpitation, apoplexy, transient ischaemic attack, elevated cholesterol, cervicalgia, hyperhidrosis, gastrointestinal bleed
bladder reten- tion	abnormal movements, acidosis, haemorrhoids, bladder retention, aspiration pneumonia, asthma, acute brain syndrome, wheeze, nervous tension, osteoporosis, gall bladder, drug withdrawal, cerebral atrophy, hernia hiatal, head injury, difficulty in walking, haematochezia
petechia	peliosis, hypomagnesaemia, leucocytosis, back injury, duodenal ulcer, excoriation, ek-bom syndrome, sunburn, deafness, leukaemia, dermatitis medicamentosa, attempted suicide, breast pain, osteomalacia, pyelonephritis, decreased libido, phlebitis, ataxia, dysphasia, dental abscess, gingival pain, herpes simplex, abnormal eeg, excessive thirst, nystagmus, hydronephrosis, cardiac valvulopathy, cirrhosis, bleeding gums, petit mal, choking, antinuclear antibody positive, diabetic neuropathy, interstitial nephritis, disseminated lupus erythematosus, agranulocytoses, petechia, hydrocephalus, acne, hyperuricaemia, fecal occult blood, status epilepticus

Continued on next page

Table S3 – continued from previous page

GO	GO description
hyperaesthesia	apnea, strabismus, esophageal spasm, erosive gastritis, pneumonia staphylococcal, pleurisy, glossodynia, nasal sinus congestion, near syncope, erythema multiforme, mycobacterium tuberculosis infection, belching, azotaemia, colitis ischemic, sneezing, abnormal vision, diabetic acidosis, wrist fracture, polyp, amyotrophy, adrenal insufficiency, laryngitis, metabolic encephalopathy, hemiplegia, blood pressure abnormal, wound dehiscence, femoral neck fracture, sleep walking, pain pelvic, otitis media, chest infection, gynaecomastia, esophageal ulcer, hyperalimentation, macula lutea degeneration, fractured pelvis nos, hypernatraemia, syncope vasovagal, streptococcal infection, failure to thrive, hepatitis toxic, endocrine disorder, dermatitis exfoliative, shoulder pain, lymphoma, bone fracture, subarachnoid haemorrhage, blepharoptosis, palsies, myelodysplasia, atrioventricular block, easy bruisability, eye infection, labyrinthitis, disorder retinal, pericarditis, aortic aneurysm, hyperaesthesia, furuncle, birth defect, lymphedema, humerus fracture, anosmia, vitamin b 12 deficiency, cholecystitis chronic, bladder cancer
bruise	drug hypersensitivity, hypoglycaemia, hallucination, blurred vision, muscle spasm, cellulitis, abdominal pain upper, insomnia, dyspepsia, extremity pain, arteriosclerotic heart disease, fainting, incontinence, diabetes, bronchitis, joint swelling, dysarthria, amnesia, burning sensation, bruise, cardiac disease
iron deficiency anaemia	flatulence, peritonitis, cerebral infarct, acute pancreatitis, enlarged spleen, conjunctivitis, tinnitus, clotting, eye pain, cognitive disorder, psychoses, nasopharyngitis, rhinitis, hypochlorhaemia, small intestinal obstruction, arteriosclerosis, fibromyalgia, cholecystitis, extrasystoles ventricular, ild, nocturia, iron deficiency anaemia, bronchial spasm, hernia, lung neoplasms, facial palsy, disorder renal, hyperlipaemia, multiple sclerosis, bone spur, allergic rhinitis, carpal tunnel, fatty liver, dental caries, acute bronchitis, hepatic failure, movement disorder, rib fracture, haematemeses, aortic stenosis, coughing blood, carotid artery stenosis, disorder peripheral vascular

Continued on next page

Table S3 – continued from previous page

GO	GO description
haematoma	pain in throat, decreased body temperature, adynamic ileus, acute respiratory distress syndrome, anaphylactic reaction, diplopia, alopecia, infection viral, dry skin, haematoma, hepatitis, cataract, spinning sensation, enlarged liver, angiitis, glaucoma, hypotension orthostatic, cerebral haemorrhage, nephrosclerosis, gingivitis, blindness, eating disorder, cardiac murmur, increased platelet count, convulsions grand mal, elevated erythrocyte sedimentation rate, blood calcium increased, dysphonia, sinus bradycardia, mouth ulcer, musculoskeletal pain, cardiac ischemia, ulcer, angioedema, tachycardia ventricular, hemiparesis, bacteraemia, ischaemia, supraventricular tachycardia
embolism pulmonary	thrombocytopenia, sepsis, eruption, still birth, erythema, narcolepsy, abnormal gait, embolism pulmonary, paraesthesia, bone marrow failure, chill, pleural effusion, infection
lung infiltration	lung infiltration, bleeding, hypoxia, metabolic acidosis, drug toxicity nos, increased white blood cell count, encephalopathy, icterus, lobar pneumonia, chronic kidney disease, hypokalaemia, cardiac enlargement, adenopathy, cardiac failure, cardiovascular collapse, cutaneous ulcer
Continued on next page	

Table S3 – continued from previous page

GO	GO description
hepatitis b	hypoglycaemia neonatal, hypercapnia, lymphocytes decreased, haemolytic anaemia, optic disc edema, idiopathic thrombocytopenic purpura, glomerulonephritis, thrombotic microangiopathy, anal fissure, deafness neurosensory, bad breath, hepatic neoplasia, neoplasia, abuse, blepharitis, cachectic, breast dysplasia, scoliosis, autonomic instability, coeliac disease, diabetic nephropathy, neuritis, bone fracture spontaneous, retinal bleeding, bone inflammation, acute myeloblastic leukemia, salivary gland inflammation, ejaculation premature, nephrotic syndrome, contracture, spondylitis, raynauds phenomenon, synovitis, atrial septal defect, obstructive uropathy, rhagades, flashing lights, external ear infection, muscle inflammation, hypophosphataemia, chronic sinusitis, atrioventricular block second degree, aspergillosis, cmv infection, polymyositis, gangrene, cardiac tamponade, joint effusion, haemorrhage intracranial, umbilical hernia, lyell, squamous cell carcinoma of skin, pancreatic cancer, liver abscess, hepatic necrosis, breakthrough bleeding, lichen planus, gastric polyps, hepatitis b, renal mass, breast disorder, hyperpigmentation, costochondritis, cheilosis, hepatorenal syndrome, ebv infection, cerebral artery embolism, malignant hypertension, post nasal drip, nodule skin, neck mass, parotitis, dermatomyositis, als, bronchiectasis, aplasia pure red cell, pleural fibrosis, serum sickness, anemia aplastic, pneumonia primary atypical, ovarian cancer, aphthous stomatitis, left ventricular hypertrophy, arthritis bacterial, meningioma, dyspnoea paroxysmal nocturnal, paraplegia, irritation skin, bradyarrhythmia, epidural abscess, myasthenia gravis, histoplasmosis, pigmentation disorder, kidney transplant, folliculitis, chronic ulcerative colitis, uterine polyp, enterocolitis, macrocytosis, irregular menstrual cycle, testicular swelling, hyperphosphatemia, sacroiliitis, pyuria, testes pain

Continued on next page

Table S3 – continued from previous page

GO	GO description
pericardial effusion	abdominal distension, blood calcium decreased, disorder lung, ascites, sinus tachycardia, aspartate aminotransferase increase, deep vein thromboses, leucopenia, abnormal lfts, fungal disease, neutropenia, pyoderma, abscess, bulging, coma, bacterial infection, mucositis oral, pericardial effusion, esophagitis, bone pain, excess potassium, pulmonary arrest
convulsion	asystole, facial flushing, confusion, tremor, drowsiness, itch, loss of consciousness, blood sodium decreased, feeling unwell, convulsion, heart rate increased, neuropathy
asthenia	difficulty breathing, arterial pressure nos decreased, anaemia, fatigue, diarrhea, asthenia, dehydration, nausea
bursitis	lipoma, breast cancer, rhinorrhea, bursitis, nasal congestion, arthritis rheumatoid, herpes zoster, colonic polyp, benign prostatic hyperplasia, renal cyst, sensory disturbance, bundle branch block left, flu, eosinophil count increased, urosepsis, sleep apnea, gastroduodenal ulcer, rotator cuff syndrome, angina, muscle disorder
Continued on next page	

Table S3 – continued from previous page

GO	GO description
platelet disorder	<p> appendectomy, intestinal perforation, stridor, fib, vocal cord paralysis, retrocol- lis, cholangitis, coated tongue, biliary tract disorder, breast cyst, superficial throm- bophlebitis, meningitides, vestibular disorder, mitral valve disease nos, malabsorption, ureteral obstruction, encephalitis, pneumonia klebsiella, transfusion reaction, abortion spontaneous, schizophrenia, odynophagia, enuresis, personality disorder, cervicitis, adenitis, proctalgia, proctitis, spondylosis, apraxia, monoclonal gammopathy, mouth bleeding, dysaesthesia, adenoma, bundle branch block, enteritis, bone marrow trans- plant, glucose intolerance, duodenal ulcer haemorrhage, carcinoma of the colon, my- ocarditis, endocarditis, defaecation urgency, thrombophlebitis, gouty arthritis, respira- tory alkalosis, stress incontinence, scleroderma, premature menopause, candidiasis of vagina, seborrhoeic keratosis, actinic keratosis, glossitis, tenosynovitis, rectal polyp, pelvic inflammatory disease, schizoaffective disorder, borderline personality disorder, nephrogenic diabetes insipidus, portal hypertension, brain neoplasm, trigger finger, allergic vasculitis, congenital heart disease, breast tenderness, acute psychosis, ke- toacidosis, agoraphobia, eye injury, erysipelas, caesarean section, pyoderma gangreno- sum, choreoathetosis, cerumen impaction, paronychia, autonomic neuropathy, coro- nary artery bypass graft, temporal arteritis, inflammatory bowel disease, gastric cancer, volvulus, vaginal discharge, haemangioma, splenic infarction, aseptic meningitis, alco- holic intoxication, causalgia, mediastinal disorder, breast swelling, sjogrens syndrome, night cramps, diabetes insipidus, gastric ulcer perforated, platelet disorder, meningi- tis viral, esophageal cancer, uterine bleeding, hyperacusia, motor retardation, chronic hepatitis, food intolerance, connective tissue disease, anisocoria, menstrual disorder, pancreatic pseudocyst, breast inflammation, bunion, viral pharyngitis, cystitis intersti- tial, asthenopia, endometriosis </p>
Continued on next page	

Table S3 – continued from previous page

GO	GO description
flank pain	flank pain, upper gastrointestinal bleeding, gastroenteritis, skin lesion, dermatitides, intervertebral disc disorder, ischias, decubitus ulcer, atrioventricular block first degree, panic attack, nightmare, night sweat, angiopathy, amentia, neuralgia, dental pain, tooth disease, lung neoplasm malignant, muscle strain, ear ache, hip fracture, tendinitis, faecal incontinence, radiculopathy, electrolyte disorder, decreased hearing, bipolar disorder, paranoia, pharyngeal inflammation, light sensitivity, trigeminal neuralgia, obesity, diverticulitis, micturition urgency, cancer, spinal compression fracture, cerebral ischaemia, gastric ulcer, face pain, reflux esophagitis, kidney pain, thyroid disease
eye swelling	pneumothorax, hot flash, disease of liver, adenocarcinoma, malnourished, septic shock, hypothyroid, bleb, colon spastic, myoglobinuria, blepharospasm, breast lump, lung fibrosis, nephrolithiasis, rhabdomyolysis, aphasia, eye swelling, bilirubinaemia, aortic regurgitation, emphysema, abnormal ecg, candidiasis of mouth, mouth pain, proteinuria, bundle branch block right, ecchymoses, drug addiction, scar, black stools, spinal fracture, colitis, ankle fracture, abnormal laboratory findings, excessive sleepiness, nodule, cyst, psoriasis, gout, orthopnea, eczema, hematoma subdural
synovial cyst	appendicitis, aphonia, nephrocalcinosis, lower gi bleeding, diverticulosis, diabetic retinopathy, aneurysm, femur fracture, psychosexual disorder, floaters, thyroid neoplasia, prostatitis, adh inappropriate, mitral valve prolapse, malignant melanoma, atrial ectopic beats, elevated prostate specific antigen, fistula, atrial flutter, atrioventricular block complete, ventricular fibrillation, periodontal disease, basal cell carcinoma, granuloma, sinus headache, dry eye, right heart failure, bacterial pneumonia, brain concussion, pharyngitis streptococcal, seroma, nail disorder, hyperkeratosis, carcinoma of prostate, detached retina, polyarthritis, cholecystitis acute, tendon injury, synovial cyst, hernia inguinal, hepatic encephalopathy, fibroids
cough	respiratory failure, edema, neumonia, anorexia, constipated, acute kidney failure, abdominal pain, loss of weight, emesis, kidney failure, cough, cardiac decompensation, body temperature increased, afib
Continued on next page	

Table S3 – continued from previous page

GO	GO description
panic disorder	cold extremities, gastroenteritis viral, bleeding vaginal, bleeding hemorrhoids, muscle paresis, blood disorder, dislocation, laceration, pulmonary mass, polyneuropathy, post traumatic stress disorder, carcinoid syndrome, atherosclerosis, joint stiffness, panic disorder, colpitis, intestinal functional disorder, embolism, duodenitis, ear infection, goiter, metabolic alkalosis, parkinson, sick sinus syndrome, hyperthyroidism, epilepsy, melanocytic naevus, cramp, elevated triglycerides, multiple myeloma, onychomycosis, adverse drug effect, polyuria, glucosuria, insulin dependent diabetes mellitus, pancreatitis relapsing, major depression, cerebral vascular disorder, ovarian cyst, hesitancy, abdominal hernia, excessive salivation, heavy menstrual period, contact dermatitis, alcohol consumption, acute sinusitis, injury of neck, essential hypertension, cholecystectomies, chest wall pain, scotoma, hepatitis c, hiccough, neurogenic bladder
incisional hernia	hoarseness, amenorrhea, arthritis infective, acne rosacea, fascitis plantar, keratitis sicca, psoriatic arthritis, adjustment disorder, postherpetic neuralgia, tardive dyskinesia, hyperparathyroidism, myelopathy, tympanic membrane perforation, animal bite, cor pulmonale, epicondylitis, ingrowing nail, dermatitis seborrheic, eyelid diseases, periodontitis, breast enlargement, altered bowel function, sarcoidosis, optic neuritis, varicose vein, incisional hernia, abnormal mammogram, carotid bruit, conversion disorder, hay fever, venous insufficiency, tension headache, keratitis, fracture nonunion, empyema, calculus ureteric, dysphemia, esophageal stenosis
mumps	perirectal abscess, soft tissue infection, cervical vertebral fracture, renal tubular acidosis, duodenal ulcer perforation, substance abuse, colitis pseudomembranous, eustachian tube disorder, galactorrhea, hyperprolactinaemia, tracheitis, vitamin d deficiency, tooth impacted, polymyalgia rheumatica, diaphragmatic hernia, cutaneous mycosis, ganglion, vitreous haemorrhage, burns second degree, allergic alveolitis, cryptococcosis, cluster headache, tracheobronchitis, xerosis, postmenopausal bleeding, mumps, tinea pedis, hypogonadism, coccydynia, corneal abrasion
End of Table	

Table S4. Commonly enriched GO terms identified by APRILE-Exp for three peptic ulcers

GO	GO description	Literature evidence
GO:0045540	Regulation of cholesterol biosynthetic process.	Cholesterol increases the resistance of <i>H. pylori</i> , and the clinical manifestations of this bacteria include peptic ulcer disease ¹⁴ . <i>Helicobacter pylori</i> is associated with dyslipidemia but not with other risk factors of cardiovascular disease.
GO:0055114	Oxidation-reduction process.	Derangements of redox homeostasis in the gastrointestinal tract are believed to lead to peptic ulcer ¹⁵ .
GO:0019369	Arachidonic acid metabolic process.	Arachidonic acid metabolites play an important role in gastrointestinal homeostasis and have a significant impact in preventing or treating peptic ulcers ¹⁶ .
GO:0033384	Geranyl diphosphate biosynthetic process.	Farnesyl diphosphate and geranyl diphosphate are both affected by the direct interaction between bisphosphonates and a biosynthetic enzyme ¹⁷ . Bisphosphonates are drugs used as treatment for osteoporosis that can cause serious inflammation and ulcers in the GI tract ¹⁸ .
GO:0006805	Xenobiotic metabolic process.	Xenobiotics are widely used in the treatment of Parkinson disease, and Parkinson patients under treatment are prone to develop peptic ulcers, hence it is thought that there is a correlation ¹⁹ .
GO:0005789	Endoplasmic reticulum membrane.	Research suggests that the endoplasmic reticulum would be the main source for autophagosomes' membrane, which is formed for autophagic processes during an ulcerative infection ²⁰ .

Continued on next page

Table S4 – continued from previous page

GO	GO description	Literature evidence
GO:0020037	Heme binding.	Some proteins that are involved in heme binding are thought to be one important virulence factor of <i>H. pylori</i> ²¹ .
GO:0005506	Iron ion binding.	Iron ion binding capacity is thought to be lower in people with <i>H. pylori</i> infection ²² .
GO:0102568	Phospholipase A2 activity consuming 1,2-dioleoylphosphatidylethanolamine.	This is directly correlated to the arachidonic acid metabolic process and to gastrointestinal ulceration ²³ .
GO:0004623	Phospholipase A2 activity.	Both this term and the previous one refer to the activity of phospholipase A2, which is a mucosal barrier breaker in peptic ulcer disease ²⁴ .
GO:0008395	Steroid hydroxylase activity.	Steroid hydroxylase is involved in steroid biosynthesis and a too high concentration of steroid can lead to gastrointestinal ulcers and bleeding ²⁵ .
End of Table		

References

1. Lebovitz, H. E. & Feinglos, M. N. Mechanism of action of the second-generation sulfonylurea glipizide. *The American journal of medicine* **75**, 46–54 (1983).
2. Zimmermann, A. E. & Katona, B. G. Lansoprazole: a comprehensive review. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* **17**, 308–326 (1997).
3. Riley, R., Parker, A., Trigg, S. & Manners, C. Development of a generalized, quantitative physicochemical model of CYP3A4 inhibition for use in early drug discovery. *Pharmaceutical research* **18**, 652–655 (2001).
4. Kjeldsen, S. E., Moan, A., Petrin, J., Weder, A. B. & Julius, S. Effects of increased arterial epinephrine on insulin, glucose and phosphate. *Blood pressure* **5**, 27–31 (1996).
5. Liamis, G., Milionis, H. & Elisaf, M. Medication-induced hypophosphatemia: a review. *QJM: An International Journal of Medicine* **103**, 449–459 (2010).
6. Moe, S. M. Disorders involving calcium, phosphorus, and magnesium. *Primary Care: Clinics in Office Practice* **35**, 215–237 (2008).
7. Karali, M. *et al.* Clinical and genetic analysis of a European cohort with pericentral retinitis pigmentosa. *International journal of molecular sciences* **21**, 86 (2020).
8. Chang, B. *et al.* A homologous genetic basis of the murine cpfl1 mutant and human achromatopsia linked to mutations in the PDE6C gene. *Proceedings of the National Academy of Sciences* **106**, 19581–19586 (2009).
9. Brazzelli, V. *et al.* Recurrence of mycosis fungoides on multiple melanocytic nevi: a case report and review of the literature. *Case reports in dermatology* **4**, 92–97 (2012).
10. Cerroni, L., Gatter, K. & Kerl, H. *An illustrated guide to skin lymphoma* (John Wiley & Sons, 2008).
11. Zhu, X. & Sly, W. Carbonic anhydrase IV from human lung. Purification, characterization, and comparison with membrane carbonic anhydrase from human kidney. *Journal of Biological Chemistry* **265**, 8795–8801 (1990).
12. Ekin, S., Berber, I., Kozat, S. & Gunduz, H. Selected trace elements and esterase activity of carbonic anhydrase levels in lambs with pneumonia. *Biological trace element research* **112**, 233–239 (2006).

13. McClintick, J. N. *et al.* Stress–response pathways are altered in the hippocampus of chronic alcoholics. *Alcohol* **47**, 505–515 (2013).
14. McGee, D. J. *et al.* Cholesterol enhances *Helicobacter pylori* resistance to antibiotics and LL-37. *Antimicrobial agents and chemotherapy* **55**, 2897–2904 (2011).
15. Cherkas, A. & Zarkovic, N. 4-Hydroxynonenal in redox homeostasis of gastrointestinal mucosa: Implications for the stomach in health and diseases. *Antioxidants* **7**, 118 (2018).
16. Isselbacher, K. The role of arachidonic acid metabolites in gastrointestinal homeostasis. *Drugs* **33**, 38–46 (1987).
17. Tsoumpira, M. K. *et al.* The inhibition of human farnesyl pyrophosphate synthase by nitrogen-containing bisphosphonates. Elucidating the role of active site threonine 201 and tyrosine 204 residues using enzyme mutants. *Bone* **81**, 478–486 (2015).
18. Cadarette, S. *et al.* Comparative gastrointestinal safety of weekly oral bisphosphonates. *Osteoporosis international* **20**, 1735–1747 (2009).
19. Carmody, R. N., Turnbaugh, P. J., *et al.* Host-microbial interactions in the metabolism of therapeutic and diet-derived xenobiotics. *The Journal of clinical investigation* **124**, 4173–4181 (2014).
20. Deen, N. S., Huang, S. J., Gong, L., Kwok, T. & Devenish, R. J. The impact of autophagic processes on the intracellular fate of *Helicobacter pylori*: more tricks from an enigmatic pathogen? *Autophagy* **9**, 639–652 (2013).
21. Worst, D., Otto, B. & De Graaff, J. Iron-repressible outer membrane proteins of *Helicobacter pylori* involved in heme uptake. *Infection and immunity* **63**, 4161–4165 (1995).
22. Qujeq, D., Sadogh, M. & Savadkahi, S. Association between *helicobacter pylori* infection and serum iron profile. *Caspian journal of internal medicine* **2**, 266 (2011).
23. Hornbeck, P. V. *et al.* PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic acids research* **43**, D512–D520 (2015).
24. Berstad, A., Berstad, K. & Berstad, A. pH-Activated Phospholipase A 2: an Important Mucosal Barrier Breaker in Peptic Ulcer Disease. *Scandinavian journal of gastroenterology* **37**, 738–742 (2002).

- 706 25. Narum, S., Westergren, T. & Klemp, M. Corticosteroids and risk of gastrointestinal bleeding: a
707 systematic review and meta-analysis. *BMJ open* 4 (2014).