GP1R-dependent estrogen signaling increases cardiac GCN5L1 expression and MCAD activity

Janet R. Manning^{1,2,3*}, Dharendra Thapa^{1,2,3,#}, Manling Zhang^{1,2,3}, Michael W. Stoner^{1,2,3}, John C. Sembrat⁴, Mauricio Rojas⁵, and Iain Scott^{1,2,3*}

¹Division of Cardiology, Department of Medicine, University of Pittsburgh, Pittsburgh, PA. ²Vascular Medicine Institute, Department of Medicine, University of Pittsburgh, Pittsburgh, PA. ³Center for Metabolism and Mitochondrial Medicine, Department of Medicine, University of Pittsburgh, Pittsburgh, PA. ⁴Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, University of Pittsburgh, Pittsburgh, PA. ⁵Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, The Ohio State University, Columbus, OH. [#]Current address: Department of Exercise Physiology, West Virginia University

***To whom correspondence should be addressed:**

Iain Scott, PhD, FAHA, FCVS	Janet Manning, PhD
Department of Medicine	Department of Medicine
University of Pittsburgh	University of Pittsburgh
BST E1259	BST E1200
200 Lothrop Street	200 Lothrop Street
Pittsburgh, PA 15261	Pittsburgh, PA 15261
Email: iain.scott@pitt.edu	Email: manningj@pitt.edu
Tel: (412) 648-7691	Tel: (412) 648-8018

ABSTRACT

Reversible lysine acetylation regulates the activity of cardiac metabolic enzymes, including those controlling fuel substrate metabolism. Mitochondrial-targeted GCN5L1 and SIRT3 have been shown to regulate the acetylation status of mitochondrial enzymes, which results in alterations to the relative oxidation rates of fatty acids, glucose, and other fuels for contractile activity. However, the role that lysine acetylation plays in driving metabolic differences between male and female hearts is not currently known. In this study, we report that estrogens induce the expression of GCN5L1 via GPER agonism in cardiac cells, which increases the enzymatic activity and acetylation status of the fatty acid oxidation enzyme medium chain acyl-CoA dehydrogenase (MCAD).

INTRODUCTION

Improved understanding of the physiological and metabolic differences between men and women may allow us to develop new therapies that can address sex-based disparities in cardiac disease treatment outcomes. Sex hormones testosterone and estrogen, as well as chromosomal effects, may contribute to sex-based differences. Pre-menopausal women exhibit increased estrogen levels relative to men and post-menopausal women, which results in greater activation of estrogen receptors in the myocardium. These are comprised of the canonical estrogen receptors alpha and beta ($ER\alpha$ and $ER\beta$), and the G-protein coupled estrogen receptor (GPER, or GPR30).¹ Canonical ERs are targeted directly to the nucleus, and interact with ER responsive elements (EREs) within the genome to regulate transcription, while GPER activation results in a cascade of posttranslational modifications in the cell that may also ultimately drive genomic responses.²⁻⁴

Estrogen receptor activation has been associated with changes in the abundance and activity of numerous enzymes involved in glucose and fatty acid energy metabolism, which result in significant sexual dimorphism in cardiac metabolic profiles. Of particular note, women exhibit greater cardiac fatty acid uptake and oxidation relative to men under both normal and pathophysiological conditions.⁵ Estrogen modulates the expression of cardiac metabolic proteins, and upregulates proteins that impact fatty acid metabolism, including PGC-1 α and acyl-CoA dehydrogenases (ACADs).⁶⁻⁸

Consequently, the presence of estrogen has a significant impact on fuel substrate utilization in the heart.

The posttranslational acetylation of non-nuclear targets has emerged as a critical regulator of metabolic activity in the heart. In mitochondria, GCN5L1 and SIRT3 have been reported to increase and decrease, respectively, the acetylation status of enzymes that metabolize fatty acids and glucose.^{9–15} However, sex differences in the acetylation of metabolic proteins in cardiac mitochondria have not been investigated. We hypothesized that differences in the expression of GCN5L1 and SIRT3 between men and women may change the acetylation status and activity of enzymes involved in glucose and fatty acid metabolism, and that estrogen signaling may drive this process.

The studies presented here demonstrate that mitochondrial protein acetylation is increased in female mice relative to males, which is associated with sex-dependent elevations in GCN5L1 abundance. In addition, we show that estrogen directly increases GCN5L1 expression in human-derived cardiac cells, and that GCN5L1 is decreased in the hearts of postmenopausal women relative to younger women. The primary mechanism for estrogen-mediated GCN5L1 upregulation is identified as GPER activation, through a transcription-independent pathway. Finally, we determine that estrogen-dependent acetylation of MCAD is dependent on GCN5L1, and that loss of GCN5L1 results in diminished MCAD activity.

METHODS

Human tissues

Fresh human cardiac tissue samples were collected from the left ventricles of organ donors deemed not suitable for transplant, under a protocol approved by the University of Pittsburgh Committee for Oversight of Research and Clinical Training (CORID). Tissues were flash-frozen and stored at -80 °C until processing. Post menopause: range = 65-86 years, median = 69 years, N = 7. Pre menopause: range = 22-39 years, median = 36 years, N = 5.

Animal care and use

All housing and experiments in mice were conducted in accordance with the guidelines established by the National Institutes of Health, and approved by the University of Pittsburgh Institutional Animal Care and Use Committee. Male and female C57BL/6J mice (aged 8-10 weeks) were purchased from The Jackson Laboratory, and maintained on a regular chow diet with a 12 h light/12 h dark light cycle.

Cell culture and drug treatments

AC16 cells (a proliferating cell line derived from human cardiomyocytes¹⁶) were purchased from Millipore. Stable GCN5L1 knockdown was generated as previously described.¹⁷ Cells were treated with 10 nM 17 β -estradiol (E2), ICI 182, 780 (Fulvestrant; an ER α and ER β antagonist with GPER agonist activity¹⁸⁻²⁰), G-1 (a selective GPER

agonist), G-36 (a selective GPER antagonist), MG-132 (a 26S protease inhibitor), and/or cycloheximide (CHX; a translation inhibitor).

Mitochondrial Isolation

Mitochondrial fractions were purified from tissue and cells using the Qproteome Mitochondrial Isolation Kit (Qiagen) according to the manufacturer's instructions. Briefly, samples were homogenized in cold Lysis Buffer, and centrifuged at 1000 *g*. Supernatant containing the cytosolic fraction was discarded, and the pellet was re-suspended and processed in cold Disruption Buffer by shearing through a 25 *g* needle and syringe. Samples were centrifuged at 1000 *g*, the supernatant was collected, and centrifuged again at 6000 *g* to pellet the mitochondrial fraction. The pellet was washed in Storage Buffer, and then used for subsequent immunoblot or MCAD activity studies as described below.

Immunoblotting

Tissue, cells, or purified mitochondria were lysed in 1% CHAPS buffer. Protein was quantitated using a BioDrop μ LITE analyzer (BioDrop), and equal amounts were loaded on a 12% SDS-PAGE gel, before transfer to nitrocellulose membranes. Membranes were blocked using Odyssey blocking buffer and incubated in primary antibodies overnight (α Tubulin, 1:1000, Cell Signaling; SIRT3, 1:1000, Cell Signaling; MCAD 1:1000, Cell Signaling; GCN5L1 1:500, generated as previously described²¹), followed by incubation at room temperature with fluorescent secondary antibodies for 1 h (800

nm anti-rabbit, LiCor). Bands were visualized using an Odyssey Imager, and quantitated using Image Studio Lite v 5.2 (LiCor).

Quantitative RT-PCR

RNA was isolated from tissue or cells using RNEasy kit (Qiagen). RNA was quantified and 500 ng-1000 ng was used to generate cDNA using Maxima Reverse Transcriptase (Thermo Fisher). Quantitative PCR was performed using SYBR-Green (Thermofisher) and primers for GCN5L1 or SIRT3. GAPDH or PPIA were used for normalization.

MCAD activity assays

MCAD activity was assessed using a DCPIP/PES-based assay as previously described. Briefly, DCPIP (50 μ M), PES (2 mM), NEM (0.2 mM), KCN (0.4 mM), Triton X-100 (0.10%), and lysate were added to ice cold potassium phosphate buffer (0.1 M). MCAD substrate octanoyl-CoA was added to a final concentration of 40 μ M, and then warmed to 37 °C for 5 min. Absorbance was read at 600 nm, and then normalized to protein concentration.

Statistical Analysis

Statistical analyses were performed using GraphPad Prism 8.3. Student's t-tests were used for simple comparisons between groups. One-way Analyses of Variance (ANOVA) was used to compare more than two groups, followed by post-hoc Student's t-tests. For studies examining multiple time-points a two-way ANOVA was used with post-hoc

Sidak's multiple comparisons tests. A *P* value <0.05 was regarded as significant. All data are represented as the mean ± SEM.

RESULTS

Mitochondrial acetylation and GCN5L1 expression is increased in the hearts of female mice compared to male mice

To determine whether mitochondrial protein acetylation status is different between sexes, we isolated mitochondria from the hearts of male and female C57BL/6J mice, and immunoblotted for acetylated lysine residues. We observed a modest but significant increase in the intensity of bands in female mice compared to male mice (**Figure 1A**). We next examined whether the increase in mitochondrial acetylation is associated with changes in the expression of proteins that regulate mitochondrial protein acetylation. No changes were observed in the expression of the mitochondrial-targeted deacetylase SIRT3 (**Figure 1B**). However, a significant increase in both GCN5L1 mRNA and protein was observed (**Figures 1C and 1D**). Based on these data, we conclude that increased acetylation in female cardiac mitochondria is driven by increased GCN5L1 abundance.

Estrogen increases GCN5L1 expression in human cardiomyocytes via GPER

To determine if changes in GCN5L1 abundance are present in a clinically relevant setting, we analyzed heart tissue obtained from female patients of pre- and post-

menopausal age. Immunoblotting revealed that cardiac tissues from women after menopause, when estrogen levels are lower, have a significantly lower GCN5L1 protein abundance (**Figure 2A**). As estrogen has been reported to mediate several of the sex differences observed in human myocardial tissue,⁴ we next determined whether estrogen induces GCN5L1 expression directly. Treatment of AC16 cells (derived from human ventricular cardiomyocytes)¹⁶ with 17- β estradiol (E2) resulted in significantly increased levels of GCN5L1 protein (**Figure 2B**). To determine whether signaling for increased GCN5L1 expression was through canonical estrogen receptors, we incubated AC16 cells with ICI 182,780 (Fulvestrant), a potent inhibitor of ER α and ER β . Surprisingly, rather than blocking GCN5L1 induction, ICI 182,780 additively increased GCN5L1 levels, and produced a robust increase in GCN5L1 even in the absence of E2 (**Figure 2C**). ICI 182,780, in addition to blocking ER α and ER β , also has been reported to act as a partial agonist for the G-protein coupled estrogen receptor (GPER).^{19,22,23} We therefore examined whether GPER played a role in estrogen-mediated GCN5L1 induction using the GPER agonist G-1, and the GPER antagonist G-36. Incubation with G-1 significantly increased GCN5L1 protein abundance, while G-36 blocked E2-mediated GCN5L1 expression (**Figure 2D**). From these data, we conclude that GCN5L1 abundance is increased by estrogen exposure via GPER-mediated signaling.

Estrogen drives GCN5L1 translation

To understand the mechanism of GPER-induced GCN5L1 elevations, we monitored mRNA levels in AC16 cells after exposure to E2 or G-1 using qPCR. Surprisingly, no

change in mRNA was observed at any of the time points measured, suggesting that E2 control of GCN5L1 expression occurs downstream of transcription (**Figure 3A**). To determine whether GCN5L1 protein elevation was due to an estrogen-induced reduction in protein degradation, the effects of 26S proteasomal inhibitor MG132 on GCN5L1 expression were evaluated. It was expected that if estrogen signaling elevates GCN5L1 levels by reducing protease activity, a blockade of protease activity would normalize protein levels in vehicle-treated cells, and the difference observed in E2-treated cells would disappear. Data showed that this was not the case, and rather MG132 amplified the increase in protein observed in the presence of E2 or G-1 (**Figure 3B**). We next examined whether estrogen alters GCN5L1 mRNA translation. To test this hypothesis, E2 and G-1 treated AC16 cells were incubated with the translational inhibitor cycloheximide (CHX). CHX treatment effectively normalized GCN5L1 levels, indicating that differences in protein expression may be attributed to GCN5L1 translational regulation (**Figure 3C**).

Estrogen-induced acetylation and activation of MCAD is reduced when GCN5L1 levels are depleted

Cardiac GCN5L1 has been reported to mediate the acetylation and activation of acyl-CoA dehydrogenases (ACADs), which mediate fatty acid breakdown.^{12,24} Among the ACADs, MCAD has been repeatedly identified as a target regulated by estrogens in the heart.⁶⁻⁸ We therefore tested whether GCN5L1 may link estrogen receptor agonism to increases in MCAD acetylation and activity. Estrogen treatment resulted in increased

214 MCAD activity in AC16 cells, which was blocked in GCN5L1-depleted cells (**Figure 4A**).
 215 When GCN5L1 was silenced, MCAD acetylation was significantly reduced in E2 and G-
 216 1 treated cells, relative to control cells under the same conditions (**Figure 4B**). These
 217 data suggest that E2- and GPER agonist-induced MCAD activation occurs via GCN5L1.
 218

DISCUSSION

Here we determined that mitochondrial protein acetylation, and GCN5L1 expression, are elevated in female mouse hearts compared to male. We find that estrogen upregulates GCN5L1 via GPER, and pharmacological block of GPER ablates induction. GCN5L1 is also elevated in pre-menopausal women, where estrogen levels are higher, compared to post-menopausal women. E2 and G-1 do not alter GCN5L1 gene transcription or proteasomal degradation; instead translational blockade prevents GCN5L1 induction. Finally, we determine that the loss of GCN5L1 blocks estrogen-mediated acetylation and activation of MCAD. These data point to a significant role for GCN5L1 in estrogen-mediated regulation of cardiac fuel metabolism (summarized in **Figure 4C**).

Significant differences in cardiac physiology and pathology between men and women are well established. Pre-menopausal women are largely protected from cardiovascular disease (CVD) compared to men, but this advantage is reduced with age.²⁵ Estrogen loss has been suggested to be a major mediator of this effect. Postmenopausal women become more susceptible to left ventricular diastolic dysfunction, and hormone replacement therapy mitigates this effect.²⁶ Ovariectomized mice and rats are similarly more susceptible to insults to the heart, including pressure overload,²⁷ AngII induced hypertrophy,²⁸ and diabetes-associated myofilament sensitization to calcium.^{29,30} Ovariectomized mice exhibit a faster onset of obesity-driven heart failure, and show earlier signs of cardiac mitochondrial dysfunction, including elevated ROS production

and swelling.³¹ Mice lacking estrogen receptors are found to be more sensitive to IR injury and hypertensive cardiomyopathy.^{28,32} Although less well-studied, testosterone also plays a role in driving sexual dimorphism between male and female individuals. Since estrogen was sufficient to reproduce an increase in GCN5L1 production, we have not evaluated the effects of testosterone. However, we cannot discount the possibility of an effect, and further studies are required to make this determination.

In recent years, GPER has taken a central role in our understanding of how estrogen impacts cardiac function and resiliency.³³ GPER activation in cardiomyocytes lacking classical ER α and ER β receptors was reported to alter intracellular calcium influx.³⁴ In addition, GPER activation is associated with protection from ischemia-reperfusion injury, dependent on PI3K activation.³⁵ Agonism of GPER with G-1 is reported to protect estrogen-deficient rats from LV remodeling.³⁶ G-1 is also reported to inhibit the opening of the mitochondrial membrane permeability pore (mPTP),³⁷ and reduces the upregulation of inflammatory cytokines TNF- α , IL-1 β , and IL-6.³⁸ Our earlier observation that GCN5L1 protects the heart from I/R injury¹⁷ raises the possibility that GPER-mediated upregulation of GCN5L1 may be an additional mechanism by which estrogen protects the heart.

The acetylation status of several mitochondria-localized proteins has been reported to impact their function and stability. A key driver of acetylation status in mitochondria is the deacetylase SIRT3, which is expressed in myocardial tissue, and has been shown to regulate the function of several mitochondrial proteins involved in oxidative

phosphorylation, membrane integrity, and redox homeostasis (See recent review by Chen et al.³⁹). However, no difference in cardiac SIRT3 expression was identified between male and female mice, despite altered mitochondrial acetylation levels, suggesting that dimorphic effects in acetylation are due to GCN5L1 only.

GPER agonism drives translation and production of GCN5L1 protein, but at no time point after GPER activation was GCN5L1 message elevated. A recent publication by Lv et al. suggests that GCN5L1 abundance is regulated post-transcriptionally in diabetic kidney cells.⁴⁰ We did not observe evidence that G-1 blocks the degradation of GCN5L1 by the 26S proteasome. Interestingly, there are reports of GCN5L1 acting as a translational coactivator to ER α in HeLa cells, binding directly to both the receptor and the corepressor element MTA1.⁴¹ That study did not evaluate a direct interaction between GCN5L1 and GPER, and although a mechanism by which GCN5L1 might interact directly with GPER was briefly considered, our immunoprecipitation studies did not support the direct binding of GPER to GCN5L1 (data not shown).

We demonstrate here that GCN5L1 is required for estrogen and GPER agonism to upregulate the acetylation and activity of medium-chain acyl-CoA dehydrogenase (MCAD) in cardiac-derived cells. MCAD is a mitochondria-localized enzyme that catalyzes the rate-limiting step in the β -oxidation of medium chain fatty acids, the α,β -dehydrogenation of fatty acyl-CoA. MCAD plays a key role in the progression of myocardial metabolic dysregulation induced by heart failure. TAC-mediated metabolic changes include the downregulation of MCAD,⁴² and gene delivery of MCAD to the

heart protects against pressure overload-induced pathological remodeling.⁴³ Subjecting mice to a high-fat diet was reported to increase contractile recovery after MI, concurrent with an increase in MCAD activity.⁴⁴ Our laboratory and others have shown that high-fat feeding increases the activity of multiple ACADs through increased acetylation in response to GCN5L1 expression.¹² In addition, MCAD has been reported previously to be regulated estrogen; E2 has been reported to upregulate or preserve cardiac MCAD expression via PGC-1.^{6–8} Studies of skeletal muscle biopsies report that women express higher levels of MCAD,⁴⁵ and MCAD expression was reported to increase in men after treatment with E2.⁴⁶ The studies reported here show for the first time that estrogen may also impact through posttranslational acetylation mediated by GCN5L1 expression.

In summary, we demonstrate here for the first time that GCN5L1 is upregulated by estrogen signaling in both mouse and human myocardium, and in cultured cardiomyocytes. The mechanism of upregulation is identified as GPER-mediated signaling. MCAD is identified as a target of acetylation and activation in the presence of E2 or GPER agonists, which is dependent on GCN5L1 expression. These findings shed new light on the role that posttranslational acetylation, mediated by GCN5L1, may play in the differences observed between men and women in cardiac metabolism.

307 **ACKNOWLEDGEMENTS**

308

309 This work was supported by NIH/NHLBI R01 (HL132917 & HL147861) research grants
310 to IS, and a UPMC CMRF Research Grant to JRM.

Figure Legends

Figure 1: Mitochondrial protein acetylation and GCN5L1 are upregulated in female

mice. A. Immunoblotting of mitochondrial lysate fractions from male and female mice demonstrate that females exhibit modestly but significantly higher levels of total protein acetylation. N = 10, B. GCN5L1, but not SIRT3, mRNA is significantly increased in female hearts relative to males. C. GCN5L1 protein levels are significantly increased in the myocardium of female mice compared to male mice. N = 10, * = $p < 0.05$, *** = $p < 0.001$ vs. male.

Figure 2: Estrogen drives GCN5L1 expression via GPER A. Human tissues from female patients of pre-menopausal (PRE) or post-menopausal (POST) age. B. Human derived AC16 cells incubated with E2 show an increase in GCN5L1. C. GCN5L1 immunoblotting after incubation for 24 hours with E2 and or the ERa/ERb antagonist ICI 182, 780. N = 4, * = $p < 0.05$, **** = $p < 0.0001$ vs. vehicle. D. GCN5L1 is elevated after incubation for 24 hours with E2 and the GPER agonist G-1, and is blocked in the presence of GPER antagonist G-36. N = 5-6, * = $p < 0.05$, ** = $p < 0.01$ vs. vehicle.

Figure 3: Estrogen promotes GCN5L1 translation. A. Expression of GCN5L1 mRNA levels determined by qPCR after treatment with E2 or G-1. N = 3, ** = $p < 0.01$ vs. vehicle. B. GCN5L1 levels are significantly elevated by blocking 26S degradation, suggesting that GPER agonism does not increase GCN5L1 via impaired proteolysis. N

= 4, * = $p < 0.05$, *** = $p < 0.001$ vs. control. C. GCN5L1 induction is blocked by CHX, a translation inhibitor. N = 4 * = $p < 0.05$, *** = $p < 0.001$ vs. control.

Figure 4 : GCN5L1 is required for estrogen-mediated MCAD acetylation and activation. A. E2 and G-1 raise MCAD activity, while GCN5L1 shRNA knockdown blocks this effect. N = 8. B. Acetylation of MCAD is significantly reduced in the presence of E2 and G-1 when GCN5L1 is absent. N = 4 * = $p < 0.05$, ** = $p < 0.01$ vs. control, † = $p < 0.01$ vs. vehicle. C. Schematic representing the hypothesized mechanism of action of estrogen on GCN5L1 in cardiomyocytes.

- 342 1. Aryan, L. *et al.* The role of estrogen receptors in cardiovascular disease.
343 *International Journal of Molecular Sciences* **21**, 1–26 (2020).
- 344 2. Meyer, M. R., Prossnitz, E. R. & Barton, M. The G protein-coupled estrogen
345 receptor GPER/GPR30 as a regulator of cardiovascular function. *Vascul.*
346 *Pharmacol.* **55**, 17–25 (2011).
- 347 3. Menazza, S. & Murphy, E. The Expanding Complexity of Estrogen Receptor
348 Signaling in the Cardiovascular System. *Circ. Res.* **118**, 994–1007 (2016).
- 349 4. Ueda, K., Adachi, Y., Liu, P., Fukuma, N. & Takimoto, E. Regulatory Actions of
350 Estrogen Receptor Signaling in the Cardiovascular System. *Frontiers in*
351 *Endocrinology* **10**, 909 (2020).
- 352 5. Mather, K. J. & DeGrado, T. R. Imaging of myocardial fatty acid oxidation.
353 *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **1861**, 1535–1543 (2016).
- 354 6. Schreiber, S. N., Knutti, D., Brogli, K., Uhlmann, T. & Kralli, A. The transcriptional
355 coactivator PGC-1 regulates the expression and activity of the orphan nuclear
356 receptor estrogen-related receptor α (ERR α). *J. Biol. Chem.* **278**, 9013–9018
357 (2003).
- 358 7. Hsieh, Y.-C. *et al.* Inhibition of cardiac PGC-1 α expression abolishes ER β
359 agonist-mediated cardioprotection following trauma-hemorrhage. *FASEB J.* **20**,
360 1109–1117 (2006).
- 361 8. Kim, M. S., Shigenaga, J. K., Moser, A. H., Feingold, K. R. & Grunfeld, C.
362 Suppression of estrogen-related receptor α and medium-chain acyl-coenzyme A
363 dehydrogenase in the acute-phase response. *J. Lipid Res.* **46**, 2282–2288 (2005).
- 364 9. Kendrick, A. A. *et al.* Fatty liver is associated with reduced SIRT3 activity and

- mitochondrial protein hyperacetylation. *Biochem. J.* **433**, 505–14 (2011).
10. Alrob, O. A. *et al.* Obesity-induced lysine acetylation increases cardiac fatty acid oxidation and impairs insulin signalling. *Cardiovasc. Res.* **103**, 485–97 (2014).
11. Koentges, C. *et al.* SIRT3 deficiency impairs mitochondrial and contractile function in the heart. *Basic Res Cardiol* **110**, 36 (2015).
12. Thapa, D. *et al.* Acetylation of mitochondrial proteins by GCN5L1 promotes enhanced fatty acid oxidation in the heart. *Am. J. Physiol. Heart Circ. Physiol.* **313**, H265–H274 (2017).
13. Manning, J. R. *et al.* Loss of GCN5L1 in cardiac cells disrupts glucose metabolism and promotes cell death via reduced Akt/mTORC2 signaling. *Biochem. J.* **476**, 1713–1724 (2019).
14. Thapa, D. *et al.* Adropin regulates pyruvate dehydrogenase in cardiac cells via a novel GPCR-MAPK-PDK4 signaling pathway. *Redox Biol.* **18**, 25–32 (2018).
15. Carnevale, I. *et al.* SIRT1-SIRT3 Axis Regulates Cellular Response to Oxidative Stress and Etoposide. *J. Cell. Physiol.* **232**, 1835–1844 (2017).
16. Davidson, M. M. *et al.* Novel cell lines derived from adult human ventricular cardiomyocytes. *J. Mol. Cell. Cardiol.* **39**, 133–147 (2005).
17. Manning, J. R. *et al.* Cardiac-specific deletion of GCN5L1 restricts recovery from ischemia-reperfusion injury. *J. Mol. Cell. Cardiol.* **129**, 69–78 (2019).
18. Robertson, J. F. ICI 182,780 (Fulvestrant)--the first oestrogen receptor down-regulator--current clinical data. *Br. J. Cancer* **85 Suppl 2**, 11–4 (2001).
19. Thomas, P., Pang, Y., Filardo, E. J. & Dong, J. Identity of an estrogen membrane receptor coupled to a G protein in human breast cancer cells. *Endocrinology* **146**,

624–632 (2005).

20. Filardo, E. J., Quinn, J. A., Bland, K. I. & Frackelton, J. Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol. Endocrinol.* **14**, 1649–1660 (2000).
21. Scott, I., Webster, B. R., Li, J. H. & Sack, M. N. Identification of a molecular component of the mitochondrial acetyltransferase programme: a novel role for GCN5L1. *Biochem. J.* **443**, 655–661 (2012).
22. Filardo, E. J., Quinn, J. A., Bland, K. I. & Frackelton, J. Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol. Endocrinol.* **14**, 1649–1660 (2000).
23. Prossnitz, E. R. & Barton, M. Signaling, physiological functions and clinical relevance of the G protein-coupled estrogen receptor GPER. *Prostaglandins and Other Lipid Mediators* **89**, 89–97 (2009).
24. Thapa, D. *et al.* Cardiomyocyte-specific deletion of GCN5L1 in mice restricts mitochondrial protein hyperacetylation in response to a high fat diet. *Sci. Rep.* **10**, (2020).
25. Cho, L. *et al.* Summary of Updated Recommendations for Primary Prevention of Cardiovascular Disease in Women: JACC State-of-the-Art Review. *J. Am. Coll. Cardiol.* **75**, 2602–2618 (2020).
26. Pines, A. *et al.* Menopause-induced changes in doppler-derived parameters of aortic flow in healthy women. *Am. J. Cardiol.* **69**, 1104–1106 (1992).

27. Mori, T. *et al.* Enhanced cardiac inflammation and fibrosis in ovariectomized hypertensive rats: A possible mechanism of diastolic dysfunction in postmenopausal women. *Hypertens. Res.* **34**, 496–502 (2011).
28. Pedram, A. *et al.* Estrogen inhibits cardiac hypertrophy: Role of estrogen receptor- β to inhibit calcineurin. *Endocrinology* **149**, 3361–3369 (2008).
29. Bupha-Intr, T., Laosiripisan, J. & Wattanapermpool, J. Moderate intensity of regular exercise improves cardiac {SR} {Ca}²⁺ uptake activity in ovariectomized rats. *J. Appl. Physiol.* **107**, 1105–1112 (2009).
30. Thawornkaiwong, A., Pantharanontaga, J. & Wattanapermpool, J. Hypersensitivity of myofilament response to Ca²⁺ in association with maladaptation of estrogen-deficient heart under diabetes complication. *Am. J. Physiol. Integr. Comp. Physiol.* **292**, R844–R851 (2007).
31. Sivasinprasasn, S. *et al.* Obese-insulin resistance accelerates and aggravates cardiometabolic disorders and cardiac mitochondrial dysfunction in estrogen-deprived female rats. *Age (Omaha)*. **37**, (2015).
32. Pelzer, T. *et al.* Increased mortality and aggravation of heart failure in estrogen receptor- β knockout mice after myocardial infarction. *Circulation* **111**, 1492–1498 (2005).
33. Groban, L. *et al.* Female Heart Health: Is GPER the Missing Link? *Frontiers in Endocrinology* **10**, (2020).
34. Dennis, M. K. *et al.* Identification of a GPER/GPR30 antagonist with improved estrogen receptor counterselectivity. *J. Steroid Biochem. Mol. Biol.* **127**, 358–66 (2011).

35. Deschamps, A. M. & Murphy, E. Activation of a novel estrogen receptor, GPER, is cardioprotective in male and female rats. *Am. J. Physiol. Heart Circ. Physiol.* **297**, H1806-13 (2009).
36. Wang, H. *et al.* G protein-coupled estrogen receptor (GPER) deficiency induces cardiac remodeling through oxidative stress. *Transl. Res.* **199**, 39–51 (2018).
37. Bopassa, J. C., Eghbali, M., Toro, L. & Stefani, E. A novel estrogen receptor GPER inhibits mitochondria permeability transition pore opening and protects the heart against ischemia-reperfusion injury. *Am. J. Physiol. Heart Circ. Physiol.* **298**, H16-23 (2010).
38. Weil, B. R. *et al.* Signaling via GPR30 protects the myocardium from ischemia/reperfusion injury. *Surgery* **148**, 436–443 (2010).
39. Chen, J., Chen, S., Zhang, B. & Liu, J. SIRT3 as a potential therapeutic target for heart failure. *Pharmacological Research* **165**, 105432 (2021).
40. T, L. *et al.* General Control of Amino Acid Synthesis 5-Like 1-Mediated Acetylation of Manganese Superoxide Dismutase Regulates Oxidative Stress in Diabetic Kidney Disease. *Oxid. Med. Cell. Longev.* **2021**, (2021).
41. Mishra, S. K. *et al.* MICOA, a novel metastasis-associated protein 1 (MTA1) interacting protein coactivator, regulates estrogen receptor-alpha transactivation functions. *J. Biol. Chem.* **278**, 19209–19 (2003).
42. Wu, J. *et al.* Variations in Energy Metabolism Precede Alterations in Cardiac Structure and Function in Hypertrophic Preconditioning. *Front. Cardiovasc. Med.* **7**, (2020).
43. Bernardo, B. C. *et al.* Gene delivery of medium chain acyl-coenzyme A

dehydrogenase induces physiological cardiac hypertrophy and protects against pathological remodelling. *Clin. Sci.* **132**, 381–397 (2018).

44. Rennison, J. H. *et al.* Enhanced acyl-CoA dehydrogenase activity is associated with improved mitochondrial and contractile function in heart failure. *Cardiovasc. Res.* **79**, 331–340 (2008).

45. Maher, A. C., Akhtar, M., Vockley, J. & Tarnopolsky, M. A. Women Have Higher Protein Content of β -Oxidation Enzymes in Skeletal Muscle than Men. *PLoS One* **5**, e12025 (2010).

46. Maher, A. C., Akhtar, M. & Tarnopolsky, M. A. Men supplemented with 17β -estradiol have increased β -oxidation capacity in skeletal muscle. *Physiol. Genomics* **42**, 342–347 (2010).

Figure 1: Mitochondrial protein acetylation and GCN5L1 are upregulated in female mice. A. Immunoblotting of mitochondrial lysate fractions from male and female mice demonstrate that females exhibit modestly but significantly higher levels of total protein acetylation. N=10, B. GCN5L1, but not SIRT3, mRNA is significantly increased in female hearts relative to males. C. GCN5L1 protein levels are significantly increased in the myocardium of female mice compared to male mice. N=10, *p<0.05, ***p<0.001 vs. male

Figure 2: Estrogen drives GCN5L1 expression via GPER A. Human tissues from female patients of pre-menopausal (PRE) or post-menopausal (POST) age. B. Human derived AC16 cells incubated with E2 show an increase in GCN5L1. GCN5L1 is not upregulated through canonical ER agonism. GCN5L1 immunoblotting after incubation for 24 hours with E2 and or the ER α /ER β antagonist ICI 182, 780. N=4, * p <0.05, **** p <0.0001 vs. vehicle. D. GCN5L1 is elevated after incubation for 24 hours with E2 and the GPER agonist G-1, and is blocked in the presence of GPER antagonist G-36. N=5-6, * p <0.05, ** p <0.01 vs. vehicle.

Figure 3: Estrogen promotes GCN5L1 translation. A. expression of GCN5L1 mRNA levels determined by qPCR after treatment with E2 or G-1. N=3 **p<0.01 vs. vehicle. B. GCN5L1 levels are significantly elevated by blocking 26S degradation, suggesting that GPER agonism does not increase GCN5L1 via impaired proteolysis. N=4, *p<0.05, ***p<0.001 vs. control. C. GCN5L1 induction is blocked by CHX, a translation inhibitor. N=4 *p<0.05, ***p<0.001 vs. control.

Figure 4: GCN5L1 is required for estrogen-mediated MCAD acetylation and activation A. E2 and G-1 raise MCAD activity, while loss of GCN5L1 blocks this effect. N=8 B. Acetylation of MCAD is significantly reduced in the presence of E2 and G-1 when GCN5L1 is absent. N=4 *p<0.05, **p<0.01 vs. control, †p<0.01 vs. vehicle. C. Schematic representing the hypothesized mechanism of action of estrogen on GCN5L1 in cardiomyocytes