TatA and TatB generate a hydrophobic mismatch that is important for function and assembly of the Tat translocon in *Escherichia coli*

Short title: The hydrophobic mismatch generated by Tat systems

Denise Mehner-Breitfeld¹, Michael T. Ringel¹, Daniel Alexander Tichy^{1,2}, Laura J. Endter², Kai Steffen Stroh², Heinrich Lünsdorf³, Herre Jelger Risselada², and Thomas Brüser^{1*}

¹Institute of Microbiology, Leibniz Universität Hannover, Hannover, Germany

²Institute for Theoretical Physics, Georg August University Göttingen, Göttingen, Germany

³Helmholtz Centre of Infection Research, Braunschweig, Germany

* To whom correspondence should be addressed: brueser@ifmb.uni-hannover.de

Author contributions:

Conceptualization: Thomas Brüser

Formal analysis: Denise Mehner-Breitfeld, Kai Steffen Stroh, Heinrich Lünsdorf

Funding acquisitions: Thomas Brüser

Investigation: Denise Mehner-Breitfeld, Michael T. Ringel, Daniel Alexander Tichy, Heinrich Lünsdorf

Methodology: Denise Mehner-Breitfeld, Michael T. Ringel, Heinrich Lünsdorf, Herre Jelger Risselada

Supervision: Laura J. Endter, Kai Steffen Stroh, Herre Jelger Risselada, Thomas Brüser

Validation: Laura J. Endter, Kai Steffen Stroh

Visualization: Denise Mehner-Breitfeld, Kai Steffen Stroh, Heinrich Lünsdorf, Thomas Brüser

Writing – original draft: Thomas Brüser

Writing – review & editing: Denise Mehner-Breitfeld, Michael T. Ringel, Daniel Alexander Tichy, Laura J.

Endter, Kai Steffen Stroh, Heinrich Lünsdorf, Herre Jelger Risselada, Thomas Brüser

24 Abstract

25   The Tat system has the unique purpose to translocate folded proteins across energy-transducing
 26 membranes. It occurs in bacteria and archaea, as well as in eukaryotic organelles of bacterial origin. In
 27 the bacterial model system *Escherichia coli*, the three components TatA, TatB, and TatC assemble to
 28 functional translocons. TatA and TatB both possess an N-terminal transmembrane helix (TMH) that is
 29 followed by an amphipathic helix (APH). The TMHs of TatA and TatB generate a hydrophobic mismatch
 30 with only 12 consecutive hydrophobic residues that span the membrane. We shortened or extended this
 31 stretch of hydrophobic residues in either TatA, TatB, or both, and analyzed effects on transport
 32 functionality and translocon assembly. The wild type length functioned best but was not an absolute
 33 requirement, as some variation was clearly tolerated. Defects of shortenings or extensions were enhanced
 34 by simultaneous mutations in TatA and TatB, indicating partial compensations of mutations in TatA by
 35 wild type TatB or vice versa. Length variation in TatB destabilized TatBC-containing complexes,
 36 revealing that the 12-residues-length is important for Tat component interactions and translocon
 37 assembly. To also address potential effects on TatA associations, we characterized these by metal tagging
 38 transmission electron microscopy and carried out molecular dynamics simulations. In these simulations,
 39 interacting short TMHs of larger TatA assemblies were thinning the membrane together with laterally
 40 aligned tilted APHs that generated a deep V-shaped groove. The conserved length of 12 hydrophobic
 41 residues may thus not only be important for translocon interactions, but also for a membrane
 42 destabilization during Tat transport. If this is the case, the specific short length could be a compromise
 43 between functionality and proton leakage minimization.

44 Introduction

45 Tat systems serve the purpose to transport folded proteins across energy-transducing membranes in
 46 bacteria, archaea, plastids, and some mitochondria [1–3]. They minimally consist of two components,
 47 TatA and TatC [4], but three-component TatABC systems are very common and found in the model Tat
 48 systems of *Escherichia coli* and plant plastids [5]. TatA and TatB are structurally similar and
 49 evolutionary related [6]. Both are membrane-anchored by a 13-15 residues long N-terminal
 50 transmembrane helix (TMH) that contains only 12 consecutive hydrophobic residues. This helix is
 51 connected to an amphipathic helix (APH) via a short hinge at the cytoplasmic surface of the membrane
 52 [7–10] (Fig 1A). The APH is followed by a negatively charged patch of residues in TatA [11], and more
 53 C-terminal regions are neither conserved nor functionally essential [12]. TatC has six transmembrane
 54 domains with both termini on the cytoplasmic side [13,14]. TatB interacts tightly with TatC, whereas
 55 TatA gradually dissociates during purifications [15–17]. From studies employing cross-linking and
 56 fluorescent protein tagging, it is known that the TatA interaction with TatBC changes during the
 57 translocation cycle, and larger TatA assemblies at TatABC complexes are believed to enable transport
 58 [18–23]. A subpopulation of TatA protomers interacts with TatBC in a way that is not affected by
 59 mutations in TatC that abolish substrate-binding [15], and resting-state-contacts of TatA to TatC have
 60 been identified [24]. TatA assemblies undergo substrate-induced conformational changes that relate to
 61 Tat transport [25,26], and the TatABC components rearrange during the translocation cycle [27,28].

Fig 1: Constructs for the analysis of the 12-residues hydrophobic stretch in TatA and TatB. (A) Sequence logos of the functionally important region of TatA and TatB. Highlighted are the 12-residues hydrophobic stretch in the transmembrane helix of TatA and TatB (yellow), the preceding polar position (bluish purple), and the amphipathic helix (turquoise). The sequence logos have been obtained using enoLOGOS [29] based on sequence alignments by Clustal Omega [30] of sequences from a wide range of organisms, including α -, β -, γ -, δ -, and ϵ -proteobacteria, firmicutes, actinobacteria, cyanobacteria, chlorobi, chloroflexaceae, thermodesulfobacteria, aquificae, and archaea. (B) TatA and TatB variants used in this study. The color code is the same as in (A). Inserted Leu residues are in red. (C) Carbonate wash analysis for the assessment of membrane integration of the TatA and TatB variants used in this study. TatA or TatB variants were detected by SDS-PAGE/Western blotting in the untreated membrane fraction (M), supernatant after carbonate wash (CS), and membranes after carbonate wash (CM). Note that all analyzed TatA and TatB variants are membrane inserted, but TatB variants are more stably membrane inserted than TatA variants, most likely due to their tighter interaction with TatC.

Tat-dependently transported proteins possess an N-terminal signal peptide that contains a highly conserved eponymous twin-arginine motif [31,1]. This motif is recognized by TatBC complexes that catalyze the membrane insertion of the signal peptide [32], whereas TatA is primarily required for the translocation step [18]. TatA shows a high tendency to self-associate [33–36], but high-resolution structures are only available for detergent-solubilized TatA [7,8,10], and the mode of TatA self-interactions in membranes is still unclear. Spin-labeling studies indicated that the transmembrane helices of TatA laterally interact [37], but only circular arrangements of aligned TatA have been suggested [8], and the orientation of the APH has not been included in the latest models [38]. Interestingly, a dimeric structure of detergent-solubilized TatA indicated that also APHs of neighboring TatA protomers laterally interact [10].

TMHs need to span the ca. 3 nm thick hydrocarbon core of membranes, and about 20 hydrophobic residues could in principle do this, but TMHs are in average 24.0 ± 5.6 residues long [39], which is generally due to flexing or tilting of helices in membranes [40]. It is therefore very unusual that TatA and TatB have membrane anchors with TMHs of 13-15 residues that contain only 12 consecutive hydrophobic residues, and that the length of this 12-residues stretch is strictly conserved from archaea to bacteria (Fig 1A). As this short length does not suffice to span a membrane lipid bilayer of normal thickness, the TMHs of TatA and TatB generate a hydrophobic mismatch. It has been found that the TMH of TatA per se can destabilize membranes [25], but potential effects of helix shortenings or extensions have never been systematically analyzed. We therefore changed the hydrophobic mismatch of TatA and TatB by generating helix shortenings and extensions, and investigated effects on Tat functionality and translocon assembly. Results indicate that both components tolerate smaller shortenings or extensions, but the natural 12 consecutive hydrophobic residues appear to function best. We demonstrate that, in case of TatB, this hydrophobic mismatch is important for the stable interaction of

TatB with TatC, and therefore the short helices most likely trigger also the association of TatA with TatBC. To also analyze potential effects on the so far uncharacterized larger TatA assemblies that are present at Tat translocons during the translocation cycle, we visualized such assemblies in whole cells by metal-tagging transmission electron microscopy (METTEM), and carried out MD simulations. As a result of these simulations, we found that the short length of the TMH, together with the APH, has the potential to locally thin and destabilize the membrane. As such an effect could be the basis for membrane permeabilization by Tat systems, which would require a delicate compromise between membrane stability and permeabilization, this might explain the strict conservation of the short length of the TMHs in TatA, TatB, and the TatAB-interacting helices of TatC.

Results

Tat systems tolerate some shortening and extension of the stretch of 12 consecutive hydrophobic residues in the TMHs of TatA and TatB

To analyze the potential role of the short stretch of 12 consecutive hydrophobic residues in the TMHs of TatA and TatB, we shortened the hydrophobic helix by removing 1, 3, or 5 residues in TatA, and 1 or 3 residues in TatB. We also extended the helix by 1, 2, 3, 4, 5, or 6 leucine residues in TatA, and by 3 or 4 leucine residues in TatB (Fig 1B). In the following, the nomenclature of the constructs is facilitated for the reader, with “A+1” standing for the construct with one Leu inserted into the helix of TatA, and “B-3” standing for the construct with three hydrophobic residues deleted in the helix of TatB, as examples. To ensure the correct ratio of the Tat proteins, all genetic constructs were generated with the *tatABC* operon under control of its natural promoter in the Tat complementation vector pABS-*tatABC* [41]. We used the *tat*-deletion strain DADE [42] in combination with these vectors, which permitted functional analyses as well as BN-PAGE detections of the Tat complexes. All TatA and TatB variants were stably produced, and carbonate washes revealed that all of them were membrane-integrated (Fig

108 1C). A significant portion of all TatA variants was released from the membranes, which is already known
 109 for TatA and may be due to the presence of these proteins in destabilized membrane regions [25]. The
 110 amount of TatA that was released by carbonate washes was higher in case of shortened TMHs (A-1, A-
 111 3, and A-5, Fig 1C). As TatA can destabilize membranes, these results may indicate that shorter
 112 membrane anchors enhance the membrane-destabilizing effect of TatA. TatB and its variants were more
 113 stably membrane-integrated, which is expected as TatB tightly associates with TatC. We then analyzed
 114 Tat functionality with the described Tat systems. As Tat-deficient strains become SDS-sensitive, which
 115 relates to cell wall defects due to Tat-requirement for transport of the cell wall amidases AmiA and AmiC
 116 [43], we assessed Tat functionality by monitoring SDS resistance of the respective strains (Fig 2A).

Fig 2: Physiological Tat functionality assays with mutated Tat systems. Complementation of SDS sensitivity (A) and chain formation (B) phenotypes of the Tat deficient strain DADE by Tat systems with indicated single or combined TatA and TatB variants. Controls: ‘no Tat’, empty vector control; ‘wt’, wild type Tat system.

Only the A-5 construct was inactive among the TatA variants with shortenings, and only the A+6 construct was inactive among the extension variants. The A-5 deletion variant showed higher sensitivity towards SDS than the non-complemented *tat* deletion strain, indicating that the A-5 variant was not only non-functional but also had an additional negative physiological effect.

In case of TatB, the B-1 and B-3 deletion-variants were both active in this assay, indicating that shorter hydrophobic helices in principle suffice to establish functional Tat translocons. This was unexpected as TatB tightly interacts with TatC for function, and we therefore expected less tolerance. The Tat system also tolerated the 3-residues extension in TatB, but not anymore the 4-residues extension. As TatA and TatB possibly could have overlapping functions and therefore could complement defects of each other, we also tested combinations of single and triple deletions in TatA and TatB, and triple extensions in TatA and TatB. Notably, the combined deletions were functional, indicating that indeed the deletions did not inactivate the proteins. The combined extension (A+3/B+3) showed a markedly reduced SDS resistance, indicating that some functional overlap had masked the effect of single +3 extensions. Nevertheless, the partial SDS resistance proofed residual activity of A+3 and B+3 variants.

A second way to monitor functionality is the chain formation phenotype, which is similarly based on the absence of AmiA and AmiC in the periplasm of Tat-deficient strains. If these amidases are not transported into the periplasm, the murein is not efficiently hydrolyzed between separating cells, resulting in chains of cells. In full agreement with the SDS-sensitivity measurements, the chain formation phenotype was only observed in those strains that were also SDS-sensitive (Fig 2B). The strain with the A+3/B+3 combination, which had reduced but not abolished SDS resistance, showed no chain formation phenotype, indicating that sufficient amidases were transported for cell separation and confirming the residual activity of the Tat system with the +3 extensions in TatA and TatB. In line with that, the SDS-sensitive B+4 strain showed no obvious chain formation, indicating that transported amidases sufficed for cell separation but not for SDS resistance.

141 To recognize weaker effects on protein transport, we then carried out biochemical analyses of the
 142 transport of the Tat model substrate HiPIP, which is an iron-sulfur cluster containing protein that strictly
 143 requires the Tat system for transport [15]. Subcellular fractions were prepared from exponentially
 144 growing cells and analyzed by SDS-PAGE/Western blotting using antibodies specifically recognizing
 145 HiPIP (Fig 3). We used a vector for low-level constitutive HiPIP production, which results in complete
 146 translocation of HiPIP into the periplasm in case of fully functional Tat systems [44]. In this assay, the
 147 mature periplasmic HiPIP band indicates transport, and any precursor in the cytoplasm indicates reduced
 148 translocation relative to the fully functional Tat system. The wild type Tat system, which served as
 149 positive control, showed a strong band of transported mature HiPIP in the periplasm and no accumulation
 150 of precursor in the cytoplasm. Fractionation controls showed that there was no leakage of cytoplasm into
 151 the periplasmic fraction, and the only faint mature HiPIP band in the cytoplasm indicated little proteolytic
 152 cleavage of the signal peptide or residual periplasm in the cytoplasmic fraction. The Tat deficient empty
 153 vector control strain did not show any transport and instead the bands that are indicative for cytoplasmic
 154 accumulation of HiPIP, which are a strong cytoplasmic precursor band and a band that is due to partial
 155 degradation of the signal peptide. All TatA variants with shortenings showed detectable cytoplasmic
 156 precursor bands, with only very little precursor accumulating in A-1 variants and larger quantities
 157 accumulating in A-3 variants. The A-5 variant was inactive, with no detectable mature protein in the
 158 periplasm. The TatA variants with extensions showed slightly less transport in the A+1 variant and only
 159 very little cytoplasmic precursor in the A+2, A+3, and A+4 variants, but a sudden almost complete block
 160 of transport with the A+5 variant. The A+6 variant was inactive. With the TatB variants, we observed
 161 little accumulation with the B-1 and B-3 variants, indicating only a minor defect, whereas the B+3 variant
 162 strongly accumulated HiPIP in the cytoplasm and the B+4 variant hardly transported HiPIP anymore.
 163 Interestingly, the combinations A-1/B-1, A-3/B-3, and A+3/B+3 all showed a stronger effect than the
 164 single variants, indicating that the defects were additive, or that there was indeed some partial

165 compensation of defects of TatA by TatB or vice versa detectable in this semi-quantitative assay. Such
166 effects were more difficult to detect by the apparently less sensitive amidase-based assays, in which few
167 transported enzymes may often suffice for physiological complementations (compare with the SDS
168 sensitivity assay in Fig 2A).

Fig 3: Biochemical Tat functionality assays with mutated Tat systems. Transport analysis by SDS-PAGE/Western blot detection of the precursor (pre, band at 14 kDa) and mature (mat, band at 11 kDa) forms of the Tat substrate HiPIP in subcellular fractions of cells containing the indicated single TatA or TatB variants or combinations thereof (upper blots). Biotin carboxyl carrier protein (BCCP, band at 22 kDa) was detected as fractionation control (lower blots). Cytoplasmically accumulating precursor is marked by asterisks. Note that any detection of mature protein in the periplasmic fraction indicates functional transport, whereas accumulation of precursor in the cytoplasm is only observed in cases of a partial or complete defect in transport. C, cytoplasm; M, membrane; P, periplasm.

169 Together, the activity measurements indicate that the 12-residues length of the hydrophobic helix
170 in wild type TatA and TatB results in highest transport efficiency. In case of TatA, shortenings by up to
171 three residues and extensions by up to four residues are tolerated, and some residual transport is even
172 detectable with a five-residues extension. TatB similarly tolerates shortenings and truncations of three
173 residues, with only little residual transport detectable in case of the four-residues extended B+4 variant.

Variation in the TatB TMH length, not in the TatA TMH length, interferes with TatBC complex stability

It is well-described that TatB interacts with the short helix 5 of TatC, whereas TatA can interact with the similarly short helix 6 of TatC [38,28]. It was therefore possible that functional effects of length variations in the hydrophobic TMH of TatA or TatB are simply due to disturbed interactions with TatC. To examine this aspect, we analyzed the stability of the known TatC-containing complexes [27].

Membrane fractions were solubilized by digitonin and analyzed by BN-PAGE/Western blotting to detect Tat complexes TC1 and TC2 (Fig 4). While the complexes were unaffected by the length variations in TatA (Fig 4A), the B+3 and B+4 variants caused disassembly of the complexes (Fig 4B). In case of B-1, only the TC2 band was affected and changed to a very diffuse band, whereas the TC1 band remained unaffected. With B-3, the diffuse band of the size of TC2 was the only remaining complex, and TC1 had completely disappeared. Some disassembly of Tat complexes is always observed and has been described previously [15]. Together, these data clearly indicate a severe effect of the TatB variants on TatBC interaction and consequently on TatBC complex stability. As also the functional TatB variants showed this destabilizing effect, sufficient TatBC complexes must have been present in living cells to support Tat transport with this expression system.

Fig 4: Analysis of Tat complexes in strains with mutated Tat systems. (A) BN-PAGE/Western blot detection of Tat complex 1 (TC1) and 2 (TC2) in strains containing Tat systems with indicated TatA variants (A) or with TatB variants alone or in combination with TatA variants (B). Blots were developed with antibodies against TatB (upper blots), or TatC (lower blots). The position without the TC1 band in B-3 Tat systems is highlighted by open circles. B+3 and B+4 extensions have more general destabilizing effects. For direct comparison, the A+3 system is analyzed in parallel with B+3 and the A+3/B+3 combination. Positions of TC1, TC2, and molecular weight markers (440 kDa, 669 kDa) are indicated on the left.

METTEM-analysis of TatA clusters in the cytoplasmic membrane at wild type expression level

In the above shown BN-PAGE analyses, we could only detect effects of TatB variants on the Tat complexes, which is not surprising as most TatA dissociates from the detectable TatBC-containing complexes upon detergent solubilization, and only a regular ladder of multiple homooligomeric TatA associations is detected that does not permit analyses of the TatBC-bound TatA [33,34]. To reveal potential effects of the mutations on TatA, the lack of structural information in principle could be overcome by MD simulations. However, to create a reasonable starting point for MD simulations, we required some information about the organization of the large TatA assemblies at translocation sites. So far, it is only known from thylakoidal cross-linking analyses that at least up to 16 TatA protomers can be cross-linked at active translocation sites over a 4-minutes time span [22], which is evidence for an association of this number of protomers. Another analysis had indirectly calculated a median of 25 interacting TatA-YFP protomers in such complexes, based on mobility and fluorescence intensity measurements [23]. The pixel size of 50 nm in these analyses certainly could not resolve single protomers or even whole TatA assemblies. We thus first aimed to examine the proposed larger TatA associations in the *E. coli* system by a higher-resolution method. To our knowledge, electron microscopy is the only currently available technique that can give sufficient positional resolution within cells. We chose metal-tagging transmission electron microscopy (METTEM) for these analyses, using metallothionein-gold tags [45–50]. Metallothionein-gold tags have the advantage of directly labeling the protein of interest with rather small EM-detectable electron-dense domains. In initial experiments with abundant recombinantly produced Tat systems, we found that three metallothioneins in row (= MT3) permitted clearly visible electron-dense granules of +/- 1 nm in diameter in *E. coli* cells, and the detected particles could be identified as gold by PEELS analysis (Fig 5). Note that the recombinant protein levels that have been used for the PEELS analysis resulted in large clusters of interconnected associations that sometimes

214 deformed the membrane, but this measurement only served to proof that the gold label of the MT3 tags
215 had worked (Fig 5BC), and no other conclusion is drawn here.

Fig 5: PEELS analysis of TatB-MT3-gold.

(A) Survey micrograph of the analyzed cell containing the recombinant TatAB(MT3)C system as expressed by pABS-*tatAB(MT3)C*, with TatB-MT3-gold visible in clusters at the cell pole. (B) Electron spectroscopic element map at the Au-O23 edge Emax at 66 eV, energy-slit width: 8 eV, superimposed in red to the image of the cell pole shown in (A). Energy-settings and corresponding energy-slit width for the jump-ratio acquisition are indicated in C (W1; Emax). (C) PEEL-spectrum of the MT3-gold containing cell pole (inset: measuring area), detecting gold at the Au-O23 edge (red graph) and an energy maximum, which is in accordance to the Au-reference PEELS (green graph) from the EELS-atlas [51].

216 Having the method established, we attempted to localize the Tat components at wild type expression
217 levels in strains that had the modified *tat* operons chromosomally integrated. With TatB-MT3 and TatC-
218 MT3, no gold was detectable at wild type level, but it was possible to detect TatA-MT3 in these
219 experiments, which is likely due to the higher abundance of TatA relative to TatB and TatC in *E. coli*,
220 which are present in an estimated 50:2:1 ratio, respectively [52]. The wild type level of the TatA-MT3-
221 abundance was confirmed by SDS-PAGE/Western blot analysis (Fig 6A). Substrates interact with the
222 cytoplasmic face of TatA [25], and it is known that YFP tags (ca. 26 kDa) at the cytoplasmically located
223 C-terminus of TatA inactivate TatA in the TatABC system of *E. coli* when produced at wild type level

224 [21]. Similarly, we found that our MT3 tag (ca. 21 kDa) rendered TatA almost inactive (Fig 6B), as a
 225 slight activity was likely due to some degradation to mature size (Fig 6A). As there so far do not exist
 226 any protein tags that can be fused to TatA without compromising activity at wild type levels, this result
 227 was not surprising. Importantly, the tag did not prevent TatA-TatA interactions, and the number of TatA
 228 protomers in these associations was comparable with the number concluded from cross-linking of active
 229 non-tagged TatA (see below, 22). Therefore, the effect on activity was most likely due to steric
 230 inhibitions of substrate interactions, as substrates need to pass through the TatA assemblies. In any case,
 231 these analyses can be taken as first direct visualization of wild-type level TatA assemblies in any
 232 organism.

233 TatA-MT3 formed gold assemblies near cell poles of irregular shape when seen in quasi top-view
 234 projection and roughly normal to the electron beam (Fig 6C-H). The mean length of the gold-assemblies
 235 was $20 \text{ nm} \pm 6.0 \text{ nm}$ ($N = 40$, min = 10 nm, max = 37 nm). The width of the assemblies was about 10 nm.
 236 To determine the number of TatA-MT3 protomers in such assemblies, we counted the number of MT3-
 237 gold particles in 35 regions containing MT3-gold, and calculated from this the average number of 17 ± 6
 238 MT3-gold particles per 200 nm^2 , which is the dimension of an average TatA-MT3-assembly.
 239 Interestingly, several of these assemblies visually appeared to be in contact, suggesting that they were
 240 somehow interdependent or connected (Fig 6E-H), which might be due to a clustering with TatBC
 241 complexes. Note that the MT3-gold intensity was variable, most likely due to individual kinetics of gold
 242 aggregate formation within MT3 tags, which may depend on the local environment (see supplemental
 243 Fig S1).

Fig 6: Detection of TatA-MT3-gold at wild type expression level. (A) Detection of TatA by SDS-PAGE/Western blot analysis in subcellular fractions of strains with MT3-tags at TatA, TatB, or TatC as encoded by single copy chromosomally integrated *tatABC*-strep operons. Wild type *E. coli* and the Tat-deficient strain DADE are analyzed for comparison. Positions of molecular weight markers are indicated on the left. C, cytoplasm; M, membrane; P, periplasm. (B) SDS-sensitivity measurements of strains with TatA(MT3)BC, TatABC (wt), or no Tat system. (C) and (D) show overview images of cells with gold-labeled TatA-MT3-assemblies. Areas boxed in the respective overviews are shown below at higher magnification (lower edge = 100 nm). (E) and (G) are original images, and (F) and (H) were topographically 'relief'-filtered to discriminate MT3-gold granules in pseudo-3D within TatA-MT3 assemblies. White 1 nm-dots (encircled in E,F,G,H) have been included as a direct scale for 1nm MT3-gold-granules.

244 Sometimes, TatA-MT3-gold-assemblies appeared in longer linear arrangements in the membrane.
 245 However, tilting of the sample revealed these arrangements to result from superposition of individual
 246 assemblies when they are oriented in parallel to the electron beam within the ultrathin section (Fig 7).

Fig 7: TEM-Tilting analysis of polar TatA assemblies. A seemingly very long TatA association, detectable by TEM of TatA-MT3-gold at a cell pole (middle), has been analyzed in the same cell by tilting the sample from -30° (left) to $+32^\circ$ (right). The tilted cell revealed that individual TatA-MT3 gold assemblies had been transformed into a single longer association due to superprojection of the assemblies when oriented parallel to the electron beam (tilt -2°).

247 In conclusion, TatA-MT3 formed longitudinal assemblies of irregular shape that were usually positioned
 248 near cell poles, the subcellular location of Tat translocons that has been observed in previous
 249 fluorescence-based studies [20,53]. The number of subunits in such assemblies agreed with previous
 250 calculations [23,22]. An often observed close clustering of several of these TatA assemblies suggested
 251 their indirect interaction, possibly mediated by TatBC, which explains the previous detection of a
 252 significant number of larger associations by fluorescence microscopy that was limited to 50 nm pixel
 253 resolution [23].

MD simulation indicates membrane thinning and lipid disorder generation by the short TMHs and aligned APHs of TatA clusters

We then examined such linear TatA assemblies by 1500 ns MD simulations with 22 TatA protomers in the cytoplasmic membrane lipid bilayer, which is in the upper range of the observed TatA assemblies and therefore might reflect functional associations at Tat translocons. The MD simulations were challenged by the fact that, in living cells, larger TatA assemblies require TatBC interactions for their formation [20,21]. As no TatBC complex structure is known, and as we therefore could not include TatBC complexes in the simulations, we attempted to overcome this problem by placing linear arrangements of TatA protomers in close proximity with opposite APH-orientation. We had four reasons to expect a linear arrangement of TatA in the TatA assemblies: (1) Our METTEM analyses had detected longitudinal assemblies (see Fig 6), (2) an NMR structure of a TatA dimer showed laterally aligned APHs, which is only possible in linear arrangements [10], (3) circularly arranged TatA protomers had been reported to collapse within 40 ns in MD simulations, unless the circles were filled with a lipid bilayer that would need to be removed to permit translocation [8], and (4) a linear arrangement would permit regular interactions between TMHs of two oppositely oriented TatA protomers. This approach was successful, as the transmembrane helices readily moved into staggered positions, and the resulting complex remained stable over time in the 1500 ns simulation (Fig 8A). Most APHs laterally contacted neighboring APHs, with some interactions interrupted (Fig 8A), which agrees with the NMR evidence for lateral APH association [10], and these aligned APHs formed an angle of in average 109° with the TMH, which agrees with solid state NMR analyses on *Bacillus subtilis* TatA [54].

Also, the hinge was positioned deep in the membrane, which agrees with accessibility analysis data [25]. The aligned APHs formed a deep V-shaped groove at the membrane surface (Fig 8B, see S1 video). As a result of this constellation, the membrane was thinned to about half its natural thickness at the hinge region. A view on the lipids in this simulation revealed that the lipid head groups were excluded from

278 areas of aligned APHs, and the acyl groups were therefore disordered and partly oriented in parallel to
 279 the APHs. The lipid disorder in the membranes could be quantified employing the lipid tail order
 280 parameter S_{zz} (Fig 8C). The data confirmed that the lipid order was lowered in the environment of the
 281 APHs.

Fig 8: MD simulations with 22 TatA protomers of the wild type and the variants A-5 and A+6. (A) Top view from the cytoplasmic side on the 22 TatA wt simulation. (B) Cross sections through TatA complexes, with the cytoplasm above and the periplasm below the membrane cross sections. TMH, yellow; APH, cyan; Q8, turquoise; hinge region, dark grey; N- and C-termini, grey; lipids, light and dark (head groups) brown. (C) Lipid tail order parameter for lipids in the vicinity of the TMH (associated) and lipids in the bulk phase (non-associated). Bars indicate 95% confidence intervals. The shown coarse grained model of DPPE illustrates tail beads and the orientation of an example tail bond vector (b) relative to the bilayer

normal (n), which is the basis for the order parameter. The other beads in the model are ethanolamine and phosphate (brown), and glycerol (light brown).

282 We then carried out the simulations with the TatA variants A-5, A-3, A-1, A+1, A+2, A+4, and A+6. As
 283 now expected, the shortenings resulted in further thinning of the membrane, whereas the extensions of
 284 the TMH resulted in a thicker membrane, which becomes evident when comparing the simulations A-5,
 285 wt, and A+6 (Fig 8B). Accordingly, the lipid order further decreased with a shorter TMH and increased
 286 with a longer TMH (Fig 8C). The TatA simulations therefore had added a potential new aspect to the
 287 role of the short length of the TMH, as they suggested that this exact length might not only be relevant
 288 for interactions (as shown for TatB; see Fig 4) but also for membrane destabilization, which
 289 mechanistically could be important (see discussion).

290 As the MD simulations had indicated that large TatA assemblies can thin the membrane, we addressed
 291 whether the number of TatA protomers in the association could be relevant for membrane thinning. We
 292 repeated the simulations with 15, 10, and 5 TatA protomers, and found that the membrane-thinning effect
 293 required the lateral association of multiple APHs, which was the reason why only associations of >10
 294 TatA protomers efficiently formed the V-shaped groove in the membrane (Fig 9AB). In case of only 5
 295 protomers, the lateral associations of APHs were too few to impose a V-shaped groove on the membrane
 296 lipids, which is why such small associations had less effect on the membrane bilayer, as quantified by
 297 the tail order parameter (Fig 9C, see S2 video).

Shortening of the TatA TMH can affect cellular growth

The MD simulations suggested a second potential reason for the exact TMH length, which is the compromise between a mechanistically required membrane destabilization and an energetically required membrane stability. It has already been shown several years ago in *in vitro* experiments, that TatA can destabilize the membrane and that substrate-binding to TatA enhances this membrane destabilization,

303 resulting in proton leakage [25]. We thus wondered whether a too short TMH could result in extreme
 304 membrane destabilization *in vivo*, which could represent a selective pressure for maintaining the
 305 minimum of 12 consecutive hydrophobic residues in TatA (and consequently also in TatB and the short
 306 helices 5 and 6 in TatC, which need to assemble with TatA at the translocation site). As proton leakage
 307 might affect growth, we analyzed growth of all strains used in our study (Fig 10).

Fig 10: Growth analysis of strains containing the Tat systems analyzed in this study. (A) A-5, B-3, but also A+6 inhibit cellular growth. For comparison, strains with the wild type Tat system (wt) and without a Tat system (no Tat) are included, as well as the A+5 strain, which does not show a growth phenotype, although A+6 shows a clear growth phenotype. (B) Comparison of the growth phenotype of strains with all Tat systems used in this study, using the OD₆₀₀ at the 3:36 h time point, as this time point is right before the end of the exponential growth of the fastest growing strains. For the strains shown in (A), the same color code is used. Error bars indicate the standard deviation of three cultures. Note that the OD₆₀₀ was recorded by a 96-well multititer plate reader with 5.3 mm initial path length.

308 Growth curves indicated a delay of growth by A-5 and B-3, which may have been caused by cellular
 309 adaptations to stressed membranes (Fig 10). A likely cellular adaptation would be the Psp response, as
 310 Psp system components are known to stabilize membranes [55], and they have been shown to interact
 311 with Tat systems [56,57]. Interestingly, also A+6, not A+5, caused a clear growth phenotype, which may
 312 be due to proton leakage as result of an influence on structures of other membrane proteins.

Discussion

Helices that pass the bacterial cytoplasmic membrane have to traverse a hydrophobic environment of about 3 nm [58]. However, the TMH of TatA is predicted to consist of only 15 residues [10], and that of TatB consists of only 13 residues [9], and in these helices there is only a stretch of 12 consecutive hydrophobic residues. A helix of 12 residues has a length of 1.8 nm, which does not suffice to span a lipid bilayer and therefore causes a hydrophobic mismatch. Such mismatches often have functional implications, such as the regulation of the mechanosensitive channel MscL or the aquaporin GlpF [58,59], and hydrophobic mismatches can drive membrane protein interactions [60]. This study now indicates that hydrophobic mismatches are also important for Tat functionality and Tat component interactions. The length of 12 residues is not a strict requirement for Tat transport but rather an optimal length, as little elongations and extensions of the TMH only reduce but not completely abolish function and interactions (Fig 2 and 3). In case of TatB, our data show that all length variations affect TatBC complex stability (Fig 4), and activity reductions are therefore likely due to weaker TatBC interactions. It is interesting that specific variations of the TMH in TatB can have selective effects on the different complexes, which might relate to conformational equilibria of the Tat complexes, such as previously described for mutations in TatC [27]. Observations of residual activity with extended or shortened hydrophobic stretches in TatB are suggestive for significant flexibility of interactions within TatBC complexes, which likely are not rigid but rather dynamic oligomers. Also in case of TatA, a compromised interaction with TatB and/or TatC could contribute to the observed transport defects. Interestingly, the combination of extensions or shortenings in TatA and TatB showed more severe defects than the mutations in single components, indicating that there exists some detectable functional overlap between TatA and TatB. The TatA TMH can occupy the same TatC binding site as TatB [28], which may account for this functional overlap, and as the TatC interaction is likely important also for TatA, length variation

336 in the TMH of TatA is expected to influence activity of the system, as does the length variation in the
337 TMH of TatB.

338 As BN-PAGE is not helpful to address potential effects on TatA associations, which is due to the multiple
339 associations that TatA forms in the presence of detergent, we sought to model by MD simulations such
340 TatA associations with wild type or mutated membrane anchors. Before doing any MD simulation, we
341 required a more direct evidence for oligomeric TatA assemblies in *E. coli*, and the METTEM analysis
342 served that purpose, showing that TatA-MT3 can indeed form large associations of in average 17 +/- 6
343 protomers, which in turn can cluster to even larger associations, suggesting bridging interactions with
344 other components, such as TatBC. In the end, our METTEM data fully agree with the cross-linking data
345 of the thylakoid system, in which interacting components had been trapped over minutes [22], and they
346 are also in remarkable agreement with the mean number of 25 protomers that have been calculated from
347 fluorescence intensity and mobility measurements of TatA-YFP foci [23]. The latter fluorescence study
348 also reported much larger TatA-YFP associations (at least up to 100 TatA-YFP), which may correspond
349 to the somehow interconnected TatA assemblies in our METTEM analyses. Bridged TatA assemblies
350 likely diffuse together and therefore were not distinguishable from individual TatA assemblies within
351 the 50 nm pixels in these fluorescence analyses [23]. In contrast, METTEM resolved the 1 nm MT3-gold
352 tags, and therefore had a higher resolution than the current limit of ~20 nm in super-resolution
353 fluorescence techniques [61]. To our knowledge, this is the first application for METTEM in localization
354 of membrane protein associations in any organism.

355 Having directly seen and characterized wild type level TatA associations in cells, we first simulated a
356 larger cluster with 22 subunits, which was well in the size range of typical TatA assemblies in METTEM
357 analysis (see supplemental Fig S1) and should be in the range of sizes of TatA associations at TatBC
358 complexes. These simulations readily yielded stable TatA associations, with TMDs interacting with each

other on a staggered position. The assemblies were rather flexible, often showing small kinks due to some irregularities in the arrangements.

Most importantly, the MD simulations revealed that a potential second function of the short TMH could be membrane destabilization. The lipid tail order parameter clearly indicates a correlation of TMH length and membrane destabilization (Fig 8C). Also important is the finding that, in these MD simulations, a larger association of TatAs is required for an efficient membrane destabilization, and that the generation of the V-shaped groove comes along with generation of membrane curvature (Fig 9). The simulations fully agree with published data: The APHs laterally interact, as previously evidenced by NMR [10], the TMHs laterally interact, as suggested by previous spin-labeling EPR studies [37], the hinge domain is positioned deep in the membrane, as supported by in fluorescence quencher accessibility studies [25], and the APHs have a tilted orientation in the membrane, as seen in solid state NMR studies with *Bacillus subtilis* TatA [54]. It is noteworthy that the average angle of the APH relative to the TMH in the simulations is near 109°, which is in good agreement with the angle determined by solid state NMR [54].

Considering the obvious mechanistic tolerance of some length variation (Figs 2 and 3), the strict conservation of the short length of 12 consecutive hydrophobic residues in TatA and TatB from bacteria, archaea, and plastids (Fig 1) is unexpected, especially if only the functionality of protein-protein interactions is assumed to generate the selective pressure for the specific length. However, if membrane destabilization is a second function of these short TMHs, which may be of fundamental importance for the transport mechanism [62], then the hydrophobic mismatch as generated by the strict 12-residue-length could be a delicate compromise between transport efficiency and proton leakage. Our MD simulations clearly show that the short TMHs and aligned tilted APHs of TatA have the potential to destabilize membranes, and the simulations of the TatA assemblies with extended or shortened TMHs show clear effects of such alterations on membrane thickness and lipid order (Fig 8). Proton-tight membranes are of key importance for energy metabolism, and bacteria have (often multiple) protective

383 systems, which compensate for membrane destabilizations. First growth experiments already indicate
384 that adaptation processes are required in response to A-5 and B-3 systems, and effects of minor changes
385 in TMH length on the cellular energetization may be similar but too small to be readily detectable under
386 the experimental conditions (Fig 9). One such adaptive system might be the Psp response system, which
387 is known to stabilize membranes [55] and which already has been shown to interact with TatA [56].
388 In light of the results of this study, we would like to put forward the hypothesis that TatA, TatB and TatC
389 together catalyze the membrane passage of folded proteins across locally destabilized membranes. The
390 “membrane weakening and pulling” hypothesis proposed that this process could depend on membrane
391 destabilization by multiple aligned TatA TMHs [63], which was later supported by an experimental
392 demonstration of membrane destabilization, MD simulations and recently by electrochromic shift
393 measurements [8,25,62]. A conformational switch of the TatA APH in response to substrate binding has
394 been first demonstrated in thylakoids and then in the *E. coli* system, and has been suggested to contribute
395 to this membrane destabilization [25,26].
396 Hydrophobic mismatches are not only generated by the TMHs of TatA and TatB, but also by the short
397 TMHs 5 and 6 of TatC. Notably, the TMHs of TatA and TatB are both shown to interact with these two
398 TatC helices in a resting state [24,38,28], which is likely the reason for the shown importance of the
399 specific short length of the TatB TMH for stable TatBC associations (Fig 4). As substrate binding has
400 been shown to trigger the association of TatA [22,64,20,21], as TatA association has been reported to
401 precede the translocation of TatBC-bound Tat substrates [18,19], as substrate has been shown to trigger
402 membrane permeabilization by TatA *in vitro* [25], and as TatA assemblies have the capacity to destabilize
403 the membrane (Fig 8), we would like to suggest that larger TatA assemblies may form at TatABC
404 complexes to enable Tat transport (see Fig 11). In principle, such an interaction would explain why most
405 TatA is lost during purification of TatBC complexes [15], and this model also explains how a simple
406 conformational shift of TatC helices 5 and 6 in response to substrate binding may enhance the

407 hydrophobic mismatch at the TatBC core to induce the binding of a large number of TatA protomers to
 408 the translocation site, which induces membrane destabilization where it is needed (Fig 11).

Fig 11: Potential substrate-induced assembly of membrane-weakening TatA-clusters at TatBC core complexes. Top view on the cytoplasmic face of the membrane, only positions of TMHs and APHs are schematically depicted. TMHs of TatA, TatB, and TatC are labeled A, B, or C, respectively. In case of TatC, which has 6 TMHs, the helix number is also indicated. The destabilized membrane region is highlighted by a reddish background. Only one docking site for TatA at the oligomeric TatBC core complex is exemplified in the scheme. The substrate that induces the conformational switch is only mentioned but not shown for clarity reasons. The color code for the TMHs and APHs is taken from Fig 1. See text for details.

409 The model also gives an idea about the region of membrane destabilization, which is essentially the
 410 region underneath the APHs. There are two plausible pathways through the destabilized membrane: (1)
 411 The two staggered rows of TatA, which are vis-à-vis positioned in the cluster, could open and permit the
 412 passage of the Tat substrate, or (2) the TMHs remain in the center of the association and the substrate
 413 passes through the thinnest point of the destabilized membrane close to the center of the TatA cluster.
 414 As Tat substrates can be large in diameter, it is likely that both happens: TMHs may lose their contact
 415 and APHs may dissociate to permit the passage of globular proteins. In such a situation of disorder, it
 416 could well be that TatA exchanges with TatB at its TatC binding site, especially when the system is
 417 saturated with transported substrates [28]. It is also possible, that an opening of TatBC complexes can

contribute to the TatA recruitment, such as proposed previously [38,24,2]. Our simulations show that TatA can form stable clusters with two interacting rows of TMHs in the center. From the energetic point of view, MD simulations indicate that there will not be any stable ring shaped TatA assembly with an aqueous hole (8). Instead, the hydrophobic mismatch can be expected to trigger associations of the TMHs, which preferentially should be linear to accommodate aligned APHs at the membrane surface that contribute to membrane destabilization (Figs 8 and 9). During the passage, transient aqueous holes would be formed that would cause uncontrolled proton flux for a very short time-span. Such a proton leakage would agree with the finding that about 80,000 protons pass through the membrane per transported protein [65], albeit proton flux does not energize the transport [66,67]. Although it is clear from these studies that the electric potential is required at certain steps of the pathway, one of which may well be the proposed conformational transition of the TatBC core complex that induces the recruitment of TatA (which is PMF-dependent; 20,21), the driving force for the transport remains to be clarified. In principle, it would be possible that the transmembrane signal peptide insertion as catalyzed by TatBC complexes [32] constitutes this driving force, as the signal peptide remains bound to the TatBC complex during transport of the mature domain [68]. Future experiments surely will shed further light on this highly fascinating transport system.

Materials and Methods

Bacterial strains and growth conditions - The *E. coli* $\Delta tatABCD \Delta tatE$ strain DADE [42], which is a derivative of the strain MC4100 [69], was used for localization and complementation studies. *E. coli* XL1-Blue Mrf^r Kan (Stratagene, La Jolla, California, USA) or *E. coli* DH5 α λpir^+ were used for cloning. Unless otherwise stated, bacteria were grown aerobically at 37°C on LB medium (1% (w/v) tryptone, 1% (w/v) NaCl, 0.5% (w/v) yeast extract) in the presence of the appropriate antibiotics (25 μ g/ml chloramphenicol, 100 μ g/ml ampicillin, 12.5 μ g/ml tetracycline, 15 μ g/ml kanamycin). For subcellular

fractionations, all cultures were normalized to an OD₆₀₀ of 1.0. For growth curves, the OD₆₀₀ was measured in 15-min intervals with cultures grown at 37°C in shaking 96-well plates (culture volume: 200 µl), using the SpectraMax iD3 Microplate Reader (Molecular Devices, LLC., San José, California, USA).

Genetic Methods and Plasmids - For analyses of Tat functionality and for optimization of the MT3-gold-labeling method, the TatABC components were constitutively overproduced using the natural *P_{tatA}* promoter and *tatABC* operon organization in plasmid pABS-*tatABC* and derivatives thereof [41]. The shortenings and extensions of the hydrophobic helices in TatA and TatB were carried out by QuikChange mutagenesis. Primers are listed in Table 1.

Table 1: Primers used in this study¹.

Mutagenesis Primers ^{2,3}	Sequence (5'→3')
Deletions	
<i>tatA</i> -1-F	CAGTATTTGGCAGTTAATTATTGCCGTCATCG
<i>tatA</i> -3-F	GTATTTGGCAGTTAGCCGTCATCGTTG
<i>tatA</i> -5-F	GGTATCAGTATTTGGCAGTTAATCGTTGTACTGCTTTTTGGC
<i>tatB</i> -1-F	CGGTTTTAGCGAACTGTTGGTGTTTCATCATC
<i>tatB</i> -3-F	GTTTTAGCGAACTGTTTCATCATCGGCC
Leucine insertions	
<i>tatA</i> +1-F	CAGTATTTGGCAGTTATTGCTTATTATTGCCGTCATCGTTG
<i>tatA</i> +2-F	CAGTATTTGGCAGTTATTGCTTCTCATTATTGCCGTCATCGTTG
<i>tatA</i> +3-F	GGCAGTTATTGCTTCTCTTAATTATTGCCGTCATCG
<i>tatA</i> +4-F	GGCAGTTATTGCTTCTCTTATTGATTATTGCCGTCATCG
<i>tatA</i> +5-F	GTTATTGCTTCTCTTATTGCTTATTATTGCCGTCATCGTTG
<i>tatA</i> +6-F	GTTATTGCTTCTCTTATTGCTTCTCATTATTGCCGTCATCGTTG
<i>tatB</i> +3-F	CGGTTTTAGCGAACTGCTGCTCCTCTATTGGTGTTTCATCATC
<i>tatB</i> +4-F	CGGTTTTAGCGAACTGCTGCTCCTCTGCTATTGGTGTTTCATCATC
Fusion Primers	
MT-I-F	CGACCGGGATCCAATTCGAGCTCGAACAACAACAAC
MT-R	TTATCGGCGCGCTTGTAACGACGGCCAGTGC
pABS- <i>tatB</i> -BamHI-F	CCCGAGGGATCCGATAAAGAGCAGGTGTAATCC
pABS- <i>tatC</i> -ClaI-R	GGCCGTATCGATCTATTCTTCAGTTTTTTCGCTTTCTGCTTC
pABS- <i>tatA</i> -MT3-R	GTTGTTGTTGTTTCGAGCTCGAATTAGATCCACCTGCTCTTATCGTGGCGCTTCG
pABS- <i>tatABC</i> -Sall-F	AGGGCATCGGTCGACGCTCTCC
pABS- <i>tatB</i> -MT3	GTTGTTGTTGTTTCGAGCTCGAATTAGATCCCGGTTTATCACTCGACGAAGGGGAAG
pABS- <i>tatC</i> -F	CCGAGCGGATCCTCGTCGAGTGATAAACCGTAAAC
Primers for cloning of pAH120-Ptat-<i>tatABC</i>-strep	
<i>tatA</i> -NdeI-F	GAACACATATGGGTGGTATCAGTATTTGGC
<i>tatC</i> -BamHI-R	AAGGATCCTTCTTCAGTTTTTTCGCTTTCTGC

¹Primer sequences are given in 5'-3' direction. ²Corresponding reverse primers covered the same sequence.

³Templates for larger deletions or insertions were the constructs with smaller deletions or insertions.

452 To analyze MT3-gold labeled Tat components at wild type level, the *tatABC* operon and its derivatives
 453 were cloned into pAH120-Ptat-*tatABC*-strep (see primer table), which is a *tatABC*-containing derivative
 454 of pAH120-Ptat-*tatA*-strep- Δ BamHI [70]. Briefly, the MT3-tag (including the linker) was amplified from
 455 the pMAL-c2x-MT3 [50] using primers MT-I-F and MT-R, cut with BglII/BssHII, and ligated into the
 456 backbone of the plasmid pAH120-Ptat-*tatABC*-strep cut with BamHI/BssHII. This resulted in pAH120-
 457 Ptat-*tatABC*-MT3, which was then cut with NdeI/BamHI and ligated into the corresponding sites of
 458 pABS-*tatABC*-H6 [27], resulting in pABS-*tatABC*-MT3. To construct pABS-*tatA*-MT3, *tatA* was
 459 amplified from pABS-*tatABC*-MT3 using primers pABS-*tatA*-MT3-R and pABS-*tatABC*-SalI-F, the
 460 fragment was cut with SalI/SacI and ligated into the plasmid backbone obtained by restriction of pABS-
 461 *tatABC*-H6 with the same enzymes. The same procedure was followed to construct pABS-*tatAB*-MT3,
 462 except that the primers utilized were pABS-*tatB*-MT3-R and pABS-*tatABC*-SalI-F. To append the *tatBC*-
 463 genes to the *tatA*-MT3-construct, these were amplified from the pABS-Ptat-*tatABC*-MT3 plasmid using
 464 primers pABS-*tatB*-BamHI-F and pABS-*tatC*-ClaI-R, the fragment was cut with BamHI/ClaI and ligated
 465 into the respective sites of the pABS-Ptat-*tatA*-MT3 backbone. The same steps were taken to construct
 466 the pABS-*tatAB*-MT3-*tatC* plasmid, only that primers pABS-*tatC*-F and pABS-*tatC*-ClaI-R were used
 467 and the fragment was ligated into the backbone of pABS-Ptat-*tatAB*-MT3. Finally, both the *tatA*-MT3-
 468 BC and *tatAB*-MT3-*tatC* constructs were transferred back into pAH120 by restriction of pAH120-Ptat-
 469 *tatABC*-MT3, pABS-*tatA*-MT3-*tatBC* and pABS-*tatAB*-MT3-*tatC* with NdeI/XbaI, and ligation of the
 470 *tat*-fragments into the pAH120-backbone. Vectors were integrated into the chromosomal lambda
 471 attachment site according to the protocol of Haldiman and Wanner [71]. pRK-*hip* was used for
 472 constitutive low-level expression of the *hip* gene from its own promotor [44].

473 **Biochemical methods** - Subcellular fractionations into periplasm, membranes, and cytoplasm were
 474 performed with 50 mL cultures as described previously [70]. SDS-PAGE analysis was carried out by
 475 standard methods [72]. BN-PAGE was performed as described previously [34], but without β -

mercaptoethanol in the buffers. TatABC complexes were solubilized with 1% digitonin. For immunoblots, proteins were semi-dry blotted on nitrocellulose membranes and blots were developed using antibodies directed against synthetic C-terminal peptides of TatA, TatB, TatC or polyclonal rabbit serum against purified HiPIP, using the ECL system (GE Healthcare, Solingen, Germany) for signal detection. Horseradish peroxidase (HRP)-conjugated goat anti rabbit antibody (Roth, Karlsruhe, Germany) served as secondary antibody. Biotin carboxyl carrier protein (BCCP) was detected by horseradish peroxidase-coupled Strep-Tactin (IBA, Göttingen, Germany). The chain formation phenotype was assessed by phase contrast microscopy. SDS sensitivity was determined by aerobic growth in LB medium containing 4% (w/v) sodium dodecyl sulfate (SDS), using the quotient of the OD₆₀₀ with/without SDS after 3 h of growth [43]. For carbonate washes of membranes, samples were treated as described before [25].

Electron microscopy – Sample preparation - Bacterial cells (20 ml culture, 5 % inoculum) were grown at 37°C at 140 rpm to OD₆₀₀ ~0.5, then 0.4 ml of an aqueous 25 mM tetrachloroauric acid solution were added, and growth was continued for 90 min at 37°C and 120 rpm. Cells were sedimented (10 min, 3,820 x g, 4°C), resuspended in 1.5 ml 100 mM HEPES pH 7.0, 90 mM sucrose (buffer A) and incubated for 5 min at room temperature (RT). Cells were sedimented again (2 min, RT, 6,800 x g) and the incubation/resuspension was repeated. Cells were then resuspended in 1 ml buffer A containing 2.5 % glutaraldehyde, incubated for > 20 min at RT, washed with 1 ml buffer A, sedimented (2 min, 6,177 x g, RT), and washed twice in 100 mM HEPES pH 6.9, 90 mM sucrose, 10 mM MgCl₂, 10 mM CaCl₂, sedimented again and embedded either in SPURR epoxy resin [73] or in LR-White resin [74]. 70 nm ultrathin sections were cut using an ultramicrotome with a diamond knife (Ultracut; Leica Biosystems, Nussloch, Germany) and picked up with neoprene-coated Cu-grids (300 mesh, hexagonal). *TEM image registration and processing* - Unstained samples were analyzed in the elastic bright-field mode with an in-column energy-filter transmission electron microscope (Libra 120 plus, Zeiss, Oberkochen,

Germany). The energy-selecting slit aperture was set to 15 eV energy-width, and images were recorded with a cooled bottom-mount 2 x 2 k CCD (SharpEye, Tröndle, Wiesenmoor, Germany) at nominal magnifications from x 12,500 to x 50,000 near Gaussian or at 500 to 1,500 nm underfocus. Data acquisition and spectrum/image-analysis were done using the iTEM software package (Olympus Soft Imaging Solutions Ltd.). Images were further processed with the Corel Draw Graphics Suite X7 (Corel Ltd.) and ImageJ Fiji software [75]. *Parallel Electron Energy-Loss Spectroscopy (PEELS)* - PEELS was done with unstained 35 nm ultrathin sections of epoxy resin-embedded *E. coli* strain DADE/pABS-*tatAB(MT3)*-C with MT3-gold as the only electron dense intracellular material. PEELS registration was performed in the energy range from 30 to 139 eV, specific for the Au-O₂₃ edge, with a 100 µm spectrum entrance aperture and 5 acquisition cycles per record. Spectrum magnification was set to 200-fold, and the illumination aperture to 0.16 mrad with 4 µA emission current; exposure time: 0.2 s, nominal magnification: x 31,500. *Electron spectroscopic element mapping* Data acquisition was performed on well-defined Au-MT3-clusters and the energy-selective slit was set to 8 eV width. Jump-ratio images were registered at 66 eV (= E_{max}) and 38 eV (= 1st window). These energy-settings were taken from PEEL spectrum data of the MT3-gold assemblies of interest. Exposure time: 2 s, illumination aperture: 0.63 mrad, nominal magnification: x 31,500.

MD simulation - Molecular dynamics simulations were performed using the GROMACS simulation package, version 2019.5 [76]. The MARTINI v2.1 coarse-grained force field [77–79] was applied to simulate lipids, peptides, and solvent. In all simulations, the system was coupled to a constant temperature bath by v-rescale algorithm [80] with a relaxation time of 1.0 ps. Simulations were performed at 320 K. Periodic boundary conditions were applied to simulate bulk behavior. The time step used in the simulation was 20 fs. The dielectric constant in the simulations was $\epsilon_T = 15$. The pressure was coupled to 1 bar employing the Parrinello-Rahman barostat [81] with a relaxation time of 12.0 ps. Unless stated otherwise, the TatA membrane system consisted of 22 TatA proteins in a lipid bilayer with 75 %

phosphatidylethanolamine, 19 % phosphatidylglycerol and 6 % cardiolipin, which is a usual ratio of these lipids in the cytoplasmic membrane of *E. coli* [82]. For the MD simulations of TatA associations, a coarse-grained TatA model up to residue Glu47 was constructed based on the known NMR structure (PDB 2MN7, 10). To model the linear assembly of TatA proteins observed in the TEM experiments, 22 copies of the equilibrated TatA structure were placed vis-à-vis in two rows of 11 subunits, and inserted in the membrane via deletion of all overlapping membrane lipids. The overall dimension of these 22-mers, about 20 nm in length, closely match to the ones observed in METTEM. This system was then equilibrated for 500 ns to enable adaption of the membrane. The TMHs readily associated in a staggered way with all TatA proteins included in the linear complex. For all simulations, the TatA sequences up to position Lys49 were used, as C-terminal truncation analyses have shown that a TatA of exactly that length is still functional [12]. MODELLER 9.24 [83] was used to create the TatA variants. The resulting data were evaluated by MDAnalysis 0.20.1 [84]. The lipid tail order parameter $S_{zz} = \frac{1}{2} (3(\mathbf{b} \cdot \mathbf{n})^2 - 1)$ was determined to quantify the average orientation of tail bond vectors $\mathbf{b}$ with respect to the bilayer normal $\mathbf{n}$ [85]. Order parameter time series were checked for equilibration and decorrelated using the pymbar package [86–88]. Angles between the TMH and the APH were calculated between two vectors connecting the center of masses (COM) of residues Gly21, Thr22, and Lys23 for the hinge either with the COM of residues Ala13, Val14, and Ile15 for the TMH, or with the COM of residues Ile28, Gly29, and Ser30 for the APH (wild type numbering). Images of the simulations were generated using NGLview v2.7.7 [89].

Acknowledgements

We thank Sybille Traupe and Katrin Gunka (LUH, Hannover) for excellent technical assistance. We gratefully acknowledge the skillful work and experience in electron microscopic sample preparation by Inge Kristen, Sabine Schmidt and Ina Schleicher (HZI, Braunschweig).

References

1. Hou B, Brüser T (2011) The Tat-dependent protein translocation pathway. *Biomol Concepts* 2: 507–523. DOI:10.1515/BMC.2011.040.
2. Cline K (2015) Mechanistic Aspects of Folded Protein Transport by the Twin Arginine Translocase (Tat). *J Biol Chem* 290: 16530–16538. DOI:10.1074/jbc.R114.626820.
3. Petru M, Wideman J, Moore K, Alcock F, Palmer T et al. (2018) Evolution of mitochondrial TAT translocases illustrates the loss of bacterial protein transport machines in mitochondria. *BMC Biol* 16: 141. DOI:10.1186/s12915-018-0607-3.
4. Jongbloed JD, Grieger U, Antelmann H, Hecker M, Nijland R et al. (2004) Two minimal Tat translocases in *Bacillus*. *Mol Microbiol* 54: 1319–1325. DOI:10.1111/j.1365-2958.2004.04341.x.
5. Müller M, Klösigen RB (2005) The Tat pathway in bacteria and chloroplasts. *Mol Membr Biol* 22: 113–121. DOI:10.1080/09687860500041809.
6. Yen M-R, Tseng Y-H, Nguyen EH, Wu L-F, Saier MH (2002) Sequence and phylogenetic analyses of the twin-arginine targeting (Tat) protein export system. *Arch Microbiol* 177: 441–450. DOI:10.1007/s00203-002-0408-4.
7. Hu Y, Zhao E, Li H, Xia B, Jin C (2010) Solution NMR Structure of the TatA Component of the Twin-Arginine Protein Transport System from Gram-Positive Bacterium *Bacillus subtilis*. *J Am Chem Soc.* DOI:10.1021/ja1053785.
8. Rodriguez F, Rouse SL, Tait CE, Harmer J, Riso A de et al. (2013) Structural model for the protein-translocating element of the twin-arginine transport system. *Proc Natl Acad Sci U S A* 110: E1092–101. DOI:10.1073/pnas.1219486110.
9. Zhang Y, Wang L, Hu Y, Jin C (2014) Solution structure of the TatB component of the twin-arginine translocation system. *Biochim Biophys Acta* 1838: 1881–1888. DOI:10.1016/j.bbame.2014.03.015.
10. Zhang Y, Hu Y, Li H, Jin C (2014) Structural basis for TatA oligomerization: an NMR study of *Escherichia coli* TatA dimeric structure. *PloS one* 9: e103157. DOI:10.1371/journal.pone.0103157.
11. Warren G, Oates J, Robinson C, Dixon AM (2009) Contributions of the transmembrane domain and a key acidic motif to assembly and function of the TatA complex. *J Mol Biol* 388: 122–132. DOI:10.1016/j.jmb.2009.02.060.
12. Lee PA, Buchanan G, Stanley NR, Berks BC, Palmer T (2002) Truncation analysis of TatA and TatB defines the minimal functional units required for protein translocation. *J Bacteriol* 184: 5871–5879. DOI:10.1128/JB.184.21.5871-5879.2002.
13. Rollauer SE, Tarry MJ, Graham JE, Jaaskelainen M, Jager F et al. (2012) Structure of the TatC core of the twin-arginine protein transport system. *Nature* 492: 210–214. DOI:10.1038/nature11683.
14. Behrendt J, Standar K, Lindenstrauss U, Brüser T (2004) Topological studies on the twin-arginine translocase component TatC. *FEMS Microbiol Lett* 234: 303–308. DOI:10.1016/j.femsle.2004.03.048.
15. Behrendt J, Brüser T (2014) The TatBC complex of the Tat protein translocase in *Escherichia coli* and its transition to the substrate-bound TatABC complex. *Biochemistry* 53: 2344–2354. DOI:10.1021/bi500169s.
16. Bolhuis A, Mathers JE, Thomas JD, Barrett CM, Robinson C (2001) TatB and TatC form a functional and structural unit of the twin-arginine translocase from *Escherichia coli*. *J Biol Chem* 276: 20213–20219. DOI:10.1074/jbc.M100682200.
17. Behrendt J, Lindenstrauss U, Brüser T (2007) The TatBC complex formation suppresses a modular TatB-multimerization in *Escherichia coli*. *FEBS Lett* 581: 4085–4090. DOI:10.1016/j.febslet.2007.07.045.
18. Cline K, Mori H (2001) Thylakoid Δ pH-dependent precursor proteins bind to a cpTatC-Hcf106 complex before Tha4-dependent transport. *J Cell Biol* 154: 719–729. DOI:10.1083/jcb.200105149.
19. Mori H, Cline K (2002) A twin arginine signal peptide and the pH gradient trigger reversible assembly of the thylakoid Δ pH/Tat translocase. *J Cell Biol* 157: 205–210. DOI:10.1083/jcb.200202048.
20. Rose P, Fröbel J, Graumann PL, Müller M (2013) Substrate-dependent assembly of the Tat

- translocase as observed in live *Escherichia coli* cells. PLoS one 8: e69488.
DOI:10.1371/journal.pone.0069488.
21. Alcock F, Baker MAB, Greene NP, Palmer T, Wallace MI et al. (2013) Live cell imaging shows reversible assembly of the TatA component of the twin-arginine protein transport system. Proc Natl Acad Sci U S A 110: E3650-9. DOI:10.1073/pnas.1306738110.
22. Dabney-Smith C, Cline K (2009) Clustering of C-Terminal Stromal Domains of Tha4 Homo-Oligomers during Translocation by the Tat Protein Transport System. Mol Biol Cell. DOI:10.1091/mbc.e08-12-1189.
23. Leake MC, Greene NP, Godun RM, Granjon T, Buchanan G et al. (2008) Variable stoichiometry of the TatA component of the twin-arginine protein transport system observed by in vivo single-molecule imaging. Proc Natl Acad Sci U S A 105: 15376-15381. DOI:10.1073/pnas.0806338105.
24. Aldridge C, Ma X, Gerard F, Cline K (2014) Substrate-gated docking of pore subunit Tha4 in the TatC cavity initiates Tat translocase assembly. J Cell Biol 205: 51-65. DOI:10.1083/jcb.201311057.
25. Hou B, Heidrich ES, Mehner-Breitfeld D, Brüser T (2018) The TatA component of the twin-arginine translocation system locally weakens the cytoplasmic membrane of *Escherichia coli* upon protein substrate binding. J Biol Chem 293: 7592-7605. DOI:10.1074/jbc.RA118.002205.
26. Aldridge C, Storm A, Cline K, Dabney-Smith C (2012) The chloroplast twin arginine transport (Tat) component, Tha4, undergoes conformational changes leading to Tat protein transport. J Biol Chem 287: 34752-34763. DOI:10.1074/jbc.M112.385666.
27. Geise H, Heidrich ES, Nikolin CS, Mehner-Breitfeld D, Brüser T (2019) A Potential Late Stage Intermediate of Twin-Arginine Dependent Protein Translocation in *Escherichia coli*. Front Microbiol 10. DOI:10.3389/fmicb.2019.01482.
28. Habersetzer J, Moore K, Cherry J, Buchanan G, Stansfeld PJ et al. (2017) Substrate-triggered position switching of TatA and TatB during Tat transport in *Escherichia coli*. Open biology 7. DOI:10.1098/rsob.170091.
29. Workman CT, Yin Y, Corcoran DL, Ideker T, Stormo GD et al. (2005) enoLOGOS: a versatile web tool for energy normalized sequence logos. Nucleic acids research 33: W389-92. DOI:10.1093/nar/gki439.
30. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K et al. (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Molecular Systems Biology 7: 539. DOI:10.1038/msb.2011.75.
31. Berks BC (1996) A common export pathway for proteins binding complex redox cofactors. Mol Microbiol 22: 393-404. DOI:10.1046/j.1365-2958.1996.00114.x.
32. Fröbel J, Rose P, Lausberg F, Blümmel A-S, Freudl R et al. (2012) Transmembrane insertion of twin-arginine signal peptides is driven by TatC and regulated by TatB. Nat Commun 3: 1311. DOI:10.1038/ncomms2308.
33. Oates J, Barrett CML, Barnett JP, Byrne KG, Bolhuis A et al. (2005) The *Escherichia coli* twin-arginine translocation apparatus incorporates a distinct form of TatABC complex, spectrum of modular TatA complexes and minor TatAB complex. J. Mol. Biol. 346: 295-305. DOI:10.1016/j.jmb.2004.11.047.
34. Richter S, Brüser T (2005) Targeting of unfolded PhoA to the TAT translocon of *Escherichia coli*. J Biol Chem 280: 42723-42730. DOI:10.1074/jbc.M509570200.
35. Gohlke U, Pullan L, McDevitt CA, Porcelli I, Leeuw E de et al. (2005) The TatA component of the twin-arginine protein transport system forms channel complexes of variable diameter. Proc Natl Acad Sci U S A 102: 10482-10486. DOI:10.1073/pnas.0503558102.
36. Pettersson P, Patrick J, Jakob M, Jacobs M, Klösigen RB et al. (2021) Soluble TatA forms oligomers that interact with membranes: Structure and insertion studies of a versatile protein transporter. Biochim Biophys Acta 1863: 183529. DOI:10.1016/j.bbamem.2020.183529.
37. White GF, Schermann SM, Bradley J, Roberts A, Greene NP et al. (2010) Subunit organization in the TatA complex of the twin arginine protein translocase. a site-directed EPR spin labeling study. J Biol Chem 285: 2294-2301. DOI:10.1074/jbc.M109.065458.
38. Alcock F, Stansfeld PJ, Basit H, Habersetzer J, Baker MA et al. (2016) Assembling the Tat protein translocase. Elife 5. DOI:10.7554/eLife.20718.
39. Baeza-Delgado C, Marti-Renom MA, Mingarro I (2013) Structure-based statistical analysis of transmembrane helices. Eur Biophys J 42: 199-207. DOI:10.1007/s00249-012-0813-9.
40. Holt A, Killian JA (2010) Orientation and dynamics of transmembrane peptides: the power of

- 658 simple models. *Eur Biophys J* 39: 609–621. DOI:10.1007/s00249-009-0567-1.
- 659 41. Berthelmann F, Brüser T (2004) Localization of the Tat translocon components in *Escherichia coli*.
- 660 *FEBS Lett* 569: 82–88. DOI:10.1016/j.febslet.2004.05.054.
- 661 42. Wexler M, Sargent F, Jack RL, Stanley NR, Bogsch EG et al. (2000) TatD is a cytoplasmic protein
- 662 with DNase activity. No requirement for TatD family proteins in sec-independent protein export. *J*
- 663 *Biol Chem* 275: 16717–16722. DOI:10.1074/jbc.M000800200.
- 664 43. Ize B, Stanley NR, Buchanan G, Palmer T (2003) Role of the *Escherichia coli* Tat pathway in outer
- 665 membrane integrity. *Mol Microbiol* 48: 1183–1193. DOI:10.1046/j.1365-2958.2003.03504.x.
- 666 44. Brüser T, Yano T, Brune DC, Daldal F (2003) Membrane targeting of a folded and cofactor-
- 667 containing protein. *Eur J Biochem* 270: 1211–1221. DOI:10.1046/j.1432-1033.2003.03481.x.
- 668 45. Mercogliano CP, DeRosier DJ (2006) Gold nanocluster formation using metallothionein: mass
- 669 spectrometry and electron microscopy. *J. Mol. Biol.* 355: 211–223.
- 670 DOI:10.1016/j.jmb.2005.10.026.
- 671 46. Bouchet-Marquis C, Pagratis M, Kirmse R, Hoenger A (2012) Metallothionein as a clonable high-
- 672 density marker for cryo-electron microscopy. *J. Struct. Biol.* 177: 119–127.
- 673 DOI:10.1016/j.jsb.2011.10.007.
- 674 47. Nishino Y, Yasunaga T, Miyazawa A (2007) A genetically encoded metallothionein tag enabling
- 675 efficient protein detection by electron microscopy. *J. Electron. Microsc.* (Tokyo) 56: 93–101.
- 676 DOI:10.1093/jmicro/dfm008.
- 677 48. Morphew MK, O'Toole ET, Page CL, Pagratis M, Meehl J et al. (2015) Metallothionein as a clonable
- 678 tag for protein localization by electron microscopy of cells. *J. Microsc.* 260: 20–29.
- 679 DOI:10.1111/jmi.12262.
- 680 49. Mercogliano CP, DeRosier DJ (2007) Concatenated metallothionein as a clonable gold label for
- 681 electron microscopy. *J. Struct. Biol.* 160: 70–82. DOI:10.1016/j.jsb.2007.06.010.
- 682 50. Diestra E, Fontana J, Guichard P, Marco S, Risco C (2009) Visualization of proteins in intact cells
- 683 with a clonable tag for electron microscopy. *J. Struct. Biol.* 165: 157–168.
- 684 DOI:10.1016/j.jsb.2008.11.009.
- 685 51. Reimer L, Zepke U, Moesch J, Schulze-Hillert S, Ross-Messemer M et al. (1992) EELSpectroscopy. A
- 686 Reference Handbook of Standard Data for Identification and Interpretation of Electron Energy
- 687 Loss Spectra and for Generation of Electron Spectroscopic Images.: Carl Zeiss, Oberkochen,
- 688 Germany.
- 689 52. Jack RL, Sargent F, Berks BC, Sawers G, Palmer T (2001) Constitutive expression of *Escherichia*
- 690 *coli* *tat* genes indicates an important role for the twin-arginine translocase during aerobic and
- 691 anaerobic growth. *J Bacteriol* 183: 1801–1804. DOI:10.1128/JB.183.5.1801-1804.2001.
- 692 53. Berthelmann F, Mehner D, Richter S, Lindenstrauss U, Lünsdorf H et al. (2008) Recombinant
- 693 expression of *tatABC* and *tatAC* results in the formation of interacting cytoplasmic TatA tubes in
- 694 *Escherichia coli*. *J Biol Chem* 283: 25281–25289. DOI:10.1074/jbc.M707757200.
- 695 54. Walther TH, Grage SL, Roth N, Ulrich AS (2010) Membrane alignment of the pore-forming
- 696 component TatA(d) of the twin-arginine translocase from *Bacillus subtilis* resolved by solid-state
- 697 NMR spectroscopy. *J Am Chem Soc* 132: 15945–15956. DOI:10.1021/ja106963s.
- 698 55. Kobayashi R, Suzuki T, Yoshida M (2007) *Escherichia coli* phage-shock protein A (PspA) binds to
- 699 membrane phospholipids and repairs proton leakage of the damaged membranes. *Mol Microbiol*
- 700 66: 100–109. DOI:10.1111/j.1365-2958.2007.05893.x.
- 701 56. Mehner D, Osadnik H, Lünsdorf H, Brüser T (2012) The Tat system for membrane translocation of
- 702 folded proteins recruits the membrane-stabilizing Psp machinery in *Escherichia coli*. *J Biol Chem*
- 703 287: 27834–27842. DOI:10.1074/jbc.M112.374983.
- 704 57. Bernal-Cabas M, Miethke M, Antelo-Varela M, Aguilar Suárez R, Neef J et al. (2020) Functional
- 705 association of the stress-responsive LiaH protein and the minimal TatAyCy protein translocase in
- 706 *Bacillus subtilis*. *Biochim Biophys Acta Mol Cell Res* 1867: 118719.
- 707 DOI:10.1016/j.bbamcr.2020.118719.
- 708 58. Katsuta H, Sawada Y, Sokabe M (2019) Biophysical Mechanisms of Membrane-Thickness-
- 709 Dependent MscL Gating: An All-Atom Molecular Dynamics Study. *Langmuir* 35: 7432–7442.
- 710 DOI:10.1021/acs.langmuir.8b02074.
- 711 59. Jensen MØ, Mouritsen OG (2004) Lipids do influence protein function-the hydrophobic matching
- 712 hypothesis revisited. *Biochim Biophys Acta* 1666: 205–226.
- 713 DOI:10.1016/j.bbamem.2004.06.009.
- 714 60. Windisch D, Ziegler C, Grage SL, Bürck J, Zeitler M et al. (2015) Hydrophobic Mismatch Drives the

- Interaction of E5 with the Transmembrane Segment of PDGF Receptor. *Biophys J* 109: 737–749. DOI:10.1016/j.bpj.2015.07.022.
61. Garlick E, Thomas SG, Owen DM (2021) Super-Resolution Imaging Approaches for Quantifying F-Actin in Immune Cells. *Front. Cell Dev. Biol.* 9: 676066. DOI:10.3389/fcell.2021.676066.
62. Asher AH, Theg SM (2021) Electrochromic shift supports the membrane destabilization model of Tat-mediated transport and shows ion leakage during Sec transport. *Proc Natl Acad Sci USA* 118: e2018122118. DOI:10.1073/pnas.2018122118.
63. Brüser T, Sanders C (2003) An alternative model of the twin arginine translocation system. *Microbiol Res* 158: 7–17. DOI:10.1078/0944-5013-00176.
64. Dabney-Smith C, Mori H, Cline K (2006) Oligomers of Tha4 organize at the thylakoid Tat translocase during protein transport. *J Biol Chem* 281: 5476–5483. DOI:10.1074/jbc.M512453200.
65. Alder NN, Theg SM (2003) Energetics of protein transport across biological membranes. a study of the thylakoid Δ pH-dependent/cpTat pathway. *Cell* 112: 231–242. DOI:10.1016/s0092-8674(03)00032-1.
66. Braun NA, Davis AW, Theg SM (2007) The chloroplast Tat pathway utilizes the transmembrane electric potential as an energy source. *Biophys J* 93: 1993–1998. DOI:10.1529/biophysj.106.098731.
67. Bageshwar UK, Musser SM (2007) Two electrical potential-dependent steps are required for transport by the *Escherichia coli* Tat machinery. *J Cell Biol* 179: 87–99. DOI:10.1083/jcb.200702082.
68. Gerard F, Cline K (2006) Efficient twin arginine translocation (Tat) Pathway transport of a precursor protein covalently anchored to its initial cpTatC binding site. *J Biol Chem* 281: 6130–6135. DOI:10.1074/jbc.M512733200.
69. Casadaban MJ (1976) Transposition and fusion of the lac genes to selected promoters in *Escherichia coli* using bacteriophage lambda and Mu. *J Mol Biol* 104: 541–555. DOI:10.1016/0022-2836(76)90119-4.
70. Taubert J, Hou B, Risselada HJ, Mehner D, Lünsdorf H et al. (2015) TatBC-independent TatA/Tat substrate interactions contribute to transport efficiency. *PloS one* 10: e0119761. DOI:10.1371/journal.pone.0119761.
71. Haldimann A, Wanner BL (2001) Conditional-replication, integration, excision, and retrieval plasmid-host systems for gene structure-function studies of bacteria. *J. Bacteriol.* 183: 6384–6393. DOI:10.1128/JB.183.21.6384-6393.2001.
72. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680–685. DOI:10.1038/227680a0.
73. Lünsdorf H, Strömpl C, Osborn AM, Bennisar A, Moore ERB et al. (2001) [26] Approach to analyze interactions of microorganisms, hydrophobic substrates, and soil colloids leading to formation of composite biofilms, and to study initial events in microbiogeological processes. *Methods Enzymol* 336: 317–IN14. DOI:10.1016/s0076-6879(01)36598-9.
74. Fernández de Castro I, Fernández JJ, Barajas D, Nagy PD, Risco C (2017) Three-dimensional imaging of the intracellular assembly of a functional viral RNA replicase complex. *J Cell Sci* 130: 260–268. DOI:10.1242/jcs.181586.
75. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M et al. (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9: 676–682. DOI:10.1038/nmeth.2019.
76. Lindahl, Abraham, Hess, van der Spoel (2019) GROMACS 2019.5 Source code: Zenodo.
77. Marrink SJ, Vries AH de, Mark AE (2004) Coarse Grained Model for Semiquantitative Lipid Simulations. *J. Phys. Chem. B* 108: 750–760. DOI:10.1021/jp036508g.
78. Marrink SJ, Risselada HJ, Yefimov S, Tieleman DP, Vries AH de (2007) The MARTINI force field: coarse grained model for biomolecular simulations. *J. Phys. Chem. B* 111: 7812–7824. DOI:10.1021/jp071097f.
79. Monticelli L, Kandasamy SK, Periole X, Larson RG, Tieleman DP et al. (2008) The MARTINI Coarse-Grained Force Field: Extension to Proteins. *J Chem Theory Comput* 4: 819–834. DOI:10.1021/ct700324x.
80. Bussi G, Donadio D, Parrinello M (2007) Canonical sampling through velocity rescaling. *J Chem Phys* 126: 14101. DOI:10.1063/1.2408420.
81. Parrinello M, Rahman A (1981) Polymorphic transitions in single crystals: A new molecular dynamics method. *J Appl Phys* 52: 7182–7190. DOI:10.1063/1.328693.

- 772 82. Raetz CR (1978) Enzymology, genetics, and regulation of membrane phospholipid synthesis in
773 *Escherichia coli*. Microbiol Rev 42: 614–659. DOI:10.1128/mr.42.3.614-659.1978.
- 774 83. Webb B, Sali A (2016) Comparative Protein Structure Modeling Using MODELLER. Curr Protoc
775 Protein Sci 86: 2.9.1-2.9.37. DOI:10.1002/cpp.20.
- 776 84. Michaud-Agrawal N, Denning EJ, Woolf TB, Beckstein O (2011) MDAAnalysis: a toolkit for the
777 analysis of molecular dynamics simulations. J. Comput. Chem. 32: 2319–2327.
778 DOI:10.1002/jcc.21787.
- 779 85. MacDermaid CM, Kashyap HK, DeVane RH, Shinoda W, Klauda JB et al. (2015) Molecular dynamics
780 simulations of cholesterol-rich membranes using a coarse-grained force field for cyclic alkanes. J
781 Chem Phys 143: 243144. DOI:10.1063/1.4937153.
- 782 86. Chodera JD, Swope WC, Pitera JW, Seok C, Dill KA (2007) Use of the Weighted Histogram Analysis
783 Method for the Analysis of Simulated and Parallel Tempering Simulations. J Chem Theory Comput
784 3: 26–41. DOI:10.1021/ct0502864.
- 785 87. Shirts MR, Chodera JD (2008) Statistically optimal analysis of samples from multiple equilibrium
786 states. J Chem Phys 129: 124105. DOI:10.1063/1.2978177.
- 787 88. Chodera JD (2016) A Simple Method for Automated Equilibration Detection in Molecular
788 Simulations. J Chem Theory Comput 12: 1799–1805. DOI:10.1021/acs.jctc.5b00784.
- 789 89. Nguyen H, Case DA, Rose AS (2018) NGLview-interactive molecular graphics for Jupyter
790 notebooks. Bioinformatics (Oxford, England) 34: 1241–1242.
791 DOI:10.1093/bioinformatics/btx789.
792

793 Supporting information

794 **S1 Fig. Detailed qualitative aspects of TatA-MT3-gold labeled assemblies in *E. coli* at wildtype level.** (A)
795 Survey view of an individual cell with a TatA-MT3 cluster, labeled with electron-dense gold (squared frame) in
796 the subpolar region. (B) Detailed view of the area framed in (A), showing the TatA-MT3 assemblies of the cluster
797 at higher magnification. Note the electron-dense granules of individual TatA-MT3-gold tags, which are about 1 –
798 1.5 nm in size. An isoscale scheme of the MD-simulated TatA assembly of 22 protomers (arrow, see Fig 8A) is
799 included for comparison. Note that the MT3 domains are 69 residues apart from the modeled APHs (green). (C) A
800 tilted, 3D-surface plot shows the range of gray level variance above the cytoplasmic background within the
801 assemblies, which is indicative for MT3-gold. High peaks with black caps represent gold granules, as they can be
802 readily recognized in (B). The range of different peak heights from high to low as seen in (C) indicates a variability
803 of gold-loading of MT3 tags during the incubation of the cells with tetrachloroauric acid, possibly caused by
804 differences in the local environment and unequal accessibility of gold.

805 **S1 Video. Cross-section video of the 1.500 ns simulation of a TatA-assembly consisting of 25**
 806 **protomers.** Note the V-shaped groove formed by the association of the short TMHs in conjunction with
 807 the aligned APHs.

808

809 **S2 Video. Cross-section video of the 1.500 ns simulation of a TatA-assembly consisting of 5**
 810 **protomers.** Note that 5 TatA protomers do not suffice to significantly thin the membrane.