Local accessory gene sharing drives lineage-specific acquisition of antimicrobial resistance in Egyptian *Campylobacter* spp.

Shaimaa F. Mouftah^{1*}, Ben Pascoe^{2,3*}, Jessica K. Calland², Evangelos Mourkas², Naomi Tonkin², Charlotte Lefèvre^{2§}, Danielle Deuker^{2§§}, Sunny Smith², Harry Wickenden², Matthew D. Hitchings⁴, Samuel K. Sheppard^{2,5#}, Mohamed Elhadidy^{1,6#}

¹Biomedical Sciences Program, University of Science and Technology, Zewail City of Science and Technology, Giza, Egypt; ²The Milner Centre of Evolution, University of Bath, Claverton Down, Bath, UK; ³Chiang Mai University, Chiang Mai, Thailand; ⁴Swansea University Medical School, Swansea University, Swansea, UK; ⁵Department of Zoology, University of Oxford, Oxford, UK; ⁶Department of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt

#Authors to which correspondence should be addressed: s.k.sheppard@bath.ac.uk; melhadidy@zewailcity.edu.eg

*These authors contributed equally.

§Present address: Division of Virology, Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, United Kingdom.

§§Present address: Nuffield Department of Medicine, The Jenner Institute, University of Oxford, Oxford, United Kingdom.

Keywords: *Campylobacter* / Population structure / Accessory genome / Antimicrobial resistance

Repositories: Short read data generated in this study are available on the NCBI Sequence Read Archive, associated with BioProject PRJNA576513 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA576513>).

27 Abstract

28 *Campylobacter* is the most common cause of bacterial gastroenteritis worldwide and diarrheal disease
 29 is a major cause of child morbidity, growth faltering and mortality in low- and middle-income
 30 countries (LMICs). Despite evidence of high incidence and differences in disease epidemiology, there
 31 is limited genomic data from studies in developing countries. In this study, we characterised the
 32 genetic diversity and accessory genome content of a collection of *Campylobacter* isolates from Cairo,
 33 Egypt. In total, 112 *Campylobacter* isolates were collected from broiler carcasses (n=31), milk and
 34 dairy products (n=24) and patients (n=57) suffering from gastroenteritis. Among the most common
 35 sequence types (STs) we identified were the globally disseminated, host generalist ST-21 clonal
 36 complex (CC21) and the poultry specialist CC206, CC464 and CC48. Notably, CC45 and the cattle-
 37 specialist CC42 were under-represented with a total absence of CC61. Comparative genomics were
 38 used to quantify core and accessory genome sharing among isolates from the same country compared
 39 to sharing between countries. Lineage-specific accessory genome sharing was significantly higher
 40 among isolates from the same country, particularly CC21 which demonstrated greater local
 41 geographical clustering. In contrast, no geographic clustering was noted in either the core or accessory
 42 genomes of the CC828, suggesting a highly admixed population. A greater proportion of *C. coli*
 43 isolates were multidrug resistant (MDR) compared to *C. jejuni*. This is a significant public health
 44 concern as MDR food chain pathogens are difficult to treat and often pose increased mortality risk
 45 demanding enhanced prevention strategies in the Egyptian market to combat such a threat.

46

47 Impact statement

48 *Campylobacter* is the leading bacterial cause of gastroenteritis worldwide and despite high incidence
 49 in low- and middle-income countries, where infection can be fatal, culture-based isolation is rare and
 50 the genotypes responsible for disease are seldom identified. Here, we sequenced the genomes of a
 51 collection of isolates from clinical cases and potential infection reservoirs from Cairo in Egypt and
 52 characterised their genetic diversity. Among the most common genotypes we identified were globally
 53 disseminated lineages implicated in human disease worldwide, including the host generalist ST-21
 54 clonal complex (CC21) and the poultry specialist genotypes CC206, CC464 and CC48. Notably
 55 however, some other globally common genotypes were under-represented or entirely absent from our
 56 collection, including those from cattle-specialist lineages, CC42 and CC61. By focussing on specific
 57 lineages, we demonstrate that there is increased accessory genome sharing in specific clonal
 58 complexes. This increased local sharing of genes may have contributed to a greater proportion of *C.*
 59 *coli* isolates possessing antimicrobial resistance determinants that suggest they could be multidrug
 60 resistant (MDR). This is a significant public health concern as MDR food chain pathogens are
 61 difficult to treat and often pose increased mortality risk demanding enhanced prevention strategies.

62

63 Data summary

64 Short read data are available on the NCBI Sequence Read Archive, associated with BioProject
 65 PRJNA576513 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA576513>). Assembled genomes,
 66 supplementary material and additional analysis files are available from FigShare:
 67 <https://doi.org/10.6084/m9.figshare.9956597>. Phylogenetic trees can be visualised and manipulated
 68 on Microreact for *C. jejuni* (https://next.microreact.org/project/Cjejuni_Egypt) and *C. coli*
 69 (https://next.microreact.org/project/Ccoli_Egypt) separately, or combined Cairo and Oxford data with
 70 additional PopPunk network clustering (<https://microreact.org/project/Campy-Egypt>).

71

72 Introduction

73 Diarrheal disease is a major cause of child morbidity, growth faltering and mortality in low- and
 74 middle-income countries (LMICs) (McCormick and Lang, 2016; Platts-Mills and Kosek, 2014).
 75 *Campylobacter* is the most common cause of bacterial gastroenteritis worldwide (Kaakoush et al.,
 76 2015) and typically human campylobacteriosis is commonly diagnosed as a disease associated with
 77 consumption of contaminated food, especially poultry (Nichols et al., 2012; Sheppard et al., 2009).
 78 Extremely high incidence in LMICs, high exposure rates (Lee et al., 2013) and endemism among
 79 young children suggests a different epidemiology (Kaakoush et al., 2015; Lanata et al., 2013; J. Liu et
 80 al., 2016). Frequent or chronic (re)infection is allied to significant morbidity, cognitive development
 81 impairment, and even death (Coker, 2002; Crofts et al., 2018; Kirk et al., 2018; Reed et al., 1996). In
 82 Egypt, campylobacteriosis is common and a leading cause of paediatric diarrhoea, with an incidence
 83 of 1.2 episodes per year (ElGendy et al., 2018; Rao, 2001) with up to 85% of children infected in their
 84 first year (Liu et al., 2012). Despite the high frequency of reported cases of *Campylobacter*-associated
 85 diarrhoea in Egypt (ElGendy et al., 2018), there are no detailed surveillance studies on the dominant
 86 sequence types and proliferation of genotypes associated with the onset of post-infectious sequelae,
 87 such as irritable bowel syndrome (PI-IBS), Guillain-Barré syndrome (GBS) or Miller
 88 Fisher syndrome (Wierzba et al., 2008).

89
 90 *Campylobacter* species are often part of the gut microbiota of various wild and farmed animals
 91 leading to frequent contamination of human food products (Asuming-Bediako et al., 2019; Waite and
 92 Taylor, 2015). In Egypt, farming practices can lack adequate biosecurity and regulation. Only limited
 93 studies have reported the prevalence and distribution of *Campylobacter* in Egyptian
 94 campylobacteriosis cases (Kaakoush et al., 2015) and little is known of the dominant source reservoirs
 95 driving infection and transmission. In Europe, potential source reservoirs have been identified through
 96 source attribution studies, with poultry products regarded as the primary source of infection (Facciola
 97 et al., 2017; Mossong et al., 2016; Sheppard et al., 2009; Thépault et al., 2018). Host-adaptation of
 98 *Campylobacter* to a wide-range of hosts is reflected in its population structure (Colles and Maiden,

2012; Dearlove et al., 2016; Griekspoor et al., 2013; Méric et al., 2018; Sheppard et al., 2014), with many lineages common in human infection able to infect multiple host species. These host generalist lineages include *C. jejuni* ST-21, ST-45 clonal complexes and the *C. coli* ST-828 complex (Dearlove et al., 2016; Mossong et al., 2016). Other genotypes are only found in a single reservoir species, often associated with global poultry or cattle production. Host specialist clonal complexes common in human disease includes the poultry-associated ST-353, ST354 and ST257 (Berthenet et al., 2019; Sheppard et al., 2009) and cattle specialist ST-61 (French et al., 2005; Mourkas et al., 2019).

Human infection in developed countries is usually sporadic and self-limiting, not requiring treatment with antibiotics. However global rates of antimicrobial resistance are rising (Mourkas et al., 2019; Zhao et al., 2016) in line with other Gram negative gastrointestinal pathogens (Tam et al., 2012; CDC, 2020). Widespread agricultural usage has driven the proliferation of tetracycline resistance through its use as a growth promoter (Abdi Hachesoo et al., 2014; Inglis et al., 2019). In particular, *C. coli* has shown an ability to acquire erythromycin resistance genes from other species (Mourkas et al., 2019). This has not been explored for Egyptian *Campylobacter* isolates, where agricultural antibiotic usage is poorly regulated (Dahshan et al., 2015) and self-medication for gastrointestinal disease is common (Abd El-Tawab et al., 2018; Sabry et al., 2014). Global differences in the use of quinolones is likely responsible for the geographical differences observed in quinolone resistance (Luangtongkum et al., 2009; Pascoe et al., 2017; Zollner-Schwetz and Krause, 2015).

We have sequenced 112 *Campylobacter* isolates collected from patients and food of animal source (i.e., broiler chicken carcasses and dairy products) in Cairo over a year to determine the most prevalent *Campylobacter* genotypes causing disease in Egypt. By screening the genome content, including known AMR determinants we provide a better understanding of the local population structure to guide disease intervention in Egypt. This study provides a basis for considering complex transmission networks in LMICs and highlights the role of globally transmitted *Campylobacter* lineages and the emergence of (horizontally acquired) antimicrobial resistance.

Methods

Ethical approval

The study represents a retrospective study that involved sequencing the genomes of a historical strain collection and no patient data collection was involved in this study. Ethical approval was granted from the respective ethics committee in the Egyptian central directorate of research and health development before conducting the study.

Isolate collection

In total, 112 *Campylobacter* isolates were collected in Cairo, Egypt from September 2017 to December 2018, including 31 isolates from broiler carcasses, 24 isolates from milk and dairy products, and 57 clinical isolates. Clinical isolates were recovered from stool samples of patients admitted to hospitals in downtown Cairo suffering from gastroenteritis symptoms. A questionnaire was distributed to all admitted patients requesting details on clinical presentation (e.g., duration of illness, symptoms, medication prescribed), dietary record of the previous 2 weeks, including consumption of specific or undercooked meats, unpasteurized milk, exposure to animal manure or faeces, and any retail outlets commonly used by patients for food consumption prior to the onset of illness. A random sampling approach was then used to include food samples from stores in the study region that were commonly listed in the questionnaire.

Sample culturing and whole genome sequencing

The isolation and enumeration of *Campylobacter* strains from different food matrices was performed according to the ISO 10272-1 (Enrichment Method; Detection of *Campylobacter* spp. after Selective Enrichment). All isolates were sub-cultured from -80°C frozen stocks onto Mueller-Hinton agar (Oxoid, United Kingdom). Plates were incubated at $42 \pm 1^{\circ}\text{C}$ under anaerobic conditions using AnaeroGen™ 2.5L Sachets (Oxoid, United Kingdom). Genomic DNA was extracted from 112 Egyptian isolates using the QIAamp DNA Mini Kit (QIAGEN, Crawley, UK), according to manufacturer's instructions and DNA concentrations were quantified using a Nanodrop

spectrophotometer before genome sequencing using an Illumina MiSeq (California, USA). Nextera XT libraries (Illumina, California, USA) were prepared following manufacturer's protocols and short paired-end reads were sequenced using 2 × 300 bp paired end v3 reagent kit (Illumina).

Genome datasets

Genomes were assembled *de novo* using SPAdes (version 3.8.0; Bankevich et al. 2012). The average number of contigs was 72 (range: 12–471) for an average total assembled sequence size of 1.70 Mbp (range: 1.56–1.86). The average N50 contig length (L50) was 14,577 (range: 3,794–55,912) and the average GC content was 30.8 % (range: 30.5–31.6). Short read data are available on the NCBI short read archive (SRA), associated with BioProject PRJNA576513. Assembled genomes and supplementary material are available from FigShare (doi:10.6084/m9.figshare.9956597; individual accession numbers and assembled genome statistics in **Supplementary Table S1**). We augmented our collection by assembling a context dataset of previously published isolates (n=204) to represent the known diversity of *C. jejuni* and *C. coli* (Calland et al., 2020; Sheppard et al., 2010, 2013, 2014). In addition, we also compared our single city survey with a previously published survey from Oxford in the UK (n=874 isolates collected over 1 year; Cody et al. 2012). Isolate genomes were archived in BIGSdb and MLST sequence types (STs) derived through BLAST comparison with the pubMLST database (Dingle et al., 2001; Jolley et al., 2018; Jolley and Maiden, 2010; Sheppard et al., 2012). Simpson's index of ST diversity was calculated for the Cairo and Oxford datasets using the equation:

$$D = 1 - \frac{\sum n(n-1)}{N(N-1)}$$

Where *n* is the number of isolates of each sequence type and *N* is the total number of isolates (Grundmann et al., 2001).

Core and accessory genome characterisation

Alignments were made from concatenated gene sequences of all core genes (found in ≥95% isolates) using MAFFT (version 7; Katoh and Standley 2013) on a gene-by-gene basis. Separate maximum-likelihood phylogenies were constructed with a GTR+I+G substitution model and ultra-fast

bootstrapping (1000 bootstraps) (Hoang et al., 2018) implemented in IQ-TREE (version 1.6.8; Nguyen et al. 2015) for *C. jejuni* (n=1,048) and *C. coli* (n=132) and visualized on Microreact (https://next.microreact.org/project/Cjejuni_Egypt; https://next.microreact.org/project/Ccoli_Egypt) (Argimón et al., 2016).

All unique genes present in at least one isolate (the pangenome) were identified by automated annotation using PROKKA (version 1.13; Seemann 2014) followed by PIRATE, a pangenomics tool that allows for orthologue gene clustering in bacteria (Bayliss et al., 2019). We defined genes in PIRATE using a wide range of amino acid percentage sequence identity thresholds for Markov Cluster algorithm (MCL) clustering (45, 50, 60, 70, 80, 90, 95, 98). Genes in the pangenome were ordered initially using the NCTC 11168 reference followed by the order defined in PIRATE based on gene synteny and frequency (Gundogdu et al., 2007; Pascoe et al., 2019). As described previously, a matrix was produced summarizing the presence/absence and allelic diversity of every gene in the pangenome list, with core genes defined as present in 95% of the genomes and accessory genes as present in at least one isolate (**Supplementary table S2**) (Méric et al., 2014). Pairwise core and accessory genome distances were compared using PopPunk (version 2.2.0; Lees et al. 2019) which uses pairwise nucleotide k-mer comparisons to distinguish shared sequence and gene content to identify divergence of the accessory genome in relation to the core genome. A two-component Gaussian mixture model was used to construct a network to define clusters, comparable to other *Campylobacter* studies (Components: 41; density 0.0579; transitivity: 0.9518; score: 0.8907) (Pascoe et al. 2020).

Core genome variation between isolates was quantified by calculating the pairwise average nucleotide identity (ANI) of all (n=112+874) *Campylobacter* genomes using FastANI v.1.058 (Jain et al., 2018). The gene presence matrix produced by PIRATE was used to generate a heatmap of shared pairwise accessory genome genes. Averages were calculated for within and between country comparisons in addition to focussed analysis on the ST21 (*C. jejuni*) and ST828 (*C. coli*) clonal complexes. Antimicrobial resistance genes and putative virulence genes were detected through comparison with

207 reference nucleotide sequences using ABRicate (version 0.8) (<https://github.com/tseemann/abricate>)
 208 and the NCBI database (Chen et al., 2005; NCBI Resource Coordinators, 2013). Point mutations
 209 related to antibiotic resistance genes were identified by PointFinder (Zankari et al., 2017) using the
 210 STAR-AMR software package (<https://github.com/phac-nml/staramr>) (**Supplementary table S3**).
 211

Results

Globally circulating genotypes among Egyptian Campylobacter isolates

We sequenced and characterized a collection of *Campylobacter* spp. isolates (n=112) from clinical cases, broiler carcasses and dairy products collected over a 14-month sampling period in Cairo, Egypt (**Figure 1A; Supplementary table S1**). Isolate genotypes were compared with all genomes deposited in the pubMLST database (97,012 profiles, data accessed 17th February 2020) and ranked according to how frequently they were found associated with human disease (**Figure 1B**). Egyptian *C. jejuni* isolates belonged to 15 clonal complexes (CCs) with a diverse assemblage of STs. Nearly half of the isolates (n = 29, 47%) were from common lineages, isolated many times before and recorded in pubMLST (>50 MLST profiles; **Figure 1B**), including the globally disseminated lineages of ST-21CC (n=37; 41%), ST-206 CC (n=10; 11%) and ST-464CC (n=7; 8%) the most abundant. Several other poultry-associated clonal complexes, which are common in human disease (Berthenet et al., 2019; Sheppard et al., 2009), including ST-353 (n=3, 3.2%), ST-354 (n=4, 4.3%) and ST-257 (n=4, 4.3%) were identified. Other globally disseminated lineages were found less often in Egypt (n≤2) i.e., ST-460 (n=2), ST-1034 (n=2), ST-42 (n=1), ST-45 CC (n=1), ST-573 (n=1), ST-574 (n=1) and ST-658 (n=1) (Colles et al., 2010; Olkkola et al., 2016).

Local sequence types

Comparison with a collection representing the known genetic diversity of *C. jejuni* and *C. coli* identified some common STs (>1,000 profiles in pubMLST) that were completely absent in our Egyptian collection, i.e., ST-53, ST-829 (*C. coli*), ST-22, ST-61, ST-51, ST-1068 (*C. jejuni*) (**Figure 1CD**). Two isolates belonging to ST-1287CC, a genotype that has previously been isolated from poultry and the environment (Magnússon et al., 2011), was observed exclusively among our Egyptian isolates, yet absent in UK and genetic context datasets. Furthermore, there were also some STs belonging to ST-21CC that were found in Egyptian isolate collection (n=>3) that are rare in global collections (<100 profiles in pubMLST), i.e., ST-1519 (n=4), ST-3769 (n=3). It was also observed that more *C. coli* was found among Egyptian clinical isolates than is typically observed, specifically the *C. coli* lineage ST-828 CC 90.4% *C. coli* isolates (19/21) belonged to the ST-828 CC within the Egyptian dataset and two *C. coli* isolates with unassigned CC of sequence types, ST-7951 and ST-1681. Three rare STs belonging to ST-828 CC were exclusively found in Egypt dataset which are ST-1058 (n=1), ST-1059 (n=1), and ST-7950 (n=1).

Increased sharing of accessory genes contributes to a local gene pool

Our Egyptian dataset was compared directly with a previously published study of a single city, ~1-year survey from Oxford in the UK (Cody et al., 2012). Both populations were similarly diverse, specifically there were 50 STs (16 CCs) among the Egyptian isolate collection, with a Simpson's diversity index of 0.817, compared to 205 STs (32 CCs) among the Oxford collection of genomes (Simpson's diversity index = 0.895; **Figure 1CD**). We used PIRATE to construct a pan-genome of all Egyptian and Oxford isolates (n=986). Consistent with other studies, we identified an open pangenome, meaning that the number of genes in the pangenome continues to increase with each additionally sequenced isolate. Accessory genes represented nearly three-quarters of the pangenome (3,410 genes; 74% of pangenome) with a quarter of the genes identified (1,225, 26%) considered core genes present in 95% or more of the isolates. Pairwise comparison of the core nucleotide sequence (%ANI) and accessory genome sharing of all isolates reflected the clonal frame, with clusters of closely related isolates sharing a large percentage of ANI (**Figure 2AB**). Direct comparison between

the Oxford and Cairo datasets suggested an increase in within-country, local accessory gene sharing (Figure 2CD). The structured clustering of pairwise comparisons of shared accessory genes suggested that this may vary between lineages and visualization of the differences in the distribution of pairwise genomic distances with PopPUNK also pointed towards lineage-specific shared gene pools (Figure 2E). Host generalist clonal complex isolates clustered closer together than the more isolated host-specific isolates. This included the two most common clonal complexes identified in our Cairo collection, ST-21CC and ST-828CC, which were investigated further (Figure 2F).

Locally diverged sequence types within the globally disseminated ST-21 clonal complexes

As one might expect of within lineage (clonal complex) comparisons, all ST-21CC isolates shared more than 99% core genome nucleotide identity and shared more accessory genes than the population average (852 genes; Figure 2CF) and significantly more genes were shared between isolates from the same country (*t*-test with Welch correction; $p < 0.0001$). A maximum-likelihood phylogeny of all CC21 isolates ($n=251$), the most common clonal complex identified in our collection from Cairo, identified geography-specific clusters of isolates (Figure 3A). These clones also clustered together when visualizing the distribution of pairwise genomic distances with PopPUNK (Figure 3B). While some specific STs were common in both Oxford and Cairo (ST21 and ST50), others were much more common in one specific location, e.g., ST-53 in Oxford, and ST-1519 and ST-3769 in Cairo (Figure 3C). There was also evidence that some lineages had enhanced AMR (Figure 3D). While the ST-50 genotype is very common and has been reported more than 3,900 times in pubMLST from 40 countries, this among the first reports from Africa. In both Oxford and Cairo datasets, ST-50 was often predicted to be MDR. ST-21 is also very common, with more than 4,000 reports from 33 countries in pubMLST but was much less likely to be MDR. Four isolates of the Cairo specific ST-3769 also represented a high proportion of MDR (Figure 3E).

Extensive multi-drug resistance in local C. coli sequence types

Greater admixture was noted between UK and Egyptian ST-828CC isolates than for ST-21CC – no geographic clustering was observed in either the core or accessory genomes (Figure 4AB). However,

only ST-827 was common in both datasets (**Figure 4C**). Several STs were found in the Oxford dataset that were not identified in Cairo, including the frequently isolated STs -829, -828, -855, 962, -1145 and -5734. Several lineages were highly resistant to lincosamides, with more than half the isolates from ST-828, ST-830 and ST-872 predicted to be resistant (**Figure 4D**). All isolates from ST828 and ST-872 were also predicted to be resistant to chloramphenicol. Overall, *C. coli* isolates (6 of 105, 5.7%) were far more likely to be considered MDR than *C. jejuni* isolates (6 of 876, 0.68%) and ST-828 complex isolates from Cairo (2 of 19, 10.5%) demonstrated much higher rates of MDR than in Oxford (3 of 77, 3.8%; **Figure 4E**).

Antimicrobial resistance genes are distributed across isolates

In characterization of the resistome, each isolate genome was screened for the presence of genes associated with AMR. In Egypt, for *C. jejuni*, the average number of AMR genes per isolate was 6.66, comparable to 6.52 for *C. coli*. In Egypt, the presence of the *tet(O)* gene, conferring tetracycline resistance, was higher in *C. coli* than *C. jejuni* (76% and 43% respectively). This pattern contrasts with Oxford where 41.7% of *C. jejuni* but only 35.3% of *C. coli* isolates were found to harbor *tet(O)*. Whilst a low proportion of Egyptian isolates (6.7%) contained the *blaOXA-61* gene, associated with β -lactam resistance, alternative alleles including *blaOXA-450* and *blaOXA-605* were abundant. In respect to lineage association with genes, in Egypt the ST-21 clonal complex had a high prevalence of genes associated with β -lactam resistance (particularly the *blaOXA-193*, *blaOXA-450* and *blaOXA-605* alleles). The *blaOXA-465* allele was closely related to ST-1034. Furthermore, *blaOXA-61* was closely associated with ST-48 (**Figure 3D**). All of these patterns were reflected amongst the Oxford isolates. However, numerous genes (including *aadE*, *Ant6-la* and *blaOXA-451*) were found amongst distant lineages. The multi-drug efflux pump encoded by a three-gene operon (*cmeABC*) was abundant amongst isolates (n=87,74%) – although an absence of the repressor gene *cmeR* in *C. coli* was observed.

Whilst the average number of resistance genes per isolate was comparable for *C. jejuni* in Egypt, this analysis indicated that *C. coli* held a greater breadth of genes across classes of antimicrobials. Hence,

the proportion of MDR isolates, considered when an isolate is resistant to at least three classes, was 28% for *C. coli* compared to 1% for *C. jejuni* (EFSA, 2021). The majority (88%) of MDR isolates in Egypt were *C. coli*, despite *C. coli* representing about a fifth of the dataset. In other words, a greater proportion of *C. coli* isolates were MDR. In Oxford, half of MDR isolates were *C. coli*, whilst in this case representing less than one tenth of the dataset. The *C. jejuni* isolates that were MDR, were all host generalists – ST-21, ST-48 or ST-206. Amongst Egyptian isolates, genes including *aad9*, *aadE*, *aadE-Cc*, *ant(6)-Ia*, *aph(2'')-If*, *aph(3')-III* and *aph(3')-IIIa* associated with aminoglycoside resistance, were almost exclusively associated with *C. coli*, particularly MDR *C. coli*. This association was not as strong in Oxford. Regarding specific genes and host associations, aminoglycoside resistance-associated genes were infrequent amongst isolates from chicken or dairy products. *ant(6)-Ia* for example, was solely found in human samples. In turn, few isolates from chicken and dairy products were MDR (only 12.5% of MDR isolates was from chicken).

Discussion

Diarrheal disease is a major threat to human health and the second leading cause of death in children under five years' old LMICs (Lanata et al., 2013). *Campylobacteriosis* is a major cause of diarrheal disease worldwide (Amour et al., 2016; ElGendy et al., 2018; Lee et al., 2013) but, despite the potential importance, little is known about *Campylobacter* in countries where it potentially poses the greatest health risk. As studies begin to take a worldview of *Campylobacter* epidemiology and transmission (Mottet and Tempio, 2017), we describe globally disseminated agriculture-associated disease-causing lineages based on core and accessory genome content, with evidence that local accessory genome sharing driving acquisition of AMR genes in specific lineages.

The Egyptian *Campylobacter* isolates included a diverse set of STs, including common disease-causing lineages and regional STs, that have rarely been reported from other parts of the world. Industrialized agriculture globalization has dispersed livestock worldwide (Mottet and Tempio, 2017), expanding the geographical range of *C. jejuni*. This is evident in the Egyptian collection as two of the most predominant genotypes belonged to the ST-21 and ST-206 clonal complexes (Figure 1D). These two host generalist clonal complexes have been extensively reported worldwide and frequently isolated from various reservoir hosts, including human clinical samples (Berthenet et al., 2019; Dingle et al., 2001; Grove-White et al., 2011; Mossong et al., 2016; Sheppard et al., 2009; Suerbaum et al., 2001). The ST21-CC exhibits considerable genome plasticity with a clear association with several virulence genes and resistance to various antimicrobial agents (Aksomaitiene et al., 2019; Gripp et al., 2011; Habib et al., 2010; Wieczorek et al., 2017; T. Zhang et al., 2016). Poultry-associated clonal complexes, ST-206, ST-464, ST-48, ST-257 and ST-354 were also common among the Egyptian isolates, all of which are among the most prevalent clonal complexes isolated in Europe (Colles et al., 2011; Elhadidy et al., 2018; Fiedoruk et al., 2019).

Further comparison of isolate genotypes collected in Cairo with a large global collection revealed the absence of certain lineages, most notably the lack of the cattle-associated genotype, ST-61 (Dingle et

al., 2002; Mourkas et al., 2020). There was only one isolate, of dairy product origin, that could be attributed to a cattle-specialist clonal complex (ST-42), which is unexpected as several (n=24) isolates were sampled from dairy products. *Campylobacter* isolates from cattle have predominantly been sampled from meat, milk products and fecal sources (n=2,726 in pubMLST; Kwan et al., 2008; Mourkas et al., 2020; Epping et al., 2021). Suggesting that dairy products isolates might represent a different source population in Egypt.

There were also no isolates belonging to the ST-22 CC, a particularly high risk lineage which is commonly found among patients with post-infectious complications of campylobacteriosis, such as GBS and IBS (Revez et al., 2011; Peters et al., 2021). Although one isolate in our collection was from ST-45 CC, this host generalist clonal complex is often one of the most commonly isolates lineages in clinical surveillance studies worldwide (De Haan et al., 2010; Sheppard et al., 2009; Shin et al., 2013; Sopwith et al., 2008). Notably however, it is often absent (or under-represented) in studies conducted in LMICs (Pascoe et al., 2020; Sarhangi et al., 2021). This is consistent with observations from other LMICs, where local differences in disease epidemiology are reflected by the absence of common *Campylobacter* lineages, and the presence of rare or unique sequence types (Graham et al., 2016; Pascoe et al., 2020; Prachantasena et al., 2016; P. Zhang et al., 2020). Among our Egyptian isolates the ST-1287 clonal complex (n=2) has been reported less than 4 times from other parts of the world (Colles et al., 2011; de Haan et al., 2010; Ramonaite et al., 2014; P. Zhang et al., 2020).

Geographical differences have been noted in ST-21CC (Kärenlampi et al., 2007; Kovanen et al., 2014; Olkkola et al., 2016; Pascoe et al., 2017; Wallace et al., 2021). ST-21 CC isolates are among the most common *C. jejuni* genotypes isolated worldwide, with one quarter of *C. jejuni* isolates recorded in the pubMLST database are ST21 CC. Isolates of the ST-50 sequence type (n= 3,915) alone have been sampled from 6 continents and 44 countries, although this will be their first report from Africa (Jolley et al., 2018). Our Egyptian ST-50 isolates do cluster together on a ML phylogeny of ST-21 CC isolates and away from the Oxford ST-21 CC when grouped by PopPunk. Two sequence types were unique to Egypt, ST-1519 and ST-3769, with nearly 10% of the ST-3769 isolates were

MDR. A slightly greater proportion of the Egyptian ST-50 isolates were also MDR, although this sequence type has been observed to be MDR in other parts of the world (Elhadidy et al., 2020).

The *C. coli* ST-828 clonal complex did not show as much geographical segregation, and when grouping our Egyptian isolates by core and accessory genome distances they clustered with the UK isolates, despite several STs being isolated in only one of the datasets. STs found in the Egyptian dataset were more often MDR than UK isolates, and overall *C. coli* from Cairo were far more MDR than *C. coli* isolates from developed countries (Du et al., 2018; Gharbi et al., 2018; Mourkas et al., 2019). The most compelling clarification for such abundance could be that *C. coli* of ST-828 CC have a great recombination potential besides the accumulation of *C. jejuni* DNA throughout the genome of this lineage which could have led to the acquisition of multiple AMR genes (Sheppard et al., 2008, 2013).

Overall, there is a clear evidence of local sharing and recent acquisition of accessory gene content of AMR genes within the Egyptian isolates. Specifically, pairwise clustering of isolates by core and accessory genome distances recapitulated clusters according to ST and clonal complex (**Figure 2**), however most Egyptian isolates were more tightly clustered than the Oxford dataset, consistent with shared acquisition of accessory genes. Overall, ANI and shared accessory genes were similar between Oxford and Egyptian isolates (per isolate), however the two most common clonal complexes found in our Cairo dataset demonstrated greater sharing of accessory genes, indicative of a shared gene pool. Our study suggested that while geographical partitioning doesn't impact the composition of the core genome, represented by the shared STs and CCs, the accessory genome is influenced. Within the Egyptian isolates, the most prevalent *C. jejuni* genotypes (ST-21CC and ST-206CC) showed clear evidence of transmission of MDR determinants among lineages. Multiple factors could influence this, such as livestock and food production practices and the segregation of MDR isoaltes. However, selective pressure for MDR is clearly attributable to antibiotic usage and potentially zoonotic transmissions as well as the rate of horizontal gene transfer (Fiedoruk et al., 2019). Our study provides evidence to support programs aimed at improved antibiotic stewardship in clinical and

409 veterinary settings. With strict control measures, and an understanding of transmission of strains from
410 animal reservoirs through the food production chain, it may be possible to reduce contamination with
411 MDR *Campylobacter* in Egypt.
412

Author statements

Author contributions

SM, JKC, BP, ME and SKS designed the study and wrote the paper.

JKC, BP, EM, GF, CL, HW performed genomic analysis.

BP, CL and MDH sequenced and assembled genomes.

All authors contributed and approved the final manuscript.

Conflict of interest

All authors declare no conflict of interest.

Funding information

This study was partially funded by Zewail City internal research grant fund (ZC 004-2019) and joint ASRT/BA research grant (Project number 1110) awarded to ME. We would like to thank the British Academy of Medical Sciences for funding the research stay of ME at the Milner Centre of Evolution, University of Bath through the Daniel Turnberg Travel Fellowship. Research and computation were supported by the Medical Research Council (MR/L015080/1), the Wellcome Trust (088786/C/09/Z) and the Biotechnology and Biological Sciences Research Council (BB/I02464X/1 and BB/R003491/1).

Ethical approval

The study represents a retrospective study that involved sequencing the genomes of a historical strain collection and no patient data collection was involved in this study. Ethical approval was granted from the respective ethics committee in the Egyptian central directorate of research and health development before conducting the study.

Acknowledgements

References

- Abd El-Tawab, A. A., Ammar, A. M., Ahmed, H. A., El Hofy, F. I., and Hefny, A. A. (2018). Fluoroquinolone resistance and gyrA mutations in *Campylobacter jejuni* and *Campylobacter coli* isolated from chicken in Egypt. *Journal of Global Antimicrobial Resistance*, 13, 22–23.
<https://doi.org/10.1016/j.jgar.2018.02.019>
- Abdi Hachesoo, B., Khoshbakht, R., Sharifi Yazdi, H., Tabatabaei, M., Hosseinzadeh, S., and Asasi, K. (2014). Tetracycline Resistance Genes in *Campylobacter jejuni* and *C. coli* Isolated From Poultry Carcasses. *Jundishapur Journal of Microbiology*, 7(8). <https://doi.org/10.5812/jjm.12129>
- Achtman, M. (2008). Evolution, population structure, and phylogeography of genetically monomorphic bacterial pathogens. *Annual Review of Microbiology*, 62, 53–70.
<https://doi.org/10.1146/annurev.micro.62.081307.162832>
- Aksomaitiene, J., Ramonaite, S., Tamuleviciene, E., Novoslavskij, A., Alter, T., and Malakauskas, M. (2019). Overlap of Antibiotic Resistant *Campylobacter jejuni* MLST Genotypes Isolated From Humans, Broiler Products, Dairy Cattle and Wild Birds in Lithuania. *Frontiers in Microbiology*, 10.
<https://doi.org/10.3389/fmicb.2019.01377>
- Amour, C., Gratz, J., Mduma, E., Svensen, E., Rogawski, E. T., McGrath, M., Seidman, J. C., McCormick, B. J., Shrestha, S., Samie, A., Mahfuz, M., Qureshi, S., Hotwani, A., Babji, S., Trigos, D. R., Lima, A. M., Bodhidatta, L., Bessong, P., Ahmed, T., ... Etiology, Risk Factors, and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development Project (MAL-ED) Network Investigators. (2016). Epidemiology and Impact of Campylobacter Infection in Children in 8 Low-Resource Settings: Results From the MAL-ED Study. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*, 63(9), 1171–1179.
<https://doi.org/10.1093/cid/ciw542>
- Argimón, S., Abudahab, K., Goater, R. J. E., Fedosejev, A., Bhai, J., Glasner, C., Feil, E. J., Holden, M. T. G., Yeats, C. A., Grundmann, H., Spratt, B. G., and Aanensen, D. M. (2016). Microreact: Visualizing and sharing data for genomic epidemiology and phylogeography. *Microbial Genomics*, 2(11), e000093.
<https://doi.org/10.1099/mgen.0.000093>
- Asakura, H., Brüggemann, H., Sheppard, S. K., Ekawa, T., Meyer, T. F., Yamamoto, S., and Igimi, S. (2012). Molecular evidence for the thriving of *Campylobacter jejuni* ST-4526 in Japan. *PloS One*, 7(11), e48394. <https://doi.org/10.1371/journal.pone.0048394>

- 470 Asuming-Bediako, N., Parry-Hanson Kunadu, A., Abraham, S., and Habib, I. (2019). Campylobacter at the
471 Human–Food Interface: The African Perspective. *Pathogens*, 8(2), 87.
472 <https://doi.org/10.3390/pathogens8020087>
- 473 Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko,
474 S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G., Alekseyev, M.
475 A., and Pevzner, P. A. (2012). SPAdes: A New Genome Assembly Algorithm and Its Applications to
476 Single-Cell Sequencing. *Journal of Computational Biology*, 19(5), 455–477.
477 <https://doi.org/10.1089/cmb.2012.0021>
- 478 Bayliss, S. C., Thorpe, H. A., Coyle, N. M., Sheppard, S. K., and Feil, E. J. (2019). PIRATE: A fast and scalable
479 pangenomics toolbox for clustering diverged orthologues in bacteria. *GigaScience*, 8(10).
480 <https://doi.org/10.1093/gigascience/giz119>
- 481 Berthenet, E., Thépault, A., Chemaly, M., Rivoal, K., Ducournau, A., Buissonnière, A., Bénéjat, L., Bessède,
482 E., Mégraud, F., Sheppard, S. K., and Lehours, P. (2019). Source attribution of *Campylobacter jejuni*
483 shows variable importance of chicken and ruminants reservoirs in non-invasive and invasive French
484 clinical isolates. *Scientific Reports*, 9(1), 8098. <https://doi.org/10.1038/s41598-019-44454-2>
- 485 Calland, J. K., Pascoe, B., Bayliss, S. C., Mourkas, E., Berthenet, E., Thorpe, H. A., Hitchings, M. D., Feil, E.
486 J., Blaser, M. J., Falush, D., and Sheppard, S. K. (2020). Quantifying bacterial evolution in the wild: A
487 birthday problem for Campylobacter lineages. *BioRxiv*, 2020.12.02.407999.
488 <https://doi.org/10.1101/2020.12.02.407999>
- 489 CDC. (2020, March 13). *Antibiotic-resistant Germs: New Threats*. Centers for Disease Control and Prevention.
490 <https://www.cdc.gov/drugresistance/biggest-threats.html>
- 491 Chen, L., Yang, J., Yu, J., Yao, Z., Sun, L., Shen, Y., and Jin, Q. (2005). VFDB: A reference database for
492 bacterial virulence factors. *Nucleic Acids Research*, 33(suppl_1), D325–D328.
493 <https://doi.org/10.1093/nar/gki008>
- 494 Cody, A. J., McCarthy, N. M., Wimalaratna, H. L., Colles, F. M., Clark, L., Bowler, I. C. J. W., Maiden, M. C.
495 J., and Dingle, K. E. (2012). A Longitudinal 6-Year Study of the Molecular Epidemiology of Clinical
496 Campylobacter Isolates in Oxfordshire, United Kingdom. *Journal of Clinical Microbiology*, 50(10),
497 3193–3201. <https://doi.org/10.1128/JCM.01086-12>
- 498 Coker, A. O. (2002). Human Campylobacteriosis in Developing Countries. *Emerging Infectious Diseases*, 8(3),
499 237–243. <https://doi.org/10.3201/eid0803.010233>

500 Colles, F. M., Ali, J. S., Sheppard, S. K., McCarthy, N. D., and Maiden, M. C. J. (2011). *Campylobacter*
501 populations in wild and domesticated Mallard ducks (*Anas platyrhynchos*): *Campylobacter* in ducks.
502 *Environmental Microbiology Reports*, 3(5), 574–580. [https://doi.org/10.1111/j.1758-](https://doi.org/10.1111/j.1758-2229.2011.00265.x)
503 2229.2011.00265.x

504 Colles, F. M., and Maiden, M. C. J. (2012). *Campylobacter* sequence typing databases: Applications and future
505 prospects. *Microbiology*, 158(11), 2695–2709. <https://doi.org/10.1099/mic.0.062000-0>

506 Colles, F. M., McCarthy, N. D., Sheppard, S. K., Layton, R., and Maiden, M. C. J. (2010). Comparison of
507 *Campylobacter* populations isolated from a free-range broiler flock before and after slaughter.
508 *International Journal of Food Microbiology*, 137(2–3), 259–264.
509 <https://doi.org/10.1016/j.ijfoodmicro.2009.12.021>

510 Comas, I., Coscolla, M., Luo, T., Borrell, S., Holt, K. E., Kato-Maeda, M., Parkhill, J., Malla, B., Berg, S.,
511 Thwaites, G., Yeboah-Manu, D., Bothamley, G., Mei, J., Wei, L., Bentley, S., Harris, S. R., Niemann,
512 S., Diel, R., Aseffa, A., ... Gagneux, S. (2013). Out-of-Africa migration and Neolithic coexpansion of
513 *Mycobacterium tuberculosis* with modern humans. *Nature Genetics*, 45(10), 1176–1182.
514 <https://doi.org/10.1038/ng.2744>

515 Crofts, A. A., Poly, F. M., Ewing, C. P., Kuroiwa, J. M., Rimmer, J. E., Harro, C., Sack, D., Talaat, K. R.,
516 Porter, C. K., Gutierrez, R. L., DeNearing, B., Brubaker, J., Laird, R. M., Maue, A. C., Jaep, K.,
517 Alcala, A., Tribble, D. R., Riddle, Mark. S., Ramakrishnan, A., ... Trent, M. S. (2018). *Campylobacter*
518 *jejuni* transcriptional and genetic adaptation during human infection. *Nature Microbiology*, 3(4), 494–
519 502. <https://doi.org/10.1038/s41564-018-0133-7>

520 Dahshan, H., Abd-Elall, A. M. M., Megahed, A. M., Abd-El-Kader, M. A., and Nabawy, E. E. (2015).
521 Veterinary antibiotic resistance, residues, and ecological risks in environmental samples obtained from
522 poultry farms, Egypt. *Environmental Monitoring and Assessment*, 187(2), 2.
523 <https://doi.org/10.1007/s10661-014-4218-3>

524 De Haan, C. P. A., Kivistö, R., Hakkinen, M., Rautelin, H., and Hänninen, M. L. (2010). Decreasing trend of
525 overlapping multilocus sequence types between human and chicken *Campylobacter jejuni* isolates over
526 a decade in finland. *Applied and Environmental Microbiology*. <https://doi.org/10.1128/AEM.00581-10>

527 Dearlove, B. L., Cody, A. J., Pascoe, B., Méric, G., Wilson, D. J., and Sheppard, S. K. (2016). Rapid host
528 switching in generalist *Campylobacter* strains erodes the signal for tracing human infections. *The ISME*
529 *Journal*, 10(3), 721–729. <https://doi.org/10.1038/ismej.2015.149>

530 Dingle, K. E., Colles, F. M., Ure, R., Wagenaar, J. A., Duim, B., Bolton, F. J., Fox, A. J., Wareing, D. R. A.,
531 and Maiden, M. C. J. (2002). Molecular Characterization of *Campylobacter jejuni* Clones: A Basis for
532 Epidemiologic Investigation. *Emerging Infectious Diseases*, 8(9), 949–955.
533 <https://doi.org/10.3201/eid0809.020122>

534 Dingle, K. E., Colles, F. M., Wareing, D. R. A., Ure, R., Fox, A. J., Bolton, F. E., Bootsma, H. J., Willems, R. J.
535 L., Urwin, R., and Maiden, M. C. J. (2001). Multilocus Sequence Typing System for *Campylobacter*
536 *jejuni*. *Journal of Clinical Microbiology*, 39(1), 14–23. <https://doi.org/10.1128/JCM.39.1.14-23.2001>

537 Du, Y., Wang, C., Ye, Y., Liu, Y., Wang, A., Li, Y., Zhou, X., Pan, H., Zhang, J., and Xu, X. (2018). Molecular
538 Identification of Multidrug-Resistant *Campylobacter* Species From Diarrheal Patients and Poultry Meat
539 in Shanghai, China. *Frontiers in Microbiology*, 9. <https://doi.org/10.3389/fmicb.2018.01642>

540 ElGendy, A., Sainato, R., Kuroiwa, J., Poly, F., Riddle, M. S., Guerry, P., and Porter, C. K. (2018).
541 Epidemiology of *Campylobacter* Infections among Children in Egypt. *The American Journal of*
542 *Tropical Medicine and Hygiene*, 98(2), 581–585. <https://doi.org/10.4269/ajtmh.17-0469>

543 Elhadidy, M., Ali, M. M., El-Shibiny, A., Miller, W. G., Elkhatib, W. F., Botteldoorn, N., and Dierick, K.
544 (2020). Antimicrobial resistance patterns and molecular resistance markers of *Campylobacter jejuni*
545 isolates from human diarrheal cases. *PLOS ONE*, 15(1), e0227833.
546 <https://doi.org/10.1371/journal.pone.0227833>

547 Elhadidy, M., Miller, W. G., Arguello, H., Álvarez-Ordóñez, A., Duarte, A., Dierick, K., and Botteldoorn, N.
548 (2018). Genetic Basis and Clonal Population Structure of Antibiotic Resistance in *Campylobacter*
549 *jejuni* Isolated From Broiler Carcasses in Belgium. *Frontiers in Microbiology*, 9, 1014.
550 <https://doi.org/10.3389/fmicb.2018.01014>

551 Epping, L., Walther, B., Piro, R. M., Knüver, M.-T., Huber, C., Thürmer, A., Flieger, A., Fruth, A., Janecko, N.,
552 Wieler, L. H., Stingl, K., and Semmler, T. (2021). Genome-wide insights into population structure and
553 host specificity of *Campylobacter jejuni*. *Scientific Reports*, 11(1), 10358.
554 <https://doi.org/10.1038/s41598-021-89683-6>

555 Facciola, A., Riso, R., Avventuroso, E., Visalli, G., Delia, S. A., and Laganà, P. (2017). CAMPYLOBACTER:
556 FROM MICROBIOLOGY TO PREVENTION. *Journal of Preventive Medicine and Hygiene*, Vol 58,
557 E79-92 Pages. <https://doi.org/10.15167/2421-4248/JPMH2017.58.2.672>

558 Falush, D., Wirth, T., Linz, B., Pritchard, J. K., Stephens, M., Kidd, M., Blaser, M. J., Graham, D. Y., Vacher,
559 S., Perez-Perez, G. I., Yamaoka, Y., Mégraud, F., Otto, K., Reichard, U., Katzwitsch, E., Wang, X.,

560 Achtman, M., and Suerbaum, S. (2003). Traces of human migrations in *Helicobacter pylori*
561 populations. *Science (New York, N.Y.)*, 299(5612), 1582–1585.
562 <https://doi.org/10.1126/science.1080857>

563 Fiedoruk, K., Daniluk, T., Rozkiewicz, D., Oldak, E., Prasad, S., and Swiecicka, I. (2019). Whole-genome
564 comparative analysis of *Campylobacter jejuni* strains isolated from patients with diarrhea in
565 northeastern Poland. *Gut Pathogens*, 11. <https://doi.org/10.1186/s13099-019-0313-x>

566 French, N., Barrigas, M., Brown, P., Ribiero, P., Williams, N., Leatherbarrow, H., Birtles, R., Bolton, E.,
567 Fearnhead, P., and Fox, A. (2005). Spatial epidemiology and natural population structure of
568 *Campylobacter jejuni* colonizing a farmland ecosystem. *Environmental Microbiology*, 7(8), 1116–
569 1126. <https://doi.org/10.1111/j.1462-2920.2005.00782.x>

570 Gagneux, S., and Small, P. M. (2007). Global phylogeography of *Mycobacterium tuberculosis* and implications
571 for tuberculosis product development. *The Lancet. Infectious Diseases*, 7(5), 328–337.
572 [https://doi.org/10.1016/S1473-3099\(07\)70108-1](https://doi.org/10.1016/S1473-3099(07)70108-1)

573 Gharbi, M., Béjaoui, A., Ben Hamda, C., Jouini, A., Ghedira, K., Zrelli, C., Hamrouni, S., Aouadhi, C.,
574 Bessoussa, G., Ghram, A., and Maaroufi, A. (2018). Prevalence and Antibiotic Resistance Patterns of
575 *Campylobacter* spp. Isolated from Broiler Chickens in the North of Tunisia. *BioMed Research*
576 *International*, 2018, 1–7. <https://doi.org/10.1155/2018/7943786>

577 Gillespie, I. A., O'Brien, S. J., Frost, J. A., Adak, G. K., Horby, P., Swan, A. V., Painter, M. J., Neal, K. R., and
578 the *Campylobacter* Sentinel Surveillance System Collaborators. (2002). A Case-Case Comparison of
579 *Campylobacter coli* and *Campylobacter jejuni* Infection: A Tool for Generating Hypotheses. *Emerging*
580 *Infectious Diseases*, 8(9), 937–942. <https://doi.org/10.3201/eid0809.010817>

581 Graham, J. P., Vasco, K., and Trueba, G. (2016). Hyperendemic *Campylobacter jejuni* in guinea pigs (*Cavia*
582 *porcellus*) raised for food in a semi-rural community of Quito, Ecuador: *Campylobacter jejuni* in
583 guinea pigs. *Environmental Microbiology Reports*, 8(3), 382–387. [https://doi.org/10.1111/1758-](https://doi.org/10.1111/1758-2229.12396)
584 2229.12396

585 Griekspoor, P., Colles, F. M., McCarthy, N. D., Hansbro, P. M., Ashhurst-Smith, C., Olsen, B., Hasselquist,
586 D., Maiden, M. C. J., and Waldenström, J. (2013). Marked host specificity and lack of
587 phylogeographic population structure of *Campylobacter jejuni* in wild birds. *Molecular Ecology*, 22(5),
588 1463–1472. <https://doi.org/10.1111/mec.12144>

589 Gripp, E., Hlahla, D., Didelot, X., Kops, F., Maurischat, S., Tedin, K., Alter, T., Ellerbroek, L., Schreiber, K.,
590 Schomburg, D., Janssen, T., Bartholomäus, P., Hofreuter, D., Woltemate, S., Uhr, M., Brenneke, B.,
591 Grüning, P., Gerlach, G., Wieler, L., ... Josenhans, C. (2011). Closely related *Campylobacter jejuni*
592 strains from different sources reveal a generalist rather than a specialist lifestyle. *BMC Genomics*.
593 <https://doi.org/10.1186/1471-2164-12-584>

594 Grove-White, D. H., Leatherbarrow, A. J. H., Cripps, P. J., Diggle, P. J., and French, N. P. (2011). Molecular
595 epidemiology and genetic diversity of *Campylobacter jejuni* in ruminants. *Epidemiology and Infection*.
596 <https://doi.org/10.1017/S0950268810002736>

597 Grundmann, H., Hori, S., and Tanner, G. (2001). Determining Confidence Intervals When Measuring Genetic
598 Diversity and the Discriminatory Abilities of Typing Methods for Microorganisms. *Journal of Clinical*
599 *Microbiology*, 39(11), 4190–4192. <https://doi.org/10.1128/JCM.39.11.4190-4192.2001>

600 Gundogdu, O., Bentley, S. D., Holden, M. T., Parkhill, J., Dorrell, N., and Wren, B. W. (2007). Re-annotation
601 and re-analysis of the *Campylobacter jejuni* NCTC11168 genome sequence. *BMC Genomics*, 8(1), 162.
602 <https://doi.org/10.1186/1471-2164-8-162>

603 Gürtler, M., Alter, T., Kasimir, S., and Fehlhaber, K. (2005). The importance of *Campylobacter coli* in human
604 campylobacteriosis: Prevalence and genetic characterization. *Epidemiology and Infection*, 133(6),
605 1081–1087. <https://doi.org/10.1017/S0950268805004164>

606 Habib, I., Uyttendaele, M., and De Zutter, L. (2010). Survival of poultry-derived *Campylobacter jejuni* of
607 multilocus sequence type clonal complexes 21 and 45 under freeze, chill, oxidative, acid and heat
608 stresses. *Food Microbiology*, 27(6), 829–834. <https://doi.org/10.1016/j.fm.2010.04.009>

609 Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., and Vinh, L. S. (2018). UFBoot2: Improving the
610 Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*, 35(2), 518–522.
611 <https://doi.org/10.1093/molbev/msx281>

612 Inglis, G. D., Gusse, J. F., House, K. E., Shelton, T. G., and Taboada, E. N. (2019). Tetracycline Resistant
613 *Campylobacter jejuni* Subtypes Emanating from Beef Cattle Administered Non-Therapeutic
614 Chlortetracycline are Longitudinally Transmitted within the Production Continuum but are Not
615 Detected in Ground Beef. *Microorganisms*, 8(1), 23. <https://doi.org/10.3390/microorganisms8010023>

616 Islam, Z., van Belkum, A., Wagenaar, J. A., Cody, A. J., de Boer, A. G., Sarker, S. K., Jacobs, B. C., Talukder,
617 K. A., and Endtz, H. P. (2014). Comparative population structure analysis of *Campylobacter jejuni*
618 from human and poultry origin in Bangladesh. *European Journal of Clinical Microbiology and*

619 *Infectious Diseases: Official Publication of the European Society of Clinical Microbiology*, 33(12),
620 2173–2181. <https://doi.org/10.1007/s10096-014-2184-x>

621 Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T., and Aluru, S. (2018). High throughput
622 ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature Communications*,
623 9(1), 5114. <https://doi.org/10.1038/s41467-018-07641-9>

624 Jolley, K. A., Bray, J. E., and Maiden, M. C. J. (2018). Open-access bacterial population genomics: BIGSdb
625 software, the PubMLST.org website and their applications. *Wellcome Open Research*, 3, 124.
626 <https://doi.org/10.12688/wellcomeopenres.14826.1>

627 Jolley, K. A., and Maiden, M. C. (2010). BIGSdb: Scalable analysis of bacterial genome variation at the
628 population level. *BMC Bioinformatics*, 11(1), 595. <https://doi.org/10.1186/1471-2105-11-595>

629 Kaakoush, N. O., Castañó-Rodríguez, N., Mitchell, H. M., and Man, S. M. (2015). Global Epidemiology of
630 *Campylobacter* Infection. *Clinical Microbiology Reviews*, 28(3), 687–720.
631 <https://doi.org/10.1128/CMR.00006-15>

632 Kärenlampi, R., Rautelin, H., Schönberg-Norio, D., Paulin, L., and Hänninen, M.-L. (2007). Longitudinal
633 Study of Finnish *Campylobacter jejuni* and *C. coli* Isolates from Humans, Using Multilocus Sequence
634 Typing, Including Comparison with Epidemiological Data and Isolates from Poultry and Cattle.
635 *Applied and Environmental Microbiology*, 73(1), 148–155. <https://doi.org/10.1128/AEM.01488-06>

636 Katoh, K., and Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:
637 Improvements in Performance and Usability. *Molecular Biology and Evolution*, 30(4), 772–780.
638 <https://doi.org/10.1093/molbev/mst010>

639 Kirk, K. F., Méric, G., Nielsen, H. L., Pascoe, B., Sheppard, S. K., Thorlacius-Ussing, O., and Nielsen, H.
640 (2018). Molecular epidemiology and comparative genomics of *Campylobacter concisus* strains from
641 saliva, faeces and gut mucosal biopsies in inflammatory bowel disease. *Scientific Reports*, 8(1), 1902.
642 <https://doi.org/10.1038/s41598-018-20135-4>

643 Kivistö, R. I., Kovanen, S., Skarp-de Haan, A., Schott, T., Rahkio, M., Rossi, M., and Hänninen, M.-L. (2014).
644 Evolution and comparative genomics of *Campylobacter jejuni* ST-677 clonal complex. *Genome*
645 *Biology and Evolution*, 6(9), 2424–2438. <https://doi.org/10.1093/gbe/evu194>

646 Kovanen, S. M., Kivistö, R. I., Rossi, M., Schott, T., Kärkkäinen, U.-M., Tuuminen, T., Uksila, J., Rautelin, H.,
647 and Hänninen, M.-L. (2014). Multilocus sequence typing (MLST) and whole-genome MLST of

648 Campylobacter jejuni isolates from human infections in three districts during a seasonal peak in
649 Finland. *Journal of Clinical Microbiology*, 52(12), 4147–4154. <https://doi.org/10.1128/JCM.01959-14>

650 Kwan, P. S. L., Birtles, A., Bolton, F. J., French, N. P., Robinson, S. E., Newbold, L. S., Upton, M., and Fox, A.
651 J. (2008). Longitudinal Study of the Molecular Epidemiology of Campylobacter jejuni in Cattle on
652 Dairy Farms. *Applied and Environmental Microbiology*, 74(12), 3626–3633.
653 <https://doi.org/10.1128/AEM.01669-07>

654 Lanata, C. F., Fischer-Walker, C. L., Olascoaga, A. C., Torres, C. X., Aryee, M. J., Black, R. E., and for the
655 Child Health Epidemiology Reference Group of the World Health Organization and UNICEF. (2013).
656 Global Causes of Diarrheal Disease Mortality in Children <5 Years of Age: A Systematic Review.
657 *PLoS ONE*, 8(9), e72788. <https://doi.org/10.1371/journal.pone.0072788>

658 Lee, G., Pan, W., Peñataro Yori, P., Paredes Olortegui, M., Tilley, D., Gregory, M., Oberhelman, R., Burga, R.,
659 Chavez, C. B., and Kosek, M. (2013). Symptomatic and Asymptomatic Campylobacter Infections
660 Associated with Reduced Growth in Peruvian Children. *PLoS Neglected Tropical Diseases*, 7(1).
661 <https://doi.org/10.1371/journal.pntd.0002036>

662 Lees, J. A., Harris, S. R., Tonkin-Hill, G., Gladstone, R. A., Lo, S. W., Weiser, J. N., Corander, J., Bentley, S.
663 D., and Croucher, N. J. (2019). Fast and flexible bacterial genomic epidemiology with PopPUNK.
664 *Genome Research*, 29(2), 304–316. <https://doi.org/10.1101/gr.241455.118>

665 Liu, J., Platts-Mills, J. A., Juma, J., Kabir, F., Nkeze, J., Okoi, C., Operario, D. J., Uddin, J., Ahmed, S., Alonso,
666 P. L., Antonio, M., Becker, S. M., Blackwelder, W. C., Breiman, R. F., Faruque, A. S. G., Fields, B.,
667 Gratz, J., Haque, R., Hossain, A., ... Houpt, E. R. (2016). Use of quantitative molecular diagnostic
668 methods to identify causes of diarrhoea in children: A reanalysis of the GEMS case-control study.
669 *Lancet (London, England)*, 388(10051), 1291–1301. [https://doi.org/10.1016/S0140-6736\(16\)31529-X](https://doi.org/10.1016/S0140-6736(16)31529-X)

670 Liu, L., Johnson, H. L., Cousens, S., Perin, J., Scott, S., Lawn, J. E., Rudan, I., Campbell, H., Cibulskis, R., Li,
671 M., Mathers, C., and Black, R. E. (2012). Global, regional, and national causes of child mortality: An
672 updated systematic analysis for 2010 with time trends since 2000. *The Lancet*, 379(9832), 2151–2161.
673 [https://doi.org/10.1016/S0140-6736\(12\)60560-1](https://doi.org/10.1016/S0140-6736(12)60560-1)

674 Luangtongkum, T., Jeon, B., Han, J., Plummer, P., Logue, C. M., and Zhang, Q. (2009). Antibiotic resistance in
675 Campylobacter: Emergence, transmission and persistence. *Future Microbiology*, 4(2), 189–200.
676 <https://doi.org/10.2217/17460913.4.2.189>

677 Magnússon, S. H., Guðmundsdóttir, S., Reynisson, E., Rúnarsson, Á. R., Harðardóttir, H., Gunnarson, E.,
678 Georgsson, F., Reiersen, J., and Marteinson, V. T. (2011). Comparison of *Campylobacter jejuni*
679 isolates from human, food, veterinary and environmental sources in Iceland using PFGE, MLST and
680 fla-SVR sequencing. *Journal of Applied Microbiology*, 111(4), 971–981.
681 <https://doi.org/10.1111/j.1365-2672.2011.05100.x>

682 McCormick, B. J. J., and Lang, D. R. (2016). Diarrheal disease and enteric infections in LMIC communities:
683 How big is the problem? *Tropical Diseases, Travel Medicine and Vaccines*, 2(1), 11.
684 <https://doi.org/10.1186/s40794-016-0028-7>

685 Méric, G., McNally, A., Pessia, A., Mourkas, E., Pascoe, B., Mageiros, L., Vehkala, M., Corander, J., and
686 Sheppard, S. K. (2018). Convergent Amino Acid Signatures in Polyphyletic *Campylobacter jejuni*
687 Subpopulations Suggest Human Niche Tropism. *Genome Biology and Evolution*, 10(3), 763–774.
688 <https://doi.org/10.1093/gbe/evy026>

689 Méric, G., Yahara, K., Mageiros, L., Pascoe, B., Maiden, M. C. J., Jolley, K. A., and Sheppard, S. K. (2014). A
690 Reference Pan-Genome Approach to Comparative Bacterial Genomics: Identification of Novel
691 Epidemiological Markers in Pathogenic *Campylobacter*. *PLOS ONE*, 9(3), e92798.
692 <https://doi.org/10.1371/journal.pone.0092798>

693 Mohan, V., Stevenson, M., Marshall, J., Fearnhead, P., Holland, B. R., Hotter, G., and French, N. P. (2013).
694 *Campylobacter jejuni* colonization and population structure in urban populations of ducks and starlings
695 in New Zealand. *MicrobiologyOpen*, 2(4), 659–673. <https://doi.org/10.1002/mbo3.102>

696 Moodley, Y., Linz, B., Yamaoka, Y., Windsor, H. M., Breurec, S., Wu, J.-Y., Maady, A., Bernhöft, S.,
697 Thiberge, J.-M., Phuanukoonnon, S., Jobb, G., Siba, P., Graham, D. Y., Marshall, B. J., and Achtman,
698 M. (2009). The peopling of the Pacific from a bacterial perspective. *Science (New York, N.Y.)*,
699 323(5913), 527–530. <https://doi.org/10.1126/science.1166083>

700 Mossong, J., Mughini-Gras, L., Penny, C., Devaux, A., Olinger, C., Losch, S., Cauchie, H.-M., van Pelt, W.,
701 and Ragimbeau, C. (2016). Human *Campylobacteriosis* in Luxembourg, 2010–2013: A Case-Control
702 Study Combined with Multilocus Sequence Typing for Source Attribution and Risk Factor Analysis.
703 *Scientific Reports*, 6(1), 20939. <https://doi.org/10.1038/srep20939>

704 Mottet, A., and Tempio, G. (2017). Global poultry production: Current state and future outlook and challenges.
705 *World's Poultry Science Journal*, 73(2), 245–256. <https://doi.org/10.1017/S0043933917000071>

706 Mourkas, E., Florez-Cuadrado, D., Pascoe, B., Calland, J. K., Bayliss, S. C., Mageiros, L., Méric, G.,
707 Hitchings, M. D., Quesada, A., Porrero, C., Ugarte-Ruiz, M., Gutiérrez-Fernández, J., Domínguez,
708 L., and Sheppard, S. K. (2019). Gene pool transmission of multidrug resistance among *Campylobacter*
709 from livestock, sewage and human disease. *Environmental Microbiology*, 21(12), 4597–4613.
710 <https://doi.org/10.1111/1462-2920.14760>

711 Mourkas, E., Taylor, A. J., Méric, G., Bayliss, S. C., Pascoe, B., Mageiros, L., Calland, J. K., Hitchings, M. D.,
712 Ridley, A., Vidal, A., Forbes, K. J., Strachan, N. J. C., Parker, C. T., Parkhill, J., Jolley, K. A., Cody,
713 A. J., Maiden, M. C. J., Kelly, D. J., and Sheppard, S. K. (2020). Agricultural intensification and the
714 evolution of host specialism in the enteric pathogen *Campylobacter jejuni*. *Proceedings of the National*
715 *Academy of Sciences*, 117(20), 11018–11028. <https://doi.org/10.1073/pnas.1917168117>

716 NCBI Resource Coordinators. (2013). Database resources of the National Center for Biotechnology
717 Information. *Nucleic Acids Research*, 41(D1), D8–D20. <https://doi.org/10.1093/nar/gks1189>

718 Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. (2015). IQ-TREE: A fast and effective
719 stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and*
720 *Evolution*, 32(1), 268–274. <https://doi.org/10.1093/molbev/msu300>

721 Nichols, G. L., Richardson, J. F., Sheppard, S. K., Lane, C., and Sarran, C. (2012). *Campylobacter*
722 epidemiology: A descriptive study reviewing 1 million cases in England and Wales between 1989 and
723 2011. *BMJ Open*, 2(4), e001179. <https://doi.org/10.1136/bmjopen-2012-001179>

724 Olkkola, S., Nykäsenoja, S., Raulo, S., Llärena, A.-K., Kovanen, S., Kivistö, R., Myllyniemi, A.-L., and
725 Hänninen, M.-L. (2016). Antimicrobial Resistance and Multilocus Sequence Types of Finnish
726 *Campylobacter jejuni* Isolates from Multiple Sources. *Zoonoses and Public Health*, 63(1), 10–19.
727 <https://doi.org/10.1111/zph.12198>

728 Pascoe, B., Méric, G., Yahara, K., Wimalaratna, H., Murray, S., Hitchings, M. D., Sproston, E. L., Carrillo, C.
729 D., Taboada, E. N., Cooper, K. K., Huynh, S., Cody, A. J., Jolley, K. A., Maiden, M. C. J., McCarthy,
730 N. D., Didelot, X., Parker, C. T., and Sheppard, S. K. (2017). Local genes for local bacteria: Evidence
731 of allopatry in the genomes of transatlantic *Campylobacter* populations. *Molecular Ecology*, 26(17),
732 4497–4508. <https://doi.org/10.1111/mec.14176>

733 Pascoe, B., Schiaffino, F., Murray, S., Méric, G., Bayliss, S. C., Hitchings, M. D., Mourkas, E., Calland, J. K.,
734 Burga, R., Yori, P. P., Jolley, K. A., Cooper, K. K., Parker, C. T., Olortegui, M. P., Kosek, M. N., and
735 Sheppard, S. K. (2020). Genomic epidemiology of *Campylobacter jejuni* associated with asymptomatic

736 pediatric infection in the Peruvian Amazon. *PLOS Neglected Tropical Diseases*, 14(8), e0008533.
737 <https://doi.org/10.1371/journal.pntd.0008533>

738 Pascoe, B., Williams, L. K., Calland, J. K., Meric, G., Hitchings, M. D., Dyer, M., Ryder, J., Shaw, S., Lopes,
739 B. S., Chintoan-Uta, C., Allan, E., Vidal, A., Fearnley, C., Everest, P., Pachebat, J. A., Cogan, T. A.,
740 Stevens, M. P., Humphrey, T. J., Wilkinson, T. S., ... Sheppard, S. K. (2019). Domestication of
741 *Campylobacter jejuni* NCTC 11168. *Microbial Genomics*, 5(7). <https://doi.org/10.1099/mgen.0.000279>

742 Peters, S., Pascoe, B., Wu, Z., Bayliss, S. C., Zeng, X., Edwinston, A., Veerabadhran-Gurunathan, S., Jawahir,
743 S., Calland, J. K., Mourkas, E., Patel, R., Wiens, T., Decuir, M., Boxrud, D., Smith, K., Parker, C. T.,
744 Farrugia, G., Zhang, Q., Sheppard, S. K., and Grover, M. (2021). *Campylobacter jejuni* genotypes are
745 associated with post-infection irritable bowel syndrome in humans. *Communications Biology*, 4(1), 1–
746 17. <https://doi.org/10.1038/s42003-021-02554-8>

747 Platts-Mills, J. A., and Kosek, M. (2014). Update on the burden of *Campylobacter* in developing countries.
748 *Current Opinion in Infectious Diseases*, 27(5), 444–450.
749 <https://doi.org/10.1097/QCO.0000000000000091>

750 Prachantasena, S., Charununtakorn, P., Muangnoicharoen, S., Hankla, L., Techawal, N., Chaveerach, P.,
751 Tuitemwong, P., Chokesajjawatee, N., Williams, N., Humphrey, T., and Luangtongkum, T. (2016).
752 Distribution and Genetic Profiles of *Campylobacter* in Commercial Broiler Production from Breeder to
753 Slaughter in Thailand. *PloS One*, 11(2), e0149585. <https://doi.org/10.1371/journal.pone.0149585>

754 Ramonaite, S., Kudirkienė, E., Tamulevičienė, E., Levinienė, G., Malakauskas, A., Götz, G., Alter, T., and
755 Malakauskas, M. (2014). Prevalence and genotypes of *Campylobacter jejuni* from urban environmental
756 sources in comparison with clinical isolates from children. *Journal of Medical Microbiology*.
757 <https://doi.org/10.1099/jmm.0.072892-0>

758 Rao, M. R. (2001). Pathogenicity and Convalescent Excretion of *Campylobacter* in Rural Egyptian Children.
759 *American Journal of Epidemiology*, 154(2), 166–173. <https://doi.org/10.1093/aje/154.2.166>

760 Reed, R. P., Friedland, I. R., Wegerhoff, F. O., and Khoosal, M. (1996). *Campylobacter* bacteremia in children:
761 *The Pediatric Infectious Disease Journal*, 15(4), 345–348. [https://doi.org/10.1097/00006454-](https://doi.org/10.1097/00006454-199604000-00012)
762 199604000-00012

763 Revez, J., Rossi, M., Ellström, P., Haan, C. de, Rautelin, H., and Hänninen, M.-L. (2011). Finnish
764 *Campylobacter jejuni* Strains of Multilocus Sequence Type ST-22 Complex Have Two Lineages with
765 Different Characteristics. *PLOS ONE*, 6(10), e26880. <https://doi.org/10.1371/journal.pone.0026880>

766 Sabry, N. A., Farid, S. F., and Dawoud, D. M. (2014). Antibiotic dispensing in Egyptian community
767 pharmacies: An observational study. *Research in Social and Administrative Pharmacy*, 10(1), 168–
768 184. <https://doi.org/10.1016/j.sapharm.2013.03.004>

769 Sarhangi, M., Bakhshi, B., and Peeraeyeh, S. N. (2021). High prevalence of *Campylobacter jejuni* CC21 and
770 CC257 clonal complexes in children with gastroenteritis in Tehran, Iran. *BMC Infectious Diseases*,
771 21(1), 108. <https://doi.org/10.1186/s12879-021-05778-5>

772 Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*, 30(14), 2068–2069.
773 <https://doi.org/10.1093/bioinformatics/btu153>

774 Sheppard, S. K., Cheng, L., Méric, G., de Haan, C. P. A., Llaena, A.-K., Martinen, P., Vidal, A., Ridley, A.,
775 Clifton-Hadley, F., Connor, T. R., Strachan, N. J. C., Forbes, K., Colles, F. M., Jolley, K. A., Bentley,
776 S. D., Maiden, M. C. J., Hänninen, M.-L., Parkhill, J., Hanage, W. P., and Corander, J. (2014). Cryptic
777 ecology among host generalist *Campylobacter jejuni* in domestic animals. *Molecular Ecology*, 23(10),
778 2442–2451. <https://doi.org/10.1111/mec.12742>

779 Sheppard, S. K., Dallas, J. F., MacRae, M., McCarthy, N. D., Sproston, E. L., Gormley, F. J., Strachan, N. J. C.,
780 Ogden, I. D., Maiden, M. C. J., and Forbes, K. J. (2009). *Campylobacter* genotypes from food animals,
781 environmental sources and clinical disease in Scotland 2005/6. *International Journal of Food*
782 *Microbiology*, 134(1–2), 96–103. <https://doi.org/10.1016/j.ijfoodmicro.2009.02.010>

783 Sheppard, S. K., Dallas, J. F., Strachan, N. J. C., MacRae, M., McCarthy, N. D., Wilson, D. J., Gormley, F. J.,
784 Falush, D., Ogden, I. D., Maiden, M. C. J., and Forbes, K. J. (2009). *Campylobacter* Genotyping to
785 Determine the Source of Human Infection . *Clinical Infectious Diseases*.
786 <https://doi.org/10.1086/597402>

787 Sheppard, S. K., Dallas, J. F., Wilson, D. J., Strachan, N. J. C., McCarthy, N. D., Jolley, K. A., Colles, F. M.,
788 Rotariu, O., Ogden, I. D., Forbes, K. J., and Maiden, M. C. J. (2010). Evolution of an Agriculture-
789 Associated Disease Causing *Campylobacter coli* Clade: Evidence from National Surveillance Data in
790 Scotland. *PLoS ONE*, 5(12), e15708. <https://doi.org/10.1371/journal.pone.0015708>

791 Sheppard, S. K., Didelot, X., Jolley, K. A., Darling, A. E., Pascoe, B., Méric, G., Kelly, D. J., Cody, A., Colles,
792 F. M., Strachan, N. J. C., Ogden, I. D., Forbes, K., French, N. P., Carter, P., Miller, W. G., McCarthy,
793 N. D., Owen, R., Litrup, E., Egholm, M., ... Falush, D. (2013). Progressive genome-wide
794 introgression in agricultural *Campylobacter coli*. *Molecular Ecology*, 22(4), 1051–1064.
795 <https://doi.org/10.1111/mec.12162>

Sheppard, S. K., Jolley, K. A., and Maiden, M. C. J. (2012). A Gene-By-Gene Approach to Bacterial Population Genomics: Whole Genome MLST of *Campylobacter*. *Genes*, 3(2), 261–277. <https://doi.org/10.3390/genes3020261>

Sheppard, S. K., McCarthy, N. D., Falush, D., and Maiden, M. C. J. (2008). Convergence of *Campylobacter* Species: Implications for Bacterial Evolution. *Science*, 320(5873), 237–239. <https://doi.org/10.1126/science.1155532>

Shin, E., Oh, Y., Kim, M., Jung, J., and Lee, Y. (2013). Antimicrobial Resistance Patterns and Corresponding Multilocus Sequence Types of the *Campylobacter jejuni* Isolates from Human Diarrheal Samples. *Microbial Drug Resistance*, 19(2), 110–116. <https://doi.org/10.1089/mdr.2012.0099>

Sopwith, W., Birtles, A., Matthews, M., Fox, A., Gee, S., Painter, M., Regan, M., Syed, Q., and Bolton, E. (2008). Identification of Potential Environmentally Adapted *Campylobacter jejuni* Strain, United Kingdom. *Emerging Infectious Diseases*, 14(11), 1769–1773. <https://doi.org/10.3201/eid1411.071678>

Suerbaum, S., Lohrengel, M., Sonnevend, A., Ruberg, F., and Kist, M. (2001). Allelic diversity and recombination in *Campylobacter jejuni*. *Journal of Bacteriology*. <https://doi.org/10.1128/JB.183.8.2553-2559.2001>

Tam, C. C., O'Brien, S. J., Tompkins, D. S., Bolton, F. J., Berry, L., Dodds, J., Choudhury, D., Halstead, F., Iturriza-Gomara, M., Mather, K., Rait, G., Ridge, A., Rodrigues, L. C., Wain, J., Wood, B., Gray, J. J., and the IID2 Study Executive Committee. (2012). Changes in Causes of Acute Gastroenteritis in the United Kingdom Over 15 Years: Microbiologic Findings From 2 Prospective, Population-Based Studies of Infectious Intestinal Disease. *Clinical Infectious Diseases*, 54(9), 1275–1286. <https://doi.org/10.1093/cid/cis028>

The European Union Summary Report on Antimicrobial Resistance in zoonotic and indicator bacteria from humans, animals and food in 2018/2019. (2021). *EFSA Journal* 2021; 19(4):6490, 179 pp. <https://doi.org/10.2903/j.efsa.2021.6490>

Thépault, A., Rose, V., Quesne, S., Poezevara, T., Béven, V., Hirchaud, E., Touzain, F., Lucas, P., Méric, G., Mageiros, L., Sheppard, S. K., Chemaly, M., and Rivoal, K. (2018). Ruminant and chicken: Important sources of campylobacteriosis in France despite a variation of source attribution in 2009 and 2015. *Scientific Reports*, 8(1), 9305. <https://doi.org/10.1038/s41598-018-27558-z>

Waite, D. W., and Taylor, M. W. (2015). Exploring the avian gut microbiota: Current trends and future directions. *Frontiers in Microbiology*, 6. <https://doi.org/10.3389/fmicb.2015.00673>

- Wallace, R. L., Cribb, D. M., Bulach, D. M., Ingle, D. J., Joensen, K. G., Nielsen, E. M., Leekitcharoenphon, P.,
Stingl, K., and Kirk, M. D. (2021). *Campylobacter jejuni* ST50, a pathogen of global importance: A
comparative genomic analysis of isolates from Australia, Europe and North America. *Zoonoses and
Public Health*. <https://doi.org/10.1111/zph.12853>
- Wieczorek, K., Denis, E., Lachtara, B., and Osek, J. (2017). Distribution of *Campylobacter jejuni* multilocus
sequence types isolated from chickens in Poland. *Poultry Science*. <https://doi.org/10.3382/ps/pew343>
- Wierzba, T. F., Abdel-Messih, I. A., Gharib, B., Baqar, S., Hendaui, A., Khalil, I., Omar, T. A., Khayat, H. E.,
Putnam, S. D., Sanders, J. W., Ng, L.-K., Price, L. J., Scott, D. A., and Frenck, R. R. (2008).
Campylobacter Infection as a Trigger for Guillain-Barré Syndrome in Egypt. *PLoS ONE*, 3(11), e3674.
<https://doi.org/10.1371/journal.pone.0003674>
- Zankari, E., Allesøe, R., Joensen, K. G., Cavaco, L. M., Lund, O., and Aarestrup, F. M. (2017). PointFinder: A
novel web tool for WGS-based detection of antimicrobial resistance associated with chromosomal
point mutations in bacterial pathogens. *Journal of Antimicrobial Chemotherapy*, 72(10), 2764–2768.
<https://doi.org/10.1093/jac/dkx217>
- Zhang, P., Zhang, X., Liu, Y., Jiang, J., Shen, Z., Chen, Q., and Ma, X. (2020). Multilocus Sequence Types and
Antimicrobial Resistance of *Campylobacter jejuni* and *C. coli* Isolates of Human Patients From
Beijing, China, 2017–2018. *Frontiers in Microbiology*, 11. <https://doi.org/10.3389/fmicb.2020.554784>
- Zhang, T., Luo, Q., Chen, Y., Li, T., Wen, G., Zhang, R., Luo, L., Lu, Q., Ai, D., Wang, H., and Shao, H.
(2016). Molecular epidemiology, virulence determinants and antimicrobial resistance of
Campylobacter spreading in retail chicken meat in Central China. *Gut Pathogens*.
<https://doi.org/10.1186/s13099-016-0132-2>
- Zhao, S., Tyson, G. H., Chen, Y., Li, C., Mukherjee, S., Young, S., Lam, C., Folster, J. P., Whichard, J. M., and
McDermott, P. F. (2016). Whole-Genome Sequencing Analysis Accurately Predicts Antimicrobial
Resistance Phenotypes in *Campylobacter* spp. *Applied and Environmental Microbiology*, 82(2), 459–
466. <https://doi.org/10.1128/AEM.02873-15>
- Zollner-Schwetz, I., and Krause, R. (2015). Therapy of acute gastroenteritis: Role of antibiotics. *Clinical
Microbiology and Infection*, 21(8), 744–749. <https://doi.org/10.1016/j.cmi.2015.03.002>

Tables and Figures

Figure 1: (A) Demographic data for Cairo, Egypt from which we collected *Campylobacter* spp. isolates (n=112; red circles) from clinical cases, broiler carcasses and dairy products collected over a 14-month sampling period. Our collection was compared to a similar published survey from Oxford, UK (n=874; green circles; Cody et al. 2012) and isolates from pubMLST.org (n=204; grey circles) for additional genetic context. (B) Clonal complexes (CCs) of isolates collected from Cairo were ranked according to the frequency in our local dataset and how often they have been sampled from human disease isolates (data from pubMLST; <https://pubmlst.org/>). Alignments were made from concatenated gene sequences of all core genes (found in $\geq 95\%$ isolates) using MAFFT (version 7; Katoh and Standley 2013) on a gene-by-gene basis. Separate maximum-likelihood phylogenies were constructed with a GTR+I+G substitution model and ultra-fast bootstrapping (1000 bootstraps) (Hoang et al., 2018) implemented in IQ-TREE (version 1.6.8; Nguyen et al. 2015) for (C) *C. jejuni* (n=1,048) and (D) *C. coli* (n=132) and visualized on Microreact (https://next.microreact.org/project/Cjejuni_Egypt; https://next.microreact.org/project/Ccoli_Egypt) (Argimón et al., 2016).

Figure 2: (A) Core genome variation between isolates was quantified by calculating the pairwise average nucleotide identity (ANI) of all UK and Oxford *Campylobacter* genomes (n=112+874) using FastANI v.1.058 (Jain et al., 2018). (B) The ANI for each isolate was estimated and averages compared within and between countries. (C) The gene presence matrix produced by PIRATE was used to generate a heatmap of shared pairwise accessory genome genes. (D) Averages were calculated for within and between country. (E) Clustering of pairwise core and accessory genome distances were compared using PopPunk. Interactive visualisation on Microreact: <https://microreact.org/project/Campy-Egypt>. (F) Comparisons of within and between country ANI and accessory gene sharing were also analysed for our two most common Egyptian lineages, ST21 (*C. jejuni*) and ST828 (*C. coli*) clonal complexes.

Figure 3: (A) Sub-tree of all Egyptian and UK ST21 clonal complex (CC21) isolates (n=251). Common sequence types are annotated and ST50 (yellow) and ST21 (green) are highlighted. (B) Within clonal complex clustering of pairwise core and accessory genome distances with PopPunk. (C) Prevalence of the most common sequence types found within CC21. (D) Prevalence of AMR determinants grouped by antibiotic class for each CC21 ST. (E) Prevalence of MDR isolates (AMR determinants for three or more antibiotic classes) in CC21 STs.

Figure 4: (A) Sub-tree of all Egyptian and UK ST828 clonal complex (CC828) isolates (n=94). Common sequence types are annotated and ST827 (orange) is highlighted. (B) Within clonal complex clustering of pairwise core and accessory genome distances with PopPunk. (C) Prevalence of the most common sequence types found within CC828. (D) Prevalence of AMR determinants grouped by antibiotic class for each CC828 ST. (E) Prevalence of MDR isolates (AMR determinants for three or more antibiotic classes) in CC828 STs.

Supplementary information

Supplementary table 1: Summary of isolate collection data and genome statistics

Supplementary table 2: Summary PIRATE core and accessory genome statistics.

Supplementary table 3: Summary of AMR genes identified by comparison with the NCBI database and pointfinder.

A

B

D

C

E

A

B

D

C

E