Inactivation of the entire Arabidopsis group II GH3s confers tolerance to salinity and drought

Rubén Casanova-Sáez^{1,2*}, Eduardo Mateo-Bonmati^{1,2,3}, Jan Šimura¹, Aleš Pěnčík⁴, Ondřej Novák^{1,4}, Paul Staswick⁵, and Karin Ljung^{1*}

¹Umeå Plant Science Centre (UPSC), Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, 901 83 Umeå, Sweden

⁴Laboratory of Growth Regulators, Faculty of Science, Palacký University and Institute of Experimental Botany of the Czech Academy of Sciences, Šlechtitelů 27, Olomouc, Czech Republic

⁵Department of Agronomy and Horticulture, University of Nebraska, Lincoln, NE, USA

*Corresponding authors: Rubén Casanova-Sáez (ruben.casanova.saez@slu.se) and Karin Ljung (karin.ljung@slu.se)

²These authors contributed equally to this work.

³Current address: John Innes Centre, NR4 7UA, Norwich, United Kingdom

Running head: Blocking IAA GH3s confers tolerance to water deprivation

Keywords: Arabidopsis, auxin, GH3, drought, salinity, stress tolerance

ORCID (R.C.-S.): 0000-0001-5683-7051

Twitter (R.C.-S.): @rcasanovasaez

ORCID (E.M.-B.): 0000-0002-2364-5173

Twitter (E.M.-B.): @emateobonmati

ORCID (J.Š.): 0000-0002-1567-2278

ORCID (A.P.): 0000-0002-1314-2249

ORCID (O.N.): 0000-0003-3452-0154

ORCID (P.S.): 0000-0003-2798-0275

ORCID (K.L.): 0000-0003-2901-189X

Twitter (K.L.): @GroupLjung

ABSTRACT

Indole-3-acetic acid (IAA) controls a plethora of developmental processes. Thus, regulation of their levels is of great relevance for plant performance. Cellular IAA concentration depends on the combined result of its transport, biosynthesis and various redundant pathways to inactivate IAA, including oxidation and conjugation. Group II members of the GRETCHEN HAGEN 3 (GH3) gene family code for acyl acid amido synthetases catalysing the conjugation of IAA to amino acids. However, the high level of functional redundancy among them has hampered thorough analysis of their roles in plant development. In this work, we generated an *Arabidopsis gh3.1,2,3,4,5,6,9,17 (gh3oct)* mutant to knock-out the group II GH3 pathway. The *gh3oct* plants had an improved root architecture, were more tolerant to osmotic stresses due to locally increased IAA levels and were more drought tolerant. IAA metabolite quantification in *gh3oct* plants suggested the existence of additional GH3-like enzymes in IAA metabolism. Moreover, our data suggested that oxIAA production depends, at least partly, on the GH3 pathway. Targeted stress-hormone analysis further suggested an involvement of ABA in the differential response to salinity of *gh3oct* plants. Taken together, our data provide new insights into the roles of group II GH3s in IAA metabolism and hormone-regulated plant development.

INTRODUCTION

Regulation of the levels and distribution of the major auxin indole-3-acetic acid (IAA) is of pivotal importance for plant development because it regulates multiple processes, including embryo, seed, root and leaf development, meristem maintenance, shoot branching and responses to environmental stresses (Casanova-Saez & Voss, 2019; Leftley *et al.*, 2021). Spatiotemporal fluctuations of IAA levels in response to external and internal stimuli drive plant developmental responses by triggering signalling cascades that result in modulation of cell division, differentiation and growth (Gallei *et al.*, 2020).

Control of IAA levels and distribution within plant cells and tissues is achieved by intercellular and intracellular transport mechanisms (Band, 2021; Hammes *et al.*, 2021) and spatiotemporally regulated IAA biosynthesis and inactivation (Zhao, 2018; Casanova-Saez *et al.*, 2021). IAA metabolic inactivation primarily involves oxidation processes and the formation of IAA-sugar and IAA-amino acid (IAA-aa) conjugates (Casanova-Saez *et al.*, 2021). IAA oxidation is facilitated by the angiosperm-related DIOXYGENASE FOR AUXIN OXIDATION (DAO) proteins (Zhao *et al.*, 2013; Porco *et al.*, 2016; Zhang *et al.*, 2016; Takehara *et al.*, 2020). The formation of oxIAA is an irreversible IAA inactivation process that operates to remove excess auxin, as inferred from higher oxIAA formation detected at sites of IAA maxima (Pencik *et al.*, 2013). Formation of ester-linked IAA conjugates with glucose (IAA-glc) is a reversible reaction known to regulate IAA levels and homeostasis. Several members of the UDP-glucosyltransferase superfamily have been shown to mediate the conjugation of both IAA and oxIAA to glucose *in vivo* (Mateo-Bonmati *et al.*, 2021).

IAA is also inactivated by conjugation to amino acids via amide bonds, a reaction catalysed by members of the GRETCHEN HAGEN 3 (GH3) family of acyl acid amido synthetases (Staswick *et al.*, 2005). Most of these conjugates, including IAA-leucine, IAA-alanine and IAA-phenylalanine, can be hydrolysed to free IAA, and therefore serve as IAA storage forms (LeClere *et al.*, 2002; Rampey *et al.*, 2004). However, IAA irreversibly conjugates to aspartate (IAA-Asp) and glutamate (IAA-Glu) during catabolism (Ostin *et al.*, 1998; Rampey *et al.*, 2004). IAA-Asp and IAA-Glu catabolites have been found at higher levels compared with reversible IAA-aa conjugates in different plant species (Kowalczyk & Sandberg, 2001; Kojima *et al.*, 2009; Pencik *et al.*, 2009). Several IAA-Asp metabolites, including 6-OH-IAA-Asp and oxIAA-Asp, have been identified in the last few decades (Ostin *et al.*, 1998; Kai *et al.*, 2007). Recently, the IAA oxidase DAO1 was found to additionally operate in the GH3 pathway by mediating the formation of oxIAA-Asp from IAA-Asp in tobacco and Arabidopsis (Müller *et al.*, 2021).

Based on sequence homology and substrate specificity studies, 19 Arabidopsis GH3

proteins have been classified into three functionally diverse groups, of which group II consists of eight GH3 members with the ability to mediate the conjugation of IAA to amino acids, i.e. GH3.1, GH3.2, GH3.3, GH3.4, GH3.5, GH3.6, GH3.9 and GH3.17 (Staswick *et al.*, 2002; Staswick *et al.*, 2005). These group II GH3s are conserved across land plants (Terol *et al.*, 2006).

Even though all Arabidopsis group II GH3s can conjugate IAA to different amino acids, substrate preferences exist among them. For instance, GH3.17 preferentially conjugates IAA to Glu, GH3.3 and GH3.4 preferentially conjugate IAA to Asp, and GH3.2, GH3.5 and GH3.6 show a similar predilection for Asp and Glu (Staswick *et al.*, 2005; Brunoni *et al.*, 2019). A range of amino acid substrate specificities has also been documented for GH3 orthologs in other plant species, e.g. *Physcomitrella patens* (Ludwig-Muller *et al.*, 2009), *Vitis vinifera* (Bottcher *et al.*, 2011) and *Picea abies* (Brunoni *et al.*, 2020). Group II GH3s are also known to be promiscuous regarding the acyl acid substrate. GH3.3, GH3.5 and GH3.6 have been found to modulate jasmonic acid homeostasis by mediating its conjugation to amino acids (Gutierrez *et al.*, 2012). GH3.5 has been further shown to mediate the conjugation of the auxin phenylacetic acid, and the benzoates salicylic acid and benzoic acid (Westfall *et al.*, 2016).

Although specific roles have been reported for individual group II GH3 members (Khan & Stone, 2007; Park *et al.*, 2007; Du *et al.*, 2012; Zheng *et al.*, 2016; Di Mambro *et al.*, 2017; Kirungu *et al.*, 2019), their overlapping expression domains and functions in IAA inactivation have likely masked additional roles of these GH3s in IAA metabolism and plant development. In the present study, we generated a group II GH3 octuple knockout to bypass the redundancy in GH3-mediated IAA metabolism. The mutant plants showed pleiotropic high-auxin-related phenotypes, such as longer hypocotyls and petioles, as well as a more branched root system with no apparent penalty on primary root growth. Phenotypic, physiological and targeted-hormonomic analyses further indicated that group II GH3s redundantly modulate IAA-dependent salinity tolerance and that they collectively participate in the response to drought. We also obtained evidence supporting a connection between the GH3 pathway and oxIAA production. Finally, our data suggested that additional GH3 or GH3-like genes might operate in IAA conjugation to amino acids.

MATERIALS AND METHODS

Plant material and culture conditions

Seeds from the Arabidopsis wild-type accession Col-0 and Ler and from the *gh3* insertional lines (Table S1) were obtained from the Nottingham Arabidopsis Stock Centre

(<http://arabidopsis.info>) and from the Arabidopsis Biological Resource Center (<https://abrc.osu.edu>). The presence and position of all insertions were confirmed by PCR amplification using gene-specific primers together with the insertion-specific primers Ds5¹-1 (for SGT lines), 3'-dSpm (for SM lines) and LBb1.3 (for SALK lines) (see Tables S1 and S2).

Arabidopsis seeds were surface sterilized with a bleach solution (40% vol/vol commercial bleach in dH₂O and 0.002% Triton X-100) for 8 min and then washed four times with sterile deionized water. Seeds were stratified for a minimum of 2 days and then sown under sterile conditions on square petri dishes containing half-strength Murashige & Skoog salt mixture (Duchefa, M0221), 1% sucrose, 0.05% MES hydrate (Sigma, M2933) and 0.8% plant agar (Duchefa, P1001) with the pH adjusted to 5.7 with potassium hydroxide. Flowering plants were grown in pots containing a 3:1 mixture of organic soil and vermiculite. All plants were grown, unless otherwise stated, under long-day conditions (16 h light, 8 h darkness) at 22 ± 1°C under cool white fluorescent light (150 µmol photons m⁻² s⁻¹). Flowering time was recorded as the number of rosette leaves at bolting under long- and short-day (8 h light, 16 h darkness) conditions.

Chemical treatments and drought experiments

Salinity and osmotic treatments were performed on half-strength MS plates supplemented with sodium chloride, D-sorbitol or D-mannitol before autoclaving. The inhibitors of YUCCA-mediated IAA biosynthesis yucasin [5-(4-chlorophenyl)-4H-1,2,4-triazole-3-thiol] and PPBo (4-phenoxyphenylboronic acid), kindly provided by Prof. Ken-ichiro Hayashi, were prepared in DMSO stock solutions. IAA treatments were performed on MS plates supplemented with indole-3-acetic acid from a stock solution in absolute ethanol. Yucasin, PPBo and IAA were added to the autoclaved MS media before pouring into plates. To mimic drought conditions *in vitro*, we prepared half-strength MS plates with a reduced water potential as described previously (Verslues *et al.*, 2006). Solidified MS medium was perfused with an overlay solution containing poly(ethylene glycol) (PEG8000; Sigma, P2139). Control plates were perfused with an overlay MS solution lacking PEG8000. For drought treatment on soil-grown plants, stratified seeds were planted in pots containing 40 g of soil mixture. Initially, 1 litre of water was added to each tray. Irrigation was then withheld until both *gh3oct* and wild-type plants displayed dryness symptoms. The weight of the pots was recorded to estimate and compare the water loss in the pots for each genotype.

Genome-wide identification of T-DNA and transposon insertions

Around 14 µg of nuclear-enriched DNA was purified from 1 g of *gh3oct* seedlings using a previously described protocol (Hanania *et al.*, 2004). Whole-genome sequencing of the sample was performed at BGI Hong Kong using a BGISEQ-500 sequencing platform. 47 million 150-bp-long reads were obtained, reaching a 59X genome depth. Trimmed fastq files were used to map the different insertions using Easymap software (Lup *et al.*, 2021). Raw reads were deposited in the Short Read Archive (SRA) with the code SRX8771538.

Gene expression analyses

For expression analyses, RNA was isolated using the Total RNA Purification Kit (Norgen). DNA was removed using the RNase-Free DNase I Kit (Norgen). First-strand cDNA synthesis was performed with the iScript cDNA Synthesis Kit (BioRad). *ACTIN2* gene was used as an internal control for relative expression quantification. Four biological replicates (each being a pool of several plants) were analysed in triplicate. qPCR reactions were performed in 10 µl reactions containing 4 µl of LightCycler 480 SYBR Green I Master (Roche), 4 µl of PCR-grade water (Roche), 1 µl of the corresponding primer pair (10 µM each) and 1 µl of the cDNA template. The primers used are listed in Table S2. Quantification of relative gene expression was performed using the comparative C_T method ($2^{-\Delta\Delta C_t}$) (Schmittgen & Livak, 2008) on a CFX384 Touch Real-Time PCR Detection System (BioRad).

Quantification of IAA and IAA metabolites

Extraction and purification of the targeted compounds (IAA, oxIAA, IAA-Asp, IAA-Glu, IAA-glc, oxIAA-glc) were performed according to (Novak *et al.*, 2012). Briefly, 10 mg of frozen material per sample was homogenized using a bead mill (27 Hz, 10 min, 4°C; MixerMill, Retsch GmbH, Haan, Germany) and extracted in 1 ml of 50 mM sodium phosphate buffer containing 0.1% sodium diethyldithiocarbamate and a mixture of ¹³C₆ isotopically labelled internal standards (Olchemim, Olomouc, Czech Republic). After centrifugation (20 000 g, 15 min, 4°C), the supernatant was transferred into new Eppendorf tubes. The pH was then adjusted to 2.5 with 1 M HCl and samples were immediately applied to preconditioned solid-phase extraction columns (Oasis HLB, 30 mg of 1 ml; Waters Inc., Milford, MA, USA). After sample application, each column was rinsed with 2 ml 5% methanol. Compounds of interest were subsequently eluted with 2 ml of 80% methanol. UHPLC-MS/MS analysis was performed according to the method described in (Pencik *et al.*, 2018), using an LC-MS/MS system consisting of a 1290 Infinity Binary LC System coupled to a 6490 Triple Quad LC/MS System with Jet Stream and Dual Ion Funnel technologies (Agilent Technologies, Santa Clara, CA, USA). The

quantification was carried out in Agilent MassHunter Workstation Quantitative Analysis software (Agilent Technologies, Santa Clara, CA, USA).

Quantification of JA, JA-Ile, SA and ABA

Samples were extracted, purified and analysed according to a method described in (Simura *et al.*, 2018). Briefly, around 20 mg of frozen material per sample was homogenized and extracted in 1 mL of ice-cold 50% aqueous acetonitrile (v/v) containing a mixture of [¹³C], [¹⁵N] or [²H] isotopically labelled internal standards using a bead mill (27 Hz, 10 min, 4°C; MixerMill, Retsch GmbH, Haan, Germany) and sonicator (3 min, 4°C; Ultrasonic bath P 310 H, Elma, Germany). After centrifugation (14 000 RPM, 15 min, 4°C), the supernatant was purified as follows. A solid-phase extraction column Oasis HLB (30 mg 1 cc, Waters Inc., Milford, MA, USA) was conditioned with 1 ml of 100% methanol and 1 ml of deionized water (Milli-Q, Merck Millipore, Burlington, MA, USA). After the conditioning steps, each sample was loaded on a SPE column and the flow-through fraction was collected as well as the 1 ml 30% aqueous acetonitrile (v/v) elution fraction. Samples were evaporated to dryness using a SpeedVac SPD111V (Thermo Scientific, Waltham, MA, USA). Prior to LC-MS analysis, samples were dissolved in 40 µL of 30% acetonitrile (v/v) and transferred to insert-equipped vials. Mass spectrometry analysis of targeted compounds was performed using an UHPLC-ESI-MS/MS system comprising a 1290 Infinity Binary LC system coupled to a 6490 Triple Quad LC/MS system with Jet Stream and Dual Ion Funnel technologies (Agilent Technologies, Santa Clara, CA, USA). The quantification was carried out in Agilent MassHunter Workstation Quantitative Analysis software (Agilent Technologies, Santa Clara, CA, USA).

Phenotypic and statistical analyses

For root and hypocotyl phenotyping, vertically grown plates were imaged using Epson Perfection V600 photo scanners. For petiole length determination, horizontally grown plates were photographed from above. Lengths were measured from scaled images using FIJI software (Schindelin *et al.*, 2012).

Statistically significant differences between mean values from wild type and mutants were analysed by a two-tailed Student's *t*-test. Multiple comparisons of genotypes, treatments and times were performed by one-way ANOVA followed by Tukey's post hoc test to a 95% confidence level. Analyses were carried out in GraphPad Prism 6.01.

Accession numbers

ACTIN2 (At3g18780), *GH3.1* (At2g14960), *GH3.2* (At4g37390), *GH3.3* (At2g23170), *GH3.4*

(At1g59500), *GH3.5* (At4g27260), *GH3.6* (At5g54510), *GH3.9* (At2g47750), *GH3.17* (At1g28130).

RESULTS

Arabidopsis group II GH3s are required for vegetative and reproductive development

Group II GH3 enzymes (*GH3.1*, *GH3.2*, *GH3.3*, *GH3.4*, *GH3.5*, *GH3.6*, *GH3.9* and *GH3.17*) are known to redundantly modulate IAA levels by catalysing the formation of IAA-amino acid conjugates, which are inactive IAA forms (Staswick *et al.*, 2005; Chen *et al.*, 2010; Porco *et al.*, 2016). Due to functional redundancy, single *gh3* mutants often show mild or no phenotypes (Staswick *et al.*, 2005). To bypass such redundancy, we generated a *gh3.1,2,3,4,5,6,9,17* octuple mutant (hereafter referred to as *gh3oct*) by crossing the individual insertional lines (Fig. S1a, Table S1). To verify that *gh3oct* was an octuple knock-out, we performed RT-qPCR analysis. We showed that expression of the full-length *GH3* transcript was completely abolished by the insertions in the *gh3oct* mutant, with just marginal expression of *GH3.9* in reproductive tissues (Fig. S1b). Because the average number of T-DNAs in insertional lines has been estimated to be 2.1 (Wilson-Sanchez *et al.*, 2014), the *gh3oct* mutant genome likely harboured several additional insertions that could have jeopardized our interpretations on the GH3-related functions. To rule out this possibility, we sequenced the *gh3oct* genome and followed a tagged-sequence strategy to map the positions of the insertions. We confirmed the presence of eight insertions within the corresponding *GH3* coding sequences and found two extra insertions at intergenic regions (Fig. S2). Due to their position, the latter were considered unlikely to contribute to the *gh3oct* phenotypes.

We then explored the developmental consequences of a group-II *GH3* knock-out. We found that primary roots from *gh3oct* mutant seedlings were comparable in length to those of the wild type, whereas lateral root density was notably increased (Fig. 1a,b,j,k). Such a root phenotype was observed along the vegetative phase (Fig. 1j,k). The rosettes from *gh3oct* seedlings showed classical high-auxin phenotypes, such as epinastic cotyledons (Fig. 1a,b) and elongated hypocotyls (Fig. 1a,b,l) and petioles (Fig. 1c,d,m). The *gh3oct* mutant also showed photoperiod-independent early flowering (Fig. 1e, S3). Most of the siliques in the *gh3oct* stems did not elongate properly (Fig. 1i), likely causing a delayed inflorescence arrest (Ware *et al.*, 2020), and thus making *gh3oct* plants taller (Fig. 1h). We observed the formation of several aberrant flowers that developed into siliques with unfused valves in the *gh3oct* mutant (Fig. 1f,g), although this phenotype showed incomplete penetrance. Taken together, our data indicate that the *gh3oct* mutant is very likely an octuple GH3 knock-out at the seedling

stage and that functional group II GH3s redundantly control the progress and timing of several developmental processes throughout the plant's life cycle.

IAA metabolism in the absence of functional group II GH3s

Due to the involvement of group II GH3s in IAA metabolism, we explored the concentrations of IAA, the irreversible IAA conjugates IAA-Asp and IAA-Glu, and the IAA inactive forms oxIAA, oxIAA-glc and IAA-glc in shoots and roots from Col-0 and the *gh3oct* mutant. According to the high-auxin phenotypes displayed by *gh3oct* plants (Fig. 1), IAA contents were found to be increased in *gh3oct* seedling shoots (Fig. 2). However, total IAA in root tissues was found at wild type levels (Fig. 2). Despite these modest differences in IAA contents, the roots of *gh3oct* seedlings were hypersensitive to exogenous IAA (Fig. S4). As expected for a total knock-out of the IAA-amido synthetase functions, no detectable IAA-Asp was found in *gh3oct* plants (Fig. 2). Surprisingly, decreased yet still detectable amounts of IAA-Glu were found in shoots and unchanged IAA-Glu levels were detected in roots from *gh3oct* seedlings (Fig. 2). The contents of the oxidative catabolites oxIAA and oxIAA-glc were diminished in *gh3oct* shoots and roots (Fig. 2), whereas no differences in the content of IAA-glc were found (Fig. 2). Overall, our data suggested that *gh3oct* plants have a reduced ability to inactivate IAA and that IAA inactivation, by means of conjugation to Glu, could still occur in the absence of functional group-II GH3 members. Our data additionally indicated a connection between the production of IAA-aa conjugates and IAA oxidative catabolites.

***gh3oct* mutant tolerance to salinity and osmotic stress**

GH3-mediated IAA metabolism has previously been suggested to modulate abiotic stress responses in plants (Park *et al.*, 2007; Du *et al.*, 2012; Kirungu *et al.*, 2019). Therefore, we focused on the involvement of group II GH3s in the response to salinity. When examining transcriptomic datasets from Arabidopsis plants exposed to NaCl, we found that Arabidopsis group II GH3s are responsive to salt treatments (Fig. S5). To corroborate this response under our experimental conditions, we used RT-qPCR to determine the relative expression of each GH3 gene to NaCl in seedling roots. We found all root-expressed group II GH3s to be upregulated upon NaCl application (Fig. 3a), suggesting that these enzymes might play a role in the response to salinity.

Salinity is well-known to cause general arrest in plant growth (Zhao *et al.*, 2020), and a decreased length of the primary root in salt-stressed Arabidopsis plants is documented (Smolko *et al.*, 2021). Thus, we used the growth of the primary root as an indicator of the

tolerance to salinity stress. Compared to Col-0 wild type, *gh3oct* plants were more tolerant to 50 mM NaCl (Fig. 3b,c). The *gh3oct* mutant also grew better than Col-0 at higher NaCl concentrations (Fig. 3c). Besides NaCl, we found that the *gh3oct* mutant exhibited higher tolerance to other osmotic stressors, such as sorbitol and mannitol, over the vegetative phase (Fig. S6). Thus, the data suggested that genetic disruption of the group II *GH3s* confers salinity tolerance, likely as part of a general osmotolerant mechanism.

Tolerance to salinity in the *gh3oct* mutant is related to endogenous IAA content

To investigate the relevance of the elevated endogenous IAA content in *gh3oct* plants for the tolerance to salinity, we co-treated *gh3oct* seedlings with both NaCl and IAA synthesis inhibitors. As shown in Fig. 3d, the higher tolerance to NaCl of the *gh3oct* mutant, in terms of a higher root growth ratio, was abolished by addition of either yucasin or PPBo, which are two inhibitors of YUCCA enzymes that cause a decrease in plant IAA levels by competing with the IAA precursor IPyA (Nishimura *et al.*, 2014; Kakei *et al.*, 2015). To rule out the possibility that the decreased root growth ratio was caused by the effect of the IAA synthesis inhibitors on root growth rather than the tolerance to salinity, we inspected the effects of yucasin and PPBo on the root growth of *gh3oct* seedlings in the absence of NaCl. We found that, at the employed concentrations, none of the inhibitors caused a decrease in the length of the root, whereas both decreased the lateral root density of the *gh3oct* mutant (Fig. 3e,f).

We next explored the contents of IAA and IAA metabolites and the tolerance to salinity of the *gh3* single mutants. IAA levels were found to be comparable to the wild type in all *gh3* mutant shoots and roots, except for *gh3.1*, which showed decreased IAA levels in roots, and *gh3.2*, which showed increased IAA levels in shoots but decreased levels in roots (Fig. S7). The IAA-Asp concentration was slightly decreased in *gh3.3* shoots and *gh3.6* roots, while a reduced concentration of IAA-Glu was detected in *gh3.1* and *gh3.2* roots (Fig. S7). Levels of oxIAA were decreased in *gh3.2* shoots but were comparable to the wild type in the other *gh3* mutant shoots and roots (Fig. S7). Under our experimental conditions, only *gh3.5* plants showed salt-tolerant root growth (Fig. S8a,b). However, different from the *gh3oct* mutant, the tolerance to salinity of *gh3.5* plants was not suppressed by decreasing endogenous IAA levels (Fig. S8c). Taken together, the results indicated that the tolerance to salinity stress of the *gh3oct* mutant is related to their higher endogenous IAA contents.

***gh3oct* mutant tolerance to drought stress**

Salinity and drought are initially perceived by the plant roots as an osmotic stress. Therefore, it is not uncommon for salt-tolerant genotypes to also exhibit a degree of tolerance to drought stress and *vice versa* (Uddin *et al.*, 2016; Lamers *et al.*, 2020). To test whether *gh3oct* plants better tolerated drought, we first grew Col-0 and *gh3oct* seedlings on media plates with a reduced water potential by using polyethylene glycol (PEG) perfusion (Verslues *et al.*, 2006). The root growth ratio of *gh3oct* plants grown on PEG-perfused plates was higher than that of the wild type (Fig. 4a), suggesting that the *gh3oct* mutant was more tolerant to drought stress when grown *in vitro*. Because *gh3.5* plants also showed more tolerant growth under salinity, we investigated the growth ratio of *gh3.5* roots in PEG-perfused media. In contrast to the *gh3oct* mutant, *gh3.5* plants were as sensitive as the wild type to PEG-mediated reduction of the water potential (Fig. S8d).

Finally, we explored whether the drought tolerance of the *gh3oct* mutant could be reproduced when grown on soil. We planted stratified Col-0 and *gh3oct* seeds in well-watered soil pots and then withheld irrigation until the plants manifested drought symptoms. As shown in Fig. 4b, *gh3oct* mutant plants showed milder drought symptoms and a higher survival rate after re-watering (80% in *gh3oct* vs. 53% in Col-0). To rule out that pots containing *gh3oct* plants could retain more water, we compared the weight of well-watered to that of dry pots (before drought symptoms appeared) for both genotypes. As shown in Fig. 4c, we found no differences between Col-0 and *gh3oct* pots. Thus, our results further supported that the *gh3oct* mutant was more tolerant to soil drought.

Hormonal landscape in *gh3oct* plants under salinity

In addition to IAA, group II GH3 members have been shown to conjugate amino acids to other phytohormones, such as jasmonic acid (JA) and salicylic acid (SA) (Gutierrez *et al.*, 2012; Westfall *et al.*, 2016). Given the involvement of GH3s in the response to salinity, we quantified the endogenous contents of these phytohormones in plants grown in control and NaCl-supplemented media. We found higher IAA levels in *gh3oct* mutant seedlings grown at 0 and 75 mM NaCl, whereas the IAA concentration was comparable to Col-0 at 150 mM NaCl (Fig. 5). Levels of JA, the JA bioactive form JA-Ile and SA were similar in Col-0 and *gh3oct* seedlings under both control and salinity conditions (Fig. 5). JA and JA-Ile levels increased in both Col-0 and *gh3oct* plants grown on NaCl-supplemented media (Fig. 5). Because NaCl stress is known to trigger an increase in levels of abscisic acid (ABA) (Simura *et al.*, 2018), we also quantified levels of this phytohormone. ABA content was higher in *gh3oct* seedlings

grown in the control medium, but this difference disappeared in seedlings that germinated under salinity (Fig. 5).

Next, we explored changes in endogenous levels of IAA, JA, JA-Ile, SA and ABA in response to salinity. For that, we transferred Col-0 and *gh3oct* seedlings to media plates supplemented with 150 mM NaCl and quantified the phytohormone concentrations at 6 and 24 h upon exposure to NaCl. IAA levels were higher in *gh3oct* seedlings at all inspected times under control conditions, but they became comparable to those of Col-0 in response to 150 mM NaCl (Fig. S9). This result is in accordance with the similar IAA contents in Col-0 and *gh3oct* seedlings grown at 150 mM NaCl concentration (Fig. 5). Levels of JA and JA-Ile were neither affected by genotype nor by salinity (Fig. S9). Levels of SA were significantly lower in *gh3oct* seedlings after 6 h of exposure to salinity, although this difference vanished after 24 h (Fig. S9). Endogenous levels of ABA prominently increased in both genotypes in response to salinity. However, the ABA content was higher in the *gh3oct* mutant after 24 h of exposure to NaCl (Fig. S9). Because higher ABA levels were also found in *gh3oct* plants grown under control conditions (Fig. 5) and ABA is known to modulate the response to osmotic stresses, e.g. salinity, by mediating stomatal closure (Hedrich & Shabala, 2018), we hypothesized a higher ability of the *gh3oct* mutant to deal with water loss. However, our observations indicated that the water loss rate in *gh3oct* detached rosettes was indistinguishable from that of the wild type (Fig. S10).

In summary, we found altered levels of IAA and ABA under control conditions and of SA and ABA in response to salinity in *gh3oct* mutant plants.

DISCUSSION

IAA metabolism and plant development without functional group II GH3s

Conjugation of IAA to amino acids is an important metabolic pathway regulating levels of this bioactive auxin and is performed by group II GH3 proteins (Staswick *et al.*, 2005). Studies on single *gh3* mutants have demonstrated the relevance of IAA conjugation for plant developmental processes, such as root meristem size control (Di Mambro *et al.*, 2017) and hypocotyl elongation (Zheng *et al.*, 2016). To identify additional roles of the group II GH3s in auxin metabolism and plant development, we generated an octuple group II *GH3* mutant. Our phenotypical analyses of *gh3oct* mutant plants supported redundant involvement of group II GH3s in lateral root, shoot and fruit development, as well as in vegetative-to-reproductive transition. While our data were not sufficient to support a direct causality between IAA metabolism and early flowering or aberrant fruit development in *gh3oct* plants, the observed

increased lateral root density and longer hypocotyls and petioles are well-known high-auxin phenotypes (Sugawara *et al.*, 2009; Gao *et al.*, 2020). Our analyses found that the IAA content was only slightly increased in *gh3oct* shoots but at wild-type levels in roots. This suggests that the high-auxin phenotypes displayed by *gh3oct* plants might arise from highly spatiotemporally localized IAA increases, which do not remarkably alter the total IAA content in shoot and root tissues where other mechanisms of IAA control may be more important.

We previously reported a sextuple *GH3* knockout mutant in which 6 group II *GH3*s (*GH3.1-6*) were disrupted (Porco *et al.*, 2016). No detectable levels of IAA-Asp were found in the *gh3* sextuple mutant, whereas conjugation of IAA to Glu was upregulated. This was likely attributable to the activity of *GH3.17*, which is highly specific for the formation of IAA-Glu (Staswick *et al.*, 2005; Brunoni *et al.*, 2019). Therefore, one might expect that formation of both IAA-Asp and IAA-Glu would be abolished after knocking out all group II *GH3*s. However, the presence of detectable IAA-Glu content in *gh3oct* seedlings, which displayed null expression of group-II *GH3* genes, strongly suggests that additional *GH3* or *GH3*-like enzymes work in IAA inactivation by means of conjugation to glutamate in Arabidopsis.

DAO proteins have been shown to regulate IAA levels by mediating its metabolic inactivation through oxidation (Zhao *et al.*, 2013; Porco *et al.*, 2016; Zhang *et al.*, 2016). Levels of the oxidative IAA catabolite oxIAA are known to rapidly increase in response to endogenous or exogenous IAA increases (Kubes *et al.*, 2012; Pencik *et al.*, 2013). However, despite showing locally increased IAA levels, as inferred from their multiple high-auxin phenotypes and hypersensitivity to IAA, reduced levels of oxIAA and oxIAA-glc were observed in *gh3oct* plants. DAOs and *GH3*s have been suggested to establish parallel and redundant pathways for IAA inactivation, with the *GH3* pathway playing a compensatory function, as supported by increased and decreased conjugate levels in the *DAO1* mutant and overexpressing plants, respectively (Porco *et al.*, 2016; Zhang *et al.*, 2016). This model has recently been challenged by the finding that IAA-Asp is an *in vivo* substrate for *DAO1* oxidase, which additionally accounts for the formation of oxIAA-Asp conjugates (Müller *et al.*, 2021). In a context where DAOs function downstream of group II *GH3*s, the reduced levels of oxIAA and oxIAA-glc in *gh3oct* plants found in the present work suggest that a significant fraction of the plant oxIAA pool derives from oxIAA-aa conjugates.

Cooperative involvement of group II *GH3*s in responses to salinity and drought

Salinity and drought are major constraints on plant growth, and therefore represent serious threats to agriculture and forestry. Data from several studies over the past decade support a

link between auxin and plant responses to salinity (Wang *et al.*, 2009; Liu *et al.*, 2015; Korver *et al.*, 2018; Fu *et al.*, 2019) and drought stress (Zhang *et al.*, 2009; Zhang *et al.*, 2012; Shi *et al.*, 2014; Jung *et al.*, 2015; Zhang *et al.*, 2020). Spatiotemporal auxin distribution and auxin sensing modulate adaptive growth responses to NaCl stress (Wang *et al.*, 2009; Liu *et al.*, 2015). Recently, Aux/IAA auxin co-receptors were shown to be required for plant tolerance to water deficit, thus suggesting a central role for auxin in integrating drought signals to elaborate genetic and physiological responses (Shani *et al.*, 2017). Although the pathways that mediate auxin-mediated responses to salinity and drought are only just being elucidated, it appears that auxin may mediate tolerance to water deprivation by regulating several adaptive responses leading to stress avoidance. These include (i) adaptive root growth (Leftley *et al.*, 2021), (ii) ROS scavenging (Fu *et al.*, 2019), (iii) regulation of stomatal apertures (Salehin *et al.*, 2019), and (iv) crosstalk with ABA and other hormones involved in the responses to salinity and drought (Sun & Li, 2014; Yu *et al.*, 2020; Salvi *et al.*, 2021).

Our work revealed that *gh3oct* mutant plants showed more tolerant growth under salinity, and that this tolerance was related to endogenous IAA levels. Tolerance to salinity was also higher in *gh3.5* mutant plants, although this effect appeared to be unaffected by endogenous IAA levels. GH3.5 is well-known to mediate the conjugation of other plant hormones, including SA and JA (Gutierrez *et al.*, 2012; Westfall *et al.*, 2016), suggesting that the salt-tolerant growth of *gh3.5* roots may rather be related to the SA or JA pathways. The increased root branching of *gh3oct* plants without penalty for primary root growth, produced a more robust root system, which might contribute to the better performance of *gh3oct* plants under water stress conditions.

In addition to root system architecture, the increased tolerance of *gh3oct* plants to salinity and drought might be driven by increased ABA content. ABA is a stress-related hormone whose levels increase very rapidly upon salt and drought stress sensing and is well-known for its direct involvement in the response to osmotic stresses, primarily by mediating stomatal closure (Hedrich & Shabala, 2018), although other stomata-independent mechanisms have been reported (Thalman *et al.*, 2016). The increased ABA content but wild-type water loss rate in *gh3oct* plants grown in control conditions indeed rule out a predisposition for *gh3oct* plants to tolerate water deficit based on lower transpiration, but suggests that other ABA-mediated mechanisms might assist *gh3oct* plants to cope with this stress.

Conclusions

In this work, we present a group-II *GH3* knock-out as a useful genetic tool to research metabolic pathways regulating levels of the main auxin IAA. It has long been accepted that group II GH3 members carry out conjugation of IAA to amino acids. Our metabolic analyses in *gh3oct* plants indicated that additional GH3s or GH3-like proteins might modulate IAA levels by means of conjugation to glutamate. Additionally, our data suggested that the major IAA catabolite oxIAA is, at least in part, produced from the GH3 pathway. Examination of mutant *gh3oct* plants also revealed that plant tolerance to water deprivation stresses, such as salinity and drought, is modulated by redundant auxin conjugation. Identifying as many components as possible of a plant's responses to salinity and drought is important for suggesting targets to guide the design of chemical and/or gene-editing approaches directed to enhance stress tolerance in plants. While the mechanisms behind the auxin-related tolerance to salinity and drought remain to be elucidated, the present work supports strategies based on salinity- and drought-inducible disruption of group II GH3s as a profitable tool to engineer plants that better tolerate water stress.

FIGURE LEGENDS

Figure 1. Phenotypes of the *gh3oct* mutant. (a-e) Col-0 and *gh3oct* plants at (a,b) 7, (c,d) 20 and (e) 23 days after stratification (das). (f,g) Aberrant reproductive structures occasionally formed by *gh3oct* mutant from latest inflorescences. (h) Plant height of Col-0 and *gh3oct* flowering plants. (i) Stems from Col-0 and the *gh3oct* mutant. White arrowheads indicate siliques that did not elongate further in the *gh3oct* mutant. (j,k) Primary root length (j) and lateral root density (k) determined at 5, 7, 10 and 12 das. (l) Hypocotyl length determined at 10 das. (m) Petiole length determined at 15 das. Scale bars indicate (a,b) 2 mm, (c,d) 1 cm, (f,g) 2 mm, and (i) 2 cm. Asterisks indicate values statistically different from Col-0 in a two-tailed Student's *t*-test [$*p<0.05$, $****p<0.0001$; (j,k) $n \geq 17$, (l) $n \geq 83$, and (m) $n = 20$].

Figure 2. Levels of IAA and IAA metabolites in shoots and roots of the *gh3oct* mutant. Concentration (expressed in picomoles per gram of fresh weight) of indole-3-acetic acid (IAA), indole-3-acetyl-aspartate (IAA-Asp), indole-3-acetyl-glutamate (IAA-Glu), 2-oxindole-3-acetic acid (oxIAA), 1-O-(2-oxindole-3-ylacetyl)- β -D-glucose (oxIAA-glc) and 1-O-indole-3-ylacetyl- β -D-glucose (IAA-glc) in shoots and roots from 7-day-old Col-0 (black bars) and *gh3oct* (red bars) seedlings. Error bars indicate standard deviation. Asterisks indicate values significantly different from Col-0 in a two-tailed Student's *t*-test ($*p<0.05$, $***p<0.001$; $n = 5$).

Figure 3. Group II GH3s are upregulated upon salt treatment and the *gh3oct* mutant showed IAA-dependent tolerance to salinity. (a) Transcriptional response of group II GH3 genes from 10-day-old Col-0 roots determined by qRT-PCR 6 hours after transferring to MS media supplemented with 0, 75 or 150 mM NaCl. No expression of GH3.4 and GH3.9 was detected in any sample. Error bars indicate standard deviation. (b) Seedlings from 7-day-old Col-0 and *gh3oct* transferred at day 5 to control MS plates or MS plates supplemented with 50 mM NaCl. (c) Primary root length of Col-0 and *gh3oct* seedlings grown for 7 days at different concentrations of NaCl. (d) Root growth ratio of 4-day-old seedlings transferred for another 3 days to media plates containing 50 mM NaCl, 50 mM NaCl + 20 μ M of the YUCCA inhibitor yucasin or 50 mM NaCl + 1 μ M of the YUCCA inhibitor PPBo (4-phenoxyphenylboronic acid). To calculate the ratio, the length of the root from seedlings grown on NaCl, NaCl + yucasin and NaCl + PPBo plates was divided by the mean root length of seedlings grown on control MS, MS + yucasin and MS + PPBo plates, respectively. (e, f) Root length (e) and lateral root density (f) in 7-day-old *gh3oct* mutant seedlings grown on control MS plates, and on plates supplemented with 20 μ M of yucasin or 1 μ M of PPBo. Asterisks indicate (a) Δ CT values significantly different from MS ($n = 4$), (c,d) values significantly different from Col-0 ($n \geq 11$), (e,f) values significantly different from MS ($n \geq 17$) in a two-tailed Student's *t*-test ($*p<0.05$; $**p<0.01$, $***p<0.001$, $****p<0.0001$).

Figure 4. *gh3oct* plants are tolerant to drought. (a) Root growth ratio of Col-0 and *gh3oct* seedlings in media with lowered water potential. 4-day-old seedlings were transferred to plates perfused with a solution of PEG8000 at the indicated concentration and grown for another 6 days. To calculate the ratio, the root length from seedlings in PEG-containing plates was divided by the mean root length of seedlings grown on control plates (perfused with liquid MS). Asterisks indicate statistical significantly different growth ratios from Col-0 in a two-tailed Student's *t*-test ($***p<0.001$; $n \geq 25$). (b) Tolerance of Col-0 and *gh3oct* plants to drought stress on soil. The image corresponds to 3-week-old plants on dry soil. Survival rate after re-watering is indicated at the bottom and represents the mean percentage from 3 independent experiments with 27, 36 and 23 plants per genotype. (c) Soil pot weight in the drought experiment. Pots were prepared with 40 g of soil mixture, watered and weighed at the beginning of the experiment (initial). Pots were re-weighed when the soil was noticeably dry and before plants started showing drought symptoms (before drought). No significant differences were found between Col-0 and *gh3oct* pots in a two-tailed Student's *t*-test.

Figure 5. Phytohormone levels from Col-0 and the *gh3oct* mutant grown under different NaCl concentrations. Concentrations, expressed in picomoles per gram of fresh weight, of indole-3-acetic acid (IAA), jasmonic acid (JA), jasmonic acid-isoleucine (JA-Ile), salicylic acid (SA) and abscisic acid (ABA) in Col-0 and *gh3oct* seedlings were determined 7 days after germination on plates containing 0, 75 or 150 mM NaCl. Error bars indicate standard deviation. Asterisks indicate values significantly different from the corresponding values of Col-0 in a two-tailed Student's *t*-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; $n = 4$). <LOD: below the limit of detection.

SUPPORTING INFORMATION

Figure S1. The *gh3oct* mutant is an octuple knock-out.

Figure S2. The *gh3oct* mutant harbours two extra T-DNA insertions at intergenic regions.

Figure S3. The *gh3oct* is a photoperiod-independent early flowering mutant.

Figure S4. Root growth in the *gh3oct* mutant is hypersensitive to IAA.

Figure S5. Transcriptional response of group-II *GH3* genes to salinity according to published datasets.

Figure S6. Root growth in the *gh3oct* mutant is tolerant to different osmolytes.

Figure S7. Levels of IAA and IAA metabolites in shoots and roots from group-II *gh3* single mutants.

Figure S8. Salinity and drought tolerance of *gh3* single mutants.

Figure S9. Levels of IAA and stress-related phytohormones in response to salinity in the *gh3oct* mutant.

Figure S10. Col-0 and *gh3oct* plants lose water at the same rate.

Table S1. Insertion lines combined to obtain the *gh3oct* mutant.

Table S2. Primer sets used in this work.

ACKNOWLEDGMENTS

Research in the laboratory of Karin Ljung is supported by grants from the Swedish Foundation for Strategic Research (Vinnova), the Knut and Alice Wallenberg Foundation (KAW), the Swedish research councils VR and Formas. E.M.-B. (JCK-1811) held postdoctoral fellowship from Kempestiftelserna. A.P. and O.N. were financially supported by the Ministry of Education, Youth and Sports of the Czech Republic (European Regional Development Fund-Project “Plants as a tool for sustainable global development” No. CZ.02.1.01/0.0/0.0/16_019/0000827). We also acknowledge the Swedish Metabolomics Centre (<http://www.swedishmetabolomicscentre.se/>) for access to instrumentation.

AUTHOR CONTRIBUTIONS

E.M.-B., R.C.-S. and K.L. conceived and designed the research; E.M.-B. and R.C.-S. performed most of the experiments; J.Š. and A.P. performed hormone analyses; P.S. made the *gh3oct* mutant; R.C.-S. prepared the manuscript draft; E.M.-B. and R.C.-S. wrote the manuscript with input from all authors. This research was supported by funds to K.L.

REFERENCES

- Band LR. 2021.** Auxin fluxes through plasmodesmata. *New Phytol* **231**(5): 1686-1692.
- Bottcher C, Boss PK, Davies C. 2011.** Acyl substrate preferences of an IAA-amido synthetase account for variations in grape (*Vitis vinifera* L.) berry ripening caused by different auxinic compounds indicating the importance of auxin conjugation in plant development. *J Exp Bot* **62**(12): 4267-4280.
- Brunoni F, Collani S, Casanova-Saez R, Simura J, Karady M, Schmid M, Ljung K, Bellini C. 2020.** Conifers exhibit a characteristic inactivation of auxin to maintain tissue homeostasis. *New Phytol* **226**(6): 1753-1765.
- Brunoni F, Collani S, Simura J, Schmid M, Bellini C, Ljung K. 2019.** A bacterial assay for rapid screening of IAA catabolic enzymes. *Plant Methods* **15**: 126.
- Casanova-Saez R, Mateo-Bonmati E, Ljung K. 2021.** Auxin Metabolism in Plants. *Cold Spring Harb Perspect Biol* **13**(3).
- Casanova-Saez R, Voss U. 2019.** Auxin Metabolism Controls Developmental Decisions in Land Plants. *Trends Plant Sci* **24**(8): 741-754.
- Chen Q, Westfall CS, Hicks LM, Wang S, Jez JM. 2010.** Kinetic basis for the conjugation of auxin by a GH3 family indole-acetic acid-amido synthetase. *J Biol Chem* **285**(39): 29780-29786.
- Di Mambro R, De Ruvo M, Pacifici E, Salvi E, Sozzani R, Benfey PN, Busch W, Novak O, Ljung K, Di Paola L, et al. 2017.** Auxin minimum triggers the developmental switch from cell division to cell differentiation in the Arabidopsis root. *Proc Natl Acad Sci U S A* **114**(36): E7641-E7649.
- Du H, Wu N, Fu J, Wang S, Li X, Xiao J, Xiong L. 2012.** A GH3 family member, OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and cold tolerance in rice. *J Exp Bot* **63**(18): 6467-6480.
- Fu Y, Yang Y, Chen S, Ning N, Hu H. 2019.** Arabidopsis IAR4 Modulates Primary Root Growth Under Salt Stress Through ROS-Mediated Modulation of Auxin Distribution. *Front Plant Sci* **10**: 522.
- Gallei M, Luschnig C, Friml J. 2020.** Auxin signalling in growth: Schrodinger's cat out of the bag. *Curr Opin Plant Biol* **53**: 43-49.
- Gao Y, Dai X, Aoi Y, Takebayashi Y, Yang L, Guo X, Zeng Q, Yu H, Kasahara H, Zhao Y. 2020.** Two homologous INDOLE-3-ACETAMIDE (IAM) HYDROLASE genes are required for the auxin effects of IAM in Arabidopsis. *J Genet Genomics* **47**(3): 157-165.
- Gutierrez L, Mongelard G, Flokova K, Pacurar DI, Novak O, Staswick P, Kowalczyk M, Pacurar M, Demailly H, Geiss G, et al. 2012.** Auxin controls Arabidopsis adventitious root initiation by regulating jasmonic acid homeostasis. *Plant Cell* **24**(6): 2515-2527.
- Hammes UZ, Murphy AS, Schwechheimer C. 2021.** Auxin Transporters-A Biochemical View. *Cold Spring Harb Perspect Biol*.
- Hanania U, Velcheva M, Sahar N, Perl A. 2004.** An improved method for isolating high-quality DNA from *Vitis vinifera* nuclei. *Plant Molecular Biology Reporter* **22**(2): 173-177.
- Hedrich R, Shabala S. 2018.** Stomata in a saline world. *Curr Opin Plant Biol* **46**: 87-95.
- Jung H, Lee DK, Choi YD, Kim JK. 2015.** OsIAA6, a member of the rice Aux/IAA gene family, is involved in drought tolerance and tiller outgrowth. *Plant Sci* **236**: 304-312.
- Kai K, Horita J, Wakasa K, Miyagawa H. 2007.** Three oxidative metabolites of indole-3-acetic acid from Arabidopsis thaliana. *Phytochemistry* **68**(12): 1651-1663.

- Kakei Y, Yamazaki C, Suzuki M, Nakamura A, Sato A, Ishida Y, Kikuchi R, Higashi S, Kokudo Y, Ishii T, et al. 2015. Small-molecule auxin inhibitors that target YUCCA are powerful tools for studying auxin function. *Plant J* **84**(4): 827-837.
- Khan S, Stone JM. 2007. Arabidopsis thaliana GH3.9 influences primary root growth. *Planta* **226**(1): 21-34.
- Kirungu JN, Magwanga RO, Lu P, Cai X, Zhou Z, Wang X, Peng R, Wang K, Liu F. 2019. Functional characterization of Gh_A08G1120 (GH3.5) gene reveal their significant role in enhancing drought and salt stress tolerance in cotton. *BMC Genet* **20**(1): 62.
- Kojima M, Kamada-Nobusada T, Komatsu H, Takei K, Kuroha T, Mizutani M, Ashikari M, Ueguchi-Tanaka M, Matsuoka M, Suzuki K, et al. 2009. Highly sensitive and high-throughput analysis of plant hormones using MS-probe modification and liquid chromatography-tandem mass spectrometry: an application for hormone profiling in *Oryza sativa*. *Plant Cell Physiol* **50**(7): 1201-1214.
- Korver RA, Koevoets IT, Testerink C. 2018. Out of Shape During Stress: A Key Role for Auxin. *Trends Plant Sci* **23**(9): 783-793.
- Kowalczyk M, Sandberg G. 2001. Quantitative analysis of indole-3-acetic acid metabolites in Arabidopsis. *Plant Physiol* **127**(4): 1845-1853.
- Kubes M, Yang H, Richter GL, Cheng Y, Mlodzinska E, Wang X, Blakeslee JJ, Carraro N, Petrasek J, Zazimalova E, et al. 2012. The Arabidopsis concentration-dependent influx/efflux transporter ABCB4 regulates cellular auxin levels in the root epidermis. *Plant J* **69**(4): 640-654.
- Lamers J, van der Meer T, Testerink C. 2020. How Plants Sense and Respond to Stressful Environments. *Plant Physiol* **182**(4): 1624-1635.
- LeClere S, Tellez R, Rampey RA, Matsuda SP, Bartel B. 2002. Characterization of a family of IAA-amino acid conjugate hydrolases from Arabidopsis. *J Biol Chem* **277**(23): 20446-20452.
- Leftley N, Banda J, Pandey B, Bennett M, Voss U. 2021. Uncovering How Auxin Optimizes Root Systems Architecture in Response to Environmental Stresses. *Cold Spring Harb Perspect Biol*.
- Liu W, Li RJ, Han TT, Cai W, Fu ZW, Lu YT. 2015. Salt stress reduces root meristem size by nitric oxide-mediated modulation of auxin accumulation and signaling in Arabidopsis. *Plant Physiol* **168**(1): 343-356.
- Ludwig-Muller J, Julke S, Bierfreund NM, Decker EL, Reski R. 2009. Moss (*Physcomitrella patens*) GH3 proteins act in auxin homeostasis. *New Phytol* **181**(2): 323-338.
- Lup SD, Wilson-Sanchez D, Andreu-Sanchez S, Micol JL. 2021. Easymap: A User-Friendly Software Package for Rapid Mapping-by-Sequencing of Point Mutations and Large Insertions. *Front Plant Sci* **12**: 655286.
- Mateo-Bonmati E, Casanova-Saez R, Simura J, Ljung K. 2021. Broadening the roles of UDP-glycosyltransferases in auxin homeostasis and plant development. *New Phytol*.
- Müller K, Dobrev PI, Pěňčík A, Hošek P, Vondráková Z, Filepová R, Malínská K, Brunoni F, Helusová L, Moravec T, et al. 2021. DIOXYGENASE FOR AUXIN OXIDATION 1 catalyzes the oxidation of IAA amino acid conjugates. *Plant Physiology*.
- Nishimura T, Hayashi K, Suzuki H, Gyohda A, Takaoka C, Sakaguchi Y, Matsumoto S, Kasahara H, Sakai T, Kato J, et al. 2014. Yucasin is a potent inhibitor of YUCCA, a key enzyme in auxin biosynthesis. *Plant J* **77**(3): 352-366.
- Novak O, Henykova E, Sairanen I, Kowalczyk M, Pospisil T, Ljung K. 2012. Tissue-specific profiling of the Arabidopsis thaliana auxin metabolome. *Plant J* **72**(3): 523-536.

- Ostin A, Kowalczyk M, Bhalerao RP, Sandberg G. 1998.** Metabolism of indole-3-acetic acid in Arabidopsis. *Plant Physiol* **118**(1): 285-296.
- Park JE, Park JY, Kim YS, Staswick PE, Jeon J, Yun J, Kim SY, Kim J, Lee YH, Park CM. 2007.** GH3-mediated auxin homeostasis links growth regulation with stress adaptation response in Arabidopsis. *J Biol Chem* **282**(13): 10036-10046.
- Pencik A, Casanova-Saez R, Pilarova V, Zukauskaitė A, Pinto R, Micol JL, Ljung K, Novak O. 2018.** Ultra-rapid auxin metabolite profiling for high-throughput mutant screening in Arabidopsis. *J Exp Bot* **69**(10): 2569-2579.
- Pencik A, Rolcik J, Novak O, Magnus V, Bartak P, Buchtik R, Salopek-Sondi B, Strnad M. 2009.** Isolation of novel indole-3-acetic acid conjugates by immunoaffinity extraction. *Talanta* **80**(2): 651-655.
- Pencik A, Simonovik B, Petersson SV, Henykova E, Simon S, Greenham K, Zhang Y, Kowalczyk M, Estelle M, Zazimalova E, et al. 2013.** Regulation of auxin homeostasis and gradients in Arabidopsis roots through the formation of the indole-3-acetic acid catabolite 2-oxindole-3-acetic acid. *Plant Cell* **25**(10): 3858-3870.
- Porco S, Pencik A, Rashed A, Voss U, Casanova-Saez R, Bishopp A, Golebiowska A, Bhosale R, Swarup R, Swarup K, et al. 2016.** Dioxygenase-encoding AtDAO1 gene controls IAA oxidation and homeostasis in Arabidopsis. *Proc Natl Acad Sci U S A* **113**(39): 11016-11021.
- Rampey RA, LeClere S, Kowalczyk M, Ljung K, Sandberg G, Bartel B. 2004.** A family of auxin-conjugate hydrolases that contributes to free indole-3-acetic acid levels during Arabidopsis germination. *Plant Physiol* **135**(2): 978-988.
- Salehin M, Li B, Tang M, Katz E, Song L, Ecker JR, Kliebenstein DJ, Estelle M. 2019.** Auxin-sensitive Aux/IAA proteins mediate drought tolerance in Arabidopsis by regulating glucosinolate levels. *Nat Commun* **10**(1): 4021.
- Salvi P, Manna M, Kaur H, Thakur T, Gandass N, Bhatt D, Muthamilarasan M. 2021.** Phytohormone signaling and crosstalk in regulating drought stress response in plants. *Plant Cell Rep.*
- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, et al. 2012.** Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**(7): 676-682.
- Schmittgen TD, Livak KJ. 2008.** Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* **3**(6): 1101-1108.
- Shani E, Salehin M, Zhang Y, Sanchez SE, Doherty C, Wang R, Mangado CC, Song L, Tal I, Pisanty O, et al. 2017.** Plant Stress Tolerance Requires Auxin-Sensitive Aux/IAA Transcriptional Repressors. *Curr Biol* **27**(3): 437-444.
- Shi H, Chen L, Ye T, Liu X, Ding K, Chan Z. 2014.** Modulation of auxin content in Arabidopsis confers improved drought stress resistance. *Plant Physiol Biochem* **82**: 209-217.
- Simura J, Antoniadi I, Siroka J, Tarkowska D, Strnad M, Ljung K, Novak O. 2018.** Plant Hormonomics: Multiple Phytohormone Profiling by Targeted Metabolomics. *Plant Physiol* **177**(2): 476-489.
- Smolko A, Bauer N, Pavlovic I, Pencik A, Novak O, Salopek-Sondi B. 2021.** Altered Root Growth, Auxin Metabolism and Distribution in Arabidopsis thaliana Exposed to Salt and Osmotic Stress. *Int J Mol Sci* **22**(15).
- Staswick PE, Serban B, Rowe M, Tiryaki I, Maldonado MT, Maldonado MC, Suza W. 2005.** Characterization of an Arabidopsis enzyme family that conjugates amino acids to indole-3-acetic acid. *Plant Cell* **17**(2): 616-627.

- Staswick PE, Tiryaki I, Rowe ML. 2002.** Jasmonate response locus JAR1 and several related Arabidopsis genes encode enzymes of the firefly luciferase superfamily that show activity on jasmonic, salicylic, and indole-3-acetic acids in an assay for adenylation. *Plant Cell* **14**(6): 1405-1415.
- Sugawara S, Hishiyama S, Jikumaru Y, Hanada A, Nishimura T, Koshiba T, Zhao Y, Kamiya Y, Kasahara H. 2009.** Biochemical analyses of indole-3-acetaldoxime-dependent auxin biosynthesis in Arabidopsis. *Proc Natl Acad Sci U S A* **106**(13): 5430-5435.
- Sun J, Li C 2014.** Cross Talk of Signaling Pathways Between ABA and Other Phytohormones. In: Zhang D-P ed. *Abscisic Acid: Metabolism, Transport and Signaling*. Dordrecht: Springer Netherlands, 243-253.
- Takehara S, Sakuraba S, Mikami B, Yoshida H, Yoshimura H, Itoh A, Endo M, Watanabe N, Nagae T, Matsuoka M, et al. 2020.** A common allosteric mechanism regulates homeostatic inactivation of auxin and gibberellin. *Nat Commun* **11**(1): 2143.
- Terol J, Domingo C, Talon M. 2006.** The GH3 family in plants: genome wide analysis in rice and evolutionary history based on EST analysis. *Gene* **371**(2): 279-290.
- Thalmann M, Pazmino D, Seung D, Horrer D, Nigro A, Meier T, Kolling K, Pfeifhofer HW, Zeeman SC, Santelia D. 2016.** Regulation of Leaf Starch Degradation by Abscisic Acid Is Important for Osmotic Stress Tolerance in Plants. *Plant Cell* **28**(8): 1860-1878.
- Uddin MN, Hossain MA, Burritt DJ. 2016.** Salinity and drought stress: Similarities and differences in oxidative responses and cellular redox regulation. In *Water Stress and Crop Plants: A Sustainable Approach*; Ahmad, P., Ed.; John Wiley & Sons: Singapore: 86–101.
- Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu JK. 2006.** Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* **45**(4): 523-539.
- Wang Y, Li K, Li X. 2009.** Auxin redistribution modulates plastic development of root system architecture under salt stress in Arabidopsis thaliana. *J Plant Physiol* **166**(15): 1637-1645.
- Ware A, Walker CH, Simura J, Gonzalez-Suarez P, Ljung K, Bishopp A, Wilson ZA, Bennett T. 2020.** Auxin export from proximal fruits drives arrest in temporally competent inflorescences. *Nat Plants* **6**(6): 699-707.
- Westfall CS, Sherp AM, Zubieta C, Alvarez S, Schraft E, Marcellin R, Ramirez L, Jez JM. 2016.** Arabidopsis thaliana GH3.5 acyl acid amido synthetase mediates metabolic crosstalk in auxin and salicylic acid homeostasis. *Proc Natl Acad Sci U S A* **113**(48): 13917-13922.
- Wilson-Sanchez D, Rubio-Diaz S, Munoz-Viana R, Perez-Perez JM, Jover-Gil S, Ponce MR, Micol JL. 2014.** Leaf phenomics: a systematic reverse genetic screen for Arabidopsis leaf mutants. *Plant J* **79**(5): 878-891.
- Yu Z, Duan X, Luo L, Dai S, Ding Z, Xia G. 2020.** How Plant Hormones Mediate Salt Stress Responses. *Trends Plant Sci* **25**(11): 1117-1130.
- Zhang J, Lin JE, Harris C, Campos Mastrotti Pereira F, Wu F, Blakeslee JJ, Peer WA. 2016.** DAO1 catalyzes temporal and tissue-specific oxidative inactivation of auxin in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* **113**(39): 11010-11015.
- Zhang Q, Li J, Zhang W, Yan S, Wang R, Zhao J, Li Y, Qi Z, Sun Z, Zhu Z. 2012.** The putative auxin efflux carrier OsPIN3t is involved in the drought stress response and drought tolerance. *Plant J* **72**(5): 805-816.

- Zhang SW, Li CH, Cao J, Zhang YC, Zhang SQ, Xia YF, Sun DY, Sun Y. 2009.** Altered architecture and enhanced drought tolerance in rice via the down-regulation of indole-3-acetic acid by TLD1/OsGH3.13 activation. *Plant Physiol* **151**(4): 1889-1901.
- Zhang Y, Li Y, Hassan MJ, Li Z, Peng Y. 2020.** Indole-3-acetic acid improves drought tolerance of white clover via activating auxin, abscisic acid and jasmonic acid related genes and inhibiting senescence genes. *BMC Plant Biol* **20**(1): 150.
- Zhao C, Zhang H, Song C, Zhu J-K, Shabala S. 2020.** Mechanisms of Plant Responses and Adaptation to Soil Salinity. *The Innovation* **1**(1): 100017.
- Zhao Y. 2018.** Essential Roles of Local Auxin Biosynthesis in Plant Development and in Adaptation to Environmental Changes. *Annu Rev Plant Biol* **69**: 417-435.
- Zhao Z, Zhang Y, Liu X, Zhang X, Liu S, Yu X, Ren Y, Zheng X, Zhou K, Jiang L, et al. 2013.** A role for a dioxygenase in auxin metabolism and reproductive development in rice. *Dev Cell* **27**(1): 113-122.
- Zheng Z, Guo Y, Novak O, Chen W, Ljung K, Noel JP, Chory J. 2016.** Local auxin metabolism regulates environment-induced hypocotyl elongation. *Nat Plants* **2**: 16025.

Casanova-Sáez et al. Figure 2

■ Col-0 ■ *gh3oct*

Casanova-Sáez et al. Figure 5