Title:

Temporal transcriptional response of *Candida glabrata* during macrophage infection reveals a multifaceted transcriptional regulator CgXbp1 important for macrophage response and drug resistance.

Maruti Nandan Rai¹, Chirag Parsania¹, Rikky Rai¹, Niranjana Shirgaonkar¹, Kaeling Tan^{1,2}, and Koon Ho Wong^{1,3,4*}

¹Faculty of health sciences, University of Macau, Avenida da Universidade, Taipa, Macau SAR, China.

²Gene Expression, Genomics and Bioinformatics core, Faculty of Health Sciences, University of Macau, Avenida da Universidade, Taipa, Macau SAR of China.

³Institute of Translational Medicine, Faculty of Health Sciences, University of Macau, Avenida da Universidade, Taipa, Macau SAR of China.

⁴MoE Frontiers Science Center for Precision Oncology, University of Macau.

Maruti Nandan Rai - mnrai@illinois.edu

Chirag Parsania - yb57653@connect.um.edu.mo

Rikky Rai – rrail@ufl.edu

Niranjana Shirgaonkar - niranjana.shirgaonkar@gmail.com

Kaeling Tan - kaelington@um.edu.mo

Koon Ho Wong - koonhowong@um.edu.mo

*For correspondence, please contact koonhowong@um.edu.mo.

Abstract

Candida glabrata can thrive inside macrophages and tolerate high levels of azole antifungals. These innate abilities render infections by this human pathogen a clinical challenge. How *C. glabrata* reacts inside macrophages and what is the molecular basis of its drug tolerance are not well understood. Here, we mapped genome-wide RNA polymerase II (RNAPII) occupancy in *C. glabrata* to delineate its transcriptional responses during macrophage infection in high temporal resolution. RNAPII profiles revealed dynamic *C. glabrata* responses to macrophage with genes of specialized pathways activated chronologically at different times of infection. We identified an uncharacterized transcription factor (CgXbp1) important for the chronological macrophage response, survival in macrophages, and virulence. Genome-wide mapping of CgXbp1 direct targets further revealed its multi-faceted functions, regulating not only virulence-related genes but also genes associated with drug resistance. Finally, we showed that CgXbp1 indeed also affects azole resistance. Overall, this work presents a powerful approach for examining host-pathogen interaction and uncovers a novel transcription factor important for *C. glabrata*'s survival in macrophages and drug tolerance.

Introduction

Phagocytes such as macrophages constitute the first line of host immune defence against invading pathogens (Brown, 2011; Erwig and Gow, 2016). The ability to escape or survive phagocytic attacks is fundamental to the virulence of pathogens (Seider *et al*, 2010; Erwig & Gow, 2016). *Candida* species are prominent opportunistic fungal pathogens with an associated mortality rate of ~29-60% among immunocompromised population (Bongomin *et al.*, 2017; Lamothe *et al.*, 2018). *Candida albicans* is responsible for most Candidiasis infections, although recent studies indicate an epidemiological shift in Candidiasis with an upsurge in infections caused by *Candida glabrata* (Benedict *et al*, 2017; Lamothe *et al*, 2018). Relative to other fungal species including *C. albicans*, *C. glabrata* is more resistant to antifungal drugs like fluconazole and can survive and proliferate inside immune cells (Seider *et al.*, 2011; Rai *et al.*, 2012). Thus far, details about how *C. glabrata* survives, adapts and proliferates in phagocytes and the basis for its intrinsically high azole resistance are still not clearly understood.

Genome-wide transcriptomic studies have been performed to map the response of *Candida species* during macrophage infection (Lorenz & Fink, 2001; Rubin-Bejerano *et al*, 2003; Lorenz *et al*, 2004; Kaur *et al*, 2007; Rai *et al*, 2012), but the insights gained into the infection process so far lack temporal resolution, centring mostly on the late stages of the pathogen-host interactions. We reason that the immediate and early pathogen response is pivotal for survival and adaptation in the host, while responses during later stages reflect strategies for growth and proliferation. Therefore, delineating the whole episode of pathogen response, instead of just a snapshot, during infection is fundamental to understanding pathogenesis. However, conventional transcriptomic analysis involving mRNAs are not suitable for dissecting dynamic temporal transcriptional changes, as measurements of mRNA levels are convoluted by transcript stabilities (Tan and Wong, 2019).

Here, we applied the powerful Chromatin Immuno-precipitation followed by Next Generation Sequencing (ChIP-seq) method against elongating RNA Polymerase II (RNAPII) to map *C. glabrata* transcriptional responses during macrophage infection. We show that *C. glabrata* responds to macrophage infection by mounting chronological transcriptional responses. Based on the expression pattern, we identified many candidate transcriptional regulators including a novel transcription factor, CgXbp1, for the macrophage response. Deletion of CgXbp1 led to accelerated transcriptional activation of genes associated with multiple biological processes during interaction with macrophages. We further demonstrate that CgXbp1 is a multifaceted transcription factor directly binding to ~10% of *C. glabrata* genes including many involves in the pathogenesis process and drug resistance. *CgXBPI* deletion resulted in attenuated survival in host macrophages, diminished virulence in the *Galleria mellonella* model of Candidiasis, and elevated resistance to antifungal drug fluconazole. Overall, our work uncovers an important novel transcription factor for *C. glabrata*'s survival in macrophages and antifungal drug resistance.

Results

Mapping high temporal resolution transcriptional responses of *C. glabrata* during macrophage infection

To understand how *C. glabrata* survives macrophage phagocytosis, we applied ChIP-seq against elongating RNAPII in a time-course experiment after 0.5, 2, 4, 6, and 8 h of THP-1 macrophage infection to map genome-wide transcription responses of *C. glabrata* during different stages of THP-1 macrophage infection (Figure 1A). As expected, genes known to be induced by macrophage phagocytosis (e.g. tricarboxylic acid [TCA] cycle, glyoxylate bypass, and iron homeostasis genes (Kaur *et al*, 2007; Rai *et al*, 2012)) had significant RNAPII occupancies at their gene bodies specifically but not at inter-genic regions (Figure 1B, Figure

1-figure supplement 1A). In addition, the ChIP-seq data also revealed temporal gene expression information. For example, the ATP synthesis gene *CgCYC1* was dramatically up-regulated immediately (0.5 h) upon macrophage internalisation, while the TCA cycle gene *CgCIT2* and glyoxylate bypass gene *CgICL1* were induced slightly later at 2 h and their transcription levels decreased subsequently (4-6 h) (Figure 1B). In contrast, an opposite transcription pattern (e.g. gradual increasing and peaking at later stages) was observed for *CgFTR1*, *CgTRR1*, and *CgMT-I*, which are involved in iron uptake, biofilm formation, and sequestration of metal ions, respectively (Figure 1B). Therefore, the RNAPII ChIP-seq approach can effectively capture a high temporal resolution gene expression landscape in *C. glabrata* during macrophage infection.

***C. glabrata* mounts dynamic, temporal and chronological transcription responses during macrophage infection**

Systematic analysis of actively transcribed genes revealed that approximately 30% of *C. glabrata* genes (n = 1,589, Supplementary File 1) were constitutively transcribed (n = 511, Figure 1-figure supplement 1B) or temporally induced (n = 1,078, Figure 1-figure supplement 1C) during macrophage infection. Temporally induced genes were further classified into six groups according to their transcription pattern using *k*-means clustering. The overall transcriptional response was highly diverse with each group exhibiting a unique temporal transcriptional pattern (Figure 1C). Interestingly, while some genes were induced immediately (0.5 h, Group 1, n = 181) upon internalisation by macrophages, transcriptional induction of over 80% of genes (Group 2-6, n = 897) did not happen until later (2-8 h). Besides, their expression patterns were highly variable, illustrating the complex and dynamic nature of *C. glabrata* transcriptional response during macrophage infection.

Gene Ontology (GO) analysis revealed chronological activation of different biological processes during the infection process (Figure 1D, Supplementary File 2). In the immediate response (0.5 h), genes (Group 1, n = 181) were significantly enriched in processes such as adhesion, responses to copper ion and nitrogen compound, positive regulation of nuclear export in response to glucose starvation, lipid oxidation, and ATP synthesis (Figure 1D). This indicated that *C. glabrata* experiences nutrient and energy deprivation immediately upon entry to macrophages (Figure 1E). Alternatively, the induction of ATP biosynthesis genes may reflect a strong demand for energy by *C. glabrata* to deal with the host's attacks and/or to adapt to the host microenvironment. Subsequently (2 h post phagocytosis), *C. glabrata* underwent a major metabolic remodelling presumably to prepare for growth and generate energy, as reflected by the next wave of transcriptional induction for genes (Group 2, n = 171) involved in the TCA cycle, biosynthesis of inosine 5' monophosphate (IMP), carboxylic acid, amino acid, nucleotide, and precursor for metabolite and energy (Figures 1C&D). In addition, cell cycle arrest and DNA damage checkpoint genes (*CgMEC3*, *CgGLC7*, *CAGL0G07271g*, and *CAGL0A04213g*) were also strongly induced at this early stage, and *C. glabrata* cells were indeed arrested at the G1-S phase cell cycle after macrophage engulfment (Figure 1-figure supplement 2). It is noteworthy that many genes and pathways previously shown to be critical for *C. glabrata* virulence (Kaur *et al.*, 2005; Rai *et al.*, 2012; Kasper *et al.*, 2015) such as adherence, response to DNA damage, oxidative stress, autophagy, TCA cycle, amino acid biosynthesis, and iron homeostasis (Figure 1-figure supplement 3A-G), were markedly induced at the early stages (0.5 and 2 h). Therefore, virulence-centric biological processes were among the most immediate *C. glabrata* responses upon macrophage phagocytosis (Figure 1E), implying the importance of the early transcriptional response towards its adaptation and survival in macrophages.

During the next stage of infection (2-4 h), *C. glabrata* continued to actively transcribe genes associated with carbon metabolism, DNA repair and pathogenesis (Group 3, Figures 1C&D), suggesting that the invading pathogen was still trying to achieve metabolic homeostasis and to counter macrophage internal milieu. Genes required for chromatin assembly and modification were also significantly induced at this stage (Figure 1-figure supplement 4A&B), supporting an earlier report about the involvement of chromatin remodelling during the infection process (Rai *et al.*, 2012). Towards the later phase of this stage (4 h), genes for responses to different stresses (e.g. oxidative, chemical stress, and osmolarity) and resistance thermo-tolerance and oxidative stress (e.g. trehalose biosynthesis and pentose phosphate pathway, respectively) become maximally induced (Group 4, Figures 1C&D). The induction of these stress response pathway genes towards the end of metabolic remodelling is somewhat unexpected, as it suggests that phagocytic attacks (e.g. ROS) against *C. glabrata* might not have occurred until the later phase. However, as shown above, the observations that DNA repair and damage response genes were already upregulated at 2 h indicate that cells had already experienced the attacks. These findings collectively suggest that *C. glabrata* elicits a coordinated stage-wise response during infection; first adapting to macrophage nutrient microenvironment before overcoming phagocytic attacks (Figure 1E). Interestingly, a family of sterol uptake genes (known as *TIR* [Tip1-related]) displayed concerted transcription activation at the end of this stage (4 h) (Figure 1-figure supplement 5). In *Saccharomyces cerevisiae*, *TIR* genes are activated and required for growth under anaerobic condition (Abramova *et al.*, 2001). Given that sterols are an essential component of the cell membrane and that ergosterol biosynthesis is an oxygen-dependent process (Joffrion and Cushion, 2010), the up-regulation of the *TIR* genes indicates an experience of oxygen deprivation and a need for sterols (presumably for proliferation) by *C. glabrata*.

Towards the late stage (6-8 h), genes required for biofilm formation, iron homeostasis, both of which play critical roles in the pathogenesis process (Seider *et al*, 2014; Rodrigues *et al*, 2017), became maximal induced (Group 5, Figures 1C&D). As biofilm formation involves cell growth and proliferation, this observation potentially suggests that the cells are preparing for growth, and this is consistent with the concomitant induction of the iron homeostasis genes that are also necessary for proliferation. Altogether, the overall results revealed details into the dynamic stage-wise responses of *C. glabrata* during macrophage infection (Figure 1E).

Identification of potential transcriptional regulators of early temporal response

We next attempted to identify the potential transcriptional regulators for the chronological transcriptional response. Remarkably, more than 25% of *C. glabrata* transcription factor (TF) genes (n = 53) were expressed during macrophage infection (Table 1), with 39 TF genes showing a temporal induction pattern (Figure 2A). Of note, eleven TFs (Aft1, Ap1, Ap5, Haa1, Hap4, Hap5, Msn4, Upc2, Yap3, Yap6 and Yap7) are known to either bind or control some of the macrophage infection-induced genes (Supplementary File 3) as reported by PathoYeasttract (Monteiro *et al*, 2020), providing strong support to the identified TFs being responsible for the observed temporal transcriptional response.

As the early response is likely to have influential effects on infection outcome, we focused on the four candidate TFs in Group 1 (induction at 0.5 h); the genes *CAGL0F00561g*, *CAGL0G02739g*, *CAGL0L03157g* and *CAGL0J04400g* are uncharacterized and annotated as the *Saccharomyces cerevisiae* orthologue of *RPA12*, *XBPI*, *DAL80*, and *HAP3*, respectively. Interestingly, three of the yeast orthologues (*RPA12*, *XBPI*, *DAL80*) have negative roles in transcription (Mai & Breeden, 1997; Marzluf, 1997; Yadav *et al*, 2016). Transcription regulatory network analysis by PathoYeasttract (Monteiro *et al.*, 2020) further showed that the

orthologues of ~35% macrophage infection-induced genes (n = 375 out of 1,078, respectively; Figure 2-figure supplement 1A, Supplementary File 4) are targets of Xbp1 in *S. cerevisiae* including a significant number of TF genes (n = 14; Figure 2-figure supplement 1B). In contrast, a much smaller set of orthologous genes (~ 7%, n = 72, Figure 2-figure supplement 1A) is annotated as being *S. cerevisiae* Hap3 targets, while no information was available on the PathoYeast database (Monteiro *et al.*, 2020) for the other two repressors. These results suggest that the chronological transcriptional response upon macrophage phagocytosis involves the interplays between transcriptional repressors and activators and that the protein encoded by *CAGL0G02739g* (hereafter referred to as *CgXBPI*) likely plays a central role in orchestrating the overall response.

CgXbp1 is crucial for the chronological transcriptional response during macrophage infection

We next deleted the *CgXBPI* gene and analysed the transcriptional response of the *Cgxbp1Δ* mutant to macrophages. RNAPII ChIP-seq time course analysis showed that a similar number of genes were transcribed in the mutant during macrophage infection (1,471 versus 1,589 genes in *Cgxbp1Δ* and wildtype, respectively) (Supplementary File 5) and ~90% of the transcribed genes are common between wildtype and the mutant (Figure 2-figure supplement 1C), suggesting that CgXbp1 has little effect on the overall set of genes transcribed during macrophage infection. Notably, the *Cgxbp1Δ* mutant had a significantly higher number of genes activated at the earliest infection time point (0.5 h, Figure 2B, Supplementary File 5) as compared to wildtype (Figure 1C); e.g., 369 genes showed accelerated expression in the *Cgxbp1Δ* mutant, while 162 and 109 genes had an unchanged or delayed gene expression profile (Figure 2C).

Systematic GO analysis revealed multiple biological processes enriched among the genes with precocious activation in the *Cgxbp1Δ* mutant (0-0.5 h). They include processes like energy generation, chromatin assembly, cellular respiration and metabolism pathways such as TCA cycle, acetate catabolism, and amino acid, carboxylic acid, nucleotide and trehalose biosynthesis (Figure 3-figure supplement 1, Supplementary File 6). On the other hand, cell adhesion, host response and biofilm formation genes, which were up-regulated in wildtype cells during the late infection stage did not happen in the *Cgxbp1Δ* mutant within the 8 h infection duration examined (Figure 3-figure supplement 1).

Given that remodelling of carbon and nitrogen metabolism is crucial for the survival of fungal pathogens inside phagocytic cells (Lorenz and Fink, 2001; Rubin-Bejerano *et al.*, 2003; Rai *et al.*, 2012; Seider *et al.*, 2014), we closely examined the expression patterns of TCA cycle and amino acid biosynthesis genes in wildtype and the *Cgxbp1Δ* mutant during macrophage infection. In wildtype cells, most genes of these two metabolic pathways were temporally induced with the maximal induction at 2h (Figures 3A&B). By contrast, the induction of these genes was advanced to 0.5 h (Figures 3A&B) and their overall expression levels were significantly higher (1.5 to 14.6 folds) in the mutant compared to wildtype, suggesting that CgXbp1 negatively regulates their expression (i.e., acting as a repressor) upon macrophage infection. Overall, the above results demonstrate that CgXbp1 is critical for the chronological transcriptional response of *C. glabrata* during macrophage infection.

Loss of *CgXBPI* diminishes *C. glabrata* proliferation in human macrophages and attenuates virulence in the *Galleria mellonella* model of candidiasis

To examine if the altered transcriptional response in the *Cgxbp1Δ* mutant affects the survival of *C. glabrata* cells in macrophages, we compared the ability of wildtype and the *Cgxbp1Δ* mutant to survive in THP-1 macrophages. PMA-differentiated THP-1 macrophages

were infected by wildtype and *Cgxbp1*Δ cells, and colony forming unit (CFU) assay was performed to determine the number of viable phagocytosed *C. glabrata* cells at 2, 8, and 24 hours post macrophage infection. No significant difference in CFUs between wildtype and *Cgxbp1*Δ cells was observed at 2 h (Figure 4-figure supplement 1), suggesting similar phagocytosis efficiency of THP-1 macrophages for the two strains. At 8 and 24 h post-infection, wildtype cells exhibited ~ 1.6 and 5.1-fold increase in CFUs compared to that at 2h. Although the *Cgxbp1*Δ mutant was able to proliferate inside macrophages, it displayed significantly lower CFUs (~20%) at both time points (1.3 and 3.9-fold) (Figure 4A). The reductions were rescued in the *Cgxbp1-pXBP1* complemented strain (Figure 4A). These results indicate that CgXbp1 is important for *C. glabrata* proliferation within macrophages.

We next examined the virulence of the wildtype and *Cgxbp1*Δ strains using the *Galleria mellonella* model of *Candida* infection (Jacobsen, 2014). We infected *G. mellonella* larvae with the wildtype, *Cgxbp1*Δ, and complemented strains, and monitored the morbidity and mortality of infected larvae over seven days. Although worms injected with wildtype or *Cgxbp1*Δ *C. glabrata* cells (but not phosphate buffered saline [PBS]) turned dark grey within 4-6 h of infection due to melanin formation, which is a moth response to *C. glabrata* infection, and eventually died (Figure 4B), larvae injected with *Cgxbp1*Δ cells have a consistently slower mortality rate by ~20-30% compared to larvae infected by wildtype cells, suggesting that the loss of Xbp1 function attenuated the virulence (Figure 4B). The attenuated virulence was rescued in the complemented strain (Figure 4B). Therefore, CgXbp1 is important for the survival of *C. glabrata* in human macrophages and virulence in the *in vivo* infection model.

CgXbp1 targets the hierarchy of gene regulatory networks of diverse biological processes

The importance of CgXbp1 towards *C. glabrata*'s response during macrophage and *in vivo* infection models prompted us to characterize its direct genome-wide roles in more details. We attempted ChIP-seq analysis against CgXbp1^{MYC} during macrophage infection but failed to obtain consistent high-quality binding profiles from biological repeats (Pearson correlation $r = 0.63$), presumably due to dynamic responses of *C. glabrata* within the ever-changing macrophage microenvironment during macrophage infection. We, therefore, turned to a defined laboratory condition for the characterization. In *S. cerevisiae*, Xbp1 (ScXbp1) encodes a global transcription repressor that regulates 15% of genes during quiescence (Mai and Breeden, 1997; Miles *et al.*, 2013). Pairwise amino acid sequence alignment showed reasonable conservation between ScXbp1 and CgXbp1, especially at the DNA binding domain and a few small scattered regions (Figure 5-figure supplement 1). Western blot analysis demonstrated that CgXbp1^{MYC} protein expression was significantly induced during macrophage infection (Figure 5A) as well as after 3 and 4 days of growth during which cells have already entered the stationary phase (Figure 5B). The sequence conservation and expression pattern indicate that CgXbp1 also plays roles during quiescence in addition to macrophage response.

Next, we set out to identify CgXbp1^{MYC} genome-wide targets using ChIP-seq during the quiescence phase (day 4 culture). Unlike the attempt for macrophage infection, ChIP-seq against CgXbp1^{MYC} during quiescence produced highly correlated biological replicates (Pearson correlation $r = 0.95$) (Figure 5-figure supplement 2A&B). Distinct CgXbp1^{MYC} bindings were observed throughout the genome, while the input DNA control exhibited only background signals (Figures 5C&D). MACS2 analysis (Zhang *et al.*, 2008; Feng *et al.*, 2012) identified 519 CgXbp1^{MYC} binding sites with 420 located at the promoter (i.e., within 2 kb upstream of the translation start site) of 384 annotated genes (Figure 5E, Supplementary Files 7&8). Two over-represented DNA binding motifs (STVCN₇TCT [where S represents G or C,

V represents T or C] and TCGAG) were identified by *de novo* motif discovery using MEME suite (Bailey *et al.*, 2009) (Figure 5F). While the TCGAG motif is similar to the consensus recognition sequence of *S. cerevisiae* Xbp1 ([TCGA], Mai & Breeden, 1997), the STVCN₇TCT sequence appears to be composed of two halves with the second half showing some resemblance to the DNA binding motif of *S. cerevisiae* Azf1 (Figure 5-figure supplement 3A&B). Interestingly, the two motifs have different occurrence among the target promoters bound by CgXbp1^{MYC} with the STVCN₇TCT motif occurring approximately three times more frequent than the TCGAG sequence (e.g., n = 233 versus n = 82, respectively) (Figure 5F, Supplementary File 9).

The direct functions of CgXbp1 were determined using GO analysis and were significantly associated with major biological processes important for host infection such as “transport”, “response to stress”, “carbon and nitrogen metabolism”, and “biofilm formation” (Table 2, Supplementary File 10). These functions are consistent with the above findings that CgXbp1 is important for *C. glabrata* response and survival in macrophages. More importantly, CgXbp1^{MYC} bindings were found at ~10% of transcription regulators (n = 57 out of 623 genes annotated as transcription regulators), including eight regulators of various stresses (e.g., oxidative stress, osmotic stress, nutrient, pH, acetate, chemical and salts) and ten TF genes involved in carbon catabolite regulation (Figure 5G, Supplementary File 11). Therefore, CgXbp1 exerts its controls over key biological processes through regulating the hierarchy of gene regulatory networks.

CgXbp1 is a negative regulator of fluconazole resistance

Several CgXbp1^{MYC} bound genes (*CgPDH1*, *CgPDR13*, *CgQDR2*, *CgAQRI* and *CgTPO1*; Figure 6A, Supplementary File 10) are important for *C. glabrata* drug resistance (Hallstrom *et al.*, 1998; Miyazaki *et al.*, 1998; Costa *et al.*, 2016, 2013; Pais *et al.*, 2016).

Therefore, we examined whether CgXbp1 affects the resistance of *C. glabrata* to the antifungal fluconazole. Serial dilution spotting assay on solid media showed that *Cgxbp1Δ* mutant had higher resistance to fluconazole compared to wildtype (Figure 6B), and the resistance was restored to the wildtype level in the complemented strain (Figure 6B). Importantly, the altered resistance is not due to an intrinsic difference in growth rate between the two strains, as demonstrated by their indistinguishable growth rates in the absence of drug in liquid media (Figure 6C). However, in presence of fluconazole (64 μg/mL), the *Cgxbp1Δ* mutant was able to grow faster and to a higher density than wildtype (Figure 6C). It is interesting to note that a biphasic growth curve was observed for both strains in the presence of fluconazole, suggesting the existence of two populations of cells (sensitive versus resistant). Consistently, we also noted from the spotting assay relatively more resistant colonies in the *Cgxbp1Δ* mutant as compared to wildtype. To confirm this, we performed a CFU assay by plating an equal number of exponentially growing wildtype, *Cgxbp1Δ* mutant and complemented cells on YPD medium with or without fluconazole (64 μg/mL). The *Cgxbp1Δ* mutant displayed ~8-fold higher CFUs on fluconazole compared to that of wildtype and the complemented strain (Figure 6D), demonstrating the loss of CgXbp1 function led to a larger population of resistant cells. In conclusion, these results demonstrate that CgXbp1 is a negative regulator of fluconazole resistance in *C. glabrata*.

Discussion

C. glabrata is well known for its abilities to withstand azole antifungal drugs and to survive and grow inside phagocytic immune cells (Kaur *et al.*, 2005; Rai *et al.*, 2012). Through mapping genome-wide RNAPII occupancy, this work delineates the temporal transcriptional response of *C. glabrata* upon entry into macrophages, offering insights into the events occurring at different stages of macrophage infection. Our result reveals that ~30%

of *C. glabrata* genes are transcribed during the adaptation, survival and growth inside the alien macrophage microenvironments. At the most immediate response (0-0.5 h), *C. glabrata* activates adherence related genes to initiate adhesion to host surfaces. Concurrently, *C. glabrata* elicits specific responses to the nutrient-limiting microenvironment inside macrophages. This immediate response was followed by *C. glabrata* efforts to deal with oxidative and DNA-damage stresses (0-2 h), and at this time the phagocytosed *C. glabrata* cells were arrested at the G1-S phase of the cell cycle (Figure 1-figure supplement 2B). Subsequently (2-4 h), *C. glabrata* undergoes transcriptional remodeling to adjust its carbon metabolism, presumably to generate energy for future challenges, growth and/or proliferation. Consistently, processes necessary for growth such as global transcription, ribosome biogenesis and copper and iron ion homeostasis were activated around this time (2-6 h). These transcriptional activities indicate that *C. glabrata* are well-adapted to the macrophage microenvironment at this stage of infection. Lastly (8 h), *C. glabrata* induced genes associated with cell proliferation and biofilm formation, implying that they have overcome macrophage attacks and are ready to grow and divide.

It is noteworthy that despite sensing starvation within the first 0.5 h upon macrophage engulfment, *C. glabrata* does not activate alternate carbon catabolic pathways until 2 h. In addition, gene expression and translation-related genes show the lowest transcription levels (i.e., RNAPII occupancy) at this immediate stage (0.5 h) relative to the other time points (Group 6 genes in Figures 1C&D), indicating global suppression of gene expression in *C. glabrata* upon macrophage phagocytosis. A recent study showed that the fungal pathogen *Cryptococcus neoformans* also down-regulate translation during exposure to oxidative stress and suggested that translation suppression may facilitate the degradation of irrelevant transcripts during stress to facilitate proper gene expression and survival (Leipheimer *et al.*, 2019). A similar strategy may be employed by *C. glabrata* in response to the stresses

experienced upon macrophage phagocytosis. Another non-mutually exclusive possibility is that the suppression of gene expression helps *C. glabrata* to reserve energy and resources for coping with the hostile, nutrient-limiting macrophage environment.

Transcriptional responses are determined by the overall TFs activities in a cell. The RNAPII profiles revealed a panel of 53 TF genes expressed during macrophage infection. In particular, 39 of them were temporally induced and are promising candidate regulators for the stage-specific transcription responses. Of note, several uncharacterized TFs (CgXbp1, CgDal80 and CgRpa12) were strongly activated at the earliest infection stage. In *S. cerevisiae*, the orthologues of these TFs play negative regulatory roles (i.e., transcriptional repressors (Mai and Breeden, 1997; Marzluf, 1997; Yadav *et al.*, 2016)), suggesting the importance of transcriptional repression in shaping the overall transcriptional response to macrophages (Figure 7). This was confirmed by the precocious transcriptional activation of a large number of genes in the *Cgxbp1Δ* mutant during macrophage infection. In addition, the ChIP-seq experiment revealed that CgXbp1 directly binds to the promoter of many TFs including 10 carbon catabolite regulators (Figure 5G, Supplementary File 11), suggesting that CgXbp1 indirectly represses the activation of many gene regulatory networks. This probably explains the delayed activation of the carbon catabolic pathway genes. Therefore, CgXbp1 exerts two-fold regulation; directly controlling downstream effector genes and indirectly affecting gene networks of diverse pathways via their hierarchies (i.e., TFs). Our overall findings suggest a regulatory model in which global transcriptional repression is established at the early infection stage to withhold transcriptional activation of certain genes whose functions are only required at later stages (Figure 7).

It is noteworthy that the *S. cerevisiae* orthologues of the other two uncharacterized proteins CgDal80 and CgRpa12 negatively regulate genes involved in nitrogen and lipid metabolism, respectively (Hofman-Bang, 1999; Yadav *et al.*, 2016). We postulate that these

repressors also act in the same fashion as CgXbp1 to delay induction of other groups of genes (such as nitrogen and lipid metabolism genes) that are not necessary for the immediate response but are required at later stages of macrophage infection and/or for proliferation. Repression may be relieved by protein turnover of the repressors and/or through enhanced activation by transcriptional activators. Therefore, the interplays between transcriptional activators and repressors (Figure 7) are crucial in shaping the dynamic transcriptional response during macrophage infection.

An attempt to identify CgXbp1 genome-wide binding targets during macrophage infection was made, however, we only managed to obtain one good quality dataset. Nevertheless, analysis of the CgXbp1 ChIP-seq result during quiescence (4-day-old culture) showed that CgXbp1 targets ~10% of all *C. glabrata* genes. Among the CgXbp1 binding sites, the motifs TCGAG and STVCN₇TCT were significantly enriched. The former motif is similar to the recognition motif of ScXbp1 (TCGA), which was identified through promoter analysis and *in vitro* gel retardation and footprint experiments using recombinant ScXbp1 (Mai and Breeden, 1997; Miles *et al.*, 2013), and is likely bound by CgXbp1 as a monomer or homo-dimer. Interestingly, the latter motif (STVCN₇TCT) was found at a higher frequency (~3 fold) than the common TCGAG motif from the CgXbp1^{MYC} binding sites, suggesting that CgXbp1 can also form a dimer with another transcription factor that recognizes the STVCN₇TCT sequence and that this hetero-dimer controls a larger number of genes than by CgXbp1 alone.

The CgXbp1^{MYC} ChIP-seq data also confirms the transcriptional phenotype of the *Cgxbp1Δ* mutant demonstrating that CgXbp1 is a pivotal regulator of many processes that contributes to the survival inside phagocytic cells (Lorenz and Fink, 2001; Rubin-Bejerano *et al.*, 2003; Rai *et al.*, 2012; Seider *et al.*, 2014). In addition, the data further revealed CgXbp1 as an important regulator of drug resistance, linking *C. glabrata* macrophage response and

drug resistance. Therefore, this work uncovers a multifaceted transcriptional regulator important for the dynamic responses during macrophage infection and antifungal drug resistance in the human fungal pathogen *C. glabrata*.

Methods

Culture conditions for *C. glabrata* and THP-1 macrophages

C. glabrata strain BG2 was used as the wildtype in all experiments. A single colony of *C. glabrata* strains were cultured overnight (14-16 h) in YPD medium at 30°C and 200 rpm in a shaker incubator. To obtain quiescence *C. glabrata* cells, cultures were grown in YPD medium at 30°C and 200 rpm for four days. The THP-1 cell line was obtained from ATCC (TIB 202). THP-1 cells were grown in RPMI medium supplemented with 20 mM glutamine, antibiotic (penicillin-streptomycin, 1X), and 10% heat-denatured serum at 37°C with 5% CO₂ in a cell culture incubator.

C. glabrata infection of macrophages for RNAPII ChIP-seq

Macrophage infection assays were done as described previously (Rai *et al.*, 2013). THP-1 monocytes were grown till 80% confluence, harvested, and re-suspended in RPMI medium at a cell density of 1 x 10⁶ cells/mL. For macrophage differentiation, Phorbol-12-myristate 13-acetate (PMA) was added to the THP-1 monocytes at a final concentration of 16 nM. Approximately 10 million cells were seeded in 100 mm culture dishes and incubated for 12 hours at 37°C with 5% CO₂ in a cell culture incubator. Subsequently, the culture medium was replaced with fresh pre-warmed complete RPMI medium to remove PMA, and cells were allowed to recover in the absence of PMA for 12 hours. Macrophage differentiation and adherence were confirmed under the microscope. Overnight grown *C. glabrata* cells were harvested, washed with PBS and finally suspended in complete RPMI medium at a density of

10⁸ yeasts/ml. To infect THP-1 macrophages, 500 µL yeast cell suspension (5x10⁷ yeast cells) was added to each culture dish containing differentiated THP-1 macrophages at a MOI of 5:1. Post 0.5 h macrophage infection, THP-1 macrophages were crosslinked using formaldehyde at a final concentration of 1% for 20 minutes before 1.5 mL of 2.5 M glycine (a final concentration of 320 nM) was added to stop the crosslinking reaction. For the remaining time points (2 h, 4 h, 6 h and 8 h), culture dishes were washed gently with PBS three times to remove non-phagocytosed yeast cells, and the medium was replaced with fresh pre-warmed RPMI medium. The infected culture was further incubated until the indicated infection times before formaldehyde crosslinking as described above, infected macrophage cultures were harvested and washed three times with ice-cold TBS before storing in -80°C freezer till chromatin extraction.

ChIP and Illumina sequencing library preparation

Chromatin was prepared using a previously described protocol (Fan et al., 2008) with modifications. Briefly, the infected macrophage cell pellet was resuspended in 400 µL FA lysis buffer and 10 µL of 100 mM PMSF solution in the presence of 100 µL equivalent zirconium beads and lysed using six 3-min cycles at maximum speed in a Bullet Blender® (Next Advance) homogeniser with at least 3 min of cooling on ice in between each cycle. Cell lysate was transferred to a new 1.5 mL tube and centrifuged at 2500 g for 5 min in a microcentrifuge. The supernatant was discarded, and the resultant pellet was re-suspended in 500 µL FA lysis buffer, and then transferred to a 2 mL screw-cap tube. Sonication was carried to shear the crosslinked chromatin (cycles of 10 sec on and 15 sec off sonication for a total of 30 min sonication time), and chromatin was stored in the -80 °C freezer until use. Chromatin immuno-precipitation was carried out using 2 µL of a commercially available anti-RNA polymerase II subunit B1 phospho-CTD Ser-5 antibody (Millipore, clone 3E8, cat.

no. 04-1572) for RNAP II, and anti-MYC tag antibody (Santa Cruz, cat. no. 9E10) for CgXbp1^{MYC}. The sample was gently mixed on an end-to-end rotator at room temperature for 1.5 h, and 10 µL of packed protein A sepharose beads (GE Healthcare cat. no. 17-0618-01) were then added. The mixture was further incubated at room temperature for another 1.5 h with gentle mixing. Immuno-precipitated material was washed twice with FA lysis buffer (150 mM NaCl), and once with FA lysis buffer (500 mM NaCl), LiCl wash buffer and TE buffer before elution in 100 µL of elution buffer, as described previously (Wong and Struhl, 2011). Eluted DNA was decrosslinked at 65°C overnight and purified using the QIAGEN PCR purification kit (cat. no. 28104). Sequencing library was generated using a multiplex Illumina sequencing protocol (Wong *et al.*, 2013) and sequenced using the Illumina HiSeq2500 platform at the Genomics and Single Cells Analysis Core facility at the University of Macau.

Bioinformatics and ChIP-seq data analyses

Raw fastq sequences were quality-checked using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) aligned against the *C. glabrata* reference genome (CBS138_s02-m07-r06) using bowtie2 (Langmead and Salzberg, 2012). To visualise the ChIP-seq data on the IGV (integrated genome viewer (Thorvaldsdóttir *et al.*, 2012)), aligned reads were processed by MACS2 (Zhang *et al.*, 2008), and BigWig files were generated using ‘bedSort’ and ‘bedGraphToBigWig’ commands from UCSC Kent utils (Kent *et al.*, 2010). Samtools (version 1.9) was used to index the resultant BAM file and check for alignment statistics. For RNAPII ChIP-seq analysis, elongating RNAPII occupancy was measured by first counting the number of reads over the gene body for all annotated genes (n = 5,311) and then normalising to gene length and sequencing depth using an in-house Perl script, and was expressed as normalised RNAPII ChIP-seq read counts. RNAPII occupancies

for all genes were ranked from high to low as shown in Figure 1-figure supplement 1, and manually inspected on the IGV to empirically determine a filtering cut-off that can reliably identify genes with significant and true RNAPII ChIP-seq signals. The normalised RNAPII ChIP-seq read counts values ≥ 12 and ≥ 25 were determined for wildtype and the *Cgxbp1* Δ mutant, respectively. These values are approximately three times higher than the ChIP-seq signal at background regions (3.2 and 7.0 for wildtype and the *Cgxbp1* Δ mutant, respectively). To ensure that lowly expressed but transcriptionally induced genes were not missed, we searched for genes with the high standard deviation among RNAPII binding signals across the five time points and empirically determined a cut-off ($SD \geq 2.25$ and ≥ 4.00 for wildtype and the *Cgxbp1* Δ mutant, respectively) that includes most, if not all, genes with significant active transcription and/or changes in its level across the time course. This standard deviation-based approach identified 68 and 38 additional genes for wildtype and the *Cgxbp1* Δ mutant, respectively. The lists of shortlisted genes for wildtype and the *Cgxbp1* Δ strains are given in Supplementary Files 1 and 5, respectively. Heatmap, k-means clustering and correlation plots were generated using an online tool FungiExpresZ (<https://cparsania.shinyapps.io/FungiExpresZ/>). GO-term enrichment, and GO slim mapping analyses were performed on the *Candida* genome database (Skrzypek *et al.*, 2017) (<http://www.candidagenome.org/GOContents.shtml>) and FungiDB (Stajich *et al.*, 2012) (<https://fungidb.org/fungidb/>). Transcription regulatory networks between transcription factors and their target genes (Figure 2-figure supplement 1A&B) were generated by the ‘Rank by TF’ function of PathoYeasttract (Monteiro *et al.*, 2020) (<http://pathoyeasttract.org/cglabrata/formrankbytf.php>) using published regulatory information of *S. cerevisiae*. For CgXbp1 ChIP-seq analysis, peak calling was done by MACS2 (Feng *et al.*, 2012) using the parameters [macs2 pileup --extsize 200] and then normalized [macs2 bdgopt]. Peaks obtained from MACS2 were assigned to the nearest gene

promoter using an in-house script. ChIP signal intensity at the 200 bp flanking regions of the peak summit from both replicates was used to determine correlation between the biological replicates (Figure 1-figure supplement 6, Figure 5-figure supplement 2).

Generation of the CgXbp1^{MYC}, Cgxbp1Δ and Cgxbp1Δ-pXBPI complemented strains

CgXbp1^{MYC} strain was generated as described previously (Qin *et al.*, 2019). Briefly, a transformation construct was generated using 1kb of 5' and 3' fragments, flanking the stop codon of *CAGL0G02739g* gene, with a 'MYC-hph' cassette between the two fragments. The 5' fragment for CgXbp1^{MYC} strain was amplified using primers 5'-ATATCGAATTCCTGCAGCCCTCCATGGTACATTGCAAAAC-3', and 5'-TTAATTAACCCGGGGATCCGCACATTCTCTTGAAGATGGG-3' from *CAGL0G02739g* gene, and 3' fragment was same as for *Cgxbp1Δ* mutant. Hygromycin resistant yeast colonies were selected and tagging was confirmed PCR and Sanger sequencing. To create the *Cgxbp1Δ* mutant, 1 kb of 5' and 3' flanking regions of the *CAGL0G02739g* gene were amplified using PCR with the primers 5'-ATATCGAATTCCTGCAGCCCGGCCAACCCCACTTCGAGGA-3' and 5'-TTAATTAACCCGGGGATCCGTTAGTGATTTTGTAGTATGG-3' for the 5' flanking region and 5'-GTTTAAACGAGCTCGAATTCTCAAACATAATATAGTCATC-3' and 5'-CTAGAACTAGTGGATCCCCCGAGAAGTTTTGGGTTGTACG-3' for the 3' flanking region. A transformation construct was created as described previously (Qin *et al.*, 2019) using the 'hph' cassette encoding hygromycin resistance as the selectable marker and used to transform the *C. glabrata* wildtype strain (BG2). Hygromycin resistant yeast colonies were checked for deletion of the *CgXBPI* gene using PCR with the primers from gene internal regions (5'-TGGTGCTTTGGACGCTACAT-3' & 5'-TCATCGCAAAGCAATTGGACA-3'). To generate the complemented strain, *CgXBPI* ORF was first amplified using the

forward (5'- GAATTCATGAGACTCACAGACTCGCCGCT-3') and reverse (5'- GTCGACTTACACATTCTCTTGAAGATGGGT-3') primers from *C. glabrata* genomic DNA, digested with *EcoRI* and *SalI*, and cloned between *EcoRI* and *SalI* restriction sites of a CEN/ARS episomal plasmid, pCN-PDC1. The resultant plasmid, pXBP1, carrying *CgXBP1* ORF was transformed into *Cgxbp1Δ* mutant, and the resultant transformed complemented strain was selected on YPD plates carrying NAT (100μg/mL).

Cell cycle analysis of intracellular *C. glabrata* cells in THP-1 macrophages

For cell cycle analyses, THP-1 macrophages were infected with *C. glabrata* cells in 24-well cell culture plate as described above. In control wells, we inoculated an equal number of *C. glabrata* cells to RPMI medium. Post 2 h incubation, *C. glabrata* cells were harvested and washed twice with 1 mL PBS. Next, harvested cells were fixed by re-suspending them in 1 mL of 70% ethanol, followed by incubation at room temperature on a rotator for 60 minutes. Fixed cells were pelleted and re-suspended in 1 mL of 50 mM sodium citrate (pH 7.0), and were sonicated for 15 seconds at 30% power to re-suspend cell aggregates. Subsequently, samples were treated with RNase cocktail (0.3 μL, Ambion cat. no. AM2286) at 37°C for 1 h to remove RNA, washed with PBS, and stained with propidium iodide (PI) for 1 hour. Cells were then passed through a 40 μm membrane filter and were analysed on the BD Accuri C6 flow cytometer (excitation: 488nm Laser, filter: 585/40, and detector: FL2).

***C. glabrata* infection of macrophages for determining viability using colony forming unit assay**

Macrophage fungi infection assays were done as described earlier (Rai *et al.*, 2013). To prepare macrophages for infection assay, THP-1 monocytes were grown till 80% confluence, harvested, and resuspended to a cell density of 10⁶ cells/ml in complete RPMI

medium. Phorbol-13-myristyl-acetate (PMA) was added to the cell suspension to 16 nM final concentration, mixed well, and 1 million cells were seeded in each well of a 24 well cell culture plate. Cells were incubated for 12 hours in a cell culture incubator, the medium was replaced with fresh pre-warmed complete RPMI medium, cells were allowed to recover from PMA stress for 12 hours. Macrophage differentiation and adherence were confirmed under the microscope. Overnight grown *C. glabrata* cells were harvested, washed with PBS and adjusted to 0.1 OD⁶⁰⁰ and resuspended in complete RPMI medium. 100 µL yeast cell suspension was added to each well of the 24 well culture plate containing differentiated THP-1 macrophages. Post 2-hour co-incubation, wells were washed three times with PBS to remove non-phagocytosed yeast cells and the medium was replaced. At indicated time post-infection, the supernatant was aspirated out from the wells, and macrophages were lysed in sterile water and incubated for 5 minutes for lysing the macrophages. Lysates containing fungal cells were collected, diluted appropriately in PBS, and plated on YPD mediums. Plates were incubated for two days at 30°C and colonies were counted after 48 hours. The viability of *C. glabrata* cells was determined by comparing the colony forming units.

***Galleria mellonella* infection assay for virulence analyses**

Indicated *C. glabrata* strains were grown in YPD medium overnight, washed with PBS thrice, and resuspended in PBS to a final cell density of 10⁸ cells/ml. Next, 20 µL of this cell suspension carrying 2 x 10⁶ *C. glabrata* cells were used to infect *G. mellonella* larvae. The infection was carried out three independent times, each on 16 to 20 larvae. An equal volume of PBS was injected into the control set of larvae. Infected larvae were transferred to a 37°C incubator, and monitored for melanin formation, morbidity and mortality for the next seven days at every 24 hours. The number of live and dead larvae was noted for seven days, and the percentage of *G. mellonella* larvae survival was calculated.

Serial dilution spotting assay

C. glabrata strains were grown in YPD medium for 14-16 h at 30°C under continuous shaking at 200 rpm. Cells were harvested from 1 ml culture, washed with PBS, and were diluted to an OD₆₀₀ of 1. Next, five ten-fold serial dilutions were prepared from an initial culture of 1 OD₆₀₀. Subsequently, 3 µL of each dilution was spotted on YPD plates with or without fluconazole (32 & 64 µg/mL). Plates were incubated at 30°C and images were captured after 2-8 days of incubation.

Growth curve analyses

A single colony of the indicated strains was inoculated to liquid YPD medium and grown for 14-16 h. The overnight grown culture was used to inoculate to YPD medium with or without 64 µg/mL fluconazole at an initial OD₆₀₀ of 0.1 in a 96-well culture plate. The culture plate was transferred to a 96 well-plate reader, Cytation3, set at 30°C and 100 rpm. The absorbance of cultures was recorded at OD₆₀₀ nm at regular intervals of 30 minutes over a period of 48 h. Absorbance values were used to plot the growth curve.

Protein extraction and western blotting

For protein extraction from macrophage-internalized *C. glabrata* cells, THP-1 macrophages were infected as described above. At the indicated time post-infection, macrophages were lysed in sterile chilled water, and phagocytosed *C. glabrata* cells were recovered and washed with 1X TBS buffer, transferred into 1.5 ml microcentrifuge tubes, and stored at -80°C until use. *C. glabrata* cell pellets were resuspended in 1X lysis buffer (50 mM HEPES, pH 7.5; 200 mM NaOAc, pH 7.5; 1 mM EDTA, 1 mM EGTA, 5 mM MgOAc, 5% Glycerol, 0.25% NP-40, 3 mM DTT and 1 mM PMSF) supplemented with protease inhibitor

cocktail (Roche). Zirconium beads equivalent to 100 μ L volume was added in microcentrifuge tubes and resuspended cells were lysed by 6 rounds of bead beating on a bullet blender. The sample was centrifuged at 12,000g at 4°C for 10 minutes. Supernatant was carefully transferred to a new tube, and the resultant protein sample was quantified using Biorad protein assay kit (DC protein assay kit, cat. no. 5000116), and stored in -80°C freezer. For western analysis, 25 μ g of protein samples were resolved on 12% SDS-PAGE gel and blotted on methanol activated PVDF membrane (350 mA, 75 minutes in cold room). PVDF membrane was transferred to 5% fat-free milk prepared in 1X TBST for blocking and incubated for 1 hour. Membranes were probed with appropriate primary (anti c-MYC antibody, Santa Cruz, cat. no. 9E10 and anti-Histone H3 antibody, Abcam, cat. no. ab1791) and secondary (goat anti-mouse IgG, Merck Millipore, cat. no. AP124P) antibodies, and Blots were developed by chemiluminescence based ECL western detection kit (GE Healthcare, cat. no. RPN2236) on ChemidocTM gel imaging system.

Data availability

RNAPII ChIP-seq and CgXbp1^{MYC} ChIP-seq data are available from the NCBI SRA database under the accession number PRJNA665114 and PRJNA743592, respectively.

References

- Abramova N, Sertil O, Mehta S, Lowry C V. 2001. Reciprocal regulation of anaerobic and aerobic cell wall mannoprotein gene expression in *Saccharomyces cerevisiae*. *J Bacteriol* **183**:2881–2887. doi:10.1128/JB.183.9.2881-2887.2001
- Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS. 2009. MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**:W202-8. doi:10.1093/nar/gkp335
- Benedict K, Richardson M, Vallabhaneni S, Jackson BR, Chiller T. 2017. Emerging issues, challenges, and changing epidemiology of fungal disease outbreaks. *Lancet Infect Dis* **17**:e403–e411. doi:https://doi.org/10.1016/S1473-3099(17)30443-7
- Bongomin F, Gago S, Oladele RO, Denning DW. 2017. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J fungi (Basel, Switzerland)* **3**. doi:10.3390/jof3040057
- Brown GD. 2011. Innate antifungal immunity: the key role of phagocytes. *Annu Rev Immunol* **29**:1–21. doi:10.1146/annurev-immunol-030409-101229
- Costa C, Henriques A, Pires C, Nunes J, Ohno M, Chibana H, Sá-Correia I, Teixeira M. 2013. The dual role of *Candida glabrata* drug/H⁺ antiporter CgAqr1 (ORF *CAGL0J09944g*) in antifungal drug and acetic acid resistance. *Front Microbiol* **4**:170. doi:10.3389/fmicb.2013.00170
- Costa C, Ribeiro J, Miranda IM, Silva-Dias A, Cavaleiro M, Costa-de-Oliveira S, Rodrigues AG, Teixeira MC. 2016. Clotrimazole Drug Resistance in *Candida glabrata* Clinical Isolates Correlates with Increased Expression of the Drug/H(+) Antiporters CgAqr1, CgTpo1_1, CgTpo3, and CgQdr2. *Front Microbiol* **7**:526. doi:10.3389/fmicb.2016.00526
- Erwig LP, Gow NAR. 2016. Interactions of fungal pathogens with phagocytes. *Nat Rev*

632 *Microbiol* **14**:163–176. doi:10.1038/nrmicro.2015.21

633 Fan X, Lamarre-Vincent N, Wang Q, Struhl K. 2008. Extensive chromatin fragmentation
634 improves enrichment of protein binding sites in chromatin immunoprecipitation
635 experiments. *Nucleic Acids Res* **36**:e125. doi:10.1093/nar/gkn535

636 Feng J, Liu T, Qin B, Zhang Y, Liu XS. 2012. Identifying ChIP-seq enrichment using
637 MACS. *Nat Protoc* **7**:1728–1740. doi:10.1038/nprot.2012.101

638 Hallstrom TC, Katzmann DJ, Torres RJ, Sharp WJ, Moye-Rowley WS. 1998. Regulation of
639 transcription factor Pdr1p function by an Hsp70 protein in *Saccharomyces cerevisiae*.
640 *Mol Cell Biol* **18**:1147–1155. doi:10.1128/mcb.18.3.1147

641 Hofman-Bang J. 1999. Nitrogen catabolite repression in *Saccharomyces cerevisiae*. *Mol*
642 *Biotechnol* **12**:35–73. doi:10.1385/MB:12:1:35

643 Jacobsen ID. 2014. *Galleria mellonella* as a model host to study virulence of *Candida*.
644 *Virulence*. doi:10.4161/viru.27434

645 Joffrion TM, Cushion MT. 2010. Sterol biosynthesis and sterol uptake in the fungal pathogen
646 *Pneumocystis carinii*. *FEMS Microbiol Lett* **311**:1–9. doi:10.1111/j.1574-
647 6968.2010.02007.x

648 Kasper L, Seider K, Hube B. 2015. Intracellular survival of *Candida glabrata* in
649 macrophages: immune evasion and persistence. *FEMS Yeast Res* **15**:fov042.
650 doi:10.1093/femsyr/fov042

651 Kaur R, Domergue R, Zupancic ML, Cormack BP. 2005. A yeast by any other name:
652 *Candida glabrata* and its interaction with the host. *Curr Opin Microbiol* **8**:378–384.
653 doi:10.1016/j.mib.2005.06.012

654 Kaur R, Ma B, Cormack BP. 2007. A family of glycosylphosphatidylinositol-linked aspartyl
655 proteases is required for virulence of *Candida glabrata*. *Proc Natl Acad Sci USA*
656 **104**:7628–7633. doi:10.1073/pnas.0611195104

657 Kent WJ, Zweig AS, Barber G, Hinrichs AS, Karolchik D. 2010. BigWig and BigBed:
658 Enabling browsing of large distributed datasets. *Bioinformatics* **26**:2204–2207.
659 doi:10.1093/bioinformatics/btq351

660 Lamoth F, Lockhart SR, Berkow EL, Calandra T. 2018. Changes in the epidemiological
661 landscape of invasive candidiasis. *J Antimicrob Chemother* **73**:i4–i13.
662 doi:10.1093/jac/dkx444

663 Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
664 **9**:357–359. doi:10.1038/nmeth.1923

665 Leipheimer J, Bloom ALM, Campomizzi CS, Salei Y, Panepinto JC. 2019. Translational
666 Regulation Promotes Oxidative Stress Resistance in the Human Fungal Pathogen
667 *Cryptococcus neoformans*. *MBio* **10**. doi:10.1128/mBio.02143-19

668 Lorenz MC, Bender JA, Fink GR. 2004. Transcriptional response of *Candida albicans* upon
669 internalization by macrophages. *Eukaryot Cell* **3**:1076–1087. doi:10.1128/EC.3.5.1076-
670 1087.2004

671 Lorenz MC, Fink GR. 2001. The glyoxylate cycle is required for fungal virulence. *Nature*
672 **412**:83–86. doi:10.1038/35083594

673 Mai B, Breeden L. 1997. Xbp1, a stress-induced transcriptional repressor of the
674 *Saccharomyces cerevisiae* Swi4/Mbp1 family. *Mol Cell Biol* **17**:6491–6501.
675 doi:10.1128/mcb.17.11.6491

676 Marzluf GA. 1997. Genetic regulation of nitrogen metabolism in the fungi. *Microbiol Mol*
677 *Biol Rev* **61**:17–32.

678 Miles S, Li L, Davison J, Breeden LL. 2013. Xbp1 Directs Global Repression of Budding
679 Yeast Transcription during the Transition to Quiescence and Is Important for the
680 Longevity and Reversibility of the Quiescent State. *PLOS Genet* **9**:1–14.
681 doi:10.1371/journal.pgen.1003854

682 Miyazaki H, Miyazaki Y, Geber A, Parkinson T, Hitchcock C, Falconer DJ, Ward DJ,
683 Marsden K, Bennett JE. 1998. Fluconazole resistance associated with drug efflux and
684 increased transcription of a drug transporter gene, PDH1, in *Candida glabrata*.
685 *Antimicrob Agents Chemother* **42**:1695–1701.

686 Monteiro PT, Oliveira J, Pais P, Antunes M, Palma M, Cavalheiro M, Galocha M, Godinho
687 CP, Martins LC, Bourbon N, Mota MN, Ribeiro RA, Viana R, Sá-Correia I, Teixeira
688 MC. 2020. YEASTRACT+: a portal for cross-species comparative genomics of
689 transcription regulation in yeasts. *Nucleic Acids Res* **48**:D642–D649.
690 doi:10.1093/nar/gkz859

691 Pais P, Costa C, Pires C, Shimizu K, Chibana H, Teixeira MC. 2016. Membrane Proteome-
692 Wide Response to the Antifungal Drug Clotrimazole in *Candida glabrata*: Role of the
693 Transcription Factor CgPdr1 and the Drug/H⁺ Antiporters CgTpo1_1 and CgTpo1_2.
694 *Mol Cell Proteomics* **15**:57–72. doi:10.1074/mcp.M114.045344

695 Qin L, Li A, Tan K, Guo S, Chen Y, Wang F, Wong KH. 2019. Universal plasmids to
696 facilitate gene deletion and gene tagging in filamentous fungi. *Fungal Genet Biol*
697 **125**:28–35. doi:10.1016/j.fgb.2019.01.004

698 Rai MN, Balusu S, Gorityala N, Dandu L, Kaur R. 2012. Functional Genomic Analysis of
699 *Candida glabrata*-Macrophage Interaction: Role of Chromatin Remodeling in
700 Virulence. *PLoS Pathog* **8**. doi:10.1371/journal.ppat.1002863

701 Rai MN, Borah S, Bairwa G, Balusu S, Gorityala N, Kaur R. 2013. Establishment of an in
702 vitro system to study intracellular behavior of *Candida glabrata* in human THP-1
703 macrophages. *J Vis Exp*.

704 Rodrigues CF, Rodrigues ME, Silva S, Henriques M. 2017. *Candida glabrata* Biofilms: How
705 Far Have We Come? *J fungi (Basel, Switzerland)* **3**. doi:10.3390/jof3010011

706 Rubin-Bejerano I, Fraser I, Grisafi P, Fink GR. 2003. Phagocytosis by neutrophils induces an

707 amino acid deprivation response in *Saccharomyces cerevisiae* and *Candida albicans*.
708 *Proc Natl Acad Sci USA* **100**:11007–11012. doi:10.1073/pnas.1834481100

709 Seider K, Brunke S, Schild L, Jablonowski N, Wilson D, Majer O, Barz D, Haas A, Kuchler
710 K, Schaller M, Hube B. 2011. The facultative intracellular pathogen *Candida glabrata*
711 subverts macrophage cytokine production and phagolysosome maturation. *J Immunol*
712 **187**:3072–3086. doi:10.4049/jimmunol.1003730

713 Seider K, Gerwien F, Kasper L, Allert S, Brunke S, Jablonowski N, Schwarzmüller T, Barz
714 D, Rupp S, Kuchler K, Hube B. 2014. Immune evasion, stress resistance, and efficient
715 nutrient acquisition are crucial for intracellular survival of *Candida glabrata* within
716 macrophages. *Eukaryot Cell* **13**:170–183. doi:10.1128/EC.00262-13

717 Seider K, Heyken A, Lüttich A, Miramón P, Hube B. 2010. Interaction of pathogenic yeasts
718 with phagocytes: survival, persistence and escape. *Curr Opin Microbiol* **13**:392–400.
719 doi:10.1016/j.mib.2010.05.001

720 Skrzypek MS, Binkley J, Binkley G, Miyasato SR, Simison M, Sherlock G. 2017. The
721 Candida Genome Database (CGD): incorporation of Assembly 22, systematic
722 identifiers and visualization of high throughput sequencing data. *Nucleic Acids Res*
723 **45**:D592–D596. doi:10.1093/nar/gkw924

724 Stajich JE, Harris T, Brunk BP, Brestelli J, Fischer S, Harb OS, Kissinger JC, Li W, Nayak
725 V, Pinney DF, Stoeckert CJJ, Roos DS. 2012. FungiDB: an integrated functional
726 genomics database for fungi. *Nucleic Acids Res* **40**:D675–81. doi:10.1093/nar/gkr918

727 Tan K, Wong KH. 2019. RNA polymerase II ChIP-seq-a powerful and highly affordable
728 method for studying fungal genomics and physiology. *Biophys Rev*.
729 doi:10.1007/s12551-018-00497-9

730 Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2012. Integrative Genomics Viewer (IGV):
731 high-performance genomics data visualization and exploration. *Brief Bioinform* **14**:178–

732 192. doi:10.1093/bib/bbs017

733 Wong KH, Jin Y, Moqtaderi Z. 2013. Multiplex Illumina sequencing using DNA barcoding.

734 *Curr Protoc Mol Biol* **Chapter 7**:Unit 7.11. doi:10.1002/0471142727.mb0711s101

735 Wong KH, Struhl K. 2011. The Cyc8-Tup1 complex inhibits transcription primarily by

736 masking the activation domain of the recruiting protein. *Genes Dev* **25**:2525–2539.

737 doi:10.1101/gad.179275.111

738 Yadav KK, Singh N, Rajvanshi PK, Rajasekharan R. 2016. The RNA polymerase I subunit

739 Rpa12p interacts with the stress-responsive transcription factor Msn4p to regulate lipid

740 metabolism in budding yeast. *FEBS Lett*. doi:10.1002/1873-3468.12422

741 Zhang Y, Liu T, Meyer CA, Eeckhoutte J, Johnson DS, Bernstein BE, Nusbaum C, Myers

742 RM, Brown M, Li W, Liu XS. 2008. Model-based analysis of ChIP-Seq (MACS).

743 *Genome Biol* **9**:R137. doi:10.1186/gb-2008-9-9-r137

744

Acknowledgements

We thank members of the Wong laboratory for their valuable comments throughout the study. We acknowledge the services and technical supports from the Genomics and Single Cell Analysis Core and the Drug and Development Core of the Faculty of Health Sciences at the University of Macau. This work was performed in part at the High-Performance Computing Cluster (HPCC), which is supported by the Information and Communication Technology Office (ICTO) of the University of Macau. We thank Lakhansing Pardeshi and Zhengqiang Miao, for Bioinformatics supports and Jacky Chan for technical supports on the HPC. This work was supported by the Research Services and Knowledge Transfer Office of the University of Macau (Grant number: MYRG2019-00099-FHS) and the Collaborative Research Fund Equipment Grant (C5012-15E) from the Research Grant Council, Hong Kong Government.

Author contributions

M.N.R. and K.H.W conceived the study, designed experiments and interpreted data. M.N.R. and R.R. performed experiments. M.N.R, C.P. and N.S. performed Bioinformatics analysis. K.T. performed Illumina sequencing and provided technical supports. K.T. and K.H.W. provided funding and resources. M.N.R. and K.H.W. wrote the manuscript.

Competing interest declaration

The authors declare no conflict of interest.

Figure Legends

Figure 1: *C. glabrata* mounts a dynamic chronological transcriptional response upon macrophage infection. (A) A schematic diagram showing the overall methodology used in this study. (B) Genome browser views of RNAPII ChIP-seq signals on *CgCYC1*, *CgCIT2*, *CgICL1*, *CgFTR1*, *CgTRR1* and *CgMT-I* genes at indicated time points. Numbers in the square brackets indicate the y-axis scale range of normalized RNAPII ChIP-seq signal used for the indicated genes across different datasets. (C) A heatmap showing temporal expression patterns of transcribed genes in *C. glabrata* during 0.5 to 8 h macrophage infection in a time-course experiment. The colour scale represents the Z-score of the normalized RNAPII ChIP-seq signal. (D) A table showing significantly enriched GO biological processes (P value $\leq$ 0.05) for the six groups of temporally transcribed genes. (E) A schematic diagram showing *C. glabrata* transcriptional responses (broadly classified into early, intermediate and late stages) during macrophage infection.

Figure 1-figure supplement 1: High-resolution RNAPII ChIP-seq can capture genome-wide active and temporally induced transcription activities in *C. glabrata* during macrophage infection. (A) A heatmap displaying RNAPII ChIP-seq signal over the gene body and 200 bp upstream and downstream regions for all *C. glabrata* genes after 0.5, 2, 4, 6 and 8 h of THP-1 macrophage infection. The colour scale represents the normalized RNAPII ChIP-seq signal. (B) Genome browser screenshots showing RNAPII ChIP-seq signal on selected constitutively transcribed genes. (C) Genome browser screenshots showing RNAPII ChIP-seq signal on selected temporally induced *C. glabrata* genes. Numbers in the square brackets indicate the y-axis scale range of normalized RNAPII ChIP-seq signal used for the indicated genes across different datasets.

Figure 1-figure supplement 2: *C. glabrata* undergoes cell cycle arrest upon macrophage phagocytosis. (A) Heatmaps showing the expression pattern for cell cycle and DNA damage checkpoint genes during macrophage infection. The colour scale represents the Z-score of the normalized RNAPII ChIP-seq signal. (B) A density plot displaying the distribution of *C. glabrata* cells at different cell cycle stages based on FACS analysis at 2 h after THP-1 macrophage infection.

Figure 1-figure supplement 3: Virulence-centric biological processes are temporally activated in *C. glabrata* at different stages of macrophage infection. (A-G) Heatmaps displaying the expression pattern for genes associated with biological processes (A) Adhesion, (B) DNA repair, (C) Response to oxidative stress, (D) Autophagy, (E) TCA cycle, (F) Amino acid biosynthesis and (G) Iron homeostasis during THP-1 macrophage infection. The colour scale represents the Z-score of the normalized RNAPII ChIP-seq signal.

Figure 1-figure supplement 4: Genes encoding histones or proteins involved in chromatin modification and remodelling are transcriptionally induced in *C. glabrata* during the early stages of macrophage infection. (A&B) Heatmaps showing the expression pattern for (A) histone H2A, H2B, H3, and H4 genes and putative (B) chromatin and histone modifiers genes during THP-1 macrophage infection. The colour scale represents the Z-score of the normalized RNAPII ChIP-seq signal.

Figure 1-figure supplement 5: TIR family genes for sterol uptake displaying co-expression during macrophage infection. Genome browser screenshots showing RNAPII ChIP-seq profile on putative TIR family genes for sterol uptake during THP-1 macrophage infection.

Figure 1-figure supplement 6: Correlation between independent biological repeats of the RNAPII ChIP-seq experiment for wildtype and the *Cgxbp1*Δ mutant. (A&B)

Scatterplots showing RNAPII ChIP-seq signals between two independent biological repeats for (A) wildtype and (B) the *Cgxbp1*Δ mutant at the indicated times. The correlation coefficient (r) for each comparison is presented.

Figure 2: CgXbp1 is central in orchestrating the dynamic transcriptional response of *C. glabrata* during macrophage infection. (A) A heatmap showing temporal expression patterns of *C. glabrata* transcription factor genes transcribed during THP-1 macrophage infection. Colour scale represents the Z-score of the normalized RNAPII ChIP-seq signal. (B) A heatmap showing temporal expression patterns of transcribed genes in the *Cgxbp1*Δ mutant during 0.5 to 8 h THP-1 macrophage infection in a time-course experiment. (C) An UpSet plot showing the number of genes induced at the indicated time points in WT and the *Cgxbp1*Δ mutant during THP-1 macrophage infection.

Figure 2-figure supplement 1: CgXbp1 is a key transcription regulator of the temporal transcriptional response of *C. glabrata* during macrophage infection. (A) A regulatory network of Xbp1 and Hap3 for the orthologue of macrophage infection-induced genes in *S. cerevisiae* based on published regulatory information available on the PathoYeasttract database. Green, red, and black arrows indicate positive, negative, and unspecified regulation, respectively. Solid and dashed lines represent DNA binding or expression-based evidence, respectively. (B) A regulatory network of CgXBPI for a subset of infection-induced TF genes based on published regulatory information available on the PathoYeasttract database. (C)

Venn diagram of actively transcribing genes in wildtype and the *Cgxbp1*Δ mutant during macrophage infection.

Figure 3: Loss of *CgXBPI* affects the expression level and timing of multiple genes of diverse physiological pathways upon macrophage phagocytosis. (A & B) Heat maps showing transcription activities of genes belonging to (A) TCA and (B) amino acid biosynthesis during THP1 macrophage infection in wildtype and the *Cgxbp1*Δ mutant.

Figure 3-figure supplement 1: *CgXbp1* is essential for the chronological transcriptional response of *C. glabrata* during macrophage infection. A table summarising the enrichment of GO-terms among the transcribed *C. glabrata* genes at the indicated time points during THP-1 macrophage infection by wildtype and the *Cgxbp1*Δ mutant. “Y” indicates a statistically significant enrichment (P value < 0.05) in a given GO term, while a blank box means no significant enrichment.

Figure 4: Loss of *CgXBPI* affects *C. glabrata* proliferation in human macrophages and attenuates virulence in the *Galleria mellonella* model of candidiasis. (A) Bar chart of CFUs obtained from *C. glabrata* cells harvested from THP-1 macrophages at indicated time points. Error bars represent the standard error of the mean (±SEM) from 3 independent experiments. Statistical significance was determined by two-sided unpaired Student’s *t*-test, *P value ≤ 0.05, **P value ≤ 0.01. **(B)** Cumulative survival curve of *G. mellonella* larvae infected with indicated *C. glabrata* strains. At least 16 larvae were used in each of three independent infection experiments. The graph represents the per cent survival of larvae infected with the indicated strains from three independent infection experiments. Statistical significance was determined by a two-sided unpaired Student’s *t*-test, **P value ≤ 0.01.

Figure 4-figure supplement 1: Phagocytosis rate is not affected by CgXbp1 deletion. Bar diagram displaying colony forming units obtained before macrophage infection and 2 h post-infection. Error bars represent mean $\pm$ standard deviation from three independent experiments.

Figure 5: CgXbp1 binds to a large number of genomic locations *in vivo*. (A & B) Western blot analysis of CgXbp1 expression (A) during THP1 macrophage infection and (B) at exponential (Log) and quiescence phases (Day 2, 3 and 4) (Figure 5-Source data). (C & D) Representative genome-browser screenshots showing CgXbp1^{MYC} ChIP-seq signal on (C) a chromosomal region and (D) the promoter of select genes. (E) A Heat map of ChIP-seq signal on promoters of CgXbp1 target genes. The colour scale indicates normalized ChIP-seq signal on 3 kb upstream and downstream flanking regions from the transcription start site (TSS) of the target genes. (F) Consensus DNA binding motifs enriched from CgXbp1 bound promoter sequences in CgXbp1 target genes by MEME analysis. (G) Genome-browser screenshots displaying CgXbp1 binding on *CgMIG1*, *CgADRI*, *CgCST6*, and *CgHAP4* promoters during the quiescence phase.

Figure 5-figure supplement 1: Protein sequence comparison between CgXbp1 and ScXbp1. Pairwise sequence alignment of amino acid sequences of *C. glabrata* ORF *CAGL0G02739g* and *S. cerevisiae* Xbp1. Black and grey shaded amino acid represents identical and similar amino acids, respectively. Boxed amino acid sequence marks the DNA binding domain.

Figure 5-figure supplement 2: Biological replicates from two independent experiments for CgXbp1^{MYC} ChIP-seq displayed a good correlation. (A) Scatterplot showing correlation of ChIP-seq signal intensity at CgXbp1^{MYC} peak summits and 200 bp flanking regions between two independent biological repeats during quiescence phase. (B) Heatmap showing ChIP-seq signals on CgXbp1^{MYC} peak summits and 1500 bp flanking regions for two independent biological repeats and the input control.

Figure 5-figure supplement 3: The two enriched motifs identified from CgXbp1 binding sites show good homology with *S. cerevisiae* Xbp1 and Azf1 DNA binding recognition motifs. (A&B) TOMTOM suite identifies two motifs in *S. cerevisiae* transcription factors, Xbp1 (A) and Azf1 (B), conserved and similar to CgXbp1.

Figure 6: CgXbp1 regulates fluconazole resistance in *C. glabrata*. (A) Genome-browser screenshots showing CgXbp1^{MYC} binding on the promoter of genes encoding drug efflux pumps, *CgPDH1*, *CgPDR13*, and *CgQDR2*. (B) Serial dilution spotting assay on YPD medium in the presence of different fluconazole concentrations (0, 32 or 64 µg/mL). (C) Growth curve of wildtype and *Cgxbp1Δ* mutant in YPD medium in the presence or absence of fluconazole (64 µg/mL). Error bars represent mean ± SEM from four independent experiments. (D) Bar graph displaying CFUs/mL obtained for indicated strains on YPD agar plates in the presence or absence of fluconazole (64 µg/mL) post 3 days of spread plating. Error bars represent mean ± SEM from three independent experiments. Statistical significance was determined by a two-sided unpaired Student's *t*-test, *P value ≤ 0.05.

Figure 7: A schematic diagram illustrating a proposed model for the regulation of transcriptional responses in *C. glabrata* during macrophage infection. A model for the

915 regulation of the chronological transcriptional responses of *C. glabrata* during macrophage
916 infection. The model was created in BioRender.com.
917

918 **Table 1:** Constitutively transcribed or temporally induced *C. glabrata* transcription factor
 919 genes during macrophage infection.

920

Table 1: Constitutively transcribed or temporally induced <i>C. glabrata</i> transcription factor genes during macrophage infection				
Temporally induced				
Group number	Cg common name	Cg ORF name	Sc common name	Sc gene desc
Group:1:(N=4)	CAGL0G02739g	CAGL0G02739g	XBP1	XhoI site-Binding Protein
Group:1:(N=4)	CAGL0L03157g	CAGL0L03157g	DAL80	Degradation of Allantoin
Group:1:(N=4)	CAGL0J04400g	CAGL0J04400g	HAP3	Heme Activator Protein
Group:1:(N=4)	CAGL0F00561g	CAGL0F00561g	RPA12	RNA Polymerase A
Group:2:(N=4)	CAGL0K06413g	CAGL0K06413g	STP1	Species-specific tRNA Processing
Group:2:(N=4)	CAGL0E00737g	CAGL0E00737g	HMO1	High MOBility group (HMG) family
Group:2:(N=4)	MET28	CAGL0K08668g	MET28	METHionine
Group:2:(N=4)	CAGL0J03608g	CAGL0J03608g	HCM1	High-Copy suppressor of Calmodulin
Group:3:(N=5)	RTG1	CAGL0C05335g	RTG1	ReTroGrade regulation
Group:3:(N=5)	CAGL0J01177g	CAGL0J01177g	ABF1	ARS-Binding Factor 1
Group:3:(N=5)	CAGL0K04543g	CAGL0K04543g	SPT4	SuPpressor of Ty's
Group:3:(N=5)	HAP4	CAGL0K08624g	HAP4	Heme Activator Protein
Group:3:(N=5)	CAGL0G07249g	CAGL0G07249g	YHP1	Yeast Homeo-Protein
Group:4:(N=4)	CAGL0L07480g	CAGL0L07480g	NRG2	Negative Regulator of Glucose-controlled genes
Group:4:(N=4)	MIG1	CAGL0A01628g	MIG1	Multicopy Inhibitor of GAL gene expression
Group:4:(N=4)	CAGL0G08646g	CAGL0G08646g	POG1	Promoter Of Growth
Group:4:(N=4)	CAGL0K02145g	CAGL0K02145g	COM2	Cousin of Msn2
Group:5:(N=17)	RME1	CAGL0K04257g	RME1	Regulator of MEiosis
Group:5:(N=17)	CAGL0M07634g	CAGL0M07634g	SOK2	Suppressor Of Kinase
Group:5:(N=17)	CAGL0M01716g	CAGL0M01716g	TEC1	Transposon Enhancement Control
Group:5:(N=17)	CAGL0F07909g	CAGL0F07909g	TBS1	ThiaBendazole Sensitive
Group:5:(N=17)	UPC2B	CAGL0F07865g	UPC2	UPTake Control
Group:5:(N=17)	ZAP1	CAGL0J05060g	ZAP1	Zinc-responsive Activator Protein
Group:5:(N=17)	CAGL0C02519g	CAGL0C02519g	MIG3	Multicopy Inhibitor of Growth
Group:5:(N=17)	HAP5	CAGL0K09900g	HAP5	Heme Activator Protein
Group:5:(N=17)	CAGL0E04312g	CAGL0E04312g	STP2	protein with similarity to Stp1p
Group:5:(N=17)	CAGL0B03421g	CAGL0B03421g	HAP1	Heme Activator Protein
Group:5:(N=17)	HAA1	CAGL0L09339g	HAA1	Homolog of Ace1 Activator
Group:5:(N=17)	GAT1	CAGL0K07634g	GAT1	Transcriptional activator of nitrogen catabolite repression genes
Group:5:(N=17)	YAP6	CAGL0M08800g	YAP6	Yeast homolog of AP-1
Group:5:(N=17)	GLM6	CAGL0J01595g	#N/A	#N/A
Group:5:(N=17)	AFT1	CAGL0H03487g	AFT1	Activator of Ferrous Transport
Group:5:(N=17)	YAP3b	CAGL0M10087g	#N/A	#N/A
Group:5:(N=17)	CAGL0E03762g	CAGL0E03762g	RIM101	Regulator of IME2
Group:6:(N=5)	AP5	CAGL0K08756g	YAP5	Yeast AP-1
Group:6:(N=5)	GCN4	CAGL0L02475g	GCN4	General Control Nonderepressible
Group:6:(N=5)	CAGL0E05566g	CAGL0E05566g	TYE7	Ty1-mediated Expression
Group:6:(N=5)	RPN4	CAGL0K01727g	RPN4	Regulatory Particle Non-ATPase
Group:6:(N=5)	CAGL0C01551g	CAGL0C01551g	TOS8	Target Of Sbf
Constitutively transcribed				
Group	Cg common name	Cg ORF name	Sc common name	Sc gene desc
Constitutively transcribed	PHO2	CAGL0L07436g	PHO2	PHOSphate metabolism
Constitutively transcribed	AP1	CAGL0H04631g	YAP1	Yeast AP-1
Constitutively transcribed	CAGL0M04983g	CAGL0M04983g	MBF1	Multiprotein Bridging Factor
Constitutively transcribed	MSN4	CAGL0M13189g	MSN4	Multicopy suppressor of SNF1 mutation
Constitutively transcribed	CAGL0E00891g	CAGL0E00891g	STB3	Sin Three Binding protein
Constitutively transcribed	CAD1	CAGL0F03069g	CAD1	CADmium resistance
Constitutively transcribed	CAGL0A04257g	CAGL0A04257g	TOD6	Twin Of Dot6p
Constitutively transcribed	CAGL0I08635g	CAGL0I08635g	BUR6	Bypass UAS Requirement
Constitutively transcribed	YAP7	CAGL0F01265g	YAP7	Yeast AP-1
Constitutively transcribed	CAGL0L02013g	CAGL0L02013g	IXR1	Intrastrand cross (X)-link Recognition
Constitutively transcribed	CAGL0M01474g	CAGL0M01474g	NCB2	Negative Cofactor B
Constitutively transcribed	CAGL0F06259g	CAGL0F06259g	ARG80	ARGINine requiring
Constitutively transcribed	SWI5	CAGL0E01331g	SWI5	SWItching deficient
Constitutively transcribed	CAGL0M09955g	CAGL0M09955g	SFP1	Split Finger Protein

921

Table 2: Table of significantly enriched and non-redundant GO-terms for biological processes among CgXbp1 target genes during quiescence phase, grouped based on biological functions.

	GO Term	P value	Genes in the background	CgXbp1 bound genes
Transport	transmembrane transport	1.19E-06	320	47
	carboxylic acid transport	3.13E-04	69	14
	acetate transport	1.40E-03	4	3
	xenobiotic transport	2.48E-03	9	4
	oxygen transport	5.16E-03	2	2
	fluconazole transport	5.16E-03	2	2
	drug transport	8.20E-03	12	4
	ammonium transport	1.04E-02	7	3
	amino acid transport	1.19E-02	44	8
	glucose transmembrane transport	1.57E-02	8	3
Response to stress	response to nutrient	6.93E-04	58	12
	cellular response to chemical stress	1.36E-03	184	25
	response to nitrosative stress	1.46E-03	8	4
	response to acetate	3.31E-03	5	3
	cellular response to oxidative stress	5.09E-03	130	18
	response to osmotic stress	1.07E-02	99	14
	response to oxidative stress	1.13E-02	151	19
	response to salt stress	1.70E-02	38	7
	response to pH	1.76E-02	85	12
	response to acidic pH	4.99E-02	12	3
Metabolism	carbohydrate metabolic process	1.50E-05	184	30
	carbohydrate biosynthetic process	2.16E-04	59	13
	glucan biosynthetic process	4.38E-04	27	8
	carbon catabolite regulation of transcription	4.89E-04	41	10
	glutamine family amino acid biosynthetic process	6.23E-04	22	7
	nitrogen utilization	9.73E-04	12	5
	regulation of glycolytic process	1.47E-02	3	2
	tricarboxylic acid cycle	3.51E-02	26	5
Biofilm	adhesion of symbiont to host	4.38E-04	27	8
	fungal-type cell wall organization or biogenesis	7.39E-04	227	30
	positive regulation of cell growth	1.12E-03	24	7
	biofilm formation	1.67E-03	81	14
	biological adhesion	1.06E-02	70	11

Figure 1

Figure 2

Figure 3

929

930

Figure 4

Figure 5

Figure 6

Figure 7

937

938 **Figure 1-figure supplement 1**

Figure 1-figure supplement 2

942

943 **Figure 1-figure supplement 3**

Figure 1-figure supplement 4

Figure 1-figure supplement 5

A

B

950

951 **Figure 1-figure supplement 6**

952

A

B

C

Figure 2-figure supplement 1

GO-term	WT			Cgxbp1Δ		
	0.5h	2h	4h	0.5h	2h	4h
small molecule metabolic process		Y	Y	Y	Y	
generation of precursor metabolites and energy		Y	Y	Y		
ATP metabolic process		Y	Y	Y		
energy derivation by oxidation of organic compounds		Y	Y	Y		
aerobic respiration		Y	Y	Y		
ribonucleotide biosynthetic process		Y		Y	Y	
purine-containing compound biosynthetic process		Y		Y	Y	
ribose phosphate biosynthetic process		Y		Y	Y	
purine nucleotide biosynthetic process		Y		Y	Y	
oxidation-reduction process		Y	Y	Y		
ATP synthesis coupled electron transport		Y		Y		
lysine biosynthetic process		Y		Y		
cellular amino acid biosynthetic process		Y		Y		
ion transmembrane transport		Y		Y		
small molecule biosynthetic process		Y		Y		
alpha-amino acid biosynthetic process		Y		Y		
carboxylic acid biosynthetic process		Y		Y		
histidine biosynthetic process		Y		Y		
transmembrane transport		Y		Y		
aspartate family amino acid biosynthetic process		Y		Y		
tricarboxylic acid cycle		Y		Y		
acetate catabolic process		Y	Y	Y		
nucleoside triphosphate biosynthetic process		Y	Y	Y		
carbon utilization		Y	Y	Y		
ribonucleoside triphosphate biosynthetic process		Y	Y	Y		
trehalose biosynthetic process			Y	Y		
response to oxidative stress			Y	Y		
response to water			Y	Y		
carbohydrate biosynthetic process			Y	Y		
drug metabolic process		Y	Y			
'de novo' IMP biosynthetic process		Y		Y	Y	
nucleoside phosphate biosynthetic process		Y		Y	Y	
ribonucleoside monophosphate biosynthetic process		Y		Y	Y	
fumarate metabolic process		Y		Y	Y	
nucleotide biosynthetic process		Y		Y	Y	
nucleosome assembly		Y	Y	Y	Y	
monocarboxylic acid metabolic process		Y	Y			

956

957 **Figure 3-figure supplement 1**

958

959 **Figure 4-figure supplement 1**

```

CAGL0G02739g  -----MRLTDSPLDDYQRLCFSQSMDLKD-----RVRHQTYKSNLAPT  38
XBP1  MKYPAFSINS DTVHLTDNPLDDYQRLYLVSVDL-RDSPPASFSAGLNIRKVNYKSSIA--  57

CAGL0G02739g  MHS LVSHLKPQLNHSSMTAADSKSDKG-----LMCFEYQFPDLVAQSQBTADSNTHI  90
XBP1  -----AQFTHPNFIIISARDAGNGE EAAAQNV LNCFEYQFPNL-----QTIQSLVHE  103

CAGL0G02739g  QYESGNRPYGLNAETDTPTAHRMSTEPSDGALDATLLLPELQQT---SPRLTGHKRGY  146
XBP1  QTL-----LSQLASSATPHSALHLHDKNILMGKIILPSRSNKTTPVSAASP-TKQEKKAL  155

CAGL0G02739g  SSPSGPNGT-TISKNQQFRISNQSN--THKFINNNCILWTFNFDKNEPASSENFVFTTG  202
XBP1  STASRENATSSLTKNQQFKLTMDHNLINDKLINPNNCVIWVSH-----DSGYVEMTG  207

CAGL0G02739g  IWRLYQDVMKGLIQIPRKGKTS-KERQDFCQEFDFIMSNCFCDQLKTAKDQSDPV---  258
XBP1  IWRLYQDVMKGLINLPRGDSVSTSQQQFFCKAEFEKILSFCFYNHSSFTSESSSVLSS  267

CAGL0G02739g  DYSRKFFHRRRSTGSI SFTQKTKSNNSVTSEN SGNTHSTSNSSVNYLDFHWFDISEKVR SQ  318
XBP1  STSSPFKRRTSTGS-----TFLDANASSSSSTSSSTQANNYIDFWHNNIKPELRDL  316

CAGL0G02739g  IFEQFKQHLEKDRNVDCSTIPKAE--EYIQRIRGGYIKIQGTWVPWYIAKLICIRFCFPI  376
XBP1  ICQSYKDFLINELGPDQIDLPNLNPNANFTKRIRGGYIKIQGTWLPMEISRLLCLRFCFPI  376

CAGL0G02739g  RYLLVPIFGEQFPVEECENYFFNVHMTNLKEFLENEHLINTVRRRQKSTG-----LATPPF  431
XBP1  RYFLVPIFGPD FPKDCESWYL-AH-----QNVTFASSTTGAGAATAATAAA  421

CAGL0G02739g  LTSSTFGNGSIDSPFSDEREEMEISPFHKYSNNLSFNYSPRMRKRFRSDNNLSPMSPSTK  491
XBP1  NTSTNFTSTAV-----ARPRQKPRPRPRQRSTSMSSHKAQK  457

CAGL0G02739g  LAFRKTPPIREEPEPYEHYTS GFSTIDN-KLSTN--NTVKEETHPYRKRA LSWSYTQNTS  548
XBP1  LVIEDALP-----SFDSEFVENLGLSSNDKNFIKKNS---KRQKSSSTYTSQTS  501

CAGL0G02739g  QRTKLPPISSLIESINWKTNSDHVNVSEPTPTVQTL SKLASFYTTTRGHKYSYPSNIC---  605
XBP1  S-----PI-----GPRDPTVQILSNLASFYNTGHRYSYPGNIYIPQ  538

CAGL0G02739g  -TVSSNEHNK LASPVKNSNMETFLTARENQTGSHHYNHI-----TEPHMKHNYIRQRD  657
XBP1  QRYSLPPPNNQLSSPQRQLN-----YTYDHLHPVPSQYQSP--RHYNVPSSP  582

CAGL0G02739g  MVPRVISDSLYPDLYKENKVD FICLRNQTHLQENV-----  693
XBP1  IAP---APPTFPQ-PYGDDHYHFLKYASEVYKQNNQRPAHNTNTNMDTSFSPRANNSLNN  638

CAGL0G02739g  -----  693
XBP1  FKFKTNSKQ  647

```

Figure 5-figure supplement 1

A

B

Figure 5-figure supplement 2

Supplementary Files

Supplementary File 1 - List of actively transcribing genes in wildtype *C. glabrata* upon macrophage infection

Supplementary File 2 - List of GO-terms enriched from temporally induced genes in wildtype *C. glabrata* in response to macrophage infection

Supplementary File 3 – Gene regulatory associations between indicated TFs and the macrophage infection-induced genes reported in the PathoYeasttract database

Supplementary File 4 – List of orthologues for the macrophage infection-induced TF and non-TF genes of *C. glabrata* previously shown to have a regulatory association with Xbp1 or Hap3 in *S. cerevisiae* obtained from the PathoYeasttract database

Supplementary File 5 – List of actively transcribing genes in *Cgxbp1Δ* mutant upon macrophage infection

Supplementary File 6 - List of GO-terms enriched from temporally induced genes in *Cgxbp1Δ* in response to macrophage infection

Supplementary File 7 - MACS2 output metadata displaying the coordinates of CgXbp1 binding sites during quiescence phase

Supplementary File 8 - List of CgXbp1 target genes displaying CgXbp1 binding during quiescence phase

991

992 **Supplementary File 9** - List of CgXbp1 target genes with two consensus DNA binding
993 motifs

994

995 **Supplementary File 10** - List of GO-terms for biological processes enriched from CgXbp1
996 targets in quiescence phase

997

998 **Supplementary File 11** – List of TF-encoding genes with CgXbp1 binding at their promoters
999

1000 **Additional files**

1001

1002 **Figure 5-Source data** – Western blot original image files used in Figures 5A & B